Sexual selection does not increase the rate of compensatory adaptation to a

mutation influencing a secondary sexual trait in *Drosophila melanogaster*

Christopher H. Chandler[1,2], Anna Mammel[1,3], Ian Dworkin[1,4]

[1]Department of Integrative Biology, BEACON Center for the Study of Evolution in

Action, Michigan State University, East Lansing, MI, United States

[2]Department of Biological Sciences, State University of New York at Oswego, Oswego,

NY, United States

[3]Department of Cell, Developmental & Cancer Biology, Oregon Health and Science

University, Portland, OR, United States

[4]Department of Biology, McMaster University, Hamilton, Ontario, Canada

Abstract

Theoretical work predicts that sexual selection can enhance natural selection,

increasing the rate of adaptation to new environments and helping purge harmful

mutations. While some experiments support these predictions, remarkably little work

has addressed the role of sexual selection on compensatory adaptation—populations'

ability to compensate for the costs of deleterious alleles that are already present. We

tested whether sexual selection, as well as the degree of standing genetic variation,

affect the rate of compensatory evolution in experimental populations of *Drosophila*

melanogaster. These populations were fixed for a spontaneous mutation causing mild

abnormalities in the male sex comb, which is important for mating success; in some

populations, the opportunity for mate choice was limited by experimentally enforced

monogamy. We first showed that this mutation is deleterious, and that its expression is

influenced by genetic background variation. Although populations did show some

compensatory adaptation with respect to reduction in the morphological abnormalities in

the sex comb, the amount of phenotypic suppression of the morphological defects did

not depend on the opportunity for sexual selection. Sexual selection, therefore, may not

always enhance natural selection; instead, the interaction between these two forces

may depend on additional factors.

Introduction

Sexual selection has important impacts on many aspects of how organisms evolve, including on speciation rates and the degree of sexual dimorphism (e.g., Masta and Maddison 2002; Ellis and Oakley 2016). While it was once considered that sexual selection may act independently or even antagonistically to other components of natural selection (e.g. viability and fecundity), recent work has shown that these can operate concordantly. For instance, sexual selection may influence how organisms respond to selective pressures in the short term, influencing how quickly populations adapt to novel environments, in particular when the population begins at a distance from an optimum (Long *et al.* 2012). Additionally, some models predict that sexual selection should help populations filter out harmful mutations more rapidly than selection on other fitness components (viability and fecundity selection) alone (Agrawal 2001). This prediction is based partly on the observation that sexual displays are often correlated with overall condition. Any mutation that reduces an organism's nonsexual fitness is therefore also likely to affect its mating success (Rowe and Houle 1996; Chandler *et al.* 2013) or even its success in sperm competition (Clark *et al.* 2012). Sexual selection against these mutations, usually in males, should therefore enhance natural selection against them.

Empirical support for this scenario has been mixed. In some studies testing these predictions, evidence supported a role for sexual selection in purging deleterious mutations or accelerating adaptation (Radwan 2004; Sharp and Agrawal 2008; Hollis *et al.* 2009; Jarzebowska and Radwan 2010; McGuigan *et al.* 2011; Long *et al.* 2012; Almbro and Simmons 2014; Lumley *et al.* 2015; Grieshop *et al.* 2016; Jacomb *et al.* 2016). However, a handful of studies also contradict these predictions (Hollis and Houle

2011; Plesnar *et al.* 2011; Arbuthnott and Rundle 2012; Cabral and Holland 2014). The

different outcomes in at least some of those cases may be due to the costs of sexual

conflict, which is often confounded with sexual selection; any benefits from sexual

selection's ability to purge deleterious mutations in those cases may have been

canceled out by the reduction in fecundity caused by sexual conflict, such as male-

induced harm (Hollis and Houle 2011; Arbuthnott and Rundle 2012; Chenoweth *et al.*

2015). Some studies also focused on large-effect mutations or exposed populations to

strong mutagens (Hollis and Houle 2011; Plesnar *et al.* 2011; Arbuthnott and Rundle

2012; Cabral and Holland 2014), so they may not be representative of how selection

acts in nature, where most segregating deleterious mutations are more likely to have

milder effects.

Of course, deleterious mutations are not always purged by selection; they can

also increase in frequency and occasionally become fixed via drift, hitchhiking or

antagonistic pleiotropy, especially if their effects on fitness are only mildly deleterious

(and in populations with a small effective population size). In those cases where the

deleterious alleles are difficult for selection to purge, alleles at other loci that

compensate epistatically for the fitness costs of these fixed deleterious alleles may be

favored by selection. There is good evidence of compensatory adaptation in both

microbial and multicellular organisms. For instance, alleles conferring antibiotic

resistance are sometimes costly in the absence of antibiotics, but compensatory

mutations can reduce those costs (Reynolds; Maisnier-Patin *et al.* 2002; Comas *et al.*

2012). In the blowfly, diazinon resistance via alleles at the *Rop-1* gene had negative

pleiotropic effects, increasing fluctuating asymmetry, but these effects were ultimately

compensated by modifiers (McKenzie and Clarke 1988; Davies *et al.* 1996).

Additionally, sex chromosome dosage compensation could also be considered a form of

compensatory adaptation, having likely evolved in response to loss-of-function

mutations on Y or W chromosomes (Charlesworth 1978).

Although sexual selection has received a lot of attention as a possible influence

on the rate of purging of deleterious mutations, the role of sexual selection in

compensatory evolution remains largely unexplored. Nevertheless, we might similarly

predict that sexual selection can also accelerate compensatory adaptation, especially if

sexual displays are condition dependent. In one study (Pischedda and Chippindale

2005), the *nub^1* mutation, which drastically reduces the size of the wing, resulting in an

inhibition of males' ability to generate courtship songs, was fixed in experimental

populations of *Drosophila melanogaster*. As predicted, this mutation had greater fitness

costs in males than it did in females, but males also showed greater compensatory

fitness recovery over 180 generations (albeit without compensating for the effects on

wing morphology directly), providing some support that sexual selection may enhance

compensatory adaptation. However, this study was not replicated (only a single

lineage), since the *nub^1* populations were originally generated for other purposes.

Clearly, more study is needed on whether sexual selection can speed up compensatory

adaptation.

In this study, we address the question of whether sexual selection can impact the

rate of compensatory evolution using experimental evolution in *Drosophila*

melanogaster. We chose a mutation in the *sex combs distal* gene (*scd^1*) (Boube *et al.*

1997; Randsholt and Santamaria 2008), a spontaneous, partially penetrant mutation

affecting the development of the male sex comb, a structure that is critical for male

mating success (Ng and Kopp 2008) and rapidly evolving across *Drosophila* species

(Atallah *et al.* 2009, 2012; Kopp 2011; Malagón *et al.* 2014). In addition to further

mapping and characterizing the effects of this mutation across different wild type

genetic backgrounds, we find that despite a general compensatory response, we

observed no evidence that sexual selection influenced the rate of compensatory

evolution for the sex comb phenotypes.

Methods

Mapping scd[1]—

sex combs distal[1] (*scd[1]*) is a spontaneous X-linked allele resulting in ectopic sex comb

bristles on the second tarsal segment of the prothoracic leg in males (Boube *et al.*

1997). In the base stock, it has incomplete penetrance, with only about 70% of males

showing the ectopic sex comb bristles, and no visible phenotype in females. In our

populations, males also sometimes exhibited minor defects in the primary sex comb,

such as a gap or partially untransformed bristles (Figure 1).

The identity of the gene and molecular lesion of this allele are unknown, although

some previous recombination mapping suggested it was near 1-30.6, and that a local

duplication of the 8C-9B region of the X chromosome onto the Y (*DP(1:Y)FF*), could

partially rescue the phenotype of *scd[1]* (Santamaria and Randsholt 1995; Randsholt and

Santamaria 2008). We attempted to further fine map the gene through duplication

mapping. Virgin female flies of strain BDSC 5070 (*y[1] scd[1] ras[1] v[1] f[1]*) were crossed to

males of strains carrying duplicated segments of the X chromosome translocated onto

the Y chromosome or chromosome III (Table 1; Cook *et al.* 2010; Venken *et al.* 2010),

and the male offspring were scored for the presence of mutant phenotypes, such as the

ectopic sex comb or disruptions in the primary sex comb. Assuming scd^1 is a recessive

loss-of-function allele, if the duplicated segment contains a functional wild-type copy of

the *scd* gene, then the mutant phenotype would be rescued and no male offspring from

these crosses would show sex comb defects. Because scd^1 is only partially penetrant,

we scored numerous male offspring from each cross.

Influence of genetic background on penetrance and expressivity—

To determine the extent of genetic variation for the penetrance and expressivity

of scd^1, we crossed virgin female y^1 scd^1 ras^1 v^1 f^1 flies to males of a randomly chosen

subset of Drosophila Genetic Reference Panel lines (Mackay *et al.* 2012). When the

adult F1 offspring eclosed, we fixed specimens in 70% ethanol, and then mounted male

prothoracic legs in 70% glycerol and scored them for the presence of ectopic sex combs

on the second tarsal segment, abnormalities in the primary sex comb, and primary sex

comb tooth number. These crosses only examine each wild-type genetic background in

a heterozygous state, and thus it is expected that this will underestimate the actual

degree of background dependence, as recessive effects of alleles in each background

will not be captured.

Introgression of scd^1—

To generate populations for experimental evolution, we introgressed the scd^1

mutation into FVW, a domesticated lab-maintained population founded from flies

collected from Fenn Valley, MI in 2010. The FVW population was maintained in

population cages with 10 bottles for egg-laying each generation for approximately 10

generations prior to beginning introgressions (Chari *et al.* 2017).

To begin the introgression (Supplementary Figure 1A), virgin y^1 scd^1 ras^1 v^1 f^1

females were crossed to FVW males. The heterozygous F1 females were then

backcrossed to FVW males. From the F2 offspring, we selected males showing the scd^1

phenotype, which were thus hemizygous for scd^1, but with phenotypically wild-type

eyes. We then crossed these males to virgin FVW females, to obtain female offspring

heterozygous for scd^1 in a partial FVW background. We then crossed these females to

FVW males, and selected males with ectopic sex combs, but not yellow bodies. Each

male from these crosses thus carries an independently derived X chromosome with

scd^1 in a random FVW background.

We obtained eight such males and crossed them to FVW females. From these

crosses, we obtained virgin females heterozygous for scd^1 in an FVW background. The

first of these virgin scd^1/scd^+ females to emerge were crossed with FVW males to obtain

more scd^1 males with a mostly FVW background. The later-emerging scd^1/scd^+ females

were kept isolated at cooler temperatures (18°C) until the scd^1 males from the previous

cross emerged. We then crossed the scd^1/scd^+ females to the scd^1 males. Finally, we

set up sib matings among the offspring of these crosses, using only males hemizygous

for scd^1 (showing a sex comb phenotype) and females of unknown genotype (either

scd^1/scd^+ or scd^1/scd^1). Of those crosses, we kept those in which the mother was

inferred to be homozygous for scd^1, in which nearly all male progeny displayed scd^1

phenotypes. This allowed us to establish a homozygous scd^1 line with an FVW genetic

background, which we designated as *scd**. *scd** carried at least four independently

derived X chromosomes with *scd^1* in an FVW background (Supplementary Figure 1A).

To introduce further genetic diversity (from the FVW population) into *scd**, *scd**

males were crossed to wild-type FVW females to obtain heterozygous *scd^1* females with

additional genetic material from the FVW background. These females were then

backcrossed to FVW males. Five replicate backcrosses were set up in culture bottles,

each with 25-30 FVW males and 25-30 females, heterozygous for *scd^1* and for alleles

from the FVW background. We then selected males with the *scd^1* phenotype, and

backcrossed them to virgin *scd** females, in six replicate bottles each containing 20-25

males and 20-25 females, to maintain *scd^1* while introducing additional genetic diversity

from the FVW population. This whole cycle was then repeated once to establish the

*scd*** base population for experimental evolution (Supplementary Figure 1B).

Fitness effects of scd^1—

To test whether the *scd^1* allele was deleterious, we tracked changes in the

frequency of the *scd^1* phenotype in polymorphic populations with the *scd^1* allele at 0.7

initial frequency. We initiated four replicate populations, each consisting of 70 *scd***

males, 70 *scd*** females, 30 FVW males, and 30 FVW females. Populations were

placed in population cages with four culture bottles for mating and oviposition for five

days, after which the flies were discarded and the bottles transferred to fresh cages at

24°C. After adult flies began emerging, they were allowed to mate for three to four days.

The old bottles were then removed, and fresh bottles were placed in the cage for egg

laying. After two days of egg laying, the flies were discarded and the bottles moved to

fresh cages. This cycle was repeated for a total of nine generations.

For each of the first five generations, and at generation nine, we scored male sex

comb phenotypes. 50 males were picked randomly, and the first prothoracic legs from

each male were mounted on glass slides in 70% glycerol/PBS to check for the presence

of an ectopic second sex comb and other abnormalities. While this does not give an

exact measurement of the frequency of the scd^1 allele because of this allele's

incomplete penetrance (though penetrance is almost complete in the FVW background;

see below), it should provide a reasonable proxy. Even though reductions in the

frequency of the mutant phenotype could also be driven by selection for suppressor

alleles, this should still give an indication of whether or not the scd^1 phenotype is

deleterious.

Experimental evolution—

To test whether sexual selection influences the rate of compensatory adaptation,

we set up two treatments. In the low sexual selection (LSS) treatment, we removed

sexual selection by enforcing monogamous mating. Each generation, we set up 100

vials, each containing one male and one virgin female. After a three-day interaction

period, we anesthetized the flies using CO_2, discarded males, and placed the females in

a population cage with four bottles containing culture media for egg laying. After four

days, the bottles were removed and incubated at 24˚C. When adult flies began eclosing,

we selected virgins for the next generation. Thus, while this treatment did preclude mate

choice, it still allowed for fecundity and viability selection (Arbuthnott and Rundle 2012).

In the high sexual selection (HSS) treatment, we followed a similar protocol

except allowed the opportunity for sexual selection. Each generation, 100 males and

100 virgin females were allowed to interact in a population cage, along with an open

culture bottle for food and moisture. After the three-day interaction period, we placed the

cage in a refrigerator to knock the flies out, and then we sorted males and females.

Males were discarded, and females were placed in fresh cages with four culture bottles

for a four-day egg-laying period. After egg laying, females were discarded, and the

bottles were placed in an environmental chamber at 24°C until adults began emerging,

at which point we collected virgins for the next generation.

Additionally, we also set up a treatment with low levels of genetic variation (LV)

to test whether compensatory adaptation is limited when segregating genetic variation is

diminished; in other words, testing whether the mutational target size of the

compensatory response was large enough that *de novo* mutations could contribute in

the time frame of the experimental evolution regime. In this treatment, each population

was established from the offspring of a single-pair mating between a randomly chosen

virgin *scd*** female and a randomly chosen *scd*** male. These populations were kept

under the same regime as the HSS treatment.

Finally, we set up a wild-type control (WTC) treatment using wild-type FVW flies.

WTC populations were also maintained under the same regime as the HSS treatment.

These provide a control for lab domestication and unknown aspects of the experimental

protocol.

All populations were initiated using randomly selected *scd*** flies (see above),

except for the LV treatments as described. We set up three replicate populations of

each treatment except for WTC, in which we performed two replicates. Experimental

evolution was conducted for a total of 24 generations. We assayed male sex comb

phenotypes as described earlier at generations 1, 7, 13, 19, and 24, using 30 randomly

selected males from each population at each time point.

To test how male sex comb traits changed over the course of experimental

evolution, we fit generalized linear mixed models using the glmmTMB (Hadfield 2010)

and lme4 (Bates *et al.* 2015) packages in R. For sex comb tooth number for both

primary and ectopic/secondary combs, we assumed a Poisson distribution and used a

log link function. The model included generation, treatment, and their interaction as

fixed effects; we also included individual fly, and in some cases replicate population

nested within treatment, as random effects (some models failed to converge when

replicate nested within treatment was included as a random effect). Additionally we also

tested for lineage specific zero-inflation in the data, but found no evidence for this, so

excluded this to reduce the number of fit parameters. To test whether the frequency of

defects in the primary sex comb changed over time, we fit a mixed logistic model (sex

comb defects present/absent), again with generation, treatment, and their interaction as

fixed effects, and replicate nested within treatment, as well as individual fly, as random

effects.

Results

Mapping scd^1—

Though we could not map *scd^1* to a specific gene, we were able to further narrow

down its location to an ~85-kb (cytological region 8F8-9A1) region on the X

chromosome through duplication mapping (Table 1, Figure 2). This region contains only

two complete annotated protein-coding genes, *btd* and *Sp1* (both of which influence

aspects of leg development and morphogenesis), and three annotated long non-coding

RNAs, CR42657, CR44016, and CR53498. Interestingly, the two parent strains carrying

the duplications that rescued scd^1 had smaller than average sex combs in the absence

of the scd^1 mutation (Table 1), similar to a past study involving this mutation (Randsholt

and Santamaria 2008), suggesting that the scd gene product is a suppressor of sex

comb development.

Influence of genetic background on penetrance and expressivity—

When females of the original scd^1 stock strain were crossed to males of various

wild-type strains to generate males that were hemizygous for scd^1 and heterozygous for

different genetic backgrounds, the penetrance and expressivity of scd^1 varied widely

(Figure 3) demonstrating segregating variation for them. The progenitor scd^1 strain from

the Bloomington stock center had the lowest penetrance (frequency of flies exhibiting an

ectopic sex comb on the second tarsal segment) and among the lowest expressivity

(number of teeth in the ectopic sex comb). Some of the other genetic backgrounds,

even in a heterozygous state, resulted in nearly complete penetrance for scd^1, including

the outbred population (FVW) for experimental evolution (described below).

Interestingly, the FVW outbred population only showed intermediate levels of

expressivity of the mutant phenotype, consistent with segregating variation in this

population. Overall this result suggests genetic background has a strong impact on the

phenotypic expression on scd^1. It also suggests that the partial penetrance initially

observed in the stock may reflect the accumulation of suppressor/compensatory

mutations in the base stock center strain itself.

Fitness effects of scd^1—

The frequency of male flies exhibiting the scd^1 phenotype decreased, on

average, across five generations of experimental evolution in populations polymorphic

for scd^1 (Figure 4). With a starting allele frequency of 0.7 the frequency decreased to an

average frequency of 0.4 (across the multiple replicates) in males. These results are

consistent with the mutation having moderate deleterious effects.

Experimental evolution—

In the populations carrying the scd^1 allele, defects such as gaps in the sex comb

were observed occasionally, and at significantly higher frequencies in the High Sexual

Selection (HSS) populations, and marginally significant frequencies in the Low Sexual

Selection (LSS) populations, than in the wild-type populations. The frequency of these

gaps appeared to decrease in the HSS populations across the 24 generations of the

experiment, although the interaction between generation and treatment was not

significant (Figure 5B; Table 2). In addition, the ectopic sex combs induced by the scd^1

mutation became smaller on average in both the HSS and LSS treatments (Figure 5C;

Table 3), losing on average ~0.5 teeth across the 24 generations of experimental

evolution. Again, we saw no evidence for differences in rate of compensation between

the EE treatments with and without sexual selection. For the Low genetic Variation (LV)

treatment, no significant change over time was observed, as expected (generation

effect = 0.0017, SD = 0.0044, p = 0.70). No significant changes were observed in the

primary sex comb tooth number across 24 generations in any of the experimental

treatments (Figure 5A; Table 4).

Discussion

Some, but not all, previous work has found that sexual selection may facilitate

populations in purging deleterious mutations (Radwan 2004; Hollis *et al.* 2009;

Jarzebowska and Radwan 2010) or accelerating rate of adaptation (Jacomb *et al.* 2016;

Parrett and Knell 2018). Few studies, however, have addressed whether sexual

selection may facilitate compensatory adaptation, where populations evolve traits to

compensate for the fitness costs of deleterious mutations. While compensatory

adaptation itself is well documented in other systems (Reynolds; Maisnier-Patin *et al.*

2002; Estes *et al.* 2011; Chandler *et al.* 2012; Comas *et al.* 2012; Chari *et al.* 2017), in

this experiment, we found no effect of the sexual selection regime on the rate of

compensatory adaptation (with respect to the sex comb) to a mutation that causes

morphological abnormalities in male sex combs in experimental *Drosophila* populations.

This is perhaps surprising, because our experiments show that the scd^1 mutation is

deleterious (Figure 4), and given the importance of the *Drosophila* sex comb for male

mating success (Ng and Kopp 2008), we expected that the fitness costs of this mutation

would involve male sexual fitness. Thus, we predicted that the costs of this mutation

would be higher in the HSS treatment, in which there was a high opportunity for female

mate choice, than in the LSS treatment, with reduced opportunity for sexual selection. It

is possible that this mutation has effects on other aspects of fitness in males or females

(viability, fecundity), but unfortunately our experiments did not directly measure specific

fitness components. Even so, theoretical work predicts that sexual selection should act

in concert with natural selection because of condition dependence (Whitlock and

Agrawal 2009); that is, mutations that reduce nonsexual fitness should also reduce

mating success, since sexual displays are often indicators of overall condition. While

some work has supported this prediction, our findings add to a growing body of work

suggesting that this is not always the case (Hollis and Houle 2011; Plesnar *et al.* 2011;

Arbuthnott and Rundle 2012, 2014; Cabral and Holland 2014; Power and Holman 2015;

Chenoweth *et al.* 2015). It is possible that the relatively minor degree of compensation

observed here made it difficult to detect differences between treatments.

Even though sexual selection did not impact the rate of compensatory evolution,

we did observe evidence of weak compensatory adaptation for the scd^1 mutation in both

the HSS and LSS treatments with respect to phenotypic suppression of the mutant

phenotypic effects. On average, the ectopic sex combs lost about half a tooth (starting

with a mean of ~ 3.5 teeth) over the course of 24 generations in these populations; in

other words, the expressivity of the mutation declined slightly. One possible explanation

for the similar response in both the high and low sexual selection treatments is simply

that the compensatory response (in terms of phenotypic suppression) was sufficiently

weak that any subtle difference between these treatments would be difficult to detect

given our design.

As expected, we did not observe any significant trend in the LV treatment, in

which populations experienced genetic bottlenecks prior to beginning the experiment

(LV populations were treated the same way as HSS populations). Combined with the

observation that genetic background has strong influences on the penetrance and

expressivity of this mutation (Figure 3), this suggests that compensatory adaptation

relies heavily on the presence of standing genetic variation, rather than new mutations.

An interesting side note is that the initial strain (obtained from the Drosophila stock

center) carrying scd^1 has among the lowest penetrance/expressivity for this mutation of

all the genetic backgrounds that we tested. This may suggest that the stock strain has

already undergone compensatory adaptation, and that alleles suppressing the

phenotypic expression of scd^1 had become fixed throughout the maintenance of this

stock (which were subsequently removed when we outcrossed the mutation), though

unfortunately we do not have any data on how long the scd^1 stock strain has been

maintained.

While we were unable to map scd^1 to a specific gene, we were able to localize it

to an ~85 kb region containing only a few candidates. The only protein-coding candidate

genes in this region, *Sp1* and *btd*, both have known roles in leg development (Estella

and Mann 2010), but are not specifically known to influence sex comb development.

There are also three long non-coding RNAs in this region (CR42657, CR44016, and

CR43498). Interestingly, all three of these RNAs show evidence of male-specific

expression in modENCODE RNA-seq data available on FlyBase (Graveley *et al.* 2011;

Brown *et al.* 2014). However, CR44016 shows expression only at very low levels and

only in adult males, not pupae or larvae, suggesting it is unlikely to be involved in the

development of sex combs. CR42657 and CR43498 both show expression in pupae

and/or larvae, as well as adult males (but not adult females); however, these RNAs

seem to be expressed in the testis and accessory gland and not other tissues (though

expression in legs specifically was not assessed in the modENCODE dataset). This

suggests that these male-specific reproductive tissues may be driving these sex-specific

expression patterns, not a role in sex comb development. Further work is necessary to

identify the molecular nature of scd^1.

There are a number of important limitations to point out about our study. First,

much of the focus was on compensation by suppression of the phenotypic effects of the

scd^1 mutation on the sex combs directly. While we observed similar levels of phenotypic

compensation with both our high and low sexual selection treatments (LSS and HSS), it

is possible that compensatory evolution differed with respect to the fitness components

(viability, fecundity, and sexual/mating components), which were not evaluated. Thus

we limit our interpretation to the effects on morphological compensation/suppression,

recognizing that we cannot rule out differential patterns of compensatory response for

fitness *per se*. Indeed, this pattern has been observed previously (Pischedda and

Chippindale 2005; Chari *et al.* 2017). Additionally, this experiment was performed over a

relatively short time period (25 generations); if we continued the experiment over a

longer period, subtle differences in the rate of morphological compensation may have

become apparent.

In summary, we found evidence of moderate compensatory adaptation to a

deleterious mutation. However, while compensatory adaptation did depend on the

presence of standing genetic variation, it did not depend on the opportunity for sexual

selection, in spite of the affected phenotype's known role in mating. This adds to a

growing body of studies suggesting that sexual selection does not always enhance

natural selection. Future work should tease apart when and why sexual and natural

selection act in concert and when they are likely to operate differently (Martínez-Ruiz

and Knell 2017).

Acknowledgments

We thank Shelby Hemker and Cody Porter for assistance in data collection and fly maintenance. The BEACON Center for the Study of Evolution in Action (NSF DBI-0939454) provided support for this work. ID was supported by funds from NSERC (Canada) Discovery and Discovery Accelerator awards.

Figure Legends

Figure 1. (A) Wild-type *Drosophila* leg showing a normal male sex comb (black arrow).

(B) scd^1 leg showing the primary sex comb (black arrow) as well as a smaller ectopic

sex comb on the second tarsal segment (white arrow).

Figure 2. Duplication mapping of scd^1. Purple bars represent duplications that rescued

the scd^1 mutant phenotype; pink bars represent duplications that failed to rescue scd^1.

(A) Entire region of the X chromosome in which duplications were tested. (B) Close-up

of the putative scd^1-containing region (red box). Because the duplication carried by

strain 30333 is sufficient to rescue the scd^1 phenotype, we hypothesize that the *scd*

gene must lie entirely within this region; at the same time, the neighboring duplications

(30522 and 30334) did not rescue the scd^1 phenotype, so *scd* must lie at least partially

outside of these regions.

Figure 3. (A) Penetrance and (B) expressivity of the scd^1 mutation varies among

different wild type genetic backgrounds. In these experiments, stock scd^1 females were

crossed to males of different wild-type strains, and phenotypes were scored in F1 males

(which were hemizygous for scd^1 and heterozygous for a different wild genetic

background). Penetrance was measured as the proportion of males showing an ectopic

sex comb on the second tarsal segment of the prothoracic leg, while expressivity was

measured as the number of sex comb teeth on the ectopic sex comb. Error bars

indicate 95% confidence intervals.

Figure 4. The scd^1 mutation is deleterious. Average frequency of scd^1 over time, across

4 replicate populations, each initialized with 70% scd^1 males. Error bars indicate 95%

confidence intervals.

428   **Figure 5.** Plots of model fits for sex comb traits in experimental evolution populations.

429   (A) Number of teeth in the primary sex comb across 24 generations of experimental

430   evolution in all experimental treatments. (B) Proportion of male flies with defects in the

431   primary sex comb in all four treatments. (C) Number of teeth in ectopic/secondary sex

432   combs in males in the HSS and LSS treatments.

References

Agrawal, A. F., 2001 Sexual selection and the maintenance of sexual reproduction.

Nature 411: 692–695.

Almbro, M., and L. W. Simmons, 2014 Sexual Selection Can Remove an Experimentally

Induced Mutation Load. Evolution 68: 295–300.

Arbuthnott, D., and H. D. Rundle, 2014 Misalignment of natural and sexual selection

among divergently adapted Drosophila melanogaster populations. Animal

Behaviour 87: 45–51.

Arbuthnott, D., and H. D. Rundle, 2012 Sexual Selection Is Ineffectual or Inhibits the

Purging of Deleterious Mutations in Drosophila Melanogaster. Evolution 66:

2127–2137.

Atallah, J., N. H. Liu, P. Dennis, A. Hon, and E. W. Larsen, 2009 Developmental

constraints and convergent evolution in Drosophila sex comb formation.

Evolution & Development 11: 205–218.

Atallah, J., H. Watabe, and A. Kopp, 2012 Many ways to make a novel structure: a new

mode of sex comb development in Drosophilidae. Evolution & Development 14:

476–483.

Bates, D., M. Mächler, B. Bolker, and S. Walker, 2015 Fitting Linear Mixed-Effects

Models Using lme4. Journal of Statistical Software 67: 1–48.

Boube, M., C. Benassayag, L. Seroude, and D. L. Cribbs, 1997 Ras1-Mediated

Modulation of Drosophila Homeotic Function in Cell and Segment Identity.

Genetics 146: 619–628.

Brown, J. B., N. Boley, R. Eisman, G. E. May, M. H. Stoiber *et al.*, 2014 Diversity and

dynamics of the *Drosophila* transcriptome. Nature 512: 393–399.

Cabral, L. G., and B. Holland, 2014 Courtship Song Does Not Increase the Rate of

Adaptation to a Thermally Stressful Environment in a Drosophila melanogaster

Laboratory Population. PLOS ONE 9: e111148.

Chandler, C. H., G. E. Chadderdon, P. C. Phillips, I. Dworkin, and F. J. Janzen, 2012

Experimental Evolution of the Caenorhabditis Elegans Sex Determination

Pathway. Evolution 66: 82–93.

Chandler, C. H., C. Ofria, and I. Dworkin, 2013 Runaway Sexual Selection Leads to

Good Genes. Evolution 67: 110–119.

Chari, S., C. Marier, C. Porter, E. Northrop, A. Belinky *et al.*, 2017 Compensatory

evolution via cryptic genetic variation: Distinct trajectories to phenotypic and

fitness recovery. bioRxiv 200725.

Charlesworth, B., 1978 Model for evolution of Y chromosomes and dosage

compensation. Proceedings of the National Academy of Sciences 75: 5618–

5622.

Chenoweth, S. F., N. C. Appleton, S. L. Allen, and H. D. Rundle, 2015 Genomic

Evidence that Sexual Selection Impedes Adaptation to a Novel Environment.

Current Biology 25: 1860–1866.

Clark, S. C. A., N. P. Sharp, L. Rowe, and A. F. Agrawal, 2012 Relative Effectiveness of

Mating Success and Sperm Competition at Eliminating Deleterious Mutations in

Drosophila melanogaster (S. R. Proulx, Ed.). PLoS ONE 7: e37351.

Comas, I., S. Borrell, A. Roetzer, G. Rose, B. Malla *et al.*, 2012 Whole-genome

sequencing of rifampicin-resistant *Mycobacterium tuberculosis* strains identifies

compensatory mutations in RNA polymerase genes. Nature Genetics 44: 106–

110.

Cook, R. K., M. E. Deal, J. A. Deal, R. D. Garton, C. A. Brown *et al.*, 2010 A New

Resource for Characterizing X-Linked Genes in Drosophila melanogaster:

Systematic Coverage and Subdivision of the X Chromosome With Nested, Y-

Linked Duplications. Genetics 186: 1095–1109.

Davies, A. G., A. Y. Game, Z. Chen, T. J. Williams, S. Goodall *et al.*, 1996 Scalloped

wings Is the Lucilia cuprina Notch Homologue and a Candidate for the Modifier of

Fitness and Asymmetry of Diazinon Resistance. Genetics 143: 1321–1337.

Ellis, E. A., and T. H. Oakley, 2016 High Rates of Species Accumulation in Animals with

Bioluminescent Courtship Displays. Current Biology 26: 1916–1921.

Estella, C., and R. S. Mann, 2010 Non-Redundant Selector and Growth-Promoting

Functions of Two Sister Genes, buttonhead and Sp1, in Drosophila Leg

Development (D. L. Stern, Ed.). PLoS Genetics 6: e1001001.

Estes, S., P. C. Phillips, and D. R. Denver, 2011 Fitness Recovery and Compensatory

Evolution in Natural Mutant Lines of C. Elegans. Evolution 65: 2335–2344.

Graveley, B. R., A. N. Brooks, J. W. Carlson, M. O. Duff, J. M. Landolin *et al.*, 2011 The

developmental transcriptome of *Drosophila melanogaster*. Nature 471: 473–479.

Grieshop, K., J. Stångberg, I. Martinossi-Allibert, G. Arnqvist, and D. Berger, 2016

Strong sexual selection in males against a mutation load that reduces offspring

production in seed beetles. Journal of Evolutionary Biology 29: 1201–1210.

Hadfield, J. D., 2010 MCMC Methods for Multi-Response Generalized Linear Mixed

Models: The MCMCglmm R Package. Journal of Statistical Software 33: 1–22.

Hollis, B., J. L. Fierst, and D. Houle, 2009 Sexual Selection Accelerates the Elimination

of a Deleterious Mutant in Drosophila Melanogaster. Evolution 63: 324–333.

Hollis, B., and D. Houle, 2011 Populations with elevated mutation load do not benefit

from the operation of sexual selection. Journal of Evolutionary Biology 24: 1918–

1926.

Jacomb, F., J. Marsh, and L. Holman, 2016 Sexual selection expedites the evolution of

pesticide resistance. Evolution 70: 2746–2751.

Jarzebowska, M., and J. Radwan, 2010 Sexual Selection Counteracts Extinction of

Small Populations of the Bulb Mites. Evolution 64: 1283–1289.

Kopp, A., 2011 Drosophila sex combs as a model of evolutionary innovations. Evolution

& Development 13: 504–522.

Long, T. A. F., A. F. Agrawal, and L. Rowe, 2012 The Effect of Sexual Selection on

Offspring Fitness Depends on the Nature of Genetic Variation. Current Biology

22: 204–208.

Lumley, A. J., Ł. Michalczyk, J. J. N. Kitson, L. G. Spurgin, C. A. Morrison *et al.*, 2015

Sexual selection protects against extinction. Nature 522: 470–473.

Mackay, T. F. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles *et al.*, 2012 The

Drosophila melanogaster Genetic Reference Panel. Nature 482: 173–178.

Maisnier-Patin, S., O. G. Berg, L. Liljas, and D. I. Andersson, 2002 Compensatory

adaptation to the deleterious effect of antibiotic resistance in Salmonella

typhimurium. Molecular Microbiology 46: 355–366.

Malagón, J. N., A. Ahuja, G. Sivapatham, J. Hung, J. Lee *et al.*, 2014 Evolution of

Drosophila sex comb length illustrates the inextricable interplay between

selection and variation. Proceedings of the National Academy of Sciences 111:

E4103–E4109.

Martínez-Ruiz, C., and R. J. Knell, 2017 Sexual selection can both increase and

decrease extinction probability: reconciling demographic and evolutionary

factors. Journal of Animal Ecology 86: 117–127.

Masta, S. E., and W. P. Maddison, 2002 Sexual selection driving diversification in

jumping spiders. Proceedings of the National Academy of Sciences 99: 4442–

4447.

McGuigan, K., D. Petfield, and M. W. Blows, 2011 Reducing Mutation Load Through

Sexual Selection on Males. Evolution 65: 2816–2829.

McKenzie, J. A., and G. Clarke, 1988 Diazinon Resistance, Fluctuating Asymmetry and

Fitnessin the Australian Sheep Blowfly, Lucilia cuprina. Genetics 120: 213–220.

Ng, C. S., and A. Kopp, 2008 Sex Combs are Important for Male Mating Success in

Drosophila melanogaster. Behav Genet 38: 195.

Parrett, J. M., and R. J. Knell, 2018 The effect of sexual selection on adaptation and

extinction under increasing temperatures. Proceedings of the Royal Society B:

Biological Sciences 285: 20180303.

Pischedda, A., and A. Chippindale, 2005 Sex, mutation and fitness: asymmetric costs

and routes to recovery through compensatory evolution. Journal of Evolutionary

Biology 18: 1115–1122.

Plesnar, A., M. Konior, and J. Radwan, 2011 The role of sexual selection in purging the

genome of induced mutations in the bulb mite. Evolutionary Ecology Research

13: 209–216.

Power, D. J., and L. Holman, 2015 Assessing the alignment of sexual and natural

selection using radiomutagenized seed beetles. Journal of Evolutionary Biology

28: 1039–1048.

Radwan, J., 2004 Effectiveness of sexual selection in removing mutations induced with

ionizing radiation. Ecology Letters 7: 1149–1154.

Randsholt, N. B., and P. Santamaria, 2008 How Drosophila change their combs: the

Hox gene Sex combs reduced and sex comb variation among Sophophora

species. Evolution & Development 10: 121–133.

Reynolds, M. G. Compensatory Evolution in Rifampin-Resistant Escherichia coli. 11.

Rowe, L., and D. Houle, 1996 The lek paradox and the capture of genetic variance by

condition dependent traits. Proceedings of the Royal Society of London. Series

B: Biological Sciences 263: 1415–1421.

Santamaria, P., and N. B. Randsholt, 1995 Characterization of a region of the X

chromosome of Drosophila including multi sex combs (mxc), a Polycomb group

gene which also functions as a tumour suppressor. Molec. Gen. Genet. 246:

282–290.

Sharp, N. P., and A. F. Agrawal, 2008 Mating Density and the Strength of Sexual

Selection Against Deleterious Alleles in Drosophila Melanogaster. Evolution 62:

857–867.

Venken, K. J. T., E. Popodi, S. L. Holtzman, K. L. Schulze, S. Park *et al.*, 2010 A

Molecularly Defined Duplication Set for the X Chromosome of Drosophila

melanogaster. Genetics 186: 1111–1125.

Whitlock, M. C., and A. F. Agrawal, 2009 Purging the Genome with Sexual Selection:

Reducing Mutation Load Through Selection on Males. Evolution 63: 569–582.