Molecular estimation of neurodegeneration pseudotime in older brains

Sumit Mukherjee[1,2], Christoph Preuss[3], Suman Jayadev[4], Gwenn Garden[4], Anna K Greenwood[1], Solveig K Sieberts[1], Phillip L De Jager[5,6], Nilufer Eretkin-Taner[7,8], Gregory W Carter[3], Lara M Mangravite[1], Benjamin A Logsdon[1,*]

[1]Sage Bionetworks, Seattle, WA, USA.
[2]Microsoft, Redmond, WA, USA[†].
[3]The Jackson Laboratory, Bar Harbor, ME, USA.
[4]Department of Neurology, University of Washington, Seattle, WA, USA.
[5]Center for Translational & Computational Neuroimmunology, Department of Neurology, Columbia University Irving Medical Center, New York City, USA
[6]Taub Institute, Columbia University Irving Medical Center, New York City, USA
[7]Mayo Clinic Florida, Department of Neurology, Jacksonville, FL, USA.
[8]Mayo Clinic Florida, Department of Neuroscience, Jacksonville, FL, USA.

[†]Current Address.

*Correspondence:
Benjamin A Logsdon, PhD
Sage Bionetworks
2901 3[rd] Avenue Suite 330
Seattle, WA 98121
ben.logsdon@sagebionetworks.org

Abstract
Therapeutic treatments for late-onset Alzheimer's disease (LOAD) are hindered by an incomplete

understanding of the temporal molecular changes that lead to disease onset and progression. Here, we

evaluate the ability of manifold learning to develop a molecular model for the unobserved temporal

disease progression from RNA-Seq data collected from human postmortem brain samples collected

within the ROS/MAP and Mayo Clinic RNA-Seq studies of the AMP-AD consortium. This approach

defines a cross-sectional ordering across samples based on their relative similarity in RNA-Seq profiles

and uses this information to define an estimate of molecular disease stage – or disease progression

pseudotime - for each sample. This transcriptional estimate of disease progression is strongly concordant

with burden of tau pathology (Braak score, $P = 1.0 \times 10^{-5}$), amyloid pathology (CERAD score, $P = 1.8 \times 10^{-5}$

$^{-5}$), and cognitive diagnosis ($P = 3.5 \times 10^{-7}$) of LOAD. Further, the disease progression estimate

recapitulates known changes in cell type abundance and impact of genes that harbor known AD risk loci.

Samples estimated to reside early in disease progression were enriched for control and early stage AD

cases, and demonstrated changes in basic cellular functions. Samples estimated to reside late in disease

progression were enriched for late-stage AD cases, and demonstrated changes in known disease processes

including neuroinflammation and amyloid pathology. We also identified a set of control samples with

late-stage estimated disease progression who also showed compensatory changes in genes involved in

affected pathways are protein trafficking, splicing, regulation of apoptosis, and prevention of amyloid

cleavage. In summary, we present a disease specific method for ordering patients based on their LOAD

disease progression from CNS transcriptomic data.

Introduction

Late-onset Alzheimer's disease (LOAD) is a devastating illness with no effective disease modifying

therapy, owing to a 99.6% failure of clinical trials[1]. There is a growing consensus that the most effective

treatments will intervene early in disease progression and halt disease pathophysiological processes prior

to conversion to LOAD[2]. In addition, there is increasing recognition that LOAD may in fact be a

spectrum of related diseases that have similar clinical and neuropathological manifestations[3,4]. Devising

successful therapeutic strategies will likely require targeting potentially diverse early-stage disease

processes that occur prior to a high burden of neuropathology or cognitive impairment.

Current approaches to identify AD affected individuals include *in vivo* measures of the pathological

hallmarks of disease – amyloid, tau, and neurodegeneration – via CSF biomarkers for amyloid and tau[5],

positron emission tomography for amyloid and tau (PET)[6], and structural and functional MRI of

neurodegeneration. Cognitive assessments are used to estimate disease burden[7], although measurable

cognitive impairment generally indicates a sustained burden of neuropathology and advanced

neurodegeneration. Based on biomarker studies of AD, by the time cognitive decline becomes detectable,

neuropathological changes of AD have already occurred, first in Aß and subsequently in tau related

measures[8] and therefore cannot be used to select patients for early disease stage studies. Furthermore,

while these measures of disease progression capture the overall increase in burden of pathology and

cognitive decline, they do not necessarily identify the dysfunctional molecular mechanisms that lead to

neuropathology and cognitive decline. There are likely many independent patient specific molecular

pathways present at an early stage in disease that then contribute to later stage disease progression[9,10].

This motivates the need to identify these early stage molecular mechanisms driving disease progression.

The Accelerating Medicines Project for Alzheimer's Disease (AMP-AD) consortia have generated

genome-wide transcriptomics of post-mortem brain tissue from patients across a broad range of

Alzheimer's disease neuropathological progression – including individuals with various stages of AD

neuropathology and those who lack AD neuropathology, but who may in face harbor early stage disease

molecular processes. We therefore sought to chart the molecular progression of the disease as reflected in

the aggregate behavior of the CNS transcriptome across these individuals. While standard approaches

such as differential expression or coexpression analyses have proven informative[11–15], these analyses do

not infer the relative stage of disease progression or identify distinct disease subtypes. Here we propose

an approach to analyze population level RNA-seq data from post-mortem brain tissue to learn a tree

structured progression (**Figure 1**) that represents distinct sub-types of disease and the relative progression

of disease across patients. With this approach, we identify potentially generalizable trajectories of LOAD

across heterogeneous patient populations at all stages of disease. Furthermore, we characterize molecular

pathways that define disease stages – a potential source of new biomarkers and therapeutic interventions

for early-stage disease processes along multiple different disease trajectories.

To learn the molecular disease staging and neuropathologic progression tree we use a manifold learning

method[16]. Manifold learning refers to a group of algorithms that aim to recover the low dimensional

subspace underlying a high dimensional dataset. Previous authors use manifold learning to estimate

disease progression from neuroimaging data[17] and to study lineage commitment of cells during

differentiation from single cell RNA-seq (scRNA-seq)[18–21]. To our knowledge, manifold learning has not

been used to estimate disease progression and/or disease stages from bulk RNA-seq data derived from

post-mortem tissue. Henceforth, we refer to manifold learning, lineage inference interchangeably in

reference to the construction of the inference of a neuropathologic progression tree. We demonstrate that

these tools can estimate the disease staging and progression tree (**Figure 2**) from bulk RNA-Seq data

collected from post-mortem brain tissues in a case/control cohort. Moreover, these trees show clear

LOAD staging, enable the study of cell type specific effects of LOAD, and allow the identification of

genetic factors driving disease progression.

Results

Unsupervised manifold learning distinguishes pathologically defined LOAD from control

We first quantify the bulk RNA-Seq data from the ROS/MAP and Mayo Clinic cohorts into gene counts

and remove any batch effects introduced due to sequencing runs using standard count normalization (see

Methods). The data from the ROS/MAP cohort is sampled from the dorsolateral prefrontal cortex

(DLPFC), and the data from the Mayo Clinic cohort is sampled from the temporal cortex (TCX). The full

pipeline we use for RNA-Seq data generation and quality control was recently reported[22]. The entire

transcriptome comprises many genes which do not have measurable expression or vary across

case/control samples, which we remove in order to reduce the noise in manifold learning[19]. To do this, we

first perform differential expression analysis between case/control samples separately for each study and

retain genes that reach an FDR of 0.10. To test if this biased the disease lineage inference, we also

perform manifold learning using only genes with high variance across samples, and we see a strong

concordance with disease lineages inferred with differentially expressed genes (**Figure S4**).   We infer the

disease lineage for each brain region on this subset of retained genes (**Figure 2A-B**). Furthermore, we

observe strong evidence of sex heterogeneity when performing the manifold learning approach, and find

that the manifolds inferred for female only samples are much more robust than for male samples. This

matches previous observations concerning disease specific sex heterogeneity[22]. We only show results for

manifolds inferred on female samples.

We first visualize the clinical diagnosis of the samples on the inferred disease staging tree to verify that

there is indeed separation of AD patients across the tree. To determine if inferred tree structure is an

accurate model of disease progression, we introduce the notion of disease pseudotime which is the

geodesic distance along the tree from an inferred initial point to the point of interest as a quantitative

linear measure of LOAD stage. We scale this estimated disease pseudotime to lie in the range $[0,1]$ to

make the effects comparable between the two studies (and brain regions). We show that for LOAD cases

compared to controls there is a significant association ($P = 0.02$ in Mayo and $P = 2.0 \times 10^{-6}$ in ROS/MAP,

logistic regression) between the estimated pseudotime and AD case/control status (**Figure 2C**).

We test whether genes in loci that have been implicated in genome wide association studies of LOAD are

associated with inferred disease pseudotime. We use the highest ranked LOAD GWAS genes (60 genes

in total)[23], **Table S1**, and compute the correlation between their expression and inferred pseudotime

(**Figure 2D**). When compared to the background of all genes, we see that there is a significant increase in

positive correlation with disease pseudotime for implicated LOAD GWAS genes (P-value: $7.3x10^{-5}$ in

Mayo and $5.6x10^{-3}$ in ROS/MAP). Furthermore, this does not appear to be driven by a small subset of

outlier genes, but by the majority of the distribution of LOAD GWAS genes. The fact that AD GWAS

loci genes have expression associations with pseudotime likely implies that the AD risk variants at these

are also eQTL as previously shown[24–27] and/or are members of co-expression networks that are

differentially expressed in AD[13,28].

To further explore the relationship between inferred disease stage and LOAD, we test for its association

with neuropathological and clinical measures of LOAD severity, namely: i) Braak score, ii) CERAD

score, and iii) cognitive diagnosis. The ROSMAP study has numeric scores for these categories available

as covariates for each sample. Braak is a semi-quantitative measure that increases with tau pathology [29]

and CERAD is a semi-quantitative measure of density of neuritic plaques[30]. We overlay these scores on

the inferred manifold for the DLPFC brain region (**Figure 3A**). We observe a progressive increase in tau,

amyloid, and cognitive burden as we traverse the inferred disease manifold (**Figure 3A**). This is further

quantified by characterizing the relationship between branches of the inferred manifold and Braak,

CERAD, and cognitive diagnosis (**Figure 3B**). We observe significant associations between pseudotime

and Braak score (P=$1.0x10^{-5}$), CERAD score (P=$1.8x10^{-5}$), and cognitive diagnosis (P=$3.5x10^{-7}$).

Inferred staging recapitulates known biology of AD

To demonstrate that the inferred disease pseudotime recapitulates known biology of LOAD, we test for

association between inferred disease stage and both the cellular response to disease and the genetics of the

disease. A prominent hypothesis in AD is that the effects of the disease vary across different brain cell

types, specifically neurons and glial subtypes. Current understanding of the cell biology of the disease

implicates progressive neuronal loss and increase in gliosis[31]. To test if the inferred pseudotime aligns

with existing cell type specific hypotheses regarding AD, we first selected from the genes used in lineage

construction the marker genes for four key cell types: neurons, astrocytes, microglia, and

oligodendrocytes based on a previously published brain cell atlas[32] (**Table S2**). We then calculate the

normalized mean expression for the marker genes of each cell type and fitted a linear model to the mean

expression with disease pseudotime as the dependent variable. We find that, in both studies, the cell

specific marker gene levels show a statistically significant linear dependence on pseudotime (**Table S3**).

Fitted effects recapitulate known neuropathologic changes which occur in AD, namely: i) a reduction in

the neuronal populations as AD progresses, and ii) an increase in expression associated with activation of

microglia, astrocytes, and oligodendrocytes as AD progresses (**Figure S5**).

Next, we test for association between assigned lineage state in ROS/MAP (DLPFC) and Mayo (TCX) and

APOE e4 status (**Figure S6**). For reference, the inferred trees for TCX and DLPFC each resolve into 6

branches (**Figure 4A,S7**). Carriers of the APOE e4 allele are significantly enriched on the State 4 branch

in TCX (P-value = 0.027, unadjusted), and suggestively enriched on the State 5 branch (P-value = 0.06,

unadjusted), compared to the State 1 branch (logistic regression).

Genetic factors associated with inferred disease staging

Lineage inference of LOAD transcriptomes provides a quantitative measure of disease progression for

genetic associate testing, and the significantly greater correlation between pseudotime and gene

expression for known LOAD risk genes (**Figure 2D**) suggests that the observed differences in disease

trajectories are influenced by genetic factors. To test this hypothesis, we perform single variant analysis

using whole-genome sequencing data for 305 patients from the ROS/MAP and 131 patients from the

Mayo cohort. Despite the limited sample size, resulting in lack of statistical power to discover genome-

wide significant associations, multiple variants reach a genome-wide suggestive threshold of $p < 1 \times 10^{-5}$

(**Table S4**).   We do not see evidence of population stratification in the analysis (**Figure S8-S9**). Notably,

the most significant association with pseudotime for the ROS/MAP cohort is observed at the *PTPRD*

locus (rs7870388, $p = 1.31 \times 10^{-6}$) (**Figure S10, Table S4**). The *PTRPD* locus is associated with the

susceptibility to neurofibrillary tangle independent of amyloid deposition in the ROS/MAP cohort[33]. For

the Mayo Clinic cohort, known LOAD variants in the *APOE* (rs6857, p = 9.18 x 10^{-6}) and *BIN1*

(rs62158731, p= 4.68 x 10^{-5}) loci overlap with variants associated with inferred disease stage (**Figure**

S10, Table S4)[34]. When comparing our association results for inferred disease stage with summary

statistics from a large-scale case-control approach, we identify multiple variants which have been

previously associated with LOAD in the IGAP cohort (**Table S5**). Furthermore, we identify several

potential novel candidate genes associated with inferred disease stage (*ADAMTS14*, *IL7*, *MAN2B1*) linked

to immune and lysosomal storage function (**Figure S10**,**Table S4**). *IL-7* has been proposed as an

inflammatory biomarker for LOAD that correlates with disease outcome and severity[35]. *ADAMTS14* is

part of a locus that has been previously linked with Alzheimer susceptibility and plays an important role

in the regulation of immune function via TGF-beta signaling.

New disease insights identified from inferred disease lineages

Another important direction of study in the field of Alzheimer's is the identification of disease subtypes,

which has so far predominantly been done using imaging data[36]. The branches of the inferred disease trees

provide a new transcriptomic-based approach to identify disease subtypes. In both brain regions and in

two separate cohorts, there were two distinct early-lineage branches corresponding to predominantly

control samples, which we interpret as different initial paths towards the disease. Similarly, both brain

regions feature several distinct branches with predominantly LOAD samples (**Figure 2A-B**).

Branch-specific differential expression patterns. To study the genes and pathways specific to each branch,

we perform a branch-specific differential expression analysis with an ANOVA model using the branches

with the highest proportion of controls as the reference branch for DLPFC (**Table S6**) and TCX (**Table**

S7). We see many genes are differentially expressed between the control branch and branches that are

enriched in the affected individuals (**Table S8**). Next, we performed an enrichment analysis on each of

these differentially expressed gene sets with the enrichR[37] package for Gene Ontology[38] annotations

(**Methods**). The results of this enrichment analysis for DLPFC and TCX tissues (**Table S9-S10**). Only

gene sets with significant enrichment are shown (FDR adjusted p-value < 0.05). Overall, we see a pattern

of loss of expression of basic cell biology mechanisms in early-stage branches including RNA splicing,

mitochondria function, protein transport, and DNA repair. Late-stage branches were characterized by

increased immune response (e.g. TGFb/WNT signaling) and apoptotic activity (**Table 1**).

While studying the different branches in the two brain regions, we observe a branch (branch 5) that

corresponds to a group of predominantly neuropathological control samples from the Mayo RNA-seq

cohort that were in close proximity to a branch with predominantly LOAD samples (branch 4) on the

inferred disease lineage (**Figure 4A**). However, most of the samples on branch 5 are neuropathological

controls as defined by the Mayo diagnostic criteria. We bi-cluster the mean expression of genes in each

branch and the branches themselves (**Methods**). This clustering analysis (**Figure 4B**) shows that the

closest branch to this potentially disease resistant branch contains the highest proportion of AD samples.

While the stage proximity implies some transcriptomic similarity between these controls and nearby

cases, we also see a secondary cluster of genes with increased expression in the resistant state while

having reduced expression in all other states. We perform an enrichment analysis on this set of genes and

find significant GO terms corresponding to: protein transport (GO:0015031), regulation of mRNA

splicing, via spliceosome (GO:0048024), negative regulation of apoptotic process (GO:0043066), and

regulation of amyloid-beta clearance (GO:1900221) (Cluster4, **Table S11**)**.** It is possible that these

potentially disease resistant individuals have compensatory mechanisms which suppress the hallmarks of

disease despite sharing gene expression patterns with pathologically affected individuals. These

observations are preliminary, and would need to be replicated in a second cohort to verify the existence of

a disease resistant gene expression signature.

Discussion

Here we proposed a novel approach to infer the Alzheimer's disease severity and disease subtypes in an

unsupervised manner from post-mortem bulk RNA-seq data that gets directly at the challenge of

identifying the temporal progression of disease in the disease resident tissue. Our strategy utilized a

manifold learning approach to infer a disease progression tree from cross-sectionally collected patient

samples from two different brain regions. The underlying assumption of our approach is that the inferred

disease progression from cross sectional samples serves as a proxy for the unobserved progression of the

disease across subtypes of LOAD. We validated this hypothesis through comparisons with

neuropathological measures of disease stage severity and against known cell type specific effects caused

by the disease. Furthermore, this approach provides clues to better understanding the molecular

heterogeneity of disease by identifying specific pathways that are dysregulated in subsets of patients at

different disease stages. This opens up the possibility of better patient stratification and precision

medicine.

We observed that different biological processes vary as a function of inferred disease stage, and that

early-stage disease processes potentially include RNA-splicing, mitochondrial function, and protein

transport – implicating multiple basic cell biology mechanisms as potential early stage disease processes

for further study in relevant model systems. Additionally, the manifold learning method identified 6

potential subtypes of LOAD from RNA-seq (i.e. branches) suggesting the LOAD populations should be

stratified by better biomarkers with tailored treatment strategies. To identify and test these stratifications

future studies should focus on longitudinal cohorts of patients with rich molecular and imaging data to be

able to identify biomarkers that can accurately and precisely stratify patients into the underlying

molecular subtypes in terms of the molecular characteristics of their transcriptome and different relative

stages of disease. Furthermore, we observe a potential disease resistant sub-type of patients. This disease

resistance should be tested in disease model systems, to identify if neuropathological readouts can be

modified by altering the function of the pathways identified in our analysis (e.g. APP processing, RNA

splicing, apoptosis, protein trafficking). While this preliminary observation needs to be validated in

another cohort, it has the potential to be a novel source of hypotheses concerning new therapeutic

development. Specifically, for constructing better combination therapy hypotheses that may confer

neuroprotection, even in patients that are mildly affected by disease.

LOAD is a complex and heterogeneous disease encompassing a broad spectrum of clinical symptoms.

Disease progression can vary widely between patients leading to different rates of cognitive decline.

Several lines of evidence suggest that these differences in progression are modified by multiple genetic

factors affecting the transition from one pathological state to another[39,40]. However, it has remained

difficult to assess the role of genetic variants affecting disease trajectories by case-control approaches

alone. Here, we showed that our novel expression trait pseudotime might be used as a molecular

phenotype to identify known and novel AD loci associated with different disease progression states across

AD patients. Despite a limited sample size, we identified previously associated AD candidate loci in the

ROSMAP (*PTPRD*) and Mayo (*BIN1, APOE*) cohorts with suggestive significance ($p < 1 \times 10^{-5}$).

Variants in *PTPRPD* have been associated with the susceptibility to neurofibrillary tangles, independent

of amyloid burden. This is in line with the results from the differential gene expression analysis of

pseudotime branches showing an enrichment of molecular pathways implicated in TAU pathology.

Furthermore, our analysis revealed several novel loci linked to immune function (*ADAMTS14*, *IL7*) and

neurotransmitter signaling (*CHRM2*, *CHRM3*) processes associated with disease pseudotime (**Table S4**).

Future studies will be needed to replicate these findings in independent cohorts of LOAD and validate the

role of candidate genes in LOAD related disease progression by first identifying peripheral biomarkers

that correspond to this molecular definition of disease stage, and then testing for GWAS association with

that disease stage. Subsequent results can improve functional interpretation by linking candidate genes

with ordered pathological processes.

Methods

RNA sequencing

The details of the sample collections, postmortem sample characteristics, the tissue and RNA

preparations, the library preparations and sequencing technology and parameters, and sample quality

control filters are provided in previously published work[41,42]. Furthermore, details of the bioinformatic

pipeline used to generate count level data has been previously described[22]. Briefly, reads were aligned to

the GENCODE24 (GRCh38) reference genome with STAR[43], and gene counts generated using the

HTSeq algorithm[44]. Genes that had more than one counts per million total reads total reads in at least

50% of samples in each tissue and diagnosis category were used for further analysis.

Differential Expression analysis on Mayo and ROS/MAP cohorts

For gene filtering we used false discovery rate of 0.05 from the previously published differential

expression analysis of Mayo and ROS/MAP RNA seq data[22]. Briefly, case control status was harmonized

across the Mayo and ROS/MAP cohorts, where controls were defined as individuals with a low burden of

amyloid and tau based on CERAD and Braak scores, and cases with a high burden. Furthermore in

ROS/MAP, clinical diagnosis was also used with controls having to have no cognitive impairment, and

cases have probably AD[22]. Differential expression analysis was run on suitably normalized data – using

conditional quantile normalization to account for variation in gene length and GC content, removing

sample outliers, covariate identification adjustment, with sampling abundance confidence estimated using

a weighted linear model with the voom-limma package[22,45,46]. A fixed/mixed effect linear model is used

to fit the differential expression model on the normalized data[22].

Manifold learning for LOAD

Manifold learning refers to a group of machine learning algorithms that recover a low dimensional

subspace underlying a high dimensional dataset. Manifold learning approaches are typically used in

datasets or applications where data samples lie on an underlying low dimensional latent space (e.g. a tree,

a line, a curved plane). The low dimensional space is learned via a projection from the high dimensional

space of the observed data (e.g. RNA-seq profiles across hundreds of patient samples) down to a low

dimensional space with suitable regularization constraints to enforce smoothness and the structural

constraints of the low dimensional space. (**Figure 1A**). Due to the necessary assumption of an underlying

latent subspace, manifold learning is commonly used in applications where it is known that the observed

data is obtained from a progression of some kind; e.g., i) to infer the temporal ordering of a sequence of

images, or ii) to infer the approximate lineage of cells in a differentiation trajectory using single cell

RNA-Seq data (**Figure 1B-C**).

Here, we repurpose methods originally developed for learning cell lineage using scRNA-Seq data, to infer

the staging of Alzheimer's disease (AD) using bulk RNA-Seq data from post-mortem brain samples with

known AD diagnosis status. Since bulk RNA-Seq has many of the same sampling and distributional

properties as scRNA-Seq, we observe that scRNA-Seq methods are applicable with no additional

modifications. As such, we use the DDRTree manifold learning approach available in the Monocle 2 R

package [19]. However, we also show that the estimated staging of disease is quite similar across some of

the other common methods used for scRNA-Seq lineage estimation (**Figures S1-S3**).

The RNA-Seq data used in this study was generated from post-mortem brain homogenate samples, and

obtained from two separate studies that are a part of the Accelerating Medicines Partnership in

Alzheimer's Disease (AMP-AD) consortium, namely: i) the Religious Orders Study and the Memory and

Aging Project (ROSMAP) [47,48], and ii) the Mayo RNA-seq study[49]. For this paper, we focused our

analysis on the temporal cortex (TCX) and dorsolateral prefrontal cortex (DLPFC) tissue samples. Within

the Mayo RNA-seq study the TCX samples are derived from individuals neuropathologically defined as

either aged controls, LOAD cases, Progressive Supranuclear Palsy (PSP) cases, or pathological aging

(PA) cases [49].   The ROSMAP study is a prospective longitudinal cohort of an aging population, and has

samples from participants with clinical and neuropathological diagnoses of LOAD[42], aged controls, and

individuals with mild cognitive impairment. Furthermore, the results presented in the main paper are

from female samples only, as we observed significant sex differences in the transcriptomic data consistent

with current knowledge of sex differences in LOAD[50,51], making a common analysis of both sexes

untenable.

Manifold learning using Discriminative Dimensionality Reduction Tree (DDRTree)

DDRTree is a manifold learning algorithm that infers a smooth low dimensional manifold by an approach

called reverse graph embedding. Briefly, the algorithm simultaneously learns a non-linear projection to a

latent space where the points lie on a spanning tree. A reverse embedding is also simultaneously learned

from the latent space to the high dimensional data. Mathematically, the DDRTree algorithm can be posed

as the following optimization problem:

$$\min_{W,Z,B,Y,R} \sum_{i=1}^{N} ||x_i - W z_i||^2 + \frac{\lambda}{2} \sum_{k,k'} b_{k,k'} ||W y_k - W y_{k'}||^2 + \gamma \left[\sum_{k=1}^{K} \sum_{i=1}^{N} r_{i,k} ||z_i - y_k||^2 + \sigma r_{i,k} \log(r_{i,k}) \right]$$

$$s.t. \quad B \; represents \; a \; spanning \; tree,$$

$$W^T W = I, r_{i,k} \geq 0, \sum_{k=1}^{K} r_{i,k} = 1$$

Here, $\{z_i\}_{i=1}^{N} \in \mathbb{R}^{genes}$ represents RNA-Seq data from each patient sample, $\{z_i\}_{i=1}^{N} \in \mathbb{R}^2$ represents the

latent representation of each sample as inferred by the algorithm, $\{y_k\}_{k=1}^{K}$ represents the centers of

clusters in the dataset, $W \in \mathbb{R}^{2 \times genes}$ represents an inverse mapping from the latent space to the high

dimensional space of RNA-Seq data, $B \in \mathbb{R}^{K \times K}$ represents a spanning tree on which the centers of the

clusters lie and $R \in \mathbb{R}^{N \times K}$ captures the soft clustering information of samples in the dataset. The first

term of the optimization problem is responsible for learning a low dimensional representation of the data

such that an inverse mapping exists to the high dimensional data points, the second term learns the tree

structure of the points and the third term learns a soft clustering for the latent dimension points as well as

the centers of the clusters. Despite the non-convexity of the problem, each individual optimization

variable can be solved for efficiently using alternative minimization as described previously[52]. This

algorithm was implemented using the Monocle package in R[19]. The code to infer the lineage in Mayo is

available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/TCX_GenerateMonocleDS_new.R, and code used

to infer the lineage in ROSMAP is available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/DLPFC_GenerateMonocleDS_new.R.

Branch assignment and pseudotime calculation for samples

Branch assignment and pseudotime calculation was also performed using the *Monocle* package using

techniques described previously[19]. Briefly, pseudotime is calculated by first identifying a root point on

one of the two ends of the maximum diameter path in the tree. Then the pseudotime of each point is

calculated by projecting it to its closest point on the spanning tree and calculating the geodesic distance to

the root point. Assigning samples to branches is done by first identifying the branches of the spanning

tree and then assigning samples to the branch on which their projection to the spanning tree lies on.

Association of pseudotime with AD status, hallmarks of Alzheimer's disease, and cognitive diagnosis

We test for association between disease pseudotime and AD case or control status with logistic regression

with AD case or control status as the outcome and inferred pseudotime as the dependent variable in both

the Mayo and ROS/MAP studies. We test for association between pseudotime and hallmarks of disease

in the ROS/MAP studies for both Braak (measure of tau pathology) score and CERAD score (measure of

amyloid pathology) with an ordinal logistic regression model, with the neuropath score as the ordered

outcome, and pseudotime as the dependent variable. Finally, we test for association between disease

pseudotime and cognitive diagnosis for the following ordered clinical diagnoses of no cognitive

impairment, mild cognitive impairment, and probable Alzheimer's disease with an ordinal logistic

regression model. All code for running these association tests is available https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/paper_figures.Rmd.

Inferring cell type specific expression patterns given marker gene expression as a function of pseudotime

List of marker genes for different major cell types in the brain was curated from a previously published

brain cell expression signature study[32]. The marker gene list was then pruned to include only genes that

were included in lineage construction. Each gene's expression as a function of pseudotime was then

obtained by smoothing using a smoothing spline of degree of freedom = 3 and normalized to lie in [0,1].

The smoothing was done to remove the effects of technical noise introduced due to RNA-Seq and the

normalization was done since the absolute expression levels of genes might be very different from each

other. The smoothed and normalized expression of marker genes for each category were then averaged to

obtain the average marker gene expression as a function of pseudotime.   A linear model was used to test

for association between average expression of a given cell type expression signature and pseudotime.

Association between GWAS loci and correlation with pseudotime

To test for association between pseudotime and LOAD GWAS genes, we computed the Spearman's

correlation between each gene's expression and pseudotime in the Mayo and ROS/MAP studies. Next,

we identified a set of genes implicated in AD GWAS loci in the International Genetics of Alzheimer's

Project (IGAP)[23]. We treated the set of genes described in Tables 1-3 of that study as high quality

candidate AD GWAS genes[23]. We test for a difference between the correlation with pseudotime of

background of all other genes and the IGAP AD genes using a linear model, and see a significant increase

in correlation between gene expression and pseudotime in both the Mayo and ROS/MAP study for AD

GWAS genes.

Branch specific differential expression analysis

We perform a state specific differential expression analysis using a one-way ANOVA model in both the

Mayo and ROS/MAP studies. The branch with the highest proportion of AD controls is defined as the

reference branch for all analyses. We use Tukey's honest significant difference method to compute P-

values for the test for change in expression of a given gene compared to the reference branch. Genes are

grouped based on their branch and direction of change in expression for further downstream pathway

enrichment analyses. Code to run analyses are available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/DLPFC_DE_Anova.R for ROS/MAP and here

https://github.com/Sage-Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/TCX_DE_Anova.R for

Mayo.

Estimating branch specific gene expression signatures

Branch specific expression signature was obtained by first calculating the average normalized expression

for all genes in each state/branch. This was followed by performing a bi-clustering using the pheatmap

package in R which uses hierarchical clustering on both samples and genes. We also used the pheatmap

package to visualize the state specific expression signatures.

Gene set enrichment analyses

For each branch specific differential expression gene set (DEGs) in both Mayo and ROS/MAP we

perform a gene set enrichment analysis against Gene Ontology pathways using the enrichR R package.

Only pathways with FDR < 0.05 are reported. The code we used to run the ROS/MAP DEG enrichments

are available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/lineage.Rmd, the code we used to run the Mayo

DEG enrichments are available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/lineageTCX.Rmd, and the code we used to run the

branch specific gene expression signature pathway enrichments is available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/resilience.Rmd.

Whole-genome sequencing

Whole-genome sequencing was performed at the New York Genome Center for all individuals from the

ROS/MAP and Mayo cohorts. Detailed information for both data sets can be accessed via synapse

(DOI:10.7303/syn2580853). Briefly, 650ng of genomic DNA from whole blood was sheared using a

Covaris LE220 sonicator. DNA fragments underwent bead-based size selection and were subsequently

end-repaired, adenylated, and ligated to Illumina sequencing adapters. Libraries were sequenced on an

Illumina HiSeq X sequencer using 2 x 150bp cycles. Paired-end reads were aligned to the GRCh37

(hg19) human reference genome using the Burrows-Wheeler Aligner (BWA-MEM v0.7.8) and processed

using the GATK best-practices workflow [53,54]. This included marking of duplicate reads by the use of

Picard tools v1.83, local realignment around indels, and base quality score recalibration (BQSR) via

Genome Analysis Toolkit (GATK v3.4.0). Joint variant calling files (vcfs) for whole-genome sequencing

data for the Mayo and ROS/MAP cohort were obtained through the AMP-AD knowledge portal

(www.synapse.org/#!Synapse:syn10901595).

Single variant association with pseudotime in two independent cohorts

Likelihood ratio tests within a linear regression framework were used to model the relationship between

the quantitative expression trait pseudotime and genetic variants in 436 AD cases. Genome-wide genetic

association analysis was performed for 305 female patients in the ROS/MAP cohort and 131 female

patients in the Mayo cohort for which both genotyping and post-mortem RNA-seq data was available. An

efficient mixed model approach, implemented in the EMMAX software suite, was used to account for

potential biases and cryptic relatedness among individuals [55]. Only variants with MAF > 0.05, genotyping

call rates > 95%, minimum sequencing depth of 20 reads and Hardy-Weinberg equilibrium $p > 10^{-4}$ were

considered for analysis. Quantile-quantile plots (**Figure S8-S9**) for the test statistics showed no

significant derivation between expected and observed p-values, highlighting that there is no consistent

differences across cases and controls except for the small number of significantly associated variants.

451 Furthermore, the genomic inflation factor (lambda) was determined to be 0.99 for the Mayo and 0.98 for

452 the ROS/MAP single variant association tests. This highlights that potential confounding factors, such as

453 population stratification have been adequately controlled.

References

1.   Cummings, J. L., Morstorf, T. & Zhong, K. Alzheimer's disease drug-development pipeline: few candidates, frequent failures. *Alzheimers. Res. Ther.* **6**, 37 (2014).

2.   Cummings, J. L., Doody, R. & Clark, C. Disease-modifying therapies for Alzheimer disease: Challenges to early intervention. *Neurology* **69**, 1622–1634 (2007).

3.   Ferreira, D. *et al.* Distinct subtypes of Alzheimer's disease based on patterns of brain atrophy: longitudinal trajectories and clinical applications. *Sci. Rep.* **7**, 46263 (2017).

4.   Bredesen, D. E. Metabolic profiling distinguishes three subtypes of Alzheimer's disease. *Aging (Albany NY)* **7**, 595 (2015).

5.   Brier, M. R. *et al.* Tau and Aβ imaging, CSF measures, and cognition in Alzheimer's disease. *Sci. Transl. Med.* **8**, 338ra66--338ra66 (2016).

6.   Gordon, B. A. *et al.* The relationship between cerebrospinal fluid markers of Alzheimer pathology and positron emission tomography tau imaging. *Brain* **139**, 2249–2260 (2016).

7.   Dichgans, M. *et al.* METACOHORTS for the study of vascular disease and its contribution to cognitive decline and neurodegeneration: An initiative of the Joint Programme for Neurodegenerative Disease Research. *Alzheimer's Dement.* **12**, 1235–1249 (2016).

8.   Jack, C. R. *et al.* Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. *Lancet Neurol.* **12**, 207–216 (2013).

9.   Au, R., Piers, R. J. & Lancashire, L. Back to the future: Alzheimer's disease heterogeneity revisited. *Alzheimer's Dement. (Amsterdam, Netherlands)* **1**, 368–370 (2015).

10.   Carrasquillo, M. M. *et al.* Late-onset Alzheimer's risk variants in memory decline, incident mild cognitive impairment, and Alzheimer's disease. *Neurobiol. Aging* **36**, 60–7 (2015).

11.   Zhang, B. *et al.* Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. *Cell* **153**, 707–20 (2013).

12.   Mostafavi, S. *et al.* A molecular network of the aging human brain provides insights into the

pathology and cognitive decline of Alzheimer's disease. *Nat. Neurosci.* **21**, 811–819 (2018).

13.   Conway, O. J. *et al.* ABI3 and PLCG2 missense variants as risk factors for neurodegenerative

diseases in Caucasians and African Americans. *Mol. Neurodegener.* **13**, 53 (2018).

14.   Allen, M. *et al.* Conserved brain myelination networks are altered in Alzheimer's and other

neurodegenerative diseases. *Alzheimers. Dement.* **14**, 352–366 (2018).

15.   Allen, M. *et al.* Divergent brain gene expression patterns associate with distinct cell-specific tau

neuropathology traits in progressive supranuclear palsy. *Acta Neuropathol.* **136**, 709–727 (2018).

16.   Bengio, Y., Courville, A. & Vincent, P. Representation learning: A review and new perspectives.

IEEE Trans. Pattern Anal. Mach. Intell. **35**, 1798–1828 (2013).

17.   Wolz, R., Aljabar, P., Hajnal, J. V. & Rueckert, D. Manifold Learning for Biomarker Discovery in

MR Imaging. in 116–123 (Springer, Berlin, Heidelberg, 2010). doi:10.1007/978-3-642-15948-

0_15

18.   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by

pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32**, (2014).

19.   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat. Methods*

14, 979–982 (2017).

20.   Rosenberg, A. B. *et al.* Single-cell profiling of the developing mouse brain and spinal cord with

split-pool barcoding. *Science (80-.).* **360**, 176–182 (2018).

21.   Mukherjee, S., Zhang, Y., Fan, J., Seelig, G. & Kannan, S. Scalable preprocessing for sparse

scRNA-seq data exploiting prior knowledge. *Bioinformatics* **34**, i124--i132 (2018).

22.   Logsdon, B. *et al.* Meta-analysis of the human brain transcriptome identifies heterogeneity across

human AD coexpression modules robust to sample collection and methodological approach.

bioRxiv 510420 (2019).

23.   Kunkle, B. W. *et al.* Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk

loci and implicates Aβ, tau, immunity and lipid processing. *Nat. Genet.* **51**, 414–430 (2019).

24.   Allen, M. *et al.* Novel late-onset Alzheimer disease loci variants associate with brain gene

506 expression. *Neurology* **79**, 221–8 (2012).

507 25. Allen, M. *et al.* Late-onset Alzheimer disease risk variants mark brain regulatory loci. *Neurol.*

508 *Genet.* **1**, e15 (2015).

509 26. Allen, M. *et al.* Association of MAPT haplotypes with Alzheimer's disease risk and MAPT brain

510 gene expression levels. *Alzheimers. Res. Ther.* **6**, 39 (2014).

511 27. Carrasquillo, M. M. *et al.* A candidate regulatory variant at the TREM gene cluster associates with

512 decreased Alzheimer's disease risk and increased TREML1 and TREM2 brain gene expression.

513 *Alzheimers. Dement.* **13**, 663–673 (2017).

514 28. Sims, R. *et al.* Rare coding variants in PLCG2, ABI3, and TREM2 implicate microglial-mediated

515 innate immunity in Alzheimer's disease. *Nat. Genet.* **49**, 1373–1384 (2017).

516 29. Braak, H., Alafuzoff, I., Arzberger, T., Kretzschmar, H. & Del Tredici, K. Staging of Alzheimer

517 disease-associated neurofibrillary pathology using paraffin sections and immunocytochemistry.

518 *Acta Neuropathol.* **112**, 389–404 (2006).

519 30. Mirra, S. S. *et al.* The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part

520 II. Standardization of the neuropathologic assessment of Alzheimer's disease. *Neurology* **41**, 479–

521 86 (1991).

522 31. De Strooper, B. & Karran, E. The Cellular Phase of Alzheimer's Disease. *Cell* **164**, 603–615

523 (2016).

524 32. Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and

525 vascular cells of the cerebral cortex. *J. Neurosci.* **34**, 11929–47 (2014).

526 33. Chibnik, L. B. *et al.* Susceptibility to neurofibrillary tangles: role of the PTPRD locus and limited

527 pleiotropy with other neuropathologies. *Mol. Psychiatry* **23**, 1521 (2018).

528 34. Lambert, J.-C. *et al.* Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for

529 Alzheimer's disease. *Nat. Genet.* **45**, 1452 (2013).

530 35. Janelidze, S. *et al.* CSF biomarkers of neuroinflammation and cerebrovascular dysfunction in early

531 Alzheimer disease. *Neurology* **91**, e867--e877 (2018).

36.   Whitwell, J. L. *et al.* Neuroimaging correlates of pathologically defined subtypes of Alzheimer's

disease: a case-control study. *Lancet Neurol.* **11**, 868–877 (2012).

37.   Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool.

BMC Bioinformatics **14**, 128 (2013).

38.   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology

Consortium. *Nat. Genet.* **25**, 25–9 (2000).

39.   Dong, A. *et al.* Heterogeneity of neuroanatomical patterns in prodromal Alzheimer's disease: links

to cognition, progression and biomarkers. *Brain* **140**, aww319 (2016).

40.   Wang, X. *et al.* Genetic determinants of disease progression in Alzheimer's disease. *J. Alzheimers.*

Dis. **43**, 649–55 (2015).

41.   Allen, M. *et al.* Human whole genome genotype and transcriptome data for Alzheimer's and other

neurodegenerative diseases. *Sci. Data* **3**, (2016).

42.   De Jager, P. L. *et al.* A multi-omic atlas of the human frontal cortex for aging and Alzheimer's

disease research. *Sci. Data* **5**, 180142 (2018).

43.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

44.   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-throughput

sequencing data. *Bioinformatics* **31**, 166–169 (2015).

45.   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear model

analysis tools for RNA-seq read counts. *Genome Biol.* **15**, (2014).

46.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and

microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

47.   A Bennett, D. *et al.* Overview and findings from the rush Memory and Aging Project. *Curr.*

Alzheimer Res. **9**, 646–663 (2012).

48.   Bennett, D. A. *et al.* Religious orders study and rush memory and aging project. *J. Alzheimer's*

Dis. 1–28 (2018).

49.   Allen, M. *et al.* Human whole genome genotype and transcriptome data for Alzheimer's and other

neurodegenerative diseases. *Sci. data* **3**, 160089 (2016).

50.   Ferretti, M. T. *et al.* Sex differences in Alzheimer disease—the gateway to precision medicine.

Nat. Rev. Neurol. 1 (2018).

51.   Deming, Y. *et al.* Sex-specific genetic predictors of Alzheimer's disease biomarkers. *Acta*

Neuropathol. **136**, 857–872 (2018).

52.   Mao, Q., Wang, L., Goodison, S. & Sun, Y. Dimensionality Reduction Via Graph Structure

Learning. in *Proceedings of the 21th ACM SIGKDD International Conference on Knowledge*

Discovery and Data Mining - KDD '15 765–774 (ACM Press, 2015).

doi:10.1145/2783258.2783309

53.   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-

generation DNA sequencing data. *Genome Res.* **20**, 1297–303 (2010).

54.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics **25**, 1754–60 (2009).

55.   Kang, H. M. *et al.* Variance component model to account for sample structure in genome-wide

association studies. *Nat. Genet.* **42**, 348–354 (2010).

Acknowledgements

This work was supported by NIA grants U54AG054345 and RF1AG057443. The ROSMAP Study data were provided by the Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago. Data collection was supported through funding by NIA grants P30AG10161, R01AG15819, R01AG17917, R01AG30146, R01AG36836, U01AG32984, U01AG46152, the Illinois Department of Public Health, and the Translational Genomics Research Institute. Mayo RNAseq Study data were provided by the following sources: The Mayo Clinic Alzheimer's Disease Genetic Studies, led by Dr. Nilufer Ertekin-Taner and Dr. Steven G. Younkin, Mayo Clinic, Jacksonville, FL using samples from the Mayo Clinic Study of Aging, the Mayo Clinic Alzheimer's Disease Research Center, and the Mayo Clinic Brain Bank. Data collection was supported through funding by NIA grants P50 AG016574, R01 AG032990, U01 AG046139, R01 AG018023, U01 AG006576, U01 AG006786, R01 AG025711, R01 AG017216, R01 AG003949, NINDS grant R01 NS080820, CurePSP Foundation, and support from Mayo Foundation. Study data includes samples collected through the Sun Health Research Institute Brain and Body Donation Program of Sun City, Arizona. The Brain and Body Donation Program is supported by the National Institute of Neurological Disorders and Stroke (U24 NS072026 National Brain and Tissue Resource for Parkinson's Disease and Related Disorders), the National Institute on Aging (P30 AG19610 Arizona Alzheimer's Disease Core Center), the Arizona Department of Health Services (contract 211002, Arizona Alzheimer's Research Center), the Arizona Biomedical Research Commission (contracts 4001, 0011, 05-901 and 1001 to the Arizona Parkinson's Disease Consortium) and the Michael J. Fox Foundation for Parkinson's Research. MSBB data were generated from postmortem brain tissue collected through the Mount Sinai VA Medical Center Brain Bank and were provided by Dr. Eric Schadt from Mount Sinai School of Medicine.

Author Contributions

S.M. and B.A.L. designed the study. S.M. and B.A.L. performed the analyses. S.M., C.P., S.J., G.G.,

A.K.G., S.K.S., P.L.D.J., N.E.T., G.W.C., L.M.M., and B.A.L. wrote the manuscript.

Tables

Table 1 - Representative significant Gene Ontology pathway enrichments (FDR<0.05) of differentially

expressed genes for each branch (FDR < 0.05).   Differential expressed genes are identified with an

ANOVA analysis, with Branch 1 as the reference.

Brain Region	Direction	Branch	Representative Enriched Gene Ontology Terms
TCX	Down	2	prespliceosome (GO:0071010), mitochondrial electron transport, cytochrome c to oxygen (GO: 0006123)
		3	negative regulation of microtubule polymerization or depolymerization (GO:0031111)
		4	mitochondrial electron transport, NADH to ubiquinone (GO: 0006120), spliceosomal tri-snRNP complex (GO:0097526), negative regulation of microtubule depolymerization (GO:0007026)
		5	axon (GO:0030424), protein kinase C activity (GO:0004697),
		6	gamma-tubulin large complex (GO:0000931), U1 snRNP (GO:0005685), mitochondrial respiratory chain complex IV (GO:0005751), response to cadmium ion (GO:0046686)
	Up	2	
		3	fatty acid elongase activity (GO:0009922), ubiquitin protein ligase activity (GO:0061630)
		4	transforming growth factor beta-activated receptor activity (GO:0005024), hippo signaling (GO:0035329), regulation of extrinsic apoptotic signaling pathway via death domain receptors (GO: 1902041), regulation of DNA repair (GO: 0006282)
		5	regulation of apoptotic process (GO:0042981), leptin mediated signaling pathway (GO:0033210), negative regulation of hippo signaling (GO:0035331), small GTPase binding (GO:0031267)
		6	extracellular ligand-gated ion channel activity (GO:0005230), integral component of mitochondrial inner membrane (GO:0031305)
DLPFC	Down	2	DNA repair (GO:0006281), intracellular protein transport (GO:0006886)
		3	mismatch repair complex binding (GO:0032404)
		4	
		5	mitochondrial respiratory chain complex assembly (GO: 0033108)
		6	
	Up	2	racemase and epimerase activity (GO: 0016857)
		3	racemase and epimerase activity (GO: 0016857)
		4	vesicle mediated transport (GO: 0016192)
		5	NuRD complex (GO: 0016581)
		6	microtuble motor activity (GO:0003777), AP-2 adaptor complex binding (GO:0035612)

Figure 1 - Overview of manifold learning for unraveling staging in Alzheimer's disease. A) Illustration of

steps in manifold learning. B) A common application of manifold learning used in computer vision to

order temporally untagged images into sequences. C) Illustration of lineage inference process for LOAD.

RNA-seq samples with different disease diagnoses were pooled, batch normalized, and a smooth

manifold was learned for each brain region across individuals (each point is an individual).

Figure 2 - Manifold learning accurately infers disease states and stages from RNA-seq samples. A)

Estimated disease progression trees from temporal cortex (TCX) and B) dorsolateral prefrontal cortex

(DLPFC) brain regions showing localization of identified LOAD samples on particular branches. C)

Distribution of pseudotime for AD cases and controls for both DLPFC and TCX. D) Distribution of

expression correlation with pseudotime for both LOAD GWAS genes and non-LOAD GWAS genes.

Figure 3 - Manifold learning replicates existing measures of staging in LOAD in DLPFC samples. A)

Samples colored by 3 different external measures of LOAD staging namely: Braak Score (tau pathology),

CERAD Score (amyloid pathology) and Cognitive Diagnosis (Clinical measure of disease severity).

Black lines denote inferred lineages. B) Distribution of samples by inferred stage for different distinct

stages in each of the three methods of measuring LOAD severity. Inferred disease stages generally

corresponded with all methods, and Cognitive diagnosis demonstrated the strongest alignment.

Figure 4 – Disease resistant state. A) The inferred manifold from the TCX region with samples colored

by their inferred disease subtype/state. State 5 (dots, circled) lies at the late end of the disease trajectory,

indicating a strong disease-like transcriptomic phenotype, yet most samples in the group did not have

pathologically diagnosed AD (Figure 2A). We hypothesize this group represents a disease resistant state

to the disease. B) Biclustering results of average expression from each disease state, with increased

expression of a gene cluster unique to State 5.