Molecular estimation of neurodegeneration pseudotime in older brains

Sumit Mukherjee[1,2*], Laura Heath[1*], Christoph Preuss[3], Suman Jayadev[4], Gwenn A. Garden[4], Anna K Greenwood[1], Solveig K Sieberts[1], Phillip L De Jager[5,6], Nilufer Ertekin-Taner[7,8], Gregory W Carter[3], Lara M Mangravite[1], Benjamin A Logsdon[1,9,10]

[1]Sage Bionetworks, Seattle, WA, USA.
[2]Microsoft, Redmond, WA, USA[†].
[3]The Jackson Laboratory, Bar Harbor, ME, USA.
[4]Department of Neurology, University of Washington, Seattle, WA, USA.
[5]Center for Translational & Computational Neuroimmunology, Department of Neurology, Columbia University Irving Medical Center, New York City, USA
[6]Taub Institute, Columbia University Irving Medical Center, New York City, USA
[7]Mayo Clinic Florida, Department of Neurology, Jacksonville, FL, USA.
[8]Mayo Clinic Florida, Department of Neuroscience, Jacksonville, FL, USA.
[9]Cajal Neuroscience, Seattle, WA, USA[†].

[†]Current Address.
[*]These authors contributed equally to this work.

[10]Correspondence:
Benjamin A Logsdon, PhD
Cajal Neuroscience
1616 Eastlake Ave E #208
Seattle, WA 98102
ben@cajalneuro.com

Abstract

The temporal molecular changes that lead to disease onset and progression in Alzheimer's

disease are still unknown. Here we develop a temporal model for these unobserved molecular

changes with a manifold learning method applied to RNA-Seq data collected from human

postmortem brain samples collected within the ROS/MAP and Mayo Clinic RNA-Seq studies.

We define an ordering across samples based on their similarity in gene expression and use this

ordering to estimate the molecular disease stage – or disease pseudotime - for each sample.

Disease pseudotime is strongly concordant with the burden of tau (Braak score, $P = 1.0\times10^{-5}$),

Aβ (CERAD score, $P = 1.8\times10^{-5}$), and cognitive diagnosis ($P = 3.5\times10^{-7}$) of LOAD. Early stage

disease pseudotime samples are enriched for controls and show changes in basic cellular

functions. Late stage disease pseudotime samples are enriched for late stage AD cases and show

changes in neuroinflammation and amyloid pathologic processes. We also identify a set of late

stage pseudotime samples that are controls and show changes in genes enriched for protein

trafficking, splicing, regulation of apoptosis, and prevention of amyloid cleavage pathways. In

summary, we present a method for ordering patients along a trajectory of LOAD disease

progression from brain transcriptomic data.

Introduction

Late-onset Alzheimer's disease (LOAD) is a devastating illness with no effective disease

modifying therapy, owing to a 99.6% failure of clinical trials[1]. There is a growing consensus that

the most effective treatments will intervene early in disease progression and halt disease

pathophysiological processes prior to conversion to LOAD[2]. In addition, there is increasing

recognition that LOAD may in fact be a spectrum of related diseases that have similar clinical

and neuropathological manifestations[3,4]. Devising successful therapeutic strategies will likely

require targeting potentially diverse early-stage disease processes that occur prior to a high

burden of neuropathology or cognitive impairment.

Current approaches to identify AD affected individuals include *in vivo* measures of the

pathological hallmarks of disease – amyloid, tau, and neurodegeneration – via CSF biomarkers

for amyloid and tau[5], positron emission tomography for amyloid and tau (PET)[6], and structural

and functional MRI of neurodegeneration. Cognitive assessments are used to estimate disease

burden[7], although measurable cognitive impairment generally indicates a sustained burden of

neuropathology and advanced neurodegeneration. Based on biomarker studies of AD, by the

time cognitive decline becomes detectable, neuropathological changes of AD have already

occurred, first in Aß and subsequently in tau related measures[8] and therefore cannot be used to

select patients for early disease stage studies. Furthermore, while these measures of disease

progression capture the overall increase in burden of pathology and cognitive decline, they do

not necessarily identify the dysfunctional molecular mechanisms that lead to neuropathology and

cognitive decline. There are likely many independent patient specific molecular pathways

present at an early stage in disease that then contribute to later stage disease progression[9,10]. This

motivates the need to identify these early stage molecular mechanisms driving disease

progression.

The Accelerating Medicines Project for Alzheimer's Disease (AMP-AD) consortia have

generated genome-wide transcriptomics of post-mortem brain tissue from patients across a broad

range of Alzheimer's disease neuropathological progression – including individuals with various

stages of AD neuropathology and those who lack AD neuropathology, but who may in fact

harbor early stage disease molecular processes. We therefore sought to chart the molecular

progression of the disease as reflected in the aggregate behavior of the brain transcriptome across

these individuals. While standard approaches such as differential expression or coexpression

analyses have proven informative[11–15], these analyses do not infer the relative stage of disease

progression or identify distinct disease subtypes. Here we propose an approach to analyze

population level RNA-seq data from post-mortem brain tissue to learn a tree structured

progression (**Figure 1**) that represents distinct sub-types of disease and the relative progression

of disease across patients. With this approach, we identify potentially generalizable trajectories

of LOAD across heterogeneous patient populations at all stages of disease. Furthermore, we

characterize molecular pathways that define disease stages – a potential source of biomarkers

and therapeutic interventions for early-stage disease processes along multiple different disease

trajectories.

To learn the molecular disease staging and neuropathologic progression tree we use a manifold

learning method[16]. Manifold learning refers to a group of algorithms that aim to recover the low

dimensional subspace underlying a high dimensional dataset. Previous authors use manifold

learning to estimate disease progression from neuroimaging data[17] and to study lineage

commitment of cells during differentiation from single cell RNA-seq (scRNA-seq)[18–21]. To our

knowledge, manifold learning has not been used to estimate disease progression and/or disease

stages from bulk RNA-seq data derived from post-mortem brain tissue. Henceforth, we refer to

manifold learning and lineage inference interchangeably in reference to the construction of a

disease progression tree. We demonstrate that these tools can estimate the disease staging and

progression tree (**Figure 2**) from bulk RNA-Seq data collected from post-mortem brain tissues in

a case/control cohort. Moreover, these trees show clear LOAD staging, enable the study of cell

type specific effects of LOAD, and allow the identification of genetic factors driving disease

progression.

Results

Manifold learning distinguishes pathologically defined LOAD from control

We first quantify the bulk RNA-Seq data from the ROS/MAP and Mayo Clinic cohorts into gene

counts and remove any batch effects introduced due to sequencing runs using standard count

normalization (see **Methods**). The data from the ROS/MAP cohort is sampled from the

dorsolateral prefrontal cortex (DLPFC), and the data from the Mayo Clinic cohort is sampled

from the temporal cortex (TCX). Patient clinical characteristics are reported in **Supplementary**

Table 1 and described in Methods. The full pipeline we use for RNA-Seq data generation and

quality control was recently reported[22]. The entire transcriptome comprises many genes which do

not have measurable expression or vary across case/control samples, which we remove in order

to reduce the noise in manifold learning[19]. To do this, we first perform differential expression

analysis between case/control samples separately for each study and retain genes that reach an

FDR of 0.10. To test if this biased the disease lineage inference, we also perform manifold

learning using only genes with high variance across samples, and we see a strong concordance

with disease lineages inferred with differentially expressed genes (**Supplementary Figure 1**).

Changing the significance threshold for the differential expression analysis to FDR < 0.01 did

not materially change these results (**Supplementary Figure 2**). We infer the disease lineage for

each brain region on this subset of retained genes (**Figure 2A-B**). Adjusting for post-mortem

interval (PMI) (**Supplementary Figure 3A, Supplementary Figure 4A**), 10 principal

components from a principal component analysis of genotype data to account for ancestry effects

(**Supplementary Figures 3B, Supplementary Figure 4B**), RNA integrity number (RIN)

(**Supplementary Figure 3C, Supplementary Figure 4C**), or all of these variables

(**Supplementary Figure 3D, Supplementary Figure 4D**) did not materially change the

overarching ordering of patients for either the TCX or DLPFC regions. Furthermore, to assess

the general robustness of the results we apply leave one out cross validation to infer disease

pseudotime for both DLFPC and TCX brain regions and find strong correlations between

lineages inferred with each sample removed, and the lineage for the entire sample set

(**Supplementary Figure 5**).

We visualize the clinical diagnosis of the samples on the inferred disease staging tree to verify

that there is indeed separation of AD patients across the tree. To determine if inferred tree

structure is an accurate model of disease progression, we introduce the notion of disease

pseudotime which is the geodesic distance along the tree from an inferred initial point to the

point of interest as a quantitative linear measure of LOAD stage. We scale this estimated disease

pseudotime to lie in the range $[0,1]$ to make the effects comparable between the two studies (and

brain regions). We show that for LOAD cases compared to controls there is a significant

association (P = 0.02 in Mayo and P= 2.0×10^{-6} in ROS/MAP, logistic regression) between the

estimated pseudotime and AD case/control status (**Figure 2C**). These effects are not abrogated

by adjusting for RIN, PMI, or ancestry in either tissue (**Supplementary Figure 6A-D,**

Supplementary Table 2). To assess whether the association between inferred disease

pseudotime is a phenomena in only the Mayo RNAseq and ROS/MAP RNA-seq data, we also

apply the lineage inference approach to expression array data from the Mayo eQTL study[23]

(**Methods**). These samples are derived from a completely independent set of donors than the

Mayo RNA-seq study[24]. Similarly, we restrict to only female samples, and test for an association

between inferred disease pseudotime and disease status (**Supplementary Figure 7A-B**). We see

a significant association between disease pseudotime and neuropathological AD diagnosis (P =

2.2×10^{-8}).

Furthermore, we observe strong evidence of sex heterogeneity when performing the manifold

learning approach and find that the manifolds inferred for female only samples show stronger

association with pseudotime than for male samples. This matches previous observations

concerning disease specific sex heterogeneity[22]. As such, we do not see as statistically

significant of an association between pseudotime and disease diagnosis in male samples (P =

0.040 in Mayo and P= 0.11 in ROS/MAP, logistic regression, **Supplementary Figure 8A-D**).

Similarly, the association between pseudotime and amyloid, tau, and cognitive diagnosis is

attenuated in male samples in ROS/MAP (**Supplementary Figure 9A-D**). For the combined

samples we see moderate evidence of disease association with pseudotime (P=0.003 in Mayo and

P=0.003 in ROS/MAP, logistic regression, **Supplementary Figure 10A-D**). The association

with neuropathological measures of disease is more robust in the combined sample

(**Supplementary Figure 11A-F**), but not as strong as in females only, hence we restrict to

female only analyses for all subsequent reported results.

We test whether genes in loci that have been implicated in genome wide association studies of

LOAD are associated with inferred disease pseudotime. We use the prioritized LOAD GWAS

genes[25], **Supplementary Table 3**, and compute the correlation between their expression and

inferred pseudotime (**Figure 2D**). When compared to the background of all genes, we see that

there is a significant increase in positive correlation with disease pseudotime for implicated

LOAD GWAS genes (P-value: 7.3×10^{-5} in Mayo and 5.6×10^{-3} in ROS/MAP). This effect is

robust to adjustments for PMI, RIN, or ancestry (**Supplementary Figure 12A-D**). Furthermore,

this does not appear to be driven by a small subset of outlier genes, but by the majority of the

distribution of LOAD GWAS genes. The fact that AD GWAS loci genes have expression

associations with pseudotime likely implies that the AD risk variants at these are also eQTL as

previously shown[26–29] and/or are members of co-expression networks that are differentially

expressed in AD[13,30].

To further explore the relationship between inferred disease stage and LOAD, we test for its

association with neuropathological and clinical measures of LOAD severity, namely: i) Braak

score, ii) CERAD score, and iii) cognitive diagnosis. The ROS/MAP study has numeric scores

for these categories available as covariates for each sample. Braak is a semi-quantitative measure

that increases with tau pathology[31] and CERAD is a semi-quantitative measure of density of

neuritic plaques[32]. We overlay these scores on the inferred manifold for the DLPFC brain region

(**Figure 3A**). We observe a progressive increase in tau, amyloid, and cognitive burden as we

traverse the inferred disease manifold (**Figure 3A**). This is further quantified by characterizing

the relationship between branches of the inferred manifold and Braak, CERAD, and cognitive

diagnosis (**Figure 3B**). We observe significant associations between pseudotime and Braak

score (P=1.0×10^{-5}), CERAD score (P=1.8×10^{-5}), and cognitive diagnosis (P=3.5×10^{-7}). In

ROS/MAP, adjustment for Braak score when fitting the DDRTree method attenuates the

association between pseudotime and disease states (P-value: 0.214, **Supplementary Figure**

13A-B), though there is still evidence of association with cognitive diagnosis (P-value: 0.03,

Supplementary Figure 13C-D). In the Mayo RNA-seq study we have Braak score and Thal

Amyloid scores for only a subset of samples but observe a similar pattern as in ROS/MAP

(**Supplementary Figure 14A-B**) for the samples that we do have data. There is a significant

association between Braak score and pseudotime (P-value: 5×10^{-5}) as well as Thal amyloid (P-

value: 1.7×10^{-5}) within this subset with available neuropathology data.

Comparison to other unsupervised learning approaches. We compare the manifold learning

approach to other unsupervised learning approaches including principal component analysis

(PCA), t-distributed Stochastic Neighbor Embedding (tSNE)[33], and Uniform Manifold

Approximation and Projection (UMAP)[34]. Correlations between the first two dimensions of each

of these approaches and the DDRTree learned pseudotimes are shown in **Supplementary Figure**

15A-F, Supplementary Figure 16A-F for DLPFC and TCX brain tissues respectively. We see

the strongest correlations between PCA1 and UMAP2 and pseudotime in both data-sets,

increasing our confidence that the overarching ordering of patients along a disease pseudotime is

a robust characteristic of the disease progression as reflected in the gene expression changes as a

function of disease, and not dependent on the underlying manifold learning approach. This is

further supported by inspecting the association between these approaches and Braak, CERAD,

and cognitive diagnosis (cogdx) scores for the DLPFC tissue (**Supplementary Table 4**).

Furthermore, the manifold learning approaches (DDRTree and UMAP) have much stronger

associations with Braak, CERAD, and cogdx scores than either PCA or tSNE. In fact, UMAP

has been proposed for lineage inference[35], and when we apply UMAP with lineage inference

using Monocle3, we observe similar results (**Supplementary Figures 17-18**) as with DDRTree

and Monocle2 (**Figure 2-3**), though the inferred pseudotimes from Monocle3 are not quite as

significant as the association with UMAP2 or from Monocle 2 with Braak, CERAD, and cogdx

(**Supplementary Table 4**).

Inferred staging recapitulates known biology of AD.

To demonstrate that the inferred disease pseudotime recapitulates known biology of LOAD, we

test for association between inferred disease stage and both the cellular response to disease and

the genetics of the disease. A prominent hypothesis in AD is that the effects of the disease vary

across different brain cell types, specifically neurons and glial subtypes. Current understanding

of the cell biology of the disease implicates progressive neuronal loss and increase in gliosis[36].

To test if the inferred pseudotime aligns with existing cell type specific hypotheses regarding

AD, we first selected from the genes used in lineage construction the marker genes for four key

cell types: neurons, astrocytes, microglia, and oligodendrocytes based on a previously published

brain cell atlas[37] (**Supplementary Table 5**). We then calculate the normalized mean expression

for the marker genes of each cell type and fitted a linear model to the mean expression with

disease pseudotime as the dependent variable. We find that, in both studies, the cell specific

marker gene levels show a statistically significant linear dependence on pseudotime

(**Supplementary Table 6**). Fitted effects recapitulate known neuropathologic changes which

occur in AD, namely: i) a reduction in the neuronal populations as AD progresses, and ii) an

increase in expression associated with activation of microglia, astrocytes, and oligodendrocytes

as AD progresses (**Figure 4**).

Next, we test for association between assigned lineage state in ROS/MAP (DLPFC) and Mayo

(TCX) and APOE e4 status (**Supplementary Figure 19**). For reference, the inferred trees for

TCX and DLPFC each resolve into 6 branches (**Figure 5A,S20**). Carriers of the APOE e4 allele

are significantly enriched on the State 4 branch in TCX (P-value = 0.027, unadjusted), and

suggestively enriched on the State 5 branch (P-value = 0.06, unadjusted), compared to the State

1 branch (logistic regression). Similarly, in the Mayo eGWAS study, when we perform an

ordinal logistic regression of APOE e4 dosage and disease pseudotime we see a significant

positive association as a function of pseudotime (P-value = $6.9\text{x}10^{-4}$, **Supplementary Figure**

7C).

Genetic factors associated with inferred disease staging

Lineage inference of LOAD transcriptomes provides a quantitative measure of disease

progression for genetic associate testing, and the significantly greater correlation between

pseudotime and gene expression for known LOAD risk genes (**Figure 2D**) suggests that the

observed differences in disease trajectories are influenced by genetic factors. To test this

hypothesis, we perform single variant analysis using whole-genome sequencing data for 305

patients from the ROS/MAP and 131 patients from the Mayo cohort. Despite the limited sample

size, resulting in lack of statistical power to discover genome-wide significant associations,

multiple variants reach a genome-wide suggestive threshold of $p < 1\text{x}10^{-5}$ (**Supplementary**

Table 7). We do not see evidence of population stratification in the analysis (**Supplementary**

Figure 21-22). Notably, the most significant association with pseudotime for the ROS/MAP

cohort is observed at the *PTPRD* locus (rs7870388, p = $1.31 \text{ x } 10^{-6}$) (**Supplementary Figure 23,**

Supplementary Table 7). The *PTRPD* locus is associated with the susceptibility to

neurofibrillary tangle independent of amyloid deposition in the ROS/MAP cohort[38]. For the

Mayo Clinic cohort, known LOAD variants in the *APOE* (rs6857, p = 9.18 x 10^{-6}) and *BIN1*

(rs62158731, p= 4.68 x 10^{-5}) loci overlap with variants associated with inferred disease stage

(**Supplementary Figure 23, Supplementary Table 7**)[39]. When comparing our association

results for inferred disease stage with summary statistics from a large-scale case-control

approach, we identify multiple variants which have been previously associated with LOAD in

the IGAP cohort (**Supplementary Table 8**). Furthermore, we identify several genes associated

with inferred disease stage (*ADAMTS14*, *IL7*, *MAN2B1*) linked to immune and lysosomal storage

function (**Supplementary Figure 23**,**Supplementary Table 7**). *IL-7* has been proposed as an

inflammatory biomarker for LOAD that correlates with disease outcome and severity[40].

ADAMTS14 is part of a locus that has been previously linked with Alzheimer susceptibility and

plays an important role in the regulation of immune function via TGF-beta signaling.

New disease insights identified from inferred disease lineages. Another important direction of

study in the field of Alzheimer's is the identification of disease subtypes, which has so far

predominantly been done using imaging data[41]. The branches of the inferred disease trees

provide a transcriptomic-based approach to identify disease subtypes. In both brain regions and

in two separate cohorts, there were two distinct early-lineage branches corresponding to

predominantly control samples, which we interpret as different initial paths towards the disease.

Similarly, both brain regions feature several distinct branches with predominantly LOAD

samples (**Figure 2A-B**).

Branch-specific differential expression patterns. To study the genes and pathways specific to

each branch, we perform a branch-specific differential expression analysis with an ANOVA

model using the branches with the highest proportion of controls as the reference branch for

DLPFC (**Supplementary Table 9**) and TCX (**Supplementary Table 10**). We see many genes

are differentially expressed between the control branch and branches that are enriched in the

affected individuals (**Supplementary Table 11**). We test for overlap between the differentially

expressed gene sets between the two studies (**Supplementary Figure 24**), and find significant

overlaps in branches enriched for late stage disease cases, especially between up-regulated genes

in State 6 of DLPFC and up regulated genes in State 5 of TCX (P-value: 4.1×10^{-108}, OR: 4.5,

Fisher's exact test), as well as genes that are up-regulated in State 6 of DLPFC and in State 4 of

TCX (P-value: 1.8×10^{-14}, OR: 1.9, Fisher's exact test), and more modestly for genes that are

down regulated in State 6 of DLPFC and down regulated in State 3 of TCX (P-value: 1.1×10^{-6},

OR: 1.6, Fisher's exact test).   Next, we performed an enrichment analysis on each of these

differentially expressed gene sets with the enrichR[42] package for Gene Ontology[43] annotations

(**Methods**). The results of this enrichment analysis for DLPFC and TCX tissues

(**Supplementary Tables 12-13**). Only gene sets with significant enrichment are shown (FDR

adjusted p-value < 0.05). Overall, we see a pattern of loss of expression of basic cell biology

mechanisms in early-stage branches including RNA splicing, mitochondria function, protein

transport, and DNA repair. Late-stage branches were characterized by increased immune

response (e.g. TGFb/WNT signaling) and apoptotic activity (**Table 1-2**).

While studying the different branches in the two brain regions, we observe a branch (branch 5)

that corresponds to a group of predominantly neuropathological control samples from the Mayo

RNA-seq cohort that were in close proximity to a branch with predominantly LOAD samples

(branch 4) on the inferred disease lineage (**Figure 5A**). However, most of the samples on branch

5 are neuropathological controls as defined by the Mayo diagnostic criteria. We bi-cluster the

mean expression of genes in each branch and the branches themselves (**Methods**). This

clustering analysis (**Figure 5B**) shows that the closest branch to this potentially disease resistant

branch contains the highest proportion of AD samples. While the stage proximity implies some

transcriptomic similarity between these controls and nearby cases, we also see a secondary

cluster of genes with increased expression in the resistant state while having reduced expression

in all other states. We perform an enrichment analysis on this set of genes and find significant

GO terms corresponding to protein transport (GO:0015031), regulation of mRNA splicing, via

spliceosome (GO:0048024), negative regulation of apoptotic process (GO:0043066), and

regulation of amyloid-beta clearance (GO:1900221) (Cluster4, **Supplementary Table 14**)**. It is

possible that these potentially disease resistant individuals have compensatory mechanisms

which suppress the hallmarks of disease despite sharing gene expression patterns with

pathologically affected individuals.

To replicate this observation, we perform a differential expression analysis on individuals in the

Mayo eGWAS study where we consider individuals that are in the top quintile of pseudotime but

are classified as neuropathological controls as resistant individuals (**Supplementary Figure 7D**,

Methods). To test if these individuals also have a similar resistant molecular endophenotype, we

compare the overlap between various differential expressed gene sets derived from these

resistant individuals and the gene sets identified in the biclustering of the Mayo RNA-seq data

(**Supplementary Figure 25**). We observe that there is a highly statistically significant overlap

between genes that are upregulated in these Mayo eGWAS resistant individuals (P-value:

2.6×10^{-51}, OR: 2.9, Fisher's exact test), and the Cluster 4 genes that are upregulated in the branch

5 Mayo RNA-seq samples (**Supplementary Figure 25**).

Discussion

Here we proposed an approach to infer the Alzheimer's disease severity and disease subtypes in

an unsupervised manner from post-mortem bulk RNA-seq data that gets directly at the challenge

of identifying the temporal progression of disease in the disease resistant tissue. Our strategy

utilized a manifold learning approach to infer a disease progression tree from cross-sectionally

collected patient samples from two different brain regions. The underlying assumption of our

approach is that the inferred disease progression from cross sectional samples serves as a proxy

for the unobserved progression of the disease across subtypes of LOAD. We validated this

hypothesis through comparisons with neuropathological measures of disease stage severity and

against known cell type specific effects caused by the disease. While one could argue that the

method is merely classifying patients as either disease cases or controls based on expression

signatures of the hallmarks of disease, we see at least three advantages of this approach beyond

that interpretation. First the application of this method appears to produce a more quantitative

measure of disease state than strictly neuropathological assessments - as born out through the

identification of distinct genetic loci that replicate based on IGAP summary statistics. This

suggests that it may be adding information related to other aspects of disease such as the effect of

neuroinflammation or neuronal injury. In addition we see evidence of neuropathological controls

that are disease resistant given their molecular state in two independent studies – which would

not be detectable with standard neuropathological or clinical assessments – and could provide

important molecular clues to mechanisms of disease resistance. Finally, there is the potential

that specific pathways associated with early stage disease processes can be characterized which

is desperately needed for hypothesis generation in the field. Furthermore, this approach provides

clues to better understanding the molecular heterogeneity of disease by identifying specific

pathways that are dysregulated in subsets of patients at different disease stages. This opens up

the possibility of better patient stratification and precision medicine.

We observed that different biological processes vary as a function of inferred disease stage, and

that early-stage disease processes - include RNA-splicing, mitochondrial function, and protein

transport - implicate multiple basic cell biology mechanisms as potential early stage disease

processes for further study in relevant model systems. Additionally, the manifold learning

method identified 6 subtypes of LOAD from RNA-seq (i.e. branches) suggesting the LOAD

populations should be stratified by better biomarkers with tailored treatment strategies. To

identify and test these stratifications future studies could focus on longitudinal cohorts of patients

with rich molecular and imaging data to identify biomarkers that can accurately and precisely

stratify patients into the underlying molecular subtypes at different relative stages of disease.

Furthermore, we observe a disease resistant sub-type of patients. This disease resistance should

be tested in disease model systems, to identify if neuropathological readouts can be modified by

altering the function of the pathways identified in our analysis (e.g. APP processing, RNA

splicing, apoptosis, protein trafficking). While this preliminary observation needs to be validated

in another cohort, it has the potential to be a source of hypotheses for therapeutic development.

Specifically, for constructing better combination therapy hypotheses that may confer

neuroprotection, even in patients that are mildly affected by disease.

LOAD is a complex and heterogeneous disease encompassing a broad spectrum of clinical

symptoms. Disease progression can vary widely between patients leading to different rates of

cognitive decline. Several lines of evidence suggest that these differences in progression are

modified by multiple genetic factors affecting the transition from one pathological state to

another[44,45]. However, it has remained difficult to assess the role of genetic variants affecting

disease trajectories by case-control approaches alone. Here, we showed that our expression trait

pseudotime might be used as a molecular phenotype to identify AD loci associated with different

disease progression states across AD patients. Despite a limited sample size, we identified

previously associated AD candidate loci in the ROSMAP (*PTPRD*) and Mayo (*BIN1, APOE*)

cohorts with suggestive significance ($p < 1 \times 10^{-5}$). Variants in *PTPRPD* have been associated

with the susceptibility to neurofibrillary tangles, independent of amyloid burden. This is in line

with the results from the differential gene expression analysis of pseudotime branches showing

an enrichment of molecular pathways implicated in TAU pathology. Furthermore, our analysis

revealed several loci linked to immune function (*ADAMTS14*, *IL7*) and neurotransmitter

signaling (*CHRM2*, *CHRM3*) processes associated with disease pseudotime (**Supplementary**

Table 4). Future studies will be needed to replicate these findings in independent cohorts of

LOAD and validate the role of candidate genes in LOAD related disease progression by first

identifying peripheral biomarkers that correspond to this molecular definition of disease stage,

and then testing for GWAS association with that disease stage. Subsequent results can improve

functional interpretation by linking candidate genes with ordered pathological processes.

Methods

ROS/MAP and Mayo RNAseq study population characteristics. Detailed descriptions of cohort

and patient characteristics included in this study can be found in previously published work[46,47].

Patient characteristics included in this study are summarized in **Supplementary Table 1**,

stratified by sex. In brief: for Mayo samples, AD diagnosis was performed according to

NINCDS-ADRDA criteria (probable or possible AD); control individuals had Braak NFT stage

≤ 3, CERAD score < 2.5, and lacked other pathologic diagnoses; Path.Aging are individuals who

lacked any pathologic diagnoses and had Braak NFT ≤ 3 and CERAD score ≥ 2. Progressive

supranuclear palsy (PSP) individuals were diagnosed neuropathologically by a single

neuropathologist. (For further details, see Allen et al 2016). For ROSMAP samples, AD

diagnosis was according to NIA Reagan criteria, which combines neuropathology and clinical

data; Control individuals had no signs of cognitive impairment; and Other individuals had MCI,

mixed pathology, or other form of dementia. Age at death was collected for all patients in the

ROSMAP study, though age at first AD diagnosis had a high degree of missingness and thus was

not used as a variable in follow-up analyses (see **Supplementary Table 1**). Braak stage

indicates the measure of severity of NFT pathology. Stages I and II indicate NFTs confined

mainly to the entorhinal region of the brain, stages III and IV indicate involvement of limbic

regions, and V and VI indicate moderate to severe neocortical involvement. CERAD score is a

semiquantitative measure of neuritic plaques. 1=definite AD; 2=probable AD; 3=possible AD;

4=No AD. Cognitive diagnostic category (as determined by neurologist): 1) NCI; 2) MCI and no

other cause of CI; 3) MCI and another cause of CI; 4) AD and no other cause of CI; 5) AD and

another cause of CI; 6) Other primary cause of dementia[48]. Thal amyloid stages: phases of

amyloid deposition. 0) no amyloid; 1) isocortical phase; 2) limbic phase; 3) basal ganglia phase;

4) basal forebrain and midbrain phase; 5) pons/medulla oblongata and cerebellum phase. The list

of differentially expressed genes used to create the Monocle objects were based on results which

included the whole data set.

RNA sequencing. The details of the sample collections, postmortem sample characteristics, the

tissue and RNA preparations, the library preparations and sequencing technology and

parameters, and sample quality control filters are provided in previously published work[46,49]. For

the bioinformatic pipeline to produce gene level counts we applied a standard pipeline[22] where

sequencing reads were aligned to the GENCODE24 (GRCh38) reference genome with STAR[50],

and gene counts generated using the HTSeq algorithm[51]. Genes that had more than one counts

per million total reads in at least 50% of samples in each tissue and diagnosis category were used

for further analysis.

Differential Expression analysis on Mayo and ROS/MAP cohorts. For gene filtering we used

false discovery rate of 0.05 from the previously published differential expression analysis of

Mayo and ROS/MAP RNA seq data[22]. Briefly, case control status was harmonized across the

Mayo and ROS/MAP cohorts, where controls were defined as individuals with a low burden of

amyloid and tau based on CERAD and Braak scores, and cases with a high burden. Furthermore

in ROS/MAP, clinical diagnosis was also used with controls having to have no cognitive

impairment, and cases have probably AD[22]. Differential expression analysis was run on suitably

normalized data – using conditional quantile normalization to account for variation in gene

length and GC content, removing sample outliers, covariate identification adjustment, with

sampling abundance confidence estimated using a weighted linear model with the voom-limma

package[22,52,53]. A fixed/mixed effect linear model is used to fit the differential expression model

on the normalized data[22].

Manifold learning for LOAD. Manifold learning refers to a group of machine learning algorithms

that recover a low dimensional subspace underlying a high dimensional dataset. Manifold

learning approaches are typically used in datasets or applications where data samples lie on an

underlying low dimensional latent space (e.g. a tree, a line, a curved plane). The low

dimensional space is learned via a projection from the high dimensional space of the observed

data (e.g. RNA-seq profiles across hundreds of patient samples) down to a low dimensional

space with suitable regularization constraints to enforce smoothness and the structural constraints

of the low dimensional space. (**Figure 1A**). Due to the necessary assumption of an underlying

latent subspace, manifold learning is commonly used in applications where it is known that the

observed data is obtained from a progression of some kind; e.g., i) to infer the temporal ordering

of a sequence of images, or ii) to infer the approximate lineage of cells in a differentiation

trajectory using single cell RNA-Seq data (**Figure 1B-C**).

Here, we repurpose methods originally developed for learning cell lineage using scRNA-Seq

data, to infer the staging of Alzheimer's disease (AD) using bulk RNA-Seq data from post-

mortem brain samples with known AD diagnosis status. Since bulk RNA-Seq has many of the

same sampling and distributional properties as scRNA-Seq, we observe that scRNA-Seq

methods are applicable with no additional modifications. As such, we use the DDRTree manifold

learning approach available in the Monocle2 R package[54]. However, we also show that the

estimated staging of disease is quite similar across some of the other common methods used for

scRNA-Seq lineage estimation (**Supplementary Figures 26-28**) including Monocle1[18] and

diffusion pseudotime (DPT)[55].

The RNA-Seq data used in this study was generated from post-mortem brain homogenate

samples, and obtained from two separate studies that are a part of the Accelerating Medicines

Partnership in Alzheimer's Disease (AMP-AD) consortium, namely: i) the Religious Orders

Study and the Memory and Aging Project (ROSMAP)[56,57], and ii) the Mayo RNA-seq study[49].

For this paper, we focused our analysis on the temporal cortex (TCX) and dorsolateral prefrontal

cortex (DLPFC) tissue samples. Within the Mayo RNA-seq study the TCX samples are derived

from individuals neuropathologically defined as either aged controls, LOAD cases, Progressive

Supranuclear Palsy (PSP) cases, or pathological aging (PA) cases[49].   The ROSMAP study is a

prospective longitudinal cohort of an aging population, and has samples from participants with

clinical and neuropathological diagnoses of LOAD[46], aged controls, and individuals with mild

cognitive impairment. Furthermore, most results presented in the main paper are from female

samples only unless indicated otherwise, as we observed significant sex differences in the

transcriptomic data consistent with current knowledge of sex differences in LOAD[58,59].   For

replication, we also consider microarray data generated from the Illumina DASL gene expression

platform from the Mayo eGWAS study from TCX for N=186 patients, of which 108 were

neuropathologically confirmed AD Cases and 78 were controls[23]. Probes were mapped to genes

using BioMart. Data was adjusted for plate using ordinary least squares regression prior to

manifold learning.

Manifold learning using Discriminative Dimensionality Reduction Tree (DDRTree). DDRTree is

a manifold learning algorithm that infers a smooth low dimensional manifold by an approach

called reverse graph embedding. Briefly, the algorithm simultaneously learns a non-linear

projection to a latent space where the points lie on a spanning tree. A reverse embedding is also

simultaneously learned from the latent space to the high dimensional data. The DDRTree

algorithm can be posed as the following optimization problem:

$$\sum_{i=1}^{N} ||\mathbf{x_i} - \mathbf{Wz_i}||^2 + \frac{\lambda}{2}\sum_{k,k\prime} b_{k,k\prime}||\mathbf{Wy_k} - \mathbf{Wy_{k\prime}}||^2 + \gamma\left[\sum_{k=1}^{K}\sum_{i=1}^{N} r_{i,k}||\mathbf{z_i} - \mathbf{y_k}||^2 + \sigma r_{i,k}\log{(r_{i,k})}\right]$$

$$\text{s.t.}\quad \mathbf{B} \text{ represents a spanning tree,}$$

$$\mathbf{W^T W} = \mathbf{I}, r_{i,k} \geq 0, \sum_{k=1}^{K} r_{i,k} = 1 \qquad\qquad (1)$$

Here, $\{\mathbf{x_i}\}_{i=1}^{N} \in R^{genes}$ represents RNA-Seq data from each patient sample, $\{\mathbf{z_i}\}_{i=1}^{N} \in R^2$

represents the latent representation of each sample as inferred by the algorithm, $\{\mathbf{y_k}\}_{k=1}^{K}$

represents the centers of clusters in the dataset, $\mathbf{W} \in R^{2 \times genes}$ represents an inverse mapping

from the latent space to the high dimensional space of RNA-Seq data, $\mathbf{B} \in R^{K \times K}$ represents a

spanning tree on which the centers of the clusters lie and $\mathbf{R} \in R^{N \times K}$ captures the soft clustering

information of samples in the dataset. The first term of the optimization problem is responsible

for learning a low dimensional representation of the data such that an inverse mapping exists to

the high dimensional data points, the second term learns the tree structure of the points and the

third term learns a soft clustering for the latent dimension points as well as the centers of the

clusters. Despite the non-convexity of the problem, each individual optimization variable can be

solved for efficiently using alternative minimization[60]. This algorithm was implemented using

the Monocle package in R[19]. When fitting the Monocle objects we also considered various

adjustments to the expression data prior to manifold learning for 10 principal components of

genetic ancestry, RNA integrity number, postmortem interval, age, Braak score - among other

potential confounds. The code to infer the lineage in MayoRNAseq is available here

https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/TCX_GenerateMonocleDS_new.R, and

code used to infer the lineage in ROSMAP is available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/DLPFC_GenerateMonocleDS_new.R.

Code to perform analysis on Mayo eGWAS study is available here: https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_ben_april_2020/mayo_egwas_GenerateMo

nocleDS_new.R.

Branch assignment and pseudotime calculation for samples. Branch assignment and pseudotime

calculation was also performed using the *Monocle* package[19]. Briefly, pseudotime is calculated

by first identifying a root point on one of the two ends of the maximum diameter path in the tree.

Then the pseudotime of each point is calculated by projecting it to its closest point on the

spanning tree and calculating the geodesic distance to the root point. Assigning samples to

branches is done by first identifying the branches of the spanning tree and then assigning

samples to the branch on which their projection to the spanning tree lies on. Robustness of

pseudotime was assessed with leave-one-out cross validation by dropping one sample at a time,

running the DDRTree method with Monocle, and then computing the absolute value of the

correlation between the pseudotime estimated with the reduced data-set, and the pseudotime

estimated with the full data set. Alternative approaches for performing dimensionality reduction

included principal component analysis (PCA), t-stochastic neighborhood embedding (tSNE)[33],

and uniform manifold approximation and projection (UMAP)[34], which were all run on the same

data set as the DDRTree method was run in R (here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/TCX_GenerateMonocleDS_new.R and

https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/DLPFC_GenerateMonocleDS_new.R).

Association of pseudotime with AD status, hallmarks of Alzheimer's disease, and cognitive

diagnosis. We test for association between disease pseudotime and AD case or control status

with logistic regression with AD case or control status as the outcome and inferred pseudotime

as the dependent variable in both the Mayo and ROS/MAP studies. We test for association

between pseudotime and hallmarks of disease in the ROS/MAP studies for both Braak (measure

of tau pathology) score and CERAD score (measure of amyloid pathology) with an ordinal

logistic regression model, with the neuropath score as the ordered outcome, and pseudotime as

the dependent variable. Finally, we test for association between disease pseudotime and

cognitive diagnosis for the following ordered clinical diagnoses of no cognitive impairment, mild

cognitive impairment, and probable Alzheimer's disease with an ordinal logistic regression

model. All code for running these association tests is available https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/paper_figures.Rmd.

Inferring cell type specific expression patterns given marker gene expression as a function of

pseudotime. List of marker genes for different major cell types in the brain was curated from a

previously published brain cell expression signature study[37]. The marker gene list was then

pruned to include only genes that were included in lineage construction. Each gene's expression

as a function of pseudotime was then obtained by smoothing using a smoothing spline of degree

of freedom = 3 and normalized to lie in [0,1]. The smoothing was done to remove the effects of

technical noise introduced due to RNA-Seq and the normalization was done since the absolute

expression levels of genes might be very different from each other. The smoothed and

normalized expression of marker genes for each category were then averaged to obtain the

average marker gene expression as a function of pseudotime.   A linear model was used to test

for association between average expression of a given cell type expression signature and

pseudotime.

Association between GWAS loci and correlation with pseudotime. To test for association

between pseudotime and LOAD GWAS genes, we computed the Spearman's correlation

between each gene's expression and pseudotime in the Mayo and ROS/MAP studies. Next, we

considered the 60 highly prioritized genes (priority score greater than 4) identified within AD

GWAS loci by the International Genetics of Alzheimer's Project (IGAP)[25]. We test for a

difference between the correlation with pseudotime of background of all other genes and the

IGAP AD genes using a linear model and see a significant increase in correlation between gene

expression and pseudotime in both the Mayo and ROS/MAP study for AD GWAS genes.

Branch specific differential expression analysis. We perform a state specific differential

expression analysis using a one-way ANOVA model in both the Mayo and ROS/MAP studies.

The branch with the highest proportion of AD controls is defined as the reference branch for all

analyses. We use Tukey's honest significant difference method to compute P-values for the test

for change in expression of a given gene compared to the reference branch. Genes are grouped

based on their branch and direction of change in expression for further downstream pathway

enrichment analyses. Overlap between differential expressed genes was depicted using UpSet

plots[61] (**Supplementary Figure 24**). Code to run analyses are available here

https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/DLPFC_DE_Anova.R for ROS/MAP

and here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/TCX_DE_Anova.R for Mayo.

Branch specific gene expression signatures. Branch specific expression signature was obtained

by first calculating the average normalized expression for all genes in each state/branch. This

was followed by performing a bi-clustering using the pheatmap package in R (https://cran.r-

project.org/web/packages/pheatmap/index.html) which uses hierarchical clustering on both

samples and genes. We also used the pheatmap R package to visualize the state specific

expression signatures.

Disease resistant subgroup validation analysis. After identifying potentially disease resistant

individuals in the Mayo RNAseq study based on the TCX brain region (individuals from branch

5, **Figure 5A**), we considered the Mayo eGWAS TCX data, and defined neuropathological

controls with pseudotimes in the top quintile of all pseudotimes as disease resistant (n=9). We

then performed a differential expression analysis using linear regression to identify array probes

that were differentially expressed between resistant and non-resistant individuals, of which there

were more than 5000 probes that were either up or down regulated at an FDR of 0.05. Overlaps

were explored between the branch specific gene clusters from Mayo RNAseq (Figure 5B) and

these Mayo eGWAS resistance differential expressed probes using UpSet plots[61]

(**Supplementary Figure 25**).

Gene set enrichment analyses. For each branch specific differential expression gene set (DEGs)

in both Mayo RNAseq and ROS/MAP we perform a gene set enrichment analysis against Gene

Ontology pathways using the enrichR[42] R package. Only pathways with FDR < 0.05 are

reported. The code we used to run the ROS/MAP DEG enrichments are available here

https://github.com/Sage-Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/lineage.Rmd, the

code we used to run the Mayo DEG enrichments are available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/lineageTCX.Rmd, and the code we used

to run the branch specific gene expression signature pathway enrichments is available here

https://github.com/Sage-Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/resilience.Rmd.

Whole-genome sequencing. Whole-genome sequencing was performed at the New York Genome

Center for all individuals from the ROS/MAP and Mayo cohorts. Detailed information for both

data sets can be accessed via synapse (DOI:10.7303/syn2580853). Briefly, 650ng of genomic

DNA from whole blood was sheared using a Covaris LE220 sonicator. DNA fragments

underwent bead-based size selection and were subsequently end-repaired, adenylated, and ligated

to Illumina sequencing adapters. Libraries were sequenced on an Illumina HiSeq X sequencer

using 2 x 150bp cycles. Paired-end reads were aligned to the GRCh37 (hg19) human reference

genome using the Burrows-Wheeler Aligner (BWA-MEM v0.7.8) and processed using the

GATK best-practices workflow[62,63]. This included marking of duplicate reads by the use of

Picard tools v1.83, local realignment around indels, and base quality score recalibration (BQSR)

via Genome Analysis Toolkit (GATK v3.4.0). Joint variant calling files (vcfs) for whole-genome

sequencing data for the Mayo and ROS/MAP cohort were obtained through the AMP-AD

knowledge portal (DOI:10.7303/syn10901595).

Single variant association with pseudotime in two independent cohorts. Likelihood ratio tests

within a linear regression framework were used to model the relationship between the

quantitative expression trait pseudotime and genetic variants in 436 AD cases. Genome-wide

genetic association analysis was performed for 305 female patients in the ROS/MAP cohort and

131 female patients in the Mayo cohort for which both genotyping and post-mortem RNA-seq

data was available. An efficient mixed model approach, implemented in the EMMAX software

suite, was used to account for potential biases and cryptic relatedness among individuals[64]. Only

variants with MAF > 0.05, genotyping call rates > 95%, minimum sequencing depth of 20 reads

and Hardy-Weinberg equilibrium $p > 10^{-4}$ were considered for analysis. Quantile-quantile plots

(**Supplementary Figures 21-22**) for the test statistics showed no significant derivation between

expected and observed p-values, highlighting that there is no consistent differences across cases

and controls except for the small number of significantly associated variants. Furthermore, the

genomic inflation factor (lambda) was determined to be 0.99 for the Mayo and 0.98 for the

ROS/MAP single variant association tests. This highlights that potential confounding factors,

such as population stratification have been adequately controlled.

Data Availability

All source data analyzed in the study is publicly available[22–24,46]. Specifically, we use a version

of the RNA-seq data from the ROS/MAP study (DOI:10.7303/syn8456638.22) and RNA-seq

data from the Mayo RNAseq (DOI:10.7303/syn8466816.19) run through the same bioinformatic

processing pipeline[22]. The array expression data from the Mayo eGWAS study is available at

DOI:10.7303/syn3617054.1. Pseudotimes for ROS/MAP individuals are available for all

individuals, females, and males respectively (DOI:10.7303/syn23446661.2,

DOI:10.7303/syn22822695.1, DOI:10.7303/syn23446654.3). Pseudotimes for MayoRNAseq

individuals are available for all individuals, females, and males respectively

(DOI:10.7303/syn23446689.1, DOI:10.7303/syn22822691.1, DOI:10.7303/syn23446688.1).

Pseudotimes for female MayoeGWAS individuals are available (DOI:10.7303/syn22822690.1).

Source data for **Figure 2C** (DOI:10.7303/syn23246577.1), **Figure 2D**

(DOI:10.7303/syn23246583.1), **Figure 3B** (DOI: 10.7303/syn23246585.1), **Figure 5B**

(DOI:10.7303/syn22822693.1), and **Supplementary Figure 6A** (DOI:10.7303/syn23445580.2),

Supplementary Figure 6B (DOI:10.7303/syn23445582.3), **Supplementary Figure 6C**

(DOI:10.7303/syn23445583.2), **Supplementary Figure 6D** (DOI:10.7303/syn23445584.2),

Supplementary Figure 7B (DOI:10.7303/syn23246588.1), **Supplementary Figure 8C** (DOI:

10.7303/syn23446680.1), **Supplementary Figure 8D** (DOI:10.7303/syn23446682.3),

Supplementary Figure 9D-F (DOI:10.7303/syn23448900.1), **Supplementary Figure 10C**

(DOI:10.7303/syn23446681.1), **Supplementary Figure 10D** (DOI:10.7303/syn23446691.1),

Supplementary Figure 11D-F (DOI:10.7303/syn23448904.1), **Supplementary Figure 12A**

(DOI:10.7303/syn23446257.3), **Supplementary Figure 12B** (DOI:10.7303/syn23446326.2),

Supplementary Figure 12C (DOI:10.7303/syn23446331.1), **Supplementary Figure 12D**

(DOI:10.7303/syn23446332.1), **Supplementary Figure 13B** (DOI:10.7303/syn23450641.1),

Supplementary Figure 13D (DOI:10.7303/syn23448918.1), **Supplementary Figure 13E**

(DOI:10.7303/syn23452920.1), **Supplementary Figure 14B** (DOI:10.7303/syn23468302.1),

Supplementary Figure 17C (DOI:10.7303/syn23505161.1), **Supplementary Figure 17D**

(DOI:10.7303/syn23505539.1), **Supplementary Figure 18D-F** (DOI:10.7303/syn23508896.1),

Supplementary Figure 24 (DOI:10.7303/syn23246590.1), **Supplementary Figure 25**

(DOI:10.7303/syn23246594.1), **Supplementary Figure 28** (DOI:10.7303/syn23246595.1) are

also available.

Code Availability

All code is publicly available (https://github.com/Sage-Bionetworks/AMPAD_Lineage).

References to code to perform specific analyses are described in detail in the **Methods**.

Competing Interests

The Authors declare no competing interests.

References

1. Cummings, J. L., Morstorf, T. & Zhong, K. Alzheimer's disease drug-development

pipeline: few candidates, frequent failures. *Alzheimers. Res. Ther.* **6**, 37 (2014).

2. Cummings, J. L., Doody, R. & Clark, C. Disease-modifying therapies for Alzheimer

disease: Challenges to early intervention. *Neurology* **69**, 1622–1634 (2007).

3. Ferreira, D. *et al.* Distinct subtypes of Alzheimer's disease based on patterns of brain

atrophy: longitudinal trajectories and clinical applications. *Sci. Rep.* **7**, 46263 (2017).

4. Bredesen, D. E. Metabolic profiling distinguishes three subtypes of Alzheimer's disease.

Aging (Albany NY) **7**, 595 (2015).

5. Brier, M. R. *et al.* Tau and A$β$ imaging, CSF measures, and cognition in Alzheimer's

disease. *Sci. Transl. Med.* **8**, 338ra66--338ra66 (2016).

6. Gordon, B. A. *et al.* The relationship between cerebrospinal fluid markers of Alzheimer

pathology and positron emission tomography tau imaging. *Brain* **139**, 2249–2260 (2016).

7. Dichgans, M. *et al.* METACOHORTS for the study of vascular disease and its

contribution to cognitive decline and neurodegeneration: An initiative of the Joint

Programme for Neurodegenerative Disease Research. *Alzheimer's Dement.* **12**, 1235–

1249 (2016).

8. Jack, C. R. *et al.* Tracking pathophysiological processes in Alzheimer's disease: an

updated hypothetical model of dynamic biomarkers. *Lancet. Neurol.* **12**, 207–16 (2013).

9. Au, R., Piers, R. J. & Lancashire, L. Back to the future: Alzheimer's disease heterogeneity

revisited. *Alzheimer's Dement. (Amsterdam, Netherlands)* **1**, 368–370 (2015).

10. Carrasquillo, M. M. *et al.* Late-onset Alzheimer's risk variants in memory decline,

incident mild cognitive impairment, and Alzheimer's disease. *Neurobiol. Aging* **36**, 60–7

(2015).

11.   Zhang, B. *et al.* Integrated systems approach identifies genetic nodes and networks in late-

onset Alzheimer's disease. *Cell* **153**, 707–20 (2013).

12.   Mostafavi, S. *et al.* A molecular network of the aging human brain provides insights into

the pathology and cognitive decline of Alzheimer's disease. *Nat. Neurosci.* **21**, 811–819

(2018).

13.   Conway, O. J. *et al.* ABI3 and PLCG2 missense variants as risk factors for

neurodegenerative diseases in Caucasians and African Americans. *Mol. Neurodegener.*

13, 53 (2018).

14.   Allen, M. *et al.* Conserved brain myelination networks are altered in Alzheimer's and

other neurodegenerative diseases. *Alzheimers. Dement.* **14**, 352–366 (2018).

15.   Allen, M. *et al.* Divergent brain gene expression patterns associate with distinct cell-

specific tau neuropathology traits in progressive supranuclear palsy. *Acta Neuropathol.*

136, 709–727 (2018).

16.   Bengio, Y., Courville, A. & Vincent, P. Representation learning: A review and new

perspectives. *IEEE Trans. Pattern Anal. Mach. Intell.* **35**, 1798–1828 (2013).

17.   Wolz, R., Aljabar, P., Hajnal, J. V. & Rueckert, D. Manifold Learning for Biomarker

Discovery in MR Imaging. in 116–123 (Springer, Berlin, Heidelberg, 2010).

doi:10.1007/978-3-642-15948-0_15

18.   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by

pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32**, (2014).

19.   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat.*

Methods **14**, 979–982 (2017).

20. Rosenberg, A. B. *et al.* Single-cell profiling of the developing mouse brain and spinal cord

with split-pool barcoding. *Science (80-.).* **360**, 176–182 (2018).

21. Mukherjee, S., Zhang, Y., Fan, J., Seelig, G. & Kannan, S. Scalable preprocessing for

sparse scRNA-seq data exploiting prior knowledge. *Bioinformatics* **34**, i124--i132 (2018).

22. Wan, Y.-W. *et al.* Meta-Analysis of the Alzheimer's Disease Human Brain Transcriptome

and Functional Dissection in Mouse Models. *Cell Rep.* **32**, 107908 (2020).

23. Zou, F. *et al.* Brain Expression Genome-Wide Association Study (eGWAS) Identifies

Human Disease-Associated Variants. *PLoS Genet.* **8**, e1002707 (2012).

24. Allen, M. *et al.* Human whole genome genotype and transcriptome data for Alzheimer's

and other neurodegenerative diseases. *Sci. Data* **3**, (2016).

25. Kunkle, B. W. *et al.* Genetic meta-analysis of diagnosed Alzheimer's disease identifies

new risk loci and implicates Aβ, tau, immunity and lipid processing. *Nat. Genet.* **51**, 414–

430 (2019).

26. Allen, M. *et al.* Novel late-onset Alzheimer disease loci variants associate with brain gene

expression. *Neurology* **79**, 221–8 (2012).

27. Allen, M. *et al.* Late-onset Alzheimer disease risk variants mark brain regulatory loci.

Neurol. Genet. **1**, e15 (2015).

28. Allen, M. *et al.* Association of MAPT haplotypes with Alzheimer's disease risk and

MAPT brain gene expression levels. *Alzheimers. Res. Ther.* **6**, 39 (2014).

29. Carrasquillo, M. M. *et al.* A candidate regulatory variant at the TREM gene cluster

associates with decreased Alzheimer's disease risk and increased TREML1 and TREM2

brain gene expression. *Alzheimers. Dement.* **13**, 663–673 (2017).

30. Sims, R. *et al.* Rare coding variants in PLCG2, ABI3, and TREM2 implicate microglial-

741   mediated innate immunity in Alzheimer's disease. *Nat. Genet.* **49**, 1373–1384 (2017).

742 31. Braak, H., Alafuzoff, I., Arzberger, T., Kretzschmar, H. & Del Tredici, K. Staging of

743   Alzheimer disease-associated neurofibrillary pathology using paraffin sections and

744   immunocytochemistry. *Acta Neuropathol.* **112**, 389–404 (2006).

745 32. Mirra, S. S. *et al.* The Consortium to Establish a Registry for Alzheimer's Disease

746   (CERAD). Part II. Standardization of the neuropathologic assessment of Alzheimer's

747   disease. *Neurology* **41**, 479–86 (1991).

748 33. Maaten, L. van der & Hinton, G. Visualizing data using t-SNE. *J. Mach. Learn. Res.* **9**,

749   2579--2605 (2008).

750 34. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and

751   Projection for Dimension Reduction. (2018).

752 35. Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis.

753   *Nature* **566**, 496–502 (2019).

754 36. De Strooper, B. & Karran, E. The Cellular Phase of Alzheimer's Disease. *Cell* **164**, 603–

755   615 (2016).

756 37. Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons,

757   and vascular cells of the cerebral cortex. *J. Neurosci.* **34**, 11929–47 (2014).

758 38. Chibnik, L. B. *et al.* Susceptibility to neurofibrillary tangles: role of the PTPRD locus and

759   limited pleiotropy with other neuropathologies. *Mol. Psychiatry* **23**, 1521 (2018).

760 39. Lambert, J.-C. *et al.* Meta-analysis of 74,046 individuals identifies 11 new susceptibility

761   loci for Alzheimer's disease. *Nat. Genet.* **45**, 1452 (2013).

762 40. Janelidze, S. *et al.* CSF biomarkers of neuroinflammation and cerebrovascular dysfunction

763   in early Alzheimer disease. *Neurology* **91**, e867--e877 (2018).

41.   Whitwell, J. L. *et al.* Neuroimaging correlates of pathologically defined subtypes of

Alzheimer's disease: a case-control study. *Lancet Neurol.* **11**, 868–877 (2012).

42.   Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment

analysis tool. *BMC Bioinformatics* **14**, 128 (2013).

43.   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene

Ontology Consortium. *Nat. Genet.* **25**, 25–9 (2000).

44.   Dong, A. *et al.* Heterogeneity of neuroanatomical patterns in prodromal Alzheimer's

disease: links to cognition, progression and biomarkers. *Brain* **140**, aww319 (2016).

45.   Wang, X. *et al.* Genetic determinants of disease progression in Alzheimer's disease. *J.*

Alzheimers. Dis. **43**, 649–55 (2015).

46.   De Jager, P. L. *et al.* A multi-omic atlas of the human frontal cortex for aging and

Alzheimer's disease research. *Sci. Data* **5**, 180142 (2018).

47.   Allen, M. *et al.* Human whole genome genotype and transcriptome data for Alzheimer's

and other neurodegenerative diseases. *Sci. data* **3**, 160089 (2016).

48.   Schneider, J. A., Arvanitakis, Z., Bang, W. & Bennett, D. A. Mixed brain pathologies

account for most dementia cases in community-dwelling older persons. *Neurology* **69**,

2197–2204 (2007).

49.   Allen, M. *et al.* Human whole genome genotype and transcriptome data for Alzheimer's

and other neurodegenerative diseases. *Sci. Data* **3**, 160089 (2016).

50.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21

(2013).

51.   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).

52.   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear

model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, (2014).

53.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing

and microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

54.   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat.*

Methods **14**, 979–982 (2017).

55.   Haghverdi, L., Büttner, M., Wolf, F. A., Buettner, F. & Theis, F. J. Diffusion pseudotime

robustly reconstructs lineage branching. *Nat. Methods* **13**, 845–848 (2016).

56.   A Bennett, D. *et al.* Overview and findings from the rush Memory and Aging Project.

Curr. Alzheimer Res. **9**, 646–663 (2012).

57.   Bennett, D. A. *et al.* Religious orders study and rush memory and aging project. *J.*

Alzheimer's Dis. 1–28 (2018).

58.   Ferretti, M. T. *et al.* Sex differences in Alzheimer disease—the gateway to precision

medicine. *Nat. Rev. Neurol.* 1 (2018).

59.   Deming, Y. *et al.* Sex-specific genetic predictors of Alzheimer's disease biomarkers. *Acta*

Neuropathol. **136**, 857–872 (2018).

60.   Mao, Q., Wang, L., Goodison, S. & Sun, Y. Dimensionality Reduction Via Graph

Structure Learning. in *Proceedings of the 21th ACM SIGKDD International Conference*

on Knowledge Discovery and Data Mining - KDD '15 765–774 (ACM Press, 2015).

doi:10.1145/2783258.2783309

61.   Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the visualization of

intersecting sets and their properties. *Bioinformatics* **33**, 2938–2940 (2017).

62.   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing

810 next-generation DNA sequencing data. *Genome Res.* **20**, 1297–303 (2010).

811 63. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

812 transform. *Bioinformatics* **25**, 1754–60 (2009).

813 64. Kang, H. M. *et al.* Variance component model to account for sample structure in genome-

814 wide association studies. *Nat. Genet.* **42**, 348–354 (2010).

Acknowledgements

This work was supported by NIA grants U54AG054345 and RF1AG057443. The ROSMAP

Study data were provided by the Rush Alzheimer's Disease Center, Rush University Medical

Center, Chicago. Data collection was supported through funding by NIA grants P30AG10161,

R01AG15819, R01AG17917, R01AG30146, R01AG36836, U01AG32984, U01AG46152, the

Illinois Department of Public Health, and the Translational Genomics Research Institute. Mayo

RNAseq Study data were provided by the following sources: The Mayo Clinic Alzheimer's

Disease Genetic Studies, led by Dr. Nilufer Ertekin-Taner and Dr. Steven G. Younkin, Mayo

Clinic, Jacksonville, FL using samples from the Mayo Clinic Study of Aging, the Mayo Clinic

Alzheimer's Disease Research Center, and the Mayo Clinic Brain Bank. Data collection was

supported through funding by NIA grants P50 AG016574, R01 AG032990, U01 AG046139,

R01 AG018023, U01 AG006576, U01 AG006786, R01 AG025711, R01 AG017216, R01

AG003949, NINDS grant R01 NS080820, CurePSP Foundation, and support from Mayo

Foundation. Study data includes samples collected through the Sun Health Research Institute

Brain and Body Donation Program of Sun City, Arizona. The Brain and Body Donation Program

is supported by the National Institute of Neurological Disorders and Stroke (U24 NS072026

National Brain and Tissue Resource for Parkinson's Disease and Related Disorders), the

National Institute on Aging (P30 AG19610 Arizona Alzheimer's Disease Core Center), the

Arizona Department of Health Services (contract 211002, Arizona Alzheimer's Research

Center), the Arizona Biomedical Research Commission (contracts 4001, 0011, 05-901 and 1001

to the Arizona Parkinson's Disease Consortium) and the Michael J. Fox Foundation for

Parkinson's Research. MSBB data were generated from postmortem brain tissue collected

through the Mount Sinai VA Medical Center Brain Bank and were provided by Dr. Eric Schadt

from Mount Sinai School of Medicine.

Author Contributions

S.M. and B.A.L. designed the study. S.M., L.H., and B.A.L. performed the analyses. S.M.,

L.H., C.P., S.J., G.G., A.K.G., S.K.S., P.L.D.J., N.E.T., G.W.C., L.M.M., and B.A.L. wrote the

manuscript.

Tables

Table 1 - Pathway enrichments for branch specific differentially expressed genes in TCX.

Differential expressed genes are identified with a two sided Tukey's honest significant difference

test (FDR < 0.05), with Branch 1 as the reference, and pathway enrichments that are significant

from a one-sided Fisher's exact test are shown (FDR <0.05).

Direction	Branch	Representative Enriched Gene Ontology Terms
Down	2	prespliceosome (GO:0071010), mitochondrial electron transport, cytochrome c to oxygen (GO: 0006123)
Down	3	negative regulation of microtubule polymerization or depolymerization (GO:0031111)
Down	4	mitochondrial electron transport, NADH to ubiquinone (GO: 0006120), spliceosomal tri-snRNP complex (GO:0097526), negative regulation of microtubule depolymerization (GO:0007026)
Down	5	axon (GO:0030424), protein kinase C activity (GO:0004697),
Down	6	gamma-tubulin large complex (GO:0000931), U1 snRNP (GO:0005685), mitochondrial respiratory chain complex IV (GO:0005751), response to cadmium ion (GO:0046686)
Up	3	fatty acid elongase activity (GO:0009922), ubiquitin protein ligase activity (GO:0061630)
Up	4	transforming growth factor beta-activated receptor activity (GO:0005024), hippo signaling (GO:0035329), regulation of extrinsic apoptotic signaling pathway via death domain receptors (GO: 1902041), regulation of DNA repair (GO: 0006282)
Up	5	regulation of apoptotic process (GO:0042981), leptin mediated signaling pathway (GO:0033210), negative regulation of hippo signaling (GO:0035331), small GTPase binding (GO:0031267)
Up	6	extracellular ligand-gated ion channel activity (GO:0005230), integral component of mitochondrial inner membrane (GO:0031305)

853 **Table 2 -** Pathway enrichments for branch specific differentially expressed genes in DLPFC.

854 Differential expressed genes are identified with a two-sided Tukey's honest significant

855 difference test (FDR < 0.05), with Branch 1 as the reference, and pathway enrichments that are

856 significant from a one-sided Fisher's exact test are shown (FDR <0.05).

Direction	Branch	Representative Enriched Gene Ontology Terms
Down	2	DNA repair (GO:0006281), intracellular protein transport (GO:0006886)
Down	3	mismatch repair complex binding (GO:0032404)
Down	5	mitochondrial respiratory chain complex assembly (GO: 0033108)
Up	2	racemase and epimerase activity (GO: 0016857)
Up	3	racemase and epimerase activity (GO: 0016857)
Up	4	vesicle mediated transport (GO: 0016192)
Up	5	NuRD complex (GO: 0016581)
Up	6	microtuble motor activity (GO:0003777), AP-2 adaptor complex binding (GO:0035612)

Figures

Figure 1 - Overview of manifold learning for unraveling staging in Alzheimer's disease. A)

Illustration of steps in manifold learning using reverse graph embedding DDRTree method. B)

Illustration of lineage inference process for LOAD. RNA-seq samples with different disease

diagnoses were pooled, batch normalized, and a smooth manifold was learned for each brain

region across individuals (each point is an individual). Total sample numbers are indicated across

Mayo RNAseq TCX and ROS/MAP DLPFC for the different diagnoses in parentheses.

Figure 2 - Manifold learning accurately infers disease states and stages from RNA-seq samples.

A) Estimated disease progression trees from temporal cortex (TCX) and B) dorsolateral

prefrontal cortex (DLPFC) brain regions showing localization of identified LOAD samples on

particular branches. C) Distribution of pseudotime for AD cases and controls for both DLPFC

and TCX for 218 independent samples from two independent studies. D) Distribution of

expression correlation with pseudotime for both LOAD GWAS genes and non-LOAD GWAS

genes for 17446 genes from two independent studies. Box plots have lower and upper hinges at

the 25th and 75th percentiles and whiskers extending to at most 1.5xIQR (interquartile range).

Figure 3 - Manifold learning replicates existing measures of staging in LOAD in DLPFC

samples. A) Samples colored by 3 different external measures of LOAD staging namely: Braak

Score (tau pathology), CERAD Score (amyloid pathology) and Cognitive Diagnosis (Clinical

measure of disease severity). Black lines denote inferred lineages. B) Distribution of samples by

inferred stage for different distinct stages in each of the three methods of measuring LOAD

severity for 338 independent samples from one study. Inferred disease stages generally

corresponded with all methods, and Cognitive diagnosis demonstrated the strongest alignment.

Box plots have lower and upper hinges at the 25th and 75th percentiles and whiskers extending

to at most 1.5xIQR (interquartile range).

Figure 4 – Cell type gene expression signatures as a function of disease pseudotime. A) Mean

expression of cell markers for astrocytes, neurons, microglia, and oligodendrocytes as a function

of pseudotime for TCX brain region, B) mean expression of cell markers for astrocytes, neurons,

microglia, oligodendrocytes as a function of pseudotime for DLPFC (B) brain region.

Figure 5 – Disease resistant state. A) The inferred manifold from the TCX region with samples

colored by their inferred disease subtype/state. State 5 (dots, circled) lies at the late end of the

disease trajectory, indicating a strong disease-like transcriptomic phenotype, yet most samples in

the group did not have pathologically diagnosed AD (Figure 2A). We hypothesize this group

represents a disease resistant state to the disease. B) Biclustering results of average expression

from each disease state, with increased expression of a gene cluster (Cluster 4) unique to State 5.

A

Noisy high dimensional data

Learn patient level reduced dimension latent variable z, and projection W

Reduced dimensional manifold

Jointly learn spanning tree of z centroids.

Estimated smooth manifold

Iterate learning until convergence.

B

Individuals

Bulk RNA-Seq (DLPFC and TCX brain regions)

Quantify expression

Individuals

genes

Manifold Learning

Learned Disease Manifold

AD (211)

Control (150)

Other (418)

Test for associations with:
1. Alzheimer's disease
2. Neuropathology
3. Genetic variants
4. Cell type markers

Direction of disease progression