Molecular estimation of neurodegeneration pseudotime in older brains

Sumit Mukherjee[1,2*], Laura Heath[1*], Christoph Preuss[3], Suman Jayadev[4], Gwenn A. Garden[4], Anna K Greenwood[1], Solveig K Sieberts[1], Phillip L De Jager[5,6], Nilufer Ertekin-Taner[7,8], Gregory W Carter[3], Lara M Mangravite[1], Benjamin A Logsdon[1,9]

[1]Sage Bionetworks, Seattle, WA, USA.
[2]Microsoft, Redmond, WA, USA[†].
[3]The Jackson Laboratory, Bar Harbor, ME, USA.
[4]Department of Neurology, University of Washington, Seattle, WA, USA.
[5]Center for Translational & Computational Neuroimmunology, Department of Neurology, Columbia University Irving Medical Center, New York City, USA
[6]Taub Institute, Columbia University Irving Medical Center, New York City, USA
[7]Mayo Clinic Florida, Department of Neurology, Jacksonville, FL, USA.
[8]Mayo Clinic Florida, Department of Neuroscience, Jacksonville, FL, USA.

[†]Current Address.
[*]These authors contributed equally to this work.

[9]Correspondence:
Benjamin A Logsdon, PhD
Sage Bionetworks
2901 3[rd] Avenue Suite 330
Seattle, WA 98121
ben.logsdon@sagebionetworks.org

Abstract

Therapeutic treatments for late-onset Alzheimer's disease (LOAD) are hindered by an incomplete

understanding of the temporal molecular changes that lead to disease onset and progression. Here, we

evaluate the ability of manifold learning to develop a molecular model for the unobserved temporal

disease progression from RNA-Seq data collected from human postmortem brain samples collected

within the ROS/MAP and Mayo Clinic RNA-Seq studies of the AMP-AD consortium. This approach

defines a cross-sectional ordering across samples based on their relative similarity in RNA-Seq profiles

and uses this information to define an estimate of molecular disease stage – or disease progression

pseudotime - for each sample. This transcriptional estimate of disease progression is strongly concordant

with burden of tau pathology (Braak score, $P = 1.0\text{x}10^{-5}$), amyloid pathology (CERAD score, $P = 1.8\text{x}10^{-5}$), and cognitive diagnosis ($P = 3.5\text{x}10^{-7}$) of LOAD. Further, the disease progression estimate

recapitulates known changes in cell type abundance and impact of genes that harbor known AD risk loci.

Samples estimated to reside early in disease progression were enriched for control and early stage AD

cases, and demonstrated changes in basic cellular functions. Samples estimated to reside late in disease

progression were enriched for late-stage AD cases, and demonstrated changes in known disease processes

including neuroinflammation and amyloid pathology. We also identified a set of control samples with

late-stage estimated disease progression who also showed compensatory changes in genes involved in

affected pathways are protein trafficking, splicing, regulation of apoptosis, and prevention of amyloid

cleavage. In summary, we present a disease specific method for ordering patients based on their LOAD

disease progression from brain transcriptomic data.

Introduction

Late-onset Alzheimer's disease (LOAD) is a devastating illness with no effective disease modifying

therapy, owing to a 99.6% failure of clinical trials[1]. There is a growing consensus that the most effective

treatments will intervene early in disease progression and halt disease pathophysiological processes prior

to conversion to LOAD[2]. In addition, there is increasing recognition that LOAD may in fact be a

spectrum of related diseases that have similar clinical and neuropathological manifestations[3,4]. Devising

successful therapeutic strategies will likely require targeting potentially diverse early-stage disease

processes that occur prior to a high burden of neuropathology or cognitive impairment.

Current approaches to identify AD affected individuals include *in vivo* measures of the pathological

hallmarks of disease – amyloid, tau, and neurodegeneration – via CSF biomarkers for amyloid and tau[5],

positron emission tomography for amyloid and tau (PET)[6], and structural and functional MRI of

neurodegeneration. Cognitive assessments are used to estimate disease burden[7], although measurable

cognitive impairment generally indicates a sustained burden of neuropathology and advanced

neurodegeneration. Based on biomarker studies of AD, by the time cognitive decline becomes detectable,

neuropathological changes of AD have already occurred, first in Aß and subsequently in tau related

measures[8] and therefore cannot be used to select patients for early disease stage studies. Furthermore,

while these measures of disease progression capture the overall increase in burden of pathology and

cognitive decline, they do not necessarily identify the dysfunctional molecular mechanisms that lead to

neuropathology and cognitive decline. There are likely many independent patient specific molecular

pathways present at an early stage in disease that then contribute to later stage disease progression[9,10].

This motivates the need to identify these early stage molecular mechanisms driving disease progression.

The Accelerating Medicines Project for Alzheimer's Disease (AMP-AD) consortia have generated

genome-wide transcriptomics of post-mortem brain tissue from patients across a broad range of

Alzheimer's disease neuropathological progression – including individuals with various stages of AD

neuropathology and those who lack AD neuropathology, but who may in face harbor early stage disease

molecular processes. We therefore sought to chart the molecular progression of the disease as reflected in

the aggregate behavior of the CNS transcriptome across these individuals. While standard approaches

such as differential expression or coexpression analyses have proven informative[11–15], these analyses do

not infer the relative stage of disease progression or identify distinct disease subtypes. Here we propose

an approach to analyze population level RNA-seq data from post-mortem brain tissue to learn a tree

structured progression (**Figure 1**) that represents distinct sub-types of disease and the relative progression

of disease across patients. With this approach, we identify potentially generalizable trajectories of LOAD

across heterogeneous patient populations at all stages of disease. Furthermore, we characterize molecular

pathways that define disease stages – a potential source of new biomarkers and therapeutic interventions

for early-stage disease processes along multiple different disease trajectories.

To learn the molecular disease staging and neuropathologic progression tree we use a manifold learning

method[16]. Manifold learning refers to a group of algorithms that aim to recover the low dimensional

subspace underlying a high dimensional dataset. Previous authors use manifold learning to estimate

disease progression from neuroimaging data[17] and to study lineage commitment of cells during

differentiation from single cell RNA-seq (scRNA-seq)[18–21]. To our knowledge, manifold learning has not

been used to estimate disease progression and/or disease stages from bulk RNA-seq data derived from

post-mortem tissue. Henceforth, we refer to manifold learning, lineage inference interchangeably in

reference to the construction of the inference of a neuropathologic progression tree. We demonstrate that

these tools can estimate the disease staging and progression tree (**Figure 2**) from bulk RNA-Seq data

collected from post-mortem brain tissues in a case/control cohort. Moreover, these trees show clear

LOAD staging, enable the study of cell type specific effects of LOAD, and allow the identification of

genetic factors driving disease progression.

Results

Manifold learning distinguishes pathologically defined LOAD from control

We first quantify the bulk RNA-Seq data from the ROS/MAP and Mayo Clinic cohorts into gene counts

and remove any batch effects introduced due to sequencing runs using standard count normalization (see

Methods). The data from the ROS/MAP cohort is sampled from the dorsolateral prefrontal cortex

(DLPFC), and the data from the Mayo Clinic cohort is sampled from the temporal cortex (TCX). Patient

clinical characteristics are reported in **Table S1** and described in Methods. The full pipeline we use for

RNA-Seq data generation and quality control was recently reported[22]. The entire transcriptome comprises

many genes which do not have measurable expression or vary across case/control samples, which we

remove in order to reduce the noise in manifold learning[19]. To do this, we first perform differential

expression analysis between case/control samples separately for each study and retain genes that reach an

FDR of 0.10. To test if this biased the disease lineage inference, we also perform manifold learning using

only genes with high variance across samples, and we see a strong concordance with disease lineages

inferred with differentially expressed genes (**Figure S1**). Changing the significance threshold for the

differential expression analysis to FDR < 0.01 did not materially change these results (**Figure S2**). We

infer the disease lineage for each brain region on this subset of retained genes (**Figure 2A-B**). Adjusting

for post-mortem interval (PMI) (**Figure S3A, S4A**), 10 principal components from a principal component

analysis of genotype data to account for ancestry effects (**Figure S3B, S4B**), RNA integrity number

(RIN) (**Figure S3C, S4C**), or all of these variables (**Figure S3D, S4D**) did not materially change the

overarching ordering of patients for either the TCX or DLPFC regions. Furthermore, to assess the general

robustness of the results we apply leave one out cross validation to infer disease pseudotime for both

DLFPC and TCX brain regions and find strong correlations between lineages inferred with each sample

removed, and the lineage for the entire sample set (**Figure S5**).

We visualize the clinical diagnosis of the samples on the inferred disease staging tree to verify that there

is indeed separation of AD patients across the tree. To determine if inferred tree structure is an accurate

model of disease progression, we introduce the notion of disease pseudotime which is the geodesic

distance along the tree from an inferred initial point to the point of interest as a quantitative linear

measure of LOAD stage. We scale this estimated disease pseudotime to lie in the range $[0,1]$ to make the

effects comparable between the two studies (and brain regions). We show that for LOAD cases compared

to controls there is a significant association ($P = 0.02$ in Mayo and $P = 2.0 \times 10^{-6}$ in ROS/MAP, logistic

regression) between the estimated pseudotime and AD case/control status (**Figure 2C**). These effects are

not abrogated by adjusting for RIN, PMI, or ancestry in either tissue (**Figure S6A-D, Table S2**). To

assess whether the association between inferred disease pseudotime is a phenomena in only the Mayo

RNAseq and ROS/MAP RNA-seq data, we also apply the lineage inference approach to expression array

data from the Mayo eQTL study[23] (**Methods**). These samples are derived from a completely independent

set of donors than the Mayo RNA-seq study[24]. Similarly, we restrict to only female samples, and test for

an association between inferred disease pseudotime and disease status (**Figure S7A-B**). We see a

significant association between disease pseudotime and neuropathological AD diagnosis ($P = 2.2 \times 10^{-8}$).

Furthermore, we observe strong evidence of sex heterogeneity when performing the manifold learning

approach, and find that the manifolds inferred for female only samples show stronger association with

pseudotime than for male samples. This matches previous observations concerning disease specific sex

heterogeneity[22]. As such, we do not see as statistically significant of an association between pseudotime

and disease diagnosis in male samples ($P = 0.040$ in Mayo and $P = 0.11$ in ROS/MAP, logistic regression,

Figure S8A-D). Similarly, the association between pseudotime and amyloid, tau, and cognitive diagnosis

is attenuated in male samples in ROS/MAP (**Figure S9A-D**). For the combined samples we see moderate

evidence of disease association with pseudotime ($P=0.003$ in Mayo and $P=0.003$ in ROS/MAP, logistic

regression, **Figure S10A-D**). The association with neuropathological measures of disease is more robust

in the combined sample (**Figure S11A-F**), but not as strong as in females only, hence we restrict to

female only analyses for all subsequent reported results.

We test whether genes in loci that have been implicated in genome wide association studies of LOAD are

associated with inferred disease pseudotime. We use the prioritized LOAD GWAS genes[25], **Table S3**,

and compute the correlation between their expression and inferred pseudotime (**Figure 2D**). When

compared to the background of all genes, we see that there is a significant increase in positive correlation

with disease pseudotime for implicated LOAD GWAS genes (P-value: 7.3×10^{-5} in Mayo and 5.6×10^{-3} in

ROS/MAP). This effect is robust to adjustments for PMI, RIN, or ancestry (**Figure S12A-D**).

Furthermore, this does not appear to be driven by a small subset of outlier genes, but by the majority of

the distribution of LOAD GWAS genes. The fact that AD GWAS loci genes have expression

associations with pseudotime likely implies that the AD risk variants at these are also eQTL as previously

shown[26–29] and/or are members of co-expression networks that are differentially expressed in AD[13,30].

To further explore the relationship between inferred disease stage and LOAD, we test for its association

with neuropathological and clinical measures of LOAD severity, namely: i) Braak score, ii) CERAD

score, and iii) cognitive diagnosis. The ROS/MAP study has numeric scores for these categories available

as covariates for each sample. Braak is a semi-quantitative measure that increases with tau pathology[31]

and CERAD is a semi-quantitative measure of density of neuritic plaques[32]. We overlay these scores on

the inferred manifold for the DLPFC brain region (**Figure 3A**). We observe a progressive increase in tau,

amyloid, and cognitive burden as we traverse the inferred disease manifold (**Figure 3A**). This is further

quantified by characterizing the relationship between branches of the inferred manifold and Braak,

CERAD, and cognitive diagnosis (**Figure 3B**). We observe significant associations between pseudotime

and Braak score ($P=1.0 \times 10^{-5}$), CERAD score ($P=1.8 \times 10^{-5}$), and cognitive diagnosis ($P=3.5 \times 10^{-7}$). In

ROS/MAP, adjustment for Braak score when fitting the DDRTree method attenuates the association

between pseudotime and disease states (P-value: 0.214, **Figure S13A-B**), though there is still evidence of

association with cognitive diagnosis (P-value: 0.03, **Figure S13C-D**). In the Mayo RNA-seq study we

have Braak score and Thal Amyloid scores for only a subset of samples, but observe a similar pattern as

in ROS/MAP (**Figure S14A-B**) for the samples that we do have data. There is a significant association

between Braak score and pseudotime (P-value: 5×10^{-5}) as well as Thal amyloid (P-value: 1.7×10^{-5}) within

this subset with available neuropathology data.

Comparison to other unsupervised learning approaches. We compare the manifold learning approach to

other unsupervised learning approaches including principal component analysis (PCA), t-distributed

Stochastic Neighbor Embedding (tSNE)[33], and Uniform Manifold Approximation and Projection

(UMAP)[34]. Correlations between the first two dimensions of each of these approaches and the DDRTree

learned pseudotimes are shown in **Figures S15A-F, S16A-F** for DLPFC and TCX brain tissues

respectively. We see the strongest correlations between PCA1 and UMAP2 and pseudotime in both data-

sets, increasing our confidence that the overarching ordering of patients along a disease pseudotime is a

robust characteristic of the disease progression as reflected in the gene expression changes as a function

of disease, and not dependent on the underlying manifold learning approach. This is further supported by

inspecting the association between these approaches and Braak, CERAD, and cognitive diagnosis (cogdx)

scores for the DLPFC tissue (**Table S4**). Furthermore, the manifold learning approaches (DDRTree and

UMAP) have much stronger associations with Braak, CERAD, and cogdx scores than either PCA or

tSNE. In fact, UMAP has been proposed for lineage inference[35], and when we apply UMAP with lineage

inference using Monocle3, we observe similar results (**Figure S17-S18**) as with DDRTree and Monocle2

(**Figure 2-3**), though the inferred pseudotimes from Monocle3 are not quite as significant as the

association with UMAP2 or from Monocle 2 with Braak, CERAD, and cogdx (**Table S4**).

Inferred staging recapitulates known biology of AD

To demonstrate that the inferred disease pseudotime recapitulates known biology of LOAD, we test for

association between inferred disease stage and both the cellular response to disease and the genetics of the

disease. A prominent hypothesis in AD is that the effects of the disease vary across different brain cell

types, specifically neurons and glial subtypes. Current understanding of the cell biology of the disease

implicates progressive neuronal loss and increase in gliosis[36]. To test if the inferred pseudotime aligns

with existing cell type specific hypotheses regarding AD, we first selected from the genes used in lineage

construction the marker genes for four key cell types: neurons, astrocytes, microglia, and

oligodendrocytes based on a previously published brain cell atlas[37] (**Table S5**). We then calculate the

normalized mean expression for the marker genes of each cell type and fitted a linear model to the mean

expression with disease pseudotime as the dependent variable. We find that, in both studies, the cell

specific marker gene levels show a statistically significant linear dependence on pseudotime (**Table S6**).

Fitted effects recapitulate known neuropathologic changes which occur in AD, namely: i) a reduction in

the neuronal populations as AD progresses, and ii) an increase in expression associated with activation of

microglia, astrocytes, and oligodendrocytes as AD progresses (**Figure 4**).

Next, we test for association between assigned lineage state in ROS/MAP (DLPFC) and Mayo (TCX) and

APOE e4 status (**Figure S19**). For reference, the inferred trees for TCX and DLPFC each resolve into 6

branches (**Figure 5A,S20**). Carriers of the APOE e4 allele are significantly enriched on the State 4

branch in TCX (P-value = 0.027, unadjusted), and suggestively enriched on the State 5 branch (P-value =

0.06, unadjusted), compared to the State 1 branch (logistic regression). Similarly, in the Mayo eGWAS

study, when we perform an ordinal logistic regression of APOE e4 dosage and disease pseudotime we see

a significant positive association as a function of pseudotime (P-value = 6.9×10^{-4}, **Figure S7C**).

Genetic factors associated with inferred disease staging

Lineage inference of LOAD transcriptomes provides a quantitative measure of disease progression for

genetic associate testing, and the significantly greater correlation between pseudotime and gene

expression for known LOAD risk genes (**Figure 2D**) suggests that the observed differences in disease

trajectories are influenced by genetic factors. To test this hypothesis, we perform single variant analysis

using whole-genome sequencing data for 305 patients from the ROS/MAP and 131 patients from the

Mayo cohort. Despite the limited sample size, resulting in lack of statistical power to discover genome-

wide significant associations, multiple variants reach a genome-wide suggestive threshold of $p < 1 \times 10^{-5}$

(**Table S7**). We do not see evidence of population stratification in the analysis (**Figure S21-S22**).

Notably, the most significant association with pseudotime for the ROS/MAP cohort is observed at the

PTPRD locus (rs7870388, p = 1.31×10^{-6}) (**Figure S23, Table S7**). The *PTRPD* locus is associated with

the susceptibility to neurofibrillary tangle independent of amyloid deposition in the ROS/MAP cohort[38].

For the Mayo Clinic cohort, known LOAD variants in the *APOE* (rs6857, p = 9.18×10^{-6}) and *BIN1*

(rs62158731, p= 4.68 x 10^{-5}) loci overlap with variants associated with inferred disease stage (**Figure**

S23, Table S7)[39]. When comparing our association results for inferred disease stage with summary

statistics from a large-scale case-control approach, we identify multiple variants which have been

previously associated with LOAD in the IGAP cohort (**Table S8**). Furthermore, we identify several

potential novel candidate genes associated with inferred disease stage (*ADAMTS14*, *IL7*, *MAN2B1*) linked

to immune and lysosomal storage function (**Figure S23**,**Table S7**). *IL-7* has been proposed as an

inflammatory biomarker for LOAD that correlates with disease outcome and severity[40]. *ADAMTS14* is

part of a locus that has been previously linked with Alzheimer susceptibility and plays an important role

in the regulation of immune function via TGF-beta signaling.

New disease insights identified from inferred disease lineages

Another important direction of study in the field of Alzheimer's is the identification of disease subtypes,

which has so far predominantly been done using imaging data[41]. The branches of the inferred disease trees

provide a new transcriptomic-based approach to identify disease subtypes. In both brain regions and in

two separate cohorts, there were two distinct early-lineage branches corresponding to predominantly

control samples, which we interpret as different initial paths towards the disease. Similarly, both brain

regions feature several distinct branches with predominantly LOAD samples (**Figure 2A-B**).

Branch-specific differential expression patterns. To study the genes and pathways specific to each branch,

we perform a branch-specific differential expression analysis with an ANOVA model using the branches

with the highest proportion of controls as the reference branch for DLPFC (**Table S9**) and TCX (**Table**

S10). We see many genes are differentially expressed between the control branch and branches that are

enriched in the affected individuals (**Table S11**). We test for overlap between the differentially expressed

gene sets between the two studies (**Figure S24**), and find significant overlaps in branches enriched for

late stage disease cases, especially between up-regulated genes in State 6 of DLPFC and up regulated

genes in State 5 of TCX (P-value: 4.1×10^{-108}, OR: 4.5, Fisher's exact test), as well as genes that are up-

regulated in State 6 of DLPFC and in State 4 of TCX (P-value: 1.8×10^{-14}, OR: 1.9, Fisher's exact test),

and more modestly for genes that are down regulated in State 6 of DLPFC and down regulated in State 3

of TCX (P-value: 1.1×10^{-6}, OR: 1.6, Fisher's exact test).   Next, we performed an enrichment analysis on

each of these differentially expressed gene sets with the enrichR[42] package for Gene Ontology[43]

annotations (**Methods**). The results of this enrichment analysis for DLPFC and TCX tissues (**Table S12-**

S13). Only gene sets with significant enrichment are shown (FDR adjusted p-value < 0.05). Overall, we

see a pattern of loss of expression of basic cell biology mechanisms in early-stage branches including

RNA splicing, mitochondria function, protein transport, and DNA repair. Late-stage branches were

characterized by increased immune response (e.g. TGFb/WNT signaling) and apoptotic activity (**Table**

1).

While studying the different branches in the two brain regions, we observe a branch (branch 5) that

corresponds to a group of predominantly neuropathological control samples from the Mayo RNA-seq

cohort that were in close proximity to a branch with predominantly LOAD samples (branch 4) on the

inferred disease lineage (**Figure 5A**). However, most of the samples on branch 5 are neuropathological

controls as defined by the Mayo diagnostic criteria. We bi-cluster the mean expression of genes in each

branch and the branches themselves (**Methods**). This clustering analysis (**Figure 5B**) shows that the

closest branch to this potentially disease resistant branch contains the highest proportion of AD samples.

While the stage proximity implies some transcriptomic similarity between these controls and nearby

cases, we also see a secondary cluster of genes with increased expression in the resistant state while

having reduced expression in all other states. We perform an enrichment analysis on this set of genes and

find significant GO terms corresponding to: protein transport (GO:0015031), regulation of mRNA

splicing, via spliceosome (GO:0048024), negative regulation of apoptotic process (GO:0043066), and

regulation of amyloid-beta clearance (GO:1900221) (Cluster4, **Table S14**). It is possible that these

potentially disease resistant individuals have compensatory mechanisms which suppress the hallmarks of

disease despite sharing gene expression patterns with pathologically affected individuals.

To replicate this observation, we perform a differential expression analysis on individuals in the Mayo

eGWAS study where we consider individuals that are in the top quintile of pseudotime, but are classified

as neuropathological controls as resistant individuals (**Figure S7D**, **Methods**). To test if these individuals

also have a similar resistant molecular endophenotype, we compare the overlap between various

differential expressed gene sets derived from these resistant individuals and the gene sets identified in the

biclustering of the Mayo RNA-seq data (**Figure S25**). We observe that there is a highly statistically

significant overlap between genes that are upregulated in these Mayo eGWAS resistant individuals (P-

value: 2.6×10^{-51}, OR: 2.9, Fisher's exact test), and the Cluster 4 genes that are upregulated in the branch 5

Mayo RNA-seq samples (**Figure S25**).

Discussion

Here we proposed a novel approach to infer the Alzheimer's disease severity and disease subtypes in an

unsupervised manner from post-mortem bulk RNA-seq data that gets directly at the challenge of

identifying the temporal progression of disease in the disease resistant tissue. Our strategy utilized a

manifold learning approach to infer a disease progression tree from cross-sectionally collected patient

samples from two different brain regions. The underlying assumption of our approach is that the inferred

disease progression from cross sectional samples serves as a proxy for the unobserved progression of the

disease across subtypes of LOAD. We validated this hypothesis through comparisons with

neuropathological measures of disease stage severity and against known cell type specific effects caused

by the disease. While one could argue that the method is merely classifying patients as either disease

cases or controls based on expression signatures of the hallmarks of disease, we see at least three

advantages of this approach beyond that interpretation. First the application of this method appears to be

produce a more quantitative measure of disease state than strictly neuropathological assessments - as born

out through the identification of novel and distinct genetic loci that replicate based on IGAP summary

statistics. This suggests that it may be adding information related to other aspects of disease such as the

effect of neuroinflammation or neuronal injury. In addition we see evidence of neuropathological controls

that are disease resistant given their molecular state in two independent studies – which would not be

detectable with standard neuropathological or clinical assessments – and could provide important

molecular clues to mechanisms of disease resistance. Finally, there is the potential that specific pathways

associated with early stage disease processes can be characterized which is desperately needed for

hypothesis generation in the field. Furthermore, this approach provides clues to better understanding the

molecular heterogeneity of disease by identifying specific pathways that are dysregulated in subsets of

patients at different disease stages. This opens up the possibility of better patient stratification and

precision medicine.

We observed that different biological processes vary as a function of inferred disease stage, and that

early-stage disease processes potentially include RNA-splicing, mitochondrial function, and protein

transport – implicating multiple basic cell biology mechanisms as potential early stage disease processes

for further study in relevant model systems. Additionally, the manifold learning method identified 6

potential subtypes of LOAD from RNA-seq (i.e. branches) suggesting the LOAD populations should be

stratified by better biomarkers with tailored treatment strategies. To identify and test these stratifications

future studies should focus on longitudinal cohorts of patients with rich molecular and imaging data to be

able to identify biomarkers that can accurately and precisely stratify patients into the underlying

molecular subtypes in terms of the molecular characteristics of their transcriptome and different relative

stages of disease. Furthermore, we observe a potential disease resistant sub-type of patients. This disease

resistance should be tested in disease model systems, to identify if neuropathological readouts can be

modified by altering the function of the pathways identified in our analysis (e.g. APP processing, RNA

splicing, apoptosis, protein trafficking). While this preliminary observation needs to be validated in

another cohort, it has the potential to be a novel source of hypotheses concerning new therapeutic

development. Specifically, for constructing better combination therapy hypotheses that may confer

neuroprotection, even in patients that are mildly affected by disease.

LOAD is a complex and heterogeneous disease encompassing a broad spectrum of clinical symptoms.

Disease progression can vary widely between patients leading to different rates of cognitive decline.

Several lines of evidence suggest that these differences in progression are modified by multiple genetic

factors affecting the transition from one pathological state to another[44,45]. However, it has remained

difficult to assess the role of genetic variants affecting disease trajectories by case-control approaches

alone. Here, we showed that our novel expression trait pseudotime might be used as a molecular

phenotype to identify known and novel AD loci associated with different disease progression states across

AD patients. Despite a limited sample size, we identified previously associated AD candidate loci in the

ROSMAP (*PTPRD*) and Mayo (*BIN1, APOE*) cohorts with suggestive significance ($p < 1 \times 10^{-5}$).

Variants in *PTPRPD* have been associated with the susceptibility to neurofibrillary tangles, independent

of amyloid burden. This is in line with the results from the differential gene expression analysis of

pseudotime branches showing an enrichment of molecular pathways implicated in TAU pathology.

Furthermore, our analysis revealed several novel loci linked to immune function (*ADAMTS14*, *IL7*) and

neurotransmitter signaling (*CHRM2*, *CHRM3*) processes associated with disease pseudotime (**Table S4**).

Future studies will be needed to replicate these findings in independent cohorts of LOAD and validate the

role of candidate genes in LOAD related disease progression by first identifying peripheral biomarkers

that correspond to this molecular definition of disease stage, and then testing for GWAS association with

that disease stage. Subsequent results can improve functional interpretation by linking candidate genes

with ordered pathological processes.

Methods

ROS/MAP and Mayo RNAseq study population characteristics

Detailed descriptions of cohort and patient characteristics included in this study can be found in

previously published work[46,47]. Patient characteristics included in this study are summarized in **Table S1**,

stratified by sex. In brief: for Mayo samples, AD diagnosis was performed according to NINCDS-

ADRDA criteria (probable or possible AD); control individuals had Braak NFT stage ≤ 3, CERAD score

< 2.5, and lacked other pathologic diagnoses; Path.Aging are individuals who lacked any pathologic

diagnoses and had Braak NFT ≤ 3 and CERAD score ≥ 2. Progressive supranuclear palsy (PSP)

individuals were diagnosed neuropathologically by a single neuropathologist. (For further details, see

Allen et al 2016). For ROSMAP samples, AD diagnosis was according to NIA Reagan criteria, which

combines neuropathology and clinical data; Control individuals had no signs of cognitive impairment;

and Other individuals had MCI, mixed pathology, or other form of dementia. Age at death was collected

for all patients in the ROSMAP study, though age at first AD diagnosis had a high degree of missingness

and thus was not used as a variable in follow-up analyses (see **Table S1**). Braak stage indicates the

measure of severity of NFT pathology. Stages I and II indicate NFTs confined mainly to the entorhinal

region of the brain, stages III and IV indicate involvement of limbic regions, and V and VI indicate

moderate to severe neocortical involvement. CERAD score is a semiquantitative measure of neuritic

plaques. 1=definite AD; 2=probable AD; 3=possible AD; 4=No AD. Cognitive diagnostic category (as

determined by neurologist): 1) NCI; 2) MCI and no other cause of CI; 3) MCI and another cause of CI; 4)

AD and no other cause of CI; 5) AD and another cause of CI; 6) Other primary cause of dementia[48]. Thal

amyloid stages: phases of amyloid deposition. 0) no amyloid; 1) isocortical phase; 2) limbic phase; 3)

basal ganglia phase; 4) basal forebrain and midbrain phase; 5) pons/medulla oblongata and cerebellum

phase. The list of differentially expressed genes used to create the Monocle objects were based on results

which included the whole data set.

RNA sequencing

The details of the sample collections, postmortem sample characteristics, the tissue and RNA

preparations, the library preparations and sequencing technology and parameters, and sample quality

control filters are provided in previously published work[46,49]. Furthermore, details of the bioinformatic

pipeline used to generate count level data has been previously described[22]. Briefly, reads were aligned to

the GENCODE24 (GRCh38) reference genome with STAR[50], and gene counts generated using the

HTSeq algorithm[51]. Genes that had more than one counts per million total reads total reads in at least

50% of samples in each tissue and diagnosis category were used for further analysis.

Differential Expression analysis on Mayo and ROS/MAP cohorts

For gene filtering we used false discovery rate of 0.05 from the previously published differential

expression analysis of Mayo and ROS/MAP RNA seq data[22]. Briefly, case control status was harmonized

across the Mayo and ROS/MAP cohorts, where controls were defined as individuals with a low burden of

amyloid and tau based on CERAD and Braak scores, and cases with a high burden. Furthermore in

ROS/MAP, clinical diagnosis was also used with controls having to have no cognitive impairment, and

cases have probably AD[22]. Differential expression analysis was run on suitably normalized data – using

conditional quantile normalization to account for variation in gene length and GC content, removing

sample outliers, covariate identification adjustment, with sampling abundance confidence estimated using

a weighted linear model with the voom-limma package[22,52,53]. A fixed/mixed effect linear model is used

to fit the differential expression model on the normalized data[22].

Manifold learning for LOAD

Manifold learning refers to a group of machine learning algorithms that recover a low dimensional

subspace underlying a high dimensional dataset. Manifold learning approaches are typically used in

datasets or applications where data samples lie on an underlying low dimensional latent space (e.g. a tree,

a line, a curved plane). The low dimensional space is learned via a projection from the high dimensional

space of the observed data (e.g. RNA-seq profiles across hundreds of patient samples) down to a low

dimensional space with suitable regularization constraints to enforce smoothness and the structural

constraints of the low dimensional space. (**Figure 1A**). Due to the necessary assumption of an underlying

latent subspace, manifold learning is commonly used in applications where it is known that the observed

data is obtained from a progression of some kind; e.g., i) to infer the temporal ordering of a sequence of

images, or ii) to infer the approximate lineage of cells in a differentiation trajectory using single cell

RNA-Seq data (**Figure 1B-C**).

Here, we repurpose methods originally developed for learning cell lineage using scRNA-Seq data, to infer

the staging of Alzheimer's disease (AD) using bulk RNA-Seq data from post-mortem brain samples with

known AD diagnosis status. Since bulk RNA-Seq has many of the same sampling and distributional

properties as scRNA-Seq, we observe that scRNA-Seq methods are applicable with no additional

modifications. As such, we use the DDRTree manifold learning approach available in the Monocle2 R

package[54]. However, we also show that the estimated staging of disease is quite similar across some of the

other common methods used for scRNA-Seq lineage estimation (**Figures S26-S28**) including Monocle1[55]

and diffusion pseudotime (DPT)[56].

The RNA-Seq data used in this study was generated from post-mortem brain homogenate samples, and

obtained from two separate studies that are a part of the Accelerating Medicines Partnership in

Alzheimer's Disease (AMP-AD) consortium, namely: i) the Religious Orders Study and the Memory and

Aging Project (ROSMAP)[57,58], and ii) the Mayo RNA-seq study[49]. For this paper, we focused our analysis

on the temporal cortex (TCX) and dorsolateral prefrontal cortex (DLPFC) tissue samples. Within the

Mayo RNA-seq study the TCX samples are derived from individuals neuropathologically defined as

either aged controls, LOAD cases, Progressive Supranuclear Palsy (PSP) cases, or pathological aging

(PA) cases[49]. The ROSMAP study is a prospective longitudinal cohort of an aging population, and has

samples from participants with clinical and neuropathological diagnoses of LOAD[46], aged controls, and

individuals with mild cognitive impairment. Furthermore, most results presented in the main paper are

from female samples only unless indicated otherwise, as we observed significant sex differences in the

transcriptomic data consistent with current knowledge of sex differences in LOAD[59,60]. For replication,

we also consider microarray data generated from the Illumina DASL gene expression platform from the

Mayo eGWAS study from TCX for N=186 patients, of which 108 were neuropathologically confirmed

AD Cases and 78 were controls[23]. Probes were mapped to genes using BioMart. Data was adjusted for

plate using ordinary least squares regression prior to manifold learning.

Manifold learning using Discriminative Dimensionality Reduction Tree (DDRTree)

DDRTree is a manifold learning algorithm that infers a smooth low dimensional manifold by an approach

called reverse graph embedding. Briefly, the algorithm simultaneously learns a non-linear projection to a

latent space where the points lie on a spanning tree. A reverse embedding is also simultaneously learned

from the latent space to the high dimensional data. Mathematically, the DDRTree algorithm can be posed

as the following optimization problem:

$$\sum_{i=1}^{N} ||x_i - Wz_i||^2 + \frac{\lambda}{2} \sum_{k,k'} b_{k,k'}||Wy_k - Wy_{k'}||^2$$

$$+ \gamma \left[\sum_{k=1}^{K} \sum_{i=1}^{N} r_{i,k}||z_i - y_k||^2 + \sigma r_{i,k} log\ (r_{i,k}) \right]$$

$$s.t.\quad B\ represents\ a\ spanning\ tree,$$

$$W^T W = I, r_{i,k} \geq 0, \sum_{k=1}^{K} r_{i,k} = 1$$

Here, $\{z_i\}_{i=1}^{N} \in R^{genes}$ represents RNA-Seq data from each patient sample, $\{z_i\}_{i=1}^{N} \in R^2$ represents the

latent representation of each sample as inferred by the algorithm, $\{y_k\}_{k=1}^{K}$ represents the centers of

clusters in the dataset, $W \in R^{2 \times genes}$ represents an inverse mapping from the latent space to the high

dimensional space of RNA-Seq data, $B \in R^{K \times K}$ represents a spanning tree on which the centers of the

clusters lie and $R \in R^{N \times K}$ captures the soft clustering information of samples in the dataset. The first term

of the optimization problem is responsible for learning a low dimensional representation of the data such

that an inverse mapping exists to the high dimensional data points, the second term learns the tree

structure of the points and the third term learns a soft clustering for the latent dimension points as well as

the centers of the clusters. Despite the non-convexity of the problem, each individual optimization

variable can be solved for efficiently using alternative minimization as described previously[61]. This

algorithm was implemented using the Monocle package in R[19]. When fitting the Monocle objects we also

considered various adjustments to the expression data prior to manifold learning for 10 principal

components of genetic ancestry, RNA integrity number, post mortem interval, age, Braak score - among

other potential confounds. The code to infer the lineage in MayoRNAseq is available here

https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/TCX_GenerateMonocleDS_new.R, and code used

to infer the lineage in ROSMAP is available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/DLPFC_GenerateMonocleDS_new.R. Code to

perform analysis on Mayo eGWAS study is available here: https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_ben_april_2020/mayo_egwas_GenerateMonocleDS

_new.R.

Branch assignment and pseudotime calculation for samples

Branch assignment and pseudotime calculation was also performed using the *Monocle* package using

techniques described previously[19]. Briefly, pseudotime is calculated by first identifying a root point on

one of the two ends of the maximum diameter path in the tree. Then the pseudotime of each point is

calculated by projecting it to its closest point on the spanning tree and calculating the geodesic distance to

the root point. Assigning samples to branches is done by first identifying the branches of the spanning

tree and then assigning samples to the branch on which their projection to the spanning tree lies on.

Robustness of pseudotime was assessed with leave-one-out cross validation by dropping one sample at a

time, running the DDRTree method with Monocle, and then computing the absolute value of the

correlation between the pseudotime estimated with the reduced data-set, and the pseudotime estimated

with the full data set. Alternative approaches for performing dimensionality reduction included principal

component analysis (PCA), t-stochastic neighborhood embedding (tSNE)[33], and uniform manifold

approximation and projection (UMAP)[34], which were all run on the same data set as the DDRTree method

was run in R (here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/TCX_GenerateMonocleDS_new.R and

https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/DLPFC_GenerateMonocleDS_new.R).

Association of pseudotime with AD status, hallmarks of Alzheimer's disease, and cognitive diagnosis

We test for association between disease pseudotime and AD case or control status with logistic regression

with AD case or control status as the outcome and inferred pseudotime as the dependent variable in both

the Mayo and ROS/MAP studies. We test for association between pseudotime and hallmarks of disease

in the ROS/MAP studies for both Braak (measure of tau pathology) score and CERAD score (measure of

amyloid pathology) with an ordinal logistic regression model, with the neuropath score as the ordered

outcome, and pseudotime as the dependent variable. Finally, we test for association between disease

pseudotime and cognitive diagnosis for the following ordered clinical diagnoses of no cognitive

impairment, mild cognitive impairment, and probable Alzheimer's disease with an ordinal logistic

regression model. All code for running these association tests is available https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/paper_figures.Rmd.

Inferring cell type specific expression patterns given marker gene expression as a function of pseudotime

List of marker genes for different major cell types in the brain was curated from a previously published

brain cell expression signature study[37]. The marker gene list was then pruned to include only genes that

were included in lineage construction. Each gene's expression as a function of pseudotime was then

obtained by smoothing using a smoothing spline of degree of freedom = 3 and normalized to lie in [0,1].

The smoothing was done to remove the effects of technical noise introduced due to RNA-Seq and the

normalization was done since the absolute expression levels of genes might be very different from each

other. The smoothed and normalized expression of marker genes for each category were then averaged to

obtain the average marker gene expression as a function of pseudotime.   A linear model was used to test

for association between average expression of a given cell type expression signature and pseudotime.

Association between GWAS loci and correlation with pseudotime

To test for association between pseudotime and LOAD GWAS genes, we computed the Spearman's

correlation between each gene's expression and pseudotime in the Mayo and ROS/MAP studies. Next,

we considered the 60 highly prioritized genes (priority score greater than 4) identified within AD GWAS

loci by the International Genetics of Alzheimer's Project (IGAP)[25]. We test for a difference between the

correlation with pseudotime of background of all other genes and the IGAP AD genes using a linear

model, and see a significant increase in correlation between gene expression and pseudotime in both the

Mayo and ROS/MAP study for AD GWAS genes.

Branch specific differential expression analysis

We perform a state specific differential expression analysis using a one-way ANOVA model in both the

Mayo and ROS/MAP studies. The branch with the highest proportion of AD controls is defined as the

reference branch for all analyses. We use Tukey's honest significant difference method to compute P-

values for the test for change in expression of a given gene compared to the reference branch. Genes are

grouped based on their branch and direction of change in expression for further downstream pathway

enrichment analyses. Overlap between differential expressed genes was depicted using UpSet plots[62]

(**Figure S24**). Code to run analyses are available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/DLPFC_DE_Anova.R for ROS/MAP and here

https://github.com/Sage-Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/TCX_DE_Anova.R for

Mayo.

Branch specific gene expression signatures

Branch specific expression signature was obtained by first calculating the average normalized expression

for all genes in each state/branch. This was followed by performing a bi-clustering using the pheatmap

package in R (https://cran.r-project.org/web/packages/pheatmap/index.html) which uses hierarchical

clustering on both samples and genes. We also used the pheatmap R package to visualize the state

specific expression signatures.

Disease resistant subgroup validation analysis

After identifying potentially disease resistant individuals in the Mayo RNAseq study based on the TCX

brain region (individuals from branch 5, **Figure 5A**), we considered the Mayo eGWAS TCX data, and

defined neuropathological controls with pseudotimes in the top quintile of all pseudotimes as disease

resistant (n=9). We then performed a differential expression analysis using linear regression to identify

array probes that were differentially expressed between resistant and non-resistant individuals, of which

there were more than 5000 probes that were either up or down regulated at an FDR of 0.05. Overlaps

were explored between the branch specific gene clusters from Mayo RNAseq (Figure 5B) and these Mayo

eGWAS resistance differential expressed probes using UpSet plots[62] (**Figure S25**).

Gene set enrichment analyses

For each branch specific differential expression gene set (DEGs) in both Mayo RNAseq and ROS/MAP

we perform a gene set enrichment analysis against Gene Ontology pathways using the enrichR[42] R

package. Only pathways with FDR < 0.05 are reported. The code we used to run the ROS/MAP DEG

enrichments are available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/lineage.Rmd, the code we used to run the Mayo

DEG enrichments are available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/lineageTCX.Rmd, and the code we used to run the

branch specific gene expression signature pathway enrichments is available here https://github.com/Sage-

Bionetworks/AMPAD_Lineage/blob/paper_rewrites_1/resilience.Rmd.

Whole-genome sequencing

Whole-genome sequencing was performed at the New York Genome Center for all individuals from the

ROS/MAP and Mayo cohorts. Detailed information for both data sets can be accessed via synapse

(DOI:10.7303/syn2580853). Briefly, 650ng of genomic DNA from whole blood was sheared using a

Covaris LE220 sonicator. DNA fragments underwent bead-based size selection and were subsequently

end-repaired, adenylated, and ligated to Illumina sequencing adapters. Libraries were sequenced on an

Illumina HiSeq X sequencer using 2 x 150bp cycles. Paired-end reads were aligned to the GRCh37

(hg19) human reference genome using the Burrows-Wheeler Aligner (BWA-MEM v0.7.8) and processed

using the GATK best-practices workflow[63,64]. This included marking of duplicate reads by the use of

Picard tools v1.83, local realignment around indels, and base quality score recalibration (BQSR) via

Genome Analysis Toolkit (GATK v3.4.0). Joint variant calling files (vcfs) for whole-genome sequencing

data for the Mayo and ROS/MAP cohort were obtained through the AMP-AD knowledge portal

(DOI:10.7303/syn10901595).

Single variant association with pseudotime in two independent cohorts

Likelihood ratio tests within a linear regression framework were used to model the relationship between

the quantitative expression trait pseudotime and genetic variants in 436 AD cases. Genome-wide genetic

association analysis was performed for 305 female patients in the ROS/MAP cohort and 131 female

patients in the Mayo cohort for which both genotyping and post-mortem RNA-seq data was available. An

efficient mixed model approach, implemented in the EMMAX software suite, was used to account for

potential biases and cryptic relatedness among individuals[65]. Only variants with MAF > 0.05, genotyping

call rates > 95%, minimum sequencing depth of 20 reads and Hardy-Weinberg equilibrium p $>10^{-4}$ were

considered for analysis. Quantile-quantile plots (**Figure S21-S22**) for the test statistics showed no

significant derivation between expected and observed p-values, highlighting that there is no consistent

differences across cases and controls except for the small number of significantly associated variants.

Furthermore, the genomic inflation factor (lambda) was determined to be 0.99 for the Mayo and 0.98 for

the ROS/MAP single variant association tests. This highlights that potential confounding factors, such as

population stratification have been adequately controlled.

Data Availability

All data analyzed in the study is publicly available and described in detail in previous publications[22–24,46].

Specifically, we use a version of the RNA-seq data from the ROS/MAP study

(DOI:10.7303/syn8456638.22) and RNA-seq data from the Mayo RNAseq (10.7303/syn8466816.19) run

through the same bioinformatic processing pipeline[22]. The array expression data from the Mayo eGWAS

study is available at DOI:10.7303/syn3617054.1.

Code Availability

All code is publicly available (https://github.com/Sage-Bionetworks/AMPAD_Lineage). References to code to perform specific analyses are described in detail in the **Methods**.

References

1. Cummings, J. L., Morstorf, T. & Zhong, K. Alzheimer's disease drug-development pipeline: few

candidates, frequent failures. *Alzheimers. Res. Ther.* **6**, 37 (2014).

2. Cummings, J. L., Doody, R. & Clark, C. Disease-modifying therapies for Alzheimer disease:

Challenges to early intervention. *Neurology* **69**, 1622–1634 (2007).

3. Ferreira, D. *et al.* Distinct subtypes of Alzheimer's disease based on patterns of brain atrophy:

longitudinal trajectories and clinical applications. *Sci. Rep.* **7**, 46263 (2017).

4. Bredesen, D. E. Metabolic profiling distinguishes three subtypes of Alzheimer's disease. *Aging*

(Albany NY) **7**, 595 (2015).

5. Brier, M. R. *et al.* Tau and Aβ imaging, CSF measures, and cognition in Alzheimer's disease.

Sci. Transl. Med. **8**, 338ra66--338ra66 (2016).

6. Gordon, B. A. *et al.* The relationship between cerebrospinal fluid markers of Alzheimer pathology

and positron emission tomography tau imaging. *Brain* **139**, 2249–2260 (2016).

7. Dichgans, M. *et al.* METACOHORTS for the study of vascular disease and its contribution to

cognitive decline and neurodegeneration: An initiative of the Joint Programme for

Neurodegenerative Disease Research. *Alzheimer's Dement.* **12**, 1235–1249 (2016).

8. Jack, C. R. *et al.* Tracking pathophysiological processes in Alzheimer's disease: an updated

hypothetical model of dynamic biomarkers. *Lancet. Neurol.* **12**, 207–16 (2013).

9. Au, R., Piers, R. J. & Lancashire, L. Back to the future: Alzheimer's disease heterogeneity

revisited. *Alzheimer's Dement. (Amsterdam, Netherlands)* **1**, 368–370 (2015).

10. Carrasquillo, M. M. *et al.* Late-onset Alzheimer's risk variants in memory decline, incident mild

cognitive impairment, and Alzheimer's disease. *Neurobiol. Aging* **36**, 60–7 (2015).

11. Zhang, B. *et al.* Integrated systems approach identifies genetic nodes and networks in late-onset

Alzheimer's disease. *Cell* **153**, 707–20 (2013).

12. Mostafavi, S. *et al.* A molecular network of the aging human brain provides insights into the

pathology and cognitive decline of Alzheimer's disease. *Nat. Neurosci.* **21**, 811–819 (2018).

13.   Conway, O. J. *et al.* ABI3 and PLCG2 missense variants as risk factors for neurodegenerative

626   diseases in Caucasians and African Americans. *Mol. Neurodegener.* **13**, 53 (2018).

627   14.   Allen, M. *et al.* Conserved brain myelination networks are altered in Alzheimer's and other

628   neurodegenerative diseases. *Alzheimers. Dement.* **14**, 352–366 (2018).

629   15.   Allen, M. *et al.* Divergent brain gene expression patterns associate with distinct cell-specific tau

630   neuropathology traits in progressive supranuclear palsy. *Acta Neuropathol.* **136**, 709–727 (2018).

631   16.   Bengio, Y., Courville, A. & Vincent, P. Representation learning: A review and new perspectives.

632   *IEEE Trans. Pattern Anal. Mach. Intell.* **35**, 1798–1828 (2013).

633   17.   Wolz, R., Aljabar, P., Hajnal, J. V. & Rueckert, D. Manifold Learning for Biomarker Discovery in

634   MR Imaging. in 116–123 (Springer, Berlin, Heidelberg, 2010). doi:10.1007/978-3-642-15948-

635   0_15

636   18.   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by

637   pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32**, (2014).

638   19.   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat. Methods*

639   **14**, 979–982 (2017).

640   20.   Rosenberg, A. B. *et al.* Single-cell profiling of the developing mouse brain and spinal cord with

641   split-pool barcoding. *Science (80-.).* **360**, 176–182 (2018).

642   21.   Mukherjee, S., Zhang, Y., Fan, J., Seelig, G. & Kannan, S. Scalable preprocessing for sparse

643   scRNA-seq data exploiting prior knowledge. *Bioinformatics* **34**, i124--i132 (2018).

644   22.   Wan, Y.-W. *et al.* Meta-Analysis of the Alzheimer's Disease Human Brain Transcriptome and

645   Functional Dissection in Mouse Models. *Cell Rep.* **32**, 107908 (2020).

646   23.   Zou, F. *et al.* Brain Expression Genome-Wide Association Study (eGWAS) Identifies Human

647   Disease-Associated Variants. *PLoS Genet.* **8**, e1002707 (2012).

648   24.   Allen, M. *et al.* Human whole genome genotype and transcriptome data for Alzheimer's and other

649   neurodegenerative diseases. *Sci. Data* **3**, (2016).

25.   Kunkle, B. W. *et al.* Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk

loci and implicates Aβ, tau, immunity and lipid processing. *Nat. Genet.* **51**, 414–430 (2019).

26.   Allen, M. *et al.* Novel late-onset Alzheimer disease loci variants associate with brain gene

expression. *Neurology* **79**, 221–8 (2012).

27.   Allen, M. *et al.* Late-onset Alzheimer disease risk variants mark brain regulatory loci. *Neurol.*

Genet. **1**, e15 (2015).

28.   Allen, M. *et al.* Association of MAPT haplotypes with Alzheimer's disease risk and MAPT brain

gene expression levels. *Alzheimers. Res. Ther.* **6**, 39 (2014).

29.   Carrasquillo, M. M. *et al.* A candidate regulatory variant at the TREM gene cluster associates with

decreased Alzheimer's disease risk and increased TREML1 and TREM2 brain gene expression.

Alzheimers. Dement. **13**, 663–673 (2017).

30.   Sims, R. *et al.* Rare coding variants in PLCG2, ABI3, and TREM2 implicate microglial-mediated

innate immunity in Alzheimer's disease. *Nat. Genet.* **49**, 1373–1384 (2017).

31.   Braak, H., Alafuzoff, I., Arzberger, T., Kretzschmar, H. & Del Tredici, K. Staging of Alzheimer

disease-associated neurofibrillary pathology using paraffin sections and immunocytochemistry.

Acta Neuropathol. **112**, 389–404 (2006).

32.   Mirra, S. S. *et al.* The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part

II. Standardization of the neuropathologic assessment of Alzheimer's disease. *Neurology* **41**, 479–

86 (1991).

33.   Maaten, L. van der & Hinton, G. Visualizing data using t-SNE. *J. Mach. Learn. Res.* **9**, 2579--

2605 (2008).

34.   McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for

Dimension Reduction. (2018).

35.   Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis. *Nature* **566**,

496–502 (2019).

36.   De Strooper, B. & Karran, E. The Cellular Phase of Alzheimer's Disease. *Cell* **164**, 603–615

(2016).

37.   Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and

678   vascular cells of the cerebral cortex. *J. Neurosci.* **34**, 11929–47 (2014).

679   38.   Chibnik, L. B. *et al.* Susceptibility to neurofibrillary tangles: role of the PTPRD locus and limited

680   pleiotropy with other neuropathologies. *Mol. Psychiatry* **23**, 1521 (2018).

681   39.   Lambert, J.-C. *et al.* Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for

682   Alzheimer's disease. *Nat. Genet.* **45**, 1452 (2013).

683   40.   Janelidze, S. *et al.* CSF biomarkers of neuroinflammation and cerebrovascular dysfunction in early

684   Alzheimer disease. *Neurology* **91**, e867--e877 (2018).

685   41.   Whitwell, J. L. *et al.* Neuroimaging correlates of pathologically defined subtypes of Alzheimer's

686   disease: a case-control study. *Lancet Neurol.* **11**, 868–877 (2012).

687   42.   Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool.

688   *BMC Bioinformatics* **14**, 128 (2013).

689   43.   Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology

690   Consortium. *Nat. Genet.* **25**, 25–9 (2000).

691   44.   Dong, A. *et al.* Heterogeneity of neuroanatomical patterns in prodromal Alzheimer's disease: links

692   to cognition, progression and biomarkers. *Brain* **140**, aww319 (2016).

693   45.   Wang, X. *et al.* Genetic determinants of disease progression in Alzheimer's disease. *J. Alzheimers.*

694   *Dis.* **43**, 649–55 (2015).

695   46.   De Jager, P. L. *et al.* A multi-omic atlas of the human frontal cortex for aging and Alzheimer's

696   disease research. *Sci. Data* **5**, 180142 (2018).

697   47.   Allen, M. *et al.* Human whole genome genotype and transcriptome data for Alzheimer's and other

698   neurodegenerative diseases. *Sci. data* **3**, 160089 (2016).

699   48.   Schneider, J. A., Arvanitakis, Z., Bang, W. & Bennett, D. A. Mixed brain pathologies account for

700   most dementia cases in community-dwelling older persons. *Neurology* **69**, 2197–2204 (2007).

701   49.   Allen, M. *et al.* Human whole genome genotype and transcriptome data for Alzheimer's and other

702   neurodegenerative diseases. *Sci. Data* **3**, 160089 (2016).

703   50.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

704   51.   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-throughput

705   sequencing data. *Bioinformatics* **31**, 166–169 (2015).

706   52.   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear model

707   analysis tools for RNA-seq read counts. *Genome Biol.* **15**, (2014).

708   53.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and

709   microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

710   54.   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat. Methods*

711   **14**, 979–982 (2017).

712   55.   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by

713   pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32**, 381–386 (2014).

714   56.   Haghverdi, L., Büttner, M., Wolf, F. A., Buettner, F. & Theis, F. J. Diffusion pseudotime robustly

715   reconstructs lineage branching. *Nat. Methods* **13**, 845–848 (2016).

716   57.   A Bennett, D. *et al.* Overview and findings from the rush Memory and Aging Project. *Curr.*

717   *Alzheimer Res.* **9**, 646–663 (2012).

718   58.   Bennett, D. A. *et al.* Religious orders study and rush memory and aging project. *J. Alzheimer's*

719   *Dis.* 1–28 (2018).

720   59.   Ferretti, M. T. *et al.* Sex differences in Alzheimer disease—the gateway to precision medicine.

721   *Nat. Rev. Neurol.* 1 (2018).

722   60.   Deming, Y. *et al.* Sex-specific genetic predictors of Alzheimer's disease biomarkers. *Acta*

723   *Neuropathol.* **136**, 857–872 (2018).

724   61.   Mao, Q., Wang, L., Goodison, S. & Sun, Y. Dimensionality Reduction Via Graph Structure

725   Learning. in *Proceedings of the 21th ACM SIGKDD International Conference on Knowledge*

726   *Discovery and Data Mining - KDD '15* 765–774 (ACM Press, 2015).

727   doi:10.1145/2783258.2783309

728   62.   Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the visualization of

729   intersecting sets and their properties. *Bioinformatics* **33**, 2938–2940 (2017).

730   63.   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-

731   generation DNA sequencing data. *Genome Res.* **20**, 1297–303 (2010).

732   64.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.

733   *Bioinformatics* **25**, 1754–60 (2009).

734   65.   Kang, H. M. *et al.* Variance component model to account for sample structure in genome-wide

735   association studies. *Nat. Genet.* **42**, 348–354 (2010).

Acknowledgements

This work was supported by NIA grants U54AG054345 and RF1AG057443. The ROSMAP Study data

were provided by the Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago. Data

collection was supported through funding by NIA grants P30AG10161, R01AG15819, R01AG17917,

R01AG30146, R01AG36836, U01AG32984, U01AG46152, the Illinois Department of Public Health,

and the Translational Genomics Research Institute. Mayo RNAseq Study data were provided by the

following sources: The Mayo Clinic Alzheimer's Disease Genetic Studies, led by Dr. Nilufer Ertekin-

Taner and Dr. Steven G. Younkin, Mayo Clinic, Jacksonville, FL using samples from the Mayo Clinic

Study of Aging, the Mayo Clinic Alzheimer's Disease Research Center, and the Mayo Clinic Brain Bank.

Data collection was supported through funding by NIA grants P50 AG016574, R01 AG032990, U01

AG046139, R01 AG018023, U01 AG006576, U01 AG006786, R01 AG025711, R01 AG017216, R01

AG003949, NINDS grant R01 NS080820, CurePSP Foundation, and support from Mayo Foundation.

Study data includes samples collected through the Sun Health Research Institute Brain and Body

Donation Program of Sun City, Arizona. The Brain and Body Donation Program is supported by the

National Institute of Neurological Disorders and Stroke (U24 NS072026 National Brain and Tissue

Resource for Parkinson's Disease and Related Disorders), the National Institute on Aging (P30 AG19610

Arizona Alzheimer's Disease Core Center), the Arizona Department of Health Services (contract 211002,

Arizona Alzheimer's Research Center), the Arizona Biomedical Research Commission (contracts 4001,

0011, 05-901 and 1001 to the Arizona Parkinson's Disease Consortium) and the Michael J. Fox

Foundation for Parkinson's Research. MSBB data were generated from postmortem brain tissue collected

through the Mount Sinai VA Medical Center Brain Bank and were provided by Dr. Eric Schadt from

Mount Sinai School of Medicine.

Author Contributions

S.M. and B.A.L. designed the study. S.M., L.H., and B.A.L. performed the analyses. S.M., L.H., C.P.,

S.J., G.G., A.K.G., S.K.S., P.L.D.J., N.E.T., G.W.C., L.M.M., and B.A.L. wrote the manuscript.

Tables

Table 1 - Representative significant Gene Ontology pathway enrichments (FDR<0.05) of differentially

expressed genes for each branch (FDR < 0.05). Differential expressed genes are identified with an

ANOVA analysis, with Branch 1 as the reference.

Brain Region	Direction	Branch	Representative Enriched Gene Ontology Terms
TCX	Down	2	prespliceosome (GO:0071010), mitochondrial electron transport, cytochrome c to oxygen (GO: 0006123)
		3	negative regulation of microtubule polymerization or depolymerization (GO:0031111)
		4	mitochondrial electron transport, NADH to ubiquinone (GO: 0006120), spliceosomal tri-snRNP complex (GO:0097526), negative regulation of microtubule depolymerization (GO:0007026)
		5	axon (GO:0030424), protein kinase C activity (GO:0004697),
		6	gamma-tubulin large complex (GO:0000931), U1 snRNP (GO:0005685), mitochondrial respiratory chain complex IV (GO:0005751), response to cadmium ion (GO:0046686)
	Up	2	
		3	fatty acid elongase activity (GO:0009922), ubiquitin protein ligase activity (GO:0061630)
		4	transforming growth factor beta-activated receptor activity (GO:0005024), hippo signaling (GO:0035329), regulation of extrinsic apoptotic signaling pathway via death domain receptors (GO: 1902041), regulation of DNA repair (GO: 0006282)

		5	regulation of apoptotic process (GO:0042981), leptin mediated signaling pathway (GO:0033210), negative regulation of hippo signaling (GO:0035331), small GTPase binding (GO:0031267)
		6	extracellular ligand-gated ion channel activity (GO:0005230), integral component of mitochondrial inner membrane (GO:0031305)
DLPFC	Down	2	DNA repair (GO:0006281), intracellular protein transport (GO:0006886)
		3	mismatch repair complex binding (GO:0032404)
		4	
		5	mitochondrial respiratory chain complex assembly (GO: 0033108)
		6	
	Up	2	racemase and epimerase activity (GO: 0016857)
		3	racemase and epimerase activity (GO: 0016857)
		4	vesicle mediated transport (GO: 0016192)
		5	NuRD complex (GO: 0016581)
		6	microtuble motor activity (GO:0003777), AP-2 adaptor complex binding (GO:0035612)

Figure 1 - Overview of manifold learning for unraveling staging in Alzheimer's disease. A) Illustration of

steps in manifold learning using reverse graph embedding DDRTree method. B) Illustration of lineage

inference process for LOAD. RNA-seq samples with different disease diagnoses were pooled, batch

normalized, and a smooth manifold was learned for each brain region across individuals (each point is an

individual). Total sample numbers are indicated across Mayo RNAseq TCX and ROS/MAP DLPFC for

the different diagnoses in parentheses.

777
778

Figure 2 - Manifold learning accurately infers disease states and stages from RNA-seq samples. A)

Estimated disease progression trees from temporal cortex (TCX) and B) dorsolateral prefrontal cortex

(DLPFC) brain regions showing localization of identified LOAD samples on particular branches. C)

Distribution of pseudotime for AD cases and controls for both DLPFC and TCX. D) Distribution of

expression correlation with pseudotime for both LOAD GWAS genes and non-LOAD GWAS genes.

Figure 3 - Manifold learning replicates existing measures of staging in LOAD in DLPFC samples. A)

Samples colored by 3 different external measures of LOAD staging namely: Braak Score (tau pathology),

CERAD Score (amyloid pathology) and Cognitive Diagnosis (Clinical measure of disease severity).

Black lines denote inferred lineages. B) Distribution of samples by inferred stage for different distinct

stages in each of the three methods of measuring LOAD severity. Inferred disease stages generally

corresponded with all methods, and Cognitive diagnosis demonstrated the strongest alignment.

Figure 4 – Cell type gene expression signatures as a function of disease pseudotime. A) Mean expression

of cell markers for astrocytes, neurons, microglia, and oligodendrocytes as a function of pseudotime for

TCX brain region, B) mean expression of cell markers for astrocytes, neurons, microglia,

oligodendrocytes as a function of pseudotime for DLPFC (B) brain region.

Figure 5 – Disease resistant state. A) The inferred manifold from the TCX region with samples colored

by their inferred disease subtype/state. State 5 (dots, circled) lies at the late end of the disease trajectory,

indicating a strong disease-like transcriptomic phenotype, yet most samples in the group did not have

pathologically diagnosed AD (Figure 2A). We hypothesize this group represents a disease resistant state

to the disease. B) Biclustering results of average expression from each disease state, with increased

expression of a gene cluster (Cluster 4) unique to State 5.