Aggregation of CAT tails blocks their degradation and causes

proteotoxicity in *S. cerevisiae*

Cole S. Sitron[1,*], Joseph H. Park[1,2,*], Jenna M. Giafaglione[1,3], and Onn Brandman[1,†]

[1]Department of Biochemistry, Stanford University, Stanford CA, 94305. [2]Department of

Chemical & Systems Biology, Stanford University, Stanford CA, 94305. [3]Current address:

Molecular Biology Interdepartmental Program, University of California Los Angeles, Los Angeles

CA, 90095. [*]These authors contributed equally to the work. [†]Corresponding author, to whom

correspondence and material requests should be addressed: onn@stanford.edu.

Running Title: CAT tail aggregation

Abstract The Ribosome-associated Quality Control (RQC) pathway co-translationally marks

incomplete polypeptides from stalled translation with two signals that trigger their proteasome-

mediated degradation. The E3 ligase Ltn1 adds ubiquitin and Rqc2 directs the large ribosomal

subunit to append carboxy-terminal alanine and threonine residues (CAT tails). When excessive

amounts of incomplete polypeptides evade Ltn1, CAT-tailed proteins accumulate and can self-

associate into aggregates. CAT tail aggregation has been hypothesized to either protect cells by

sequestering potentially toxic incomplete polypeptides or harm cells by disrupting protein

homeostasis. To distinguish between these possibilities, we modulated CAT tail aggregation in

Saccharomyces cerevisiae with genetic and chemical tools to analyze CAT tails in aggregated

and un-aggregated states. We found that enhancing CAT tail aggregation induces proteotoxic

stress and antagonizes degradation of CAT-tailed proteins, while inhibiting aggregation reverses

these effects. Our findings suggest that CAT tail aggregation harms RQC-compromised cells

and that preventing aggregation can mitigate this toxicity.

14
Introduction

Failed rounds of translation produce incomplete, potentially toxic polypeptides that organisms

across all clades of life have evolved responses to degrade (Brandman and Hegde, 2016;

Defenouillère and Fromont-Racine, 2017; Ikeuchi et al., 2018; Joazeiro, 2019; Moore and

Sauer, 2007). In prokaryotes, the primary degradative response involves a tRNA-mRNA hybrid

molecule (tmRNA) (Moore and Sauer, 2007). The tmRNA enters stalled ribosomes, re-initiates

translation elongation with its tRNA moiety and switches the ribosome's template to its mRNA

moiety (Moore and Sauer, 2007). This prompts the ribosome to synthesize a tmRNA-encoded

tag on the incomplete polypeptide's C-terminus that marks it for proteolysis (Moore and Sauer,

2007). The eukaryotic response, called Ribosome-associated Quality Control (RQC), begins

when a set of factors recognize ribosomes that have stalled on the same mRNA and collided

into each other (Ikeuchi et al., 2019; Juszkiewicz et al., 2018; Simms et al., 2017). These factors

then split the ribosomes into their large and small subunits, leaving the incomplete polypeptide

(RQC substrate) tethered to the large subunit (Juszkiewicz and Hegde, 2017; Lyumkis et al.,

2014; Matsuo et al., 2017; Pisareva et al., 2011; Shao et al., 2013; Shoemaker et al., 2010;

Sitron et al., 2017; Sundaramoorthy et al., 2017; Tsuboi et al., 2012). The E3 ligase Ltn1 binds

to the large subunit and ubiquitylates the incomplete polypeptide, marking it for proteasomal

degradation (Bengtson and Joazeiro, 2010; Brandman et al., 2012; Defenouillère et al., 2013;

Shao et al., 2013; Verma et al., 2013).

Disruption of tmRNA or Ltn1 compromises the cell's ability to degrade incomplete polypeptides

and reduces survival under stresses that increase translational stalling (Abo et al., 2002; Choe

et al., 2016; Defenouillère et al., 2013; Kostova et al., 2017; Lytvynenko et al., 2019; Moore and

Sauer, 2005; Oussenko et al., 2005). This deficit in fitness at the cellular level has clinically-

relevant consequences. tmRNA deficiency prevents growth of some disease-causing

prokaryotes (e.g. *Mycoplasma genitalium* and *Staphylococcus aureus*) and decreases the

virulence of others (e.g. *Salmonella enterica* and *Streptococcus pneumoniae*) (Chaudhuri et al.,

2009; Glass et al., 2006; Julio et al., 2000; Mann et al., 2012). In metazoan eukaryotes,

mutations to *LTN1* or perturbations that introduce large influxes of RQC substrates lead to

neurodegeneration (Chu et al., 2009; Ishimura et al., 2014; Wu Z, Tantray I, Lim J, Chen S, Li Y,

Davis Z, Sitron C, Dong J, Gispert S, Auburger G, Brandman O, Bi X, Snyder M, Lu B., 2019).

Each of these phenotypes highlights the central role that tmRNA and Ltn1 play in maintaining

protein homeostasis and avoiding the toxicity associated with compromised co-translational

quality control.

A conserved back-up degradation pathway mediated by Rqc2 and its prokaryotic homologs

mitigates some of the toxicity associated with loss of tmRNA or Ltn1 function (Lytvynenko et al.,

2019; Sitron and Brandman, 2019). Rqc2 homologs bind the large ribosomal subunit and direct

it to elongate the incomplete polypeptide's C-terminus with either alanine ("Ala tails" in

prokaryotes) or both alanine and threonine residues ("CAT tails" in yeast) (Lytvynenko et al.,

2019; Osuna et al., 2017; Shen et al., 2015). Metazoan CAT tails may include a more diverse

repertoire of amino acids (Wu Z, Tantray I, Lim J, Chen S, Li Y, Davis Z, Sitron C, Dong J,

Gispert S, Auburger G, Brandman O, Bi X, Snyder M, Lu B., 2019). These extensions, made

without a small subunit or mRNA, act as "degrons" to mark incomplete polypeptides for

degradation by the bacterial protease ClpXP or the eukaryotic proteasome (Lytvynenko et al.,

2019; Sitron and Brandman, 2019).

8
Loss of Ltn1 function results in a buildup of CAT-tailed ("CATylated") proteins (Shen et al.,

2015). The CAT tails on these proteins can homo-typically associate and form aggregates of

CATylated proteins, which have been observed in yeast and metazoans (Choe et al., 2016;

Defenouillère et al., 2016; Wu Z, Tantray I, Lim J, Chen S, Li Y, Davis Z, Sitron C, Dong J,

Gispert S, Auburger G, Brandman O, Bi X, Snyder M, Lu B., 2019; Yonashiro et al., 2016). The

presence of threonine in yeast CAT tails increases their aggregation propensity (Yonashiro et

al., 2016). It is thereby an attractive hypothesis that CAT tail aggregation has an adaptive

function during stress. For instance, aggregation may protect the proteome by sequestering

potentially harmful undegraded RQC substrates when Ltn1 becomes overburdened.

Alternatively, it has been hypothesized that CAT tail aggregates disrupt cellular fitness by

depleting the pool of chaperones (Choe et al., 2016; Defenouillère et al., 2016; Yonashiro et al.,

2016), and thereby contribute to the proteotoxicity associated with Ltn1 failure. No study has

evaluated these hypotheses by controlling the aggregation state of CAT tails and testing

whether CAT tail aggregation plays a beneficial or harmful role during stress.

23
We used chemical and genetic approaches in *Saccharomyces cerevisiae* to modulate CAT tail

aggregation propensity in Ltn1-deficient cells and determine how aggregation affects: 1) the

function of CAT tails as degrons and 2) proteotoxic stress. We found that under conditions

favoring CAT tail aggregation (elevated Rqc2 levels), CAT tails lose their degron function and

induce proteotoxic stress. Conversely, we observed that increasing CAT tail solubility (with

guanidinium hydrochloride or by genetic perturbation of RNA Polymerase III function) restores

CAT tail-mediated degradation and reduces proteotoxic stress. Our work demonstrates that

CAT tail aggregation contributes to the toxicity of Ltn1 disruption rather than aiding in adaptation

to stress.

6
7   **Results**

8
CAT tail aggregation is associated with compromised CAT tail degradation.

We began our investigation into the effects of aggregation on CAT tail-mediated degradation by

validating the model RQC substrate RQCsub. RQCsub consists of green fluorescent protein

(GFP) attached via an inert linker including a tobacco etch virus (TEV) protease site to twelve

stall-inducing arginine CGN codons (Figure 1A) (Letzring et al., 2010). Translation of RQCsub

produces a stalled GFP-linker-arginine "arrest product" (Brandman et al., 2012). The RQCsub

arrest product accumulated at low levels in wt strains by SDS-PAGE (Figure 1A). The arrest

product increased in abundance after we compromised RQC function by deleting either of two

RQC genes: the CAT tail-elongating factor *RQC2* or the E3 ubiquitin ligase *LTN1* (Figure 1A).

This result indicates that Rqc2 and Ltn1 contribute to RQCsub degradation. While deletion of

RQC2 or *LTN1* stabilized RQCsub, the resultant protein products displayed different mobilities:

RQCsub from *rqc2Δ* cells migrated as a crisp band and that from *ltn1Δ* cells migrated as a

higher molecular weight smear (Figure 1A). This smear collapsed into a single band after

additional deletion of *RQC2* in the *ltn1Δ* strain (Figure 1A), consistent with the smear containing

CATylated RQCsub. Taken together, these results confirm that RQCsub is degraded via RQC

and CATylated by Rqc2.

25
To vary the aggregation propensity of CAT tails, we overexpressed *RQC2*, which has previously

been shown to increase CAT tail aggregation (Yonashiro et al., 2016). Overexpression of *RQC2*

with the *TDH3* promoter increased Rqc2 levels 28.4-fold relative to the native promoter, as

assessed by levels of red fluorescent protein (RFP)-tagged Rqc2 (Figure 1-figure supplement

1A). In the <u>R</u>QC2 <u>o</u>verexpression and <u>L</u>TN1 <u>d</u>eletion strain (hereafter named "*ROLD*"), a

proportion of RQCsub remained in the well rather than migrating into an SDS-PAGE gel (Figure

1A) despite prior boiling in dodecylsulfate detergent. This detergent-resistant species was more

prominent in *ROLD* than in *ltn1Δ* lysates (Figure 1-figure supplement 1B). TEV treatment, which

cleaves RQCsub's C-terminus to remove its CAT tail, enabled RQCsub purified from *ROLD*

cells to migrate as a single band into the gel (Figure 1-figure supplement 1C). The properties of

this detergent-resistant species are consistent with *RQC2* overexpression reducing the solubility

of CATylated RQCsub in the *ltn1Δ* background. Microscopy revealed that RQCsub had a diffuse

localization in the majority of *ltn1Δ* cells (<1% of cells with inclusions) but became more

punctate in *ROLD* cells (56.9% of cells with inclusions) (Figure 1B). To test whether these

inclusions depended on CATylation, we disrupted CATylation by deleting *RQC2* or

overexpressing a CATylation-incompetent *rqc2$_{aaa}$* allele (Shen et al., 2015) instead of *RQC2-WT*.

Neither of these perturbations (*rqc2Δ ltn1Δ* nor *rqc2$_{aaa}$-OE ltn1Δ*) induced the punctate RQCsub

localization we observed in *ROLD* cells (Figure 1B), confirming that formation of RQCsub

inclusions requires CATylation. These results demonstrate that the *ROLD* genetic background

can be used to enhance CAT tail aggregation.

19
We next sought a tool to inhibit CAT tail aggregation. Millimolar concentrations of the chaotrope

guanidinium hydrochloride (GuHCl) in yeast growth medium solubilize CAT tail aggregates and

prevent propagation of aggregated yeast prions (Byrne et al., 2007; Defenouillère et al., 2016;

Eaglestone et al., 2000; Ness et al., 2002). When we treated *ROLD* cells with GuHCl, we

observed that RQCsub still localized to inclusions by microscopy (Figure 1-figure supplement

1D). However, monitoring RQCsub mobility in SDS-PAGE assessed how GuHCl affected

RQCsub more sensitively than microscopy, revealing that GuHCl treatment enabled a larger

proportion of RQCsub extracted from *ROLD* cells to migrate into the resolving gel rather than

remaining in the well (Figure 1C). GuHCl thus reduces the detergent insolubility of CAT tails and

can thereby serve as a tool to inhibit CAT tail aggregation.

Single-color model RQC substrates like RQCsub, while useful for visualizing CAT tails by SDS-

PAGE and microscopy, have limited utility in degradation measurements due to noisy

expression (Sitron and Brandman, 2019). To assess how aggregation affects degradation of

CATylated RQC substrates, we utilized the two-color model RQC substrate RQCsub$_{LONG}$

(Sitron and Brandman, 2019) (Figure 1D,E). RQCsub$_{LONG}$ includes an internal expression

control, RFP followed by viral T2A peptides, at the N-terminus of GFP-linker-arginine (Figure

1E). Ribosomes translating RQCsub$_{LONG}$ produce RFP then skip formation of a peptide bond

within the T2A sequence (Donnelly et al., 2001; Szymczak and Vignali, 2005), severing RFP

from the GFP-linker-arginine RQC substrate (Figure 1E). As a result, each round of translation

produces RFP and GFP stoichiometrically, but only GFP becomes an RQC substrate (Figure

1E). When we analyzed GFP and RFP separately, we observed that each signal was lower in

ROLD than in *ltn1Δ* cells (Figure 1-figure supplement 2A). This loss of fluorescent signal was

especially evident for the expression control RFP (Figure 1-figure supplement 2A), indicative of

reduced RQCsub$_{LONG}$ translation in the *ROLD* background. This reduction in translation may

explain why we observed lower RQCsub levels in the *ROLD* background compared to *ltn1Δ* by

SDS-PAGE (Figure 1A) and highlights the need for a precise way to measure degradation that

is not confounded by changes in translation.

We then controlled for these differences in translation to quantify how increased CAT tail

aggregation (in the *ROLD* strain) affects degradation of CATylated RQCsub$_{LONG}$. The ratio of

GFP:RFP at the single-cell level serves as an internal, expression-controlled readout of

RQCsub$_{LONG}$ stability (Figure 1E). CAT tail-mediated degradation of RQCsub$_{LONG}$ in the

absence of Ltn1 requires the proteasome-associated E3 ubiquitin ligase Hul5 and the

proteasome (Sitron and Brandman, 2019). Therefore, comparing stability before and after *HUL5*

deletion quantifies CAT tail-mediated degradation (Figure 1E). RQCsub$_{LONG}$ stability did not

change after treatment of *ltn1Δ hul5Δ* or *ROLD hul5Δ* cells with the proteasome inhibitor

bortezomib (Figure 1-figure supplement 2B), supporting the requirement for Hul5 in CAT tail-

mediated degradation of RQCsub$_{LONG}$ by the proteasome. *HUL5* deletion stabilized

RQCsub$_{LONG}$ by 30.2% in *ltn1Δ* and 13.6% in *ROLD* backgrounds (Figure 1F), consistent with

RQC2 overexpression reducing CAT tail-mediated RQCsub$_{LONG}$ degradation. These results

demonstrate that the *ROLD* genetic background, which potentiates CAT tail aggregation,

impairs CAT tail-mediated degradation.

11
To ensure that it is aggregation (and not another effect of *ROLD*) that reduces CAT tail-

mediated degradation, we inhibited aggregation with GuHCl and measured RQCsub$_{LONG}$

stability . We observed that GuHCl treatment had two effects on RQCsub$_{LONG}$ degradation in

ROLD strains. First, increasing concentrations of GuHCl increased the magnitude of the effect

of *HUL5* deletion on the stability of RQCsub$_{LONG}$ (from 13.6% to 20.9% to 28.2% at 0, 5, and 10

mM GuHCl respectively; Figure 1G), suggesting that treating *ROLD* cells with GuHCl rescues

degradation of RQCsub$_{LONG}$. Second, the magnitude of bortezomib-induced RQCsub$_{LONG}$

stabilization increased with increasing GuHCl when Hul5 was intact (Figure 1-figure supplement

2C). This result indicates that the degradation that GuHCl rescued involved both Hul5 and the

proteasome. Taken together, these data imply that the aggregation-conducive *ROLD*

background impairs CAT tail-mediated degradation, and solubilization of CAT tails rescues this

impairment. We thereby posit that CAT tail aggregation antagonizes CAT tail-mediated

degradation.

CAT tails exert proteotoxic stress in their aggregated state.

To determine how aggregation affects the proteotoxicity of CAT tails, we measured proteotoxic

stress after varying CAT tail aggregation propensity (Figure 2A). Proteotoxic stress activates the

transcription factor Heat Shock Factor 1 (Hsf1), which then orchestrates transcription of

chaperones and other proteostasis-restoring factors (Åkerfelt et al., 2007; Morimoto, 2011).

Hsf1 activation scales with the magnitude of proteotoxic stress and a previously-published

reporter can measure the degree of activation (Brandman et al., 2012). This reporter consists of

GFP under the control of an artificial promoter containing four heat shock elements (Sorger and

Pelham, 1987), the consensus binding site for Hsf1 (Figure 2B). *LTN1* deletion alone activated

the Hsf1 reporter 2.07-fold relative to wt, while additional *RQC2* overexpression (*ROLD*) led to

22.6-fold activation (Figure 2B). The potent activation observed in the *ROLD* background

required *LTN1* deletion, as *RQC2* overexpression in the wt background weakly affected the

reporter (Figure 2B). Thus, the combination of *RQC2* overexpression and *LTN1* deletion, which

potentiates CAT tail aggregation (Figure 1-figure supplement 1B) (Yonashiro et al., 2016),

generates a synthetic interaction that results in elevated Hsf1 activity. This increase in Hsf1

activity is consistent with an increase in proteotoxic stress.

17
If CAT tail aggregation causes proteotoxicity, we reasoned that solubilization of CATylated

proteins by GuHCl treatment might reduce this toxicity. GuHCl treatment activated the Hsf1

reporter in a dose-dependent manner in wt, *RQC2-OE*, and *ltn1Δ* strains (Figure 2C), indicating

that GuHCl alone exerts some proteotoxic stress in cells. Despite this general activation, GuHCl

decreased Hsf1 reporter activation in *ROLD* cells (Figure 2C), suggesting that solubilization of

CAT tails alleviates the Hsf1 activation associated with the *ROLD* background. We tested the

specificity of this response by measuring the effect of GuHCl on the Hsf1 reporter in other

genotypes with high Hsf1 activity (Brandman et al., 2012) (Figure 2D). Hsf1 reporter

deactivation was unique to the *ROLD* condition, as GuHCl increased Hsf1 reporter activation in

ump1Δ, *hsp104Δ*, and *hsc82Δ* strains and had no significant effect in the *ssa2Δ* strain (Figure

2D). These results suggest that CAT tails exert proteotoxic stress more potently in their

aggregated than in their soluble state.

4
To ensure that our Hsf1 measurements reflect perturbations to proteostasis, we analyzed how

CAT tail aggregation affects cells' sensitivity to stressors. At 30^{O}C, all of the strains we

examined (wt, *RQC2* overexpression, *ltn1Δ*, and *ROLD*) grew equally well in a spot assay in the

absence and presence of 5 mM GuHCl (Figure 2E, plates 1 and 2). This result demonstrates

that neither the *ROLD* background nor GuHCl impairs growth in the absence of stress. We then

introduced two mild stressors to perturb proteostasis: the thermal stress of 37^{O}C incubation or

introduction of 50 ng/ml cycloheximide to increase translational stalling. Neither of these

stresses induced a growth defect in the *RQC2* overexpression (*RQC2-OE*) or *ltn1Δ* strains

relative to wt (Figure 2E, plates 3 and 5). By contrast, 37^{O}C incubation and 50 ng/ml

cycloheximide severely inhibited growth in the *ROLD* strain (Figure 2E, plates 3 and 5).

Increased CAT tail aggregation thus co-occurs with sensitivity to thermal and ribosome stalling

stresses. Additional GuHCl partially rescued growth of the *ROLD* strain in the presence of

cycloheximide (Figure 2E, plate 5 versus plate 6). The combination of GuHCl and 37^{O}C led to a

synthetic growth defect in all strains (Figure 2E, plate 3 versus plate 4). Despite this growth

defect, the *ROLD* strain grew comparably to wt, *RQC2-OE*, and *ltn1Δ* strains when exposed to

both GuHCl and 37^{O}C (Figure 2E, plate 4). Thus, CAT tail aggregation sensitizes cells to mild

thermal and ribosome stalling stress. The agreement with our Hsf1 measurements (Figure

2B,C) suggests that CAT tail aggregation perturbs proteostasis under the conditions we

measured rather than facilitating adaptation to stress.

24
Disruption of CAT tail aggregation by Pol III perturbation restores CAT tail

degradation and alleviates proteotoxicity.

Given that GuHCl mildly reduced CAT tail aggregation, we sought an orthogonal perturbation

that would more potently inhibit CAT tail aggregation. To find such a perturbation, we screened

for mutations that reduced Hsf1 reporter activation in the RQC-compromised *rqc1Δ rps0aΔ*

strain, which exhibits Hsf1 hyperactivation (Brandman et al., 2012). After crossing a genome-

wide collection of hypomorphs into this background, we found that all hits other than *RQC2*

(whose deletion blocks CATylation) were related to RNA Polymerase III (Pol III). These hits

included a non-essential polymerase biogenesis factor, *BUD27* (Mirón-García et al., 2013), and

essential Pol III subunits, *RPC17* and *RPC160*. Disruption of each of these by deletion or a

"decreased abundance by mRNA perturbation" allele (DAmP) (Yan et al., 2008) nearly

eliminated activation of the Hsf1 reporter in *ROLD* cells compared to wt (Figure 3-figure

supplement 1A). To assess the specificity of this effect to RQC-related Hsf1 activation, we

measured the effect of *BUD27* and *RPC17* perturbations in strains with elevated Hsf1 signaling

(Figure 3A). Perturbations to *BUD27* and *RPC17* reduced Hsf1 reporter activity in wt, *ump1Δ*,

hsp104Δ, *hsc82Δ*, and *ssa2Δ* strains (Figure 3A, black versus grey bars). However, the

magnitude of this reduction in each strain was substantially less than the 20.7-fold and 13.7-fold

reduction we observed in the *ROLD* background after disruption of *BUD27* and *RPC17*,

respectively (Figure 3A). Furthermore, Hsf1 reporter activation increased in the Pol III-perturbed

backgrounds upon deletion of *UMP1*, *HSP104*, *HSC82*, and *SSA2* relative to the Pol III-

perturbed background strain (Figure 3-figure supplement 1B). Taken together, these data

indicate that Pol III impairment reduces *ROLD*-induced Hsf1 activation without generally

preventing Hsf1 activation in other genetic backgrounds.

22
Because Pol III transcribes target genes like 5S rRNA and tRNAs that play roles in translation

(Turowski and Tollervey, 2016), we wondered whether Pol III perturbation blocks *ROLD*-induced

Hsf1 activation by perturbing CAT tail synthesis. Pol III perturbation could block CAT tail

synthesis by preventing ribosome stalling, a necessary prerequisite for the RQC pathway to

initiate. We evaluated this possibility using a quantitative stalling reporter similar to one

previously developed (Juszkiewicz and Hegde, 2017; Sundaramoorthy et al., 2017) (Figure 3-

figure supplement 1C). *BUD27* deletion did not alleviate stalling induced by (CGN)12 (the

stalling sequence in RQCsub) in wt or *ROLD* strains relative to control (Figure 3-figure

supplement 1C). Thus, the effects of Pol III perturbation on translation should not confound our

analysis of RQCsub by preventing ribosomes translating RQCsub from stalling. When we

monitored RQCsub mobility by SDS-PAGE to directly observe any changes in CAT tails, we

observed that RQCsub expressed in the *ROLD bud27Δ* strain migrated as a higher molecular

weight smear compared to the CATylation-deficient *ltn1Δ rqc2Δ* strain and similar to the

CATylation-competent *ltn1Δ* strain (Figure 3B). Thus, Pol III perturbation leaves CATylation

intact.

12
Given its ability to mitigate Hsf1 hyperactivation in the *ROLD* strain (Figure 3A;Figure 3-figure

supplement 1A), we investigated whether Pol III perturbation could inhibit CAT tail aggregation

similarly to GuHCl. *ROLD* cells carrying hypomorphic *BUD27* or *RPC17* alleles formed RQCsub

inclusions less frequently (3.57% and 4.37% relative to 56.9% in *ROLD*), consistent with a

reduction in CAT tail aggregation (Figure 3C). Additionally, *BUD27* deletion in the *ROLD*

background increased the mobility of RQCsub by SDS-PAGE, reducing the proportion that

remained in the well and increasing the proportion that migrated into the resolving gel (Figure 3-

figure supplement 1D). Pol III perturbation thus serves as another tool, in addition to GuHCl, to

inhibit CAT tail aggregation.

22
Given that GuHCl solubilized CAT tails and rescued their degradation in *ROLD* cells (Figure

1C,G), we asked whether Pol III perturbation yields the same effect. To quantify CAT tail

degradation, we measured RQCsub$_{LONG}$ stability in *ROLD* and *ROLD bud27Δ* cells (Figure 3D).

The difference in RQCsub$_{LONG}$ stability (GFP:RFP) in each of these backgrounds with Hul5

(degradation intact) and without Hul5 (degradation blocked) defined the degree of degradation

(Sitron and Brandman, 2019). We validated this approach by demonstrating that the

proteasome inhibitor bortezomib did not significantly stabilize RQCsub$_{LONG}$ in *ROLD hul5Δ* and

ROLD bud27Δ hul5Δ strains (Figure 3-figure supplement 1E), indicating that *HUL5* deletion

blocked degradation in the *ROLD* and *ROLD bud27Δ* backgrounds. *HUL5* deletion stabilized

RQCsub$_{LONG}$ in *ROLD bud27Δ* more than in *ROLD* cells (33.7% compared to 13.6%; Figure

3E). In agreement with the effect of GuHCl (Figure 1G), these data demonstrate that limiting

CAT tail aggregation by perturbing Pol III restores degradation of CATylated proteins.

9
We then used Pol III perturbation to analyze how reducing CAT tail aggregation affects

proteostasis by determining its effects on growth in the presence of stress (Figure 3D). As

previously published, *BUD27* deletion slightly lowered growth rate at 30^{O}C and strongly

impaired growth at 37^{O}C in the wt background (Figure 3F) (Deplazes et al., 2009). We observed

similar effects of *BUD27* deletion in the *ROLD* background (Figure 3F). Thus, the sensitivity to

mild thermal stress caused by Pol III impairment renders this perturbation unable to rescue heat

sensitivity in the *ROLD* background. Despite causing a growth defect in wt cells, *BUD27*

deletion enabled faster growth of *ROLD* cells in the presence of cycloheximide (Figure 3F).

Unexpectedly, the growth rate of *ROLD bud27Δ* cells upon cycloheximide exposure was greater

than that of the *bud27Δ* strain (Figure 3F). Pol III perturbation, similarly to GuHCl, thereby

reduces cells' sensitivity to increased ribosome stalling in addition to inhibiting CAT tail

aggregation. These results suggest that improving CAT tail solubility can enhance adaptation to

ribosome stall-inducing stress.

23
Discussion

We have examined how CAT tail aggregation affects the cell's ability to cope when its burden of

incomplete proteins exceeds Ltn1's ability to ubiquitylate them. In cells with compromised Ltn1

function, we modulated the aggregation propensity of CAT tails to determine how aggregation

affects the cellular capacity to handle elevated levels of incomplete proteins. Aggregation

compromised degradation of CATylated proteins (Figure 1F) and diminished cellular fitness

during stress (Figure 2E), while inhibiting aggregation reversed each of these effects (Figure

1G;Figure 2E;Figure 3E,F). Our findings suggest that CAT tail aggregation is detrimental, rather

than adaptive, in cells with diminished Ltn1 function.

7
Since their discovery, multiple functions have been attributed to CAT tails. These include a role

in assisting Ltn1 activity (Kostova et al., 2017; Osuna et al., 2017; Sitron and Brandman, 2019)

and Ltn1-independent roles such as targeting Ltn1-evading RQC substrates for degradation

(Sitron and Brandman, 2019) or serving as aggregation-inducing post-translational modifications

(Yonashiro et al., 2016). We previously presented evidence supporting an adaptive, Ltn1-

independent function for CAT tails by showing that loss of CAT tails sensitized cells to increased

ribosome stalling (cycloheximide) specifically in the absence of Ltn1 (Sitron and Brandman,

2019). Here we show that this adaptive Ltn1-independent role is unlikely to be CAT tail-

mediated aggregation, as aggregation instead increased sensitivity to cycloheximide (Figure

2E;Figure 3F). Combined with the recent finding that CAT tail degron function is shared with

prokaryotic Ala tails (Lytvynenko et al., 2019), our results suggest that marking escaped RQC

substrates for degradation is the primary function of CAT tails.

20
Our findings elevate the question of why yeast and metazoan CAT tails have retained their

threonine content, and thereby aggregation propensity, through evolution. We propose two

models to explain this. In the first model, CAT tail aggregation facilitates adaptation to an as-of-

yet untested stress with evolutionary relevance. The tools we used here to control aggregation

may prove useful in probing this model by systematically assessing the fitness benefit of CAT

tail aggregation in diverse conditions. The second model proposes that the inclusion of

threonine in CAT tails facilitates a function of CAT tails other than aggregation. For instance,

threonine incorporation could produce more potent CAT tail degrons, assist in Ltn1-mediated

ubiquitylation, or aid in mounting a protective heat shock response. To preserve threonine

content, the fitness benefit from this threonine-mediated CAT tail function would need to

outweigh any fitness deficits arising from CAT tail aggregation. Given the recent discovery of

Rqc2 homologs that incorporate different amino acids into CAT tails (Lytvynenko et al., 2019;

Wu Z, Tantray I, Lim J, Chen S, Li Y, Davis Z, Sitron C, Dong J, Gispert S, Auburger G,

Brandman O, Bi X, Snyder M, Lu B., 2019), domain-swapping between these homologs to bias

CAT tail amino acid content may assist in determining the function of threonine in CAT tails in

future studies. Broadly, understanding how the variable amino acid composition of CAT tails has

evolved to meet physiological demands in different organisms may prove a fruitful area of

research.

12
We exploited *RQC2* overexpression and Pol III disruption as tools to genetically modify CAT tail

aggregation. While these proved useful in analyzing CAT tails in their aggregated and un-

aggregated states, the underlying mechanism behind these perturbations is not the focus of this

study (nor does it influence our conclusions) and remains unclear. *RQC2* overexpression has

been proposed to enhance aggregation because endogenous Rqc2 is limiting for CATylation

(Yonashiro et al., 2016). Under such a scenario, *RQC2* overexpression would enable

CATylation of more escaped RQC substrates that could then nucleate CAT tail aggregates.

Although pleiotropic (Turowski and Tollervey, 2016), Pol III perturbation could function in the

opposite manner. By limiting translation initiation (Deplazes et al., 2009), Pol III perturbation

could decrease the total amount of CATylated proteins in the cell and thereby disfavor

nucleation of CAT tail aggregation. Further studies will be needed to decipher the mechanisms

by which these perturbations alleviate CAT tail aggregation.

25
We report here that CAT tail aggregation modifies the toxicity associated with disrupted Ltn1

function. In mice, a hypomorphic *LTN1* allele induces a progressive neurodegeneration that

shares phenotypic similarities with amyotrophic lateral sclerosis (ALS) (Chu et al., 2009). It is

possible that degenerating neurons in these *LTN1*-hypomorphic mice harbor toxic CAT tail

aggregates, as has been observed in a *Drosophila* model of Parkinson Disease (Wu Z, Tantray

I, Lim J, Chen S, Li Y, Davis Z, Sitron C, Dong J, Gispert S, Auburger G, Brandman O, Bi X,

Snyder M, Lu B., 2019). As the connection between the phenotypes in these disease models

and human disease becomes more clear, future studies can focus on strategies to mitigate

toxicity associated with compromised Ltn1. Our study serves as a proof-of-principle that

disrupting CAT tail aggregation lessens this toxicity.

9
10 **Materials and Methods**
11

12 **Yeast Strains and Culturing**

13 The parental wild-type yeast strain used in this study was BY4741. All gene deletions, genomic

14 integrations, and plasmid transformations were done in this background using standard

15 methods. For a complete list of strains used, see Supplementary Table 1.

16
Yeast cultures were grown at 30^OC in YPD or synthetic defined (SD) media with appropriate

nutrient dropouts. SD media used for growth of yeast cultures contained: 2% w/v dextrose

(Thermo Fisher Scientific, Waltham, MA), 13.4 g/L Yeast Nitrogen Base without Amino Acids

(BD Biosciences, San Jose, CA), 0.03 g/L L-isoleucine (Sigma-Aldrich, St. Louis, MO), 0.15 g/L

L-valine (Sigma-Aldrich), 0.04 g/L adenine hemisulfate (Sigma-Aldrich), 0.02 g/L L-arginine

(Sigma-Aldrich), 0.03 g/L L-lysine (Sigma-Aldrich), 0.05 g/L L-phenylalanine (Sigma-Aldrich),

0.2 g/L L-Threonine (Sigma-Aldrich), 0.03 g/L L-tyrosine (Sigma-Aldrich), 0.018 g/L L-histidine

(Sigma-Aldrich), 0.09 g/L L-leucine (Sigma-Aldrich), 0.018 g/L L-methionine (Sigma-Aldrich),

0.036 g/L L-tryptophan (Sigma-Aldrich), and 0.018 g/L uracil (Sigma-Aldrich). Media additionally

contained 5 mM or 10 mM guanidinium hydrochloride (Sigma-Aldrich) where indicated and

bortezomib (LC Laboratories, Woburn, MA) treatment lasted 4 hours.

1
2 **Plasmids**

3 Plasmids were constructed using NEBuilder HiFi DNA Assembly Master Mix (New England

4 Biolabs, Ipswich, MA). All plasmids used in this study are described in Supplementary Table 2.

5
6 Two variants of RQCsub were used that had identical construction except for the region

7 following the stall: one had a fluorescent protein (RFP) and the other had a series of affinity

8 tags. The version without the post-stall RFP was used for microscopy and the version with the

9 post-stall RFP was used in immunoblots; the two exceptions to this are that the version without

10 the post-stall RFP was used in immunoblots in Figure 1C and Figure 1-figure supplement 1B.

11
12
13 **Flow cytometry**

14 Fluorescence was measured using a BD Accuri C6 flow cytometer (BD Biosciences). Three

15 independent yeast cultures were grown to log phase overnight in appropriate SD dropout media.

16 Data analysis was performed with MATLAB (MathWorks, Natick, MA).

17
18 For measurements of the Hsf1 activity reporter, raw fluorescence values were first normalized to

19 side scatter, log_2-transformed, and then the value obtained from the wt strain was subtracted

20 from each condition. These calculations are reported as log_2 fold wt in the figures.

21
For measurements of RQCsub$_{LONG}$, a detailed description of the analysis workflow can be

found in (Sitron and Brandman, 2019). Briefly, yeast expressing the plasmid were gated

according to above-background RFP fluorescence. Then, bleedthrough from the RFP channel

was calculated using an RFP-only control strain and subtracted from signal in the GFP channel.

The resultant bleedthrough-corrected GFP:RFP values were normalized to the corresponding

untreated *hul5Δ* strain for each background. To quantify the change in GFP:RFP upon *HUL5*

deletion, the normalized value from the *HUL5* condition was subtracted from that of the *hul5Δ*

condition.

For the stalling reporter, the analysis workflow matched that of RQCsub$_{LONG}$ with the order of

the fluorophores being reversed, i.e. plasmid-expressing yeast selected based on GFP,

bleedthrough from GFP subtracted from RFP. Where indicated, corrected RFP:GFP values

were normalized to a non-stalling reporter that contained a non-stalling sequence encoding

serine and threonine (Sitron et al., 2017) in place of the stalling CGN codons.

7
Statistical Analysis

A paired t-test (using the "ttest" function in MATLAB) was used to assess whether mean

measurements differed between two different conditions, using the null hypothesis: $\mu_{condition\ 1}$ =

$\mu_{condition\ 2}$. To calculate the s.e.m. for Δ stability measurements (Fig. 1F,G;Fig. 3E), a

propagation of error formula was used: $SEM_{hul5\Delta\ -\ HUL5} = sqrt(SEM_{HUL5}^2 + SEM_{hul5\Delta}^2)$. To

analyze the significance of Δ stability measurements, a t-test for particular contrast was

performed using the "lm" and "linearHypothesis" functions in R (R Foundation for Statistical

Computing). The null hypothesis for this test was: $\mu_{hul5\Delta+\ condition\ 1} - \mu_{HUL5\ +\ condition\ 1} = \mu_{hul5\Delta+}$

$_{condition\ 2} - \mu_{HUL5\ +\ condition\ 2}$.

Immunoblots

For whole-cell immunoblots, 0.375/OD600 x mL of yeast culture (OD600 = 0.4-0.8) grown

overnight were pelleted and resuspended then boiled for 5 min at 95^{O}C in 15 µL 4x NuPage

LDS Sample Buffer (Thermo Fisher Scientific) with 5% β-mercaptoethanol.

22
For SDS-PAGE, samples were loaded into a NuPAGE Novex 4-12% Bis-Tris 1.5 mm protein gel

(Thermo Fisher Scientific) and run in MOPS buffer. Gels were semi-dry transferred to 0.45 µm

nitrocellulose membranes (Thermo Fisher Scientific) using the Trans-Blot Turbo system (Bio-

Rad, Hercules, CA). Gels were wet transferred to .45 µm nitrocellulose membranes (Thermo

Fisher Scientific) in the Trans-Blot Cell (Bio-Rad) in Towbin Buffer or semi-dry transferred using

a Trans-Blot Turbo (Bio-Rad).

3
Membranes were blocked in TBST with 5% fat-free milk (Safeway, Pleasanton, CA) for 1 hour,

incubated overnight at 4°C or at room temperature for 4 hrs in one of the following primary

antibodies: 1:3000 mouse α-GFP (MA5-15256, Thermo Fisher Scientific), 1:3000 rabbit α-

hexokinase (H2035-01, US Biological, Salem, MA), or 1:1000 rabbit α-RFP (AB233, Evrogen,

Moscow, Russia). The following secondary antibodies were then used at 1:5000 dilution: IRDye

800CW donkey anti-mouse, IRDye 800CW goat anti-rabbit, IRDye 680RD goat anti-rabbit, or

IRDye 680RD goat anti-mouse (LiCor Biosciences, Lincoln, NE). Blots were scanned on a Licor

Odyssey (LiCor Biosciences).

12
Lysate Preparation and Immunoprecipitation (IP)

Yeast were grown in SD media and harvested at OD600 0.8 to 1.0. Cells were harvested using

either 1) centrifugation followed by resuspension in IP buffer (100 mM KOAc, 10 mM MgCl2, 25

mM HEPES-KOH pH 7.4) and freezing of cell droplets in liquid nitrogen; or 2) vacuum filtration

and flash freezing in liquid nitrogen. Frozen yeast were cryogenically pulverized into powder

using a Freezer/Mill (SPEX SamplePrep, Metuchen, NJ) and stored at -80°C.

19
Frozen yeast powder was thawed and solubilized at 4°C in IP buffer supplemented with Pierce

Protease Inhibitor Tablets, EDTA-free (Thermo Fisher Scientific). Crude lysate was clarified by a

single spin at 5000 x g, incubated with GFP-Trap_A (ChromoTek, Planegg-Martinsried,

Germany) resin for 1 hour at 4°C with rotation, then washed 10 times in IP buffer. For

immunoblotting, washed GFP-Trap resin was boiled in 4X NuPAGE LDS Sample Buffer

(Thermo Fisher Scientific) with 5% β-mercaptoethanol and analyzed as described above in

"Immunoblots."

27
TEV Protease Digestion

GFP IP was performed as described above, but instead of immediate elution, the rein was

equilibrated in IP buffer with 1 mM DTT and incubated overnight at 4°C with 5 μL ProTEV Plus

(Promega, Sunnyvale, CA) in 100 μL total reaction volume. The resin was then washed with

fresh IP buffer with 1 mM DTT and boiled in 4X NuPAGE LDS Sample Buffer (Thermo Fisher

Scientific) with 5% β-mercaptoethanol and analyzed as described above.

6
Fluorescence Microscopy

Live cell fluorescence imaging was performed using two set-ups: 1) a Nikon Eclipse Ti-E

inverted fluorescence microscope (Nikon, Melville, NY) using a 100X/1.4NA oil objective lens

and μManager (Edelstein et al., 2010), and 2) a Nikon Eclipse 80i microscope using a

100X/1.4NA oil objective lens and Metamorph (Molecular Devices, San Jose, CA). Yeast were

grown in SD media to log phase, centrifuged briefly, resuspended in 5 μL fresh SD media and

mounted on a clean coverslip and slide. Images were prepared and analyzed with ImageJ

(NIH). For quantification, images were randomly acquired until N > 300 cells. Independent

cultures were imaged on different days and error represents standard deviation of 3

independent cultures.

17
Yeast spot assay

Yeast were grown to log-phase in YPD, then diluted to OD600 = 0.1. 200 μL of this diluted

culture was diluted 1:10 into YPD four times in a sterile 96-well plate (Greiner Bio-one,

Kremsmünster, Austria). 5 μL of the serial dilutions were transferred from the 96-well plate onto

YPD agar plates with or without additional 5 mM guanidinium hydrochloride (Sigma-Aldrich)

and/or 50 ng/mL cycloheximide (Sigma-Aldrich). Plates were then incubated at 30^OC or 37^OC.

Plates grown at 30^OC with or without 5 mM guanidinium hydrochloride as well as 37^OC plates

were imaged after two days. Plates containing 50 ng/mL cycloheximide with or without 5 mM

guanidinium hydrochloride grown at 30^OC were imaged after three days. Plates with 5 mM

guanidinium hydrochloride grown at 37^OC were imaged after six days.

Acknowledgements

We thank D. Pincus, D. Jarosz, R. Kopito, B. Lu, Z. Wu, E.P. Geiduschek, and the members of

the Brandman lab for helpful suggestions. We thank the Theriot and Straight labs (Stanford

University) for the use of their microscopes. We thank J. Weissman and D. Wong for facilitating

the screen for suppressors of RQC-mediated Hsf1 activation. This work was supported by

Stanford University (O.B.), the US National Institutes of Health (grant No. R01GM115968 to

O.B.), and the National Institute of General Medical Sciences of the US National Institutes of

Health (grant No. T32GM007276 to C.S.S.).

Author Contributions

12 O.B., C.S.S., and J.P. conceived of the study and designed the experiments. C.S.S., J.P., and

13 J.M.G. performed the experiments. O.B., C.S.S., and J.H.P. analyzed and interpreted the data.

14 O.B., C.S.S., and J.H.P. wrote the manuscript. O.B. supervised the study.

Competing Interests

17 The authors declare no competing interests.

References

Abo T, Ueda K, Sunohara T, Ogawa K, Aiba H. 2002. SsrA-mediated protein tagging in the
presence of miscoding drugs and its physiological role in Escherichia coli. *Genes to Cells*.
doi:10.1046/j.1365-2443.2002.00549.x
Åkerfelt M, Trouillet D, Mezger V, Sistonen LEA. 2007. Heat shock factors at a crossroad
between stress and development. *Ann N Y Acad Sci* **1113**:15–27.
Bengtson MH, Joazeiro CAP. 2010. Role of a ribosome-associated E3 ubiquitin ligase in protein
quality control. *Nature* **467**:470–473.
Brandman O, Hegde RS. 2016. Ribosome-associated protein quality control. *Nat Struct Mol Biol*
23:7–15.
Brandman O, Stewart-Ornstein J, Wong D, Larson A, Williams CC, Li G-W, Zhou S, King D,
Shen PS, Weibezahn J, Dunn JG, Rouskin S, Inada T, Frost A, Weissman JS. 2012. A
ribosome-bound quality control complex triggers degradation of nascent peptides and
signals translation stress. *Cell* **151**:1042–1054.

Byrne LJ, Cox BS, Cole DJ, Ridout MS, Morgan BJT, Tuite MF. 2007. Cell division is essential
for elimination of the yeast [PSI+] prion by guanidine hydrochloride. *Proc Natl Acad Sci U S*
A **104**:11688–11693.
Chaudhuri RR, Allen AG, Owen PJ, Shalom G, Stone K, Harrison M, Burgis TA, Lockyer M,
Garcia-Lara J, Foster SJ, Pleasance SJ, Peters SE, Maskell DJ, Charles IG. 2009.
Comprehensive identification of essential Staphylococcus aureus genes using Transposon-
Mediated Differential Hybridisation (TMDH). *BMC Genomics* **10**:291.
Choe Y-J, Park S-H, Hassemer T, Körner R, Vincenz-Donnelly L, Hayer-Hartl M, Hartl FU. 2016.
Failure of RQC machinery causes protein aggregation and proteotoxic stress. *Nature*
531:191–195.
Chu J, Hong NA, Masuda CA, Jenkins BV, Nelms KA, Goodnow CC, Glynne RJ, Wu H, Masliah
E, Joazeiro CAP, Kay SA. 2009. A mouse forward genetics screen identifies LISTERIN as
an E3 ubiquitin ligase involved in neurodegeneration. *Proc Natl Acad Sci U S A* **106**:2097–
2103.
Defenouillère Q, Fromont-Racine M. 2017. The ribosome-bound quality control complex: from
aberrant peptide clearance to proteostasis maintenance. *Current Genetics*.
doi:10.1007/s00294-017-0708-5
Defenouillère Q, Yao Y, Mouaikel J, Namane A, Galopier A, Decourty L, Doyen A, Malabat C,
Saveanu C, Jacquier A, Fromont-Racine M. 2013. Cdc48-associated complex bound to 60S
particles is required for the clearance of aberrant translation products. *Proc Natl Acad Sci U*
S A **110**:5046–5051.
Defenouillère Q, Zhang E, Namane A, Mouaikel J, Jacquier A, Fromont-Racine M. 2016. Rqc1
and Ltn1 prevent CAT-tail induced protein aggregation by efficient recruitment of Cdc48 on
stalled 60S subunits. *J Biol Chem* **291**:12245–12253.
Deplazes A, Möckli N, Luke B, Auerbach D, Peter M. 2009. Yeast Uri1p promotes translation
initiation and may provide a link to cotranslational quality control. *EMBO J* **28**:1429–1441.
Donnelly MLL, Ryan MD, Mehrotra A, Gani D, Hughes LE, Luke G, Li X. 2001. Analysis of the
aphthovirus 2A/2B polyprotein "cleavage" mechanism indicates not a proteolytic reaction,
but a novel translational effect: a putative ribosomal "skip." *Journal of General Virology*.
doi:10.1099/0022-1317-82-5-1013
Eaglestone SS, Ruddock LW, Cox BS, Tuite MF. 2000. Guanidine hydrochloride blocks a critical
step in the propagation of the prion-like determinant [PSI+] of Saccharomyces cerevisiae.
Proc Natl Acad Sci U S A **97**:240–244.
Edelstein A, Amodaj N, Hoover K, Vale R, Stuurman N. 2010. Computer Control of Microscopes
Using µManager. *Current Protocols in Molecular Biology*.
doi:10.1002/0471142727.mb1420s92
Glass JI, Assad-Garcia N, Alperovich N, Yooseph S, Lewis MR, Maruf M, Hutchison CA 3rd,
Smith HO, Venter JC. 2006. Essential genes of a minimal bacterium. *Proc Natl Acad Sci U S*
A **103**:425–430.
Ikeuchi K, Izawa T, Inada T. 2018. Recent Progress on the Molecular Mechanism of Quality
Controls Induced by Ribosome Stalling. *Front Genet* **9**:743.
Ikeuchi K, Tesina P, Matsuo Y, Sugiyama T, Cheng J, Saeki Y, Tanaka K, Becker T, Beckmann
R, Inada T. 2019. Collided ribosomes form a unique structural interface to induce Hel2–
driven quality control pathways. *EMBO J* **38**:e100276.
Ishimura R, Nagy G, Dotu I, Zhou H, -L. Yang X, Schimmel P, Senju S, Nishimura Y, Chuang JH,
Ackerman SL. 2014. Ribosome stalling induced by mutation of a CNS-specific tRNA causes
neurodegeneration. *Science* **345**:455–459.
Joazeiro CAP. 2019. Mechanisms and functions of ribosome-associated protein quality control.
Nat Rev Mol Cell Biol **20**:368–383.
Julio SM, Heithoff DM, Mahan MJ. 2000. ssrA (tmRNA) Plays a Role inSalmonella enterica
Serovar Typhimurium Pathogenesis. *J Bacteriol* **182**:1558–1563.

Juszkiewicz S, Chandrasekaran V, Lin Z, Kraatz S, Ramakrishnan V, Hegde RS. 2018. ZNF598
Is a Quality Control Sensor of Collided Ribosomes. *Mol Cell* **72**:469–481.e7.
Juszkiewicz S, Hegde RS. 2017. Initiation of Quality Control during Poly(A) Translation Requires
Site-Specific Ribosome Ubiquitination. *Mol Cell* **65**:743–750.e4.
Kostova KK, Hickey KL, Osuna BA, Hussmann JA, Frost A, Weinberg DE, Weissman JS. 2017.
CAT-tailing as a fail-safe mechanism for efficient degradation of stalled nascent
polypeptides. *Science* **357**:414–417.
Letzring DP, Dean KM, Grayhack EJ. 2010. Control of translation efficiency in yeast by codon–
anticodon interactions. *RNA* **16**:2516–2528.
Lytvynenko I, Paternoga H, Thrun A, Balke A, Müller TA, Chiang CH, Nagler K, Tsaprailis G,
Anders S, Bischofs I, Maupin-Furlow JA, Spahn CMT, Joazeiro CAP. 2019. Alanine Tails
Signal Proteolysis in Bacterial Ribosome-Associated Quality Control. *Cell* **178**:76–90.e22.
Lyumkis D, dos Passos DO, Tahara EB, Webb K, Bennett EJ, Vinterbo S, Potter CS, Carragher
B, Joazeiro CAP. 2014. Structural basis for translational surveillance by the large ribosomal
subunit-associated protein quality control complex. *Proceedings of the National Academy of*
Sciences **111**:15981–15986.
Mann B, van Opijnen T, Wang J, Obert C, Wang Y-D, Carter R, McGoldrick DJ, Ridout G, Camilli
A, Tuomanen EI, Rosch JW. 2012. Control of virulence by small RNAs in Streptococcus
pneumoniae. *PLoS Pathog* **8**:e1002788.
Matsuo Y, Ikeuchi K, Saeki Y, Iwasaki S, Schmidt C, Udagawa T, Sato F, Tsuchiya H, Becker T,
Tanaka K, Ingolia NT, Beckmann R, Inada T. 2017. Ubiquitination of stalled ribosome
triggers ribosome-associated quality control. *Nat Commun* **8**:159.
Mirón-García MC, Garrido-Godino AI, García-Molinero V, Hernández-Torres F, Rodríguez-
Navarro S, Navarro F. 2013. The prefoldin bud27 mediates the assembly of the eukaryotic
RNA polymerases in an rpb5-dependent manner. *PLoS Genet* **9**:e1003297.
Moore SD, Sauer RT. 2007. The tmRNA system for translational surveillance and ribosome
rescue. *Annu Rev Biochem* **76**:101–124.
Moore SD, Sauer RT. 2005. Ribosome rescue: tmRNA tagging activity and capacity in
Escherichia coli. *Mol Microbiol* **58**:456–466.
Morimoto RI. 2011. The heat shock response: systems biology of proteotoxic stress in aging and
disease. *Cold Spring Harb Symp Quant Biol* **76**:91–99.
Ness F, Ferreira P, Cox BS, Tuite MF. 2002. Guanidine Hydrochloride Inhibits the Generation of
Prion "Seeds" but Not Prion Protein Aggregation in Yeast. *Mol Cell Biol* **22**:5593–5605.
Osuna BA, Howard CJ, Kc S, Frost A, Weinberg DE. 2017. In vitro analysis of RQC activities
provides insights into the mechanism and function of CAT tailing. *Elife* e27949.
Oussenko IA, Abe T, Ujiie H, Muto A, Bechhofer DH. 2005. Participation of 3′-to-5′
exoribonucleases in the turnover of Bacillus subtilis mRNA. *J Bacteriol* **187**:2758–2767.
Pisareva VP, Skabkin MA, Hellen CUT, Pestova TV, Pisarev AV. 2011. Dissociation by Pelota,
Hbs1 and ABCE1 of mammalian vacant 80S ribosomes and stalled elongation complexes.
EMBO J **30**:1804–1817.
Shao S, von der Malsburg K, Hegde RS. 2013. Listerin-dependent nascent protein ubiquitination
relies on ribosome subunit dissociation. *Mol Cell* **50**:637–648.
Shen PS, Park J, Qin Y, Li X, Parsawar K, Larson MH, Cox J, Cheng Y, Lambowitz AM,
Weissman JS, Brandman O, Frost A. 2015. Protein synthesis. Rqc2p and 60S ribosomal
subunits mediate mRNA-independent elongation of nascent chains. *Science* **347**:75–78.
Shoemaker CJ, Eyler DE, Green R. 2010. Dom34:Hbs1 promotes subunit dissociation and
peptidyl-tRNA drop-off to initiate no-go decay. *Science* **330**:369–372.
Simms CL, Yan LL, Zaher HS. 2017. Ribosome Collision Is Critical for Quality Control during No-
Go Decay. *Mol Cell* **68**:361–373.e5.
Sitron CS, Brandman O. 2019. CAT tails drive degradation of stalled polypeptides on and off the
ribosome. *Nat Struct Mol Biol* **26**:450–459.

Sitron CS, Park JH, Brandman O. 2017. Asc1, Hel2, and Slh1 couple translation arrest to
        nascent chain degradation. *RNA* **23**:798–810.
3   Sorger PK, Pelham HR. 1987. Purification and characterization of a heat-shock element binding
        protein from yeast. *EMBO J* **6**:3035–3041.
5   Sundaramoorthy E, Leonard M, Mak R, Liao J, Fulzele A, Bennett EJ. 2017. ZNF598 and
        RACK1 Regulate Mammalian Ribosome-Associated Quality Control Function by Mediating
        Regulatory 40S Ribosomal Ubiquitylation. *Mol Cell* **65**:751–760.e4.
8   Szymczak AL, Vignali DAA. 2005. Development of 2A peptide-based strategies in the design of
        multicistronic vectors. *Expert Opin Biol Ther* **5**:627–638.
10 Tsuboi T, Kuroha K, Kudo K, Makino S, Inoue E, Kashima I, Inada T. 2012. Dom34:hbs1 plays a
        general role in quality-control systems by dissociation of a stalled ribosome at the 3' end of
        aberrant mRNA. *Mol Cell* **46**:518–529.
13 Turowski TW, Tollervey D. 2016. Transcription by RNA polymerase III: insights into mechanism
        and regulation. *Biochem Soc Trans* **44**:1367–1375.
15 Verma R, Oania RS, Kolawa NJ, Deshaies RJ. 2013. Cdc48/p97 promotes degradation of
        aberrant nascent polypeptides bound to the ribosome. *Elife* **2**:e00308.
17 Wu Z, Tantray I, Lim J, Chen S, Li Y, Davis Z, Sitron C, Dong J, Gispert S, Auburger G,
        Brandman O, Bi X, Snyder M, Lu B. 2019. MISTERMINATE Mechanistically Links
        Mitochondrial Dysfunction with Proteostasis Failure. *Mol Cell* **75**:835–848.
20 Yan Z, Costanzo M, Heisler LE, Paw J, Kaper F, Andrews BJ, Boone C, Giaever G, Nislow C.
        2008. Yeast Barcoders: a chemogenomic application of a universal donor-strain collection
        carrying bar-code identifiers. *Nature Methods*. doi:10.1038/nmeth.1231
23 Yonashiro R, Tahara EB, Bengtson MH, Khokhrina M, Lorenz H, Chen K-C, Kigoshi-Tansho Y,
        Savas JN, Yates JR, Kay SA, Craig EA, Mogk A, Bukau B, Joazeiro CAP. 2016. The
        Rqc2/Tae2 subunit of the ribosome-associated quality control (RQC) complex marks
        ribosome-stalled nascent polypeptide chains for aggregation. *Elife* **5**:e11794.

Figure 1. Aggregation compromises CAT tail degradation. (**A**) Left, whole cell immunoblots (IBs) of lysates containing the model RQC substrate RQCsub (schematic above). Right, IB of RQCsub immunoprecipitated (IPed) from *RQC2* overexpression and *LTN1* deletion (*ROLD*) lysate. Arrow indicates molecular weight of RQCsub without CAT tails. Asterisk denotes the full-length RQCsub protein product, produced when ribosomes translate through the stall sequence (region past the stall not pictured in schematic). GFP, green fluorescent protein. OE, overexpression. (**B**) Fluorescence microscopy of cells expressing RQCsub with percentages of cells containing observable GFP inclusions reported below. (**C**) IB of RQCsub expressed in *ROLD* cultures grown in guanidinium hydrochloride (GuHCl). Arrows as in **A**. (**D**) Schematic of experimental approach for measuring the effect of aggregation of CAT tail-mediated degradation. (**E**) Left, schematic of expression-controlled model RQC substrate RQCsub_LONG. Center, definitions of "stability," "Δ stability," and "CAT tail-mediated degradation." (**F**) and (**G**) Δ stability measurements of RQCsub_LONG expressed in *ltn1Δ* and *ROLD* cells after *HUL5* deletion to assess CAT tail-mediated degradation. P-values above bars are derived from t-tests for particular contrast, measuring how significantly different *HUL5* deletion-induced stabilization is between either *ltn1Δ* and *ROLD* backgrounds in **F** or between different GuHCl concentrations in **G**. Error bars represent the standard error of the mean (s.e.m.).

Figure 1-figure supplement 1. *RQC2* overexpression and guanidinium hydrochloride are tools to control CAT tail aggregation. (**A**) Flow cytometry of cells expressing Rqc2-RFP to assess the degree of Rqc2 overproduction by the *TDH3* promoter. Error bars represent s.e.m. from three independent cultures. (**B**) Immunoblot (IB) of RQCsub expressed in *ltn1Δ* compared to *ROLD*. (**C**) IB of RQCsub immunoprecipitated (IPed) from *ltn1Δ* and *ROLD* lysates with and without tobacco etch virus (TEV) protease treatment. Arrow denotes molecular weight of non-CATylated RQCsub. Asterisk indicates the full-length RQCsub protein product, produced when ribosomes translate through the stall sequence (region past the stall not pictured in schematic). (**D**) Microscopy of RQCsub expressed in *ROLD* grown in various concentrations of guanidinium hydrochloride (GuHCl).

Figure 1-figure supplement 2. Supporting data for the effect of aggregation of CAT tail-mediated degradation. (**A**) Flow cytometry of cells expressing RQCsub$_{LONG}$. Error bars indicate s.e.m. from three independent cultures. (**B**) and (**C**) Additional data to support Figure 1F,G. Stability measurements of RQCsub$_{LONG}$ expressed in *ROLD* cells grown in indicated GuHCl concentrations with additional bortezomib treatment to inhibit the proteasome and *HUL5* deletion to measure CAT tail-mediated degradation. Error bars as in **A**. P-values are indicated above bars. Thick lines indicate paired t-tests, probing the significance of bortezomib (btz)-induced stabilization. Thin lines denote t-tests for particular contrast, measuring how significantly different *HUL5* deletion-induced stabilization is under different conditions.

Figure 2. Aggregated CAT tails are proteotoxic. (**A**) Schematic of experimental approach for assessing how aggregation influences the effect of CAT tails on proteostasis. (**B**), (**C**), and (**D**) Flow cytometry of cells containing an integrated reporter for Hsf1 activation (schematic above). Error bars represent s.e.m. from three independent cultures. P-values from indicated paired t-tests are indicated by lines between bars. HSE, heat shock element. (**E**) Spot assay of strains grown at 30°C, 37°C, and in 50 ng/ml cycloheximide with and without the presence of 5 mM GuHCl. Plate numbers indicated below images.

Figure 3. Pol III perturbation inhibits CAT tail aggregation, increases degron activity, and restores proteostasis. (**A**) Flow cytometry of cells containing an integrated reporter for Hsf1 activation. Error bars indicate s.e.m. from three independent cultures. P-values from paired t-tests indicated above bars. (**B**) IB of RQCsub expressed in indicated strains. (**C**) Fluorescence microscopy of cells expressing RQCsub. Percentages of cells containing GFP inclusions reported below micrographs. (**D**) Schematic of experimental strategy for measuring how disruption of aggregation modulates CAT tail-mediated degradation and CAT tails' effect on proteostasis. (**E**) Δ stability measurements of RQCsub_LONG expressed in indicated strains after *HUL5* deletion to quantify CAT tail-mediated degradation. Error bars as in **A**. P-value indicates the result of a t-test for particular contrast. (**F**) Spot assay of cells grown under indicated conditions.

Figure 3-figure supplement 1. Effects of Pol III perturbation on Hsf1 activation, stalling, and CAT tail degron activity. (**A**) Flow cytometry of cells containing an integrated reporter for Hsf1 activation. Error bars indicate s.e.m. from three independent cultures. P-values from paired t-tests indicated above bars. (**B**) Flow cytometry of Pol III-perturbed cells containing an integrated reporter for Hsf1 activation. These data are also contained in Figure 3A, but are reordered here to simplify comparisons within two Pol III-perturbed genetic backgrounds. Error bars as in **A**. (**C**) Left, schematic of a stalling reporter similar to that used in Juszkiewicz and Hegde 2017 and Sundaramoorthy et al. 2017. Right, schematic of stalling reporter with the same (CGN)12 stalling sequence contained in RQCsub or a non-stalling (Ser-Thr)6 sequence. Error bars as in **A**. (**D**) IB of RQCsub expressed in *ROLD* compared to *ROLD bud27Δ*. (**E**) Additional data to support Figure 3E. Stability measurements of RQCsub_LONG expressed in indicated strains with bortezomib (btz) treatment to inhibit the proteasome and *HUL5* deletion to block CAT tail-mediated degradation. Error bars indicate s.e.m. from three independent cultures. P-values are given above bars. Results of paired t-tests measuring the significance of bortezomib-induced stabilization are indicated with thick lines. The result of a t-test for particular contrast is indicated with thin lines; this assesses how significantly different *HUL5* deletion-induced stabilization is in *ROLD* compared to *ROLD bud27Δ*.

Supplementary Table 1. Yeast strains used in this study.

Strain Number	Description
yOB199	BY4741
yOB259	BY4741 *ltn1Δ::kanMX*
yOB454	BY4741 *rqc2Δ::kanMX*
yOBj004	BY4741 *ltn1Δ::kanMX rqc2Δ::natMX*
yOB518	BY4741 *pRQC2::HIS3-pTDH3*
yOB551	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX*
yOB255	BY4741 *ura3::HSE-EmGFP*
yOB549	BY4741 *ura3::HSE-EmGFP rqc2Δ::kanMX*
yOB550	BY4741 *ura3::HSE-EmGFP ltn1Δ::kanMX*
yOBj245	BY4741 *ura3::HSE-EmGFP rqc2Δ::kanMX ltn1Δ::natMX*
yOB515	BY4741 *ura3::HSE-EmGFP pRQC2::HIS3-pTDH3*
yOBj185	BY4741 *ura3::HSE-EmGFP pRQC2::HIS3-pTDH3 ltn1Δ::natMX*
yOB1118	BY4741 *RQC2-WT::LEU ltn1Δ::hphMX*
yOB2825	BY4741 *RQC2-WT::LEU ltn1Δ::hphMX hul5Δ::natMX*
yOB3812	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX hul5Δ::hphMX*
yOBj482	BY4741 *ura3::HSE-EmGFP ump1Δ::natMX*
yOBj538	BY4741 *ura3::HSE-EmGFP ump1Δ::natMX bud27Δ::kanMX*
yOB738	BY4741 *ura3::HSE-EmGFP ump1Δ::natMX rpc17-DAmP::natMX*
yOB736	BY4741 *ura3::HSE-EmGFP hsp104Δ::natMX*
yOB808	BY4741 *ura3::HSE-EmGFP hsp104Δ::natMX bud27Δ::kanMX*
yOB809	BY4741 *ura3::HSE-EmGFP hsp104Δ::natMX rpc17-DAmP::kanMX*
yOBj542	BY4741 *ura3::HSE-EmGFP hsc82Δ::natMX*
yOB818	BY4741 *ura3::HSE-EmGFP hsc82Δ::natMX bud27Δ::kanMX*
yOB722	BY4741 *ura3::HSE-EmGFP hsc82Δ::natMX rpc17-DAmP::kanMX*
yOBj563	BY4741 *ura3::HSE-EmGFP ssa2Δ::natMX*
yOB819	BY4741 *ura3::HSE-EmGFP ssa2Δ::natMX bud27Δ::kanMX*
yOB725	BY4741 *ura3::HSE-EmGFP ssa2Δ::natMX rpc17-DAmP::kanMX*
yOBj536	BY4741 *ura3::HSE-EmGFP bud27Δ::kanMX*
yOB710	BY4741 *ura3::HSE-EmGFP rpc17-DAmP::kanMX*
yOBj495	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX bud27Δ::kanMX*
yOBj556	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX rpc17-DAmP::kanMX*
yOBj536	BY4741 *ura3::HSE-EmGFP bud27Δ::kanMX*
yOBj185	BY4741 *ura3::HSE-EmGFP pRQC2::HIS3-pTDH3 ltn1Δ::natMX bud27Δ::kanMX*
yOBj556	BY4741 *ura3::HSE-EmGFP pRQC2::HIS3-pTDH3 ltn1Δ::natMX rpc17-DAmP::kanMX*
yOBj626	BY4741 *ura3::HSE-EmGFP pRQC2::HIS3-pTDH3 ltn1Δ::natMX rpc160-DAmP::kanMX*
yOB3812	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX hul5Δ::hphMX bud27Δ::kanMX*

Supplementary Table 2. Plasmids used in this study.

Plasmid Name	Description	Selection Marker
RQCsub	CEN/ARS pTDH3-GFP_TEV_R12_RFP	*URA3*
RQCsub (no RFP)	CEN/ARS pTDH3-GFP_TEV_R12_3xHA_3xMYC_3xFLAG	*URA3*
RQCsubLONG	CEN/ARS pTDH3-mKate2-2xT2A-GFP_TEVlonglinker_R12-3xMYC	*URA3*
pRS315	CEN/ARS empty vector	*LEU2*
pJP014	CEN/ARS pRQC2-RQC2WT	*LEU2*
pJP030	CEN/ARS pTDH3-rqc2aaa (D9A, D98A, R99A)	*LEU2*
pJP024	CEN/ARS pRQC2-RQC2WT-mKate2	*LEU2*
pJP055	CEN/ARS pTDH3-RQC2WT-mKate2	*LEU2*
Stalling Reporter	CEN/ARS pTDH3-GFP-2xT2A-TEV_R12-2xT2A-mKate2	*URA3*
Non-stalling Reporter	CEN/ARS pTDH3-GFP-2xT2A-TEV_6xST-2xT2A-mKate2	*URA3*