1   Rqc2 levels alter CAT tail composition, function, and toxicity in *S. cerevisiae*

2
Cole S. Sitron[1,*], Joseph H. Park[1,2,*], Jenna M. Giafaglione[1,3], and Onn Brandman[1,†]

[1]Department of Biochemistry, Stanford University, Stanford CA, 94305. [2]Department of

Chemical & Systems Biology, Stanford University, Stanford CA, 94305. [3]Current address:

Molecular Biology Interdepartmental Program, University of California Los Angeles, Los Angeles

CA, 90095. [*]These authors contributed equally to the work. [†]Corresponding author, to whom

correspondence and material requests should be addressed: onn@stanford.edu.

11
12

Abstract

The Ribosome-associated Quality Control (RQC) pathway targets incomplete polypeptides from stalled translation for degradation. The primary route of degradation involves ubiquitylation of ribosome-tethered incomplete polypeptides by the E3 ligase Ltn1. To safeguard cells from Ltn1 failure, the protein Rqc2 directs the ribosome to append incomplete polypeptides with carboxy-terminal alanine and threonine residues (CAT tails) that act as degrons, marking RQC-evading polypeptides for degradation. Ltn1 disruption leads to neurodegeneration in animal models yet how this becomes toxic to cells is unclear. We show here that increased Rqc2 levels alter CAT tail composition and exacerbate the toxicity of Ltn1 failure by driving CAT tail aggregation, which inhibits CAT tail degron activity and disrupts proteostasis. Guanidinium hydrochloride exposure or RNA Polymerase III perturbation reverses these Rqc2-induced effects. Our work demonstrates that increased Rqc2 levels convert CAT tails from degrons into toxic aggregates and that pharmacologic or genetic interventions can reverse this process.

Introduction

Cells must continuously surveil their proteomes to protect themselves from defective proteins. Disruption of this process causes a buildup of defective proteins, a hallmark of many neurodegenerative diseases (Hartl, 2017; Ross and Poirier, 2004). Understanding how compromised protein surveillance can become toxic to cells has been a fundamental goal of the protein quality control field.

The earliest stage of surveillance occurs when cells monitor protein synthesis to detect failed rounds of translation and degrade potentially toxic incomplete polypeptides (Brandman and Hegde, 2016; Defenouillère and Fromont-Racine, 2017; Ikeuchi et al., 2018; Joazeiro, 2019). Protein synthesis can fail to complete due to inefficient messenger RNA (mRNA) decoding (Dimitrova et al., 2009; Doma and Parker, 2006; Gamble et al., 2016; Letzring et al., 2010;

Meaux and Van Hoof, 2006), causing ribosomes translating the same mRNA to stall and

subsequently collide with each other (Ikeuchi et al., 2019; Juszkiewicz et al., 2018; Simms et al.,

2017). These collisions trigger a process called Ribosome-associated Quality Control (RQC),

which recycles these ribosomes and targets their attached incomplete polypeptides (RQC

substrates) for degradation (Bengtson and Joazeiro, 2010; Brandman et al., 2012; Defenouillère

et al., 2013; Shao et al., 2013; Verma et al., 2013). RQC begins when a set of proteins detect

the collided ribosomes and separate them into small (40S) and large (60S) subunits, leaving the

incomplete polypeptide tethered to the 60S subunit (Juszkiewicz and Hegde, 2017; Lyumkis et

al., 2014; Matsuo et al., 2017; Pisareva et al., 2011; Shao et al., 2013; Shoemaker et al., 2010;

Sitron et al., 2017; Sundaramoorthy et al., 2017; Tsuboi et al., 2012).

11
The newly separated 60S subunit recruits a complex of proteins that collaborate to ubiquitylate

the incomplete polypeptide and facilitate its degradation by the proteasome (Bengtson and

Joazeiro, 2010; Brandman et al., 2012; Defenouillère et al., 2013; Lyumkis et al., 2014; Shao et

al., 2013; Verma et al., 2013). A member of this complex, the RING E3 ubiquitin ligase Ltn1,

ubiquitylates the incomplete polypeptide while it is still attached to the ribosome (Bengtson and

Joazeiro, 2010; Brandman et al., 2012; Defenouillère et al., 2013; Lyumkis et al., 2014; Shao et

al., 2013; Shao and Hegde, 2014; Verma et al., 2013). Rqc2 stabilizes Ltn1's binding to the

ribosome (Defenouillère et al., 2013; Shao et al., 2015) and directs the 60S subunit to extend

the C-terminus of the incomplete polypeptide with alanine and threonine residues ("CAT tails")

(Osuna et al., 2017; Shen et al., 2015). This process ("CATylation") enables Ltn1 to ubiquitylate

lysine residues previously buried in the ribosome exit tunnel (Kostova et al., 2017; Sitron and

Brandman, 2019) or confined to structured regions on incomplete polypeptides (Sitron and

Brandman, 2019). Additionally, CAT tails act as degrons off the ribosome to mark polypeptides

that escape Ltn1 for proteasomal degradation (Sitron and Brandman, 2019), a function

conserved from prokaryotes (Lytvynenko et al., 2019). By targeting incomplete polypeptides for degradation, RQC protects cells from the proteotoxic consequences of failed translation.

In animal models, crippling RQC with hypomorphic *LTN1* alleles (Chu et al., 2009) or mutations that compromise the ability to respond to increased ribosome stalling (Ishimura et al., 2014) leads to neurodegeneration. Cellular models of RQC failure display phenotypes that suggest how disruption of RQC might result in this pathology. Failure of RQC after CATylation occurs, for example by mutation of *LTN1*, leads to a buildup of CATylated proteins (Shen et al., 2015). CAT tails have amyloid-like properties, which can cause CATylated proteins to aggregate and form inclusions that sequester chaperones (Choe et al., 2016; Defenouillère et al., 2016; Yonashiro et al., 2016). Indeed, inclusions of stalled proteins with C-terminal extensions added by an Rqc2 ortholog were observed in a *Drosophila* model of Parkinson's Disease (Wu et al., 2019). Despite these aberrant phenotypes, RQC failure is well-tolerated in cellular models and does not impair viability in the absence of strong additional stressors (Choe et al., 2016; Defenouillère et al., 2013; Kostova et al., 2017).

Toward the eventual goal of understanding the relationship between RQC failure and neurodegeneration, we sought to determine how RQC failure can become toxic in *Saccharomyces cerevisiae*. We found that the combination of Ltn1 disruption and elevated Rqc2 levels changed CAT tail composition, potentiated aggregation of CATylated proteins (aggregation previously observed by Yonashiro and colleagues [Yonashiro et al., 2016]), and antagonized the ability of CAT tails to act as degrons. These Rqc2-perturbed cells displayed phenotypes characteristic of proteostasis disruption such as sequestration of the chaperone Sis1 into CAT tail inclusions, induction of the heat shock response, and sensitivity to otherwise mild stressors. Thus, our work indicates that Rqc2, which ordinarily protects cells from RQC failure, increases the toxicity of RQC failure when expressed at high levels.

We additionally uncovered two interventions that reverse the toxic effects of RQC failure. First,

improving the solubility of CATylated proteins by growing cells in the presence of the protein

denaturant guanidinium hydrochloride reduced the proteostasis strain caused by increased

Rqc2 levels. Second, we identified RNA Polymerase III (Pol III) in a screen for modulators of the

heat shock response to RQC failure. Genetic disruption of Pol III depleted an insoluble species

that converted CAT tails into their aggregated form; Pol III perturbation thereby restored CAT tail

degron function and proteostasis. Our work demonstrates that interventions targeting CAT tail

aggregation can reduce the toxicity of RQC failure.

9
10 **Results**

11
12 **Elevated Rqc2 levels increase the toxicity of RQC failure**

13
To study RQC failure in *Saccharomyces cerevisiae*, we first needed a model substrate that

would enable us to visualize defects in the RQC pathway. We used a model substrate

("RQCsub") that consists of green fluorescent protein (GFP) attached via an inert linker

including a tobacco etch virus (TEV) protease site to twelve stall-inducing arginine CGN codons

(Fig. 1A) (Letzring et al., 2010). Translation of RQCsub produces a stalled GFP-linker-arginine

"arrest product" (Brandman et al., 2012). This RQCsub arrest product accumulated to low levels

in wt strains by SDS-PAGE (Fig. 1A). This arrest product increased in abundance after we

compromised RQC function by deleting two RQC genes: the CAT tail-elongating factor *RQC2* or

the E3 ubiquitin ligase *LTN1* (Fig. 1A). This result indicates that Rqc2 and Ltn1 contribute to

RQCsub degradation. While deletion of *RQC2* or *LTN1* stabilized RQCsub, the resultant protein

products displayed different mobilities; RQCsub migrated as a crisp band in *rqc2Δ* and a higher

molecular weight smear in *ltn1Δ* (Fig. 1A). This smear collapsed into a single band after

additional deletion of *RQC2* in the *ltn1Δ* strain (Fig. 1A) or cleavage of RQCsub's C-terminus

with TEV protease (Fig. 1B), consistent with the smear containing CATylated RQCsub. Taken

together, these results confirm that the model substrate RQCsub enables visualization of RQC-

mediated degradation and CATylation.

3
We hypothesized that overexpression of *RQC2*, which potentiates CAT tail aggregation

(Yonashiro et al., 2016), would increase the cellular toxicity of RQC failure. As expected, when

we overexpressed *RQC2* in the RQC-compromised *ltn1Δ* strain, a large proportion of RQCsub

remained in the well rather than migrating into an SDS-PAGE gel (Fig. 1A,B) despite prior

boiling in dodecylsulfate detergent. This detergent-resistant species was more apparent in the

RQC2 overexpression and *LTN1* deletion (hereafter referred to as "*ROLD*") strain than *ltn1Δ*

(Fig. 1B). TEV treatment, which cleaves RQCsub's C-terminus and removes its CAT tails,

enabled RQCsub to migrate as a single band into the gel (Fig. 1B). The formation of this

detergent-resistant species is consistent with *RQC2* overexpression reducing the solubility of

CATylated RQCsub in the *ltn1Δ* background. Microscopy of RQCsub under these conditions

revealed that RQCsub had a diffuse localization in the majority of *ltn1Δ* cells (<1% of cells with

inclusions) but became more punctate in *ROLD* (56.9% of cells with inclusions) (Fig. 1C). To

test whether these inclusions depended on CATylation, we disrupted CATylation by deleting

RQC2 or overexpressing a CATylation-incompetent *rqc2$_{aaa}$* allele (Shen et al., 2015) instead of

RQC2-WT. Neither of these perturbations (*rqc2Δ ltn1Δ* and *rqc2$_{aaa}$-OE ltn1Δ*) induced the

punctate RQCsub localization we observed in *ROLD* (Fig. 1C), confirming that formation of

RQCsub inclusions requires CATylation. These results indicate that *RQC2* overexpression

produces aggregation-prone CAT tails that sequester RQC substrates into detergent-insoluble

inclusions in cells with compromised Ltn1.

23
Cytosolic CAT tail aggregates recruit the Hsp40 chaperone Sis1 (Choe et al., 2016;

Defenouillère et al., 2016; Yonashiro et al., 2016), thereby depleting Sis1 from the nucleus. In

strains where RQCsub had diffuse localization (wt, *ltn1Δ*, and *ltn1Δ rqc2Δ*), there was little

colocalization of Sis1 and the substrate (Fig. 1D). In contrast, RQCsub colocalized with Sis1 in

inclusions at the periphery of most cells in the inclusion-containing *ROLD* strain (Fig. 1D). This

peripheral localization of Sis1 did not overlap with a nuclear marker, consistent with the *ROLD*

perturbation depleting Sis1 from the nucleus (Supplementary Fig. 1A). Furthermore, a larger

portion of Sis1 was retained as a detergent-resistant species in the well of an SDS-PAGE gel in

the *ROLD* condition compared to wt, *ltn1Δ*, and *ltn1Δ rqc2Δ* (Fig. 1E). Although perturbation of

LTN1 compromises RQC, these data demonstrate that *LTN1* deletion alone does not modify

Sis1 localization. However, the combination of *LTN1* deletion and increased *RQC2* expression

leads to depletion of Sis1 from the nucleus into detergent-resistant inclusions.

10
Sequestration of Sis1 and other chaperones into aggregates has been shown to impair

proteostasis (Choe et al., 2016; Defenouillère et al., 2016; Park et al., 2013; Yonashiro et al.,

2016). Proteostasis impairment activates Heat Shock Factor 1 (Hsf1), a transcription factor that

orchestrates transcription of chaperones and other proteostasis-restoring factors (Åkerfelt et al.,

2007; Morimoto, 2011). To assess the magnitude of proteostasis disruption caused by *ROLD*,

we measured Hsf1 activity using a previously published reporter (Brandman et al., 2012),

consisting of GFP under the control of an artificial promoter containing four heat shock elements

(Sorger and Pelham, 1987), the consensus binding site for Hsf1 (Fig. 2A). Deletion of *LTN1*

alone activated the Hsf1 reporter 2.07-fold relative to wt, while additional *RQC2* overexpression

(*ROLD*) led to 22.6-fold activation (Fig. 2A). The potent activation observed in *ROLD* required

LTN1 deletion, as *RQC2* overexpression in the wt background weakly affected the reporter (Fig.

2A). These data establish that *RQC2* overexpression and *LTN1* deletion have a synthetic

interaction that results in elevated Hsf1 transcriptional activity. This synergistic enhancement in

Hsf1 activity implies that increased Rqc2 levels may exert additional proteotoxic stress on RQC-

compromised cells.

To investigate how *RQC2* overexpression might increase the proteotoxicity of RQC failure, we

analyzed the composition of CAT tails under this condition. As expected, RQCsub purified from

the CATylation-competent *ltn1Δ* strain was enriched in alanine and threonine relative to the

CATylation-incompetent *ltn1Δ rqc2Δ* strain (Fig. 2B). This result confirmed that this assay could

detect changes in CAT tail composition. Unexpectedly, alanine content in RQCsub was

enriched by 31% in *ROLD* relative to *ltn1Δ* (Fig. 2B). Thus, *RQC2* overexpression, which

induces a large disruption of proteostasis in *ltn1Δ* cells, increases alanine content in CAT tails.

One reason why cells tolerate *LTN1* disruption is that CAT tails function as Ltn1-independent

degrons, thereby targeting RQC substrates that escape Ltn1 for proteasomal degradation

(Sitron and Brandman, 2019). To assess whether *RQC2* overexpression impairs this

degradation pathway, we quantified the stability of the model RQC substrate RQCsub$_{LONG}$

(Sitron and Brandman, 2019). RQCsub$_{LONG}$ enables quantification of its stability by including an

internal expression control, red fluorescent protein (RFP) followed by viral T2A peptides, at the

N-terminus of GFP-linker-arginine (Fig. 2C). Ribosomes translating RQCsub$_{LONG}$ produce RFP

then skip formation of a peptide bond within the T2A sequence (Donnelly et al., 2001; Szymczak

and Vignali, 2005), severing RFP from the GFP-linker-arginine RQC substrate (Fig. 2C). As a

result, each round of translation produces RFP and GFP stoichiometrically, but only GFP

becomes an RQC substrate (Fig. 2C). Thus, the GFP:RFP ratio serves as an internally-

controlled readout for RQCsub$_{LONG}$ stability (Fig. 2C). CAT tail degron activity on RQCsub$_{LONG}$

requires the E3 ubiquitin ligase Hul5 and the proteasome (Sitron and Brandman, 2019).

Therefore, comparing stability before and after *HUL5* deletion quantifies CAT tail degron activity

(Fig. 2C). RQCsub$_{LONG}$ stability did not change after treatment with the proteasome inhibitor

bortezomib in *ltn1Δ hul5Δ* or *ROLD hul5Δ* (Supplementary Fig. 1B), supporting the requirement

for Hul5 in CAT tail-mediated degradation of RQCsub$_{LONG}$ by the proteasome. *HUL5* deletion

stabilized RQCsub$_{LONG}$ by 30.2% in the *ltn1Δ* background and 13.6% in the *ROLD* background

(Fig. 2C), consistent with *RQC2* overexpression reducing RQCsub$_{LONG}$ degradation. These

results demonstrate that *RQC2* overexpression impairs CAT tail degron activity.

4
Given that *RQC2* overexpression in *ltn1Δ* impaired degradation of CATylated proteins and

potentiated their aggregation, we wondered whether this results in defective cellular fitness. At

30^{O}C, all of the strains we measured grew equally well in a spot assay (Fig. 2D). This result

demonstrates that *ROLD* does not impair growth in the absence of stress. We then introduced

two mild stressors to perturb proteostasis: the thermal stress of 37^{O}C incubation or introduction

of 50 ng/ml cycloheximide to increase translational stalling. Neither of these stresses induced a

growth effect in the *RQC2* overexpression (*RQC2-OE*) or *ltn1Δ* strains relative to wt (Fig. 2D).

Contrastingly, 37^{O}C incubation and 50 ng/ml cycloheximide severely inhibited *ROLD* growth

(Fig. 2D). Thus, *RQC2* overexpression sensitizes RQC-compromised cells to otherwise mild

stressors. We posit that this stress-sensitivity, in conjunction with elevated Hsf1 activation,

reflects a fragile state of proteostasis in *ROLD*.

16
17   **The denaturant guanidinium hydrochloride ameliorates the toxic effects of elevated Rqc2**

18   **levels**

19
We have observed that *ROLD* induces several phenotypes that correlate with aggregation of

CATylated proteins: co-aggregation of Sis1 (Fig. 1D,E), increased Hsf1 activity (Fig. 2A),

impaired Ltn1-independent degradation of CATylated proteins (Fig. 2C), and sensitivity to mild

stressors (Fig. 2D). These correlations prompted us to hypothesize that aggregation of

CATylated proteins causes this constellation of phenotypes. To test this hypothesis, we treated

live *ROLD* cells with guanidinium hydrochloride (GuHCl), a protein denaturant that suppresses

aggregation of CAT tails and other aggregation-prone proteins at low concentrations

(Defenouillère et al., 2016; Ness et al., 2002). Sis1 and RQCsub still localized to inclusions by

microscopy after 5 mM GuHCl treatment in *ROLD* (Supplementary Fig. 2A). However,

monitoring RQCsub mobility by SDS-PAGE assessed how GuHCl affected RQCsub and Sis1

more sensitively than microscopy, revealing that GuHCl treatment enabled a larger proportion of

both RQCsub and Sis1 extracted from *ROLD* cells to migrate into an SDS-PAGE gel rather than

remaining in the well (Fig. 3A). GuHCl thus reduces the detergent-insolubility of CATylated

proteins and Sis1 caused by *ROLD*.

7
We then tested whether improving CATylated protein solubility with GuHCl would relieve the

Hsf1 activation observed in *ROLD*. GuHCl treatment activated the Hsf1 reporter in a dose-

dependent manner in wt, *RQC2-OE*, and *ltn1Δ* strains (Fig. 3B). This result might indicate that

GuHCl, a protein denaturant, exerts some proteotoxic stress in cells. Despite this, GuHCl

decreased Hsf1 reporter activation in *ROLD* (Fig. 3B), suggesting that solubilization of CAT tails

alleviates the Hsf1 activation caused by *ROLD*. We tested the specificity of this response by

measuring the effect of GuHCl on the Hsf1 reporter in other genotypes with high Hsf1 activity

(Brandman et al., 2012) (Fig. 3C). Hsf1 reporter deactivation was unique to the *ROLD* condition,

as GuHCl increased Hsf1 reporter activation in *ump1Δ*, *hsp104Δ*, *hsc82Δ*, and *ssa2Δ* strains

(Fig. 3C). These results are consistent with aggregation of CAT tails exerting proteotoxic stress

on cells.

19
Given that short, soluble CAT tails act as Ltn1-independent degrons, we questioned whether

solubilizing aggregated CAT tails in *ROLD* might rescue deficits in CAT tail degron activity (Fig.

2C). We again compared the stability (GFP:RFP) of RQCsub$_{LONG}$ in *ROLD* relative to *ROLD*

hul5Δ, the condition where RQCsub$_{LONG}$ degradation is blocked (Sitron and Brandman, 2019),

to measure CAT tail degron activity. GuHCl had two effects on degradation of RQCsub$_{LONG}$.

First, increasing concentrations of GuHCl increased the magnitude that *HUL5* deletion stabilized

RQCsub$_{LONG}$ (from 13.6% to 20.9% to 28.2% at 0, 5, and 10 mM GuHCl; Fig. 3D), suggesting

that treating *ROLD* cells with GuHCl enhances degradation of RQCsub$_{LONG}$. Second, the

magnitude of bortezomib-induced RQCsub$_{LONG}$ stabilization increased with increasing GuHCl

when Hul5 was intact (Supplementary Fig. 2B). This result indicates that the degradation that

GuHCl enhanced involved both Hul5 and the proteasome. Taken together, these data have two

implications: 1) the insolubility of CAT tails in *ROLD* impairs CAT tail degron activity, and 2)

solubilization of CAT tails rescues this impairment.

7
We next investigated whether GuHCl could dampen the sensitivity of *ROLD* cells to mild

stressors. GuHCl did not generally affect growth (Fig. 3E, plate 1 compared to 2), but additional

GuHCl partially rescued *ROLD* growth in the presence of cycloheximide (Fig. 3E, plate 5

compared to plate 6). While the combination of GuHCl and 37^{O}C incubation led to poorer growth

of all strains relative to 37^{O}C alone, *ROLD* grew comparably to wt, *RQC2-OE*, and *ltn1Δ* when

exposed to both GuHCl and 37^{O}C (Fig. 3E, plate 3 compared to plate 4). Thus, CAT tail-

solubilizing GuHCl treatment ameliorates the stress sensitivity caused by *RQC2* overexpression

in cells with impaired RQC.

16
17   **RNA Polymerase III modulates the toxicity of RQC failure**

18
To identify pathways responsible for the proteotoxicity of RQC failure, we screened for mutants

that had weak activation of the Hsf1 reporter in an RQC-compromised strain. After crossing a

genome-wide collection of hypomorphs into the Hsf1-inducing *rqc1Δ rps0aΔ* strain (Brandman

et al., 2012), we found that all hits other than *RQC2* (whose deletion blocks CATylation) were

related to RNA Polymerase III (Pol III). These hits included a non-essential polymerase

biogenesis factor, *BUD27* (Mirón-García et al., 2013), as well as essential Pol III subunits,

RPC17 and *RPC160*. Disruption of each of these by deletion or a "<u>d</u>ecreased <u>a</u>bundance by

<u>m</u>RNA <u>p</u>erturbation" allele (DAmP) (Yan et al., 2008) nearly eliminated activation of the Hsf1

reporter in *ROLD* compared to wt (Fig. 4A). To assess the specificity of this reduction in Hsf1

reporter induction, we measured the effect of *BUD27* and *RPC17* perturbations in a panel of

strains with elevated Hsf1 signaling (Fig. 4B). Perturbations to *BUD27* and *RPC17* reduced Hsf1

reporter activity in wt, *ump1Δ*, *hsp104Δ*, *hsc82Δ*, and *ssa2Δ* strains (Fig. 4B, comparison

between black and grey bars). In each strain, the magnitude of this reduction was substantially

less than the 20.7-fold and 13.7-fold reduction we observed in *ROLD* after disruption of *BUD27*

and *RPC17*, respectively (Fig. 4B). Furthermore, Hsf1 reporter activation increased in the Pol

III-perturbed backgrounds after the *ROLD* perturbation or upon deletion of *UMP1*, *HSP104*,

HSC82, and *SSA2* (Fig. 4B; Supplementary Fig. 3A). Taken together, these data indicate that

Pol III impairment lessens proteotoxicity caused by *ROLD* more than it does for other genetic

stressors and does not generally prevent Hsf1 activation.

11
One mechanism by which Pol III disruption might alleviate proteotoxicity in *ROLD* is by

preventing ribosome stalling. Translation initiation, which involves Pol III target genes such as

5S rRNA and tRNAs (Turowski and Tollervey, 2016), modulates stalling by controlling the

amount of ribosomes on an mRNA (Juszkiewicz et al., 2018; Simms et al., 2017). When fewer

ribosomes translate an mRNA (low initiation), ribosomes stuck at stalling sequences collide less

frequently, thereby preventing recognition of stalls and subsequent RQC (Juszkiewicz et al.,

2018; Simms et al., 2017). We therefore speculated that lower levels of translation components

in Pol III-perturbed strains might reduce the generation of RQC substrates, thereby preventing

their toxic build-up. To evaluate the effect of Pol III perturbation on translation of stalling

sequences, we used a quantitative stalling reporter similar to one previously developed

(Juszkiewicz and Hegde, 2017; Sundaramoorthy et al., 2017) (Supplementary Fig. 3B). *BUD27*

deletion did not alleviate stalling induced by (CGN)12 (the stalling sequence in RQCsub) in wt or

ROLD relative to control (Supplementary Fig. 3B). Thus, the effects of Pol III perturbation on

translation should not confound our analysis of RQCsub by preventing ribosomes translating

RQCsub from stalling. We then titrated the strength of the stalling sequence by varying the

number of difficult-to-decode CGA codons (Letzring et al., 2010) it contained. In the *ROLD*

background, *BUD27* deletion had no effect on stalling induced by short stretches of CGA

codons (one or two CGA codons) or a more severe stretch (eight CGA codons); *BUD27* deletion

weakly reduced stalling on sequences that contained four or six CGA codons (Supplementary

Fig. 3C). Thus, ribosomes still stall when translating difficult-to-decode codons after Pol III

perturbation. This result suggests that prevention of stalling is not the mechanism by which Pol

III perturbation alleviates proteotoxicity in *ROLD*.

Given that stalling still occurred after disruption of Pol III, we wondered whether CATylation

remains intact under this condition. To analyze the effect of Pol III perturbation on CATylation,

we monitored RQCsub mobility by SDS-PAGE (Fig. 4C). As in *ltn1Δ*, RQCsub expressed in

ROLD bud27Δ migrated as a smear which was higher in molecular weight than RQCsub

expressed in the CATylation-deficient *ltn1Δ rqc2Δ* strain (Fig. 4C). Furthermore, amino acid

analysis revealed that RQCsub CAT tails had similar composition in *ROLD* and *ROLD bud27Δ*

cells (Supplementary Fig. 3D). Pol III perturbation thereby has no effect on CATylation in *ROLD*.

We next investigated whether a reduction in co-aggregation of CATylated proteins and Sis1

could explain the dampened Hsf1 activation observed after Pol III disruption in *ROLD*. *ROLD*

cells carrying hypomorphic *BUD27* or *RPC17* alleles formed RQCsub inclusions less frequently

(3.57% and 4.37% relative to 56.9% in *ROLD*), consistent with a reduction in CAT tail

aggregation (Fig. 4D). Similarly, *BUD27* deletion in *ROLD* prevented colocalization of RQCsub

with Sis1 (Fig. 4E) and led to stronger Sis1 nuclear localization than in *ROLD* (Supplementary

Fig. 3E). Compared to *ROLD*, a smaller proportion of Sis1 in *ROLD bud27Δ* migrated as a

detergent-resistant species during SDS-PAGE (Fig. 4F). Taken together, these results indicate

that Pol III disruption limits aggregation of both CATylated proteins and Sis1 caused by *RQC2*

overexpression in cells with compromised RQC.

Because we previously observed that solubilization of CAT tails with GuHCl improved their

degron strength, we wondered whether Pol III perturbation would have the same effect. To

quantify CAT tail degron strength, we measured degradation of RQCsub$_{LONG}$ in the *ROLD* and

ROLD bud27Δ backgrounds. The difference in RQCsub$_{LONG}$ stability (GFP:RFP) in each of

these backgrounds with (degradation intact) and without Hul5 (degradation blocked) defined the

degree of degradation (Sitron and Brandman, 2019). We validated this approach by

demonstrating that the proteasome inhibitor bortezomib did not significantly stabilize

RQCsub$_{LONG}$ in *hul5Δ* strains (Supplementary Fig. 4A), demonstrating that *HUL5* deletion still

blocked degradation. *HUL5* deletion stabilized RQCsub$_{LONG}$ in the *ROLD bud27Δ* background

more than in *ROLD* (33.7% compared to 13.6%; Fig. 4G), consistent with improved degradation.

Thus, perturbing Pol III in the *ROLD* condition improves CAT tail solubility and degradation of

CATylated proteins.

13
14 **Pol III perturbation renders cells refractory to RQC stress**

15
Because disruption of Pol III allowed for CATylation in *ROLD* but the resultant CAT tails did not

form inclusions in cells, we hypothesized that Pol III impairment depletes a factor that nucleates

aggregation of CAT tails. To test this hypothesis, we added lysate from the *ROLD* strain to

lysate from an RQCsub-expressing *ROLD bud27Δ* strain and monitored formation of the

detergent-resistant RQCsub species *in vitro* (Fig. 5A). In agreement with microscopy from these

strains (Figs. 1D and 5D), deletion of *BUD27* in *ROLD* reduced the abundance of a detergent-

resistant RQCsub species that failed to migrate into an SDS-PAGE gel (Fig. 5A). Addition of

ROLD lysate that lacked RQCsub to *ROLD bud27Δ* lysate containing RQCsub converted a

population of the substrate into its detergent-resistant form (Fig. 5A), suggesting that aggregate

nucleation had taken place. Lysate from wt cells lacked this apparent nucleation activity (Fig.

5A). We then tested whether the nucleating factor from *ROLD* was an RNA species (e.g. a Pol

III-dependent transcript) by first Rnase A-treating the *ROLD* lysate. This RNAse A-treated lysate

still stimulated the appearance of detergent-resistant RQCsub (Fig. 5A). However, the 95,000 g

supernatant of the *ROLD* lysate displayed less ability to convert RQCsub into its detergent-

resistant form compared to total *ROLD* lysate (Fig. 5A). Taken together, these data indicate that

the *ROLD* condition leads to accumulation of an insoluble species that converts CAT tails into a

detergent-insoluble form. We posit that perturbation of Pol III reduces the abundance of this

species and thereby limits CAT tail aggregation.

8
We then sought to determine whether forcing nucleation of CAT tails into inclusions would revert

Pol III-perturbed *ROLD* cells into a stressed state. To accomplish this, we first tested whether

expression of "hard-coded" CAT tails, proteins containing genetically-encoded stretches of

alanine and threonine, would nucleate aggregation of endogenous CAT tails. These proteins

were not substrates for RQC, as depletion of RQC components failed to increase levels of RFP

constructs containing hard-coded CAT tails (Supplementary Fig. 4B). RFP with a C-terminal

sequence containing a 20 amino acid hard-coded CAT tail (RFP-AT20) displayed diffuse

localization in wt and *ROLD bud27Δ* (Fig. 5B), two strains where few cells contain inclusions of

CATylated RQC substrates (Figs. 1C,4D). In the *ROLD* strain, wherein many cells carry

inclusions of CATylated RQC substrates (Fig. 1C), RFP-AT20 formed peripheral inclusions (Fig.

5B). These results suggest that a 20 amino acid-long CAT tail alone cannot nucleate

aggregation, but can form inclusions in strains with existing CAT tail aggregates. RFP attached

to a 60 amino acid-long hard-coded CAT tail (RFP-AT60) localized to inclusions in all strains we

tested (Fig. 5B). Expressing this protein in *ltn1Δ* and *ROLD bud27Δ*, strains with few

endogenous CAT tail inclusions (Figs. 1C,4D), led to colocalization of RFP-AT60 and

endogenously CATylated RQCsub in peripheral inclusions (Fig. 5C). Similarly, expression of the

non-fluorescent hard-coded CAT tail construct GST-AT60 in *ROLD bud27Δ* led to colocalization

of Sis1 and RQCsub in puncta, while the shorter GST-AT20 hard-coded CAT tail did not (Fig.

5D). Taken together, these results demonstrate that hard-coded CAT tails can force nucleation

of endogenously CATylated proteins and Sis1 into inclusions.

Using hard-coded CAT tails, we monitored whether restoring nucleation of CAT tail inclusions

disrupts proteostasis in Pol III-perturbed *ROLD* cells. We measured the activity of an Hsf1

reporter as a proxy for proteostasis, reasoning that high Hsf1 activity indicated a disruption in

proteostasis. *ROLD bud27Δ* cells displayed Hsf1 reporter activity lower than *ROLD* (Fig. 5E).

Expression of the non-fluorescent GST-AT20 hard-coded CAT tail, which did not nucleate

inclusions of RQCsub and Sis1 (Fig. 5D), did not activate the Hsf1 reporter in *ROLD bud27Δ*

cells (Fig. 5E). Surprisingly, despite its ability to induce formation of inclusions (Fig. 5D), GST-

AT60 expression failed to activate the Hsf1 reporter (Fig. 5E). These data imply that: 1) re-

localization of Sis1 from the nucleus into CAT tail inclusions does not suffice for Hsf1 activation,

and 2) formation of CAT tail aggregates does not disrupt proteostasis in the Pol III-perturbed

ROLD condition.

We reasoned that this lack of Hsf1 activation after Pol III perturbation implies that Pol III

perturbation decreases the fragility of proteostasis caused by *ROLD*. To test this hypothesis, we

analyzed cells' sensitivity to mild stress under these conditions. As previously published, *BUD27*

deletion slightly lowered growth rate at 30^{O}C and strongly impaired growth at 37^{O}C in the wt

background (Deplazes et al., 2009) as well as in *ROLD* (Fig. 5F). Thus, the sensitivity to mild

thermal stress caused by Pol III impairment renders this perturbation unable to rescue heat

sensitivity in *ROLD*. Despite causing a growth defect in wt cells, *BUD27* deletion enabled faster

growth of *ROLD* cells in the presence of cycloheximide (Fig. 5F). Unexpectedly, the growth rate

of *ROLD bud27Δ* cells upon cycloheximide exposure was faster than the *bud27Δ* strain (Fig.

5F). We thus posit that Pol III perturbation desensitizes *ROLD* cells to stress caused by

increased ribosome stalling.

Discussion

Here we discovered that the combination of high *RQC2* expression and compromised RQC function altered CAT tail composition and converted them into an aggregated form that compromised CAT tail degron activity and disrupted proteostasis (Figs. 1-2). This proteostasis-disrupted state featured re-localization of Sis1 from the nucleus into CAT tail inclusions (Fig. 1D), activation of heat shock signaling (Fig. 2A), and hypersensitivity to mild stressors (Fig. 2D). Guanidinium hydrochloride (GuHCl) treatment and Pol III perturbation ameliorated the effects of *RQC2* overexpression, restoring CAT tail degron function and decreasing proteotoxic stress (Figs. 3-5), thereby mitigating the toxicity of RQC failure.

Although CAT tails have amyloid-like character (Choe et al., 2016; Defenouillère et al., 2016; Yonashiro et al., 2016), we observed little CAT tail aggregation at endogenous Rqc2 levels (Fig. 1B). However, increased *RQC2* expression led to formation of CAT tail inclusions (Fig. 1C). These results imply that high Rqc2 levels alter the properties of CAT tails to drive their aggregation. We posit that this alteration may occur in cis or in trans. In cis, the increased alanine content in CAT tails caused by *RQC2* overexpression (Fig. 2B) could increase their amyloid-like character. Polyalanine tracts have been shown to form amyloid and render proteins detergent-insoluble (Albrecht and Mundlos, 2005; Oma et al., 2007; Rankin et al., 2000; Shinchuk et al., 2005), similarly to what we observed for CAT tails after *RQC2* overexpression (Fig. 1A,B). In trans, increased Rqc2 levels could enable CATylation of more endogenous RQC substrates that nucleate aggregation of other CATylated proteins. This explanation assumes that endogenous Rqc2 is limiting for CAT tail aggregation, as has been hypothesized previously (Yonashiro et al., 2016). In support of this trans-acting explanation, we demonstrated that trans-acting factors can nucleate CAT tail aggregation: addition of exogenous lysate converted non-aggregated CAT tails into a detergent-resistant state (Fig. 5A) and short CAT tails that did not typically aggregate formed inclusions in cells with abundant CAT tail aggregates (Fig. 5B).

1   These cis and trans mechanisms are not mutually exclusive, and could both contribute to how

2   elevated Rqc2 levels promote CAT tail aggregation.

3
In addition to potentiating CAT tail aggregation, *RQC2* overexpression in RQC-compromised

(*ltn1Δ*) cells disrupted proteostasis more than *LTN1* deletion alone (Figs. 1-2). Yet, in this same

ltn1Δ background, we showed previously that the absence of Rqc2 prevents degradation of

escaped RQC substrates and sensitizes cells to stress (Sitron and Brandman, 2019). This

dichotomy implies that cells must strike a balance in setting Rqc2 levels to avoid toxicity when

RQC fails, maintaining Rqc2 expression at a level that supports CAT tail degron function without

leading to proteostasis disruption. Rqc2 levels may thereby modify the pathology of RQC failure.

For instance, our results predict that tissues with high Rqc2 levels relative to Ltn1 or low

expression of both would be most susceptible to impaired RQC. Future studies will evaluate the

role of *RQC2* expression in the neurodegenerative pathology of RQC-compromised animals.

14
CAT tail degron activity safeguards cells from RQC failure. We show here that high levels of

aggregation caused by *RQC2* overexpression and *LTN1* deletion (*ROLD*) co-occur with

disruption of this back-up degradation pathway. This co-occurrence highlights a potential

pathologic positive feedback loop wherein impaired degradation of CATylated proteins causes

them to accumulate, increasing their aggregation, and further sequestering them from

degradation. Two possibilities exist for how *RQC2* overexpression triggers such a feedback

loop: 1) either the altered composition of CAT tails caused by increased Rqc2 (Fig. 2B) creates

weaker degrons that start to build up and aggregate, or 2) a decrease in solubility (Fig. 1A,B)

due to altered amino acid composition (Fig. 2B) or nucleation by trans-acting factors (Fig. 5A)

(e.g. other CATylated proteins) causes CAT tails to become inaccessible to degradation. The

observation that GuHCl both improved the solubility of CAT tails (Fig. 3A) and restored their

degron functionality (Fig. 3D) argues for the latter possibility, although these models are not

mutually exclusive. This inverse relationship between the function of CAT tails as degrons and

CAT tail aggregation implies that CAT tail aggregation is toxic, rather than being adaptive as

previously hypothesized (Yonashiro et al., 2016). Thus, improving the solubility of CAT tails may

prove to be an effective intervention for combating the toxicity of RQC failure.

4
We describe here two interventions that increase the solubility of CAT tails, restore CAT tail

degron function, and reduce the toxicity of RQC failure (Figs. 3-5). GuHCl is a chaotrope which

may disfavor interactions between CAT tails and thereby disrupt aggregate nucleation, as has

been shown for other chaotropic agents on amyloid-like proteins (Rubin et al., 2013). While we

posit that this may be the mechanism by which GuHCl directly improves CAT tail solubility, our

results cannot rule out other indirect mechanisms (Ness et al., 2002). Pol III impairment

reversed RQC failure toxicity more potently than GuHCl. Pol III plays many roles in cells

(Turowski and Tollervey, 2016), so perturbation of Pol III is likely pleiotropic. Our results suggest

that Pol III regulates CAT tail aggregation at the level of nucleation, as Pol III-perturbed lysates

lacked the ability to nucleate CAT tail aggregates (Fig. 5A). Investigating the mechanism by

which Pol III inhibition relieves toxicity from RQC failure and leveraging this mechanism to

develop therapies are promising areas for future research. Indeed, genetic and small-molecule

disruption of Pol III has been shown to be well tolerated in *Drosophila* and *C. elegans*, leading to

lifespan extension (Filer et al., 2017). Our work demonstrates that pharmacologic and genetic

interventions can mitigate the toxic effects of RQC failure.

20
21   **Materials and Methods**

22
23   *Yeast Strains and Culturing*

24
25   The parental wild-type yeast strain used in this study was BY4741. All gene deletions, genomic

26   integrations, and plasmid transformations were done in this background using standard

27   methods. For a complete list of strains used, see Supplementary Table 1. For a complete list of

28   plasmids used, see Supplementary Table 2.

1
Yeast cultures were grown at 30°C in YPD or synthetic defined (SD) media with appropriate

nutrient dropouts. SD media used for growth of yeast cultures contained: 2% w/v dextrose

(Thermo Fisher Scientific, Waltham, MA), 13.4 g/L Yeast Nitrogen Base without Amino Acids

(BD Biosciences, San Jose, CA), 0.03 g/L L-isoleucine (Sigma-Aldrich, St. Louis, MO), 0.15 g/L

L-valine (Sigma-Aldrich), 0.04 g/L adenine hemisulfate (Sigma-Aldrich), 0.02 g/L L-arginine

(Sigma-Aldrich), 0.03 g/L L-lysine (Sigma-Aldrich), 0.05 g/L L-phenylalanine (Sigma-Aldrich),

0.2 g/L L-Threonine (Sigma-Aldrich), 0.03 g/L L-tyrosine (Sigma-Aldrich), 0.018 g/L L-histidine

(Sigma-Aldrich), 0.09 g/L L-leucine (Sigma-Aldrich), 0.018 g/L L-methionine (Sigma-Aldrich),

0.036 g/L L-tryptophan (Sigma-Aldrich), and 0.018 g/L uracil (Sigma-Aldrich). Media additionally

contained 5 mM or 10 mM guanidinium hydrochloride (Sigma-Aldrich) where indicated and

bortezomib (LC Laboratories, Woburn, MA) treatment lasted 4 hours.

13
14 *Plasmids*

15
16 Plasmids were constructed using NEBuilder HiFi DNA Assembly Master Mix (New England

17 Biolabs, Ipswich, MA). All plasmids used in this study are described in Supplementary Table 2.

18
19 Two variants of RQCsub were used that had identical construction except for the region

20 following the stall: one had a fluorescent protein (RFP) and the other had a series of affinity

21 tags. The version without the post-stall RFP was used for microscopy and the version with the

22 post-stall RFP was used in immunoblots; the one exception to this is that the version without the

23 post-stall RFP was used in an immunoblot in Fig. 3A.

24
Hard-coded CAT tail constructs contained stretches of alanine and threonine that were encoded

by the DNA sequences that follow. AT20: 5'-

GCAACCGCAACAGCTACCGCTACAGCCACGGCGACCGCCACTGCAACAGCGACAGCGAC

TTAA-3'. AT60: 5'-

GCCGCCGCAACAACAGCCACTACCACAGCTGCCACTGCTGCGACGACCACAGCCGCAGC

GACGGCTACAGCAGCGACAGCGACTACCGCAACGGCTGCAGCGACAGCCACGGCGACTG

CAGCTACTGCCACGACGACGGCAACCGCCACGGCAACAACAACTACAACCGCGACTGCC

GCATAA-3'.

5
6 *Flow cytometry*

7
8 Fluorescence was measured using a BD Accuri C6 flow cytometer (BD Biosciences). Three

9 independent yeast cultures were grown to log phase overnight in appropriate SD dropout media.

10 Data analysis was performed with MATLAB (MathWorks, Natick, MA).

11
12 For measurements of the Hsf1 activity reporter, raw fluorescence values were first normalized to

13 side scatter, $\log_2$-transformed, and then the value obtained from the wt strain was subtracted

14 from each condition. These calculations are reported as $\log_2$ fold wt in the figures.

15
For measurements of RQCsub$_{LONG}$, a detailed description of the analysis workflow can be

found in Sitron and Brandman, 2019. Briefly, yeast expressing the plasmid were gated

according to above-background RFP fluorescence. Then, bleedthrough from the RFP channel

was calculated using an RFP-only control strain and subtracted from signal in the GFP channel.

The resultant bleedthrough-corrected GFP:RFP values were normalized to the corresponding

untreated *hul5Δ* strain for each background. To quantify the change in GFP:RFP upon *HUL5*

deletion, the normalized value from the *HUL5* condition was subtracted from that of the *hul5Δ*

condition.

24
25 For the stalling reporter, the analysis workflow matched that of RQCsub$_{LONG}$ with the order of

26 the fluorophores being reversed, i.e. plasmid-expressing yeast selected based on GFP,

27 bleedthrough from GFP subtracted from RFP. Where indicated, corrected RFP:GFP values

1   were normalized to a non-stalling reporter that contained a non-stalling sequence encoding

2   serine and threonine (Sitron et al., 2017) in place of the stalling CGN codons.

3
4   *Statistical Analysis*

5
A paired t-test (using the "ttest" function in MATLAB) was used to assess whether mean

measurements differed between two different conditions, using the null hypothesis: $\mu_{condition\ 1}$ =

$\mu_{condition\ 2}$. To calculate the s.e.m. for Δ stability measurements (Figs. 2C,3D,4G), a propagation of

error formula was used: $SEM_{hul5\Delta - HUL5} = sqrt(SEM_{HUL5}^2 + SEM_{hul5\Delta}^2)$. To analyze the significance of Δ

stability measurements, a t-test for particular contrast was performed using the "lm" and

"linearHypothesis" functions in R (R Foundation for Statistical Computing). The null hypothesis

for this test was: $\mu_{hul5\Delta+\ condition\ 1} - \mu_{HUL5\ +\ condition\ 1} = \mu_{hul5\Delta+\ condition\ 2} - \mu_{HUL5\ +\ condition\ 2}$.

13
14   *Immunoblots*

15
16   For whole-cell immunoblots, 0.375/OD600 x mL of yeast culture (OD600 = 0.4-0.8) grown

17   overnight were pelleted and resuspended then boiled for 5 min at 95^{O}C in 15 µL 4x NuPage

18   LDS Sample Buffer (Thermo Fisher Scientific) with 5% β-mercaptoethanol.

19
Samples were loaded into gels and transferred to nitrocellulose using one of two systems. First,

samples were loaded into a NuPAGE Novex 4-12% Bis-Tris 1.5 mm protein gel (Thermo Fisher

Scientific) and run in MOPS buffer. Gels were semi-dry transferred to 0.45 µm nitrocellulose

membranes (Thermo Fisher Scientific) using the Trans-Blot Turbo system (Bio-Rad, Hercules,

CA). Second, samples were loaded into hand-poured 10% Tris-Glycine protein gels with a 4%

stacking gel and run in Tris-Glycine buffer. Gels were wet transferred to .45 µm nitrocellulose

membranes (Thermo Fisher Scientific) in the Trans-Blot Cell (Bio-Rad) in Towbin Buffer.

27
28   Membranes were blocked in TBST with 5% fat-free milk (Safeway, Pleasanton, CA) for 1 hour,

29   incubated overnight at 4°C or at room temperature for 4 hrs in one of the following primary

antibodies: 1:3000 mouse α-GFP (MA5-15256, Thermo Fisher Scientific), 1:3000 rabbit α-

hexokinase (H2035-01, US Biological, Salem, MA), or 1:1000 rabbit α-RFP (AB233, Evrogen,

Moscow, Russia). The following secondary antibodies were then used at 1:5000 dilution: IRDye

800CW donkey anti-mouse, IRDye 800CW goat anti-rabbit, IRDye 680RD goat anti-rabbit, or

IRDye 680RD goat anti-mouse (LiCor Biosciences, Lincoln, NE). Blots were scanned on a Licor

Odyssey (LiCor Biosciences).

Lysate Preparation and Immunoprecipitation (IP)

Yeast were grown in SD media and harvested at OD600 0.8 to 1.0. Cells were harvested using

either 1) centrifugation followed by resuspension in IP buffer (100 mM KOAc, 10 mM MgCl2, 25

mM HEPES-KOH pH 7.4) and freezing of cell droplets in liquid nitrogen; or 2) vacuum filtration

and flash freezing in liquid nitrogen. Frozen yeast were cryogenically pulverized into powder

using a Freezer/Mill (SPEX SamplePrep, Metuchen, NJ) and stored at -80°C.

Frozen yeast powder was thawed and solubilized at 4°C in IP buffer supplemented with Pierce

Protease Inhibitor Tablets, EDTA-free (Thermo Fisher Scientific). Crude lysate was clarified by a

single spin at 5000 x g, incubated with GFP-Trap_A (ChromoTek, Planegg-Martinsried,

Germany) resin for 1 hour at 4°C with rotation, then washed 10 times in IP buffer. For

immunoblotting, washed GFP-Trap resin was boiled in 4X NuPAGE LDS Sample Buffer

(Thermo Fisher Scientific) with 5% β-mercaptoethanol and analyzed as described above in

"Immunoblots."

Amino Acid Analysis

To isolate RQCsub for amino acid analysis, a GFP IP was performed as described above,

except elution was performed with IP buffer adjusted to pH 2.0 to acid-elute proteins. Amino

acid analysis was performed by UC Davis Proteomics Core using the following protocol,

adapted from (Nakazawa and Manabe, 1992): samples were dried in a glass hydrolysis tube.

The samples were then hydrolyzed in 6N HCl, 1% phenol at 110°C for 24 hrs under vacuum.

After hydrolysis, the samples were cooled, unsealed, and dried again. The dried samples were

dissolved in 1mL sodium diluent (Pickering, Mountain View, CA) with 40 nmol/mL NorLeucine

added as an internal standard. For each run, 50 µL of sample was injected onto the ion-

exchange column. An amino acid standards solution for protein hydrolysate on the Na-based

Hitachi 8800 (Sigma,A-9906) is used to determine response factors, and thus calibrate the

Hitachi 8800 analyzer for all of the amino acids. In addition this standard has been verified

against the National Institute of Standards and Technology (NIST) standard reference material

2389a.

11
12 *TEV Protease Digestion*

13
14 GFP IP was performed as described above, but instead of immediate elution, the rein was

15 equilibrated in IP buffer with 1 mM DTT and incubated overnight at 4°C with 5 µL ProTEV Plus

16 (Promega, Sunnyvale, CA) in 100 µL total reaction volume. The resin was then washed with

17 fresh IP buffer with 1 mM DTT and boiled in 4X NuPAGE LDS Sample Buffer (Thermo Fisher

18 Scientific) with 5% β-mercaptoethanol and analyzed as described above.

19
20 *Lysate mixing*

21
Yeast lysate was prepared as described above. Crude lysate from *ROLD bud27Δ* cells

expressing RQCsub was mixed with crude lysate from *ROLD* or wt cells (lacking RQCsub) and

then incubated for 2 hours at 30°C with agitation. For RNAse treatment, *ROLD* lysate was

treated with 100 µg/mL RNAseA (Thermo Fisher Scientific) for 20 minutes at room temperature

prior to lysate mixing. To remove insoluble elements, *ROLD* lysate was centrifuged at 95,000 x

g for 20 minutes and only the upper 50% of the supernatant was used for lysate mixing. After

lysate mixing, GFP IP and immunoblot analysis was performed as described above.

Fluorescence Microscopy

3
Live cell fluorescence imaging was performed using two set-ups: 1) a Nikon Eclipse Ti-E

inverted fluorescence microscope (Nikon, Melville, NY) using a 100X/1.4NA oil objective lens

and µManager (Edelstein et al., 2010), and 2) a Nikon Eclipse 80i microscope using a

100X/1.4NA oil objective lens and Metamorph (Molecular Devices, San Jose, CA). Yeast were

grown in SD media to log phase, centrifuged briefly, resuspended in 5 µL fresh SD media and

mounted on a clean coverslip and slide. Images were prepared and analyzed with ImageJ

(NIH). For quantification, images were randomly acquired until N > 300 cells. Independent

cultures were imaged on different days and error represents standard deviation of 3

independent cultures.

13
14 *Yeast spot assay*

15
Yeast were grown to log-phase in YPD, then diluted to OD600 = 0.1. 200 µL of this diluted

culture was diluted 1:10 into YPD four times in a sterile 96-well plate (Greiner Bio-one,

Kremsmünster, Austria). 5 µL of the serial dilutions were transferred from the 96-well plate onto

YPD agar plates with or without additonal 5 mM guanidinium hydrochloride (Sigma-Aldrich)

and/or 50 ng/mL cycloheximide (Sigma-Aldrich). Plates were then incubated at 30^OC or 37^OC.

Plates grown at 30^OC with or without 5 mM guanidinium hydrochloride as well as 37^OC plates

were imaged after two days. Plates containing 50 ng/mL cycloheximide with or without 5 mM

guanidinium hydrochloride grown at 30^OC were imaged after three days. Plates with 5 mM

guanidinium hydrochloride grown at 37^OC were imaged after six days.

25
26 **Acknowledgments**

27
We thank D. Pincus, D. Jarosz, R. Kopito, B. Lu, Z. Wu, and the members of the Brandman lab

for helpful suggestions. We thank the Theriot and Straight labs (Stanford University) for the use

of their microscopes. J. Schulze (University of California at Davis, Davis, CA, USA) performed

the amino acid analysis at the UC Davis Genome Center Molecular Structure Facility. This work

was supported by Stanford University (O.B.), the US National Institutes of Health (grant No.

R01GM115968 to O.B.), and the National Institute of General Medical Sciences of the US

National Institutes of Health (grant No. T32GM007276 to C.S.S.).

Author Contributions

7
8   O.B., C.S.S., and J.H.P. conceived of the study and designed the experiments. C.S.S., J.H.P.,

9   and J.M.G. performed the experiments. O.B., C.S.S., and J.H.P. analyzed and interpreted the

10   data. O.B., C.S.S., and J.H.P. wrote the manuscript. O.B. supervised the study.

11
12 **Competing Interests**

13
14   The authors declare no competing interests.

15
References

Åkerfelt M, Trouillet D, Mezger V, Sistonen LEA. 2007. Heat shock factors at a crossroad
between stress and development. *Ann N Y Acad Sci* **1113**:15–27.
Albrecht A, Mundlos S. 2005. The other trinucleotide repeat: polyalanine expansion disorders.
Curr Opin Genet Dev **15**:285–293.
Bengtson MH, Joazeiro CAP. 2010. Role of a ribosome-associated E3 ubiquitin ligase in protein
quality control. *Nature* **467**:470–473.
Brandman O, Hegde RS. 2016. Ribosome-associated protein quality control. *Nat Struct Mol Biol*
23:7–15.
Brandman O, Stewart-Ornstein J, Wong D, Larson A, Williams CC, Li G-W, Zhou S, King D,
Shen PS, Weibezahn J, Dunn JG, Rouskin S, Inada T, Frost A, Weissman JS. 2012. A
ribosome-bound quality control complex triggers degradation of nascent peptides and
signals translation stress. *Cell* **151**:1042–1054.
Choe Y-J, Park S-H, Hassemer T, Körner R, Vincenz-Donnelly L, Hayer-Hartl M, Hartl FU.
2016. Failure of RQC machinery causes protein aggregation and proteotoxic stress.
Nature **531**:191–195.
Chu J, Hong NA, Masuda CA, Jenkins BV, Nelms KA, Goodnow CC, Glynne RJ, Wu H, Masliah
E, Joazeiro CAP, Kay SA. 2009. A mouse forward genetics screen identifies LISTERIN
as an E3 ubiquitin ligase involved in neurodegeneration. *Proc Natl Acad Sci U S A*
106:2097–2103.
Defenouillère Q, Fromont-Racine M. 2017. The ribosome-bound quality control complex: from
aberrant peptide clearance to proteostasis maintenance. *Current Genetics*.
doi:10.1007/s00294-017-0708-5

Defenouillère Q, Yao Y, Mouaikel J, Namane A, Galopier A, Decourty L, Doyen A, Malabat C, Saveanu C, Jacquier A, Fromont-Racine M. 2013. Cdc48-associated complex bound to 60S particles is required for the clearance of aberrant translation products. *Proc Natl Acad Sci U S A* **110**:5046–5051.

Defenouillère Q, Zhang E, Namane A, Mouaikel J, Jacquier A, Fromont-Racine M. 2016. Rqc1 and Ltn1 prevent CAT-tail induced protein aggregation by efficient recruitment of Cdc48 on stalled 60S subunits. *J Biol Chem* jbc–M116.

Deplazes A, Möckli N, Luke B, Auerbach D, Peter M. 2009. Yeast Uri1p promotes translation initiation and may provide a link to cotranslational quality control. *EMBO J* **28**:1429–1441.

Dimitrova LN, Kuroha K, Tatematsu T, Inada T. 2009. Nascent peptide-dependent translation arrest leads to Not4p-mediated protein degradation by the proteasome. *J Biol Chem* **284**:10343–10352.

Doma MK, Parker R. 2006. Endonucleolytic cleavage of eukaryotic mRNAs with stalls in translation elongation. *Nature* **440**:561–564.

Donnelly MLL, Ryan MD, Mehrotra A, Gani D, Hughes LE, Luke G, Li X. 2001. Analysis of the aphthovirus 2A/2B polyprotein "cleavage" mechanism indicates not a proteolytic reaction, but a novel translational effect: a putative ribosomal "skip." *Journal of General Virology*. doi:10.1099/0022-1317-82-5-1013

Edelstein A, Amodaj N, Hoover K, Vale R, Stuurman N. 2010. Computer Control of Microscopes Using μManager. *Current Protocols in Molecular Biology*. doi:10.1002/0471142727.mb1420s92

Filer D, Thompson MA, Takhaveev V, Dobson AJ, Kotronaki I, Green JWM, Heinemann M, Tullet JMA, Alic N. 2017. RNA polymerase III limits longevity downstream of TORC1. *Nature* **552**:263–267.

Gamble CE, Brule CE, Dean KM, Fields S, Grayhack EJ. 2016. Adjacent Codons Act in Concert to Modulate Translation Efficiency in Yeast. *Cell* **166**:679–690.

Hartl FU. 2017. Protein Misfolding Diseases. *Annu Rev Biochem* **86**:21–26.

Ikeuchi K, Izawa T, Inada T. 2018. Recent Progress on the Molecular Mechanism of Quality Controls Induced by Ribosome Stalling. *Front Genet* **9**:743.

Ikeuchi K, Tesina P, Matsuo Y, Sugiyama T, Cheng J, Saeki Y, Tanaka K, Becker T, Beckmann R, Inada T. 2019. Collided ribosomes form a unique structural interface to induce Hel2–driven quality control pathways. *EMBO J* e100276.

Ishimura R, Nagy G, Dotu I, Zhou H, -L. Yang X, Schimmel P, Senju S, Nishimura Y, Chuang JH, Ackerman SL. 2014. Ribosome stalling induced by mutation of a CNS-specific tRNA causes neurodegeneration. *Science*. doi:10.1126/science.1249749

Joazeiro CAP. 2019. Mechanisms and functions of ribosome-associated protein quality control. *Nat Rev Mol Cell Biol* **20**:368–383.

Juszkiewicz S, Chandrasekaran V, Lin Z, Kraatz S, Ramakrishnan V, Hegde RS. 2018. ZNF598 Is a Quality Control Sensor of Collided Ribosomes. *Mol Cell* **72**:469–481.e7.

Juszkiewicz S, Hegde RS. 2017. Initiation of Quality Control during Poly(A) Translation Requires Site-Specific Ribosome Ubiquitination. *Mol Cell* **65**:743–750.e4.

Kostova KK, Hickey KL, Osuna BA, Hussmann JA, Frost A, Weinberg DE, Weissman JS. 2017. CAT-tailing as a fail-safe mechanism for efficient degradation of stalled nascent polypeptides. *Science* **357**:414–417.

Letzring DP, Dean KM, Grayhack EJ. 2010. Control of translation efficiency in yeast by codon–anticodon interactions. *RNA*. doi:10.1261/rna.2411710

Lytvynenko I, Paternoga H, Thrun A, Balke A, Müller TA, Chiang CH, Nagler K, Tsaprailis G, Anders S, Bischofs I, Maupin-Furlow JA, Spahn CMT, Joazeiro CAP. 2019. Alanine Tails Signal Proteolysis in Bacterial Ribosome-Associated Quality Control. *Cell*. doi:10.1016/j.cell.2019.05.002

Lyumkis D, dos Passos DO, Tahara EB, Webb K, Bennett EJ, Vinterbo S, Potter CS, Carragher B, Joazeiro CAP. 2014. Structural basis for translational surveillance by the large ribosomal subunit-associated protein quality control complex. *Proceedings of the National Academy of Sciences*. doi:10.1073/pnas.1413882111

Matsuo Y, Ikeuchi K, Saeki Y, Iwasaki S, Schmidt C, Udagawa T, Sato F, Tsuchiya H, Becker T, Tanaka K, Ingolia NT, Beckmann R, Inada T. 2017. Ubiquitination of stalled ribosome triggers ribosome-associated quality control. *Nat Commun* **8**:159.

Meaux S, Van Hoof A. 2006. Yeast transcripts cleaved by an internal ribozyme provide new insight into the role of the cap and poly(A) tail in translation and mRNA decay. *RNA* **12**:1323–1337.

Mirón-García MC, Garrido-Godino AI, García-Molinero V, Hernández-Torres F, Rodríguez-Navarro S, Navarro F. 2013. The prefoldin bud27 mediates the assembly of the eukaryotic RNA polymerases in an rpb5-dependent manner. *PLoS Genet* **9**:e1003297.

Morimoto RI. 2011. The heat shock response: systems biology of proteotoxic stress in aging and disease. *Cold Spring Harb Symp Quant Biol* **76**:91–99.

Nakazawa M, Manabe K. 1992. The direct hydrolysis of proteins containing tryptophanon polyvinylidene difluoride membranes by mercaptoethanesulfonic acid in the vapor phase. *Anal Biochem* **206**:105–108.

Ness F, Ferreira P, Cox BS, Tuite MF. 2002. Guanidine Hydrochloride Inhibits the Generation of Prion "Seeds" but Not Prion Protein Aggregation in Yeast. *Mol Cell Biol* **22**:5593–5605.

Oma Y, Kino Y, Toriumi K, Sasagawa N, Ishiura S. 2007. Interactions between homopolymeric amino acids (HPAAs). *Protein Sci* **16**:2195–2204.

Osuna BA, Howard CJ, Kc S, Frost A, Weinberg DE. 2017. In vitro analysis of RQC activities provides insights into the mechanism and function of CAT tailing. *Elife* **6**. doi:10.7554/eLife.27949

Park S-H, Kukushkin Y, Gupta R, Chen T, Konagai A, Hipp MS, Hayer-Hartl M, Hartl FU. 2013. PolyQ proteins interfere with nuclear degradation of cytosolic proteins by sequestering the Sis1p chaperone. *Cell* **154**:134–145.

Pisareva VP, Skabkin MA, Hellen CUT, Pestova TV, Pisarev AV. 2011. Dissociation by Pelota, Hbs1 and ABCE1 of mammalian vacant 80S ribosomes and stalled elongation complexes. *EMBO J* **30**:1804–1817.

Rankin J, Wyttenbach A, Rubinsztein DC. 2000. Intracellular green fluorescent protein-polyalanine aggregates are associated with cell death. *Biochem J* **348 Pt 1**:15–19.

Ross CA, Poirier MA. 2004. Protein aggregation and neurodegenerative disease. *Nat Med* **10 Suppl**:S10–7.

Rubin J, Khosravi H, Bruce KL, Lydon ME, Behrens SH, Chernoff YO, Bommarius AS. 2013. Ion-specific effects on prion nucleation and strain formation. *J Biol Chem* **288**:30300–30308.

Shao S, Brown A, Santhanam B, Hegde RS. 2015. Structure and assembly pathway of the ribosome quality control complex. *Mol Cell* **57**:433–444.

Shao S, Hegde RS. 2014. Reconstitution of a minimal ribosome-associated ubiquitination pathway with purified factors. *Mol Cell* **55**:880–890.

Shao S, von der Malsburg K, Hegde RS. 2013. Listerin-dependent nascent protein ubiquitination relies on ribosome subunit dissociation. *Mol Cell* **50**:637–648.

Shen PS, Park J, Qin Y, Li X, Parsawar K, Larson MH, Cox J, Cheng Y, Lambowitz AM, Weissman JS, Brandman O, Frost A. 2015. Protein synthesis. Rqc2p and 60S ribosomal subunits mediate mRNA-independent elongation of nascent chains. *Science* **347**:75–78.

Shinchuk LM, Sharma D, Blondelle SE, Reixach N, Inouye H, Kirschner DA. 2005. Poly-(L-alanine) expansions form core beta-sheets that nucleate amyloid assembly. *Proteins* **61**:579–589.

Shoemaker CJ, Eyler DE, Green R. 2010. Dom34:Hbs1 promotes subunit dissociation and peptidyl-tRNA drop-off to initiate no-go decay. *Science* **330**:369–372.

Simms CL, Yan LL, Zaher HS. 2017. Ribosome Collision Is Critical for Quality Control during No-Go Decay. *Mol Cell* **68**:361–373.e5.

Sitron CS, Brandman O. 2019. CAT tails drive degradation of stalled polypeptides on and off the ribosome. *Nat Struct Mol Biol* **26**:450–459.

Sitron CS, Park JH, Brandman O. 2017. Asc1, Hel2, and Slh1 couple translation arrest to nascent chain degradation. *RNA* **23**:798–810.

Sorger PK, Pelham HR. 1987. Purification and characterization of a heat-shock element binding protein from yeast. *EMBO J* **6**:3035–3041.

Sundaramoorthy E, Leonard M, Mak R, Liao J, Fulzele A, Bennett EJ. 2017. ZNF598 and RACK1 Regulate Mammalian Ribosome-Associated Quality Control Function by Mediating Regulatory 40S Ribosomal Ubiquitylation. *Mol Cell* **65**:751–760.e4.

Szymczak AL, Vignali DAA. 2005. Development of 2A peptide-based strategies in the design of multicistronic vectors. *Expert Opin Biol Ther* **5**:627–638.

Tsuboi T, Kuroha K, Kudo K, Makino S, Inoue E, Kashima I, Inada T. 2012. Dom34:hbs1 plays a general role in quality-control systems by dissociation of a stalled ribosome at the 3' end of aberrant mRNA. *Mol Cell* **46**:518–529.

Turowski TW, Tollervey D. 2016. Transcription by RNA polymerase III: insights into mechanism and regulation. *Biochem Soc Trans* **44**:1367–1375.

Verma R, Oania RS, Kolawa NJ, Deshaies RJ. 2013. Cdc48/p97 promotes degradation of aberrant nascent polypeptides bound to the ribosome. *Elife* **2**:e00308.

Wu Z, Tantray I, Lim J, Chen S, Li Y, Davis Z, Citron C, Dong J, Gispert S, Auburger G, Brandman O, Bi X, Snyder M, Lu B. 2019. MISTERMINATE Mechanically Links Mitochondrial Dysfunction with Proteostasis Failure. *bioRxiv*. doi:10.1101/554634

Yan Z, Costanzo M, Heisler LE, Paw J, Kaper F, Andrews BJ, Boone C, Giaever G, Nislow C. 2008. Yeast Barcoders: a chemogenomic application of a universal donor-strain collection carrying bar-code identifiers. *Nature Methods*. doi:10.1038/nmeth.1231

Yonashiro R, Tahara EB, Bengtson MH, Khokhrina M, Lorenz H, Chen K-C, Kigoshi-Tansho Y, Savas JN, Yates JR, Kay SA, Craig EA, Mogk A, Bukau B, Joazeiro CAP. 2016. The Rqc2/Tae2 subunit of the ribosome-associated quality control (RQC) complex marks ribosome-stalled nascent polypeptide chains for aggregation. *Elife* **5**:e11794.

Figure 1. *RQC2* overexpression enhances aggregation of CATylated proteins and Sis1 in *ltn1Δ* cells. (**A**) Left, whole cell immunoblots (IBs) of lysates containing RQCsub (a model RQC substrate; schematic above). Right, immunoblot of RQCsub immunoprecipitated (IPed) from *RQC2* overexpression and *LTN1* deletion (*ROLD*) lysate. Arrow indicates molecular weight of RQCsub without CAT tails. Asterisk denotes the full-length RQCsub protein product, produced when ribosomes translate through the stall sequence (region past the stall not pictured in schematic). GFP, green fluorescent protein. OE, overexpression. (**B**) Immunoblot of RQCsub immunoprecipitated from *ltn1Δ* and *ROLD* lysates with and without tobacco etch virus (TEV) protease treatment. Arrows and asterisks as in **A**. (**C**) Fluorescence microscopy of cells expressing RQCsub with percentages of cells containing observable GFP inclusions reported below. (**D**) Fluorescence microscopy of SIS1-RFP cells expressing RQCsub. (**E**) Whole-cell immunoblot of Sis1-RFP. Asterisk denotes a non-specific band detected by the anti-RFP antibody.

Figure 2. *ROLD* induces Hsf1 activation, disrupts CAT tail composition and degron function, and sensitizes cells to stress. (**A**) Flow cytometry of cells containing an integrated reporter for Hsf1 activation (schematic above). Error bars represent the standard error of the mean (s.e.m.) derived from three independent cultures. P-values from indicated paired t-tests are indicated by lines between bars. HSE, heat shock element. (**B**) Amino acid analysis of RQCsub immunoprecipitated from *ltn1Δ*, *ltn1Δ rqc2Δ*, or *ROLD* lysates. (**C**) Left, schematic of expression-controlled model RQC substrate RQCsub_LONG. Center, definitions of "stability," "Δ stability," and "CAT tail degron activity." Right, Δ stability measurements of RQCsub_LONG expressed in *ltn1Δ* and *ROLD* cells after *HUL5* deletion to assess CAT tail degron activity. Error bars as in **A**. P-values above bars are derived from t-tests for particular contrast, measuring how significantly different *HUL5* deletion-induced stabilization is between *ltn1Δ* and *ROLD* backgrounds. (**D**) Spot assay for strains grown under indicated conditions.

Figure 3. Guanidinium hydrochloride solubilizes CAT tails and mitigates *ROLD*-induced phenotypes. (**A**) Whole cell immunoblot (IB) of *ROLD* cells expressing genomically RFP-tagged Sis1 and RQCsub with and without guanidinium hydrochloride (GuHCl) treatment. Arrow next to the anti-GFP IB indicates molecular weight of RQCsub without CAT tails. Asterisk next to the anti-RFP IB points to a non-specific band detected by the anti-RFP antibody. (**B** and **C**) Flow cytometry of cells containing an integrated reporter for Hsf1 activation with and without GuHCl treatment. Error bars represent s.e.m. of three independent cultures. Lines between bars indicate p-values from paired t-tests. (**D**) Δ stability measurements of RQCsub$_{LONG}$ to assess how growth in GuHCl affects CAT tail degron activity (defined in Fig. 2C) for *ROLD* cells. Error bars as in **B** and **C**. P-values indicated above bars are the result of a t-test for particular contrast, measuring how significantly different CAT tail degron activity is at different GuHCl concentrations. (**E**) Spot assay of strains grown at 30°C, 37°C, and in 50 ng/ml cycloheximide with and without the presence of 5 mM GuHCl. Plate numbers indicated below images.

Figure 4. Pol III perturbation reverses *ROLD*-induced effects on heat shock, CAT tail aggregation, and CAT tail degron activity. (**A** and **B**) Flow cytometry of cells containing an integrated reporter for Hsf1 activation with and without perturbations to RNA Polymerase III. Error bars indicate s.e.m. from three independent cultures. P-values from paired t-tests are indicated with lines connecting bars. (**C**) Whole-cell immunoblot of RQCsub. Arrow indicates molecular weight of RQCsub without CAT tails. (**D**) Fluorescence microscopy of cells expressing RQCsub. The percentages of cells containing inclusions are indicated below. (**E**) Fluorescence microscopy of *SIS1-RFP* cells expressing RQCsub. (**F**) Whole-cell immunoblot of Sis1-RFP. (**G**) Δ stability measurements of RQCsub$_{LONG}$ expressed in indicated strains after *HUL5* deletion to quantify CAT tail degron activity. Error bars as in **A** and **B**. P-value indicates the result of a t-test for particular contrast; this assesses how significantly different *HUL5* deletion-induced stabilization is in *ROLD* compared to *ROLD bud27Δ*.

Figure 5. Pol III perturbation renders cells resistant to nucleation of CAT tail aggregates and insensitive to increased ribosome stalling. (**A**) Upper, schematic of lysate mixing experiment. RQCsub-expressing lysate from *ROLD bud27Δ* was mixed with various RQCsub-free lysates. Lower, immunoblot of immunoprecipitated mixed lysates. Arrow and asterisk indicate molecular weight of non-CATylated RQCsub and RQCsub product derived from complete translation through the RQCsub stall sequence, respectively. (**B**) Fluorescence microscopy of cells expressing RFP with a hard-coded CAT tail appended to its C-terminus (RFP-AT20 and RFP-AT60 indicate 20 and 60 amino acid-long appendages, respectively). Quantification of cells containing inclusions indicated below images. (**C**) Fluorescence microscopy of cells co-expressing RQCsub and RFP-AT60. (**D**) Fluorescence microscopy of *ROLD bud27Δ SIS1-RFP* cells co-expressing RQCsub and GST with hard-coded CAT tails. (**E**) Flow cytometry of cells containing an integrated reporter for Hsf1 activation expressing indicated constructs. EV, empty vector. Error bars indicate s.e.m. from three independent cultures. P-values from paired t-tests indicated above bars. (**F**) Spot assay of cells grown under indicated conditions.

Figure S1. *ROLD* depletes Sis1 from the nucleus and impairs CAT tail degron activity. (**A**) Fluorescence microscopy of cells containing an RFP-tagged *SIS1* allele in their genomes and expressing a nuclear marker, bipartite (bp) SV40 NLS-tagged GFP. (**B**) Additional data to support Fig. 2C. Stability measurements of RQCsub$_{LONG}$ expressed in *ltn1Δ* and *ROLD* cells after bortezomib (btz) treatment to inhibit the proteasome and *HUL5* deletion to block CAT tail degron activity. Error bars represent s.e.m. from three independent cultures. P-values from statistical tests appear above bars. Thicker lines come from paired t-tests, measuring the significance of bortezomib-induced stabilization. Thinner lines come from t-tests for particular contrast, measuring how significantly different *HUL5* deletion-induced stabilization is between *ltn1Δ* and *ROLD* backgrounds.

Figure S2. Effects of guanidinium hydrochloride on RQCsub inclusion formation and CAT tail degron activity. (**A**) Fluorescence microscopy of *ROLD SIS1-RFP* cells expressing RQCsub in the absence or presence of 5 mM guanidinium hydrochloride (GuHCl). (**B**) Additional data to support Fig. 3D. Stability measurements of RQCsub_LONG expressed in *ROLD* cells grown in indicated GuHCl concentrations with additional bortezomib treatment to inhibit the proteasome and *HUL5* deletion to measure CAT tail degron activity. Error bars represent s.e.m. from three independent cultures. P-values are indicated above bars. Thick lines indicate paired t-tests, probing the significance of bortezomib-induced stabilization. Thin lines denote t-tests for particular contrast, measuring how significantly different *HUL5* deletion-induced stabilization is at different GuHCl concentrations.

Figure S3. Effects of Pol III perturbation on Hsf1 activation, ribosome stalling, CAT tail composition, and Sis1 localization. (**A**) Flow cytometry of Pol III-perturbed cells containing an integrated reporter for Hsf1 activation. These data are also contained in Fig. 4B, but are reordered here to simplify comparisons within two Pol III-perturbed genetic backgrounds. Error bars indicate s.e.m. from three independent cultures. (**B**) Left, schematic of a stalling reporter similar to that used in Juszkiewicz and Hegde 2017 and Sundaramoorthy et al. 2017. Right, schematic of stalling reporter with the same (CGN)12 stalling sequence contained in RQCsub or a non-stalling (Ser-Thr)6 sequence. Error bars as in **A**. (**C**) Quantification of the stalling reporter from **B** containing various lengths of CGA codon stretches rather than (CGN)12. Error bars as in **A**. (**D**) Amino acid analysis of RQCsub immuno-purified from *ROLD bud27Δ* cells compared to *ROLD*. (**E**) Fluorescence microscopy of *SIS1-RFP* cells expressing GFP with a nuclear localization sequence, bipartite SV40.

Figure S4. Effects of mutations on an RQC substrate and hard-coded CAT tail constructs. (**A**) Additional data in support of Fig. 4G. Stability measurements of RQCsub$_{LONG}$ expressed in indicated strains with bortezomib (btz) treatment to inhibit the proteasome and *HUL5* deletion to block CAT tail degron activity. Error bars indicate s.e.m. from three independent cultures. P-values are given above bars. Results of paired t-tests measuring the significance of bortezomib-induced stabilization are indicated with thick lines. The result of a t-test for particular contrast is indicated with thin lines; this assesses how significantly different *HUL5* deletion-induced stabilization is in *ROLD* compared to *ROLD bud27Δ*. (**B**) Flow cytometry of RFP with appended hard-coded CAT tails 20 (AT20) or 60 (AT60) amino acids in length. Error bars as in **A**.

Supplementary Table 1. Yeast strains used in this study.

Strain Number	Description
yOB199	BY4741
yOB259	BY4741 *ltn1Δ::kanMX*
yOB454	BY4741 *rqc2Δ::kanMX*
yOBj004	BY4741 *ltn1Δ::kanMX rqc2Δ::natMX*
yOB518	BY4741 *pRQC2::HIS3-pTDH3*
yOB551	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX*
yOBj578	BY4741 *sis1::SIS1-mKate2-hphMX*
yOBj579	BY4741 *sis1::SIS1-mKate2-hphMX ltn1Δ::kanMX*
yOBj580	BY4741 *sis1::SIS1-mKate2-hphMX ltn1Δ::kanMX rqc2Δ::natMX*
yOBj581	BY4741 *sis1::SIS1-mKate2-hphMX pRQC2::HIS3-pTDH3 ltn1Δ::natMX*
yOB255	BY4741 *ura3::HSE-EmGFP*
yOB549	BY4741 *ura3::HSE-EmGFP rqc2Δ::kanMX*
yOB550	BY4741 *ura3::HSE-EmGFP ltn1Δ::kanMX*
yOBj245	BY4741 *ura3::HSE-EmGFP rqc2Δ::kanMX ltn1Δ::natMX*
yOB515	BY4741 *ura3::HSE-EmGFP pRQC2::HIS3-pTDH3*
yOBj185	BY4741 *ura3::HSE-EmGFP pRQC2::HIS3-pTDH3 ltn1Δ::natMX*
yOB1118	BY4741 *RQC2-WT::LEU ltn1Δ::hphMX*
yOB2825	BY4741 *RQC2-WT::LEU ltn1Δ::hphMX hul5Δ::natMX*
yOB3812	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX hul5Δ::hphMX*
yOBj482	BY4741 *ura3::HSE-EmGFP ump1Δ::natMX*
yOBj538	BY4741 *ura3::HSE-EmGFP ump1Δ::natMX bud27Δ::kanMX*
yOB738	BY4741 *ura3::HSE-EmGFP ump1Δ::natMX rpc17-DAmP::natMX*
yOB736	BY4741 *ura3::HSE-EmGFP hsp104Δ::natMX*
yOB808	BY4741 *ura3::HSE-EmGFP hsp104Δ::natMX bud27Δ::kanMX*
yOB809	BY4741 *ura3::HSE-EmGFP hsp104Δ::natMX rpc17-DAmP::kanMX*
yOBj542	BY4741 *ura3::HSE-EmGFP hsc82Δ::natMX*
yOB818	BY4741 *ura3::HSE-EmGFP hsc82Δ::natMX bud27Δ::kanMX*
yOB722	BY4741 *ura3::HSE-EmGFP hsc82Δ::natMX rpc17-DAmP::kanMX*
yOBj563	BY4741 *ura3::HSE-EmGFP ssa2Δ::natMX*
yOB819	BY4741 *ura3::HSE-EmGFP ssa2Δ::natMX bud27Δ::kanMX*
yOB725	BY4741 *ura3::HSE-EmGFP ssa2Δ::natMX rpc17-DAmP::kanMX*
yOBj536	BY4741 *ura3::HSE-EmGFP bud27Δ::kanMX*
yOB710	BY4741 *ura3::HSE-EmGFP rpc17-DAmP::kanMX*
yOBj495	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX bud27Δ::kanMX*
yOBj556	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX rpc17-DAmP::kanMX*
yOBj536	BY4741 *ura3::HSE-EmGFP bud27Δ::kanMX*
yOBj185	BY4741 *ura3::HSE-EmGFP pRQC2::HIS3-pTDH3 ltn1Δ::natMX bud27Δ::kanMX*

yOBj556	BY4741 *ura3::HSE-EmGFP pRQC2::HIS3-pTDH3 ltn1Δ::natMX rpc17-DAmP::kanMX*
yOBj626	*SE-EmGFP pRQC2::HIS3-pTDH3 ltn1Δ::natMX rpc160-DAmP::kanMX*
yOBj582	BY4741 *sis1::SIS1-mKate2-hphMX pRQC2::HIS3-pTDH3 ltn1Δ::natMX bud27Δ::kanMX*
yOB3812	BY4741 *pRQC2::HIS3-pTDH3 ltn1Δ::natMX hul5Δ::hphMX bud27Δ::kanMX*

Supplementary Table 2. Plasmids used in this study.

Plasmid Name	Description	Selection Marker
RQCsub	CEN/ARS pTDH3-GFP_TEV_R12_RFP	*URA3*
RQCsub (no RFP)	CEN/ARS pTDH3-GFP_TEV_R12_3xHA_3xMYC_3xFLAG	*URA3*
RQCsubLONG	CEN/ARS pTDH3-mKate2-2xT2A-GFP_TEVlonglinker_R12-3xMYC	*URA3*
pRS315	CEN/ARS empty vector	*LEU2*
pJP014	CEN/ARS pRQC2-RQC2WT	*LEU2*
pJP030	CEN/ARS pTDH3-rqc2aaa (D9A, D98A, R99A)	*LEU2*
pJP088	CEN/ARS pTDH3-3xFLAG_mKate2_TEV_20AT	*LEU2*
pJP063	CEN/ARS pTDH3-3xFLAG_mKate2_TEV_60AT	*LEU2*
bJP084	CEN/ARS pTDH3-GST_TEV_20AT	*LEU2*
bJP083	CEN/ARS pTDH3-GST_TEV_60AT	*LEU2*
GFP-R12	CEN/ARS pTDH3-GFP_TEV_R12_3xHA_3xMYC_3xFLAG	*URA3*
pJP082	pFA6a_mKate2_tADH1-hphMX	*HYG*
bpSV40-NLS-GFP	CEN/ARS pTDH3-bpSV40NLS-GFP	*URA3*
Stalling Reporter	CEN/ARS pTDH3-GFP-2xT2A-TEV_R12-2xT2A-mKate2	*URA3*
Non-stalling Reporter	CEN/ARS pTDH3-GFP-2xT2A-TEV_6xST-2xT2A-mKate2	*URA3*
Stalling Reporter - CGA1	CEN/ARS pTDH3-GFP-2xT2A-TEV_1xCGA-2xT2A-mKate2	*URA3*
Stalling Reporter - CGA2	CEN/ARS pTDH3-GFP-2xT2A-TEV_2xCGA-2xT2A-mKate2	*URA3*
Stalling Reporter - CGA4	CEN/ARS pTDH3-GFP-2xT2A-TEV_4xCGA-2xT2A-mKate2	*URA3*
Stalling Reporter - CGA6	CEN/ARS pTDH3-GFP-2xT2A-TEV_6xCGA-2xT2A-mKate2	*URA3*
Stalling Reporter - CGA8	CEN/ARS pTDH3-GFP-2xT2A-TEV_8xCGA-2xT2A-mKate2	*URA3*