Rolling stones gather moss: Movement and longevity of moss balls on an Alaskan

glacier

Scott Hotaling[1], Timothy C. Bartholomaus[2], and Sophie L. Gilbert[3]

Affiliations:

[1] School of Biological Sciences, Washington State University, Pullman, WA, USA; ORCID =

0000-0002-5965-0986

[2] Department of Geology, University of Idaho, Moscow, ID, USA; ORCID = 0000-0002-1470-

6720

[3] College of Natural Resources, University of Idaho, Moscow, ID, USA; ORCID = 0000-0002-

9974-5146

Correspondence: Sophie L. Gilbert, College of Natural Resources, University of Idaho,

Moscow, ID, 83844, USA; Email: sophiegilbert@uidaho.edu; Phone: (208) 885-8605

Abstract:

Glaciers support diverse ecosystems that are largely comprised of microbial life. However, at

larger, macroscopic scales, glacier moss balls (sometimes called "glacier mice") develop from

impurities in the ice and represent a relatively rare biological phenomenon. These ovoid-shaped

conglomerations of dirt and moss are only found on some glacier surfaces and provide key

habitats for invertebrate colonization. Yet, despite their development and presence being widely

reported, no targeted studies of their movement and longevity have been conducted. This

knowledge gap is particularly important when considering the degree to which glacier moss

balls may represent viable, long-term biotic habitats on glaciers, perhaps complete with their

own ecological succession dynamics. Here, we describe the movement and longevity of glacier

moss balls on the Root Glacier in southcentral Alaska, USA. We show that glacier moss balls

move an average of 2.5 cm per day in herd-like fashion, and their movements are positively

correlated with glacier ablation. Surprisingly, the dominant moss ball movement direction does

not align with the prevailing wind nor downslope directions; instead, we propose that it depends

on the dominant direction of solar radiation. We also show that glacier moss balls are relatively

long-lived, with a lifespan in excess of 6 years and annual survival rates similar to large

vertebrates. Finally, we observed moss ball formation on the Root Glacier to occur within a

narrow, low albedo stripe downwind of a nunatuk, a potential key source of moss spores and/or

fine-grained sediment that interact to promote their formation.

Keywords: cryobiology, glacier mice, glacier biology, jokla-mys, Root Glacier, Wrangell-St.

Elias National Park

Introduction:

Glaciers have long been overlooked as important components of global biodiversity, but

it is now clear that they host thriving, multi-trophic ecosystems (Anesio and Laybourn-Parry

2012), supporting taxa from microbes to vertebrates (Rosvold 2016; Dial et al. 2016; Hotaling et

al. 2019; Hotaling et al. 2017b). Most biological activity on glaciers occurs within surface ice

where microorganisms take advantage of nutrients that are either wind-delivered or generated

in situ (Hotaling et al. 2017b). In addition to a nutrient input, impurities on the glacier surface can

drive the development of at least two potential "hotspots" of biological diversity on glaciers: well-

studied cryoconite holes (depressions in the ice surface caused by local melt, Anesio et al.

2017) and glacier moss balls (ovular conglomerations of moss and sediment that move on the

glacier surface, Coulson and Midgley 2012).

Under the right conditions, a small piece of rock or other impurity can set in motion the

formation of a glacier moss ball [also referred to as "jokla-mys" (Eythórsson 1951), "glacier

mice" (e.g., Coulson and Midgley 2012), or "moss cushions" (e.g., Porter et al. 2008)]. On a

local scale, glacier moss balls are typically distributed with some degree of local clustering (e.g.,

~1 glacier moss ball/m^2; Fig. 1). While more immobile moss aggregations have been observed

on glaciers elsewhere (e.g., East Africa, Uetake et al. 2014), true glacier moss balls appear to

be particularly rare, having only been described on a few geographically disparate glaciers in

Alaska (Shacklette 1966; Heusser 1972), Iceland (Eythórsson 1951), Svalbard (Belkina and

Vilnet 2015), and South America (Perez 1991). Many different moss species have been found in

glacier moss balls (Shacklette 1966; Heusser 1972; Perez 1991; Porter et al. 2008), suggesting

that they are not dependent on specific taxa, but instead their development is likely driven by

the interaction of suitable biotic (e.g., availability of moss spores) and abiotic (e.g., growth

substrate) factors. However, the specific steps and timeline underlying glacier moss ball genesis

remains unclear.

An intriguing aspect of glacier moss balls, and one that is almost certainly partially

responsible for their "glacier mice" namesake, is their movement. Though the speed and

direction of glacier moss ball travel has not been studied, it has been posited that they move by

inducing the formation of an ice pedestal, then rolling or sliding off of it (Porter et al. 2008). Moss

balls first reduce local albedo by shielding the ice beneath them from sunlight and locally

reducing the ablation rate. As the surrounding ice melts, the glacier moss ball is left on an

elevated pedestal. Eventually, a threshold is reached where the moss ball falls from its pedestal

and the process begins anew, potentially including a "flip" of the moss ball that exposes what

was previously their underside (Porter et al. 2008). However, the speed and direction of moss

ball movement has not been tested, though it has been suggested that their movements

generally track the downslope direction of their local habitat (Porter et al. 2008).

Where they do occur, glacier moss balls contribute to glacier biodiversity by offering a

thermally buffered, island-like habitat on the glacier surface that can host a wide array of

invertebrates (Coulson and Midgley 2012). On Icelandic glaciers, moss balls contain

invertebrate communities dominated by springtails (Collembola), tardigrades (Tardigrada), and

nematodes (Nematoda; Coulson and Midgley 2012). While an array of potential food resources

are available on glacier surfaces (Hotaling et al. 2017b), these are typically only exploited by

invertebrates on the margins of glaciers (e.g., springtails, spiders, grylloblattids) because

suitable on-glacier habitat is lacking (Mann et al. 1980). Glacier moss balls may therefore

provide key habitable islands on the glacier that facilitate wider resource exploitation versus

glaciers without moss balls (Coulson and Midgley 2012). It is also possible that glacier moss

balls, which have not been shown to be inhabited by larger predatory insects (e.g.,

grylloblattids) may provide prey refuge that are sufficiently removed from the typical foraging

areas of their predators. Either way, it is clear that glacier moss balls represent important habitat

for glacier-associated fauna yet basic aspects of their ecology (e.g., longevity and movement)

are unknown.

In this study, we took an integrated behavioral ecology and geophysical approach to the

study of glacier moss balls to answer three basic questions about their life history: (1) What is

the lifespan of a glacier moss ball? (2) How quickly do they move and is their movement

idiosyncratic or herd-like? (3) Are the movements of glacier moss balls linked to the ablation of

the glacier itself? The answers to these questions have implications for invertebrate fauna in

glaciated ecosystems, nutrient cycling (both directly via moss ball decomposition and indirectly

as supporting habitat for biotic communities), and feedback between glacier moss balls and

local ablation rates. Beyond biotic interactions and ecosystem dynamics, glaciers are rapidly

receding worldwide (Gardner et al. 2013; Larsen et al. 2015; Roe et al. 2017) and their

diminished extents will almost certainly affect the persistence of glacier moss balls on local and

global scales. Thus, it is important to better understand these unique micro-ecosystems before

their habitats are lost.

Fig. 1. a) Our study site (solid green square) on the Root Glacier in southcentral Alaska, USA, within
Wrangell-St. Elias National Park. Contour lines are spaced every 100 m in elevation. The dashed square
represents the field of view shown in panel (b). The inset map shows the location of the Root Glacier
(white star) within Alaska. b) Satellite image of the study site (green square) showing the confluence of
the Root and Kennicott Glaciers with the Donoho nunatak to the northwest. The image was recorded on
19 June 2013. c) A landscape view looking northwest of the study site dotted with glacier moss balls. d) A
close-up view of a glacier moss ball with the type of bracelet tag used in this study.

Materials and methods:

Study area

We conducted fieldwork over four years (July, 2009 - July, 2012) on the lowest portion of

the Root Glacier, a major tributary to the Kennicott Glacier, in the Wrangell Mountains in

Wrangell-St. Elias National Park, Alaska, USA (Fig. 1a). Our study area (61.5076° N, 142.9172°

W, ~700 m elevation) spanned a ~15 x ~40 m (600 m^2) area of glacier ice selected for its

especially high concentration of moss balls. The site has a gentle slope, dipping 3° east-

northeast (N75°E) and is found between two medial moraines (Fig. 1b), each ~100 m away.

Moss ball concentrations decrease both up- and down-glacier and are absent from the coarse-

grained (> 5 cm) rock that covers the adjacent medial moraines.

We estimated the proportion of fine-grained sediment cover on the ice within our study

area by applying image processing techniques in the Python package scikit-image (Van der

Walt et al. 2014) to two vertical photographs taken of representative ice surfaces. Pixel

brightness contrasts between ice and sediment are most distinct within the blue band of the red-

green-blue images, so we differentiated between sediment (dark pixels) and ice (bright pixels)

by binarizing the blue band with Otsu's thresholding method. We then performed a

morphological opening to diminish the influence of light-colored sediment grains set within the

otherwise dark sediment cover. Finally, we quantified the areal sediment cover as being

approximately equal to the number of dark colored pixels relative to the total number of pixels in

the binarized images.

Mark-recapture

During the summer of 2009, we tagged 30 glacier moss balls with a bracelet identifier

(Fig. 1d). Each bracelet consisted of a unique combination of colored glass beads (~2-3 mm in

diameter) threaded on aluminum wire. Bracelets were threaded through the moss ball center

and pulled snug so as to not protrude beyond the moss ball's exterior and interfere with

movement. We returned eight times during the 2009 season to re-survey moss balls and record

their movements. We followed up our initial surveys with annual visits from 2010-2012. During

each survey, we visually inspected in and around the core study area multiple times in an effort

to recapture moss balls. As part of this process, we visually inspected each moss ball in the

area for any sign of a bracelet tag. After inspection, we replaced each moss ball in the exact

location and orientation as it was found.

Moss ball movement and glacier ablation

We assessed moss ball movement over 54 days in 2009. As benchmarks for their

movement, we installed three ~1.3 cm PVC tubes into the glacier. Each stake was drilled ~60

cm into the glacier. Stakes were installed in a triangle that spanned the study area and served

two purposes. First, the stakes provided a reference against which the location of each moss

ball was measured. Second, they allowed us to measure glacier ablation (i.e., the distance the

ice surface moves vertically down) over the same study period so we could test for links

between moss ball movement and the rate of glacier ablation.

To track glacier moss ball movement, we measured the distance between re-identified

moss balls and each of the reference stakes for each visit to the site. Next, for each moss ball,

we calculated three independent positions within our field site--one for each of the three pairs of

reference stakes. We assigned the location of a surveyed moss ball to the mean of these three

relative positions and constructed a location covariance matrix for each measurement, to assign

uncertainties to surveyed locations. After diagonalizing the covariance matrix, we identified the

size (eigenvalues) and orientation (eigenvectors) of an uncertainty ellipse around each mean

location. Major and minor axes of the uncertainty ellipse were defined as twice the square root

of the eigenvalue lengths, such that each error ellipse represented a 2σ error window. Thus,

assuming independent, normal errors, we are 95% confident that the true location of each moss

ball fell within its error ellipse. While we used stakes for most of the measurement period, we

were forced to switch to washers (~5 cm in diameter) laid flat on the ice surface later in the

season, during a period when we were unable to drill the benchmark stakes sufficiently deep to

avoid melting out between visits to the study area. Before transitioning from benchmark stakes

to washers, we tested the stability of the washers to ensure that they did not slide over the ice

surface. Over a 5-day period, we did not detect significant washer movement (outside of 2σ

uncertainty). Final measurements (11 August 2009) and calculations were made relative to the

washers.

For the purposes of quantifying glacier ablation, the height of each stake above the local

ice surface was re-measured during each visit and periodically re-drilled into the ice as

necessary. Ablation reported in this study is the mean ice surface lowering rate calculated for

each of the three stakes. As an assessment of ablation uncertainty, we also calculated the

maximum deviation of any single stake's ablation rate from the overall mean.

Longevity

We sought to understand how long glacier moss balls survive, particularly across

individual seasons. We hypothesized that moss balls might survive better or worse in some

years due to variation in environmental conditions (e.g., precipitation, freeze-thaw cycles) or

random chance (e.g., a crevasse opening within a key area). Furthermore, we wanted to know

not only how likely we are to detect glacier moss balls, given that they had persisted within the

study area, but also if our detection probability varies among years. To do this, we fit capture-

recapture models of annual survival to each glacier moss ball included in the study. Because

moss balls were individually marked but were not equipped with radio-transmitters or other

devices which would allow us to know their ultimate fates, we applied Cormack-Jolly-Seber

(CJS; Lebreton et al. 1992) survival models. These CJS models develop a "capture history" of

each moss ball to estimate apparent survival (i.e., the probability that an individual is in the

population at time i and still in the population at time i+1) and probability of detection if they

persisted within our study area. Survival estimates from CJS models only represent apparent

survival because emigration cannot be estimated from survival data with unknown fates (i.e., we

did not know if a tagged moss ball had disaggregated, lost its identifying bracelet, or was no

longer in the study area). Therefore, our estimates of apparent survival are likely to

underestimate true survival (e.g., a moss ball might have lost its bracelet or moved out of the

study site). In addition, CJS models also account for imperfect detection, which in our case

would be if a moss ball persisted within our study area but was overlooked.

Using our individual moss ball annual detection data (1 = detected, 0 = not detected), we

fit four competing CJS survival models, including the null model [no effect of year on apparent

survival ($\square$) or detection probability (p); Model 1)], an effect of year on $\square$ (Model 2), an effect of

year on p (Model 3), or an effect of year on both $\square$ and p (Model 4). We then selected the

model(s) best supported by our data using Akaike's information criterion (AIC; Akaike 1998),

adjusted for small sample size (AICc; Hurvich and Tsai 1989). Our model selection approach

was based on model likelihoods and models were penalized for extra parameters to favor

parsimony.

Finally, we calculated the average life expectancy of a glacier moss ball. To do this, we

used annual survival rates based on life-table analysis (Deevey Jr 1947; Millar and Zammuto

1983), in which average life expectancy was calculated as -1/ln(Annual Survival Rate). Because

this estimation of life expectancy is quite sensitive to annual survival rate, we calculated it for

both the lowest annual survival rate and the mean annual survival rate. Thus, the true average

life expectancy might be substantially greater than the conservative values estimated here. This

framework for estimating average life expectancy does not account for potentially higher or

lower mortality rates when glacier moss balls are first forming or nearing the end of their

lifespans.

Results:

 Study area

Our study area was located on a "bare ice" glacier surface, between two medial

moraines covered by coarse-grained, angular, rock debris. However, two types of sediment

distinguish the study area surface from what would be considered clean, pure, water ice. First,

glacier moss balls were found amidst gravel and small boulders (< 30 cm diameter), spaced

every ~1 m. Second, the ice surface within the study area has an unusually pervasive, fine-

grained sediment cover, ~1-3 mm thick, which partially blankets the otherwise bare ice. Image

processing indicated that this fine sediment covers approximately 70% of the study area

surface. This low albedo sediment cover is visible in all inspected satellite imagery of the site

and first appears at lower concentrations emerging from cleaner ice ~1 km northwest of the

study site (Fig. 1b). Down-glacier of the study site, the low albedo region extends ~1.7 km as a

~300-m-wide, rounded finger that spans adjacent medial moraines, in a manner consistent with

wind-deposited dust, draping over underlying geomorphic features. Therefore, we interpreted

the southeast (135°) trend direction of this low albedo finger to be the prevailing, down-glacier,

katabatic wind direction.

During the 26 days of glacier ablation measurements, the ice surface lowered by 1.91 m

due to melt and sublimation. Ablation rates ranged from 5.8-9.6 cm per day (cm/d) between

measurement times, and averaged 7.3 cm/d.

Fig. 2. (A) Locations of surveyed glacier moss balls throughout the survey period. Most likely locations of
each moss ball are shown with small filled circles relative to an arbitrary, local grid system. Ellipses
surrounding each moss ball indicate 2 uncertainty (i.e., 95% confidence) of their location. Thin black

lines connect consecutive surveyed locations for individual moss balls. The red rectangle identifies the
location of the large-scale view in panel (B). (B) A zoomed in view of movement patterns for six glacier
moss balls (red square in A), showing their similar azimuths.

Fig. 3. (A) A comparison of glacier moss ball movements versus the dominant wind (dashed red line) and
downslope (dashed blue line) directions. Direction of each moss ball's motion between measurement
times is shown with thin gray lines, while the bold black line indicates the median direction of all glacier
moss ball movements. (B) Glacier moss ball movement versus ablation rate. Median ablation rate is
indicated with a bold red line, while the mean +/- the maximum absolute deviation from the mean are
shown with thin red lines. The median speed of glacier moss balls is shown with the bold blue line, while
the 25th and 75th percentile speeds are shown with thin blue lines. Numbers in circles along the bottom
of the plot represent the number of moss balls surveyed at each timepoint (single measurements not
indicated).

Movement

Glacier moss ball movements varied systematically over the study period, with increases
and decreases that coincided with changes in direction (Figs. 2-3). Median moss ball speed was
2.5 cm/d, but their rates varied widely throughout the season. The median speed started at 1.8
cm/d in late June, increased to 4.0 cm/d at the start of July, then slowed to 2.0 cm/d during late
July/early August. The maximum observed speed for any glacier moss ball was 7.8 cm/d during
the 5-day period from July 9-14 (excluding two outlier speeds that were more than 8 interquartile
ranges greater than the median, 14.2 and 21.0 cm/d, and which were based upon particularly
uncertain moss ball positions). The interquartile range of moss ball speeds was approximately
50% of the median speed; thus, these observed increases and decreases in speed reflect
changes in the entire population of moss balls.

The direction of glacier moss ball movements was not random. Rather, glacier moss
balls underwent clear changes in their direction of motion (i.e., azimuth) throughout the summer
season (Fig. 3a). While individual moss balls moved in many directions, when viewed in
aggregate, azimuths of the population clearly cluster over time. Early in the season, median
moss ball motion was south-southeast (165°) but over the ensuing weeks azimuths
progressively increased, such that at the end of the measurement period the median azimuth
was west-southwest (240°; Fig. 3a).

Considering speeds and azimuths together, we see the moss ball population initially
moving at 2 cm/d to the south for 9 days, then the group nearly doubles its speed to 4 cm/d
while deviating slightly to the right (towards the west). After a week at these maximum speeds,
speeds drop by 25% to 3 cm/d while also deviating 45 degrees further towards the west for five
days. During the next 5-day measurement period, speeds drop further, back to 2 cm/d while the
azimuths turn another 10-15 degrees further west. Over the final 28-day measurement period,
the azimuths remain stable, while speeds continued to fall. This decrease in speed is apparent
in the decline of the upper quartile of speeds, despite our not making sufficient new
measurements to influence the median speed.

Our fine-scale movement and ablation data allowed us to compare glacier moss ball
speeds and azimuths with potential drivers of their motion. The southern and western directions
of moss ball movement are clearly distinct from both the prevailing wind direction as inferred
from the dust plume (towards the southeast) or the downhill direction of the gently sloping ice
surface (towards the east-northeast; Fig. 3a). Instead, we find more rapid moss ball speeds are
associated with more rapid ablation; an ordinary least squares model between ablation rate and

speed indicates that, on average, for every 1 cm of surface ablation, the glacier moss balls

move horizontally 0.34 cm (Fig. 3b). However, the relationship between ablation rate and speed

is relatively weak (R^2 = 0.40). It should also be noted that during the course of our study,

participants in a program hosted by the Wrangell Mountains Center, McCarthy, Alaska, visually

confirmed the posited primary movement method described by Porter et al. (2008), when a

glacier moss ball was observed rolling off its elevated pedestal and inverting in the process.

Table 1. Apparent survival models for glacier moss balls tested in this study with their
corresponding Akaike's Information Criterion scores that have been adjusted for small sample
sizes (AICc). Relative AICc scores (ΔAICc) model weight are also given. Lower ΔAICc and
higher model weight indicate greater support for a given model. Model components: probability
of detection (p), apparent survival (ϕ).

Model	Description	AICc	ΔAICc	Weight
1	Null; No year effect on p or ϕ	107.09	1.56	0.26
2	Year effect on ϕ	105.53	0	0.58
3	Year effect on p	108.92	3.39	0.10
4	Year effect on both p and ϕ	110.25	4.72	0.05

301
302

303 **Table 2**. Estimates of the apparent survival (ϕ) and detection probability (p) of glacier moss balls
304 for the two best-fit models. Parentheses after estimates indicate 95% confidence intervals.

Model	Parameter	Estimate
1	p	0.84 (0.70-0.92)
	ϕ	0.86 (0.75-0.93)
2	p	0.82 (0.69-0.91)
	ϕ (2009-2010)	0.74 (0.55-0.87)
	ϕ (2010-2011)	0.98 (0.27-0.99)
	ϕ (2011-2012)	1.0 (0.99-1)

Longevity

We initially tagged 30 glacier moss balls in 2009. We subsequently recaptured 18 moss

balls each in 2010, 2011, and 2012 (although this was not the same 18 moss balls each year).

Recapture rates for individual glacier moss balls were highly variable with some never seen

again after the first year (n = 8) and others detected every year (n = 13). The best-fit survival

model included differing apparent survival (□) among years, but with constant detection

probability (p; Model 2; Table 1). This model received 58% of AICc weight, compared to 26% for

the null model (Model 1), and less than 10% for the other models (Models 3 & 4; Table 1). The

average annual rate of apparent survival, □, based on the null model, was 0.86 [95%

confidence interval (CI) = 0.75-0.93], and the average detection rate was 0.84 (95% CI = 0.70-

0.92). When parameterized by year, the annual apparent survival rate ranged from 0.74 in

2009-2010 to 1.0 in 2011-2012 with a particularly large 95% CI for 2010-2011 (Table 2; Fig. 4).

Fig. 4. Estimates of apparent moss ball survival (☐; dark circles) with 95% confidence intervals (thin dark
lines) from model 2, the best-fit model, which included a year effect on ☐. Year-long, bracketed time
intervals labeled on the x-axis are identified by their starting year. For instance, apparent survival for
2009-2010 is shown as 2009.
324

Our detection rate estimates may underestimate actual glacier moss ball survival for

several reasons. First, at least four glacier moss balls lost their marking bracelet after the first

year because we found the marking bracelet on the ice, separate from a moss ball. Second,

another moss ball partially obscured its bracelet by growing to cover the beads, but we were

able to detect a single bead and then delicately "excavate" the bracelet. Since we did not

destructively search glacier moss balls that did not have an obvious bracelet, it is possible that

additional instances of lost marking bracelets or growth to cover beads may have impacted our

detection. Third, between 2009 and 2010, two tagged moss balls fell inside of a shallow

crevasse within the study area. The two crevasse-bound glacier moss balls persisted, and likely

continued to photosynthesize and grow to some capacity for the remainder of the study. We

continued to check crevasses in the study area carefully, but some moss balls could have fallen

into deeper crevasses, or into shallow crevasses in a way that obscured their markings, and

therefore persisted without detection.

Our estimate of average life expectancy varied depending on whether the lowest overall

or mean annual survival rate were used. If using the lowest annual survival rate (0.74), average

life expectancy was 3.3 years (95% CI = 1.67-7.18). However, we expect this life expectancy to

be biased low to some extent, because we were only able to estimate apparent survival (e.g.,

some insecure tags fell off moss balls that likely still persisted). If using the mean annual

apparent survival rate across the entire study (0.86), average life expectancy rose to 6.63 years

(95% CI = 3.48-13.78), although this may be biased high because we did not tag any new moss

balls in years 2 and 3 (2010 and 2011), but simply re-captured existing (and therefore high

survival probability) glacier moss balls. When thinking of lifespan, it is relevant to note that we

also observed a glacier moss ball split roughly in half during the course of the study along its

intermediate axis. The moss ball had become elongated and essentially pulled apart. This

mechanism may contribute to both keeping glacier moss balls ovular as well as a form of

cloning, propagating new moss balls on the landscape.

Discussion:

Glacier moss balls are intriguing components of glacier ecosystems that integrate

physical (e.g., debris cover) and ecological (e.g., invertebrate colonization) factors into a unique

habitat type. Glacier moss balls have a global distribution, with colonies identified in Iceland,

North and South America, and Asia (Eythórsson 1951; Heusser 1972; Perez 1991; Porter et al.

2008; Coulson and Midgley 2012) and more dispersed moss aggregations on glaciers have

been described from an even broader area (e.g., Uganda, Uetake et al. 2014). Previous studies

have revealed a great deal about glacier moss ball biology (e.g., their invertebrate colonizers,

Coulson and Midgley 2012) yet their movement and longevity has remained unknown. It has

been speculated that glacier moss ball movement patterns likely follow the general downward

slope of the glacier (Porter et al. 2008) and that they represent an ephemeral habitat type on

glaciers, a factor that may limit colonization by specific invertebrate taxa (e.g., spiders; Coulson

and Midgley 2012). Our results do not align with these predictions of movement and longevity.

Rather, we show that glacier moss balls, at least on a relatively gently sloped Alaskan glacier,

exhibit relatively quick (2.5 cm/d) herd-like movements that do not align with the downward

slope of the glacier nor the dominant wind direction. Glacier moss balls are also relatively long-

lived with a mean lifespan of more than 6 years.

Movement

On the Root Glacier, glacier moss balls move relatively quickly (~2.5 cm/d) in similar

general directions and at similar general speeds. Directions of motion do not align solely with

either the downhill direction nor the direction of the prevailing wind. The rate of glacier moss ball

movements is also positively correlated, albeit weakly, with overall glacier ablation (Fig. 3b). It

appears likely that the dominant direction of solar radiation, which melts exposed ice

surrounding glacier moss balls more rapidly than the insulated ice below them (Porter et al.

2008), is the major force driving glacier moss ball movement. However, the relative

contributions of gravity in the downslope direction versus solar radiation is almost certainly

dependent on glacier steepness. Porter et al. (2008) posited a considerable effect of gravity on

glacier moss ball movement for a relatively steep (9.6°) Icelandic glacier which contrasts with

our much flatter study area on the Root Glacier (~3°). Still, regardless of steepness, differential

melt patterns create pedestals that glacier moss balls rest upon and, eventually, enough ice

melts below the moss ball causing it to fall and potentially flip (Porter et al. 2008). Assuming

glacier moss balls are, on average, ~10 cm in their intermediate axis, and their only means of

movement is melt-induced flipping driven by pedestal emergence at the rate of 6-9 cm/d, their

rates of movement would imply each glacier moss ball flips every ~2-4 days. However, we

cannot rule out alternative modes of glacier moss ball movement. Many glacier moss balls have

one side that is flattened and commonly faces down, while the other, more rounded and

vegetated side faces skyward (Shacklette 1966). Given this orientation, an alternative scenario

is that glacier moss balls also move by basal sliding over the wet glacier surface below.

One movement-related question remains puzzling: why do the azimuths of glacier moss

balls appear to shift simultaneously throughout the summer season, resulting in the moss ball

"herd" synchronously changing directions (Fig. 3a)? They begin the season moving generally

south and slowly transition towards the west. Given their independence from the dominant wind

direction and downhill direction of the glacier, we can speculate that a slow shift in the dominant

direction of solar radiation or shifting weather patterns during summer drives this pattern.

Perhaps the weather transitioned from clear mid-day skies during late June and early July

(associated with the most rapid motion and southerly azimuths), to a different weather pattern in

late July consisting of morning clouds and afternoon sun, that drives enhanced ablation on the

west sides of moss balls, and therefore preferential rolling towards the west. Furthermore, the

interaction between rate of movement and ablation may depend additionally on the degree to

which the dominant solar radiation and downslope directions align. If they are in the same

direction, glacier moss balls should move rapidly, downslope, towards the sun. However, if melt

is being driven from a direction perpendicular to the downslope direction, then glacier moss

balls will move considerably less per unit melt.

Longevity

Glacier moss balls show considerable potential to persist across multiple years as stable

ecological units. On average, 86% of the marked glacier moss balls included in this study

survived annually which translates to a lifespan of more than 6 years. This longevity is on par

with large, long-lived vertebrates (e.g., adult female ungulates in typically have annual survival
probabilities ≥ 0.85; Gaillard and Yoccoz 2003; Gaillard et al. 1998). Thus, with high rates of
survival across multiple years, and relatively high detection rates, we consider glacier moss
balls to be long-lived, rather than ephemeral, glacier features. Unlike living individual organisms
which can show senescence as they age (e.g., Loison et al. 1999), moss ball survival rates are
unlikely to decline with time in the traditional sense, nor are they likely to exhibit density
dependence in survival (e.g., Festa-Bianchet et al. 2003), however the factors that control
disaggregation may be the most important factor for moss ball longevity. At any rate, the
temporal stability of individual moss balls on the glacier surface means that they likely exist for
long enough to develop complex biotic communities, a fact supported by the distinctive
invertebrate communities inhabiting moss balls (Coulson and Midgley 2012). However, the
degree to which geographic location (e.g., distance to a glacier margin), and not persistence,
influences invertebrate colonization remains to be tested.

The limited scope of our mark-recapture data collection precludes us from drawing
conclusions about the inter-annual drivers of moss ball apparent survival. However, we can
highlight potential factors that may influence it. First, it is possible that glacier moss balls moved
more frequently out of the study area in one year versus others, perhaps due to exceptionally
clear skies (and thus higher rates of glacier ablation). Second, we observed a number of
fragmented moss balls. This fragmentation may be part of normal glacier moss ball growth
trajectories, too frequent or intense freeze thaw cycles, perhaps as a product of moss ball water
content, or some other as yet unknown factor. It is also unclear at what rate fragmented glacier
moss balls continue to move and grow, eventually developing back into mature, ovoid, full-sized
moss balls. If glacier moss balls did survive within our study area, they had an 84% probability
of being detected in a given year. This indicates that our bracelet and colored beads marking
scheme was relatively successful. However, for future studies, more robust marks should be
considered. One possibility is the use of passive integrated transponder (PIT) tags which are
commonly used for mark-recapture studies of a variety of organisms (e.g., fish; Castro-Santos
et al. 1996), and allow researchers to scan study organisms electronically rather than rely on
visual ID.
440
Genesis, growth, and disaggregation
Our combined movement and longevity analyses allow us to add new speculation about
patterns of glacier moss ball growth as well as additional evidence for previous hypotheses
regarding their genesis and disaggregation (e.g., Heusser 1972; Perez 1991). In terms of

growth, our documentation of glacier moss balls rolling over a fine-grained, wet, sedimentary

substrate is consistent with growth through the adherence of fine-grained sediment to an

existing moss ball. When a moss ball rolls from its elevated pedestal, sediment grains stick to

the moss (or potentially to the cohesive sediment itself). We visually observed such "dirty" moss

on some glacier moss balls in our study area. As the moss itself grows, this adhered sediment

may then become integrated within the fibrous material, increasing the size of the glacier moss

ball. Field observation of moss growth over and around our identification bracelets indicates that

several millimeters of growth can occur within years. However, the observation that most

bracelets were not engulfed by sediment accumulation and moss growth during our four-year

study period suggests an upper limit on moss ball growth. Consistent with a greater than 6-year

lifespan, moss ball growth to an observed size of 10 cm must take years, and given the

conservative nature of our longevity estimates, potentially much longer.

Understanding year-to-year moss ball growth, however, does not explain moss ball

genesis, nor disaggregation. It is well-established that fibrous moss provides the skeletal

structure that allows moss balls to be cohesive, ovoid structures. A source of moss spores is

there essential for initial glacier moss ball genesis (in our study, putatively, the Donoho

nunatak). The question, then, is how glacier moss balls begin to grow in the first place, and on

what substrate. (Eythórsson 1951) suggested that a "stone kernel" is likely at their centers.

However, later investigations (e.g., Shacklette 1966; Gremmen 1982) found mixed results that

largely reflected a consensus that there is no general rule about rock cores at the center of

glacier moss balls. Our exploratory testing of moss balls also indicated that some, but not all,

moss balls contained a ~1-cm gravel "kernel" at their centers. Potentially, these kernels, with

adhered fine-grained sediment, provide a growth substrate for initially wind-deposited moss

spores. In our study area, the co-occurrence of moss balls with an unusually extensive, fine-

grained "plume" of sediment cover (Fig. 1b) aligns with a similar observation by Heusser (1972)

for the Gilkey Glacier in southeastern Alaska, USA. The fine-grained sediment may be essential

proto-soil for moss ball growth and may explain the unusual density of moss ball occurrence at

our study area. The origin of this fine-grained sediment is unknown, but in satellite imagery (Fig.

1b), it appears to originate from the ice itself. It may be a volcanic ash layer emerging from the

ice after being carried down from the high, volcanic, Wrangell Mountain peaks. Once moss

growth initiates, the moss itself, and the abundant moisture of the ablating glacier surface, may

provide the necessary cohesion for the incipient moss ball formation, and continued growth

thereafter.

In this study, we identified very few glacier moss balls greater than ~15 cm on their long
axis. Generally, moss balls appear to rarely exceed ~10 cm except for rare cases in Alaska
where they have been reported up to 18 cm (Heusser 1972; Benninghoff 1955). Why glacier
moss balls in Alaska appear to grow larger than elsewhere in the world remains an open
question but, regardless of location, there appears to be some size limiting process within the
moss ball lifecycle at work. Shacklette (1966) suggested that the tensile strength of moss stems
may be the key factor controlling their size. Exceeding this tensile limit appears to occur when
the moss ball major axis grows too great relative to the intermediate axis. For instance, when a
moss ball becomes too elongated, subtle variations in ice surface topography may lead the two
ends of a moss ball to move in different directions, leading to a tear in the middle when the
moss ball's tensile strength is exceeded. We observed such a tearing-in-two of a long, linear
moss ball during the course of our study. Again, while this process applies an upper-limit to
moss ball size it also circles back to inform questions regarding the presence or absence of a
rock kernel. If the upper size limit is reached and a moss ball splits, only one of the two
remaining moss balls involved in this "cloning" process will retain the gravel kernel. This may
explain why a number of moss balls do not appear to have any coarse-grained rock at their
cores.
495
Conclusions
Since their first description nearly 75 years ago (Eythórsson 1951), a general
understanding of the physical and biological composition of glacier moss balls and their
ecological role as drivers of small-scale ecosystem development has been established. In this
study, we extended this previous work to quantify the movement and longevity of glacier moss
balls on an Alaskan glacier. In light of these results, we discussed a potential life cycle for moss
balls. We showed that glacier moss balls move relatively quickly, at a rate of centimeters per
day, and that moss balls within the surveyed colony speed up, slow down, and change direction
in synchrony. This finding suggests that moss ball motion is controlled by some broadly applied
external forcing. However, for our study, this external forcing was surprisingly not solely
associated with either the wind or downslope direction. Instead, the glacier moss ball movement
patterns we observed likely depend on a combination of the intensity of glacier ice ablation, the
direction of solar radiation, and the physical surface of the glacier (i.e., the downslope direction).
Future studies that take a similar mark-recapture approach to the study of glacier moss ball
movements on multiple glaciers throughout the world, including a range of steepness, will shed
important light on the general nature of their movements.

We also showed that glacier moss balls are long-lived, with an average life expectancy

of more than 6 years, on par with many relatively long-lived vertebrates (e.g., large mammalian

herbivores, Gaillard and Yoccoz 2003). This potential for glacier moss balls to act as relatively

stable, long-term ecological units further confirms their potential to act as key habitat for

invertebrates. Coulson and Midgley (2012) previously described invertebrate colonization of

glacier moss balls and suggested that a lack of Enchytraeidae and Aranea may be the result of

the ephemeral nature of moss balls in glacier habitats. Our results contrast with this assumption.

Instead, we postulate that selective invertebrate colonization of glacier moss balls may depend

instead on their geographic locations and their frequent movements or, as Coulson and Midgley

(2012) noted, may simply reflect the variable dispersal capacities of potential colonizers.

Given the importance of microbial diversity to carbon cycling (Anesio et al. 2009),

ecosystem function (Anesio et al. 2017; Hotaling et al. 2017a; Hotaling et al. 2017b), and even

albedo (Ganey et al. 2017), future efforts to understand the microbial ecology of glacier moss

balls could shed important light on their ecological role in glacier ecosystems. Like cryoconite,

the granular, darkly pigmented dust that accumulates on the surface of glaciers and drives

hotspots of biological activity and microbial diversity (Cook et al. 2016), glacier moss balls may

have similar value at the ecosystem scale. Indeed, glacier moss balls may act as reservoirs of

microbial diversity, seeding and re-seeding a glacier surface with key, colonizing microbial life

as they move around its surface.

Acknowledgements:

We thank the Wrangell Mountains Center for logistical support and assisting with field

measurements, and Dr. Billy Armstrong for providing the orthoimage of the study area.

References:

Akaike H (1998) Information theory and an extension of the maximum likelihood principle. In:
Selected papers of Hirotugu Akaike. Springer, pp 199-213
Anesio AM, Hodson AJ, Fritz A, Psenner R, Sattler B (2009) High microbial activity on glaciers:
importance to the global carbon cycle. Global change biology 15:955-960
Anesio AM, Laybourn-Parry J (2012) Glaciers and ice sheets as a biome. Trends Ecol Evol
27:219-225
Anesio AM, Lutz S, Chrismas NA, Benning LG (2017) The microbiome of glaciers and ice
sheets. NPJ biofilms and microbiomes 3:10
Belkina OA, Vilnet AA (2015) Some aspects of the moss population development on the
Svalbard glaciers. Czech Polar Reports 5:160-175
Benninghoff WS (1955) Jökla mýs. Journal of Glaciology 2:514-515
Castro-Santos T, Haro A, Walk S (1996) A passive integrated transponder (PIT) tag system for
monitoring fishways. Fisheries Research 28:253-261

Cook J, Edwards A, Takeuchi N, Irvine-Fynn T (2016) Cryoconite: the dark biological secret of
the cryosphere. Progress in Physical Geography 40:66-111
Coulson S, Midgley N (2012) The role of glacier mice in the invertebrate colonisation of glacial
surfaces: the moss balls of the Falljökull, Iceland. Polar Biology 35:1651-1658
Deevey Jr ES (1947) Life tables for natural populations of animals. The Quarterly Review of
Biology 22:283-314
Dial RJ, Becker M, Hope AG, Dial CR, Thomas J, Slobodenko KA, Golden TS, Shain DH (2016)
The role of temperature in the distribution of the glacier ice worm, Mesenchytraeus
solifugus (Annelida: Oligochaeta: Enchytraeidae). Arctic, Antarctic, and Alpine Research
48:199-211
Eythórsson J (1951) Correspondence. Jökla-mys. Journal of Glaciology 1:503
Festa-Bianchet M, Gaillard JM, Côté SD (2003) Variable age structure and apparent density
dependence in survival of adult ungulates. Journal of Animal Ecology 72:640-649
Gaillard J-M, Festa-Bianchet M, Yoccoz NG (1998) Population dynamics of large herbivores:
variable recruitment with constant adult survival. Trends in Ecology & Evolution 13:58-63
Gaillard J-M, Yoccoz NG (2003) Temporal variation in survival of mammals: a case of
environmental canalization? Ecology 84:3294-3306
Ganey GQ, Loso MG, Burgess AB, Dial RJ (2017) The role of microbes in snowmelt and
radiative forcing on an Alaskan icefield. Nature Geoscience 10:754
Gardner AS, Moholdt G, Cogley JG, Wouters B, Arendt AA, Wahr J, Berthier E, Hock R, Pfeffer
WT, Kaser G (2013) A reconciled estimate of glacier contributions to sea level rise: 2003
to 2009. Science 340:852-857
Gremmen N (1982) The vegetation of Subantarctic Islands Marion and Prince Edward. The
Hague, Boston, London, W. Junk Publishers,
Heusser CJ (1972) Polsters of the moss Drepanocladus berggrenii on Gilkey Glacier, Alaska.
Bulletin of the Torrey Botanical Club:34-36
Hotaling S, Finn DS, Joseph Giersch J, Weisrock DW, Jacobsen D (2017a) Climate change and
alpine stream biology: progress, challenges, and opportunities for the future. Biological
Reviews 92:2024-2045
Hotaling S, Hood E, Hamilton TL (2017b) Microbial ecology of mountain glacier ecosystems:
biodiversity, ecological connections and implications of a warming climate.
Environmental microbiology 19:2935-2948
Hotaling S, Shain DH, Lang SA, Bagley RK, Lusha M, Weisrock DW, Kelley JL (2019) Long-
distance dispersal, ice sheet dynamics, and mountaintop isolation underlie the genetic
structure of glacier ice worms. Proceedings of the Royal Society B: Biological Sciences
286:20190983
Hurvich CM, Tsai C-L (1989) Regression and time series model selection in small samples.
Biometrika 76:297-307
Larsen C, Burgess E, Arendt A, O'neel S, Johnson A, Kienholz C (2015) Surface melt
dominates Alaska glacier mass balance. Geophysical Research Letters 42:5902-5908
Lebreton J-D, Burnham KP, Clobert J, Anderson DR (1992) Modeling survival and testing
biological hypotheses using marked animals: a unified approach with case studies.
Ecological Monographs 62:67-118
Loison A, Festa-Bianchet M, Gaillard J-M, Jorgenson JT, Jullien J-M (1999) Age-specific
survival in five populations of ungulates: evidence of senescence. Ecology 80:2539-2554
Mann D, Edwards J, Gara R (1980) Diel activity patterns in snowfield foraging invertebrates on
Mount Rainier, Washington. Arctic and Alpine Research 12:359-368
Millar JS, Zammuto RM (1983) Life histories of mammals: an analysis of life tables. Ecology
64:631-635

Perez FL (1991) Ecology and morphology of globular mosses of Grimmia longirostris in the
Paramo de Piedras Blancas, Venezuelan Andes. Arctic and Alpine Research 23:133-
148
Porter P, Evans A, Hodson A, Lowe A, Crabtree M (2008) Sediment–moss interactions on a
temperate glacier: Falljökull, Iceland. Annals of Glaciology 48:25-31
Roe GH, Baker MB, Herla F (2017) Centennial glacier retreat as categorical evidence of
regional climate change. Nature Geoscience 10:95
Rosvold J (2016) Perennial ice and snow-covered land as important ecosystems for birds and
mammals. Journal of biogeography 43:3-12
Shacklette HT (1966) Unattached moss polsters on Amchitka Island, Alaska. Bryologist:346-352
Uetake J, Tanaka S, Hara K, Tanabe Y, Samyn D, Motoyama H, Imura S, Kohshima S (2014)
Novel biogenic aggregation of moss gemmae on a disappearing African glacier. PloS
one 9:e112510
Van der Walt S, Schönberger JL, Nunez-Iglesias J, Boulogne F, Warner JD, Yager N, Gouillart
E, Yu T (2014) scikit-image: image processing in Python. PeerJ 2:e453
614