Simultaneous Targeting of XPO1 and BCL2 as an Effective Treatment Strategy for Double-Hit Lymphoma

Yuanhui Liu[1,2], Nancy G. Azizian[1,2], Yaling Dou[3], Lan V. Pham[4], Yulin Li[1,2*]

[1]Center for Immunotherapy Research, Houston Methodist Cancer Center, Houston, TX 77030, USA, [2]Department of Medicine, Weill Cornell Medical College, New York, NY 10065; [3]Center for Transplantation Immunology, Houston Methodist Research Institute, Houston, TX 77030; [4]Department of Hematopathology, Division of Pathology and Laboratory Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX 77030.

Running title: Combined targeting of XPO1 and BCL2 in DHLs

Key Words: Double-hit lymphoma, MYC, BCL2, XPO1, Patient-derived xenograft, brain metastasis, targeted combination therapy,

*Corresponding author: Yulin Li, Houston Methodist Cancer Center. 6670 Bertner Avenue. Houston, TX 77030. Phone: 713-441-7350; Fax: 713-441-7438; Email: yli@houstonmethodist.org

Abstract

Double-hit lymphoma (DHL) is among the most aggressive and chemoresistant lymphoma subtypes. DHLs carry genomic abnormalities in MYC, BCL2 and/or BCL6 oncogenes. Due to simultaneous overexpression of these driver oncogenes, DHLs are highly resistant to the frontline therapies. Most DHLs overexpress both MYC and BCL2 driver oncogenes concurrently. We reasoned that simultaneous suppression of the two driver oncogenes would be more effective in eradicating DHLs than inactivation of single oncogene. XPO1 is a receptor for nuclear cytoplasmic transport of protein and RNA species. Recently, XPO1 inhibition was shown to downregulate MYC expression in several cancer cell lines. We therefore examined the role of XPO1 as a therapeutic target in suppressing MYC function and the potential synergistic effects of simultaneous suppression of XPO1 and BCL2 in the treatment of DHL. Here, we demonstrate that XPO1 inhibition abrogates MYC protein expression and induces massive tumor cell apoptosis. Combined use of XPO1 and BCL2 inhibitors is highly effective in eradicating DHL cells in cell culture. Notably, in a mouse model of DHL bearing primary tumor cells derived from lymphoma patients, combined treatment with XPO1 and BCL2 inhibitors blocks tumor progression, prevents brain metastasis, and extends host survival. Thus, our study confirms the simultaneous targeting of MYC and BCL2 driver oncogenes through the combined use of XPO1 and BCL2 inhibitors as a unique approach for the treatment of DHLs.

Introduction:

Double-hit lymphoma (DHL) is a subtype of Non-Hodgkin's Lymphoma (NHL) with genomic abnormalities in MYC and BCL2 (and less frequently BCL6), leading to the overexpression of these driver oncogenes. The prognosis for the majority of NHLs has improved significantly in the past decades due to the development of chemotherapy, targeted therapy, and immunotherapy. In comparison, DHL remains highly resistant and refractory to the first line immunochemotherapy, R-CHOP, with a 5-year overall survival rate of 30%[1,2].

The co-existence of multiple oncogenic events in DHLs, including MYC, BCL2, and BCL6, provides an opportunity for combined targeted therapy. As the two common drivers of DHL, MYC and BCL2 cooperate in lymphomagenesis and tumor maintenance. A combined therapy aimed at targeting both MYC and BCL2 may arguably be more effective than suppressing either MYC or BCL2 alone in eradicating the tumors cells[3,4]. Among several BCL2 inhibitors, ABT-199, was developed and tested in clinical trials, and approved by FDA for the treatment of chronic chronic lymphocytic leukemia[5]. In contrast, direct targeting of MYC has proven challenging due to its structural property as a transcription factor.

XPO is an adaptor of nuclear export for many protein and RNA species. Recent studies suggest that XPO1 may regulate nuclear export of mRNAs encoding several oncoproteins, such as MYC, BCL2, Cyclin D1, and PIM1[6-8]. Moreover, suppression of XPO1 with selective inhibitors of nuclear export (SINEs) has been shown to downregulate MYC expression in several tumor types[9-12]. We therefore hypothesize that tumor cells with concurrent MYC and BCL2 overexpression, such as DHL, may be effectively targeted through the combined treatment with XPO1 and BCL2 inhibitors.

Here, we demonstrate that XPO1 inhibition in DHL tumor cells abrogates MYC protein expression. Furthermore, combined suppression of XPO1 and BCL2 synergizes to induce massive cell death in DHL tumor cells *in vitro*. Most importantly, combination therapy with XPO1 and BCL2 inhibitors blocks tumor progression and metastasis, and extends survival in a mouse model bearing patient-derived DHL tumors. Our data suggests that combined targeting of XPO1/BCL2 can be a robust therapeutic strategy for DHLs carrying both MYC and BCL2 driver oncogenes.

Results:

XPO inhibition abrogates MYC protein expression and induces apoptosis in DHLs

First, we examined whether XPO1 inhibition affects MYC protein levels in DHL cell lines. Treatment with the XPO1 inhibitor, KPT8602, led to a significant decrease in MYC protein expression in the majority of DHLs in our diverse cell line panel (**Fig 1A**). Three DHL cell lines (SU-DHL4, Toledo, and SU-DHL6) were selected to further examine MYC regulation by XPO1 inhibitors. Treatment with two specific XPO1 inhibitors, KPT330 and KPT8602, points to a dose- and time- dependent downregulation of MYC protein expression in all three cell lines (**Fig 1B-C**, and **Supplementary Fig S1**). These results establish that XPO1 inhibition effectively abrogates MYC protein expression in DHL tumor cells. Decreased MYC protein level was followed by changes in gene expression of MYC downstream targets (**Fig 1D-E**). Genes known to be upregulated by MYC, including ENO1, APEX1, RPL3, RPS5, SRM and nucleolin[13], were significantly downregulated upon XPO1 inhibition. In contrast, genes reportedly suppressed by MYC, such as HBP1, P27, and P15, were upregulated upon treatment with XPO1 inhibitors. The abrogation of MYC protein expression by XPO1 inhibition was accompanied by induction of apoptosis, as manifested by the cleavage of PARP and Caspase-3 (**Supplementary Fig S1**).

We concluded that XPO1 suppression abrogates the function of MYC oncogene and induces mass apoptosis in DHL tumor cells.

XPO1 has been reported to mediate nuclear export of several mRNAs encoding oncoproteins, including MYC, BCL2, and PIM1. Therefore, XPO1 inhibition may potentially reduce MYC protein expression through the nuclear retention of MYC mRNA. To test whether XPO1 inhibition affects nuclear export of MYC mRNA, we examined the distribution of MYC transcripts by cellular fractionation followed by quantitative realtime PCR. Compartmental separation of the nuclear versus cytoplasmic fractions was well achieved as shown by quantification of GAPDH and laminin B1 proteins, and Neat1 transcript in each compartment[14] (**Supplementary Fig S2C-D**). Interestingly, MYC nuclear to cytoplasmic ratio decreased by 50% upon XPO1 inhibition, while its mRNA abundance in the whole cell extract increased by two folds (**Fig 1F-G**). The nuclear to cytoplasmic ratio of BCL2 but BCL6 increased modestly at 16 hours of posttreatment. The level and nuclear/cytoplasmic ratio of beta-actin, used as an internal control, was not affected by XPO1 inhibition (**Supplementary Fig S2A-B**). This observation indicates that XPO1 inhibition may not result in a preferential nuclear retention of MYC mRNA. Therefore, the drastic downregulation of MYC protein expression upon XPO1 inhibition cannot be explained by changes in the levels and/or transport of MYC mRNA. We further ruled out the role of protein degradation as a contributing factor, as treatment with a proteasome inhibitor, carfilzomib, did not affect MYC shutdown by XPO1 inhibition (**Supplementary Fig S2E**). Taken together, these data strongly suggest that the effective downregulation of MYC protein by XPO1 inhibition may occur at the level of protein translation.

XPO1 inhibition synergizes with BCL2 inhibition in vitro in killing DHL tumor cells

The compelling downregulation of MYC protein expression by XPO1 inhibition suggests the possibility to target MYC and BCL2 driver oncogenes in DHL by combining XPO1 and BCL2 inhibitors. To test this notion, we first determined the IC50 values for two targeted agents, ABT199 (a BCL2 inhibitor), and KPT8602 (an XPO1 inhibitor) in a panel of DHL cell lines. Among the DHL cell lines examined, some were resistant to ABT199 treatment, while others were moderately resistant to KPT8602 (**Fig 2A-B**). However, to our surprise, none of the DHL cell lines were resistant to both (**Fig 2C, Supplementary Fig S3A-B**), lending experimental support to the combinatorial use of XPO1 and BCL2 inhibitors to eradicate DHL tumor cells. Further *in vitro* examination of the drug combination using additional DHL cell lines demonstrated a strong synergy in cell death induction (**Fig 3A**), while treatment with single agent showed moderate effects. The apoptotic event resulting from the combined treatment was accompanied by enhanced PARP and Caspase 3 cleavage compared to treatment with either single agents (**Fig 3B-D**). The apoptosis phenotype induced by synergistic drug combination was further confirmed by flow cytometric analysis following 7AAD/Annexin V staining (**Fig 3E**). MCL1 is known to play an important role in regulating the sensitivity of tumor cells to BCL2 inhibitors as MCL1 overexpression is associated with resistance to ABT199[15,16]. Next, we examined whether XPO1 inhibition enhances the sensitivity of tumors cells to ABT199 by downregulating MCL1. Indeed, MCL1 downregulation was observed in 8 out of 12 DHL cell lines treated with XPO1 inhibitor KPT8602 (**Supplementary Fig 1A**). This observation may partially explain the synergistic effect between XPO1 and BCL2 inhibitors in inducing apoptosis of DHL tumor cells.

Combined targeting of XPO1 and BCL2 blocks tumor progression and spread in vivo

Encouraged by our *in vitro* results, we initiated an *in vivo* study to test the combination therapy. Primary human DHL PDX tumor cells obtained from the Public Repository of Xenografts (PRoXe) were

transplanted in NOD/SCID/IL2R gamma (NSG) mice[17]. These PDX tumor cells were labeled with firefly luciferase for bioluminescence imaging (BLI). Importantly, as part of the therapeutic regimen, a low dose of KPT8602 (7.5mg/kg, half of dose used in other studies) was used[18-20]. Additionally, at the start of the treatment, the tumor loads were at least 5 folds higher than recommended[17,21]. Despite the low drug dose and higher than recommended tumor loads, we obtained a significant therapeutic response (**Fig 4**). The combined treatment completely blocked tumor progression and significantly extended survival, while single agents were only moderately effective (**Fig 4A-C**). Importantly, BLI signals from the skulls were reduced by 22 folds indicating the drug treatment effectively blocks tumor metastasis to brain (**Fig 4B, 4D**). Thus, combined targeting of XPO1 and BCL2 is highly effective for the treatment of human DHL *in vivo*.

Discussion:

XPO1 inhibition is a potential strategy to target MYC oncogene in DHL tumors as XPO1 inhibitors effectively downregulate MYC protein expression and reprogram the gene expression of MYC downstream targets. The mechanism by which XPO1 inhibition abolishes MYC protein expression remains unknown. We found that neither the expression nor the transport of MYC transcript can explain the downregulation of MYC expression by XPO1. Protein degradation as a contributing factor was also ruled out as proteasome inhibition did not rescue the downregulation of MYC by XPO1 inhibitors. These observations point to a translational regulation of MYC expression by XPO1. MYC mRNA has very long 5' and 3'UTRs, containing regulatory motifs (such as IRES and ARE) important for translational initiation. We speculate that XPO1 may control nuclear cytoplasmic partitioning of key translation initiation factors essential for MYC protein expression. A comprehensive proteomic and genetic approach is underway to identify the *bona fide* translational regulator(s) in DHL tumor cell fractions.

In DHL tumors carrying genomic translocations of MYC and BCL2, and overexpressing the two driver oncogenes, MYC and BCL2 may be targeted simultaneously using XPO1 and BCL2 inhibitors. Accordingly, we find that concomitant suppression of XPO1 and BCL2 is effective for treatment of human DHL tumors. XPO1 and BCL2 inhibitors synergize *in vitro* to induce apoptosis in DHL tumor cells, and most importantly, block tumor progression and dissemination *in vivo*. Although our study has focused on the treatment of DHLs with genomic translocation of MYC and BCL2, the concept of simultaneously targeting multi-driver oncogenes, may be extended to other DHLs or triple-hit lymphomas (THLs) with genomic abnormalities in BCL6 oncogene. The inclusion of BCL6 inhibitors,

such as 79-6[22], in our combined therapeutic regimen could be tested in preclinical settings for the treatment of DHL/THLs.

Compared to other types of B-cell lymphomas, DHL tumors are more likely to spread to a patient's central nervous system (CNS)[23]. Brain involvement in lymphoma patients generally confers a more abysmal prognosis with a median survival of 2-5 months[23]. Our *in vivo* study with human PDX reveals that combined targeting of XPO1 and BCL2 blocks brain metastasis of DHL tumors. Thus, it may be possible to prevent tumor spread using this type of combination targeted therapy. Our finding is in concordance with a recent case report, where the use of another XPO1 inhibitor, selinexor (KPT330), was shown to restrain the CNS relapse in one DLBCL patient[24]. The role of XPO1 in brain metastasis and the notion of using XPO1 inhibitors to suppress CNS involvement of lymphoma may warrant further preclinical examination.

Materials and Methods:

Cell lines and chemical reagents

DHL cell lines RC, SU-DHL-4 (DHL4), SU-DHL-6 (DHL6), SU-DHL-10 (DHL10) and Toledo were purchased from the ATCC. DB, SU-DHL-5 (DHL5), OCI-LY19, Will2, WSU, Val and U2932 were gifts from Dr. Lan Pham at the University of Texas MD Anderson Cancer Center. Cells were cultured in RPMI 1640, supplemented with 20% FBS (except for RC, DHL4, and Toledo, supplemented with 10% FBS), and 1% penicillin/streptomycin in a 5% CO_2 humidified incubator. ABT199 (Venetoclax) was obtained from Houston Methodist Pharmacy. KPT8602 and KPT330 were purchased from Selleckchem. Carfilzomib was purchased from Cayman Chemical.

IC50 determination

Cells were plated at 2×10^4 cells in 100 μl culture medium per well in 96-well plates. Drugs were added the next day in 4 replicates. Following five days of drug treatment (except for DHL10, which was treated for three days), cell viability was assessed using the CellTiter-Glo® 2.0 Cell Viability Assay (Promega) according to the manufacturer's instructions. Data analyses and calculation of median effective dose (IC50) were carried out using GraphPad Prism 8 (GraphPad).

Western blot analysis

DHL cells were lysed on ice in CelLytic™ MT Cell Lysis Reagent (Sigma) supplemented with protease and phosphatase inhibitors. Protein concentration was determined with Pierce™ BCA Protein Assay Kit (Thermo Fisher). Protein lysates were resolved on SDS-PAGE gels and transferred to PVDF membranes. The following antibodies were used for the Western blot analysis: MYC, cleaved PARP,

Caspase 3, BCL2, XPO1, Lamin B1, GAPDH (Cell Signaling), MYC (Abcam), β-Tubulin, β-Actin (Proteintech).

Cellular Fractionation and quantitative real-time PCR

Cytoplasmic and nuclear RNA were isolated and purified using the RNA Subcellular Isolation Kit (Active Motif) following the manufacturer's protocol. RNA binding and purification was performed using RNeasy Mini Kit (Qiagen). Total RNA was extracted with RNeasy Mini Kit (Qiagen). cDNA was synthesized using the Verso cDNA Synthesis Kit (Thermo Scientific) according to the manufacturer's instructions. qPCR analysis was performed on 7500 Real Time PCR System using SYBR Green (Applied Biosystems). Transcript levels were normalized to GAPDH, and relative gene expression was determined with ddCt method. Primer sequences are available upon request.

Flow cytometry

The PE Annexin-V Apoptosis Detection Kit (BD Biosciences) was used to detect apoptosis following manufacturer's instruction. Briefly, cells were suspended in 150 μl of binding buffer and mixed with 5 μl of FITC-conjugated Annexin-V and 7-AAD, followed by incubation at room temperature for 15 min in the dark. The stained cells were analyzed using a BD FACS Fortessa flow cytometer (BD Bioscience). All flow cytometric data were analyzed with Flowjo software (Tree Star).

In vivo therapeutic studies

Animal studies, procurement of patients' samples, and experimental protocols were approved by Institutional Animal Care and Use Committee (IACUC) at Houston Methodist Research Institute. All experimental methods were performed in accordance with the relevant national and institutional

guidelines and regulations. The DHL PDX sample (DFBL-69487-V3-mCLP) was obtained from the Public Repository of Xenografts (PRoXe), which collects PDX samples from patients with informed consent. Cells (10^6) were engrafted via tail vein injection into 6- to 8-week-old NSG mice. The animals were monitored once a week by whole-body imaging on an IVIS Lumina III platform. Treatment started upon the appearance of measurable tumors. Tumor volumes were assessed from the start of the treatment 2 times per week using IVIS imaging system. The following treatment schemes were used: daily oral gavage with ABT199 (50mg/kg) and KPT8602 (7.5mg/kg) for 5 successive days, followed by 2 days off for three weeks. ABT199 tablets were dissolved in water by sonication; KPT8602 was dissolved in 0.5% methylcellulose plus 1% Tween-80 as reported[18-20]. BLI data were analyzed with the Living Image software, version 4.2 (Caliper Life Sciences). Animal studies were carried out in accordance with IACUC approval at Houston Methodist Research Institute (AUP-1117-0053).

Statistical analysis

Two-tailed Student's t test was used to analyze the quantitative PCR data for mRNA expression. Cell death rates among different treatment groups were analyzed using ANOVA with a Tukey's test. Results were presented as mean ± standard deviation. Animal survival in different groups was compared by Kaplan-Meier analysis with Log-rank (Mantel-Cox) test.

Figure Legends:

Figure 1. XPO1 inhibitors suppress MYC protein expression in DHLs

(A) Changes in MYC protein level upon treatment with KPT8602 (1μM for 24 h). Asterisk (*) indicates that DHL5 is not considered a DHL.

(B) Time-dependent modulation of MYC protein in three DHL cell lines upon exposure to KPT330 or KPT8602 for 4, 8, 12 and 24 h. Cells were treated with a drug concentration of 0.5 μM for DHL4/Toledo and 1μM for DHL6.

(C) Dose-dependent reduction of MYC protein level upon XPO1 inhibition. Cells were treated with 0 to 0.8 μM of KPT330 or KPT8602 for 24 h. β-tubulin (Tub) serves as a loading control.

(D-E) Changes in transcript levels of MYC targets upon XPO1 inhibition. DHL6 was treated with KPT8602 at 1 μM for 8 and 24 hours.

(F) Total mRNA and (G) nuclear to cytoplasmic ratio of MYC and β-tubulin transcripts in DHL6 treated with 1μM of KPT8602 for 4, 16 or 24 h. All mRNA values were normalized to GAPDH.

Figure 2. Determination of IC50 values for ABT199 and KPT8602 in a panel of DHL cell lines

(A) IC50 values for ABT199. (B) IC50 values for KPT8602.

(C) IC50 values for ABT199 and KPT8602 in a panel of DHL cell lines.

Figure 3. KPT8602 synergizes with ABT199 *in vitro* in killing DHL tumor cells

(A) Cell viability upon treatment with KPT8602 (100nM) and ABT199 (40nM for DHL4/DHL6, and 20nM for Toledo) for 48 hours. (Two-tailed test. ** indicates $P < 0.0001$; * indicates $P < 0.001$.)

(B-D) DHL cells were treated with 80nM ABT199, 100nM KPT8602 or both for 8 or 24 hours. Western blot analysis of protein lysates for MYC, cleaved PARP, Caspase3 (Casp3), BCL2 and β-Tubulin (Tub).

(E) Flow cytometric analysis of apoptosis induction in DHL4 treated with 100nM ABT199 and 50nM KPT8602 for 48 hours. Cells were stained with Annexin-V and 7-AAD.

Figure 4. *In vivo* therapeutic effects of combining XPO1 and BCL2 inhibitors

(A) Growth of PDX tumors in NSG mice as imaged by BLI. Arrows indicate the three cycles of treatment.

(B) Representative BLI images of tumor-bearing mice following drug treatments.

(C) Kaplan-Meier survival analysis of tumor-bearing mice (Log-rank Mantel-Cox test. $P<0.0001$).

(D) BLI signals from the skull of the tumor bearing animals following drug treatment. The difference of BLI signals between control and single or combination treated mice was significant (Two-tailed t test. * Control vs ABT199, $P=0.02$; ** Control vs KPT8602, $P=0.01$; *** Control vs Combination, $P=0.0008$.).

Supplementary Figure Legends

Figure S1

(A) Western blot analysis of MYC, MCL1, and PARP/Caspase 3 cleavage. A panel of DHL cell lines was treated with 1 μM KPT8602 for 24 h. Part of the data were shown in **Fig 1**.

(B) Western blot analysis of MYC and PARP/Caspase 3 cleavage upon XPO1 inhibition in three DHL cell lines treated for 4, 8, 12, and 24 hours at a drug concentration of 0.5 μM for DHL4/Toledo and 1μM for DHL6.

(C) Western blot analysis of MYC and PARP/Caspase 3 cleavage upon treatment with different concentrations of XPO1 inhibitors for 24 hours in three DHL cell lines.

Figure S2

(A) Total mRNA and (B) nuclear to cytoplasmic ratio of BCL2, BCL6 and β-tubulin (loading control) in DHL6 treated with 1μM KPT8602 for 4, 16 or 24 h. All mRNA levels were normalized to GAPDH.

(C) Quantificaiton of nuclear and cytoplasmic Neat1 mRNA levels by realtime PCR.

(D) Analysis of nuclear and cytoplasmic GAPDH (cytoplasmic marker) and Lamin B1 (nuclear marker) by Western Blot.

(E) Representative DHL cells were treated with 1 μM KPT8602 and/or 10 nM Carfilzomib for 24 hours. β-actin was used as a loading control.

Figure S3

(A) IC50 values for ABT199 in a panel of DHL cell lines.

(B) IC50 values for KPT8602 in a panel of DHL cell lines.

(C) Cell viability in DHL cells treated with KPT8602 and ABT199 for 72 hours. The IC50 values for KPT8602 were calculated in the presence of different concentrations of co-administered ABT199.

(D) Cell morphology of DHL cells treated with KPT8602 (100nM) and ABT199 (40nM for DHL4/DHL6, and 20nM for Toledo) for 48 hours.

Reference

Petrich, A. M. *et al.* Impact of induction regimen and stem cell transplantation on outcomes in double-hit lymphoma: a multicenter retrospective analysis. *Blood* **124**, 2354-2361, doi:10.1182/blood-2014-05-578963 (2014).

Hu, S. *et al.* MYC/BCL2 protein coexpression contributes to the inferior survival of activated B-cell subtype of diffuse large B-cell lymphoma and demonstrates high-risk gene expression signatures: a report from The International DLBCL Rituximab-CHOP Consortium Program. *Blood* **121**, 4021-4031; quiz 4250, doi:10.1182/blood-2012-10-460063 (2013).

Vaux, D. L., Cory, S. & Adams, J. M. Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells. *Nature* **335**, 440-442, doi:10.1038/335440a0 (1988).

Arcinas, M., Heckman, C. A., Mehew, J. W. & Boxer, L. M. Molecular mechanisms of transcriptional control of bcl-2 and c-myc in follicular and transformed lymphoma. *Cancer Res* **61**, 5202-5206 (2001).

Souers, A. J. *et al.* ABT-199, a potent and selective BCL-2 inhibitor, achieves antitumor activity while sparing platelets. *Nat Med* **19**, 202-208, doi:10.1038/nm.3048 (2013).

Volpon, L. *et al.* A biochemical framework for eIF4E-dependent mRNA export and nuclear recycling of the export machinery. *RNA* **23**, 927-937, doi:10.1261/rna.060137.116 (2017).

Topisirovic, I. *et al.* Molecular dissection of the eukaryotic initiation factor 4E (eIF4E) export-competent RNP. *EMBO J* **28**, 1087-1098, doi:10.1038/emboj.2009.53 (2009).

Marullo, R. *et al.* Abstract LB-062: XPO1 is a rational target for double and triple-hit aggressive B-cell lymphomas. *Cancer research* **75**, LB-062-LB-062, doi:10.1158/1538-7445.am2015-lb-062 (2015).

Tai, Y. T. *et al.* CRM1 inhibition induces tumor cell cytotoxicity and impairs osteoclastogenesis in multiple myeloma: molecular mechanisms and therapeutic implications. *Leukemia* **28**, 155-165, doi:10.1038/leu.2013.115 (2014).

Ranganathan, P. *et al.* XPO1 Inhibition using Selinexor Synergizes with Chemotherapy in Acute Myeloid Leukemia by Targeting DNA Repair and Restoring Topoisomerase IIalpha to the Nucleus. *Clin Cancer Res* **22**, 6142-6152, doi:10.1158/1078-0432.CCR-15-2885 (2016).

Culjkovic-Kraljacic, B. *et al.* Combinatorial targeting of nuclear export and translation of RNA inhibits aggressive B-cell lymphomas. *Blood* **127**, 858-868, doi:10.1182/blood-2015-05-645069 (2016).

Zheng, Y. *et al.* KPT-330 inhibitor of XPO1-mediated nuclear export has anti-proliferative activity in hepatocellular carcinoma. *Cancer Chemother Pharmacol* **74**, 487-495, doi:10.1007/s00280-014-2495-8 (2014).

Zeller, K. I., Jegga, A. G., Aronow, B. J., O'Donnell, K. A. & Dang, C. V. An integrated database of genes responsive to the Myc oncogenic transcription factor: identification of direct genomic targets. *Genome Biol* **4**, R69, doi:10.1186/gb-2003-4-10-r69 (2003).

Lellahi, S. M. *et al.* The long noncoding RNA NEAT1 and nuclear paraspeckles are up-regulated by the transcription factor HSF1 in the heat shock response. *The Journal of biological chemistry* **293**, 18965-18976, doi:10.1074/jbc.RA118.004473 (2018).

Choudhary, G. S. *et al.* MCL-1 and BCL-xL-dependent resistance to the BCL-2 inhibitor ABT-199 can be overcome by preventing PI3K/AKT/mTOR activation in lymphoid malignancies. *Cell Death Dis* **6**, e1593, doi:10.1038/cddis.2014.525 (2015).

16 Phillips, D. C. *et al.* Loss in MCL-1 function sensitizes non-Hodgkin's lymphoma cell lines to the BCL-2-selective inhibitor venetoclax (ABT-199). *Blood Cancer J* **5**, e368, doi:10.1038/bcj.2015.88 (2015).

17 Townsend, E. C. *et al.* The Public Repository of Xenografts Enables Discovery and Randomized Phase II-like Trials in Mice. *Cancer cell* **29**, 574-586, doi:10.1016/j.ccell.2016.03.008 (2016).

18 Vercruysse, T. *et al.* The Second-Generation Exportin-1 Inhibitor KPT-8602 Demonstrates Potent Activity against Acute Lymphoblastic Leukemia. *Clinical cancer research : an official journal of the American Association for Cancer Research* **23**, 2528-2541, doi:10.1158/1078-0432.CCR-16-1580 (2017).

19 Etchin, J. *et al.* KPT-8602, a second-generation inhibitor of XPO1-mediated nuclear export, is well tolerated and highly active against AML blasts and leukemia-initiating cells. *Leukemia* **31**, 143-150, doi:10.1038/leu.2016.145 (2017).

20 Hing, Z. A. *et al.* Next-generation XPO1 inhibitor shows improved efficacy and in vivo tolerability in hematological malignancies. *Leukemia* **30**, 2364-2372, doi:10.1038/leu.2016.136 (2016).

21 Rava, M. *et al.* Therapeutic synergy between tigecycline and venetoclax in a preclinical model of MYC/BCL2 double-hit B cell lymphoma. *Sci Transl Med* **10**, doi:10.1126/scitranslmed.aan8723 (2018).

22 Cerchietti, L. C. *et al.* A small-molecule inhibitor of BCL6 kills DLBCL cells in vitro and in vivo. *Cancer cell* **17**, 400-411, doi:10.1016/j.ccr.2009.12.050 (2010).

23 Qualls, D. & Abramson, J. S. Advances in risk assessment and prophylaxis for central nervous system relapse in diffuse large B-cell lymphoma. *Haematologica* **104**, 25-34, doi:10.3324/haematol.2018.195834 (2019).

24 Bobillo, S. *et al.* Promising activity of selinexor in the treatment of a patient with refractory diffuse large B-cell lymphoma and central nervous system involvement. *Haematologica* **103**, e92-e93, doi:10.3324/haematol.2017.181636 (2018).

Figure 1

Figure 2

Figure 3

Figure 4