# Neuronal Responses in Posterior Parietal Cortex during Learning of Implied Serial Order

**Fabian Munoz**[1,5], **Greg Jensen**[2,5], **Benjamin C. Kennedy**[3], **Yelda Alkan**[1,5], **Herbert S. Terrace**[2,4], **and Vincent P. Ferrera**[1,4,5]

[1]**Dept. of Neuroscience, Columbia University Medical Center**
[2]**Dept. of Psychology, Columbia University**
[3]**Dept. of Neurosurgery, University of Pennsylvania**
[4]**Dept. of Psychiatry, Columbia University Medical Center**
[5]**Zuckerman Mind Brain Behavior Institute, Columbia University**

## ABSTRACT

Monkeys are able to learn the implied ordering of pairs of images drawn from an ordered set, without ever seeing all of the images simultaneously, and without explicit spatial or temporal cues. Learning is evidenced by their ability to make transitive inferences (TI), which are judgments of serial order based on the transitive relation that A>B and B>C implies A>C. Because posterior parietal cortex has been shown to integrate spatial and numerical information, we recorded the activity of parietal neurons in rhesus macaques while they learned 5- and 7-item TI lists. Symbolic distance (but not ordinal position) strongly predicted decision accuracy, and learning was consistently faster as symbolic distance increased. Neurons exhibited selectivity for stimulus identity during learning, but not before or after. Furthermore, across the population of neurons, there was significant modulation of firing correlated with the relative rank of the two stimuli presented on each trial. The observed behavior during learning is best explained by a virtual workspace model, not by associative or reward mechanisms. The neural data support a role for posterior parietal cortex in representing several variables that contribute to serial learning, particularly information about the ordinal ranks of both stimuli presented during a given trial. Thus, parietal cortex can be considered as part of the neural substrate for learning and representing abstract relationships in a virtual workspace.

Keywords:     Mental workspace, model-based learning, transitive inference, lateral intraparietal area

Transitive inference (TI) refers to judgments that if A>B and B>C, then one can infer that A>C by the transitive property of ordinal rank (Halford et al., 2010). Rank efficiently encodes list order or hierarchy, as only one value is needed for each item (which scales linearly with list length), rather than encoding a separate relationship for each combination of items (which scales exponentially). A wide variety of species can learn ordered lists (Vasconcelos, 2008; Jensen, 2017). Such hierarchical knowledge is considered important in a variety of natural contexts (Moses et al., 2006; Haun et al., 2010; Gazes et al., 2013).

Several lines of evidence suggest that abstract cognitive mechanisms give rise to TI abilities (Frank et al., 2006; Kumaran and Ludwig, 2013). The strongest of these is the symbolic distance effect (SDE), in which accuracy to each pair varies directly with the distance that separates them (Terrace et al., 2003; Terrace, 2012). Recent evidence suggests that purely associative mechanisms and reward signals cannot account for these effects (Jensen et al., 2013; Kumaran, 2013; Jensen et al., 2019). Abstract representations of ordinal rank may be linked to those of spatial position (Roberts and Phelps, 1994; Taffe et al., 2004), with both utilizing a "virtual workspace." This workspace may be used across modalities, such that spatial

and non-spatial information are integrated into a common representation (Gevers et al., 2003; Walsh, 2003; Hubbard et al., 2005).

Parietal cortex has long been associated with spatial relationships, and parietal lobe damage in humans gives rise to a well-known spatial hemineglect syndrome. In parietal stroke cases (e.g. Gerstmann Syndrome), multimodality spatial and number-related deficits can co-occur (Pia et al., 2009). Parietal dysfunction involving the intraparietal sulcus (IPS) is associated with irregularities in spatial and numerical representation (Wingard et al., 2002; Molko et al., 2003). Posterior parietal cortex (PPC) has been shown to integrate spatial, temporal, and reward information (Roitman et al., 2007). There is also evidence suggesting that the same areas of parietal lobe are responsible for both spatial representation and serial learning (Acuna et al., 2002; Hinton et al., 2010; Smets et al., 2013), including TI (Gould et al., 2009; Prado et al., 2013).

IPS is also implicated in various number representation and manipulation tasks (Nieder et al., 2002; Piazza et al., 2004, 2007; Prado et al., 2010; Roitman et al., 2012). Electrophysiological studies suggest that PPC encodes numerosity (Jacob and Nieder, 2009; Sawamura et al., 2010; Viswanathan and Nieder, 2013). Lateral intraparietal (LIP) neurons exhibit features of a magnitude accumulator (Dehaene and Changeux, 1993; Sawamura et al., 2002; Nieder, 2013) and are implicated in spatial cognition, including spatial attention (Bisley and Goldberg, 2003), judging relative spatial position (Chafee et al., 2007), and probabilistic reasoning (Yang and Shadlen, 2007). LIP also appears to integrate relevant spatial, temporal, numerical, and reward signals guiding behavior (Freedman and Assad, 2009), making this region a candidate for representation of the virtual workspace.

Electrophysiology in non-human primates offers an opportunity to study serial learning without confounds of semantic encoding and language use. Macaques can learn 7-item lists within a few hundred trials. That allows their behavior to be examined with concurrent single-unit electrophysiology. Although other studies implicate the parietal lobe in serial learning in non-human primates, they have been limited by spatial, temporal, and stimulus property confounds. The present study is the first to examine the electrophysiological basis of ordinal position and distance effects during the training of tasks in which the stimulus order is entirely implicit. Evidence that the firing rates of LIP neurons correlate with ordinal position of or symbolic distance between stimuli, would support the encoding of a mental number line in LIP. Our paradigm is uniquely poised to address whether TI is accomplished by using purely associative mechanisms, and, if not, whether a higher order representation involving the parietal cortex is likely.

## MATERIALS & METHODS

### Subjects

Subjects were two male rhesus macaques (H and L), 13-14 years old and weighing 9-10 kg at time of experiment. The research was conducted in accordance with U.S. Department of Health and Human Services (National Institutes of Health) guidelines for the care and use of laboratory animals, and was approved by the Institutional Animal Care and Use Committee at Columbia University and the New York State Psychiatric Institute. Monkeys were prepared for experiments by surgical implantation of a post for head restraint and a recording chamber to give access to cortex. All surgery was performed under general anesthesia (isoflurane 1-3%) and aseptic conditions. Subjects had prior experience performing transitive inference tasks (Jensen et al., 2015).

### Visual Stimuli & Eye Movements

The subjects were seated in an upright primate chair during the experiment and responded to visual stimuli by making saccadic eye movements. Stimuli were generated and controlled by a CRS VSG2/3F

video frame buffer. Stimuli were displayed on a CRT (subject H) or LCD (subject L) monitor with a resolution of 1280 x 1024 pixels at 60 Hz. Viewing distance was 60 cm.

Visual stimuli consisted of 140-by-130 pixel color photographs (7°by 8°visual angle) and small squares that served as fixation and eye movement targets. The fixation point was a red square (0.5°visual angle).

Eye position was recorded using a monocular scleral search coil (CNC Engineering, Seattle WA) and digitized at a rate of 1 kHz (Judge et al., 1980). Subjects made choices by making eye movements from the central fixation point to target stimuli. Eye velocity was computed offline by convolving eye position with a digital filter. The filter was constructed by taking the first derivative of a temporal Gaussian, $G(t)$, such that

$$\frac{dG}{dt} = -k \cdot t \cdot \exp\left(\frac{t^2}{\tau^2}\right)$$

Here, $\tau = 8$ msec and $k$ is a constant that sets the filter gain to 1.0. This filter does not introduce a time shift between the position input and velocity output, but adds temporal uncertainty to the velocity estimates. Horizontal and vertical eye velocities ($h'(t)$ and $v'(t)$, respectively) were combined to estimate radial eye speed $r'(t)$ using the Pythagorean identity:

$$r'(t) = \sqrt{\left(h'(t)^2 + v'(t)^2\right)}$$

## Behavioral Tasks

Subjects were trained to perform three visual-oculomotor tasks: delayed visually guided saccades, delayed memory guided saccades, and transitive inference.

### Delayed visual and memory guided saccades

The purpose of this task was to estimate the spatial selectivity (receptive and movement field) of each neuron. Each trial started with a small white spot (0.5°white square) presented in the center of the video display. Once the subject fixated on this spot, a peripheral cue appeared (1.0°white square). The subject was required to hold fixation for 0.75 to 1.25 seconds. Then the central fixation target was turned off, instructing the subject to make a saccade to the peripheral cue. The eccentricity and azimuth of the peripheral cue varied from trial to trial. Targets were presented at 8, 16, or 24 locations. This allowed the experimenter to determine the spatial locations to which each neuron responded most strongly. The stimulus presented at the preferred location is hereafter referred to as falling within the cell's receptive field. The other stimulus was presented diametrically opposite relative to the center of the screen, and is referred to as falling outside the receptive field. The delay between the peripheral cue onset and the saccade allowed experimenters to distinguish between visual and movement-related neuronal activity. The memory guided saccade task was identical to the visually guided saccade task, except that the peripheral cue was presented only briefly (0.5 sec) and then turned off. During the cue presentation and the subsequent memory delay interval (0.75 to 1.25 sec), the subject continued to fixate on the center fixation target. Then the central fixation target was turned off, instructing the subject to make a saccade to the remembered location of the peripheral cue. After the saccade, the peripheral cue was turned back on to provide feedback about saccade accuracy.

### Transitive inference

Prior to the beginning of each recording session, a set of 5 or 7 photographs never before seen by the subjects were selected from a database of over 2500 images. A different set of images was used for each

**Figure 1.** Transitive inference task procedure. **A.** Example of a list of 7 random images assigned ranks A thru G. **B.** All 21 image pairs for a 7-item list, and their respective joint ranks and symbolic distances. **C.** Trials structure and timing.

session. These stimuli were randomly assigned a unique rank indicated by the letters A-E or A-G, forming an ordered list (Fig. 1A). No explicit information about stimulus rank was presented to the subjects. That is, the letter assigned to each stimulus was never displayed, nor was there any information about serial order in the spatial or temporal presentation of the stimuli.

For 5- and 7-item lists, there are 10 and 21 possible pairs, respectively. In addition to being identified in terms of which stimuli they included, each pair could be described in terms of two metrics that encode information about both stimuli. The "symbolic distance" was calculated by taking the difference between the stimulus ranks (D'Amato and Colombo, 1990). For example, since the pair BC consists of items that are adjacent in the list (i.e. ranks 2 and 3, respectively), their symbolic distance is 1, while the pair AG (ranks 1 and 7) has a symbolic distance of 6. The "joint rank" is the sum of the ranks of the two stimuli presented on each trial, and is orthogonal to symbolic distance (Jensen et al., 2017). BC would have a joint rank of 5 (2+3), while AG would have a joint rank of 8 (1+7). A diagram mapping the symbolic distance and joint rank for all pairs is presented in Figure 1B.

On each trial, the subject first fixated a target in the center of the screen (Fig. 1C). Following a random delay (0.4 sec to 1.2 sec, positively skewed with a mean of 0.5 sec), the fixation point disappeared and two stimuli appeared equidistant from the center of the screen in opposite directions. To receive a reward, subjects had to select the stimulus with the lower rank by making a saccadic eye movement to the selected stimulus within 1.5 sec of stimulus onset, and fixating for a random interval (0.4 to 0.6 sec, uniformly distributed). When this interval elapsed, auditory feedback was delivered indicating that the response was either correct (high tone, 880 Hz) and would be rewarded, or was incorrect (low tone, 440 Hz). In order to receive the reward on correct trials, subjects had to maintain fixation for another interval (0.35 to 0.65 sec, uniformly distributed), after which the screen went dark and fluid rewards (juice or water drops) were delivered.

The first phase of each session presented single stimuli from the list. A "block" of responses during this phase consisted of presentations of each of the stimuli, both in and out of the cell's receptive field.

Thus, for a 7-item list, each block was 14 trials long, randomly ordered. The first phase lasted 5 blocks (i.e. either 50 or 70 trials in total). A saccade to these stimuli always yielded a reward. These presentations were used to determine whether the cells demonstrated selectivity toward any of the individual stimuli prior to training.

For the remainder of the session, pairs of stimuli were presented. With positional counterbalancing, this resulted in blocks that were 20 or 42 trials in length. In order to receive a reward, subjects had to choose the stimulus with the lower rank. Subjects were presented with all pairs of stimuli, one randomized block at a time, for up to 20 blocks (i.e. either 400 or 840 trials in total).

If good isolation from the recorded cell was maintained past the end of the all-pairs phase, and the subject remained motivated to work, a new list of photographs was drawn from the database, and the task was repeated (with another 50 to 70 trials of single stimuli and up to 20 blocks of pairs). During a single recording session, subjects were capable of learning 1 to 4 lists in this fashion.

## Neuronal Recording

Subjects had plastic recording chambers implanted over the right hemisphere at stereotaxic coordinates (subject H: anterior 5, medial 16; subject L: posterior 6, medial 14). Single-cell activity was recorded with glass-coated tungsten electrodes (Alpha Omega) with impedances between 0.5 and 2 MΩ measured at a frequency of 1 kHz. Raw signals were amplified, digitized, and high-pass filtered. We used FHC APM and Alpha Omega SnR recording systems. Action potential waveforms were identified by a time-amplitude window discriminator (FHC preamplifier) or threshold crossings (Alpha Omega). Action potentials were converted into digital pulses that were sampled by the computer with 0.02 ms temporal resolution. Waveforms were stored for offline spike sorting using WaveSorter (Phillips, 2012) or WaveClus (Quiroga et al., 2004). However, offline spike sorting was necessary only in exceptional cases because recording was mostly restricted to cells that could be clearly isolated online.

Anatomical reconstructions of recording sites were localized using combination of recording grid coordinates, microdrive depth readings, and stereotaxic position of the chamber, and were projected on the structural MRI using custom MATLAB scripts. Figure 2 plots a reconstruction of recording sites in structural MRI, matched over a 3D template atlas of the anatomical macaque rhesus brain for area classification (Reveley et al., 2017).

## Analysis

Analysis of behavioral and neuronal data consisted mainly of the following techniques: bootstrapped permutation testing (BPT), kernel density estimation (KDE) and Gaussian process regression (GPR). BPT was used to determine whether stimulus labels were informative with respect to overall neuronal activity. KDE was used for session-level averages of specific epochs within each trial, in which fairly large numbers of observations could be mustered to obtain an optimized nonparametric estimate. GPR modeling was used to make inferences about processes that unfolded trial by trial, in order to interpolate across gaps.

Each pair of stimuli was classified according to two metrics: symbolic distance (a measure of relative position, calculated by taking the difference between the stimulus ranks) and joint rank (a measure of absolute position, calculated by taking the sum of stimulus ranks). Because these metrics are strictly independent of one another, they form a two-dimensional continuum along which performance can be analyzed (Jensen et al., 2017).

### z-transform of neural activity

To construct a common metric across recording sessions with different activity levels, the firing rate on each trial ($FR_{\text{trial}}$) was converted to a $z$-score using the mean of the firing rate ($\mu_{FR}$) and standard deviation ($\sigma_{FR}$) over the entire session:

**Figure 2.** Reconstruction of recording sites and classification of neurons. **A,B.** Anatomical reconstructions of recording sites for NHP H ($n = 58$ sites) and L ($n = 53$ sites) were based on sterotaxic MR images, chamber implantation and angles, and matched to the macaque rhesus anatomical templates (Reveley et al., 2017). The templates were adapted to overlap over the MRI images plus the recording sites. The labels of the coronal slides are numbered to relative position to EBZ (Ear Bar Zero). **C,D.** Neuronal responses averaged over all trials of the TI task for NHP H ($n = 64$) and L ($n = 53$) optimally sorted by correlation similarity. **E.** Six families of response types defined by hierarchical clustering of mean firing rate vs. time, across all task conditions. The neurons ($n = 117$) were optimally ordered using correlation similarity.

$$z(FR) = \frac{FR_{\text{trial}} - \mu_{FR}}{\sigma_{FR}}$$

### *Bootstrap testing*

Bootstrapping was used to estimate distributions of sample means. $N$ samples were randomly drawn with replacement from the original set of $N$ observations. The mean of the resampled observations was calculated. This procedure was repeated typically 10,000 times to generate the bootstrap distribution.

### Random permutation testing

When examining the relationship between neural activity and explanatory variables such as ordinal position, it is useful to ask whether the observed relationship could arise by chance given the observed neuronal activity statistics. This was done by randomly permuting ("shuffling") the explanatory variable labels among the trials. Each label occurred as many times in the shuffled data as in the original data. This procedure estimates the likelihood that apparent "tuning" for a particular explanatory variable could arise by chance. Random permutation testing could be combined with bootstrapping to obtain a distribution of apparent tuning functions.

### Kernel density estimation

Eye movement characteristics and average session rates of neural activity both used KD estimates. For eye movements (position and velocity), we used a Gaussian kernel with Silverman's optimal bandwidth (Silverman, 1986). For spike trains, we used the method described by Shimazaki and Shinomoto (2010).

### Gaussian process regression

To evaluate how estimated performance changed as a function of learning, we used Gaussian process regression (Rasmussen and Williams, 2006). GPR is a highly flexible non-linear estimation technique that is well-suited to time series analysis. Although not widely used in neurophysiology, GPR is a well-established procedure (Durbin, 1985) that has seen extensive application in other domains. It has been called a categorical analysis with an infinite number of categories, arrayed along a continuum (McElreath, 2016). One continuum of interest is time: Response accuracy for each pair changes over time. Orthogonal to time are the continua of symbolic distance and joint rank, which are also expected to influence response accuracy. GP regression is performed by estimating the extent to which every observation covaries with every other, given some prior metric for comparing the distance between observations along the continua of interest. Each observation influences the estimate for other 'nearby' observations (e.g. that occur at similar times or have similar symbolic distance) more than observations that are distant (Lucas et al., 2015). Although such an analysis is not possible using classical methods (because of irreducible uncertainty about how distance and similarity interact to give rise to the data), Bayesian methods make GPR feasible by imposing a strong prior belief that pairs at similar times and with similar symbolic distance and joint rank should display similar patterns of neural activity (Gelman et al., 2014). GPR depends on relatively few assumptions, instead allowing the data to determine the form taken by the time series estimate. The chief constraint is that Gaussian processes are presumed to be smooth (i.e. differentiable without discontinuity). Beyond this constraint, one can imagine the model estimate as the posterior distribution of the relative density of all possible smooth curves, conditioned on the data and the informative prior. Although a full Gaussian process model can be computationally prohibitive to fit (requiring runtime $O(n^3)$ for $n$ observations), we will take advantage of the "expectation propagation" approximation (Tolvanen et al., 2014) implemented in the GPstuff toolbox (Vanhatalo et al., 2013) to accelerate estimation.

## RESULTS

Two subjects (H and L) completed 141 behavioral sessions (H: 74, L: 67). The average number of trials per session was 711 (H) and 890 (L). In each session the subject learned a novel 5-item or 7-item TI list; one session = one TI list. For H, 5-item lists were used in 22 sessions, and 7-item lists for the remaining 52 sessions. For L, all 67 sessions used 7-item lists. A total of 117 neurons were recorded (H: 64, L: 53) during these sessions. Some neurons were recorded with more than one TI list (2 lists N = 38; 3 lists N = 5). Eighteen sessions contained data from more than one neuron, separated using the spike

**Figure 3.** Response accuracy (proportion correct) was modeled as a Gaussian Process with a logit link, using trial number, symbolic distance, and joint rank as predictive dimensions **A.** Probability of choosing each item during the first 500 trials. The 80% and 99% credible intervals for the mean are plotted as the dark and light shaded areas, respectively. **B.** Mean accuracy over the first 500 trials (black line), as well as accuracy sorted by symbolic distance (colored lines). **C.** Estimated accuracy for each pair at trial 250 sorted by joint rank and symbolic distance. Thick whiskers correspond to the 80% credible interval, and thin whiskers (as well as shading) correspond to the 99% credible interval. **D.** Estimated accuracy at trial 500.

sorting methods described above. The result was a total of 164 recordings (H: 84, L: 80), where each recording contains data from one neuron with a novel list.

## Behavioral Performance

Performance accuracy and response time were analyzed to determine the influence of independent variables that are related to serial order: ordinal position (aka rank), symbolic distance, and joint rank, as well as variables related to reinforcement. A total of 141 sessions were analyzed (H: 74, L: 67). For some analyses, sessions with five-item lists were excluded so that the same set of independent variables could be used for all sessions. Excluding 5-item lists resulted in 119 sessions (H: 52, L: 67). Each session started with a novel set of pictures that were arranged into a list by assigning a rank order to each picture. Figure 3 illustrates the development of both the probability of choosing a given stimulus and the probability of a correct response over the first 500 choice trials, as estimated using GP regression. Overall learning curves (Fig. 3B) showed a monotonic increase in response accuracy (Fig. 3B black line) from a baseline of 0.5 (chance) up to around 0.75. The shaded regions represent the 80% and 99% credible intervals around the estimate of the mean.

Performance consistently improved as a function of symbolic distance. Each session began with a novel set of stimuli with an initially unknown ranking, and the effect of symbolic distance on performance emerged over the course of the session, stabilizing after 250 choice trials (Fig. 3A & 3B). Accuracy was lowest for adjacent pairs, somewhat higher for distance 2 pairs, and so on (Fig. 3C). Across sessions, the correlation (Pearson's $r$) of overall performance (session average) with symbolic distance was significant for both subjects (H: $r = 0.62$, $p < 0.0001$, $N = 400$; L: $r = 0.61$, $p < 0.0001$, $N = 402$) with moderate effect sizes (H: $r^2 = 0.38$; L: $r^2 = 0.37$).

The correlation of overall performance with joint rank was significant for only one subject and effect sizes were very small for both animals (H: $r = 0.04$, $p = 0.32$, $N = 726$; L: $r = 0.15$, $p < 0.0001$, $N = 737$). Performance tended to be better for pairs that included the first and last list items, a "terminal item effect" (Fig. 3C & 3D).

Subjects were allowed up to 1.5 sec after stimulus onset to respond by making a saccadic eye movement to the target or distractor. Despite the long time-window, choice reaction times (RT, aka saccade latency)

**Figure 4.** Eye movement metrics. **A.** Latency of choice saccade relative to stimulus onset sorted by symbolic distance and joint rank. **B.** Peak eye velocity. **C.** Saccade amplitude. Thick and thin whiskers are 80% and 99% credible intervals. Note that y-axes do not go to zero as effects are very small in absolute terms.

were approximately 0.3 sec and remained in a narrow range throughout each session. Within this range, RT varied slightly with symbolic distance and joint rank. Figure 4A shows KD estimates indicating that smaller symbolic distances resulted in slower reaction times, as did smaller joint rank.

Response vigor is an index of performance that can be derived from eye movements and may be related to reward value (Reppert et al. 2015). Vigor was quantified by calculating peak saccade velocity (KD estimate), as shown in Figure 4B. Peak velocity for all pairs that included the first item (stimulus A) was very consistent (~329 degrees per second), and then increased as a function of the rank of the correct response. However, the absolute variation in peak saccade velocity was only ~2% (i.e. a range of 329 to 335 degrees per second). Figure 4C shows KD estimates of saccade amplitude, which was primarily determined by target eccentricity, and showed no significant variation with distance or joint rank.

It is possible that subjects used a model-free reinforcement learning (RL) strategy to perform the task. A hallmark of model-free RL is that choices that are rewarded are more likely to be repeated than those that are not rewarded. We investigated this by first looking at the pattern of spatial choices; if the subject was rewarded for choosing the stimulus in a particular location, would they be more or less likely to choose the same location on the next trial regardless of whether it was the correct response? There was evidence in favor of a spatial win-stay, lose-shift strategy; rewarded responses were repeated 56% of the time, significantly different from the expectation of 50% ($t$-test $p < 0.0001$, $N = 141$, Cohen $d = 0.68$). Unrewarded trials were followed by the same response location only 48% of the time ($t$-test $p < 0.002$, $N = 141$, Cohen's $d = 0.27$). This result generally agrees with a previous study of reward-based decision-making (Kubanek and Snyder, 2017), which found that monkeys were more likely to change their response after receiving a relatively small reward. Reaction time (choice saccade latency) did not vary depending on the outcome of the previous trial (mean RT difference after reward - after no reward = -0.9 ms, SD = 6.3 ms, paired $t$-test $p = 0.1$, $N = 141$).

It should be noted that a spatial choice bias is deleterious to overall performance, as the target location was randomized and provided no information about the correct choice. The presence of a spatial bias does not rule out a choice bias that is positively related to learning of serial order. This possibility was tested by asking whether reward history biased the choice probability for each of the 5 or 7 pictorial

stimuli conditioned on whether or not the previous choice of that stimulus had been rewarded. This analysis was performed for non-terminal list items (B thru D, or B thru F), for which the expected choice probability is 0.5, regardless of performance accuracy. Some conditions were excluded if the choice probability could not be calculated (i.e. the denominator was 0, which happened in 11 of 661 cases). The average choice probability following rewarded choices was 0.51 (SD = 0.17) compared to 0.51 (SD = 0.18) following unrewarded choices. The difference was not significant ($t$-test paired by session and item, $p = 0.99$, $N = 650$). Similarly, the average latency of choice saccades varied by less than 1ms between previously rewarded and unrewarded conditions (paired $t$-test $p = 0.34$, $N = 661$). The results were the same whether all choice trials in each session were included, or only the first 250 trials when learning of serial order was most rapid. These results provide no support for the idea that subjects used a model-free reinforcement learning strategy to learn or perform the task. If anything, rewarded choices were slightly less likely to be repeated than unrewarded choices.

The results for choice probability are supported by analysis of conditional error rates. If an item was presented and the subject chose correctly, he was more likely to make an error the next time that item was presented (conditional error rate = 0.18, SD = 0.08) compared to cases where the first response had been incorrect (conditional error rate = 0.14, SD = 0.14; $t$-test paired by session and item, $p < 0.001$, $N = 846$, Cohen's $d = 0.33$). The same pattern was found regardless of whether the item was presented as a target or a distractor. Hence, there was significant evidence that subjects monitored errors and adapted subsequent behavior.

**Electrophysiology**

A hierarchical clustering algorithm was used to classify neurons according to similarity of firing rate dynamics. Firing rates were first averaged across trials using KD estimation (Shimazaki and Shinomoto, 2010), to estimate firing rate across the session. These averages were then $z$-standardized, and sorted into clusters on the basis of the correlation of their activity over the course of a trial. Categories were identified by building a dendrogram of correlations using agglomerative hierarchical cluster analysis (Hastie and Tibshirani, 2010), with $r = 0.7$ as the cutoff for cluster membership. This resulted in 6 clusters, depicted in Figure 2E.

Cluster V, comprising visual neurons, contained the most recordings. An example recording session with one visual neuron is shown in Figure 5. The cell had a transient, short latency (50 msec) response to the appearance of a visual stimulus in its receptive field (Fig 5A). The neuronal response was well before the behavioral response latency (150 msec for single targets, 200 msec for choice targets). The spatial selectivity (Fig. 5B, black) was assessed on single target trials, showing a 2:1 difference between responses to stimuli inside vs. outside the receptive field. On choice trials, selectivity was assessed by sorting according to saccade direction (Fig. 5B, blue). This cell did not differentiate between saccades toward or away from the RF, and can thus be said to have visual spatial, but not motor or intentional selectivity. The visual spatial selectivity can be seen in Figure 5C (black) where firing rate has been converted to $z$-score and plotted as a function of ordinal position (rank) for the stimulus inside the receptive field. During single target presentations, variations in the response to different images can only be due to the visual properties of the stimuli, as the NHP has yet to learn the rank of each stimulus. Responses during choice trials (Fig. 5C, blue, red) are also shown as a function of the rank of the RF stimulus. A $z$-score of zero means that the response was equal to the average response across all conditions. Averaging of responses when two stimuli are present, inside and outside the RF, is discussed below. Responses on choice trials are also plotted as a function of symbolic distance (difference in rank of the two stimuli, Fig. 5D) and joint rank (Fig. 5E). This cell shows effects that were characteristic of the population: strong spatial selectivity for single targets, response averaging for two stimuli, little selectivity for ordinal position, but

**Figure 5.** Example of a recording session for one cell. **A.** Spike raster aligned on stimulus onset (vertical blue line). Each red dot (for single target trials) or black dot (for choice trials) represents an action potential. Each row is one trial. Magenta dots are saccade latencies. **B.** Firing rate vs. target position (-1 is left hemifield, 1 is right hemifield). Small dots are individual trials, large dots are averages. Black/gray corresponds to single target trials, blue corresponds to choice trials. **C.** *z*-transformed firing rate vs. ordinal position. **D.** *z*-transformed firing rate vs. symbolic distance. **E.** *z*-transformed firing rate vs. joint rank.

some modulation by symbolic distance and joint rank. These features of the data are elaborated below.

At the start of each recording session with the TI task there were five blocks (50 - 70 trials) in which the 5 or 7 pictorial stimuli were presented singly. During each trial, one stimulus appeared either inside the receptive field of the neuron ($T_{in}$) or at a location diametrically opposite to the receptive field ($T_{out}$). Neuronal responses on these "single target" trials were compared to later "decision" trials with a target ($T_{in}$) and distractor ($D_{in}$) to determine the effect of the stimulus outside the receptive field, given prior reports of surround suppression in LIP (Falkner et al., 2010; Arcizet et al., 2018). Responses (defined as spike counts within a time window of 50 to 150 msec after stimulus onset) to a single target in the receptive field ($T_{in}$) were compared to responses with a target in the receptive field and distractor at the opposite location ($T_{in} + D_{out}$). Across the population of 164 recordings, the presence of the distractor reduced the response to the target by an average of 14% or 4.2 sp/sec ($p < 0.0001$, $N = 164$, Cohen's $d = 0.36$). The response to two stimuli presented at the same time ($R_{both}$) could be expressed as a weighted average of the response to single stimuli inside ($R_{in}$) or outside ($R_{out}$) the receptive field:

$$R_{both} = k \cdot R_{in} + (1 - k) \cdot R_{out}$$

Here, k denotes the weight of the stimulus in the receptive field, which was estimated at 0.69 by linear regression. The $R_{out}$ stimulus reduced responses to the $R_{in}$ stimulus even though, at the population level, the $R_{out}$ response was not significantly different than background activity (paired $t$-test $p = 0.78$, $N = 164$). These observations establish that individual LIP neurons are influenced by stimuli at both spatial locations, consistent with Falkner et al. (2010), and providing a basis for examining the neural representation of quantities defined jointly by both stimuli, such as symbolic distance and joint rank.

### Neural activity and performance

In the TI task, all correctly completed trials were rewarded, thus reward probability was equal to percent correct. Reward-related modulation of neural activity in macaque parietal cortex was described by Platt and Glimcher (1999). It has also been reported that neuronal activity in parietal cortex is correlated with performance accuracy (Zhang et al., 2014). To test this hypothesis, we compared the average difference in activity (spike counts) between correct and incorrect trials for each recording. Three trial epochs were examined: background (200 msec prior to stimulus onset), visual (50-150 msec after stimulus onset) and presaccadic (100 msec prior to onset of choice saccade). Each of 164 recordings was subjected to a Wilcoxon (Matlab ranksum) test with a criterion of $\alpha = 0.05$. Effect sizes were measured using $\eta^2$. Few recordings showed significant effects of performance in any trial epoch and the effect sizes were tiny (background epoch: $N = 13$ (8%) significant recordings, mean $\eta^2 = 0.003$; visual $N = 15$ (9%), $\eta^2 = 0.003$; presaccadic $N = 29$ (18%), $\eta^2 = 0.04$), suggesting negligible association between neural activity and performance outcome or reward probability in the current data set.

To summarize the preceding results, we constructed peri-stimulus time histograms (PSTH's) by counting spikes in 100 msec bins spaced at 10 msec intervals. Bins were labeled with the leading edge of the time interval, hence the bin corresponding to t = 0 msec included spike times from -99 to 0 msec. The counts in each bin were averaged over recordings to estimate the mean population response. Figure 6A shows strong spatial selectivity in the population response to single targets inside (black) vs. outside (gray) the receptive field. However, spatial selectivity was not apparent when choice trials were sorted by saccade direction (Fig. 6A, dark blue = toward RF, light blue = away from RF). Similar results were obtained when activity was aligned with saccade onset (Fig. 6B).

The population results were supported by a time-resolved GLM whose explanatory variables were target location (for single target trials), saccade direction (for choice trials), and outcome (choice trials). Each explanatory variable had 2 levels and therefore a linear model was appropriate. Figures 6C and

**Figure 6.** Effects of target position, saccade direction, and outcome. **A.** Population firing rate aligned on stimulus onset, averaged over 164 recordings. Single target activity is depicted in black (inside RF) or gray (outside RF), while choice trials are depicted in blue (dark depicting choices toward the RF, and light depicting choices away from the RF). The dashed line represents stimulus onset. **B.** Firing rate aligned to saccade onset, with choice trials depicted in red. **C.** GLM beta weights in-RF/out-RF effect averaged over neural population, with activity aligned on stimulus onset. **D.** Same description as C, but aligned on saccade onset.

D confirm that stimulus position was a strong modulator of neuronal responses on single target trials.

However, saccade direction and trial outcome were relatively weak. Post-hoc ANOVAs (Kruskal-Wallis) were run for each recording and explanatory variable separately with alpha = 0.05. For target location, 71/164 (43%) recordings reached significance with an average effect size (generalized $\eta^2$) of 0.34. For saccade direction, 63/164 (38%) recordings reached significance (avg. effect size = 0.08). For outcome, 33/164 (20%) were significant (avg. effect size = 0.02).

### Neural effects of reward history

Behaviorally, we found that rewarded spatial choices were likely to be repeated (although rewarded list item choices were not). This raised the possibility that reward history might affect neuronal responses. Indeed, in a task with varying reward magnitudes, Kubanek and Snyder (2017) found that LIP neurons fired more vigorously following trials that ended with relatively small rewards. In the current data (164 recording sessions), there was little evidence that reward outcome affected neural activity on the next trial. For choice saccades toward the receptive field, firing rate during the period between stimulus onset and choice saccade was slightly elevated on the trial following a rewarded trial compared to an unrewarded trial (reward - no reward, mean firing rate difference = 0.07 sp/sec, $p = 0.75$, $N = 164$). For saccades away from the receptive field, firing rate was slightly lower (reward - no reward, mean firing rate difference = -0.19 sp/sec, $p = 0.39$, $N = 164$). These effects were negligible and suggest that, in the context of the current task, reward history was not a significant modulator of neural activity in LIP.

### Encoding of stimulus identity

Sereno and Maunsell (1998) reported visual feature selectivity in monkey posterior parietal cortex using stimuli that controlled for luminance cues. In the current study, the stimuli were not controlled for luminance, chromaticity, contrast or spatial frequency content, but were randomly selected colorful images with content that was easily recognized and discriminated. Early on in the experiments, stimuli were manipulated to have identical spatial frequency content within each of the 3 color channels, but it was difficult to recognize distinct objects within these images. Given that the subjects were asked to learn a new set of 5 or 7 images for every session, it was deemed important that they be able to easily discriminate the stimuli even if this meant sacrificing control over low-level visual features. Nevertheless, it is possible to ask if neuronal responses to the different stimuli varied reliably, and the lack of low-level control increases the expectation that such variability should be found. For this analysis, 7-item recording sessions ($n = 142$) were used. Spikes were counted within a time window 50-150 msec after stimulus onset on each trial, converted to spikes per second, and then $z$-transformed.

Trials in which only a single stimulus appeared were sorted according to the identity of that stimulus. Figures 7A and 7B show the values of z(FR) sorted by stimulus and averaged over all (142) recording sessions. The stimulus that evoked the strongest response in each session was assigned to position 1, and the other stimuli were assigned to position 2-7, maintaining their rank order. Some degree of apparent stimulus preference can arise by chance, given finite sampling. To control for this, the preceding analysis was repeated with the stimulus identities randomly shuffled among trials. There was no significant difference between the shuffled and unshuffled responses (inside RF, bootstrapped $p = 0.58$; outside RF, bootstrapped $p = 0.70$; post-hoc tests for each item were n.s.). Response differences across stimuli thus appear to be accounted for by random variability, and there is no evidence that neuronal responses reliably encoded stimulus identity during the initial phase of the task when the stimuli were presented individually.

After the first 5 blocks of each recording session, images were presented in pairs, one image inside of and the other opposite to the receptive field. As shown above (Fig. 3), TI learning was most rapid during the first 250 trials of paired presentations. During this time, there was a single best stimulus for each neuron that evoked a stronger response when presented in the receptive field compared to the "best" stimulus when stimulus identity was shuffled across trials (Fig 7C, bootstrapped difference $p < 0.001$,

**Figure 7.** Encoding of stimulus identity using recordings with 7-items lists ($n$=142). **A.** Trials with a single stimulus in RF. Mean $z$-transformed firing rate during visual epoch for original stimulus labels (blue, not shuffled) and shuffled labels (black). Light blue and grey points are estimates of the means from 10,000 bootstrap iterations. **B.** $z$-transformed firing rate during visual epoch of first 250 choice trials with best stimulus inside RF. Light blue dots are responses with original labels, grey dots are responses with shuffled labels. Dark blue and black dots are estimates of the mean from 10,000 bootstrap iterations. **C.** Same as B, but for choice trials after trial number 250.

$N = 142$). This effect was strongest during learning trials (i.e., the first 250 paired presentations, Fig 7C), but was not present during later trials, when performance reached a plateau (Fig. 7D; bootstrapped $p = 0.8$; $N = 142$). Hence, during TI learning, the visual response of LIP neurons reliably but transiently

**Figure 8.** Symbolic distance selectivity. **A.** Recordings ($n$=142) sorted by preferred symbolic distance based on z-scored firing rate. **B.** Mean normalized firing rate vs. symbolic distance. Gray lines are 95% confidence intervals for all trials based on 10,000 bootstrap iterations.

encoded stimulus identity.

### Encoding of symbolic distance and joint rank

Symbolic distance is the difference in list position (rank) of the two stimuli presented on each trial. It is a joint property that is independent of the spatial configuration of the stimuli. For each recording, $z$-scores of the average firing rate in the interval starting 50 msec after stimulus onset and ending with saccade onset were sorted by symbolic distance. Figure 8A shows the firing for each recording, sorted by the symbolic distance that was associated with the strongest response. Roughly half the recordings had the greatest activity for one of the two largest symbolic distances. Figure 8B shows normalized firing rate averaged over recordings and plotted as a function of symbolic distance. The correlation was significant when all choice trials were included ($r = 0.11$, $p = 0.02$, $N = 142$).

Joint rank is the sum of the list positions of the two items presented on each trial, and is complementary to symbolic distance. There was no systematic increase or decrease in firing as a function of joint rank. To compare the effects of symbolic distance and joint rank, the variance accounted for (VAC) was computed. For each recording, firing rate was smoothed by applying Gaussian Process Regression (GPR) using two models; one with trials, time, and symbolic distance (SD), and another with trials, time, and Joint Rank (JR). The estimation was done within 4 epochs: baseline (250 ms before stimulus onset), visual (between 5 ms to 150 ms after visual presentation), presaccadic (100 msec before saccade initiation) and postsaccadic (500 ms after saccade execution) period. Each period was divided in intervals of 10 msec to create a spike counts over time as a prior to the GPR. The posterior estimate of firing rate (mean and standard deviation) was continuous in time, trial, SD and JR. The estimated firing rate and uncertainty were then averaged

**Figure 9.** Variance accounted for by symbolic distance and joint rank, sorted by neuron class and by trial epoch. Box-and-whisker diagrams depict mean VAC for each cluster, with boxes corresponding to 80% credible intervals and whiskers corresponding to 95% credible intervals.

within the previously specified time windows. To estimate the amount of variance associated to SD or JR across the session for visual , presaccadic or postsaccadic epochs, we subtracted the baseline, and then evaluated the variance account for (VAC) using partial eta squared ($\eta_p^2$), with F-values and degrees of freedom from the Welch test for populations with unequal variance. Each $\eta_p^2$ was sorted by cell class (Fig. 2E) and in order of increasing magnitude (Fig. 9 A-F). The density of each $\eta_p^2$ was estimated by bootstrap. The median with 80% (box) and 95% (whiskers) credible intervals are shown in Figure 9. VAC values for JR are generally larger than for SD. VAC was similar across clusters, suggesting that SD and JR are represented across all cell classes.

## DISCUSSION

The transitive inference (TI) paradigm tests subjects' ability to learn the implied serial order of a set of pictorial stimuli without any explicit spatial or temporal cues. This is accomplished by presenting pairs of images drawn from a rank-ordered list of images and rewarding subjects for choosing the item that has the lower rank. Much work in humans and other animals has supported the idea that TI learning relies on a virtual workspace (reviewed by Jensen, 2017), but few studies have examined its physiological underpinnings in brain regions implicated in spatial cognition (Brunamonti et al., 2016). Here, we tested the virtual workspace hypothesis by recording neuronal activity in parietal area LIP while monkeys learned novel TI lists consisting of sets of 5 or 7 images assigned ranks that were learned by trial-and-error.

Each recording session started with a novel set of images. Subjects began at chance levels of performance, and were able to learn each list within about 250 trials. Prior research argues against

the idea that monkeys memorize the response to each pair (Treichler et al., 2007), or rely solely on the experienced reward value of each stimulus (Jensen et al., 2019). Rather, they appear to acquire abstract knowledge of the list order, and of the rule that dictates choosing the lower ranked item in each pair. In the current study, behavior was more consistent with representation and rule-based learning than with model-free reinforcement learning. Neuronal activity in parietal cortex was strongly visual and spatially selective, but also showed significant influences of cognitive variables related to list order such as item rank, symbolic distance, and joint rank. These results support the idea that parietal cortex is involved in representing abstract serial order.

We tested the contribution of model-free learning mechanisms by performing a sequential trial analysis to determine if received rewards biased subjects' behavior on subsequent trials. Reward history was found to introduce a spatial bias, but did not bias responses in favor of choosing a previously rewarded list item. It is therefore unlikely that model-free RL can account for learning. These results are in accord with a sizable literature showing that experienced reward value does not fully account for transitive inference or serial learning in multiple primate species (Terrace, 2012; Jensen et al., 2019). There was no evidence that reward history modulated neuronal responses as found by others (Kubanek and Snyder, 2017). Variations in neuronal response were only weakly correlated with trial outcome (Zhang et al., 2014).

In agreement with previous work in LIP (Gnadt and Andersen, 1988), neuronal responses to single stimuli were spatially selective with a strong contralateral visual field bias. However, when two stimuli were presented (choice trials), one inside and the other outside the receptive field, neural responses were best described as a weighted average of the responses to the individual stimuli. Averaging occurred even though the stimulus outside the receptive field, when presented alone, evoked only small increases or decreases in firing rate relative to background activity. The weighted average was biased roughly 70:30 in favor of the stimulus inside the receptive field. These results are in general accord with previous work showing that LIP neurons can be influenced by stimuli outside the receptive field (Falkner et al., 2010; Arcizet et al., 2018).

During choice trials, neurons exhibited little spatial selectivity, i.e. responses did not vary depending on whether the choice saccade was directed toward or away from the receptive field. This result is surprising given the extensive literature on target selection and decision-making in LIP (see Gold and Shadlen, 2007; Huk et al., 2017), but may be accounted for by two features of the current task. First, the decision required that subjects attend to and compare two stimuli in different locations. In many prior studies, the perceptual decision was based on a single stimulus and did not require dividing attention. Second, the decision was relatively fast - accurate choices were made within roughly 300 msec. Previous studies have reported response times of up to  900 msec (e.g. Roitman and Shadlen, 2002). Perceptual decisions and target selection are separable processes (Bennur and Gold, 2011). It is possible that, in the current task, the parietal target selection signal simply did not have sufficient time to evolve.

There was no evidence that item identity based on shape, color, or other visual features were encoded when single stimuli were presented during the initial trials of each session, in contrast with previous work showing shape selectivity in LIP (Sereno and Maunsell, 1998). This was despite substantial item-wise variation in responses, possibly driven by low-level visual properties of the stimuli that were not systematically controlled. However, this variation did not survive a permutation test, which is a fairly stringent test of reliable stimulus encoding. The lack of selectivity during single stimulus presentations argues against the possibility that any signals found during choice trials were driven by visual features of the stimuli.

During the first 250 trials of learning, there was a signal that reliably distinguished among list items. Because the neurons showed no inherent visual selectivity, it is likely that this signal was related to the learned list positions. The effect of list position disappeared after the initial learning, as accuracy reached

a plateau. Thus, the representation of item list position in parietal cortex appears to be transient.

Two variables that are defined jointly by the two stimuli presented on choice trials are symbolic distance (difference of ranks of individual stimuli) and joint rank (sum of the two ranks). There was evidence that both variables are represented in parietal cortex, with moderate effect sizes as measured by variance accounted for (effect size approximately 0.2). These effects are substantially smaller than the effects of spatial position (effect size 0.36), but much larger than the effects of trial outcome (effect size at most 0.04).

The representation of variables that arise from the comparison of spatially remote stimuli is contrary to conventional thinking about visual or visual-movement neurons with spatially localized receptive or movement fields. However, such responses are not unprecedented in the dorsal stream. Zaksas and Pasternak (2005) found effects of remote stimuli in visual MT when monkeys were required to compare motion stimuli in the contra- and ipsilateral visual fields. Moreover, models of spatial remapping around eye movements strongly suggest that individual LIP neurons receive information from across the entire visual field (Quaia et al., 1998).

Overall, our results support an emerging view of parietal cortex in which: 1) Spatial and non-spatial information is represented by the same population of neurons. 2) Non-spatial information can encode abstract quantities related to implied serial order, which obeys the law of transitivity. Finally, 3) information is integrated across visual space to support decisions based on the comparison of spatially remote stimuli. All of these properties add to growing evidence that parietal cortex likely plays a substantial role in the construction of a virtual workspace for both spatial and abstract cognitive tasks.

### Acknowledgments
We are grateful to Drew Altschul and Jessica Joiner for assistance during the early part of this project.

### Funding
This study was supported by NIH-MH081153 and NIH-MH111703 (VPF and HST).

### Author Contributions
The study was designed by GJ, VPF, and HST. Data were collected by FM, YA, and BCK. Data were analyzed by GJ, VPF, and FM. The paper was written by VPF, GJ, BCK, and FM, with input from HST and YA. FM and GJ made equal contributions.

## REFERENCES

Acuna, B. D., Eliassen, J. C., Donoghue, J. P., and Sanes, J. N. (2002). Frontal and parietal lobe activation during transitive inference in humans. *Cerebral Cortex*, 12:1312–1321.

Arcizet, F., Mirpour, K., Foster, D. J., and Bisley, J. W. (2018). Activity in LIP, but not V4, matches performance when attention is spread. *Cerebral Cortex*, 28:4195–4209.

Bennur, S. and Gold, J. I. (2011). Distinct representations of a perceptual decision and the associated oculomotor plan in the monkey lateral intraparietal area. *Journal of Neuroscience*, 31:913–921.

Bisley, J. W. and Goldberg, M. E. (2003). Neuronal activity in the lateral intraparietal area and spatial attention. *Science*, 299:81–86.

Brunamonti, E., Mione, V., Bello, F. D., Genovesio, A., and Ferraina, S. (2016). Neuronal modulation in the prefrontal cortex in a transitive inference task: Evidence of neuronal correlates of mental schema mangement. *Journal of Neuroscience*, 36:1223–1236.

Chafee, M. V., Averbeck, B. B., and Crowe, D. A. (2007). Representing spatial relationships in posterior parietal cortex: single neurons code object-referenced position. *Cerebral Cortex*, 17:2914–2932.

D'Amato, M. R. and Colombo, M. (1990). The symbolic distance effect in monkeys (*cebus apella*). *Animal Learning & Behavior*, 18:133–140.

Dehaene, S. and Changeux, J. P. (1993). Development of elementary numerical abilities: a neuronal model. *Journal of Cognitive Neuroscience*, 5:390–407.

Durbin, J. (1985). The first-passage density of a continuous gaussian process to a general boundary. *Journal of Applied Probability*, 22:99–122.

Falkner, A. L., Krishna, B. S., and Goldberg, M. E. (2010). Surround suppression sharpens the priority map in the lateral intraparietal area. *Journal of Neuroscience*, 30:12787–12797.

Frank, M. J., O'Reilly, R. C., and Curran, T. (2006). When memory fails, intuition reigns: midazolam enhances implicit inference in humans. *Psychological Science*, 17:700–707.

Freedman, D. J. and Assad, J. A. (2009). Distinct encoding of spatial and nonspatial visual information in parietal cortex. *Journal of Neuroscience*, 29:5671–5680.

Gazes, R. P., Brown, E. K., Basile, B. M., and Hampton, R. R. (2013). Automated cognitive testing of monkeys in social groups yields results comparable to individual laboratory-based testing. *Animal Cognition*, 16:445–458.

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., and Rubin, D. B. (2014). *Bayesian daya analysis, third edition*. CRC Press, Boca Raton, FL.

Gevers, W., Reynvoet, B., and Fias, W. (2003). The mental representation of ordinal sequences is spatially organized. *Cognition*, 87:B87–B95.

Gnadt, J. W. and Andersen, R. A. (1988). Memory related motor planning activity in posterior parietal cortex of macaque. *Experimental Brain Research*, 70:216–220.

Gold, J. I. and Shadlen, M. N. (2007). The neural basis of decision making. *Annual Review of Neuroscience*, 30:535–574.

Gould, R. L., Brown, R. G., Own, A. M., ffytche, D. H., and Howard, R. J. (2009). fMRI BOLD response to increasing task difficulty during successful paired associates learning. *NeuroImage*, 20:1006–1019.

Halford, G. S., Wilson, W. H., and Phillips, S. (2010). Relational knowledge: the foundation of higher cognition. *TRENDS in Cognitive Sciences*, 14:497–505.

Hastie, T. and Tibshirani, R. (2010). *The Elements of Statitical Learning*, chapter Hierarchical clustering, pages 520–528. Springer, New York, NY.

Haun, D. B., Jordan, F. M., Vallortigara, G., and Clayton, N. S. (2010). Origins of spatial, temporal and numerical cognition: insights from comparative psychology. *TRENDS in Cognitive Sciences*, 14:552–560.

Hinton, E. C., Dymond, S., von Hecker, U., and Evans, C. J. (2010). Neural correlates of relational reasoning and the symbolic distance effect: involvement of parietal cortex. *Neuroscience*, 168:138–148.

Hubbard, E. M., Piazza, M., Pinel, P., and Dehaene, S. (2005). Interactions between number and space in parietal cortex. *Nature Reviews Neuroscience*, 6:435–448.

Huk, A. C., Katz, L. N., and Yates, J. L. (2017). The role of the lateral intraparietal area in (the study of) decision making. *Annual Review of Neuroscience*, 40:349–372.

Jacob, S. N. and Nieder, A. (2009). Tuning to non-symbolic proportions in the human frontoparietal cortex. *European Journal of Neuroscience*, 30:1432–1442.

Jensen, G. (2017). Serial learning. In Call, J., Burghardt, G. M., Pepperberg, I. M., Snowdon, C. T., and Zentall, T., editors, *APA Handbook of Comparative Psychology*, pages 385–409. American Psychological Association, Washington, DC.

Jensen, G., Alkan, Y., Ferrera, V. P., and Terrace, H. S. (2019). Reward associations do not explain transitive inference performance in monkeys. *Science Advances*, In Press.

Jensen, G., Alkan, Y., Munoz, F., Ferrera, V. P., and Terrace, H. S. (2017). Transitive inference in humans

(*homo sapiens*) and rhesus macaques (*macaca mulatta*) after massed training of the last two list items. *Journal of Comparative Psychology*, 131:231–245.

Jensen, G., Altschul, D., Danly, E., and Terrace, H. S. (2013). Transfer of a serial representation between two distinct tasks by rhesus macaques. *PLOS ONE*, 8:e70285.

Jensen, G., Munoz, F., Alkan, Y., Ferrera, V. P., and Terrace, H. S. (2015). Implicit value updating explains transitive inference performance: The betasort model. *PLOS Computational Biology*, 11:e1004523.

Judge, S. J., Richmond, B. J., and Chu, F. C. (1980). Implantation of magnetic search coils for measurement of eye position: An improved method. *Vision Research*, 20:535–538.

Kubanek, J. and Snyder, L. H. (2017). Reward size informs repeat-switch decisions and strongly modulates the activity of neurons in parietal cortex. *Cerebral Cortex*, 27:447–459.

Kumaran, D. (2013). Schema-driven facilitation of new hierarchy learning in the transitive inference paradigm. *Learning & Memory*, 20:388–394.

Kumaran, D. and Ludwig, H. (2013). Transitivity performance, relational hierarchy knowledge and awareness: results of an instructional framing manipulation. *Hippocampus*, 23:1259–1268.

Lucas, C. G., Griffiths, T. L., Williams, J. J., and Kalish, M. L. (2015). A rational model of function learning. *Psychonomic Bulletin & Review*, 22:1193–1215.

McElreath, R. (2016). *Statistical Rethinking*. CRC Press, Boca Raton, FL.

Molko, N., Cachia, A., Riviere, D., Manginn, J., Bruandet, M., Bihan, D. L., Cohen, L., and Dehaene, S. (2003). Functional and structural alterations of the intraparietal sulcus in a developmental dyscalculia of genetic origin. *Neuron*, 40:847–858.

Moses, S. N., Villate, C., and Ryan, J. D. (2006). An investigation of learning strategy supporting transitive inference performance in humans compared to other species. *Neuropsychologia*, 44:1370–1387.

Nieder, A. (2013). Coding of abstract quantity by 'number neurons' of the primate brain. *Journal of Comparative Physiology A, Neuroethology, Sensory, Neural, and Behavioral Physiology*, 199:1–16.

Nieder, A., Freedman, D. J., and Miller, E. K. (2002). Representation of the quantity of visual items in the primate prefrontal cortex. *Science*, 297:1708–1711.

Phillips, M. H. (2012). Wavesorter: a new free and open-source tool for the offline analysis and classification of neural waveforms. In *2012 Neuroscience Meeting Planner*, page Program No. 508, New Orleans, LA. Society for Neuroscience.

Pia, L., Corazzini, L. L., Folegatti, A., Gindri, P., and Cauda, F. (2009). Mental number line disruption in a right-neglect patient after a left-hemisphere stroke. *Brain & Cognition*, 69:81–88.

Piazza, M., Izard, V., Pinel, P., Bihan, D. L., and Dehaene, S. (2004). Tuning curves for approximate numerosity in the human intraparietal sulcus. *Neuron*, 44:547–555.

Piazza, M., Pinel, P., Bihan, D. L., and Dehaene, S. (2007). A magnitude code common to numerosities and number symbols in human intraparietal cortex. *Neuron*, 53:293–305.

Platt, M. L. and Glimcher, P. W. (1999). Neural correlates of decision variables in parietal cortex. *Nature*, 400:233–238.

Prado, J., Mutreja, R., and Booth, J. R. (2013). Fractionating the neural substrates of transitive reasoning: task-dependent contributions of spatial and verbal representations. *Cerebral Cortex*, 23:499–507.

Prado, J., Noveck, I. A., and Henst, J. B. V. D. (2010). Overlapping and distinct neural representations of numbers and verbal transitive series. *Cerebral Cortex*, 20:720–729.

Quaia, C., Optican, L. M., and Goldberg, M. E. (1998). The maintenance of spatial accuracy by the perisaccadic remapping of visual receptive fields. *Neural Networks*, 11:1229–1240.

Quiroga, R. Q., Nadasdy, Z., and Ben-Shaul, U. (2004). Unsupervised spike detection and sorting with wavelets and superparamagnetic clustering. *Neural Computation*, 16:1661–1687.

Rasmussen, C. E. and Williams, C. K. I. (2006). *Gaussian Processes for Machine Learning*. MIT Press,

Cambridge, MA.

Reveley, C., Gruslys, S., Ye, F. Q., Glen, D., Samaha, J., Russ, B. E., Saad, Z., Seth, A. K., Leopold, D. A., and Saleem, K. S. (2017). Three-dimensional digital template atlas of the macaque brain. *Cerebral Cortex*, 27:4463–4477.

Roberts, W. A. and Phelps, M. T. (1994). Transitive inference in rats: a test of the spatial coding hypothesis. *Psychological Science*, 5:368–374.

Roitman, J. D., Brannon, E. M., and Platt, M. L. (2007). Monotonic coding of numerosity in macaque lateral intraparietal area. *PLOS Biology*, 5:e208.

Roitman, J. D., Brannon, E. M., and Platt, M. L. (2012). Representation of numerosity in posterior parietal cortex. *Frontiers in Integrative Neuroscience*, 6:Article 25.

Roitman, J. D. and Shadlen, M. N. (2002). Response of neurons in the lateral intraparietal area during a combined visual discrimination reaction time task. *Journal of Neuroscience*, 22:9475–9489.

Sawamura, H., Shima, K., and Tanji, J. (2002). Numerical representation for action in the parietal cortex of the monkey. *Nature*, 415:918–922.

Sawamura, H., Shima, K., and Tanji, J. (2010). Deficits in action selection based on numerical information after inactivation of the posterior parietal cortex in monkeys. *Journal of Neurophysiology*, 104:902–910.

Sereno, A. B. and Maunsell, J. H. (1998). Shape selectivity in primate lateral intraparietal cortex. *Nature*, 395:500–503.

Shimazaki, H. and Shinomoto, S. (2010). Kernel bandwidth optimization in spike rate estimation. *Journal of Computational Neuroscience*, 29:171–182.

Silverman, B. W. (1986). *Density estimation for statistics and data analysis*. Chapman & Hall, London, UK.

Smets, K., Gebuis, T., and Reynvoet, B. (2013). Comparing the neural distance effect derived from the non-symbolic comparison and the same-different task. *Frontiers in Human Neuroscience*, 7:Article 28.

Taffe, M. A., Weed, M. R., Gutierrez, T., David, S. A., and Gold, L. H. (2004). Modeling a task that is sensitive to dementia of the alzheimer's type: individual differences in acquisition of a visuo-spatial paired-associate learning task in rhesus monkeys. *Behavioural Brain Research*, 149:123–133.

Terrace, H. (2012). The comparative psychology of ordinal behavior. In Zentall, T. and Wasserman, E. A., editors, *Oxford Handbook of Comparative Cognition*, pages 615–651. Oxford University Press, Oxford, UK.

Terrace, H. S., Son, L. K., and Brannon, E. M. (2003). Serial expertise of rhesus macaques. *Psychological Science*, 14:66–73.

Tolvanen, V., Jylänki, P., and Vehtari, A. (2014). Expectation propagation for nonstationary heteroscedastic gaussian process regression. *2014 IEEE International Workshop on Machine Learning for Signal Processing*, pages 1–6.

Treichler, F. R., Raghanti, M. A., and Tilburg, D. N. V. (2007). Serial list linking by macaque monkeys (*Macaca mulatta*): List property limitation. *Journal of Comparative Psychology*, 121:250–259.

Vanhatalo, J., Riihimäki, J., Hartikainen, J., Jylänki, P., Tolvanen, V., and Vehtari, A. (2013). Gpstuff: Bayesian modeling with gaussian processes. *Journal of Machine Learning Research*, 14:1175–1179.

Vasconcelos, M. (2008). Transitive inference in non-human animals: An empirical and theoretical analysis. *Behavioural Processes*, 78:313–334.

Viswanathan, P. and Nieder, A. (2013). Neuronal correlates of a visual "sense of number" in primate parietal and prefrontal cortices. *Proceedings of the National Academy of Sciences of the United States of America*, 110:11187–11192.

Walsh, V. (2003). A theory of magnitude: common cortical metrics of time, space and quantity. *TRENDS in Cognitive Sciences*, 7:483–488.

Wingard, E. M., Barrett, A. M., Crucian, G. P., Doty, L., and Heilman, K. M. (2002). The gerstmann syndrome in alzheimer's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, 72:403–405.

Yang, T. and Shadlen, M. N. (2007). Probabilistic reasoning by neurons. *Nature*, 447:1075–1080.

Zaksas, D. and Pasternak, T. (2005). Area mt neurons respond to visual motion distant from their receptive fields. *Journal of Neurophysiology*, 94:4156–4167.

Zhang, M., Wang, X., and Goldberg, M. E. (2014). A spatially nonselective baseline signal in parietal cortex reflects the probability of a monkey's success on the current trial. *Proceedings of the National Academy of Sciences of the United States of America*, 111:8967–8972.