Bad and good news for ash trees in Europe: alien pest *Agrilus planipennis* has spread to the Ukraine and the south of

European Russia, but does not kill *Fraxinus excelsior* in the forests

Marina J. Orlova-Bienkowskaja

A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, 33 Leninskiy Prospect, Moscow,

Russia 119071

e-mail: marinaorlben@yandex.ru

Alexander N. Drogvalenko

V.N. Karazin Kharkiv National University, 4 Svobody Square, Kharkov, Ukraine 61022

e-mail: triplaxxx@ukr.net

Ilya A. Zabaluev

A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, 33 Leninskiy Prospect, Moscow,

Russia 119071

e-mail: fatsiccor66@mail.ru

Alexey S. Sazhnev

Papanin Institute for Biology of Inland Waters, Russian Academy of Sciences, Borok, Nekouz District, Yaroslavl

Region, Russia 152742

e-mail: sazh@list.ru

Elena Yu. Peregudova

Saint Petersburg State Forest Technical University, Institutskiy per., 5, Saint Petersburg, Russia 194021

e-mail: dinamo-l@mail.ru

Sergey G. Mazurov

Middle School of General education, Leski, Krasnoe District, Lipetsk Region, Russia 399675

e-mail: mazusergej@yandex.ru

Evgenij V. Komarov

All-Russian Scientific Research Institute of Irrigated Agriculture,

Timiryazev str., 9, Volgograd Russia 400002

e-mail: evkomarov@rambler.ru

Andrzej O. Bieńkowski

A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, 33 Leninskiy Prospect, Moscow,

Russia 119071

e-mail: bienkowski@yandex.ru

Acknowledgements

The study was supported by Russian Science Foundation, Project No 16-14-10031. We would like to thank entomologists V.V. Martynov and T.V. Nikulina (Donetsk, Ukraine), L.V. Egorov (Cheboksary, Russia), A.N. Volodchenko (Balashov, Russia), V.V. Anikin and M.V. Lavrentiev (Saratov, Russia) for the information about surveys of ash trees in different regions of European Russia and Ukraine and to Ya.N. Kovalenko for the valuable information on official phytoquarantine documents.

Abstract Emerald Ash Borer (EAB) *Agrilus planipennis* (Coleoptera: Buprestidae) native to Asia is a devastating pest of ash trees (*Fraxinus* spp.). It was first detected in Europe in Moscow in 2003. It is spreading over the country and causing serious damage to plantations of *F. pennsylvanica* introduced from North America. Our surveys of ash trees in 2017–2019 in 25 localities has shown that the range in ER has the area of 600 000 km^2, i.e. exceeds the area of Spain. We have first detected *A. planipennis* in the very south of Russia (Volgograd) and in the Ukraine (Markovka district of Lugansk Region). Therefore, now *A. planipennis* is detected in 14 regions of ER: Bryansk, Kaluga, Lipetsk, Moscow, Orel, Ryazan, Smolensk, Tambov, Tula, Tver, Vladimir, Volgograd, Voronezh and Yaroslavl as well as in Lugansk Region of the Ukraine. The only native ash species *F. excelsior* is susceptible to *A. planipennis*, but all cases of its infestation refer to the trees in urban, roadside and shelterbelt plantations planted near *F. pennsylvanica*, which is highly susceptible to the pest. Our survey of 500 *F. excelsior* trees in the large Tulskie Zaseki forest situated in the center of the pest range has not revealed any signs of infestation. There is no doubt that *A. planipennis* will appear in other European countries soon and severely damage plantations of *F. pennsylvanica* there. But the hope appeared that the pest could be less dangerous for *F. excelsior* in European forests.

Keywords Emerald Ash Borer; EAB; *Fraxinus pennsylvanica*; *Fraxinus excelsior;* invasive pest; ash

Introduction

Emerald ash borer (EAB) *Agrilus planipennis* (Coleoptera: Buprestidae) is a devastating alien pest of ash trees in European Russia and North America (Baranchikov et al. 2008; Haack et al. 2015). The native range of this wood-boring beetle occupies a restricted territory in East Asia (Orlova-Bienkowskaja and Volkovitsh 2017). In 2003 *A. planipennis* was first recorded in European Russia, namely in Moscow (Shankhiza 2007), and the severe outbreak and quick spread of the pest began (Baranchikov et al. 2008). By 2013 the pest was recorded in 9 regions of European Russia: from Yaroslavl in the north to Voronezh in the south (Straw et al 2013; Orlova-Bienkowskaja 2014a). The probabilistic model of spread has shown that in the next few years the range of *A. planipennis* can significantly expand, and the pest can appear in the neighboring countries (Orlova-Bienkowskaja and Bieńkowski 2018). The first aim of our study was to determine the current range of *A. planipennis*.

Fraxinus pennsylvanica introduced from North America is widely planted in European Russia as an ornamental and landscaping tree. It is highly susceptible to *A. planipennis* (Baranchikov et al. 2014). The overwhelming majority of detected infestations of ash trees in European Russia refer to this ash species (Baranchikov et al. 2008; Straw et al 2013; Orlova-Bienkowskaja 2014a). The only native ash species *F. excelsior* is also susceptible to the pest (Baranchikov et al. 2014), but studies on the feeding preferences of adult *A. planipennis* on different species of ash suggest that *F. excelsior* is not as susceptible as *F. pennsylvanica* (Pureswaran and Poland 2009). *Fraxinus excelsior* is rare in Moscow Region and other regions of Central Russia. So the information about infestation of *F. excelsior* by *A. planipennis* was scarce and referred only to urban plantations and artificial

shelterbelts, where *F. excelsior* was planted together with *F. pennsylvanica* (Straw et al. 2013; Baranchikov 2018).
There were no information about the impact of A. planipennis on *F. excelsior* in natural forests and, therefore, it was
a great deal of uncertainty surrounding the likely future impact of *A. planipennis* on ash in European forests (Straw
et al. 2013). So the second aim of our study was to get the information about current condition of *F. excelsior* in
natural forest stands in the regions occupied by *A. planipennis*.

Materials and Methods

In 2017, 2018 and 2019 we made surveys of ash trees in 25 localities of European Russia, Belarus and Ukraine (see
Tables 1, 2). In the most of these localities we examined artificial plantations in the cities, along motorways and
railroads an in the field shelter belts. The presence of the pest was detected by D-shaped exit holes on the stem,
larval galleries under the bark and the presence of larvae under the bark and adults in the canopy (Figs 1–4). Besides
own data we accumulated and mapped published records of *A. planipennis* and personal communications of
colleagues from several regions.
To determine the impact of *A. planipennis* on European ash *F. excelsior* we made a survey of more than
500 ash trees in the large forest Tulskie Zaseki situated in Tula Region in the center of the recent *A. planipennis*
range.
97

99 **Fig. 1** Exit hole of *Agrilus planipennis* on *Fraxinus pennsylvanica* in Volgograd (photo by E.V. Komarov).
100

101
102   **Fig. 2** Larval galleries of *Agrilus planipennis* on *Fraxinus pennsylvanica* in Volgograd (photo by E.V. Komarov).

104
Fig. 3 Adults of *Agrilus planipennis* in the pupal cells on *Fraxinus pennsylvanica* in Volgograd (photo by E.V.
Komarov).

Fig. 4 Adult of *Agrilus planipennis* on the leaf of *Fraxinus pennsylvanica* in Volgograd (photo by E.V. Komarov).

Abbreviations used in the tables:

AD – Observations by Alexander N. Drogvalenko

AS – Observations by A.S. Sazhnev

EK – Observations by Evgenij V. Komarov

EP – Observations by E. Yu. Peregudova

IZ – Observations by Ilya A. Zabaluev

MOB and AB – Observations by Marina J. Orlova-Bienkowskaja and Andrzej O. Bieńkowski.

SM – Observations by Sergey G. Mazurov

Table 1. Localities, where *Agrilus planipennis* has been detected in European Russia and the Ukraine. RU – Russia,

UA – Ukraine

Locality	Damaged trees	Kind of plantations	Year of the first detection	Lat	Lon	Sources of information
Moscow (*A. planipennis* occurs all over the city)	*F. pennsylvanica,* few *F. excelsior*	Urban	2003	55.81	37.64	(Shankhiza 2007; Izhevskii and Mozolevskaya 2010; Baranchikov et al., 2008; Straw et al., 2013; Orlova-Bienkowskaja 2014;

							Volkovitsh and Mozolevskaya 2014)
Moscow Reg.: Manikhino		*F. pennsylvanica*	Roadside	2006	55.88	36.97	(Shankhiza 2007)
Moscow Reg.: Serpukhov		*F. pennsylvanica*	Urban	2009	54.92	37.41	(Baranchikov et al. 2009)
Moscow Reg.: Mozhaisk		*F. pennsylvanica*	Urban	2009	55.51	36.01	
Moscow Reg.: Mytiszhi		*F. pennsylvanica*	Urban	2009	55.91	37.74	
Moscow Reg.: Pushkino		*F. pennsylvanica*	Urban	2009	56.01	37.84	(Baranchikov and Kurteev 2012)
Moscow Reg.: Korolev		*F. pennsylvanica*	Urban	2011	55.92	37.85	Volkovitsh and Mozolevskaya 2014
Moscow Reg.: Zelenograd		*F. pennsylvanica, few F. excelsior*	Urban	2011	56.00	37.18	(Orlova-Bienkowskaja 2014)
Smolensk Reg.: localities along the road Moscow-Vyazma		*F. pennsylvanica*	Roadside	2012	55.30	34.50	(Baranchikov and Kurteev 2012)
Moscow Reg.: Sergiev Posad		*F. pennsylvanica*	Urban	2012	56.31	38.13	
Ryazan Reg:Vysokoe		Not indicated	Not indicated	2012	54.52	39.41	(Baranchikov 2013)
Tula Reg.: Shchekino		*F. excelsior*	Roadside	2013	53.99	37.63	(Straw et al. 2013)
Tver Reg.: Zubstov		*F. pennsylvanica*	Urban	2013	56.18	34.58	
Tver Reg.: Novozavidovskiy		*F. pennsylvanica*	Roadside	2013	56.55	36.43	
Tver Reg.: Emmaus		*F. pennsylvanica*	Roadside	2013	56.78	36.09	
Moscow Reg.: Uzunovo		*F. pennsylvanica*	Along the railway	2013	54.55	38.62	(Orlova-Bienkowskaja 2014)
Moscow Reg.: Klin		*F. pennsylvanica*	Urban	2013	56.33	36.73	

Location	Species	Habitat	Year			Reference
Moscow Reg.: Staraya Kupavna	*F. pennsylvanica*	Urban	2013	55.81	38.18	
Voronezh	*F. pennsylvanica, F. excelsior*	Urban	2013	51.68	39.18	
Tula	*F. pennsylvanica, F. excelsior*	Urban	2013	54.20	37.62	
Kaluga	*F. pennsylvanica*	Urban	2013	54.52	36.26	
Tver Reg.: Konakovo	*F. pennsylvanica*	Urban	2013	56.71	36.77	
Orel	*F. pennsylvanica, F. excelsior*	Urban	2013	52.97	36.07	
Yaroslavl	*F. pennsylvanica*	Urban	2013	57.63	39.87	
Tambov Reg.: Michurinsk	*F. pennsylvanica*	Urban	2013	52.90	40.50	
Vladimir Reg.: Petushki	Not indicated	Not indicated	2013	55.93	39.46	(Baranchikov 2013)
Orel Region: Livny	Not indicated	Urban	2013	52.43	37.60	(Rosselkhoznadzor 2014)
Moscow Reg.: Solnechnogorsk	*F. pennsylvanica*	Roadside	2014	56.19	36.98	Own data (MOB and AB)
Moscow Reg.: Planernaya	*F. pennsylvanica*	Along the railway	2014	55.92	37.38	
Moscow Reg.: Povarovka	*F. pennsylvanica*	Along the railway	2014	56.06	37.08	
Smolensk Reg.: Semirechje	Not indicated	Roadside	2014	54.89	32.07	(Zvyagintsev et al. 2015)
Smolensk Reg.: Andreykovo	Not indicated	Roadside	2014	55.24	34.26	
Tver: Khiminstituta Settlement	*F. pennsylvanica*	Roadside	2015	56.80	36.03	(Peregudova 2016)
Moscow Reg.: Stupino	*F. pennsylvanica*	Urban	2017	54.89	38.10	Own data (MOB and AB)
Moscow Reg.: Kolomna	*F. pennsylvanica*	Urban	2017	55.08	38.79	
Moscow Reg.: Dedinovo	*F. pennsylvanica*	Urban	2017	55.06	39.12	
Voronezh Reg.: Talovaya	Not indicated	Not indicated	2018	51.11	40.72	(Baranchikov et al. 2018)
Lipetsk Reg.: Elets	Not indicated	Not indicated	2018	52.62	38.50	Baranchikov Y.N, oral communication

Country and locality	Examined trees	Kind of plantations	Year of survey	Lat	Lon	Sources of information
Smolensk	Not indicated	Along the railway	2018	54.80	32.03	in PREPSYS conference (Baranchikov and Seraya 2018)
Tver: Doroshiha railway station	*F. pennsylvanica*	Along the railway	2018	56.87	35.84	(Peregudova 2019)
Voronezh Reg.: Olkhovatskoe Forestry	Not indicated	Not indicated	2018	50.12	39.26	(Rosselkhoznadzor 2019)
Volgograd Reg.: Sarpinsky Island	*F. pennsylvanica*	Floodplain forest	2018	48.60	44.49	Own data (EK)
Volgograd Reg.: Volga-Akhtuba floodplain	*F. pennsylvanica*	Floodplain forest	2019	48.69	44.61	
Volgograd Reg.: Sarepta Peninsula	*F. pennsylvanica*	Floodplain forest	2019	48.53	44.51	
Volgograd City: Mira Street	*F. pennsylvanica*	Urban	2019	48.71	44.52	
Bryansk: Majskij Park	Not indicated	Urban	2019	53.31	34.31	(Rosselkhoznadzor 2019)
Moscow Reg.: Shakhovskaya	*F. pennsylvanica,* few *F. excelsior*	Urban	2019	56.03	35.51	Own data (MOB and AB)
Lipetsk Reg.: Leski	*F. pennsylvanica*	Along the railway	2019	52.88	39.00	Own data (SM)
Ukraine: Markovka District, near Starokozhij Forest	*F. pennsylvanica*	Field shelter belt	2019	49.61	39.56	Own data (AD)

122 Table 2. Localities where *A. planipennis* was not detected during the surveys in 2017, 2018 and 2019. BY - Belarus,

123 RU - Russia, UA - Ukraine

Country and locality	Examined trees	Kind of plantations	Year of survey	Lat	Lon	Sources of information
BY: Borisov	*F. pennsylvanica, F. excelsior*	Urban	2017	54.22	28.51	Own data (MOB and AB)
RU: Cheboksary	*F. pennsylvanica*	Urban	2017	56.14	47.25	L.V. Egorov, pers. comm.
RU: Saransk	*F. pennsylvanica*	Urban	2017	54.19	45.18	A.B. Ruchin, pers. comm.
RU: Tambov	*F. pennsylvanica*	Urban	2017	52.72	41.45	R.N. Ishin, pers. comm.

Location	Species	Habitat	Year	Lat	Lon	Source
RU: Millerovo	*F. pennsylvanica, F. excelsior*	Urban and roadside	2017	48.92	40.39	(Martynov et al. 2017)
RU: Shakhty	*F. pennsylvanica, F. excelsior*	Urban and roadside	2017	47.71	40.22	(Martynov et al. 2017)
RU: Rostov-na-Donu	*F. pennsylvanica, F. excelsior*	Urban and roadside	2017	47.23	39.72	(Martynov et al. 2017)
RU: Krasnodar	*F. pennsylvanica, F. excelsior*	Urban	2018	45.11	39.00	Own data (MOB and AB)
RU: Sochi	*F. excelsior*	Natural forest and urban plantations	2018	43.44	39.93	Own data (MOB and AB)
BY: Stolbtsy	*F. pennsylvanica, F. excelsior*	Urban	2018	53.48	26.74	Own data (MOB and AB)
BY: Mogilev	*F. pennsylvanica, F. excelsior*	Urban	2018	53.89	30.33	Own data (MOB and AB)
BY: Orsha	*F. pennsylvanica, F. excelsior*	Urban	2018	54.51	30.42	Own data (MOB and AB)
BY: Vitebsk	*F. pennsylvanica, F. excelsior*	Urban	2018	55.18	30.20	Own data (MOB and AB)
RU: St.-Petersburg	*F. pennsylvanica*	Urban	2018	59.93	30.28	Own data (MOB and AB)
UA: Donetsk	*F. pennsylvanica, F. excelsior*	Urban and roadside	2019	48.01	37.83	V.V. Martynov, T.V. Nikulina, pers. comm.
RU: Saratov	*F. pennsylvanica, F. excelsior*	Urban	2019	51.52	45.98	V.V. Anikin, M.V. Lavrentiev, pers. comm.
RU: Balashov	*F. pennsylvanica, F. excelsior*	Urban	2019	51.55	43.15	A.N. Volodchenko, pers. comm.
RU: Rzev	*F. pennsylvanica*	Urban	2019	56.26	34.34	Own data (MOB and AB)
RU: Pskov Reg.: Sebezh	*F. excelsior*	Urban	2019	56.30	28.47	Own data (MOB and AB)
RU: Pskov Reg.: Velikie Luki	*F. pennsylvanica*	Urban	2019	56.34	30.53	Own data (MOB and AB)
RU: Tver Reg.: Ostashkov	*F. pennsylvanica, F. excelsior*	Urban	2019	57.14	33.12	Own data (MOB and AB)
RU: Tver Reg.: Bologoye	*F. pennsylvanica, F. excelsior*	Urban	2019	57.89	34.05	Own data (MOB and AB)
RU: Tver Reg.: Vyshniy Volochek	*F. pennsylvanica*	Urban	2019	57.58	34.57	Own data (EP, MOB and AB)
RU: Yaroslavl Reg.: Borok	*F. pennsylvanica*	Urban	2019	58.05	38.25	Own data (AS)

UA: Kharkiv	*F. pennsylvanica, F. excelsior*	Urban	2019	49.99	36.26	Own data (AD)
RU: Tula Reg.: Tulskie Zaseki, Severno-Odoevskoe Forestry	*F. excelsior*	Natural forest	2019	54.01	36.78	Own data (IZ)
RU: Tula Reg.: Tulskie Zaseki, Krapivna	*F. excelsior*	Natural forest	2019	53.97	37.16	Own data (IZ)

Results

Current range

By 2019 *A. planipennis* has been detected in 14 regions of European Russia and Lugansk Region of the Ukraine

(Fig. 5). Our surveys in 2018 and 2019 have shown that *A. planipennis* has spread to the south of European Russia:

Volgograd Region and to the east of the Ukraine: Lugansk Region, but the western border of the range has not

significantly change in the last five years.

Fig. 5 Range of *Agrilus planipennis* in European Russia and the Ukraine in 2019. Blue dots – localities, where *A.*

planipennis was first detected in 2003 –2017. Red dots – localities, where it was first detected in 2018 –2019. Green

squares – localities, where it was not detected during surveys in 2018–2019. Green triangles – localities of surveys

of *Fraxinus excelsior* in the broad-leaved forest Tulskie Zaseki, where ash trees are healthy and have no signs of *A.*

planipennis infestations. Russia: BR – Bryansk Region, KA – Kaluga Region, LI – Lipetsk Region, LU – Lugansk

Region, MO – Moscow Region, OR – Orel Region, RY – Ryazan Region, SM – Smolensk Region, TA – Tambov

Region, TU – Tula Region, TV – Tver Region, VG – Volgograd Region, VL – Vladimir Region, VO – Voronezh

Region, YA – Yaroslavl Region. Kaz - Kazakhstan.

The outbreak in Volgograd Region

The survey of declined plantations of *F. pennsylvanica* (104.9 ha) in Sarpinsky Island in the Volga River (the city of

Volgograd) in October 2018 has shown that the trees have been severely damaged by *A. planipennis*. The island is

occupied with the floodplain forest, which initially consisted of oak, poplar, elm later replaced by artificially planted

green ash. In different parts of the island the degree of damage of ash trees varied from 20–30% to 90–100%. The

official phytoquarantine zone is declared in the island (Rosselkhoznadzor 2019). The decline of the ash plantations

because of the pest outbreak attracted the attention of the local TV (Vesti Volgograd 2018). Since 21 May 2019 the

mass emerging of adult *A. planipennis* in the island began. In 2019 the outbreak was also detected in the left bank of

Volga – in Volga-Akhtuba floodplain (more than 500 damaged *F. pennsylvanica*) and in Sarepta Peninsula (more

than 200 ones). Besides that, one tree damaged by *A. plannipennis* was also detected in the center of Volgograd city

(Mira Street).

The first record in the Ukraine

In 20–22 of June 2019 ash trees in Starokozjij Forest and field shelter belt in its vicinity were examined (Markovka

District of Lugansk Region of the Ukraine; 49.612958 N, 39.56344 E) (the survey was made by A.N. Drogvalenko).

The forest consists of *Quercus* sp., *Acer platanoides*, *Fraxinus excelsior* and *Pyrus* sp. Undergrowth and edges

consist of *Acer campestre, Acer tataricum, Fraxinus excelsior, F. pennsylvanica, Prunus spinosa, Prunus fruticosa*.

Forest belts consisted mainly from *F. pennsylvanica*. As a result of examination of 250 ash trees three trees of *F.*

pennsylvanica damaged by *A. planipennis* were detected. These trees were situated at the edge of the forest belts and

have the diameter of 7–10 cm. Characteristic D-shaped exit holes were situated at a height of 50-200 cm. The

infested trees had dying of upper branches, foliage density reduced (small leaves) and fewer seeds.

Records in Lipetsk region

Lipetsk region is surrounded by the regions where *A. planipennis* was found in 2013 (Orlova-Bienkowskaja 2014a).

The survey by M.J. Orlova-Bienkowskaja in the city of Gryazy in Lipetsk region did not revealed the infestation of

trees by *A. planipennis*, though the trees of *F. pennsylvanica* were in poor condition, and the species *Agrilus*

convexicollis usually associated with *A. planipennis* was found (Orlova-Bienkowskaja and Volkovitsh 2014). In

2018 *A. planipennis* was first found in Lipetsk Region - in the city of Elets (Y.N. Baranchikov, oral communication

in PREPSYS conference, 2018). In June 2019 *A. planipennis* was also found in plantations along the railway in

Leski (Lipetsk Region) (our data, survey was made by S.G. Mazurov.

The results of surveys of trees at the western and north-west border of range

In 2018 we tried to find *A. planipennis* in the east of Belarus: in the cities of Mogilev, Orsha and Vitebsk, but did

not find any signs of infestations in spite of *F. pannsylvanica* and *F. excelsior* are very common in these cities.

Therefore, the westernmost localities of the pest are in Smolensk Region: Semirechje (56 km from the border of
Belarus), where the pest was first recorded in 2014 (Zvyagintsev et al. 2015) and Smolensk (Baranchikov and
Seraya 2018). In the north-west the border of range is in Tver region: In the north-west the most remote locality is
still Zubtsov, where *A. planipennis* was found in 2013 (Straw et al., 2013). In 2019 we failed to find the pest in other
cities of Tver and Pskov Regions, even in the city of Rhev, which is only 15 km north-west to Zubtsov. The
marginal location of range in the north-western outscirt of Tver city, where *A. planipennis* was first detected in 2018
(Peregudova, 2019). It is at the distance of 19 km from Emmaus, where *A. planipennis* was detected in 2013 (Straw
et al., 2013). In the cities of Tver and Smolensk infested trees occur only in restricted small foci, though the most of
ash trees in the cities are have no signs of infestation.

The results of survey of *F. excelsior* in the natural forest Tulskie Zaseki in the center of the pest range

The large broad-leaved forest Tulskie Zaseki (65 000 ha - проверить) is situated in Tula Region in the center of
current range of *A. planipennis*. *Fraxinus excelsior* is one of the main tree species of this forest. The pest was
detected in its vicinity – in Tula and Shchekino as well as in surrounding regions: Kaluga and Orel in 2013 (Straw et
al., 2013; Orlova-Bienkowskaja M. J. 2014a). Trees of *Fraxinus pennsylvanica* are very usual in the cities of Tula
and Shchekino, and now almost all of them are destroyed by the pest (Baranchikov et al 2018b and own
observations by I.A. Zabaluev). In 19–20 of June 2019 more than 500 trees of *F. excelsior* were examined by I.A.
Zabaluev in the forest Tulskie Zaseki in two localities: near Krapivna village and in Severno-Odoevskoe Forestry
(Fig. 6). The trees were situated both in the edge and in the depth of the forest. No signs of infestation by *A.*
planipennis have been found. Ash trees are in good condition.

Fig. 6 The broad-leaved forest Tulskie Zaseki (shaded with green), where *F. excelsior* is usual is shaded with green.

The range of *A. planipennis* is shaded with gray. Green triangles - locations of survey of of *F. excelsior* trees: 1 –

Severno-Odoevskoe Forestry, 2 – vicinity of Krapivna village. Blue dots – localities, where *A. planipennis* has been

detected.

Discussion

Borders of the range

The westernmost localities are in Smolensk region: Semirechje (Zvyagintsev et al. 2015) and Smolensk

(Baranchikov and Seraya 2018). In the north-west the border of the range is in Tver Region: Zubtsov and Tver. The

northernmost locality is Yaroslavl. The border of the range in the north-east is poorly known. The marginal points in

this direction are in Vladimir, Ryazan and Tambov regions. The most eastern and the most southern locality is

Volgograd. In the south-west the border cross Voronezh Region and Bryansk regions of Russia and Lugansk Region

of the Ukraine.

The northern and north-western border of the pest range is almost the same as in 2013. The northernmost

locality - Yaroslavl is at the latitude 57.63' N, i.e. much to the north of the northernmost locality in North America

(47.31' N) (Emereld Ash Borer Info 2019) and in Asia (49.42' N) (Orlova-Bienkowskaja and Volkovitsh 2018).

Therefore, it is unknown if *A. planipennis* will spread further north - to St.-Petersburg and northern Europe, or it has

already reached the range of its potential border in the north. To answer this question the ecological model of

potential range in Europe should be done.

In the south the situation is much worse. The pest has already appeared in the Ukraine, and since there the

outbreaks of the pest in Volgograd, Bryansk and all over Voronezh Region, the pest will probably enter Ukraine and

Belarus with a broad front.

Mechanism of spread is unknown

Natural spread of *A. planipennis* populations is limited to only a few kilometers per year (Siegert et al. 2010). So the

disperse of the pest in the vast territory: 600 000 km^2 in 16 years after the first detection can be explained only by

human-assisted dispersal. The important vector of spread of this pest in North America is unintentional introduction

with firewood (Herms and McCullough 2014). But unlike the United States, ash firewood is not used in European

Russia. Ash wood is also not used for making package in Russia, since the wood of *F. excelsior* is expensive and *F.*

pennsylvanica is an ornamental and landscaping tree not used for timber. It is also doubtful that larvae can be

transported with seedlings, since *A. planipennis* in Moscow Region occur only in the trees more than 10 cm thick

(Orlova-Bienkowskaja 2014). It is suspected that the vector of dispersal is hitchhiking of adult beetles on vehicles

(Straw et al. 2013). Up to now no specimens of *A. planipennis* have been yet detected in vehicles in Russia, but

other Buprestidae sometimes occur in transport. In particular, jewel beetles have been detected in cruise ships sailing

along the Volga River (Komarov E.V., unpublished data). So it is not excluded that *A. planipennis* was introduced

from Moscow to Volgograd by ships. It is also quite possible, that *A. planipennis* appeared in Volgograd, Voronezh

and other regions remote from Moscow not as a result of spread from Moscow, but as a result of multiple

introduction events from Asia or North America. This hypothesis is supported by the fact that the focus of *A.*

planipennis in Volgograd is not young. The death of the trees on large territory indicates that *A. planipennis*

appeared there least 7–10 years ago.

The distance from Moscow to Volgograd is more than 900 km. It is more than the distance from Moscow to
Lithuania, Latvia and Estonia. Obviously, the pest can be detected in any part of Eastern Europe. Besides that, *A.*
planipennis can soon appear in Kazakhstan, since the distance between the Volgograd and the border of country is
about 150 km.
237   Since there are no infested trees of *F. excelsior* in a big forest (in Tula Region) in the center of *A.*
planipennis range, it seems, that European ash is more resistant to the pest than American species, at least in natural
forest stands. All cases of infestation of *F. excelsior* in Russia refer to the trees which grow near plantations of *F.*
pennsylvanica. *Fraxinus pennsylvanica* is highly susceptible to the pest and triggers the outbreak of the pest, which
can also attack *F. excelsior* growing near. The similar situation was observed in China: in 1964 the plantations of *F.*
americana introduced from North America triggered the outbreak of *A. planipennis* in Harbin (Wei et al. 2004). The
pest damaged both *F. americana* and native ash species in the region, but after the trees of *F. americana* were cut
off, the outbreak finished.
245   *Agrilus planipennis* has not become a forest pest in European Russia and Ukraine. It occurs only in
artificial plantations: urban plantations, along motor roads and railroads, in field shelter belts. The overvelming
majority of damaged and killed trees are *F. pennsylvanica*. The only native ash species *F. excelsior* is also
susceptible to *A. planipennis*, but there are still no evidences of massive damage of this ash species in the forests.
There is no doubt that *A. planipennis* will severely damage plantations of *F. pennsylvanica* in Europe. But there is a
hope that natural forests consisting of *F. excelsior* are more resistant to the pest.

References

Baranchikov YN (2013) EAB is the leading abbreviation for European forest protection in the first half of this
253   century. In: Selikhovkin AV, Musolin DL (eds) The Kataev Memorial Readings – 7. Pests and Diseases of
254   Woody Plants in Russia. Proceedings of the International Conference. Saint Petersburg (Russia),
255   November, 25–27, 2013. Saint Petersburg State Forest Technical University, Saint Petersburg, pp.8–9 (**in**
256   **Russian**)
257   Baranchikov YN, Demidko DA, Seraya LG (2018a) A quarter-century of emerald ash borer in Europe. In: Preparing
258   Europe for invasion by the beetles emerald ash borer and bronze birch borer, two major tree-killing pests
259   October 1–4, 2018, Vienna. Abstracts, p. 9
260   Baranchikov YN, Demidko DA, Seraya LG (2018b) Deep in the rear of the invasion: the narrow-leaved ash ash
261   *Agrilus planipennis* (Coleoptera: Buprestidae) in Tula. In: Selikhovkin AV, Musolin DL (eds) The Kataev
262   Memorial Readings – 10. Proceedings of the International Conference. Saint Petersburg (Russia), October,
263   22–25, 2018. Saint Petersburg State Forest Technical University, Saint Petersburg, pp.8–9 (**in Russian**)
264   https://doi.org/10.21266/SPBFTU.2018.KATAEV.1.
265   Baranchikov YN, Gninenko YI,Yurchenko GI (2010) Emerald ash borer in Russia: 2009 situation update. Proc. 21st
266   USDA Interagency research forum on invasive species. USDA FS APHIS: Morgantown. GTR-NRSP-75,
267   pp.66–67
268   Baranchikov YN, Kurteev VV (2012) Invasive area of the emerald ash borer in Europe. Ecological and economical
269   consequences of invasions of dendrophilous insects. Forest Institute of Siberian Branch of Russian
270   Academy of Sciences, Krasnoyarsk, pp 91–94 (**in Russian**)

Baranchikov YN, Mozolevskaya EG, Yurchenko GI, Kenis M (2008) Occurrence of the emerald ash borer (*Agrilus*
planipennis) in Russia and its potential impact on European forestry. EPPO Bull 38:233–238

Baranchikov YN, Seraya LG (2018) Smolensk is occupied by *Agrilus planipennis* (Coleoptera: Buprestidae) ten
years ago. In: Selikhovkin AV, Musolin DL (eds) The Kataev Memorial Readings – 10. Proceedings of the
International Conference. Saint Petersburg (Russia), October, 22–25, 2018. Saint Petersburg State Forest
Technical University, Saint Petersburg, p.10 (**in Russian**)
https://doi.org/10.21266/SPBFTU.2018.KATAEV.1.

Baranchikov YN, Seraya LG, Grinash MN (2014) All European ash species are susceptible to emerald ash borer
Agrilus planipennis Fairmaire (Coleoptera: Buprestidae) – a Far Eastern invader. Siberian Journal of Forest
Science 6:80–85.

Emereld Ash Borer Info (2019) http://www.emeraldashborer.info. Accessed 28 June 2019

Herms DA, McCullough DG (2014) Emerald ash borer invasion of North America: History, biology, ecology,
impacts, and management. Annual Review of Entomology 59:13–30. https://doi.org/10.1146/annurev-ento-
011613-162051

Izhevskii SS, Mozolevskaya EG (2010) *Agrilus planipennis* Fairmaire in Moscow ash trees. Russian Journal of
Biological Invasions 1(3):153–155

Martynov VV, Nikulina TV, Shokhin IV (2017) Modern distribution of invasive dendrophylous insects in Rostov
region. Subtropical and decorative gardening 63:175–182 (**in Russian**)

Orlova-Bienkowskaja MJ (2014a) Ashes in Europe are in danger: the invasive range of *Agrilus planipennis* in
European Russia is expanding. Biological Invasions 16(7):1345–1349 https://doi.org/10.1007/s10530-013-
0579-8

Orlova-Bienkowskaja MJ (2014b) European range of the emerald ash borer *Agrilus planipennis* (Coleoptera:
Buprestidae) is expanding: the pest destroys ashes in the northwest of Moscow Oblast and in part of Tver
Oblast. Russian Journal of Biological Invasions (Springer) 5(1):32–37
https://doi.org/10.1134/S2075111714010081

Orlova-Bienkowskaja MJ, Bieńkowski AO (2016) The life cycle of the emerald ash borer *Agrilus planipennis* in
European Russia and comparisons with its life cycles in Asia and North America. Agricultural and Forest
Entomology 18(2): 182–188 https://doi.org/10.1111/afe.12140

Orlova-Bienkowskaja MJ, Bieńkowski AO (2018) Modeling long-distance dispersal of emerald ash borer in
European Russia and prognosis of spread of this pest to neighboring countries within next 5 years. Ecol
Evol. 8:9295–9304. https://doi.org/10.1002/ece3.4437

Orlova-Bienkowskaja MJ, Volkovitsh MG (2014) Range expansion of *Agrilus convexicollis* in European Russia
expedited by the invasion of emerald ash borer, *Agrilus planipennis* (Coleoptera: Buprestidae). Biological
Invasions 17(2):537–544. https://doi.org/10.1007/s10530-014-0762-6

Orlova-Bienkowskaja MJ, Volkovitsh MG (2018). Are native ranges of the most destructive invasive pests well
known? A case study of the native range of the emerald ash borer, *Agrilus planipennis* (Coleoptera:
Buprestidae). Biological Invasions 20(5):1275-1286. https://doi.org/10.1007/s10530-017-1626-7

Peregudova EY (2016) The first records of buprestid beetle *Agrilus planipennis* in Tver city and *Agrilus*
convexicollis in Tver Oblast. In: Musolin DL, Selikhovkin AV (eds) Dendrobiont invertebrates and fungi
and their role in forest ecosystems Proceedings of the International Conference. Saint Petersburg (Russia),
November 23–25, 2016. Saint Petersburg State Forest Technical University, Saint Petersburg, p.82 (**in**
Russian)

Peregudova EY (2019) The emerald ash borer focus in Tver city, in the North-western border of the invasive range.
Russian Journal of Biological Invasions 2:1–7

Pureswaran DS, Poland TM (2009) Host selection and feeding preference of *Agrilus planipennis* (Coleoptera;
Buprestidae) on ash (*Fraxinus* spp.). Environ Entomol 38:757–765

Rosselkhoznadzor (2014) Rosselkhoznadzor has confirmed the information about detection of the Emerald Ash
Borer in Livny http://www.fsvps.ru/fsvps/print/press/68279.html. Accessed 26 June 2019

Rosselkhoznadzor (2019) Open Data Russia https://data.gov.ru/opendata/7708523530-carantinzone Accessed 26
June 2019

Siegert NW, McCullough DG, Williams DW, Fraser I, Poland TM (2010) Dispersal of *Agrilus planipennis*
(Coleoptera: Buprestidae) from discrete epicenters in two outlier sites. Environ Entomol 39:253–265.
https://doi.org/10.1603/EN09029

Straw NA, Williams DT, Kulinich OA, Gninenko YI (2013) Distribution, impact and rate of spread of emerald ash
borer *Agrilus planipennis* (Coleoptera: Buprestidae) in the Moscow region of Russia. Forestry 86:515–522.
https:// doi.org/10.1093/forestry/cpt031

Volkovitsh MG, Mozolevskaya EG (2014) The tenth «anniversary» of the invasion of emerald ash borer *Agrilus*
planipennis Fairm. (Coleoptera: Buprestidae) in Russia: results and prospects. Izvestia Sankt-Peterburgskoj
Lesotehnicheskoj Akademii 207:8–19 (**in Russian with English summary**)

Zvyagintsev VB, Baranov OY, Panteleev SV (2015) The prevalence of necrosis of ash branches caused by the
invasive pathogen *Hymenoscyphus fraxineus* Baral et al. in the Moscow Oblast and along the highway M1.
Current   Mycology   in   Russia   5(1),   Chapter   10:413–414   (**in   Russian**)   https://
doi.org/10.14427/cmr.2015.v.10

Vesti Volgograd (2018) https://www.youtube.com/watch?v=kV8lHe1Ddc8 Accessed 26 June 2019

Yu C (1992) *Agrilus marcopoli* Obenberger. In: Xiao G (ed) Forest Insects of China, 2nd edn. China Forestry
Publishing House, Beijing, pp.400–401 (**in Chinese**)