Guidelines and considerations for metabarcoding-based port baseline biodiversity

surveys: towards improved marine non-indigenous species monitoring

Anaïs Rey, Oihane C. Basurko, Naiara Rodriguez-Ezpeleta*

AZTI, Marine Research Division, Txatxarramendi ugartea z/g, Sukarrieta 48395 Bizkaia,

Spain

Corresponding author: nrodriguez@azti.es

Running head: Metabarcoding-based port monitoring

Summary

1. Monitoring introduction and spread of non-indigenous species via maritime

transport requires performing port biological baseline surveys. Yet, the

comprehensiveness of these surveys is often compromised by the large number

of habitats present in a port, the seasonal variability of the inhabiting

communities and the time-consuming morphological approach used for

taxonomic classification. Metabarcoding represents a promising alternative for

rapid comprehensive port biological baseline surveys, but, before this technique

can be routinely applied in this context, standardized protocols should be

developed.

2. We applied metabarcoding using two alternative barcodes (based on the

Cytochrome Oxidase I or the 18S ribosomal RNA gene) to about two hundred

port samples collected i) from diverse habitats (water column – including

environmental DNA and zooplankton, sediment and fouling structures), ii) at

different sites within the port (from inner to outer estuary), and iii) during the

four seasons of the year. Comparing the biodiversity metrics and taxonomic

composition derived from each sample group, we investigated the impact of the

different strategies for metabarcoding-based port biodiversity baseline surveys.

3. Each sampling method resulted in a distinct community profile and water

samples alone using universal primers did not produce comprehensive

macroorganismal biodiversity to substitute organismal sampling. Sampling at

different seasons and locations resulted in higher observed biodiversity, but

nestedness analyses suggested that sampling could be reduced to two seasons.

Metabarcoding also allowed to detect previously recorded non-indigenous

species as well as to reveal presence of new ones, even if in low abundance.

4. We provide the first comprehensive evaluation of metabarcoding for port

biodiversity baseline surveys from which we have derived guidelines for how,

when and where samples should be collected. These guidelines are key to

develop the cost-effective and standardizable approach needed for monitoring

non-indigenous species introduction to ports via ballast water and hull fouling,

particularly relevant with the recent entry into force of the International

Convention for the Control and Management of Ships' Ballast Water and

Sediments.

Key-words: ballast water, environmental DNA, environmental monitoring, invasive

species, metabarcoding, non-indigenous species, port biological baseline surveys

Introduction

Globalization has led to an increased maritime transportation, with more and

larger ships than ever before transferring species into ports via ballast water and hull

fouling (Seebens, Gastner, & Blasius, 2013). Among the thousands species arriving daily

(Carlton, 1999), some are Non-Indigenous Species (NIS) and can become invasive,

unbalancing the equilibrium of the recipient ecosystem (Bax, Williamson, Aguero,

Gonzalez, & Geeves, 2003). Thus, being important gateways for introduction of NIS,

ports need to be monitored to provide information required by legal frameworks aiming

at controlling biological invasions (Lehtiniemi et al., 2015).

One of these legal frameworks is the International Convention for the Control

and Management of Ships' Ballast Water and Sediments (IMO, 2004), currently in force.

This convention requires that ships treat their ballast water before its release to port,

unless they show that the risk of transferring NIS between the donor and recipient ports

is limited (David, Gollasch, & Pavliha, 2013). Such risk assessment relies on the

combination of port environmental (temperature and salinity) and biological

information. The obtention of biological information is done by cataloguing biodiversity

through Port Biological Baseline Surveys (PBBS). Given the diverse range of habitats

(such as soft sediment, water column, or exposed and sheltered artificial structures)

allowing presence of distinct organismal groups (such as benthic macrofauna, fouling

organisms or planktonic organisms), comprehensive PBBS require collecting samples

using various methods (Kraus et al., 2018) and at different spatial and temporal scales

(Lehtiniemi et al., 2015).

Global initiatives have been initiated to standardize sampling for PBBS

(HELCOM/OSPAR, 2013; Awad, Haag, Anil, & Abdulla, 2014; Kraus et al., 2018), and are

based on morphological taxonomic identification of the species found. Yet, this method

lacks accuracy for identification of larvae and eggs, developmental stages at which many

NIS are transported (Gittenberger, Rensing, Niemantsverdriet, Schrieken, & Stegenga,

2014), and relies on taxonomists who are often specialized on the local biota, but who

have limited knowledge of alien taxa (Pyšek et al., 2013) and/or on specific taxonomic

groups coexisting in a port (Bishop & Hutchings, 2011). Moreover morphological

identification is time-consuming (Mandelik, Roll, & Fleischer, 2010), especially when

considering the several tens of samples required to characterize a port. These limitations

translate into PBBS being completed several years after sample collection (Bott, 2015),

reducing the effectiveness of prevention of NIS introduction and spread control

strategies.

83   DNA metabarcoding (Taberlet, Coissac, Pompanon, Brochmann, & Willerslev,

2012), the simultaneous identification of taxa present in a complex environmental

sample based on a conserved DNA fragment, is revolutionizing traditional biodiversity

monitoring (Creer et al., 2016; Elbrecht, Vamos, Meissner, Aroviita, & Leese, 2017), and

could represent the cost-effective and possible to standardize alternative required for

PBBS (Lehtiniemi et al., 2015). Metabarcoding can be applied to DNA extracted from

bulk organismal samples (*e.g.* zooplankton) or to environmental DNA (eDNA), defined

as traces of DNA released in the environment in form of cells, feces, skin, saliva, mucus,

etc. (Shaw, Weyrich, & Cooper, 2016). The later has received particular attention lately

for its potential to cost-effectively and noninvasively survey species richness from many

ecosystems (Deiner, Fronhofer, Mächler, Walser, & Altermatt, 2016), and for being able

to detect spatially discrete communities (Jeunen et al., 2019).

Metabarcoding has been described as a promising tool for NIS monitoring

(Comtet, Sandionigi, Viard, & Casiraghi, 2015), and recent studies have highlighted the

potential of metabarcoding applied to eDNA to perform port biodiversity surveys

(Borrell, Miralles, Huu, Mohammed-Geba, & Garcia-Vazquez, 2017; Grey et al., 2018;

Lacoursière-Roussel et al., 2018). Yet, very few studies have performed port surveys

based on additional sampling substrates (Brown, Chain, Zhan, MacIsaac, & Cristescu,

2016; Zaiko et al., 2016) and only one has compared biodiversity assessments obtained

by the different substrates (Koziol et al., 2019). Additionally, few studies have

considered the spatial and temporal variability of ports (*e.g.* eDNA in arctic ports

Lacoursière-Roussel et al. (2018) and settlement plates in austral temperate port Zaiko

et al.(2016)) but none have fully evaluated their impact on all the different communities

of a port, which is critical to ensure maximum biodiversity recovery in port baseline

surveys. In sum, the design of sampling schemes for metabarcoding based PBBS that

best encompass the heterogeneity of ports concerning substrate, locality and season

has not been fully investigated to date.

Here, we have applied metabarcoding to about two hundred samples collected

from the port of Bilbao (Northern Spain) to evaluate the effect of i) using various

sampling methods to capture biodiversity found in different substrates, ii) sampling at

different sites from outer to inner estuary, iii) sampling during different seasons of the

year, and iv) using alternative genetic markers. Our results have allowed to conclude

that combining the alternative sampling substrates and universal primers allows a more

comprehensive port biodiversity assessment, that sampling from a single substrate type

or at a single season does not capture the whole port biodiversity and that the port

salinity gradient needs to be considered when looking for NIS. From our results, we have

derived a series of guidelines and considerations for PBBS that will be particularly

relevant with recent entry into force of the International Convention for the Control and

Management of Ships' Ballast Water and Sediments.

Materials and methods

Sampling

Zooplankton, sediment, fouling and water samples were collected from the port of

Bilbao in autumn 2016, winter 2017, spring 2017, and late summer 2017 at four sites

(Figs 1 and S1). Sites 1, 2 and 3 depict the outer parts of the port with busy berths and

are characterized by deep waters (10-30 m) and salinity (>30 PSU); site 4 depicts the

inner, less busy part of the port and is characterized by low water depth (6-9 m) and

lower salinity (>20 PSU). Zooplankton, sediment and fouling sampling was based on the

protocol designed by Helcom/Ospar (HELCOM/OSPAR, 2013). Zooplankton samples

were collected in vertical tows using Pairovet nets (mesh sizes of 60 and 280 μm) at

three points per site, mixing the collected material to have one sample per mesh size

per site. The collected zooplankton was grinded with a mortar until no integer organism

could be appreciated and was stored in 96% ethanol at -20 °C. Sediment samples were

collected by sieving material collected from a Van Veen grab (0.07–0.1 m^2) with a 1 mm

mesh size sieve. Retained material was processed following Aylagas et al. (2016), and

retrieved benthic macroinvertebrates were stored in 96% ethanol at -20 °C. Fouling

samples were collected by placing 15 x 15 cm polyvinyl chloride plates at 1, 3 and 7 m

depth in each site. Plates were deployed in winter and spring and recovered in spring

and late summer respectively (see Table S1). At recovery, plates were placed in

individual sterile plastic bags, soaked in 96% ethanol and thoroughly scrapped with a

scalpel to retrieve fouling organisms attached to it. The detached organisms were then

homogenized with a blender and stored in 96% ethanol at -20 °C. At each sampling site,

1 L surface water sample was collected from each sampling point with a bottle. Samples

from the same site were combined into a single sample by site. In sites 1 and 3, three

additional samples were taken at maximum depth using a Niskin bottle and combined.

Each 3 L combined water sample was filtered using a 0.45 µm Sterivex filter unit (Merck

Chemicals & Life Science), which was stored at -80 °C until further use. Additionally,

surface water samples from three sampling sites from the ports of Vigo and A Coruña,

located in Galicia, Northwestern Spain, were collected in March 2017 and filtered as

described above.

DNA extraction, library preparation and sequencing

Total genomic DNA was isolated from zooplankton samples with the DNeasy blood &

tissue kit (QIAGEN), benthic macroinvertebrate and fouling organism samples with,

respectively, PowerMax and PowerSoil DNA Isolation Kits (MOBIO) replacing the initial

bead-beating step by an overnight incubation at 56°C with proteinase K (0.4 mg/ml).

Filter samples were extracted with the DNeasy blood & tissue kit (QIAGEN) following the

"SX filters without preservation buffer" developed by Spens et al. (2017). Negative

controls were included in each batch of DNA extraction and followed the whole process.

DNA concentration was measured with the Quant-iT dsDNA HS assay kit (Thermo

Scientific) using a Qubit 2.0 Fluorometer (Life Technologies), purity was inferred from

260/280 and 260/230 absorbance ratios with the ND-1000 Nanodrop (Thermo

Scientific), and integrity was assessed by electrophoresis in 0.7% agarose. Two primer

pairs widely used to assess eukaryotic biodiversity (Hebert, Ratnasingham, & de Waard,

2003; Amaral-Zettler, McCliment, Ducklow, & Huse, 2009) were used:

mlCOIintF/dgHCO2198 ("COI primers"), targeting a 313 bp fragment of the cytochrome

oxidase I (COI) gene (Leray et al., 2013) and 1389F/1510R ("18S primers"), targeting a

variable length fragment (87 to 186 bp) of the V9 hypervariable region of the 18S rRNA

gene (Amaral-Zettler et al., 2009). DNA extracted from zooplankton, fouling organism

and filter samples were amplified with both primer pairs, whereas DNA from sediment

was only amplified with the "COI primers". PCR amplifications were performed in two

rounds. For the first PCR, 1 μl of genomic DNA (5 ng/μl) was added to a mix consisting in

5 μl of 1X Phusion Master Mix (Thermo Scientific), 0.2 μl of each primer (0.2 μM) and

3.6 μl of MilliQ water. For the "18S primers", PCR conditions consisted on an initial 3 min

denaturation step at 98 °C; followed by 25 cycles of 10 s at 98 °C, 30 s at 57 °C and 30 s

at 72 °C and finally 10 min at 72 °C. For the "COI primers", PCR conditions consisted on

an initial 3 min denaturation step at 98 °C; followed by 35 cycles of 10 s at 98 °C, 30 s at

46 °C and 45 s at 72 °C and finally 5 min at 72 °C. Negative controls were included within

each set of PCRs. For each DNA extract, three PCR amplifications were performed and

pooled. Once purified using AMPure XP beads (Beckman Coulter), the mixed PCR

products were used as template for the generation of dual-indexed amplicons in a

second PCR round following the "16S Metagenomic Sequence Library Preparation"

protocol (Illumina) using the Nextera XT Index Kit (Illumina). Multiplexed PCR products

were purified again using the AMPure XP beads, quantified using Quant-iT dsDNA HS

assay kit and a Qubit 2.0 Fluorometer (Life Technologies), normalized to equal

concentration and sequenced using the 2 x 300 paired-end MiSeq (Illumina). Reads were

demultiplexed based on their barcode sequences.

Raw read preprocessing, clustering and taxonomic assignment

After quality checking of demultiplexed paired-end reads with FASTQC (Andrews, 2010),

forward and reverse primers were removed using Cutadapt (Martin, 2011) with the

"anchored 5' adapter" and for "paired end reads" options and with the "linked adapter"

option for COI and 18S respectively. Forward and reverse reads were merged using PEAR

(Zhang, Kobert, Flouri, & Stamatakis, 2014) with a minimum sequence overlap of 217 bp

and maximum amplicon length of 313 bp and of 80 and 190 bp for COI and 18S barcodes

respectively. Merged reads with average Phred quality score lower than 20 were

removed with Trimmomatic (Bolger, Lohse, & Usadel, 2014). Using Mothur (Schloss et

al., 2009), sequences without ambiguous bases were aligned to BOLD

(https://www.boldsystems.org)   or   SILVA   ([https://www.arb-](https://www.arb-silva.de/documentation/release-132/)

[silva.de/documentation/release-132/](https://www.arb-silva.de/documentation/release-132/)) for COI and 18S respectively, and only those

covering the barcode region were kept. Chimeras, detected using *de novo* mode of

UCHIME (Edgar, Haas, Clemente, Quince, & Knight, 2011), were removed, and

remaining reads were clustered into OTUs using Swarm 2.2.1 (Mahé, Rognes, Quince,

de Vargas, & Dunthorn, 2014). "Singleton" OTUs, composed by a single read, were

removed, and the remaining OTUs were taxonomically assigned using the Naïve Bayes

Classifier (Wang, Garrity, Tiedje, & Cole, 2007) using BOLD (accessed in May 2018) or

PR2 (release 4.10.0) databases for COI and 18S respectively.

Community analyses

Apart from the complete dataset, we created two subsets: a taxa targeted through PBBS

dataset*,* including only reads classified to the class level and excluding those matching

to non-targeted groups for PBBS such as Mammalia, Aves, Insecta, Collembola,

Arachnida and all classes of Fungi; and a NICS dataset, including only reads matching

either the Non-Indigenous and Cryptogenic Species (NICS) previously detected in the

port of Bilbao (Martínez & Adarraga, 2006; Adarraga & Martínez, 2011; Butrón, Orive, &

Madariaga, 2011; Adarraga & Martínez, 2012; Zorita et al., 2013; Tajadura, Bustamante,

& Salinas, 2016) or the AquaNIS database (AquaNIS. Editorial board., 2015). Most

analyses were conducted using RStudio (Team, 2015) with *vegan* (Oksanen et al., 2013),

adespatial (Dray et al., 2018) and *indicspecies* (De Cáceres & Legendre, 2009) libraries.

Indicator species analyses (Dufrene & Legendre, 1997) were performed on 1) the taxa

targeted through PBBS dataset to identify indicator taxa of each sampling method and

2) the NICS dataset to identify non-indigenous indicator taxa of each sampling site.

These analyses were based on the IndVal index calculated as the product of the degree

of specificity (measuring the uniqueness to a sampling method or site) and the degree

of fidelity (measuring the frequency of occurrence within a sampling method or site) of

an OTU to a given sampling condition. Statistical significance of associations was

assessed by performing 10,000 permutations. The effects of season and locality on taxa

targeted through PBBS communities were tested for significance using a permutational

multivariate analysis of variance (PERMANOVA) after checking for multivariate

homogeneity of group dispersions (betadisper). PERMANOVA and betadisper were

performed on OTU abundance (Hellinger distance) and presence/absence (Jaccard

dissimilarities). The contribution of replacement (changes in OTU identity) and

nestedness (richness differences where one sample is a subset of a richer sample) to

beta diversity of taxa targeted through PBBS between seasons and between sites was

computed using the relativized nestedness index of Podani and Schmera (2011) based

on Jaccard dissimilarities matrix. For each sampling method, replacement and

nestedness were calculated between pairwise comparisons of 1) samples belonging to

the same site but sampled at different seasons (season variation of beta diversity), and

2) samples belonging to the same season but sampled at different sites (spatial variation

of beta diversity). The mean proportion of nestedness and replacement contribution to

beta-diversity between sites and between seasons was then calculated.

Results

Communities retrieved by each sampling method and genetic marker

A total of 5,718,639 and 7,055,675 COI and 18S barcode reads were kept for

analysis (Table S1). Negative controls showed little contamination (with only 19, 36 and

55 of COI and 31 and 741 18S reads belonging to taxa targeted through PBBS). The reads

corresponding to the 192 samples collected at the port of Bilbao resulted in 40,318 and

20,473 OTUs for COI and 18S respectively. For all sampling methods, rarefaction curves

of OTUs against reads approached saturation, indicating enough sequencing effort (Figs

2A and S2). The majority of the OTUs, 89% for COI and 73% for 18S, were unique to one

sampling method (Fig. 2B), and, for both barcodes, but particularly for 18S, filtered

water (aimed at retrieving eDNA from macroorganisms and microbial eukaryotes)

resulted in higher OTU richness and unique OTUs than other sampling methods (Figs 2A,

B and S2). Different sampling methods retrieved distinct biological communities as

observed with the PCA (Fig. 2C) and confirmed with PERMANOVA analyses (COI:

R^2=0.28, p-value= 9.99e-05; 18S: R^2=0.30, p-value=9.99e-05).

The percentage of reads assigned to species or higher taxonomic levels differed

between sampling methods and barcodes. The majority of the reads were assigned to

species level for zooplankton nets (COI: 84% and 18S: 75%), settlement plates (COI: 69%

and 18S: 69%) and sediment grabs (COI: 64%); for filtered water, although 51% of reads

were assigned to species level with 18S, only 8% could be assigned to a species with COI

(Fig. 3A). Filtered water analyzed with COI had also the largest proportion of reads that

could not be assigned to phylum (87%). This is possibly due to the amplification of non-

metazoan taxa, which are underrepresented in the BOLD database (Fig. S3), which was

confirmed with BLAST searches against GenBank (data not shown).

Both barcodes detected a wide range of eukaryotic groups (Fig. 3B). With COI,

the greatest majority of reads belonged to metazoans, with Arthropoda dominating all

sampling methods except sediment grabs, dominated by Annelida. With 18S, a more

diverse spectrum of taxa was retrieved, including phytoplankton and macroalgae; as

expected, the number of reads assigned to these phyla in zooplankton nets and

settlement plates was very low in comparison to filtered water.

Distribution of taxa targeted through port biological baseline survey

A total of 16,828 and 9,091 OTUs were kept as taxa targeted through PBBS for

COI and 18S respectively. From those, indicator analysis identified respectively 2,600

(15%) and 1,700 (19%) OTUs significantly associated to one of the sampling methods.

Settlement plates were associated with 1,268 (COI) and 808 (18S) OTUs, zooplankton

nets with 1,001 (COI) and 315 (18S) OTUs, and sediment grabs with 238 (COI) OTUs. Only

12 OTUs (9 of which were metazoans) were indicators of filtered water with COI while

580 (3 of which were metazoans), were associated to this sampling method with 18S.

We observed expected strong associations of some taxa to one particular sampling

method: fouling bivalves (Ostreoida, Mytilida) with settlement plates, copepods

(Calanoida, Cyclopoida) with zooplankton nets and sea urchins (Spatangoida) with

sediment grabs (Fig. 4). Yet, less obvious associations were also observed such as

barnacles (Sessilia) and two polychaeta orders (Spionida, Phyllodocida) with

zooplankton nets or dinoflagellates (Peridiniales and Gymnodiniales) with settlement

plates, illustrating the advantages of a complementary sampling approach to recover

the diversity of these taxonomic groups.

The need for complementarity of sampling methods was further confirmed by

PCA on PBBS target taxa (Fig. 5). For both barcodes, expected taxonomic differences

were found between the sampling methods: sediment grabs were characterized by the

polychaeta *Maldane glebifex* (Fig. 5B), zooplankton nets were distinguished by several

copepods (Fig. 5B, D) and settlement plates, by the encrusting *Semibalanus balanoides*

(Fig. 5D). Filtered water communities were characterized differently according to each

barcode. For COI, filtered water samples were not differentiated as a distinct group and

were in general close to those retrieved with zooplankton samples collected at the same

season (Fig. 5A). In contrast, for 18S, filtered water was different due to the presence of

the phytoplankton species *Phaeocystis globosa* (Fig. 5D); yet, when targeting only

metazoan taxa, the patterns observed with COI were similarly to the ones retrieved with

18S (Fig. S4A). Concerning filtered samples, when looking only at metazoan taxa, the

proportion of OTUs detected was largely different compared to the other methods (Fig.

S5). For instance, Decapoda orders had a low OTU diversity with filtered water (COI:

0,7%; 18S: 7%), whereas more OTU diversity could be detected for Calanoida (COI: 8%;

18S: 35%), Sabellida (COI: 15%; 18S: 28%) and Leptothecata (COI: 13%; 18S: 58%). In

general, filtered water recovered the smallest metazoan OTU diversity.

Influence of sampling seasonality and locality on detected biodiversity

Seasonal variations contributed significantly to differences in community composition of zooplankton nets and filtered water but not of sediment grabs (Table S2). Total OTU richness and unique OTU richness varied between seasons for all sampling methods (Fig. 6A, B). Spring and late summer were generally richer than autumn and winter, excepted for filtered water communities depicted with 18S. For sediment grabs and zooplankton nets with 18S, a strong proportion of OTUs found in autumn and winter were a subset of OTUs retrieved in late summer and spring. Indeed, the nestedness component of sediment grab assemblages represented between 60 to 70% of total compositional variations among these pairs of seasons (Fig. 6D). For zooplankton net assemblages, 63 and 72% of nestedness was observed between late summer and spring and between late summer and autumn respectively, suggesting that late summer recovered a majority of spring and autumn diversity (Fig. 6D). Thus, the combination of spring and late summer for sediment grabs retrieved 85% of the total OTU richness over the four seasons, while for zooplankton nets, in late summer alone 71% was retrieved (Fig. 6C, E). For filtered water and settlement plates, seasonal community changes were driven by both OTU replacement and nestedness, with similar relative contributions (Fig. 6D). The seasonal influence observed in intra-port samples was also observed between ports (Fig. S6). The differences in communities between Bilbao, A Coruña and Vigo (ports belonging to the same ecoregion but separated by over 500 km) during the same season was generally smaller than that between seasons in the same port, indicating that communities were driven by seasonality rather than location (Fig. S6 and Table S3). This pattern was more pronounced with 18S than with COI.

Locality within the Bilbao port appeared to impact on benthic assemblages (Table

S2), since sites outside the estuary (sites 1 to 3) were different from site 4 inside the

estuary; mainly, the polychaete *Maldane glebifex* was less abundant in site 4 (Fig. 5B).

Regarding zooplankton nets, site 4 was the main driver of difference in communities as,

when not considered, no significant differences between sites were observed (Table S2).

Each site harbored a similar proportion of the total OTU richness found by each sampling

method (Fig. 7C). This proportion did not exceed 60%. Indeed, OTU replacement

contributed more to community variation among sites than nestedness, especially with

sediment grab and settlement plates (Fig. 7D). OTU replacement was more important

when comparing sites from outside the estuary (sites 1, 2 and S) with site 4 inside the

estuary, while it contributed less to community variation among sites 1,2 and 3. This is

congruent with site 4 having generally more unique OTUs than the other sites (Fig. 7B).

An exception was observed for filtered water with COI, where site 4 had the lowest OTU

richness and unique OTU richness, and where nestedness contributed more to the

variation in community composition in comparison to sites 1 and 2 (Fig. 7A, B, D).

Detection of Non-Indigenous and Cryptogenic Species (NICS)

Our port baseline biodiversity survey detected 79 NICS, among which 29 were

previously recorded in the port Bilbao (Table S4). Most of the other species (43) were

previously detected NICS inside the port's Large Marine Ecoregion (LME) but, for 7 of

them (*Ammothea hilgendorfi, Bugulina fulva, Grandidierella japonica, Melita nitida,*

Neodexiospira brasiliensis, Pseudochattonella verruculosa, Tubificoides pseudogaster),

it was, to our knowledge, the first report in the LME. The indicator species analysis

performed on non-indigenous OTU revealed that site 4 was associated to the highest

non-indigenous and cryptogenic diversity with, for COI, 603 OTU indicators, and for 18S,

24 OTU indicators. These OTUs belonged to 10 NICS with *Oithona davisae*, *Acartia tonsa*

and *Allita succinea*, found by both barcodes, *Ficopomatus enigmaticus,* with 18S and

Amphibalanus eburneus, *Grandidierella japonica*, *Polydora cornuta*, *Austrominius*

modestus, *Monocorophium acherusicum*, *Xenostrobus securis,* with COI. Regarding the

three other sites, for COI, site 1 was associated to one *Clytia hemisphaerica* OTU and to

one *Balanus trigonus* OTU, site 2, to three *Clytia hemisphaerica* and four *Mytilus edulis*

OTUs and site 3, to one *Balanus trigonus* OTU. For 18S, only site 1 was associated to one

OTU, assigned to *Gymnodinium aureolum.*

Discussion

In order to be comprehensive, port baseline surveys need to have an extensive spatial

and temporal coverage so that most biodiversity is captured by the collected samples.

Unlike morphological taxonomic identification, metabarcoding allows the analysis of

hundreds of samples is a cost-effective manner (Aylagas, Borja, Muxika, & Rodríguez-

Ezpeleta, 2018), enabling for comprehensive PBBS. Metabarcoding-based PBBS can

follow a taxonomy dependent or independent approach. The first approach requires

comparing the sequences against reference databases for taxonomic assignment,

whose accuracy strongly depends on the completeness of the reference database; this

reference database is particularly relevant for detection of NICS. The second approach

consists on inferring diversity metrics from sequences to compare across samples for

which the development of standardized procedures is mandatory so that comparisons

are possible. Here we have focused on developing such procedures by evaluating

alternative strategies.

A comprehensive metabarcoding-based port baseline survey should rely on a combination of sampling methods

Most of the past attempts on using metabarcoding for port monitoring have relied on a unique sampling method (Brown et al., 2016; Zaiko et al., 2016; Borrell et al., 2017; Grey et al., 2018; Lacoursière-Roussel et al., 2018) and only one evaluated the importance of sampling methods (Koziol et al., 2019). Here, in order to define comprehensive guidelines for developing standardizable port sampling protocols, we have adapted the HELCOM/OSPAR guidelines to collect zooplankton, fouling, and sediment samples for genetic analyses. In agreement with previous work (Koziol et al., 2019), our metabarcoding analysis of the collected samples has shown that each sampling method recovered a distinct subset of the port community, and that, despite some taxa being expectedly associated to a given sampling method, the total diversity, even within specific taxonomic groups, was only recovered by combining different methods. Thus, combining sampling methods is needed to maximize biodiversity detection.

Metabarcoding of eDNA from water samples using universal primers does not provide a comprehensive view of the overall macroorganismal diversity

As an attempt to find a sampling method that would capture a large fraction of the diversity with limited effort, we also explored metabarcoding on eDNA extracted from filtered water samples. Recent applications of this technique have shown its potential to recover spatial and temporal variation of macroorganismal communities (Bista et al., 2017; O'Donnell et al., 2017), including detection of non-indigenous taxa

(Klymus, Marshall, & Stepien, 2017). Consequently, recent studies have considered

eDNA for port surveys (Borrell et al., 2017; Grey et al., 2018; Lacoursière-Roussel et al.,

2018). Yet, as suggested by other studies in coral reef sites (Djurhuus et al., 2018) and

freshwater streams (Macher et al., 2018) and recently in ports (Koziol et al., 2019), our

analyses show that, compared to bulk sample metabarcoding, eDNA metabarcoding

recovered only a subset of the metazoan diversity and did not provide additional

information on targeted groups. This suggests that, despite requiring less sampling

effort, COI and 18S based metabarcoding of eDNA obtained from filtered water cannot

be used to replace the need of obtaining bulk samples (using zooplankton nets, grabs

and settlement plates) for PBBS taxa detection. In this context, at least two alternatives

are possible to improve eDNA-based biodiversity assessments: i) increasing sequencing

depth (Grey et al., 2018) or ii) using group specific markers (Koziol et al., 2019). Yet, none

of them ensures full biodiversity recovery and both significantly increase costs. Thus, the

decision between increasing sampling depth and/or using multiple group specific

primers for eDNA and adding multiple substrate sampling needs to be carefully

considered and will be an important area of future research to optimize metabarcoding-

based port monitoring.

A comprehensive metabarcoding-based port baseline survey should include

spatiotemporal sampling

While it is expected that increased temporal coverage will retrieve more taxa,

the costs associated to extensive sampling demands an evaluation of the benefits gained

by sampling at each season. Indeed, the HELCOM/OSPAR protocol limits sampling to late

summer for sediment and fouling, and to spring and late summer for plankton. Here we

show that for sediment and fouling, spring sampling provides higher diversity than late

summer. Although sampling in late summer could be appropriate for morphological

taxonomy because of the more abundance of adult individuals, for metabarcoding,

sampling in spring is preferable because during this season i) sizes of organisms are less

variable and thus metabarcoding is less likely to under detect small organisms (Elbrecht,

Peinert, & Leese, 2017) and ii) species diversity is at its maxima due to being a high

recruitment period with abundant organisms at early life stages (Bijleveld et al., 2018).

For zooplankton diversity, our results show that the HELCOM/OSPAR approach of

sampling in spring and late summer produces the highest diversity. However, the data

obtained from filtered water were inconclusive. Although spring was the most diverse

season for COI, this was not the case for 18S, and, overall, other combinations of seasons

appeared to perform better. Previous studies have already shown that seasonal

variations are important considerations for eDNA studies (Lacoursière-Roussel et al.,

2018), but further characterizations are needed over multiple years to design an

adequate protocol for maximizing biodiversity recovery with eDNA. Our study also

highlights the influence of sampling site in recovered biodiversity, although only one of

the four sites sampled, the less saline one located at the inner part of the estuary,

showed a significantly different community composition.

Concerning spatial sampling, we observed no significant differences between

sites when using eDNA, except between the only site located in the inner estuary, and

the others and only when using zooplankton, and significant differences overall when

using sediment grabs. The proportion of the total OTUs found in a single site was low,

suggesting that spatially comprehensive sampling is crucial to recover the port´s

biodiversity. Interestingly, site 4 was not only the most different from all four sites but

was also the one that recorded the largest number of NIS. In coherence with our

findings, it has been observed that brackish environments favors NIS settlement (Zorita

et al., 2013) because these species usually support wider range of salinity (Cardeccia et

al., 2018). Thus, samples for port monitoring should include those with a wide range of

abiotic conditions and covering priority sampling, such as highly active ship berths,

potential reservoirs of newly arrived NIS (Hewitt & Martin, 2001).

Metabarcoding provides valuable information on Non-Indigenous Species

Our metabarcoding port biodiversity baseline survey detects species previously

recorded in the port of Bilbao and species known to be present in the port´s large marine

ecoregion. Importantly, it also detects presence of seven species for which no records

exist so far in this ecoregion, highlighting the potential of metabarcoding for early NICS

detection. Nonetheless, not all NICS species previously recorded in Bilbao were found

by our analyses. This might be due to these species not being present in the port at the

time of our survey, to biases of the metabarcoding process such as differential DNA

extraction, primer non-specificity, etc (Xiong, Li, & Zhan, 2016), or to database

incompleteness. Primer bias and/or database incompleteness could also explain why 35

and 42% of NICS were uniquely detected with COI and 18S respectively. Yet, from the

total species included in the AquaNIS database (n=1083), only 460 and 369 are included

in the BOLD and PR2 databases respectively. This stresses the need of increasing

reference databases and of using multiple universal primers for species detection (Grey

et al., 2018). Importantly, our study confirmed that metabarcoding can detect species

occurring at low abundance (Pochon, Bott, Smith, & Wood, 2013) as we found the non-

indigenous amphipod *Melita nitida* with only 18 reads while it took intensive surveys in

2013, 2014 and 2016 in 3 distinct sampling regions of the Bay of Biscay to record 76

individuals (Gouillieux, Lavesque, Blanchet, & Bachelet, 2016).

Guidelines and recommendations for standardized and comparable metabarcoding-

based port baseline surveys

Based on the comparative analysis of samples collected using alternative sampling

methods, at different seasons of the year and from different port locations, we have

determined the effect of each factor on derived biodiversity estimates. This has allowed

to define a series of standard guidelines that maximize recovered biodiversity while

minimizing sampling effort and which can be applied to ensure comparability among

ports:

1. Sampling site selection: Obtain information about port traffic and potentially

489   relevant abiotic conditions (*i.e,* salinity gradient) and select at least three sites,

490   including active and inactive berths and covering the salinity range (*i.e.* marine

491   to brackish).

492   2. Sampling method selection: Collect samples from, at least, zooplankton nets,

493   sediment grabs, filtered water and settlement plates (see Material and Methods

494   for detailed sampling techniques).

495   3. Sampling season selection: In temperate latitudes, collect benthic (sediment

496   grabs) and fouling (settlement plates) macrofauna at least in spring and late

497   summer, zooplankton at least in late summer and eDNA (from filtered water) at

498   least in spring.

499   4. Metabarcoding data generation: Follow standardized protocols for sample pre-

500   processing, DNA extraction, barcode amplification, using at least two universal

primers, and sequencing (see Material and Methods) and use established

procedures for read preprocessing, OTU clustering and taxonomic assignment

(see provided analysis code).

5. Non-Indigenous Species detection: Compare the list of species found in the port

with comprehensive and updated databases of Non-Indigenous Species (*i.e.*

AquaNIS) for a first indication of potential invaders presence.

These guidelines and considerations are particularly relevant due to the recent entry

into force of the International Convention for the Control and Management of Ships'

Ballast Water and Sediments as they will allow performing standardized risk

assessments to evaluate the possibility of granting an exemption for ships travelling

between specific ports.

Acknowledgments

This manuscript is a result of the Aquainvad-ED (AQUAtic INVADers: Early Detection,

Control and Management - www.aquainvad-ed.com) project funded by the European

Union (Horizon 2020 research and innovation program under the Marie Sklodowska-

Curie grant agreement no 642197). We thank the Port Authority of Bilbao for authorizing

sampling in the port and Manuel González for making the necessary arrangements for

it. We also thank Iñaki Mendibil and Elisabete Bilbao for laboratory work, Ekaitz

Erauskin, Iker Urtizberea and Lander Larrañaga Juarez for sampling, Anders Lanzen for

providing comments on the manuscript, and Carlota Bermejo, student of the

postgraduate course in Scientific Illustration of the UPV/EHU, for producing Figure 1.

This manuscript is contribution X [to be completed upon acceptance] from the Marine

Research Division of AZTI.

Authors' contributions

AR, OB and NR-E conceived the study and designed methodology; AR and OB collected the samples; AR preformed laboratory analyses; AR and NR-E analyzed the data; AR and NR-E wrote the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Data Accessibility

Raw demultiplexed MiSeq reads are available at NCBI SRA Bioproject PRJNA515494. Scripts used for OTU table generation and taxonomic assignment are available at https://github.com/rodriguez-ezpeleta/metabarcoding-pbbs.

References

Adarraga, I., & Martínez, J. (2011). First record of invasive *Theora lubrica* (Mollusca: Bivalvia: Semelidae) in the Atlantic Ocean. *Marine Biodiversity Records*, *4*. doi:10.1017/S1755267211001011

Adarraga, I., & Martínez, J. (2012). First record of the invasive brackish water mytilid *Limnoperna securis* (Lamarck, 1819) in the Bay of Biscay. *Aquatic Invasions*, *7*(2).

Amaral-Zettler, L. A., McCliment, E. A., Ducklow, H. W., & Huse, S. M. (2009). A Method for Studying Protistan Diversity Using Massively Parallel Sequencing of V9 Hypervariable Regions of Small-Subunit Ribosomal RNA Genes. *PLOS ONE*, *4*(7), e6372. doi:10.1371/journal.pone.0006372

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.

AquaNIS. Editorial board. (2015). Information system on Aquatic Non-Indigenous and

Cryptogenic   Species.   World   Wide   Web   electronic   publication.

www.corpi.ku.lt/databases/aquanis. Version 2.36+.

Awad, A., Haag, F., Anil, A. C., & Abdulla, A. (2014). *Guidance on Port Biological Baseline*

Surveys (PBBS). GEF-UNDP-IMO GloBallast Partnerships, IOI, NIO, IUCN.

Aylagas, E., Borja, Á., Muxika, I., & Rodríguez-Ezpeleta, N. (2018). Adapting

metabarcoding-based benthic biomonitoring into routine marine ecological

status   assessment   networks.   *Ecological   Indicators*,   *95*,   194–202.

doi:10.1016/j.ecolind.2018.07.044

Aylagas, E., Mendibil, I., Borja, Á., & Rodríguez-Ezpeleta, N. (2016). Marine Sediment

Sample Pre-processing for Macroinvertebrates Metabarcoding: Mechanical

Enrichment   and   Homogenization.   *Frontiers   in   Marine   Science*,   *3*.

doi:10.3389/fmars.2016.00203

Bax, N., Williamson, A., Aguero, M., Gonzalez, E., & Geeves, W. (2003). Marine invasive

alien species: a threat to global biodiversity. *Marine Policy*, *27*(4), 313–323.

doi:10.1016/S0308-597X(03)00041-1

Bijleveld, A. I., Compton, T. J., Klunder, L., Holthuijsen, S., Horn, J. ten, Koolhaas, A., …

Veer, H. W. van der. (2018). Presence-absence of marine macrozoobenthos does

not generally predict abundance and biomass. *Scientific Reports*, *8*(1), 3039.

doi:10.1038/s41598-018-21285-1

Bishop, M. J., & Hutchings, P. A. (2011). How useful are port surveys focused on target

pest identification for exotic species management? *Marine Pollution Bulletin*,

62(1), 36–42. doi:10.1016/j.marpolbul.2010.09.014

Bista, I., Carvalho, G. R., Walsh, K., Seymour, M., Hajibabaei, M., Lallias, D., … Creer, S.

(2017). Annual time-series analysis of aqueous eDNA reveals ecologically

relevant dynamics of lake ecosystem biodiversity. *Nature Communications*, *8*,

14087. doi:10.1038/ncomms14087

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for

Illumina   sequence   data.   *Bioinformatics*,   *30*(15),   2114–2120.

doi:10.1093/bioinformatics/btu170

Borrell, Y. J., Miralles, L., Huu, H. D., Mohammed-Geba, K., & Garcia-Vazquez, E. (2017).

DNA in a bottle—Rapid metabarcoding survey for early alerts of invasive species

in ports. *PLOS ONE*, *12*(9), e0183347. doi:10.1371/journal.pone.0183347

Bott, N. J. (2015). Detection of Invasive Species. In S.-K. K. Prof (Ed.), *Springer Handbook*

of Marine Biotechnology (pp. 441–453). Springer Berlin Heidelberg.

doi:10.1007/978-3-642-53971-8_16

Brown, E. A., Chain, F. J. J., Zhan, A., MacIsaac, H. J., & Cristescu, M. E. (2016). Early

detection of aquatic invaders using metabarcoding reveals a high number of non-

indigenous species in Canadian ports. *Diversity and Distributions*, *22*(10), 1045–

1059. doi:10.1111/ddi.12465

Butrón, A., Orive, E., & Madariaga, I. (2011). Potential risk of harmful algae transport by

ballast waters: The case of Bilbao Harbour. *Marine Pollution Bulletin*, *62*(4), 747–

757.

Cardeccia, A., Marchini, A., Occhipinti-Ambrogi, A., Galil, B., Gollasch, S., Minchin, D., …

Ojaveer, H. (2018). Assessing biological invasions in European Seas: Biological

traits of the most widespread non-indigenous species. *Estuarine, Coastal and*

Shelf Science, *201*, 17–28. doi:10.1016/j.ecss.2016.02.014

Carlton, J. T. (1999). The scale and ecological consequences of biological invasions in the

World's oceans. (pp. 195–212). Presented at the Population and Community

Biology Series Volume 24, Kluwer Academic Publishers.

Comtet, T., Sandionigi, A., Viard, F., & Casiraghi, M. (2015). DNA (meta) barcoding of

biological invasions: a powerful tool to elucidate invasion processes and help

managing aliens. *Biological Invasions*, *17*(3), 905–922.

Creer, S., Deiner, K., Frey, S., Porazinska, D., Taberlet, P., Thomas, W. K., … Bik, H. M.

(2016). The ecologist's field guide to sequence-based identification of

biodiversity. *Methods in Ecology and Evolution*, *7*(9), 1008–1018.

doi:10.1111/2041-210X.12574

David, M., Gollasch, S., & Pavliha, M. (2013). Global ballast water management and the

'same location' concept: a clear term or a clear issue? *Ecological Applications*,

23(2), 331–338. doi:10.1890/12-0992.1

De Cáceres, M. D., & Legendre, P. (2009). Associations between species and groups of

sites: indices and statistical inference. *Ecology*, *90*(12), 3566–3574.

Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J.-C., & Altermatt, F. (2016).

Environmental DNA reveals that rivers are conveyer belts of biodiversity

information. *Nature Communications*, *7*, 12544. doi:10.1038/ncomms12544

Djurhuus, A., Pitz, K., Sawaya, N. A., Rojas-Márquez, J., Michaud, B., Montes, E., …

Breitbart, M. (2018). Evaluation of marine zooplankton community structure

through environmental DNA metabarcoding: Metabarcoding zooplankton from

eDNA. *Limnology and Oceanography: Methods*, *16*(4), 209–221.

doi:10.1002/lom3.10237

Dray, S., Bauman, D., Blanchet, G., Borcard, D., Clappe, S., Guenard, G., … Wagner, H. H.

(2018). *adespatial: Multivariate Multiscale Spatial Analysis*. Retrieved from

https://CRAN.R-project.org/package=adespatial

Dufrene, M., & Legendre, P. (1997). Species Assemblages and Indicator Species: The

Need for a Flexible Asymmetrical Approach. *Ecological Monographs*, *67*(3), 345.

doi:10.2307/2963459

Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., & Knight, R. (2011). UCHIME improves

sensitivity and speed of chimera detection. *Bioinformatics*, *27*(16), 2194–2200.

doi:10.1093/bioinformatics/btr381

Elbrecht, V., Peinert, B., & Leese, F. (2017). Sorting things out: Assessing effects of

unequal specimen biomass on DNA metabarcoding. *Ecology and Evolution*,

7(17), 6918–6926. doi:10.1002/ece3.3192

Elbrecht, V., Vamos, E. E., Meissner, K., Aroviita, J., & Leese, F. (2017). Assessing

strengths and weaknesses of DNA metabarcoding-based macroinvertebrate

identification for routine stream monitoring. *Methods in Ecology and Evolution*,

8(10), 1265–1275. doi:10.1111/2041-210X.12789

Gittenberger, A., Rensing, M., Niemantsverdriet, P., Schrieken, N., & Stegenga, H. (2014,

November). Port of Rotterdam survey and monitoring non-native species

conform HELCOM/OSPAR protocol.

Gouillieux, B., Lavesque, N., Blanchet, H., & Bachelet, G. (2016). First record of the non-

indigenous *Melita nitida Smith*, 1873 (Crustacea: Amphipoda: Melitidae) in the

Bay of Biscay (NE Atlantic). *BioInvasions Records*, *5*(2), 85–92.

Grey, E. K., Bernatchez, L., Cassey, P., Deiner, K., Deveney, M., Howland, K. L., … Lodge,

D. M. (2018). Effects of sampling effort on biodiversity patterns estimated from

environmental DNA metabarcoding surveys. *Scientific Reports*, *8*(1).

doi:10.1038/s41598-018-27048-2

Hebert, P. D., Ratnasingham, S., & de Waard, J. R. (2003). Barcoding animal life:

cytochrome c oxidase subunit 1 divergences among closely related species.

Proceedings of the Royal Society of London B: Biological Sciences, *270*(Suppl 1),

S96–S99.

HELCOM/OSPAR. (2013). Joint HELCOM/OSPAR Guidelines for the Contracting Parties of

OSPAR and HELCOM on the granting of exemptions under International

Convention for the Control and Management of Ships' Ballast Water and

Sediments, Regulation A-4 (OSPAR Agreement 2013-09), 45 pp.

Hewitt, C. L., & Martin, R. B. (2001). Revised protocols for baseline port surveys for

introduced marine species–design considerations, sampling protocols and

taxonomic sufficiency. *Centre for Research on Introduced Marine Pests Technical*

Report, (22).

IMO. (2004). International Convention for the Control and Management of Ships' Ballast

Water and Sediments, 2004. International Maritime Organization, 13 Feb 2004,

London.

Jeunen, G.-J., Knapp, M., Spencer, H. G., Lamare, M. D., Taylor, H. R., Stat, M., …

Gemmell, N. J. (2019). Environmental DNA (eDNA) metabarcoding reveals strong

discrimination among diverse marine habitats connected by water movement.

Molecular Ecology Resources, *19*(2), 426–438. doi:10.1111/1755-0998.12982

Klymus, K. E., Marshall, N. T., & Stepien, C. A. (2017). Environmental DNA (eDNA)

metabarcoding assays to detect invasive invertebrate species in the Great Lakes.

PLOS ONE, *12*(5), e0177643. doi:10.1371/journal.pone.0177643

Koziol, A., Stat, M., Simpson, T., Jarman, S., DiBattista, J. D., Harvey, E. S., … Bunce, M.

(2019). Environmental DNA metabarcoding studies are critically affected by

substrate selection. *Molecular Ecology Resources*, *19*(2), 366–376.

doi:10.1111/1755-0998.12971

Kraus, R., Ninčević-Gladan, Ž., Auriemma, R., Bastianini, M., Bolognini, L., Cabrini, M., …

Žuljević, A. (2018). Strategy of port baseline surveys (PBS) in the Adriatic Sea.

Marine Pollution Bulletin. doi:10.1016/j.marpolbul.2018.08.067

Lacoursière-Roussel, A., Howland, K., Normandeau, E., Grey, E. K., Archambault, P.,

Deiner, K., … Bernatchez, L. (2018). eDNA metabarcoding as a new surveillance

approach for coastal Arctic biodiversity. *Ecology and Evolution*.

doi:10.1002/ece3.4213

Lehtiniemi, M., Ojaveer, H., David, M., Galil, B., Gollasch, S., McKenzie, C., … Pederson,

J. (2015). Dose of truth—Monitoring marine non-indigenous species to serve

legislative requirements. *Marine Policy*, *54*, 26–35.

doi:10.1016/j.marpol.2014.12.015

Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., … Machida, R. J.

(2013). A new versatile primer set targeting a short fragment of the

mitochondrial COI region for metabarcoding metazoan diversity: application for

characterizing coral reef fish gut contents. *Frontiers in Zoology*, *10*, 34.

doi:10.1186/1742-9994-10-34

Macher, J.-N., Vivancos, A., Piggott, J. J., Centeno, F. C., Matthaei, C. D., & Leese, F.

(2018). Comparison of environmental DNA and bulk-sample metabarcoding

using highly degenerate cytochrome c oxidase I primers. *Molecular Ecology*

Resources. doi:10.1111/1755-0998.12940

Mahé, F., Rognes, T., Quince, C., de Vargas, C., & Dunthorn, M. (2014). Swarm: robust

and fast clustering method for amplicon-based studies. *PeerJ*, *2*, e593.

doi:10.7717/peerj.593

Mandelik, Y., Roll, U., & Fleischer, A. (2010). Cost-efficiency of biodiversity indicators for

Mediterranean ecosystems and the effects of socio-economic factors. *Journal of*

Applied Ecology, *47*(6), 1179–1188. doi:10.1111/j.1365-2664.2010.01864.x

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.journal*, *17*(1), 10–12. doi:10.14806/ej.17.1.200

Martínez, J., & Adarraga, I. (2006). Programa de vigilancia y control de la introducción

de especies invasoras en los ecosistemas litorales de la costa vasca. 1. *Costa de*

Gipuzkoa, *2*.

O'Donnell, J. L., Kelly, R. P., Shelton, A. O., Samhouri, J. F., Lowell, N. C., & Williams, G.

D. (2017). Spatial distribution of environmental DNA in a nearshore marine

habitat. *PeerJ*, *5*, e3044. doi:10.7717/peerj.3044

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R. B., …

Wagner, H. (2013). Package 'vegan'. *Community Ecology Package, Version*, *2*(9).

Pochon, X., Bott, N. J., Smith, K. F., & Wood, S. A. (2013). Evaluating detection limits of

next-generation sequencing for the surveillance and monitoring of international

marine pests. *PLoS One*, *8*(9), e73935.

Podani, J., & Schmera, D. (2011). A new conceptual and methodological framework for

exploring and explaining pattern in presence - absence data. *Oikos*, *120*(11),

1625–1638. doi:10.1111/j.1600-0706.2011.19451.x

Pyšek, P., Hulme, P. E., Meyerson, L. A., Smith, G. F., Boatwright, J. S., Crouch, N. R., …

Wilson, J. R. U. (2013). Hitting the right target: taxonomic challenges for, and of,

plant invasions. *AoB PLANTS*, *5*. doi:10.1093/aobpla/plt042

Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., …

Weber, C. F. (2009). Introducing mothur: Open-Source, Platform-Independent,

Community-Supported Software for Describing and Comparing Microbial

Communities. *Applied and Environmental Microbiology*, *75*(23), 7537–7541.

doi:10.1128/AEM.01541-09

Seebens, H., Gastner, M. T., & Blasius, B. (2013). The risk of marine bioinvasion caused

by global shipping. *Ecology Letters*, *16*(6), 782–790. doi:10.1111/ele.12111

Shaw, J. L. A., Weyrich, L., & Cooper, A. (2016). Using environmental (e)DNA sequencing

for aquatic biodiversity surveys: a beginner's guide. *Marine and Freshwater*

Research. doi:10.1071/MF15361

Spens, J., Evans, A. R., Halfmaerten, D., Knudsen, S. W., Sengupta, M. E., Mak, S. S. T., …

Hellström, M. (2017). Comparison of capture and storage methods for aqueous

macrobial eDNA using an optimized extraction protocol: advantage of enclosed

filter. *Methods in Ecology and Evolution*, *8*(5), 635–645. doi:10.1111/2041-

210X.12683

Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012). Towards

next-generation biodiversity assessment using DNA metabarcoding. *Molecular*

Ecology, *21*(8), 2045–2050.

Tajadura, J., Bustamante, M., & Salinas, J. I. S. (2016). Temporal and spatial variability in

the distribution of non-indigenous zoobenthic species in the intertidal hard

substrates in the 'Abra of Bilbao' Poster ISOBAY conference 2016

https://www.ehu.eus/documents/5019180/0/Book-of-abstract-ISOBAY15.pdf.

ISOBAY Conference.

Team, Rs. (2015). RStudio: integrated development for R. *RStudio, Inc., Boston, MA URL*

Http://Www. Rstudio. Com, *42*.

Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naive Bayesian classifier for

rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied*

and Environmental Microbiology, *73*(16), 5261–5267. doi:10.1128/AEM.00062-

07

Xiong, W., Li, H., & Zhan, A. (2016). Early detection of invasive species in marine

ecosystems using high-throughput sequencing: technical challenges and possible

solutions. *Marine Biology*, *163*(6), 1–12.

Zaiko, A., Schimanski, K., Pochon, X., Hopkins, G. A., Goldstien, S., Floerl, O., & Wood, S.

A. (2016). Metabarcoding improves detection of eukaryotes from early

biofouling communities: implications for pest monitoring and pathway

management. *Biofouling*, *32*(6), 671–684. doi:10.1080/08927014.2016.1186165

Zhang, J., Kobert, K., Flouri, T., & Stamatakis, A. (2014). PEAR: a fast and accurate

Illumina Paired-End reAd mergeR. *Bioinformatics*, *30*(5), 614–620.

doi:10.1093/bioinformatics/btt593

Zorita, I., Solaun, O., Borja, A., Franco, J., Muxika, I., & Pascual, M. (2013). Spatial

distribution and temporal trends of soft-bottom marine benthic alien species

collected during the period 1989–2008 in the Nervión estuary (southeastern Bay

of Biscay). *Journal of Sea Research*, *83*, 104–110.

Figures

Figure 1. Sampling protocol depicting sampling location (Port of Bilbao), sites (1, 2, 3 and

4) and points per site (dots); and illustrating the sampling methods used and the

targeted biological communities.

Figure 2. Overall description of community detected per barcode. A: OTU accumulation

curves per sampling method. B: Venn diagrams of the number and percentage of OTUs

shared between sampling methods. C: Principal Component Analyses of the Hellinger-

transformed abundances of OTUs. Sample scores are displayed in scaling 1 with ellipses

representing the 95% confidence dispersion of each sampling method.

770 Figure 3. Taxonomic assignment per barcode and per sampling method. A: Percentage

771 of reads assigned to each taxonomic level. B: Relative abundance of reads classified to

772 at least Phylum level for the 10 most abundant phyla.

784 Figure 4. Distribution of indicator OTUs associated to each sampling method at the order

785 level (in log scale) for COI and 18S. Only orders with at least 10 indicator OTUs are shown.

Figure 5. Principal Component Analyses of Hellinger-transformed abundances of OTUs included in the *PBBS targeted taxa dataset* for COI (A-B) and 18S (C-D). A & C: Samples scores in scaling 1 with ellipses representing the 95% confidence dispersion of each sampling method. B & D: OTU scores in scaling 1 with the circle of equilibrium contribution. Only OTUs whose projected length in these two principal component axes exceeding the value of equilibrium contribution are represented.

Figure 6. Seasonal variation of alpha and beta diversity for each sampling method with

COI and 18S. A: Total OTU richness recovered at each season. B: OTU richness unique to

each season. C: Proportion of the total OTU richness detected with one season. D:

Decomposition of between-season beta diversity into replacement and nestedness

components. E: Proportion of the total OTU richness detected with two seasons.

807   Figure 7. Spatial variation of alpha and beta diversity for each sampling method with COI

808   and 18S. A: Total OTU richness recovered at each site. B: OTU richness unique to each

809   site. C: Proportion of the total OTU richness detected with one site. D: Decomposition

810   of between-site beta diversity into replacement and nestedness components