Considerations for metabarcoding-based port biological baseline surveys aimed at

marine non-indigenous species monitoring and risk-assessments

Anaïs Rey, Oihane C. Basurko, Naiara Rodriguez-Ezpeleta*

AZTI, Marine Research Division, Txatxarramendi ugartea z/g, Sukarrieta 48395 Bizkaia,

Spain

Corresponding author: nrodriguez@azti.es

Running head: Metabarcoding-based port monitoring

Abstract

1. Monitoring introduction and spread of non-indigenous species via maritime
11

    transport and performing risk assessments require port biological baseline
12

    surveys. Yet, the comprehensiveness of these surveys is often compromised by
13

    the large number of habitats present in a port, the seasonal variability and the
14

    time-consuming morphological approach used for taxonomic identification.
15

    Metabarcoding represents a promising alternative for rapid comprehensive
16

    port biological baseline surveys, but its application in this context requires
17

    further assessments.
18

2. We applied metabarcoding (based on barcodes of the Cytochrome c oxidase
19

    subunit I and of the 18S ribosomal RNA gene) to 192 port samples collected i)
20

    from diverse habitats (water column – including environmental DNA and
21

    zooplankton, sediment and fouling structures), ii) at different sites (from inner
22

    to outer estuary), and iii) during the four seasons of the year.
23

3. By comparing the biodiversity metrics derived from each sample group, we
24

    show that each sampling method resulted in a distinct community profile and
25

    that environmental DNA alone cannot substitute for organismal sampling, and
26

    that, although sampling at different seasons and locations resulted in higher
27

    observed biodiversity, operational results can be obtained by sampling selected
28

    locations and seasons.
29

4. By assessing the taxonomic composition of the samples, we show that
30

    metabarcoding data allowed the detection of previously recorded non-
31

    indigenous species as well as to reveal presence of new ones, even if in low
32

    abundance.
33

5. *Synthesis and application.* Our comprehensive assessment of metabarcoding

for port biological baseline surveys sets the basics for cost-effective,

standardized and comprehensive monitoring of non-indigenous species and for

performing risk assessments in ports. This development will contribute to the

implementation of the recently entered into force International Convention for

the Control and Management of Ships' Ballast Water and Sediments.

Key-words: ballast water, environmental DNA, environmental monitoring, invasive

species, metabarcoding, non-indigenous species, port biological baseline surveys

Introduction

Globalization has led to an increased maritime transportation, with more and larger ships than ever before transferring species into ports via ballast water and hull fouling (Seebens, Gastner, & Blasius, 2013). Among the thousands species arriving daily (Carlton, 1999), some are Non-Indigenous Species (NIS) and can become invasive, disrupting native populations (Bax, Williamson, Aguero, Gonzalez, & Geeves, 2003). Thus, being important gateways for introduction of NIS, ports need to be monitored to provide information required by legal frameworks aiming at controlling biological invasions (Lehtiniemi et al., 2015).

One of these legal frameworks is the International Convention for the Control and Management of Ships' Ballast Water and Sediments (IMO, 2004), which requires that ships treat their ballast water before its release to port, unless they show that the risk of transferring NIS between the donor and recipient ports is limited (David, Gollasch, & Pavliha, 2013). Such risk assessment requires cataloguing biodiversity through Port Biological Baseline Surveys (PBBS), which requires collecting samples using various methods (Kraus et al., 2018) and at different spatial and temporal scales (Lehtiniemi et al., 2015) given the diverse range of habitats (such as soft sediment, water column, or exposed and sheltered artificial structures) allowing presence of distinct organismal groups (such as benthic macrofauna, fouling organisms or planktonic organisms).

Global initiatives have been initiated to standardize sampling for PBBS (HELCOM/OSPAR, 2013; Awad, Haag, Anil, & Abdulla, 2014; Kraus et al., 2018) and are based on morphological taxonomic identification of the species found. Yet, this method lacks accuracy for identification of larvae and eggs, developmental stages at

which many NIS are transported (Gittenberger, Rensing, Niemantsverdriet, Schrieken,

& Stegenga, 2014) and relies on taxonomists who are often specialized on the local

biota, but who have limited knowledge of alien taxa (Pyšek et al., 2013) and/or on

specific taxonomic groups coexisting in a port (Bishop & Hutchings, 2011). Moreover

morphological identification is time-consuming (Mandelik, Roll, & Fleischer, 2010),

especially when considering the several tens of samples required to characterize a

port. These limitations translate into PBBS being completed several years after sample

collection (Bott, 2015), reducing the effectiveness of prevention of NIS introduction

and spread control strategies.

DNA metabarcoding (Taberlet, Coissac, Pompanon, Brochmann, & Willerslev,

2012), the simultaneous identification of taxa present in a complex environmental

sample based on a conserved DNA fragment, is revolutionizing traditional biodiversity

monitoring (Creer et al., 2016; Elbrecht, Vamos, Meissner, Aroviita, & Leese, 2017),

and could represent the cost-effective and possible to standardize alternative required

for PBBS (Lehtiniemi et al., 2015). Metabarcoding can be applied to DNA extracted

from bulk organismal samples (community DNA) or to environmental DNA (eDNA),

which is extraorganismal DNA released in the environment in form of cells, feces, skin,

saliva, mucus, etc. (Shaw, Weyrich, & Cooper, 2016). The later has received particular

attention lately for its potential to cost-effectively and noninvasively survey species

richness from many ecosystems (Deiner, Fronhofer, Mächler, Walser, & Altermatt,

2016), and for being able to detect spatially discrete communities (Jeunen, Knapp,

Spencer, Lamare, et al., 2019).

Metabarcoding has been extensively explored and validated as an alternative

tool to morphological identification (e.g. Aylagas et al. (2016; 2018), and described as a

promising tool for NIS monitoring (Comtet, Sandionigi, Viard, & Casiraghi, 2015).

Recent studies have highlighted the potential of metabarcoding applied to eDNA to

perform port biodiversity surveys (Borrell, Miralles, Huu, Mohammed-Geba, & Garcia-

Vazquez, 2017; Grey et al., 2018; Lacoursière-Roussel et al., 2018). Yet, very few

studies have performed port surveys based on additional sampling substrates (Brown,

Chain, Zhan, MacIsaac, & Cristescu, 2016; Zaiko et al., 2016) and only one has

compared biodiversity assessments obtained by the different substrates (Koziol et al.,

2019). Additionally, few studies have considered the spatial and temporal variability of

ports (*e.g.* eDNA in arctic ports Lacoursière-Roussel et al. (2018) and settlement plates

in austral temperate port Zaiko et al. (2016)) but none has fully evaluated the impact

of spatial and temporal variability on all the different communities of a port, which is

critical to ensure maximum biodiversity recovery in PBBS.

Here, we have applied metabarcoding to 192 samples collected from the port

of Bilbao (Northern Spain) to evaluate the effect of i) using various sampling methods

to capture biodiversity found in different substrates, ii) sampling at different sites from

outer to inner estuary, iii) sampling during different seasons of the year, and iv) using

alternative genetic markers. Our results provide relevant information for performing

non-indigenous species monitoring and risk assessments in ports in response to the

International Convention for the Control and Management of Ships' Ballast Water and

Sediments.

Materials and methods

Sampling

Zooplankton, sediment, fouling and water samples were collected from the port of Bilbao in autumn 2016, winter 2017, spring 2017, and late summer 2017 at four sites (Fig 1 and Appendix 1). Sites 1, 2 and 3 represent the outer parts of the port with busy berths and are characterized by deep waters (10-30 m) and salinity (>30 PSU); site 4 represent s the inner, less busy part of the port and is characterized by low water depth (6-9 m) and lower salinity (>20 PSU). Zooplankton, sediment and fouling sampling was based on the protocol designed by Helcom/Ospar (HELCOM/OSPAR, 2013). Zooplankton samples were collected in vertical tows using Pairovet nets (mesh sizes of 60 and 280 μm) at three points per site, mixing the collected material to have one sample per mesh size per site. The collected zooplankton was grinded with a mortar until no integer organism could be appreciated and was stored in 96% ethanol at -20 °C. Sediment samples (3 samples per site) were collected by sieving material collected from a Van Veen grab (0.07–0.1 m^2) with a 1 mm mesh size sieve. Retained material was processed following Aylagas et al. (2016) based on decantation and homogenization phases, and retrieved benthic macroinvertebrates were stored in 96% ethanol at -20 °C. Homogenization was done by blending the decanted organic material in a PHILIPS hr2095 700 W 2 L glass jar with 96% ethanol until no fragments of animals and others organic materials could be observed. During decantation and homogenization steps, all laboratory equipment used were cleaned between samples by soaking in 10% bleach for 15 min and thoroughly rinsing with deionized water. Fouling samples were collected by placing 15 x 15 cm polyvinyl chloride plates at 1, 3 and 7 m depth in a suspended array in each site. Plates were deployed in winter and

spring and recovered in spring and late summer respectively (see Table S1). At

recovery, were placed in individual sterile plastic bags, soaked in 96% ethanol and

thoroughly scrapped both sides with a scalpel to retrieve fouling organisms attached to

it. The detached organisms were then homogenized with a blender (Conair™ Waring™

Mini-sample Containers for Blender) in 96% ethanol and stored in 96% ethanol at -20

°C. Cleaning procedure was similar as the one performed for sediment samples. At

each sampling site, 1 L surface water sample was collected from each sampling point

with a bottle. Samples from the same site and from the same sampling method were

combined into a single sample by site. In sites 1 and 3, three additional samples were

taken at one meter above the bottom using a Niskin bottle and combined. Each 3 L

combined water sample was filtered using a 0.45 µm Sterivex filter unit (Merck

Chemicals & Life Science), which was stored at -80 °C until further use. Additionally, 3 L

surface water sample from three sampling sites from the ports of Vigo and A Coruña,

located in Galicia, Northwestern Spain following the same sampling procedure than for

Bilbao by combining 1 L water from three sampling point per sampling site, were

collected in March 2017 and filtered as described above.

DNA extraction, library preparation and sequencing

Total genomic DNA was isolated from zooplankton samples with the DNeasy blood &

tissue kit (QIAGEN), benthic macroinvertebrate and fouling organism samples with,

respectively, PowerMax and PowerSoil DNA Isolation Kits (MOBIO) replacing the initial

bead-beating step by an overnight incubation at 56°C with proteinase K (0.4 mg/ml).

Filter samples were extracted with the DNeasy blood & tissue kit (QIAGEN) following

the "SX filters without preservation buffer" developed by Spens et al. (2017). Negative

controls (preservation buffer) were included in each batch of DNA extraction and

followed the same procedure as all other samples. DNA concentration was measured

with the Quant-iT dsDNA HS assay kit (Thermo Scientific) using a Qubit 2.0 Fluorometer

(Life Technologies), purity was inferred from 260/280 and 260/230 absorbance ratios

with the ND-1000 Nanodrop (Thermo Scientific), and integrity was assessed by

electrophoresis in 0.7% agarose. Two primer pairs widely used to assess eukaryotic

biodiversity (Hebert, Ratnasingham, & de Waard, 2003; Amaral-Zettler, McCliment,

Ducklow, & Huse, 2009) were used: mlCOIintF/dgHCO2198 (COI primers), targeting a

313 bp fragment of the Cytochrome c oxidase subunit I (COI) gene (Meyer, 2003; Leray

et al., 2013) and 1389F/1510R (18S primers), targeting a variable length fragment (87

to 186 bp) of the V9 hypervariable region of the 18S rRNA gene (Amaral-Zettler et al.,

2009). DNA extracted from zooplankton, fouling organism and filter samples were

amplified with both primer pairs, whereas DNA from sediment was only amplified with

the COI primers. PCR amplifications were performed in two rounds. For the first PCR, 1

µl of genomic DNA (5 ng/µl) was added to a mix consisting in 5 µl of 2X Phusion Master

Mix (Thermo Scientific), 0.2 µl of each primer (0.2 µM) and 2.6 µl of MilliQ water. For

the 18S primers, PCR conditions consisted on an initial 3 min denaturation step at 98

°C; followed by 25 cycles of 10 s at 98 °C, 30 s at 57 °C and 30 s at 72 °C and finally 10

min at 72 °C. For the COI primers, PCR conditions consisted of an initial 3 min

denaturation step at 98 °C; followed by 35 cycles of 10 s at 98 °C, 30 s at 46 °C and 45 s

at 72 °C and finally 5 min at 72 °C. Negative controls of PCR (no template) were

included within each set of PCRs. For each DNA extract, three PCR amplifications were

performed and pooled. Once purified using AMPure XP beads (Beckman Coulter), the

mixed PCR products were used as template for the generation of dual-indexed

amplicons in a second PCR round following the "16S Metagenomic Sequence Library

Preparation" protocol (Illumina) using the Nextera XT Index Kit (Illumina). Multiplexed

PCR products were purified again using the AMPure XP beads, quantified using Quant-

iT dsDNA HS assay kit and a Qubit 2.0 Fluorometer (Life Technologies), normalized to

equal concentration and sequenced using the 2 x 300 paired-end MiSeq (Illumina).

Reads were demultiplexed based on their barcode sequences.

Raw read preprocessing, clustering and taxonomic assignment

After quality checking of demultiplexed paired-end reads with FASTQC (Andrews,

2010), forward and reverse primers were removed using Cutadapt (Martin, 2011) with

the anchored 5' adapter and for "paired end reads" options and with the linked

adapter option for COI and 18S respectively. Forward and reverse reads were merged

using PEAR (Zhang, Kobert, Flouri, & Stamatakis, 2014) with a minimum sequence

overlap of 217 bp and maximum amplicon length of 313 bp and of 80 and 190 bp for

COI and 18S barcodes respectively. Merged reads with average Phred quality score

lower than 20 were removed with Trimmomatic (Bolger, Lohse, & Usadel, 2014). Using

Mothur (Schloss et al., 2009), sequences without ambiguous bases were aligned to

BOLD (https://www.boldsystems.org) or SILVA (https://www.arb-

silva.de/documentation/release-132/) for COI and 18S respectively, and only those

covering the barcode region were kept. Chimeras, detected using *de novo* mode of

UCHIME (Edgar, Haas, Clemente, Quince, & Knight, 2011), were removed, and

remaining reads were clustered into OTUs using Swarm 2.2.1 with the step-by-step

aggregation clustering algorithm implemented with default settings (Mahé, Rognes,

Quince, de Vargas, & Dunthorn, 2014). SWARM algorithm does not rely on a fixed

threshold for delimiting OTUs which is pertinent when performing PBBS where highly

diverse biodiversity can be found. "Singleton" OTUs, composed by a single read, were

removed. No rarefaction was performed to avoid decreasing sensitivity by choosing an

arbitrary minimum library size (McMurdie and Holmes, 2014). The remaining OTUs

were taxonomically assigned using the Naïve Bayes Classifier (Wang, Garrity, Tiedje, &

Cole, 2007) using BOLD (accessed in May 2018) or PR2 (release 4.10.0) databases for

COI and 18S respectively.

Community analyses

Apart from the complete dataset, we created two subsets: a taxa targeted through

PBBS dataset, including only reads classified to the class level and excluding those

matching to non-targeted groups for PBBS such as Mammalia, Aves, Insecta,

Collembola, Arachnida and all classes of Fungi; and a NICS dataset, including only reads

matching either the 68 Non-Indigenous and Cryptogenic Species (NICS) previously

detected in the port of Bilbao (Martínez & Adarraga, 2006; Adarraga & Martínez, 2011,

2012; Butrón, Orive, & Madariaga, 2011; Zorita et al., 2013; Tajadura, Bustamante, &

Salinas, 2016) or the 1083 species present in the AquaNIS database (AquaNIS. Editorial

board., 2015). Most analyses were conducted using RStudio (Team, 2015) with *vegan*

(Oksanen et al., 2013), *adespatial* (Dray et al., 2018) and *indicspecies* (De Cáceres &

Legendre, 2009) libraries. Indicator species analyses (Dufrene & Legendre, 1997) were

performed on 1) the taxa targeted through PBBS dataset to identify indicator taxa of

each sampling method and 2) the NICS dataset to identify non-indigenous indicator

taxa of each sampling site. These analyses were based on the IndVal index calculated

as the product of the degree of specificity (measuring the uniqueness to a sampling

method or site) and the degree of fidelity (measuring the frequency of occurrence

within a sampling method or site) of an OTU to a given sampling condition. Statistical

significance of associations was assessed by performing 10,000 permutations. The

effects of season and locality on taxa targeted through PBBS communities were tested

for significance using a permutational multivariate analysis of variance (PERMANOVA)

after checking for multivariate homogeneity of group dispersions (betadisper).

PERMANOVA and betadisper were performed on Euclidean distances on Hellinger-

transformed OTU abundances (Hellinger distances), which are appropriate for

community ordination and clustering (Legendre & Gallagher, 2001) and on Jaccard

dissimilarities based on OTU presence/absence. The contribution of replacement

(changes in OTU identity) and nestedness (richness differences where one sample is a

subset of a richer sample) to beta diversity of taxa targeted through PBBS between

seasons and between sites was computed using the relativized nestedness index of

Podani and Schmera (2011) based on Jaccard dissimilarities matrix. For each sampling

method, replacement and nestedness were calculated between pairwise comparisons

of 1) samples belonging to the same site but sampled at different seasons (season

variation of beta diversity), and 2) samples belonging to the same season but sampled

at different sites (spatial variation of beta diversity). The mean proportion of

nestedness and replacement contribution to beta-diversity between sites and between

seasons was then calculated. For all OTUs assigned to a Non-Indigenous and

Cryptogenic Species, we blasted their representative sequences against BOLD and PR2

databases respectively for COI and 18S barcodes.

Results

Communities retrieved by each sampling method and genetic marker

A total of 5,718,639 and 7,055,675 COI and 18S barcode reads were kept for analysis (Table S1). Negative controls showed little contamination (with only 164, 66 and 213 of COI and 2031 and 163 18S quality filtered reads). The reads corresponding to the 192 samples collected at the port of Bilbao resulted in 40,318 and 20,473 OTUs for COI and 18S respectively. For all sampling methods, OTU accumulation curves of OTUs against reads approached saturation, suggesting that adding more sequencing effort would provide limited increase in diversity (Figs 2A and Appendix 2). The majority of the OTUs, 89% for COI and 73% for 18S, were unique to one sampling method (Fig. 2B), and, for both barcodes, but particularly for 18S, filtered water (aimed at retrieving eDNA from macroorganisms and microbial eukaryotes) resulted in higher OTU richness and unique OTUs than other sampling methods (Figs 2A, B and Appendix 2). Different sampling methods retrieved distinct biological communities as observed with the PCA (Fig. 2C) and confirmed with PERMANOVA analyses (COI: R^2=0.28, p-value= 9.99e-05; 18S: R^2=0.30, p-value=9.99e-05).

The percentage of reads assigned to species or higher taxonomic levels differed between sampling methods and barcodes. The majority of the reads were assigned to species level for zooplankton nets (COI: 84% and 18S: 75%), settlement plates (COI: 69% and 18S: 69%) and sediment grabs (COI: 64%); for filtered water, although 51% of reads were assigned to species level with 18S, only 8% could be assigned to a species with COI (Fig. 3A). Filtered water analyzed with COI had also the largest proportion of reads that could not be assigned to phylum (87%). This was due to the non-specific

amplification of prokaryotic and non-target eukaryotic DNAs (Appendix 3), which is

known for COI (Collins et al., 2019).

Both barcodes detected a wide range of eukaryotic groups (Fig. 3B). With COI,

the greatest majority of reads belonged to metazoans, with Arthropoda dominating all

sampling methods except sediment grabs, dominated by Annelida. With 18S, a more

diverse spectrum of taxa was retrieved, including phytoplankton and macroalgae; as

expected, the number of reads assigned to these phyla in zooplankton nets and

settlement plates was very low in comparison to filtered water.

Distribution of taxa targeted through port biological baseline survey

A total of 16,828 and 9,091 OTUs were kept as taxa targeted through PBBS for

COI and 18S respectively. From those, indicator analysis identified respectively 2,600

(15%) and 1,700 (19%) OTUs significantly associated to one of the sampling methods.

Settlement plates were associated with 1,268 (COI) and 808 (18S) OTUs, zooplankton

nets with 1,001 (COI) and 315 (18S) OTUs, and sediment grabs with 238 (COI) OTUs.

Only 12 OTUs (9 of which were metazoans) were indicators of filtered water with COI

while 580 (3 of which were metazoans), were associated to this sampling method with

18S. We observed expected strong associations of some taxa to one particular

sampling method: fouling bivalves (Ostreoida, Mytilida) with settlement plates,

copepods (Calanoida, Cyclopoida) with zooplankton nets and sea urchins (Spatangoida)

with sediment grabs (Fig. 4). Yet, less obvious associations were also observed such as

barnacles (Sessilia) and two polychaeta orders (Spionida, Phyllodocida) with

zooplankton nets or dinoflagellates (Peridiniales and Gymnodiniales) with settlement

plates, illustrating the advantages of a complementary sampling approach to recover

the diversity of these taxonomic groups.

The need for complementarity of sampling methods was further confirmed by

PCA on PBBS target taxa (Fig. 5). For both barcodes, expected taxonomic differences

were found between the sampling methods: sediment grabs were characterized by the

polychaeta *Maldane glebifex* (Fig. 5B), zooplankton nets were distinguished by several

copepods (Fig. 5B, D) and settlement plates, by the encrusting *Semibalanus balanoides*

(Fig. 5D). Filtered water communities were characterized differently according to each

barcode. For COI, filtered water samples were not differentiated as a distinct group

and were in general close to those retrieved with zooplankton samples collected at the

same season (Fig. 5A). In contrast, for 18S, filtered water was different due to the

presence of the phytoplankton species *Phaeocystis globosa* (Fig. 5D); yet, when

targeting only metazoan taxa, the patterns observed with COI were similarly to the

ones retrieved with 18S (Appendix 4). For filtered samples, the proportion of metazoan

OTUs detected was largely different compared to the other methods (Appendix 5). For

instance, Decapoda orders had a low diversity with filtered water (COI: 0,7%; 18S: 7%),

whereas more diversity could be detected for Calanoida (COI: 8%; 18S: 35%), Sabellida

(COI: 15%; 18S: 28%) and Leptothecata (COI: 13%; 18S: 58%). In general, filtered water

recovered the smallest metazoan diversity.

Influence of sampling seasonality and locality on detected biodiversity

Seasonal variations contributed significantly to differences in community

composition of zooplankton nets and filtered water but not of sediment grab (Table

S2). Total richness and unique richness varied between seasons for all sampling

methods (Fig. 6A, B). Spring and late summer were generally richer than autumn and

winter, excepted for filtered water communities represented with 18S. For sediment

grabs and zooplankton nets with 18S, a strong proportion of taxa found in autumn and

winter were a subset of taxa retrieved in late summer and spring. Indeed, the

nestedness component of sediment grab assemblages represented between 60 to 70%

of total compositional variations among these pairs of seasons (Fig. 6D). For

zooplankton net assemblages, 63 and 72% of nestedness was observed between late

summer and spring and between late summer and autumn respectively, suggesting

that late summer recovered a majority of spring and autumn diversity (Fig. 6D). Thus,

the combination of spring and late summer for sediment grabs retrieved 85% of the

total OTU richness over the four seasons, while for zooplankton nets, in late summer

alone 71% was retrieved (Fig. 6C, E). For filtered water and settlement plates, seasonal

community changes were driven by both OTU replacement and nestedness, with

similar relative contributions (Fig. 6D). The seasonal influence observed in intra-port

samples was also observed between ports (Appendix 6). The differences in

communities between Bilbao, A Coruña and Vigo (ports belonging to the same

ecoregion but separated by over 500 km) during the same season was generally

smaller than that between seasons in the same port, indicating that communities were

driven by seasonality rather than location (Appendix 6 and Table S3). This pattern was

more pronounced with 18S than with COI.

Locality within the Bilbao port appeared to impact on benthic assemblages

(Table S2), since sites outside the estuary (sites 1 to 3) were different from site 4 inside

the estuary; mainly, the polychaete *Maldane glebifex* was less abundant in site 4 (Fig.

5B). Regarding zooplankton nets, site 4 was the main driver of difference in

communities as, when not considered, no significant differences between sites were

observed (Table S2). Each site harbored a similar proportion of the total OTU richness

found by each sampling method (Fig. 7C). This proportion did not exceed 60%. Indeed,

OTU replacement contributed more to community variation among sites than

nestedness, especially with sediment grab and settlement plates (Fig. 7D). OTU

replacement was more important when comparing sites from outside the estuary

(sites 1, 2 and S) with site 4 inside the estuary, while it contributed less to community

variation among sites 1,2 and 3. This is congruent with site 4 having generally more

unique OTUs than the other sites (Fig. 7B). An exception was observed for filtered

water with COI, where site 4 had the lowest OTU richness and unique OTU richness,

and where nestedness contributed more to the variation in community composition in

comparison to sites 1 and 2 (Fig. 7A, B, D).

Detection of Non-Indigenous and Cryptogenic Species (NICS)

Our port baseline biological survey detected 79 putative NICS, among which 29

of the 68 previously recorded in the port Bilbao were found (Tables S4, 5 and 6). Most

of the other species (43) were previously detected NICS inside the port's Large Marine

Ecoregion (LME) but, for 7 of them (*Ammothea hilgendorfi, Bugulina fulva,*

Grandidierella japonica, Melita nitida, Neodexiospira brasiliensis, Pseudochattonella

verruculosa, Tubificoides pseudogaster), it was, to our knowledge, the first report in

the LME. The indicator species analysis performed on reads corresponding to non-

indigenous species revealed that site 4 was associated to the highest number of non-

indigenous and cryptogenic species, which were *Oithona davisae*, *Acartia tonsa* and

Allita succinea, found by both barcodes, *Ficopomatus enigmaticus,* found with 18S and

Amphibalanus eburneus, Grandidierella japonica, Polydora cornuta, Austrominius

modestus, Monocorophium acherusicum, Xenostrobus securis, found with COI.

Regarding the three other sites, for COI, site 1 was associated to *Clytia hemisphaerica*

and to *Balanus trigonus*, site 2, to *Clytia hemisphaerica* and *Mytilus edulis* and site 3, to

Balanus trigonus. For 18S, only site 1 was associated to *Gymnodinium aureolum.*

Discussion

Metabarcoding-based port baseline surveys require a combination of sampling

methods and should not rely solely on eDNA

Most of the past attempts on using metabarcoding for port monitoring have

relied on a single sampling method (Brown et al., 2016; Zaiko et al., 2016; Borrell et al.,

2017; Grey et al., 2018; Lacoursière-Roussel et al., 2018) and only one evaluated the

importance of using different sampling methods (Koziol et al., 2019). In agreement

with the latter, our metabarcoding analysis shows that each sampling method

recovered a distinct subset of the port community, and that, despite some taxa being

expectedly associated to a given sampling method, the total diversity, even within

specific taxonomic groups, was only recovered by combining different methods.

Interestingly, despite the documented potential of eDNA metabarcoding to capture a

large fraction of the macroorganismal diversity with limited effort (Bista et al., 2017;

O'Donnell et al., 2017), including detection of non-indigenous taxa (Klymus, Marshall,

& Stepien, 2017) and port surveys (Borrell et al., 2017; Grey et al., 2018; Lacoursière-

Roussel et al., 2018), our analyses show that, compared to bulk sample metabarcoding,

eDNA metabarcoding recovered only a subset of the metazoan diversity and did not

provide additional information on targeted groups. This confirms previous findings in

coral reef sites (Djurhuus et al., 2018) and freshwater streams (Macher et al., 2018)

and recently in ports (Koziol et al., 2019), suggesting that despite requiring less

sampling effort, COI and 18S based metabarcoding of eDNA obtained from filtered

water should not be used to replace the need of obtaining bulk samples (using

zooplankton nets, grabs and settlement plates) for PBBS taxa detection. Increasing

evidences showed that the use of universal and degenerated COI primers is leading to

non-specific amplification of prokaryotic and non-target eukaryotic DNA (Collins et al.,

2019). Yet, in this context, at least two alternatives are possible to improve eDNA-

based biodiversity assessments: i) increasing sequencing depth (Grey et al., 2018) or ii)

using group specific markers (Jeunen, Knapp, Spencer, Taylor, et al., 2019). Yet, none

of them ensures full biodiversity recovery and both significantly increase costs. Thus,

the decision between increasing sampling depth and/or using multiple group specific

primers for eDNA and adding multiple substrate sampling needs to be carefully

considered and will be an important area of future research to optimize

metabarcoding-based port monitoring.

Metabarcoding-based port baseline surveys should include spatiotemporal sampling

While it is expected that increased temporal coverage will retrieve more taxa,

the HELCOM/OSPAR protocol, a widely applied protocol developed for PBBS

(http://jointbwmexemptions.org/ballast_water_RA/apex/f?p=104:13), limits sampling

to late summer for sediment and fouling, and to spring and late summer for plankton

in order to reduce costs. Yet, so far, no studies using metabarcoding have been

performed to support this decision. Here we show that for sediment and fouling,

spring sampling provides higher diversity than late summer. Although sampling in late

summer could be appropriate for morphological taxonomy because of the more

abundance of adult individuals, for metabarcoding, sampling in spring is preferable

because during this season i) sizes of organisms are less variable and thus

metabarcoding is less likely to under detect small organisms (Elbrecht, Peinert, &

Leese, 2017) and ii) species diversity is at its maxima due to being a high recruitment

period with abundant organisms at early life stages (Bijleveld et al., 2018). For

zooplankton diversity, our results show that the HELCOM/OSPAR sampling in spring

and late summer produces the highest diversity. However, the data obtained from

filtered water were inconclusive. Previous studies have already shown that seasonal

variations are important considerations for eDNA studies (Lacoursière-Roussel et al.,

2018), but further characterizations are needed over multiple years to design an

adequate protocol for maximizing biodiversity recovery with eDNA.

Concerning spatial sampling, for all sampling methods, OTU replacement

generally contributed more to community variation among sites than nestedness,

suggesting that spatially comprehensive sampling is crucial to recover the port´s

biodiversity. Interestingly, site 4 was not only the most different from all four sites but

was also the one that recorded the largest number of NIS. In coherence with our

findings, it has been observed that brackish environments favors NIS settlement (Zorita

et al., 2013) because these species usually support wider range of salinity (Cardeccia et

al., 2018). Thus, samples for port monitoring should include those with a wide range of

abiotic conditions and covering priority sampling, such as highly active ship berths,

potential reservoirs of newly arrived NIS (Hewitt & Martin, 2001).

Metabarcoding provides valuable information on Non-Indigenous and Cryptogenic

Species (NICS)

Our metabarcoding port biological baseline survey detected NICS previously recorded

in the port of Bilbao and NICS known to be present in the port's marine ecoregion.

Importantly, it also unveiled presence of seven NICS for which no records exist so far in

this ecoregion, highlighting the potential of metabarcoding for early NICS detection.

Nonetheless, not all NICS species previously recorded in Bilbao were found by our

analyses. This might be due to these species not being present in the port at the time

of our survey, to biases of the metabarcoding process such as differential DNA

extraction, primer non-specificity, etc (Xiong, Li, & Zhan, 2016), or to database

incompleteness. Primer bias and/or database incompleteness could also explain why

35 and 42% of NICS were uniquely detected with COI and 18S respectively. Yet, from

the total species included in the AquaNIS database (n=1083), only 460 and 369 are

included in the BOLD and PR2 databases respectively. This stresses the need of

increasing reference databases and of using multiple universal primers for species

detection (Grey et al., 2018). Importantly, our study confirmed that metabarcoding can

detect species occurring at low abundance (Pochon, Bott, Smith, & Wood, 2013) as we

found the non-indigenous amphipod *Melita nitida* with only 18 reads while it took

intensive surveys in 2013, 2014 and 2016 in 3 distinct sampling regions of the Bay of

Biscay to record 76 individuals (Gouillieux, Lavesque, Blanchet, & Bachelet, 2016).

Conclusions

The Ballast Water Management Convention, entered into force in September 2017, aims at preventing the spread of non-indigenous species from one region to another (IMO, 2004). Yet, the implementation of this convention still requires technological developments for assessing compliance and granting exemptions, for which genetic methods have been suggested promising (Rey, Basurko, & Rodríguez-Ezpeleta, 2018). Based on the comparative analysis of 192 samples assessing the use of alternative sampling methods, of sampling at different seasons and at different port locations, we have demonstrated the suitability of metabarcoding for port biodiversity surveys and NIS monitoring and settled the guidelines for future studies. We show that i) combining two pairs of universal primers provides a more holistic view of the port biodiversity, ii) a combination of sampling methods is necessary to recover the different taxonomic groups, iii) environmental DNA cannot replace traditional sampling; vi) sampling should take place in spring and late summer preferably and v) spatial coverage should cover the port's salinity gradient. Considering the cost-effectiveness of metabarcoding with respect to morphological identification, these guidelines and considerations are particularly relevant for performing the risk assessment required for granting exemptions within the International Convention for the Control and Management of Ships' Ballast Water and Sediments.

Acknowledgments

This manuscript is a result of the Aquainvad-ED (AQUAtic INVADers: Early Detection, Control and Management - www.aquainvad-ed.com) project funded by the European

Union (Horizon 2020 research and innovation program under the Marie Sklodowska-

Curie grant agreement no 642197). We thank the Port Authority of Bilbao for

authorizing sampling in the port and Manuel González for making the necessary

arrangements for it. We also thank Iñaki Mendibil and Elisabete Bilbao for laboratory

work, Ekaitz Erauskin, Iker Urtizberea and Lander Larrañaga Juarez for sampling,

Anders Lanzen for providing comments on the manuscript, and Carlota Bermejo,

student of the postgraduate course in Scientific Illustration of the UPV/EHU, for

producing Figure 1. This manuscript is contribution X [to be completed upon

acceptance] from the Marine Research Division of AZTI.

Authors' contributions

AR, OB and NR-E conceived the study and designed methodology; AR and OB collected

the samples; AR preformed laboratory analyses; AR and NR-E analyzed the data; AR

and NR-E wrote the manuscript. All authors contributed critically to the drafts and

gave final approval for publication.

Data Accessibility

Raw demultiplexed MiSeq reads are available at NCBI SRA Bioproject PRJNA515494.

Scripts used for OTU table generation and taxonomic assignment are available at

https://github.com/rodriguez-ezpeleta/metabarcoding-pbbs.

References

Adarraga, I., & Martínez, J. (2011). First record of invasive *Theora lubrica* (Mollusca:

Bivalvia: Semelidae) in the Atlantic Ocean. *Marine Biodiversity Records*, *4*.

doi:10.1017/S1755267211001011

Adarraga, I., & Martínez, J. (2012). First record of the invasive brackish water mytilid

Limnoperna securis (Lamarck, 1819) in the Bay of Biscay. *Aquatic Invasions*,

7(2).

Amaral-Zettler, L. A., McCliment, E. A., Ducklow, H. W., & Huse, S. M. (2009). A Method

for Studying Protistan Diversity Using Massively Parallel Sequencing of V9

Hypervariable Regions of Small-Subunit Ribosomal RNA Genes. *PLOS ONE*, *4*(7),

e6372. doi:10.1371/journal.pone.0006372

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.

AquaNIS. Editorial board. (2015). Information system on Aquatic Non-Indigenous and

Cryptogenic Species. World Wide Web electronic publication.

www.corpi.ku.lt/databases/aquanis. Version 2.36+.

Awad, A., Haag, F., Anil, A. C., & Abdulla, A. (2014). *Guidance on Port Biological*

Baseline Surveys (PBBS). GEF-UNDP-IMO GloBallast Partnerships, IOI, NIO,

IUCN.

Aylagas, E., Mendibil, I., Borja, Á., & Rodríguez-Ezpeleta, N. (2016a). Marine Sediment

Sample Pre-processing for Macroinvertebrates Metabarcoding: Mechanical

Enrichment and Homogenization. *Frontiers in Marine Science*, *3*.

doi:10.3389/fmars.2016.00203

Aylagas, E., Borja, Á., Irigoien, X. & Rodríguez-Ezpeleta, N. (2016b). Benchmarking DNA

Metabarcoding for biodiversity-Based monitoring and Assessment. *Frontiers in*

Marine Science, 3. doi:10.3389/fmars.2016.00096

Aylagas, E., Borja, Á., Muxika, I. & Rodríguez-Ezpeleta, N. (2018). Adapting

metabarcoding-based benthic biomonitoring into routine marine ecological

status assessment networks. *Ecological Indicators, 95 (1),* 194-202

Bax, N., Williamson, A., Aguero, M., Gonzalez, E., & Geeves, W. (2003). Marine invasive

alien species: a threat to global biodiversity. *Marine Policy, 27*(4), 313–323.

doi:10.1016/S0308-597X(03)00041-1

Bijleveld, A. I., Compton, T. J., Klunder, L., Holthuijsen, S., Horn, J. ten, Koolhaas, A., …

Veer, H. W. van der. (2018). Presence-absence of marine macrozoobenthos

does not generally predict abundance and biomass. *Scientific Reports, 8*(1),

3039. doi:10.1038/s41598-018-21285-1

Bishop, M. J., & Hutchings, P. A. (2011). How useful are port surveys focused on target

pest identification for exotic species management? *Marine Pollution Bulletin,*

62(1), 36–42. doi:10.1016/j.marpolbul.2010.09.014

Bista, I., Carvalho, G. R., Walsh, K., Seymour, M., Hajibabaei, M., Lallias, D., … Creer, S.

(2017). Annual time-series analysis of aqueous eDNA reveals ecologically

relevant dynamics of lake ecosystem biodiversity. *Nature Communications, 8,*

14087. doi:10.1038/ncomms14087

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for

Illumina sequence data. *Bioinformatics, 30*(15), 2114–2120.

doi:10.1093/bioinformatics/btu170

Borrell, Y. J., Miralles, L., Huu, H. D., Mohammed-Geba, K., & Garcia-Vazquez, E. (2017).

DNA in a bottle—Rapid metabarcoding survey for early alerts of invasive

species in ports. *PLOS ONE*, *12*(9), e0183347.

doi:10.1371/journal.pone.0183347

Bott, N. J. (2015). Detection of Invasive Species. In S.-K. K. Prof (Ed.), *Springer*

Handbook of Marine Biotechnology (pp. 441–453). Springer Berlin Heidelberg.

doi:10.1007/978-3-642-53971-8_16

Brown, E. A., Chain, F. J. J., Zhan, A., MacIsaac, H. J., & Cristescu, M. E. (2016). Early

detection of aquatic invaders using metabarcoding reveals a high number of

non-indigenous species in Canadian ports. *Diversity and Distributions*, *22*(10),

1045–1059. doi:10.1111/ddi.12465

Butrón, A., Orive, E., & Madariaga, I. (2011). Potential risk of harmful algae transport

by ballast waters: The case of Bilbao Harbour. *Marine Pollution Bulletin*, *62*(4),

747–757.

Cardeccia, A., Marchini, A., Occhipinti-Ambrogi, A., Galil, B., Gollasch, S., Minchin, D., …

Ojaveer, H. (2018). Assessing biological invasions in European Seas: Biological

traits of the most widespread non-indigenous species. *Estuarine, Coastal and*

Shelf Science, *201*, 17–28. doi:10.1016/j.ecss.2016.02.014

Carlton, J. T. (1999). The scale and ecological consequences of biological invasions in

the World's oceans. (pp. 195–212). Presented at the Population and

Community Biology Series Volume 24, Kluwer Academic Publishers.

Collins, R. A., Bakker, J., Wangensteen, O. S., Soto, A. Z., Corrigan, L., Sims, D. W., …

Mariani, S. (2019). Non-specific amplification compromises environmental DNA

metabarcoding with COI. *Methods in Ecology and Evolution*, *10*(11), 1985–2001.

doi:10.1111/2041-210X.13276

Comtet, T., Sandionigi, A., Viard, F., & Casiraghi, M. (2015). DNA (meta) barcoding of

biological invasions: a powerful tool to elucidate invasion processes and help

managing aliens. *Biological Invasions*, *17*(3), 905–922.

Creer, S., Deiner, K., Frey, S., Porazinska, D., Taberlet, P., Thomas, W. K., … Bik, H. M.

(2016). The ecologist's field guide to sequence-based identification of

biodiversity. *Methods in Ecology and Evolution*, *7*(9), 1008–1018.

doi:10.1111/2041-210X.12574

David, M., Gollasch, S., & Pavliha, M. (2013). Global ballast water management and the

"same location" concept: a clear term or a clear issue? *Ecological Applications*,

23(2), 331–338. doi:10.1890/12-0992.1

De Cáceres, M. D., & Legendre, P. (2009). Associations between species and groups of

sites: indices and statistical inference. *Ecology*, *90*(12), 3566–3574.

Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J.-C., & Altermatt, F. (2016).

Environmental DNA reveals that rivers are conveyer belts of biodiversity

information. *Nature Communications*, *7*, 12544. doi:10.1038/ncomms12544

Djurhuus, A., Pitz, K., Sawaya, N. A., Rojas-Márquez, J., Michaud, B., Montes, E., …

Breitbart, M. (2018). Evaluation of marine zooplankton community structure

through environmental DNA metabarcoding: Metabarcoding zooplankton from

eDNA. *Limnology and Oceanography: Methods*, *16*(4), 209–221.

doi:10.1002/lom3.10237

Dray, S., Bauman, D., Blanchet, G., Borcard, D., Clappe, S., Guenard, G., … Wagner, H.

H. (2018). *adespatial: Multivariate Multiscale Spatial Analysis*. Retrieved from

https://CRAN.R-project.org/package=adespatial

Dufrene, M., & Legendre, P. (1997). Species Assemblages and Indicator Species: The

Need for a Flexible Asymmetrical Approach. *Ecological Monographs*, *67*(3), 345.

doi:10.2307/2963459

Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., & Knight, R. (2011). UCHIME

improves sensitivity and speed of chimera detection. *Bioinformatics*, *27*(16),

2194–2200. doi:10.1093/bioinformatics/btr381

Elbrecht, V., Peinert, B., & Leese, F. (2017). Sorting things out: Assessing effects of

unequal specimen biomass on DNA metabarcoding. *Ecology and Evolution*,

7(17), 6918–6926. doi:10.1002/ece3.3192

Elbrecht, V., Vamos, E. E., Meissner, K., Aroviita, J., & Leese, F. (2017). Assessing

strengths and weaknesses of DNA metabarcoding-based macroinvertebrate

identification for routine stream monitoring. *Methods in Ecology and Evolution*,

8(10), 1265–1275. doi:10.1111/2041-210X.12789

Gittenberger, A., Rensing, M., Niemantsverdriet, P., Schrieken, N., & Stegenga, H.

(2014, November). Port of Rotterdam survey and monitoring non-native

species conform HELCOM/OSPAR protocol.

Gouillieux, B., Lavesque, N., Blanchet, H., & Bachelet, G. (2016). First record of the non-

indigenous *Melita nitida Smith*, 1873 (Crustacea: Amphipoda: Melitidae) in the

Bay of Biscay (NE Atlantic). *BioInvasions Records*, *5*(2), 85–92.

Grey, E. K., Bernatchez, L., Cassey, P., Deiner, K., Deveney, M., Howland, K. L., … Lodge,

D. M. (2018). Effects of sampling effort on biodiversity patterns estimated from

626    environmental DNA metabarcoding surveys. *Scientific Reports*, *8*(1).

627    doi:10.1038/s41598-018-27048-2

628 Hebert, P. D., Ratnasingham, S., & de Waard, J. R. (2003). Barcoding animal life:

629    cytochrome c oxidase subunit 1 divergences among closely related species.

630    *Proceedings of the Royal Society of London B: Biological Sciences*, *270*(Suppl 1),

631    S96–S99.

632 HELCOM/OSPAR. (2013). Joint HELCOM/OSPAR Guidelines for the Contracting Parties

633    of OSPAR and HELCOM on the granting of exemptions under International

634    Convention for the Control and Management of Ships' Ballast Water and

635    Sediments, Regulation A-4 (OSPAR Agreement 2013-09), 45 pp.

636 Hewitt, C. L., & Martin, R. B. (2001). Revised protocols for baseline port surveys for

637    introduced marine species–design considerations, sampling protocols and

638    taxonomic sufficiency. *Centre for Research on Introduced Marine Pests*

639    *Technical Report*, (22).

640 IMO. (2004). International Convention for the Control and Management of Ships'

641    Ballast Water and Sediments, 2004. International Maritime Organization, 13

642    Feb 2004, London.

643 Jeunen, G.-J., Knapp, M., Spencer, H. G., Lamare, M. D., Taylor, H. R., Stat, M., …

644    Gemmell, N. J. (2019). Environmental DNA (eDNA) metabarcoding reveals

645    strong discrimination among diverse marine habitats connected by water

646    movement. *Molecular Ecology Resources*, *19*(2), 426–438. doi:10.1111/1755-

647    0998.12982

648 Jeunen, G.-J., Knapp, M., Spencer, H. G., Taylor, H. R., Lamare, M. D., Stat, M., …

649    Gemmell, N. J. (2019). Species-level biodiversity assessment using marine

environmental DNA metabarcoding requires protocol optimization and

standardization. *Ecology and Evolution*, *9*(3), 1323–1335.

doi:10.1002/ece3.4843

Klymus, K. E., Marshall, N. T., & Stepien, C. A. (2017). Environmental DNA (eDNA)

metabarcoding assays to detect invasive invertebrate species in the Great

Lakes. *PLOS ONE*, *12*(5), e0177643. doi:10.1371/journal.pone.0177643

Koziol, A., Stat, M., Simpson, T., Jarman, S., DiBattista, J. D., Harvey, E. S., … Bunce, M.

(2019). Environmental DNA metabarcoding studies are critically affected by

substrate selection. *Molecular Ecology Resources*, *19*(2), 366–376.

doi:10.1111/1755-0998.12971

Kraus, R., Ninčević-Gladan, Ž., Auriemma, R., Bastianini, M., Bolognini, L., Cabrini, M., …

Žuljević, A. (2018). Strategy of port baseline surveys (PBS) in the Adriatic Sea.

Marine Pollution Bulletin. doi:10.1016/j.marpolbul.2018.08.067

Lacoursière-Roussel, A., Howland, K., Normandeau, E., Grey, E. K., Archambault, P.,

Deiner, K., … Bernatchez, L. (2018). eDNA metabarcoding as a new surveillance

approach for coastal Arctic biodiversity. *Ecology and Evolution*.

doi:10.1002/ece3.4213

Legendre, P. & Gallagher, E. (2001). Ecologically meaningful transformations for
ordination of species data. *Oecologia, 129 (2),* 271-280

Lehtiniemi, M., Ojaveer, H., David, M., Galil, B., Gollasch, S., McKenzie, C., … Pederson,

J. (2015). Dose of truth—Monitoring marine non-indigenous species to serve

legislative requirements. *Marine Policy*, *54*, 26–35.

doi:10.1016/j.marpol.2014.12.015

Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., … Machida, R. J.

(2013). A new versatile primer set targeting a short fragment of the

mitochondrial COI region for metabarcoding metazoan diversity: application for

characterizing coral reef fish gut contents. *Frontiers in Zoology*, *10*, 34.

doi:10.1186/1742-9994-10-34

Macher, J.-N., Vivancos, A., Piggott, J. J., Centeno, F. C., Matthaei, C. D., & Leese, F.

(2018). Comparison of environmental DNA and bulk-sample metabarcoding

using highly degenerate cytochrome c oxidase I primers. *Molecular Ecology*

Resources. doi:10.1111/1755-0998.12940

Mahé, F., Rognes, T., Quince, C., de Vargas, C., & Dunthorn, M. (2014). Swarm: robust

and fast clustering method for amplicon-based studies. *PeerJ*, *2*, e593.

doi:10.7717/peerj.593

Mandelik, Y., Roll, U., & Fleischer, A. (2010). Cost-efficiency of biodiversity indicators

for Mediterranean ecosystems and the effects of socio-economic factors.

Journal of Applied Ecology, *47*(6), 1179–1188. doi:10.1111/j.1365-

2664.2010.01864.x

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.Journal*, *17*(1), 10–12. doi:10.14806/ej.17.1.200

Martínez, J., & Adarraga, I. (2006). Programa de vigilancia y control de la introducción

de especies invasoras en los ecosistemas litorales de la costa vasca. 1. *Costa de*

Gipuzkoa, *2*.

McMurdie, P. J. & Holmes, S. (2014). Waste not, Want Not: Why Rarefying Microbiome

Data is Inadmissible. P*los Computational Biology 10(4)*, e1003531.

Meyer, C. P. (2003). Molecular systematics of cowries (Gastropoda: Cypraeidae) and

diversification patterns in the tropics. *Biological Journal of the Linnean Society*,

79(3), 401–459. doi:10.1046/j.1095-8312.2003.00197.x

O'Donnell, J. L., Kelly, R. P., Shelton, A. O., Samhouri, J. F., Lowell, N. C., & Williams, G.

D. (2017). Spatial distribution of environmental DNA in a nearshore marine

habitat. *PeerJ*, *5*, e3044. doi:10.7717/peerj.3044

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R. B., …

Wagner, H. (2013). Package 'vegan'. *Community Ecology Package, Version*, *2*(9).

Pochon, X., Bott, N. J., Smith, K. F., & Wood, S. A. (2013). Evaluating detection limits of

next-generation   sequencing   for   the   surveillance   and   monitoring   of

international marine pests. *PLoS One*, *8*(9), e73935.

Podani, J., & Schmera, D. (2011). A new conceptual and methodological framework for

exploring and explaining pattern in presence - absence data. *Oikos*, *120*(11),

1625–1638. doi:10.1111/j.1600-0706.2011.19451.x

Pyšek, P., Hulme, P. E., Meyerson, L. A., Smith, G. F., Boatwright, J. S., Crouch, N. R., …

Wilson, J. R. U. (2013). Hitting the right target: taxonomic challenges for, and

of, plant invasions. *AoB PLANTS*, *5*. doi:10.1093/aobpla/plt042

Rey, A., Basurko, O. C., & Rodríguez-Ezpeleta, N. (2018). The challenges and promises

of genetic approaches for ballast water management. *Journal of Sea Research*,

133, 134–145. doi:10.1016/j.seares.2017.06.001

Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., …

Weber, C. F. (2009). Introducing mothur: Open-Source, Platform-Independent,

Community-Supported   Software   for   Describing   and   Comparing   Microbial

Communities. *Applied and Environmental Microbiology*, *75*(23), 7537–7541.

doi:10.1128/AEM.01541-09

Seebens, H., Gastner, M. T., & Blasius, B. (2013). The risk of marine bioinvasion caused

by global shipping. *Ecology Letters*, *16*(6), 782–790. doi:10.1111/ele.12111

Shaw, J. L. A., Weyrich, L., & Cooper, A. (2016). Using environmental (e)DNA

sequencing for aquatic biodiversity surveys: a beginner's guide. *Marine and*

Freshwater Research. doi:10.1071/MF15361

Spens, J., Evans, A. R., Halfmaerten, D., Knudsen, S. W., Sengupta, M. E., Mak, S. S. T., …

Hellström, M. (2017). Comparison of capture and storage methods for aqueous

macrobial eDNA using an optimized extraction protocol: advantage of enclosed

filter. *Methods in Ecology and Evolution*, *8*(5), 635–645. doi:10.1111/2041-

210X.12683

Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012). Towards

next-generation biodiversity assessment using DNA metabarcoding. *Molecular*

Ecology, *21*(8), 2045–2050.

Tajadura, J., Bustamante, M., & Salinas, J. I. S. (2016). Temporal and spatial variability

in the distribution of non-indigenous zoobenthic species in the intertidal hard

substrates in the 'Abra of Bilbao' Poster ISOBAY conference 2016

https://www.ehu.eus/documents/5019180/0/Book-of-abstract-ISOBAY15.pdf.

ISOBAY Conference.

Team, Rs. (2015). RStudio: integrated development for R. *RStudio, Inc., Boston, MA*

URL Http://Www. Rstudio. Com, *42*.

Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naive Bayesian classifier for

rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied*

743 *and Environmental Microbiology*, *73*(16), 5261–5267. doi:10.1128/AEM.00062-

744 07

745 Xiong, W., Li, H., & Zhan, A. (2016). Early detection of invasive species in marine

746 ecosystems using high-throughput sequencing: technical challenges and

747 possible solutions. *Marine Biology*, *163*(6), 1–12.

748 Zaiko, A., Schimanski, K., Pochon, X., Hopkins, G. A., Goldstien, S., Floerl, O., & Wood, S.

749 A. (2016). Metabarcoding improves detection of eukaryotes from early

750 biofouling communities: implications for pest monitoring and pathway

751 management. *Biofouling*, *32*(6), 671–684.

752 doi:10.1080/08927014.2016.1186165

753 Zhang, J., Kobert, K., Flouri, T., & Stamatakis, A. (2014). PEAR: a fast and accurate

754 Illumina Paired-End reAd mergeR. *Bioinformatics*, *30*(5), 614–620.

755 doi:10.1093/bioinformatics/btt593

756 Zorita, I., Solaun, O., Borja, A., Franco, J., Muxika, I., & Pascual, M. (2013). Spatial

757 distribution and temporal trends of soft-bottom marine benthic alien species

758 collected during the period 1989–2008 in the Nervión estuary (southeastern

759 Bay of Biscay). *Journal of Sea Research*, *83*, 104–110.

Figures legends

Figure 1. Sampling protocol representing sampling location (Port of Bilbao), sites (1, 2,

3 and 4) and points per site (dots); and illustrating the sampling methods used and the

targeted biological communities.

Figure 2. Overall description of community detected per barcode. A: OTU accumulation

curves per sampling method. B: Venn diagrams of the number and percentage of OTUs

shared between sampling methods. C: Principal Component Analyses of the Hellinger-

transformed abundances of OTUs. Sample scores are displayed in scaling 1 with

ellipses representing the 95% confidence dispersion of each sampling method.

Figure 3. Taxonomic assignment per barcode and per sampling method. A: Percentage

of reads assigned to each taxonomic level. B: Relative abundance of reads classified to

at least Phylum level for the 10 most abundant phyla.

Figure 4. Distribution of indicator OTUs associated to each sampling method at the

order level (in log scale) for COI and 18S. Only orders with at least 10 indicator OTUs

are shown.

Figure 5. Principal Component Analyses of Hellinger-transformed abundances of OTUs

included in the *PBBS targeted taxa dataset* for COI (A-B) and 18S (C-D). A & C: Samples

scores in scaling 1 with ellipses representing the 95% confidence dispersion of each

sampling method. B & D: OTU scores in scaling 1 with the circle of equilibrium

contribution. Only OTUs whose projected length in these two principal component

axes exceeding the value of equilibrium contribution are represented.

Figure 6. Seasonal variation of alpha and beta diversity for each sampling method with

COI and 18S. A: Total OTU richness recovered at each season. B: OTU richness unique

to each season. C: Proportion of the total OTU richness detected with one season. D:

Decomposition of between-season beta diversity into replacement and nestedness

components. E: Proportion of the total OTU richness detected with two seasons.

Figure 7. Spatial variation of alpha and beta diversity for each sampling method with

COI and 18S. A: Total OTU richness recovered at each site. B: OTU richness unique to

each site. C: Proportion of the total OTU richness detected with one site. D:

Decomposition of between-site beta diversity into replacement and nestedness

components

Appendix figure legends

Appendix 1: Picture of the port depicting location of each site sampled. A: Examples of

the different types of structures at each site. B: Fouling communities settled on plates

after 3 months. Picture frames are colored according to the site they were taken at.

Appendix 2: OTU accumulation curves per sample for COI and 18S barcodes.

Appendix 3: Number of filtered water reads assigned to each phylum for three percent

identity ranges when blasting the unclassified reads (*i.e.* reads not classified below the

phylum level) against NCBI. The 5 most abundant phyla per percent identity range are

represented, the remaining being grouped under "other".

Appendix 4: Principal Component Analyses of the Hellinger-transformed abundances

of OTU included in the PBBS targeted metazoan taxa dataset for 18S. A: Samples

scores in scaling 1 with ellipses representing the 95% confidence dispersion of each

sampling method. B: OTU scores in scaling 1 with the circle of equilibrium contribution.

Only OTUs whose projected length in these two principal component axes exceeding

the value of equilibrium contribution are represented.

Appendix 5: Percentage of the total OTU richness detected by each sampling method

for the 10 most diverse metazoan orders of the PBBS targeted taxa dataset for each

barcode.

Appendix 6: Principal Component Analyses (PCA) of the Hellinger-transformed

abundances of OTUs included in the PBBS targeted taxa dataset of Vigo, A Coruña and

Bilbao filtered water samples. Sample scores are displayed in scaling 1. A: PCA

performed with 18S. B: PCA performed with 18S targeting only metazoan. C: PCA

performed with COI.