1

Structural and behavioral analysis reveals that Insomniac impacts sleep by functioning as a

Cul3 adaptor

Qiuling Li, Kayla Y. Y. Lim, and Nicholas Stavropoulos[*]

NYU Neuroscience Institute, Department of Neuroscience and Physiology, New York University

School of Medicine, New York, NY 10016, USA

Keywords: sleep, Cullin-3, ubiquitin, ubiquitin ligase, KCTD2, KCTD5, KCTD17, myoclonic

dystonia

Running Title: Structural and functional dissection of Insomniac

[*] **Corresponding author**

E-mail: nicholas.stavropoulos@nyumc.org

Abstract

The *insomniac* (*inc*) gene is required for normal sleep in *Drosophila* and encodes a

conserved BTB protein that is a putative adaptor for the Cullin-3 (Cul3) ubiquitin ligase. Here

we test whether Inc serves as a Cul3 adaptor by generating mutant forms of Inc and assessing

their biochemical properties and physiological activity in vivo. We show that the N-terminal

BTB domain of Inc is necessary and sufficient for Inc self-association and interactions with

Cul3. Inc point mutations that weaken interactions with Cul3 impair the ability of Inc to rescue

the sleep deficits of *inc* mutants, indicating that Cul3-Inc binding is critical for Inc function in

vivo. Deletions of the conserved Inc C-terminus preserve Inc-Inc and Inc-Cul3 interactions but

abolish Inc activity in vivo, implicating the Inc C-terminus as an effector domain that recruits Inc

substrates. Mutation of a conserved C-terminal arginine similarly abolishes Inc function,

suggesting that this residue is vital for the recruitment or ubiquitination of Inc targets. Mutation

of the same residue in the human Inc ortholog KCTD17 is associated with myoclonic dystonia,

indicating its functional importance in Inc family members. Finally, we show that Inc assembles

into multimeric Cul3-Inc complexes in vivo and that depleting Cul3 causes accumulation of Inc,

suggesting that Inc is negatively regulated by Cul3-dependent autocatalytic ubiquitination, a

hallmark of Cullin adaptors. Our findings implicate Inc as a Cul3 adaptor and provide tools to

identify the targets of Inc family proteins that impact sleep and neurological disorders.

Introduction

Sleep is a conserved animal behavior regulated by genetic and molecular mechanisms that remain elusive. Elucidating these mechanisms is a longstanding goal in biology, given that the purpose of sleep is still not well understood and because alterations in these mechanisms may cause sleep disturbances, including those associated with neurodegenerative and neurodevelopmental disorders. Various genes that strongly influence the duration and characteristics of sleep have been identified by unbiased genetic screens in flies (Afonso et al., 2015; Cirelli et al., 2005; Koh et al., 2008; Pfeiffenberger and Allada, 2012; Rogulja and Young, 2012; Stavropoulos and Young, 2011; Toda et al., 2019), zebrafish (Chiu et al., 2016; Singh et al., 2017), worms (Iannacone et al., 2017), and mice (Funato et al., 2016). While many of these genes function in pathways governing neuronal excitability or neurotransmission (Chiu et al., 2016; Cirelli et al., 2005; Koh et al., 2008; Singh et al., 2017; M. Wu et al., 2014; M. N. Wu et al., 2010), others influence sleep by mechanisms that remain poorly understood.

insomniac (*inc*) mutations strongly curtail the duration and consolidation of sleep but do not alter its circadian regulation (Pfeiffenberger and Allada, 2012; Stavropoulos and Young, 2011). *inc* activity is required in neurons for normal sleep, and conversely, restoring *inc* solely to neurons is largely sufficient to rescue the sleep deficits of *inc* mutants (Pfeiffenberger and Allada, 2012; Stavropoulos and Young, 2011). While these findings indicate that *inc* influences sleep primarily through neurons, the molecular mechanisms by which *inc* functions remain poorly defined. *inc* encodes a conserved protein of the Bric-a-brac, Tramtrack, Broad Complex (BTB) domain superfamily (Stavropoulos and Young, 2011), which comprises more than 80 proteins in *Drosophila* (Stogios et al., 2005). The BTB domain mediates protein-protein interactions including homomeric self-associations and binding to heterologous proteins (Stogios

et al., 2005). BTB-domain containing proteins cluster into distinct subfamilies based on sequence

variation within the BTB domain and the presence of additional domains that underlie diverse

cellular functions (Stogios et al., 2005). BTB proteins include transcriptional regulators (DiBello

et al., 1991; Godt et al., 1993; Harrison and Travers, 1990), ion channels (Butler et al., 1989;

Papazian et al., 1987; Pongs et al., 1988; Wei et al., 1990), auxiliary subunits of $GABA_B$

receptors (Schwenk et al., 2010), and adaptors for the Cullin-3 (Cul3) ubiquitin ligase (Furukawa

et al., 2003; Pintard et al., 2003; Xu et al., 2003).

Indirect evidence suggests that Inc may serve as a substrate adaptor for the Cul3 ubiquitin

ligase complex (Li et al., 2017; Pfeiffenberger and Allada, 2012; Stavropoulos and Young,

2011). BTB proteins that function as Cul3 adaptors form homomultimers that bind Cul3 and

recruit substrates to Cul3 complexes for ubiquitination (Furukawa et al., 2003; Geyer et al.,

2003; Pintard et al., 2003; Xu et al., 2003). In cultured cells, Inc is able to bind Cul3

(Pfeiffenberger and Allada, 2012; Stavropoulos and Young, 2011) and to self-associate (Li et al.,

2017; Pfeiffenberger and Allada, 2012), but whether these molecular interactions occur in vivo

or are necessary for the physiological activity of Inc is unknown. Neuronal depletion of either *inc*

or *Cul3* causes short sleep (Pfeiffenberger and Allada, 2012; Stavropoulos and Young, 2011),

consistent with the notion that *Cul3* and *inc* influence sleep through a common pathway, yet the

modular nature of Cul3 complexes leaves unclear whether Cul3 impacts sleep through Inc. Cul3

assembles with tens of different BTB adaptors to ubiquitinate hundreds of substrates (Emanuele

et al., 2011), and may impinge upon sleep through multiple adaptor and substrate pathways. In

the absence of functional evidence linking Inc and Cul3 in a concerted mechanism that

influences sleep, the role of Inc as Cul3 adaptor remains speculative.

Here, we have generated mutant forms of Inc to test the hypothesis that Inc impacts sleep

by functioning as a Cul3 adaptor. Analysis of these Inc mutants in cultured cells and in vivo

reveals that Inc has the biochemical properties and molecular interactions expected of a Cul3

adaptor. The N-terminal BTB domain of Inc mediates interactions with Cul3 and homomeric

self-associations. Weakening Cul3-Inc associations impairs Inc activity in vivo, indicating that

Inc function requires the assembly of Cul3-Inc complexes. Deletion of the conserved C-terminal

domain of Inc abolishes Inc function in vivo, as does a missense mutation of a conserved,

disease-associated arginine residue in the C-terminal domain. These findings identify the Inc C-

terminus as a putative substrate binding domain and define a region within the C-terminus likely

to bind substrates. Mutation of the same residue in KCTD17, a human ortholog of Inc, is

associated with myoclonic dystonia (Mencacci et al., 2015), indicating its functional importance

across Inc family proteins. Our findings reveal that Inc influences sleep by functioning as an

adaptor for the Cul3 ubiquitin ligase complex. More broadly, our studies provide structural and

functional insights into Inc family proteins and tools to identify their substrates that impact

neuronal function and behavior.

Results

The Inc BTB domain mediates Cul3 binding and Inc homomultimerization

Inc and its orthologs have two conserved domains, an N-terminal BTB domain and a C-terminal domain unique to Inc family members (Figures 1A and S1A) (Dementieva et al., 2009; Stavropoulos and Young, 2011). The BTB domains of Cul3 adaptors mediate Cul3 binding and adaptor multimerization, while distal domains recruit substrates (Furukawa et al., 2003; Geyer et al., 2003; Pintard et al., 2003; Xu et al., 2003). To test whether Inc domains have the biochemical properties expected of a Cul3 adaptor, we generated a deletion series in which Inc is truncated progressively from its N- or C-terminus (Figure 1A). We first assessed the stability of these Inc derivatives and their associations with Cul3 by expressing epitope-tagged forms of these proteins in S2 cells and performing co-immunoprecipitations. Deleting residues preceding the BTB domain (Inc^{22-211}) did not significantly alter interactions with Cul3 or the stability of Inc (Figure 1B). In contrast, removing part of the BTB domain (Inc^{31-211}) or deleting the BTB domain entirely ($Inc^{124-211}$) abolished Cul3 binding and resulted in a destabilized Inc protein (Figure 1B). In contrast, C-terminal truncations of Inc did not alter Cul3 binding or Inc stability. Inc proteins lacking 25 C-terminal residues (Inc^{1-186}), most of the C-terminal domain (Inc^{1-156}), or all residues following the BTB domain (Inc^{1-123}) associated with Cul3 similarly to full-length Inc (Figure 1B), indicating that the Inc C-terminus is dispensable for Inc-Cul3 interactions. The isolated Inc BTB domain (Inc^{22-123}) was sufficient for Inc-Cul3 interactions, although these associations were weaker than those of Inc^{1-123} (Figure 1B), suggesting that Inc N-terminal residues contribute to Cul3 binding. Alternatively, weaker binding of Inc^{22-123} to Cul3 might reflect an inhibitory effect of fusing the 3×FLAG tag to the isolated BTB domain. Together, these results indicate that the Inc BTB domain is necessary and sufficient for Cul3 binding, a key property expected of a Cul3 adaptor.

Next, we assessed the ability of truncated Inc proteins to associate with full length Inc.

Partial deletion of the Inc BTB domain (Inc[31-211]) or its complete removal (Inc[124-211]) abolished

Inc-Inc interactions (Figure 1C). The reduction in Inc stability caused by truncating the BTB

domain (Figures 1B and 1C) suggests that Inc is an obligate homomultimer and that Inc monomers

are intrinsically unstable. In contrast, C-terminal truncations of Inc did not detectably alter Inc-Inc

associations (Figure 1C), indicating that the Inc C-terminus is dispensable for Inc multimerization.

The Inc BTB domain (Inc[22-123]) was sufficient to bind Inc (Figure 1C), albeit more weakly than

Inc[1-123], suggesting that N-terminal Inc residues contribute to Inc-Inc interactions or that

interactions of the isolated BTB domain are occluded by the 3×FLAG tag. We conclude that the

Inc BTB domain mediates Inc self-association and Cul3 binding and that the Inc C-terminus is

dispensable for these interactions.

Identification of Inc point mutants that selectively weaken Inc-Cul3 and Inc-Inc interactions

We next sought to identify Inc point mutants that selectively perturb Inc-Cul3 and Inc-Inc

interactions, in order to test the necessity of these interactions for Inc activity in vivo. Our efforts

to mutagenize Inc were informed by the crystal structure of human KCTD5, an Inc ortholog that

forms homopentamers (Dementieva et al., 2009), and by the structures of Cul3 complexed with

BTB-MATH and BTB-BACK-Kelch adaptors (Canning et al., 2013; Errington et al., 2012; Ji and

Privé, 2013), in which adaptors form homodimers and assemble with Cul3 in a 2:2 stoichiometry

(Canning et al., 2013; Errington et al., 2012; Ji and Privé, 2013). While Inc and its orthologs are

divergent from BTB-MATH and BTB-BACK-Kelch adaptors and likely bind Cul3 with a different

mechanism and stoichiometry (Balasco et al., 2014; Ji et al., 2015), comparison of these structures

suggested a region of Inc that may bind Cul3. We selected eight residues in the Inc BTB domain

that are conserved between Inc and its three human orthologs and that reside on the solvent-

accessible surface of KCTD5 (Figures S1A and S1B), reasoning that these residues may contribute

to Cul3 binding. We mutated these residues individually to alanine (F47A, D57A, D61A, F105A,

Y106A, N107A) or to oppositely charged residues (R50E, E104K), and assessed the consequences

for Inc-Inc and Inc-Cul3 interactions in cultured S2 cells. Most mutants did not detectably alter

Inc self-association or Cul3 binding (Table 1). Two mutants, F47A and F105A, weakened

interactions with Cul3 but did not significantly affect Inc self-association or stability (Figures 2A

and 2B). A double mutant combining F47A and F105A behaved similarly and selectively impaired

Cul3 binding (Figures 2A and 2B). These phenylalanine residues cluster in a hydrophobic patch

on the surface of KCTD5 that may be buried upon Cul3 binding (Figure S1B). This patch is distinct

from the interface of adjacent KCTD5 subunits, consistent with the lack of a measurable impact

of mutating F47 and F105 on Inc-Inc interactions (Figure 2B).

To identify Inc mutants that impair Inc-Inc homomultimerization, we mutated seven

conserved residues that form a network of polar and charged interactions between adjacent KCTD5

subunits (Dementieva et al., 2009) (Figures S1A and S1C). We mutated these residues to alanine

(T36A, D71A, D73A, N82A) or to oppositely charged residues (R85E, K88D, E101K) in single

and double mutant combinations. None of these single or double mutants significantly altered Inc

self-associations in S2 cells (Table 2). We reasoned that Inc may form homopentamers, like

KCTD5, and that cooperative interactions between Inc subunits might limit the effects of these

mutations. We therefore generated a triple point mutant predicted to disrupt interactions of Inc

subunits with both flanking neighbors in a putative Inc pentamer (T36A, D71A, R85E) (Figure

S1C). This triple mutant exhibited significantly weakened interactions with both Inc and Cul3 and

a markedly reduced stability (Figures 2C and 2D), resembling the consequences of deleting the

Inc BTB domain (Figures 1B and 1C). These findings support the conclusion that multimerization-

deficient Inc proteins are unstable and suggest that Inc multimerization is required for efficient

Cul3 binding. These results furthermore suggest that Cul3 binds cooperatively to adjacent Inc

subunits and that Cul3-Inc complexes have an architecture distinct from Cul3-adaptor complexes

formed by BTB-MATH and BTB-BACK-Kelch proteins (Canning et al., 2013; Errington et al.,

2012; Ji and Privé, 2013).

The assembly of Inc-Cul3 complexes in vivo is required for the function of Inc

Inc mutations that selectively impair Cul3 binding represent key tools to assess whether

Inc impacts sleep by functioning as a Cul3 adaptor. To assess the physiological activity of Inc

mutants in vivo, we generated UAS transgenes expressing 3×FLAG-Inc or -Inc point mutants that

weaken Cul3 interactions: Inc^{F47A}, Inc^{F105A}, and $Inc^{F47A/F105A}$. We integrated these transgenes at the

same genomic site (*attP2*) and backcrossed them to generate an isogenic allelic series, enabling

careful comparisons of their activity. We first assessed the expression of these proteins using *inc-*

Gal4, a driver that fully rescues the sleep deficits of *inc* mutants when used to restore *inc*

expression (Li et al., 2017; Stavropoulos and Young, 2011) and which therefore recapitulates *inc*

expression in cells relevant for sleep. Inc and Inc point mutants were expressed at similar levels

under *inc-Gal4* control in *inc* mutants (Figure 3A) or inc^+ animals (Figure S2A), indicating that

these point mutants do not alter Inc stability in vivo, recapitulating findings from cultured cells

(Figures 2A and 2B). Next, we assessed the impact of expressing Inc point mutants on sleep-wake

behavior. Because adaptor proteins deficient for Cul3 interactions might sequester substrates and

thus function as dominant negatives, we first tested whether the sleep of inc^+ animals was altered

by the expression of Inc point mutants. Animals expressing Inc point mutants under the pan-

neuronal *elav^{c155}-Gal4* control slept indistinguishably from those expressing wild-type Inc and

from control animals lacking *UAS* transgenes (Figure S2B). Thus, expression of Inc point mutants

in neurons does not elicit dominant negative effects or antagonize endogenous Inc function.

We next tested whether Inc-Cul3 interactions are required for Inc function in vivo. If so,

attenuating these interactions should impair the ability of Inc to rescue the sleep phenotypes of *inc*

mutants. While Inc expressed with *inc-Gal4* fully rescued the short sleep of *inc^1* null mutants, Inc

mutants that weaken Inc-Cul3 interactions provided only a partial rescue (Figures 3B and 3C),

supporting the interpretation that Inc activity in vivo requires binding to the Cul3 ubiquitin ligase

complex. Rescue was more strongly impaired for Inc^{F105A} and Inc$^{F47A/F105A}$ than Inc^{F47A} (Figures

3B and S3A-D). Because these Inc mutants are expressed at similar levels in vivo (Figure 3A), we

speculate that the F47A mutation might impair Cul3 binding in vivo to a lesser degree than in S2

cells. Collectively, the impaired rescuing activity of this trio of Inc mutants suggests that Inc

activity requires the assembly of Cul3-Inc complexes in vivo and that disruption of these

complexes inhibits sleep.

The conserved C-terminal domain of Inc is essential for Inc activity in vivo

While the Inc C-terminus is dispensable for Inc-Inc and Inc-Cul3 associations (Figures 1B

and 1C), its evolutionary conservation suggests an essential function. If Inc serves as a Cul3

adaptor, its C-terminus is predicted to bind substrates and recruit them to the Cul3 complex for

ubiquitination. To test whether the C-terminus is required for physiological function of Inc in vivo,

we generated *UAS* transgenes expressing 3×FLAG-tagged Inc C-terminal truncations: Inc$^{1-186}$,

Inc$^{1-156}$, and Inc$^{1-123}$. These proteins were expressed at similar levels under *inc-Gal4* control in *inc*

mutants (Figure 4A), indicating that these proteins are stable in vivo, as in cultured cells (Figures

1B and 1C). Neuronal expression of Inc C-terminal truncations using *elav^{c155}-Gal4* did not

significantly alter sleep, indicating that these proteins do not elicit dominant negative effects or

antagonize endogenous *inc* function (Figure S4).

To determine whether the Inc C-terminus is essential for Inc function, we tested whether

C-terminally truncated Inc proteins rescue the sleep deficits of *inc* mutants. Wild-type Inc

expressed under *inc-Gal4* control fully rescued the sleep deficits of *inc^1* mutants (Figures 4B and

4C). In contrast, animals expressing Inc$^{1-156}$ or Inc$^{1-123}$ slept indistinguishably from *inc* mutants,

indicating that these proteins failed to rescue the *inc* phenotype (Figures 4B and 4C). Expression

of Inc$^{1-186}$ partially rescued the sleep phenotypes of *inc* mutants as assessed by several sleep

parameters, including total sleep duration (Figure 4B), nighttime and daytime sleep (Figures S5A

and S5B), and sleep bout duration and number (Figures S5C and S5D). These findings indicate

that the Inc C-terminus is essential for Inc activity in vivo and that removing the terminal 25

residues of Inc curtails Inc function. The stability of C-terminally truncated Inc proteins and their

ability to engage normally in Inc-Cul3 and Inc-Inc interactions (Figures 1B and 1C), supports the

hypothesis that the Inc C-terminus recruits Inc binding partners including substrates whose

ubiquitination influences sleep.

Mutation of a conserved disease-associated arginine in the Inc C-terminus abolishes Inc

function and defines a region of Inc likely to recruit substrates

To further dissect the function of the Inc C-terminus, we mutated several conserved

residues whose attributes suggested they might contribute to Inc function. A missense mutation of

a conserved C-terminal arginine in the human Inc ortholog KCTD17 (R145H) is associated with

a dominant form of myoclonic dystonia, a neurological movement disorder (Mencacci et al., 2015).

We reasoned that this arginine residue might be critical for the function of Inc and its orthologs.

In particular, the analogous residue in KCTD5 (R159) resides on a lateral solvent-accessible

surface of the C-terminal domain that might bind and orient substrates for Cul3-dependent

ubiquitination (Figure S1D). We therefore generated the analogous mutation in Inc (Inc^{R135H}) to

assess whether it impacts Inc function. We also mutated three conserved Inc residues (N167, Y168,

G169) to alanine (Inc^{AAA}), reasoning that their location within a surface-exposed loop on the

bottom surface of the KCTD5 C-terminus and their conservation amid flanking non-conserved

residues (Figures S1A and S1D) might reflect an important function. In cultured S2 cells, both

Inc^{R135H} and Inc^{AAA} were stably expressed and associated with Cul3 and Inc indistinguishably from

wild-type Inc, indicating that these mutations do not alter Inc multimerization or Cul3 binding

(Figure 5).

To determine whether Inc^{R135H} and Inc^{AAA} alter the physiological activity of Inc, we

generated UAS transgenes to express these mutants in vivo. As in cultured cells, Inc^{R135H} and

Inc^{AAA} were stably expressed in vivo (Figure 6A). Expression of Inc^{R135H} and Inc^{AAA} with *elav^{c155}-

Gal4* did not alter sleep in *inc$^+$* animals, indicating that these proteins do not have dominant

negative activity or inhibit endogenous Inc function (Figure S6). Next, we tested whether Inc^{R135H}

and Inc^{AAA} were able to rescue the sleep deficits of *inc* mutants. Inc^{R135H} was unable to restore

sleep to *inc^1* mutants, indicating that the R135H mutation abolishes Inc function (Figures 6B-C

and S7A-D). In contrast, Inc^{AAA} completely rescued *inc^1* mutants and behaved indistinguishably

from wild-type Inc (Figures 6B-C and S7A-D). These data indicate that R135 is critical for Inc

function and suggest that mutation of this residue impairs the recruitment of Inc binding partners

including substrates. The analogous mutation in KCTD17 may similarly alter its ability to engage

targets, suggesting that deficient substrate ubiquitination is a cause of KCTD17-associated

myoclonic dystonia.

Inc exhibits properties of a Cul3 adaptor in neurons in vivo

Neuronal depletion of *inc* or *Cul3* shortens sleep, indicating that the activity of both genes

is required in neurons for normal sleep (Pfeiffenberger and Allada, 2012; Stavropoulos and Young,

2011). We performed a series of experiments to determine whether Inc exhibits the properties of

a Cul3 adaptor in vivo. First, we tested whether Cul3-Inc complexes assemble in vivo, by co-

expressing epitope-tagged Inc and Cul3 in neurons using the *elav^{c155}-Gal4* driver and performing

co-immunoprecipitations. We observed that Cul3 and Inc associate in neurons in vivo (Figure 7A).

Second, we tested whether Inc homomultimerizes in vivo, by co-expressing HA-Inc and Myc-Inc

in neurons and assessing their physical interactions. We observed strong self-association of Inc in

neurons (Figure 7B), indicating that Inc forms homomultimeric complexes in vivo, as in cultured

cells (Figure 1C) (Li et al., 2017; Pfeiffenberger and Allada, 2012). Third, we tested whether Inc

abundance is regulated by Cul3 in vivo. Cul3 adaptors are often regulated by autocatalytic

ubiquitination and degradation in Cul3 complexes (Djagaeva and Doronkin, 2009; Geyer et al.,

2003; Hudson and Cooley, 2010; Pintard et al., 2003; Wee et al., 2005; Zhang et al., 2005). To test

whether Inc abundance is regulated by Cul3 activity, we expressed neuronal RNAi against *Cul3*

and assessed Inc levels in head lysates. We observed an increase in Inc protein levels upon

depletion of Cul3 (Figure 7C), indicating that Cul3 negatively regulates Inc in adult neurons. qRT-

PCR revealed that neuronal Cul3 RNAi did not increase *inc* transcript levels, indicating that

elevated Inc levels arise by a post-transcriptional mechanism (Figure 7D). We conclude that Inc is

regulated endogenously in neurons by Cul3-dependent autocatalytic ubiquitination and subsequent

degradation. Together, these results indicate that Inc binds Cul3 in vivo and has the characteristics

expected of a Cul3 adaptor in neurons, a cell type through which *inc* influences sleep

(Pfeiffenberger and Allada, 2012; Stavropoulos and Young, 2011).

Discussion

The neuronal activity of Insomniac is vital for sleep, yet the molecular mechanism

underlying its function has remained speculative. Our analysis implicates Inc as a Cul3 adaptor in

vivo and provides insight into the mechanism by which Inc binds Cul3 and recruits substrates to

ultimately impact behavior. Inc has a modular organization and conserved domains that fulfill the

properties expected of a Cul3 adaptor (Furukawa et al., 2003; Geyer et al., 2003; Pintard et al.,

2003; Xu et al., 2003). The Inc BTB domain mediates the homomeric assembly of Inc and

interactions with Cul3, while the Inc C-terminus is dispensable for these interactions but is vital

for Inc function in vivo, consistent with its putative function in recruiting substrates to Cul3-Inc

complexes for ubiquitination. Inc assembles into Cul3-Inc complexes in neurons and importantly,

reducing the affinity of Inc-Cul3 associations impairs the activity of Inc in vivo. These findings

link the biochemical associations of Cul3 and Inc to prior findings that normal sleep-wake cycles

require the activity of both proteins in neurons (Pfeiffenberger and Allada, 2012; Stavropoulos and

Young, 2011). The negative regulation of Inc by Cul3 in neurons is characteristic of autocatalytic

degradation exhibited by Cul3 adaptors (Djagaeva and Doronkin, 2009; Geyer et al., 2003; Hudson

and Cooley, 2010; Pintard et al., 2003; Wee et al., 2005; Zhang et al., 2005), providing further

support for the function of Inc as a Cul3 adaptor in cells through which Inc impacts sleep.

Our data and prior studies (Balasco et al., 2014; Dementieva et al., 2009; Ji et al., 2015)

suggest that Inc forms an obligate homopentamer in which neighboring Inc subunits interact

cooperatively. Consistent with a pentameric structure for Inc, a triple mutation altering opposite

sides of Inc and thus its interactions with both neighboring subunits severely compromised Inc

multimerization. Inc multimerization is furthermore required for efficient Cul3 binding, suggesting

that Cul3 binds cooperatively to the BTB domains of neighboring Inc subunits. Such a mechanism

is consistent with the consequences of mutating conserved phenylalanine residues (F47 and F105)

near this interface, which impair Cul3 binding and Inc function in vivo. It is perhaps surprising

that Inc mutants that impair Cul3 interactions do not elicit dominant negative phenotypes when

overexpressed, although we note that these mutants retain some ability to bind Cul3. Similarly, Inc

overexpression might be expected to sequester substrates but does not alter sleep (Li et al., 2017;

Stavropoulos and Young, 2011). We speculate that robust Inc homomultimerization may buffer

Cul3-Inc complexes against perturbations in Inc abundance and contribute, along with additional

mechanisms that regulate Inc activity, to the permissive influence that Inc exerts on sleep.

Our functional analysis of the Inc C-terminus provides insight into the mechanism by

which Inc is likely to bind substrates and orient them for ubiquitination in Cul3 complexes. The

Inc C-terminus is dispensable for Cul3 binding and Inc multimerization, yet is absolutely essential

for Inc function in vivo, implicating it as a substrate recruitment domain. We furthermore

identified a conserved, surface-exposed arginine residue (R135) within the Inc C-terminus

required for Inc activity in vivo. Mutation of this residue is behaviorally indistinguishable from

deleting the Inc C-terminus and does not alter Inc stability, suggesting that it does not disrupt the

structure of the C-terminal domain. While our results do not exclude the possibility that this residue

is essential for Inc trafficking or other mechanisms that might regulate Inc activity, the simplest

possibility suggested by its position on the lateral surface of the Inc C-terminus is that it contributes

directly to binding targets that impact sleep. A missense mutation of the same residue in the human

Inc ortholog KCTD17 (R145H) is associated with myoclonic dystonia (Mencacci et al., 2015),

suggesting a similar functional importance for substrate recruitment in Inc orthologs. The partial

functional impairment of Inc caused by removing C-terminal residues (Inc$^{1-186}$) is consistent with

the possibility that they contribute to substrate binding. These residues are not resolved in the

KCTD5 crystal structure (Dementieva et al., 2009), suggesting that they are flexible in the absence

of substrates and may become structured upon substrate binding. While a parsimonious model is

that Inc directly binds substrates, our findings do not rule out more complex models of substrate

recruitment that require co-adaptor proteins, analogous to the mechanism recently described for

KLHL12, a BTB-Kelch family adaptor (McGourty et al., 2016). Further studies are required to

elucidate Inc substrates and distinguish among these models.

Our recent findings revealed that orthologs of Inc can substitute for Inc in flies and

restore sleep to *inc* mutants, suggesting that the functions and targets of Inc and its orthologs are

evolutionarily conserved (Li et al., 2017). Our present studies provide evidence for the function

of Inc as a Cul3 adaptor in vivo and the foundation for identifying Inc substrates. While

substrates for Inc orthologs have been recently described in cultured cells (Brockmann et al.,

2017; Kasahara et al., 2014; Kim et al., 2017), whether these targets or other proteins mediate the

impact of Cul3 and Inc on sleep remains unknown. Elucidating Inc substrates and the

downstream pathways will advance our understanding of how protein ubiquitination pathways

contribute to the regulation of nervous system function and mechanisms underlying behavior.

Acknowledgements

We thank Zachary Zuchowski for assistance in constructing plasmids. This work was

supported by an International Student Research Fellowship from the Howard Hughes Medical

Institute (HHMI) to Q.L., and by grants from the National Institutes of Health (NS111304), the

Mathers Foundation, Whitehall Foundation grant 2013-05-78, fellowships from the Alfred P.

Sloan and Leon Levy Foundations, a NARSAD Young Investigator Award from the Brain and

Behavior Foundation, the J. Christian Gillin, M.D. Research Award from the Sleep Research

Society Foundation, and a Career Scientist Award from the Irma T. Hirschl/Weill-Caulier Trust

to N.S.

Methods

Plasmids and molecular cloning

Vectors for expression in S2 cells were as follows:

pAc5.1–Inc-HA (pNS277) encodes Inc fused to a C-terminal 1×HA epitope

(GSYPYDVPDYA) and was generated by ligating EcoRI-XhoI digested pAc5.1-v5-HisA

backbone and an EcoRI-XhoI Inc-HA fragment from pNS273 (as described below).

pAc5.1–3×Myc-Inc (pNS351) encodes an N-terminal 3×Myc epitope

(MEQKLISEEDLGSEQKLISEEDLGSEQKLISEEDLAS) fused to Inc as previously described

(Stavropoulos and Young, 2011).

pAc5.1–3×Myc-Inc$^{22-211}$ (pNS370) was generated by ligating NheI-XhoI digested

pNS309 (Stavropoulos and Young, 2011) and the NheI-XhoI fragment liberated from the PCR

amplification product of pNS351 template and primers oNS285 and oNS315.

pAc5.1–3×Myc-Inc$^{31-211}$ (pNS371) was generated similarly to pNS370, substituting

primers oNS285 and oNS316.

pAc5.1–3×Myc-Inc$^{124-211}$ (pNS372) was generated similarly to pNS370, substituting

primers oNS285 and oNS317.

pAc5.1–3×Myc-Inc$^{1-123}$ (pNS374) was generated similarly to pNS370, substituting

primers oNS277 and oNS319.

pAc5.1–3×Myc-Inc$^{1-156}$ (pNS375) was generated similarly to pNS370, substituting

primers oNS277 and oNS320.

pAc5.1–3×Myc-Inc$^{1-186}$ (pNS376) was generated similarly to pNS370, substituting

primers oNS277 and oNS321.

pAc5.1–3×HA-Inc (pNS402) encodes an N-terminal 3×HA epitope

(MYPYDVPDYAGSYPYDVPDYAGSYPYDVPDYAAS) fused to Inc and was generated by

ligating NheI-XhoI digested pNS310 (Stavropoulos and Young, 2011) and a NheI-XhoI *inc*

fragment liberated from pNS351.

pAc5.1–3×FLAG-Cul3 (pNS403) encodes an N-terminal 3×FLAG epitope

(MDYKDDDDKGSDYKDDDDKGSDYKDDDDKAS) fused to *Drosophila* Cul3 and was

generated by ligating NheI-NotI digested pNS311 and a NheI-NotI Cul3 fragment liberated from

pNS314 (Stavropoulos and Young, 2011). pNS311 contains an N-terminal 3×FLAG tag and was

generated from pNS298, a derivative of pAc5.1/V5-HisA that contains a C-terminal 3×FLAG

tag. To construct pNS298, oligonucleotides oNS234 and oNS235 were phosphorylated,

annealed, and cloned into XhoI-XbaI digested pAc5.1/V5-HisA. To construct pNS311, EcoRI-

NotI digested pAc5.1/V5-HisA was ligated to the EcoRI-NotI fragment liberated from the PCR

amplification product of pNS298 template and primers oNS240 and oNS241.

pAc5.1–3×FLAG-Inc (pNS408) encodes an N-terminal 3×FLAG epitope

(MDYKDDDDKGSDYKDDDDKGSDYKDDDDKAS) fused to Inc and was generated by

ligating NheI-NotI digested pNS311 and a NheI-XhoI *inc* fragment liberated from pNS351.

pAc5.1–3×HA-Inc^{F47A} (pNS409) encodes an N-terminal 3×HA epitope

(MYPYDVPDYAGSYPYDVPDYAGSYPYDVPDYAAS) fused to Inc^{F47A} and was generated

by ligating NheI-XhoI digested pNS310 and an NheI-XhoI Inc^{F47A} fragment. The NheI-XhoI

Inc^{F47A} fragment was liberated from the fusion PCR amplification product of primers oNS683

and oNS684 and an equimolar mix of overlapping 5' and 3' Inc^{F47A} fragments as template. The

5' and 3' fragments were generated by PCR amplification of pNS408 template with primers

oNS683/oNS695 and oNS684/oNS694, respectively.

pAc5.1–3×HA-Inc^{R50E} (pNS410) was generated similarly to pNS409, substituting

primers oNS683/oNS697 and oNS684/oNS696 for generating the 5' and 3' Inc^{R50E} fragments,

respectively.

pAc5.1–3×HA-Inc^{D57A} (pNS411) was generated similarly to pNS409, substituting

primers oNS683/oNS699 and oNS684/oNS698 for generating the 5' and 3' Inc^{D57A} fragments,

respectively.

pAc5.1–3×HA-Inc^{D61A} (pNS412) was generated similarly to pNS409, substituting

primers oNS683/oNS701 and oNS684/oNS700 for generating the 5' and 3' Inc^{D61A} fragments.,

respectively.

pAc5.1–3×HA-Inc^{E104K} (pNS413) was generated similarly to pNS409, substituting

primers oNS683/oNS703 and oNS684/oNS702 for generating the 5' and 3' Inc^{E104K} fragments,

respectively.

pAc5.1–3×HA-Inc^{D73A} (pNS430) was generated similarly to pNS409, substituting

primers oNS683/oNS686 and oNS684/oNS685 for generating the 5' and 3' Inc^{D73A} fragments,

respectively.

pAc5.1–3×HA-Inc^{N82A} (pNS431) was generated similarly to pNS409, substituting

primers oNS683/oNS686 and oNS684/oNS685 for generating the 5' and 3' Inc^{N82A} fragments,

respectively.

pAc5.1–3×HA-Inc^{K88D} (pNS432) was generated similarly to pNS409, substituting

primers oNS683/oNS688 and oNS684/oNS687 for generating the 5' and 3' Inc^{K88D} fragments,

respectively.

pAc5.1–3×HA-Inc^{E101K} (pNS433) was generated similarly to pNS409, substituting

primers oNS683/oNS688 and oNS684/oNS687 for generating the 5' and 3' Inc^{E101K} fragments,

respectively.

pAc5.1–3×HA-Inc^{T36A} (pNS434) was generated similarly to pNS409, substituting

primers oNS683/oNS693 and oNS684/oNS689 for generating the 5' and 3' Inc^{T36A} fragments,

respectively.

pAc5.1–3×HA-Inc^{D71A} (pNS435) was generated similarly to pNS409, substituting

primers oNS683/oNS693 and oNS684/oNS689 for generating the 5' and 3' Inc^{D71A} fragments,

respectively.

pAc5.1–3×HA-Inc^{R85E} (pNS436) was generated similarly to pNS409, substituting

primers oNS683/oNS692 and oNS684/oNS691 for generating the 5' and 3' Inc^{R85E} fragments,

respectively.

pAc5.1–3×HA-Inc$^{D73A/N82A}$ (pNS414) was generated similarly to pNS409, substituting

pNS430 template and primers oNS683/oNS686 for generating the 5' Inc$^{D73A/N82A}$ fragment; and

pNS431 template and primers oNS684/oNS685 for generating the 3' Inc$^{D73A/N82A}$ fragment.

pAc5.1–3×HA-Inc$^{K88D/E101K}$ (pNS415) was generated similarly to pNS409, substituting

pNS432 template and primers oNS683/oNS688 for generating the 5' Inc$^{K88D/E101K}$ fragment; and

pNS433 template and primers oNS684/oNS687 for generating the 3' Inc$^{K88D/E101K}$ fragment.

pAc5.1–3×HA-Inc$^{T36A/D71A}$ (pNS416) was generated similarly to pNS409, substituting

pNS434 template and primers oNS683/oNS693 for generating the 5' Inc$^{T36A/D71A}$ fragment; and

pNS435 template and primers oNS684/oNS689 for generating the 3' Inc$^{T36A/D71A}$ fragment.

pAc5.1–3×HA-Inc$^{D71A/R85E}$ (pNS417) was generated similarly to pNS409, substituting

pNS435 template and primers oNS683/oNS692 for generating the 5' Inc$^{D71A/R85E}$ fragment; and

pNS436 template and primers oNS684/oNS691 for generating the 3' Inc$^{D71A/R85E}$ fragment.

pAc5.1–3×HA-Inc$^{T36A/D71A/R85E}$ (pNS418) was generated by ligating EcoRV-XhoI

digested pNS434 backbone and an EcoRV-XhoI 3×HA-Inc$^{D71A/R85E}$ fragment from pNS417.

pAc5.1–3×HA-Inc^{F105A} (pNS419) was generated similarly to pNS409, substituting

primers oNS683/oNS1123 and oNS684/oNS1122 for generating the 5' and 3' Inc^{F105A}

fragments, respectively.

pAc5.1–3×HA-Inc^{Y106F} (pNS420) was generated similarly to pNS409, substituting

primers oNS683/oNS1146 and oNS684/oNS1145 for generating the 5' and 3' Inc^{Y106F}

fragments, respectively.

pAc5.1–3×HA-Inc^{N107A} (pNS421) was generated similarly to pNS409, substituting

primers oNS683/oNS1125 and oNS684/oNS1124 for generating the 5' and 3' Inc^{N107A}

fragments, respectively.

pAc5.1–3×HA-Inc$^{F47A/F105A}$ (pNS422) was generated similarly to pNS409, substituting

pNS409 template, primers oNS683/oNS1123, and oNS684/oNS1122 for generating the 5' and 3'

Inc$^{F47A/F105A}$ fragments, respectively.

pAc5.1–3×HA-Inc^{R135H} (pNS426) was generated by ligating HindIII-XhoI digested

pNS402 backbone and a HindIII-XhoI Inc^{R135H} fragment from pNS428 (as described below).

pAc5.1–3×HA-Inc^{AAA} (pNS427) was generated similarly to pNS426, substituting a

HindIII-XhoI Inc^{AAA} fragment from pNS429 (as described below).

Vectors for *Drosophila* transgenesis were as follows:

pUASTattB–Myc-Inc (pNS346) encodes an N-terminal Myc epitope

(MEQKLISEEDLAS) fused to Inc, as previously described (Li et al., 2017).

pUASTattB–Myc-Inc^{R135H} (pNS428) was generated by ligating a HindIII-XhoI digested

pNS346 backbone and a HindIII-XhoI Inc^{R135H} fragment. The Inc^{R135H} fragment was liberated

from the fusion PCR amplification product of primers oNS1126 and oNS1127 and an equimolar

mix of overlapping 5' and 3' Inc^{R135H} fragments as template. The 5' and 3' fragments were

generated by PCR amplification of pNS346 template with primers oNS1126/oNS1555 and

oNS1127/oNS1554, respectively.

pUASTattB–Myc-Inc^{AAA} (pNS429) was generated similarly as pNS428, substituting a

HindIII-XhoI Inc^{AAA} fragment. The Inc^{AAA} fragment was generated similarly as Inc^{R135H},

substituting primers oNS1126/oNS1557 and oNS1127/oNS1556 for generating the 5' and 3'

Inc^{AAA} fragments, respectively.

pUAST–Inc-HA (pNS273) encodes Inc fused to a C-terminal HA epitope

(GSYPYDVPDYA) and was generated by three piece ligation of pUAST BglII-XhoI, a BglII-

EcoRI *inc* fragment liberated from pNS272 (Stavropoulos and Young, 2011), and an EcoRI-

XhoI HA fragment generated by phosphorylating and annealing oligonucleotides oNS191 and

oNS192.

pUASTattB–3×FLAG-Inc (pNS404) encodes an N-terminal 3×FLAG epitope

(MDYKDDDDKGSDYKDDDDKGSDYKDDDDKAS) fused to Inc and was generated by three

piece ligation of EcoRI-XhoI digested pUASTattB, an EcoRI-NheI 3×FLAG fragment liberated

from the PCR amplification product of pNS311 template and primers ACF and oNS241, and a

NheI-XhoI *inc* fragment liberated from pNS351.

pUASTattB–3×FLAG- Inc$^{1-186}$ (pNS405) was generated similar to pNS404, substituting a

NheI-XhoI Inc fragment prepared as for pNS376.

pUASTattB–3×FLAG-Inc$^{1-156}$ (pNS406) was generated similar to pNS404, substituting a

NheI-XhoI Inc fragment prepared as for pNS375.

pUASTattB–3×FLAG- Inc$^{1-123}$ (pNS407) was generated similar to pNS404, substituting a

NheI-XhoI Inc fragment prepared as for pNS374.

pUASTattB–3×FLAG-Inc^{F47A} (pNS423) was generated similar to pNS404, substituting a

NheI-XhoI Inc fragment prepared as for pNS409.

pUASTattB–3×FLAG-Inc^{F105A} (pNS424) was generated similar to pNS404, substituting a

NheI-XhoI Inc fragment prepared as for pNS419.

pUASTattB–3×FLAG-Inc$^{F47A/F105A}$ (pNS425) was generated similar to pNS404,

substituting a NheI-XhoI Inc fragment prepared as for pNS422.

Oligonucleotides

Oligonucleotides used in this work, listed 5' to 3', are as follows:

oNS98   ACTGGGATCCTGGGAGCACGAGCAGCAAAGGAG

oNS184   CCAGCCATCCGACAGCGTTGAGATC

oNS191

AATTTTGGGAATTGGATCCTACCCCTACGATGTGCCCGATTACGCCTAAC

oNS192

TCGAGTTAGGCGTAATCGGGCACATCGTAGGGGTAGGATCCAATTCCCAA

oNS234

TCGAGGCTAGCGACTACAAGGATGATGACGATAAGGGCTCCGATTACAAGGA

CGACGATGATAAGGGATCCGATTACAAGGATGATGACGACAAGTGAT

oNS235

CTAGATCACTTGTCGTCATCATCCTTGTAATCGGATCCCTTATCATCGTCGTCC

TTGTAATCGGAGCCCTTATCGTCATCATCCTTGTAGTCGCTAGCC

oNS240

ACTGGAATTCCGCGGCAACATGGACTACAAGGATGATGACGATAAGGGC

oNS241

ACTGGCGGCCGCTCCTAGGGTGCTAGCCTTGTCGTCATCATCCTTGTAATCGG

AT

oNS277   ACGTGCTAGCATGAGCACGGTGTTCATAAACTCGC

oNS283   GATCTCAACATGGAGCAGAAGCTGATCAGCGAGGAGGATCTGG

oNS284   CTAGCCAGATCCTCCTCGCTGATCAGCTTCTGCTCCATGTTGA

oNS285   ACGTGCTAGCTCGAGGGGTTGTGTGTGAATATATAGCGCGA

oNS315   ACGTGCTAGCCAGTGGGTCAAGCTGAACGTAG

oNS316   ACGTGCTAGCACCTACTTCCTCACCACAAAGACG

oNS317   ACGTGCTAGCCAGCGACCCCAAACGGACAA

oNS319   ACGTCTCGAGTTAATCCCTGTGCAGGATGCACTC

oNS320   ACGTCTCGAGTTACCTCCAGCCATCCGACAG

oNS321   ACGTCTCGAGTTATGTGCCACACTCTTTGGATACCA

oNS337   TTTTTTTTTTTTTTTTTTTTTTTVN

oNS683   ACGTGCTAGCATGAGCACGGTGTTCATAAACTCGC

544	oNS684	ACGTCTCGAGGGGTTGTGTGTGAATATATAGCGCGA
545	oNS685	CCTGATCGACAGAGCCCCCAAATACTTTGC
546	oNS686	GGCGCAGGTAGGCGAGCACGGGTGCAAA
547	oNS687	TGCGCCACGGCGACCTTGTGCTCGAT
548	oNS688	TAGAACTCAGCCTCCTTCAGGACGCCTTC
549	oNS689	CTACTTCCTCACCGCCAAGACGACGCTC
550	oNS691	GCCTACCTGATCGCCAGAGACCCCAAA
551	oNS692	CAAGCTTGCCGTGTTCCAGGTAATTGAGC
552	oNS693	TTTGGGGTCTCTGGCGATCAGGTAGGC
553	oNS694	GACCCAAATTCGGCCCTCTCCCGTCTG
554	oNS695	CAGACGGGAGAGGGCCGAATTTGGGTC
555	oNS696	TCGTTCCTCTCCGAACTGATTCAGGAGG
556	oNS697	CCTCCTGAATCAGTTCGGAGAGGAACGA
557	oNS698	CAGGAGGACTGCGCCTTGATATCAGATCG
558	oNS699	CGATCTGATATCAAGGCGCAGTCCTCCTG
559	oNS700	CGACTTGATATCAGCCCGGGACGAGAC
560	oNS701	GTCTCGTCCCGGGCTGATATCAAGTCG
561	oNS702	CCTGGAGGAGGCTAAGTTCTACAACGTGAC
562	oNS703	GTCACGTTGTAGAACTTAGCCTCCTCCAGG
563	oNS810	ATGCTACTTTTGTCGCCCATCGC
564	oNS811	CTGGGTTATCCTTGGTTTATCCTGGCCT
565	oNS1122	GGAGGCTGAGGCCTACAACGTGAC
566	oNS1123	GTCACGTTGTAGGCCTCAGCCTCC

oNS1124   GCTGAGTTCTACGCCGTGACGCAGC

oNS1125   GCTGCGTCACGGCGTAGAACTCAGC

oNS1126   CAACTGCAACTACTGAAARCRGCCAAGAAG

oNS1127   GGTAGTTTGTCCAATTATGTCACACCACAGAAG

oNS1145   GAGGCTGAGTTCTTCAACGTGACGCAGC

oNS1146   GCTGCGTCACGTTGAAGAACTCAGCCTC

oNS1554   AAGCGCGTTTATCATGTGCTGCAGTGC

oNS1555   GCACTGCAGCACATGATAAACGCGCTT

oNS1556

ATCAGCATGCAGTACACGGCCGCCGCGCCCTTCGAAAACAATGAGTTCCTG

oNS1557   ATTGTTTTCGAAGGGCGCGGCGGCCGTGTACTGCATGCTGATCAGCT

ACF   GACACAAAGCCGCTCCATCAG

attP2-5' CACTGGCACTAGAACAAAAGCTTTGGCG

RPS3A   CGAACCTTCCGATTTCCAAGAAACGC

RPS3B   ACGACGGACGGCCAGTCCTCC

Cell culture and biochemistry

S2 cells were cultured in S2 media containing 10% FBS, penicillin, and streptomycin,

and were transfected with Effectene (Qiagen) as described previously (Stavropoulos and Young,

2011). Transfections were performed in 6 well plates for ~24 hr, after which liposome-

containing media was replaced with fresh culture media. 400 ng of total DNA was used for each

transfection. For transfections involving two plasmids, an equal amount of each was used. Empty

vector lacking insert was used to equalize DNA amounts as indicated in Figures . Cells were

harvested 36-48 hr after transfections were initiated, washed twice in PBS, and lysed in ice-cold

NP40 lysis buffer (50 mM Tris pH 7.6, 150mM NaCl, 0.5% NP40) containing protease

inhibitors. Protein extracts were quantitated in duplicate (BioRad, 5000111).

For co-immunoprecipitations of truncated Inc proteins from S2 cells, 700-1000 μg total

protein was incubated overnight at 4°C with 1:100 anti-FLAG (Sigma, F1804) or anti-Myc

(Sigma, C3956) antibody. Complexes were precipitated by incubation with Gammabind G

sepharose beads (Life Technologies, 10-1243) for 1 hr at 4°C on a nutator, washed 4×5 min at

4°C with lysis buffer, and denatured in SDS sample buffer, separated on Tris SDS-PAGE gels,

and transferred to nitrocellulose. For co-immunoprecipitations of Inc point mutant proteins from

S2 cells, 450-1000 μg of total protein was immunoprecipitated overnight with 20 μl (50% slurry)

of anti-FLAG (Sigma, F2426) affinity gel at 4°C on a nutator. Samples were then washed 4×5

min at 4°C with lysis buffer, denatured in SDS sample buffer, separated on Tris SDS-PAGE

gels, and transferred to nitrocellulose. Membranes were blocked for 1-1.5 hr at room temperature

in LI-COR Odyssey buffer (LI-COR, 927-40000). Membranes were subsequently incubated in

blocking buffer containing 0.1% Tween 20 and the appropriate primary antibodies for 1-2 hr at

room temperature or 4°C overnight: rabbit anti-Myc (1:2,000, Sigma, C3956), mouse anti-FLAG

(1:2,000, Sigma, F1804), rat anti-HA (1:2,000, Roche, 11867431001), and rabbit anti-HA

(1:2,000, Bethyl Laboratories, A190-208A). After washing 4×5 min in a solution containing 150

mM NaCl, 10mM Tris pH 7.6, and 0.1% Tween 20 (TBST), membranes were incubated in the

dark for 30-60 min at room temperature with appropriate secondary antibodies, all diluted

1:15,000 or 1:30,000 in blocking buffer containing 0.1% Tween 20 and 0.01% SDS: Alexa 680

donkey anti-rabbit (Life Technologies, A10043), Alexa 680 donkey anti-mouse (Life

Technologies, A10038), Alexa 790 anti-mouse (Life Technologies, A11371), Alexa 790 anti-rat

(Jackson ImmunoResearch, 712-655-153). Membranes were then washed 4×5 min in TBST, 1×5

min in TBS, and imaged on a Li-Cor Odyssey CLx instrument.

Fly protein extracts were prepared from whole animals or from sieved heads by manual

pestle homogenization in ice-cold NP40 lysis buffer supplemented with protease inhibitors. For

co-immunoprecipitation from fly head protein extracts, complexes were immunoprecipitated

with 30 µl (50% slurry) of anti-Myc (Sigma, E6654) affinity gel for 1.5 hr at 4°C on a nutator.

Samples were then washed 4×5 min at 4°C with lysis buffer, denatured in SDS sample buffer,

separated on Tris SDS-PAGE gels and blotted as described above. Primary antibodies were

rabbit anti-Myc (1:2,000, Sigma, C3956), mouse anti-FLAG (1:2,000, Sigma, F1804), and rat

anti-HA (1:2,000, Roche, 11867431001). Secondary antibodies were Alexa 680 donkey anti-rat

(Jackson ImmunoResearch, 712-625-1533), Alexa 680 donkey anti-mouse (Life Technologies,

A10038), Alexa 790 anti-rabbit (Life Technologies, A11374). For assessing in vivo expression

of Inc proteins, 30 µg was separated on Tris-SDS-PAGE gels and blotted as described above.

Primary antibodies were rabbit anti-Myc (1:2,000, Sigma, C3956), mouse anti-FLAG (1:2,000,

Sigma, F1804), mouse anti-tubulin (1:10,000, DSHB, 12G10), and rabbit anti-tubulin (1:60,000,

VWR, 89364-004). Secondary antibodies were Alexa 680 anti-mouse (Life Technologies,

A10038) and Alexa 790 anti-rabbit (Life Technologies, A11374).

qRT-PCR

Total RNA was isolated using TRIZOL (Life Technologies, 15596-026). 5 µg of RNA

was reverse transcribed with a poly-T primer (oNS337) and SuperScript II reverse transcriptase.

qPCR was performed using SYBR Green Supermix (Bio-Rad, 1725272) and the following

primers: oNS98 and oNS184 (*inc*); oNS810 and oNS811 (*Cul3*); RPS3A and RPS3B (*rps3*). All

real-time PCR reactions were performed using a BioRad DNA Engine Opticon 2 System.

Fly stocks and transgenes

elav[c155]-*Gal4* (Lin and Goodman, 1994), *inc*[1] (Stavropoulos and Young, 2011), *inc-Gal4*

(Stavropoulos and Young, 2011), *inc*[1] *inc-Gal4* (Li et al., 2017), *attP2: UAS-Myc-Inc* (Li et al.,

2017), *attP2: UAS-3×FLAG-Inc* (Li et al., 2017), and *UAS-3×FLAG-3×HA-Cul3* (Hudson and

Cooley, 2010) are previously described. *UAS-Cul3-RNAi* is 11861R-2 obtained from the NIG-

Fly stock center. Transgenic flies generated in this study using pUASTattB-based vectors were

integrated at *attP2* (Groth et al., 2004) with phiC31 recombinase (BestGene); integration was

verified by PCR using primer attP2-5' paired with oNS277. All transgenes were backcrossed six

to eight generations to Bloomington stock 5905, an isogenic w^{1118} stock described elsewhere as

iso31 (Ryder et al., 2004).

Sleep analysis

Crosses were set with five virgin females and three males on cornmeal, agar, and

molasses food. One to four day old male flies eclosing from LD-entrained cultures raised at 25°C

were loaded in glass tubes containing cornmeal, agar, and molasses food. Animals were

monitored for 5-7 days at 25°C in LD cycles using DAM2 monitors (Trikinetics). The first 36-48

hours of data were discarded and an integral number of days of data (3-5) were analyzed using

custom Matlab code. Locomotor data were collected in 1 min bins. Sleep was defined by

locomotor inactivity for 5 min or more; all minutes within inactive periods exceeding 5 min were

assigned as sleep. This definition classifies more sleep than the definition used in our prior

studies (Li et al., 2017; Li and Stavropoulos, 2016; Stavropoulos and Young, 2011), in which

sleep is scored as beginning on the fifth minute of locomotor inactivity and the preceding four

minutes are classified as quiet wakefulness. Dead animals were excluded from analysis by a

combination of automated filtering and visual inspection of locomotor traces.

Statistics

One-way ANOVA and Tukey post-hoc tests were used for comparisons of total sleep,

daytime sleep, nighttime sleep, and sleep bout number. Nonparametric Kruskal-Wallis tests and

Dunn's post hoc tests were used for comparisons of sleep bout length. Unpaired two-sided

Student's t-tests were used for comparisons of *inc* mRNA levels in vivo.

Sequence alignments

Alignments were performed with Clustal Omega 2.1 and BOXSHADE. GenBank

accession numbers for proteins in Figure S1A are: Inc, NP_001284787; KCTD2, NP_056168;

KCTD5, NP_061865; KCTD17.3, NP_001269614; KCTD17.4, NP_001269615.

Figure Legends

Figure 1. The Inc BTB domain mediates Inc-Cul3 and Inc-Inc associations

A) Schematic of N- and C-terminally truncated Inc proteins. CTD, C-terminal domain. **B and C)**

Co-immunoprecipitation of 3×Myc-tagged Inc or Inc truncation proteins with 3×FLAG-Cul3 **(B)**

or Inc-HA **(C)** from transiently transfected S2 cells. LC, immunoglobulin light chain.

Figure 2. Identification of Inc BTB domain point mutants that specifically weaken Inc-Cul3

associations

A-D) Co-immunoprecipitation analysis of 3×HA-tagged Inc or Inc point mutants and 3×FLAG-

Cul3 **(A and C)** or 3×FLAG-Inc **(B and D)** from transiently transfected S2 cells.

Figure 3. Inc-Cul3 binding is required for Inc activity in vivo

Biochemical and behavioral analysis of *inc^1 inc-Gal4* animals expressing 3×FLAG-tagged Inc or

Inc point mutants that weaken Cul3 associations. **A)** Immunoblot analysis of whole animal lysates.

B) Total sleep duration. Mean ± SEM is shown. n = 16-58; *p < 0.01 for comparison to *inc^1 inc-*

Gal4 animals and not significantly different from wild-type controls; ‡p < 0.01 for comparisons to

inc^1 inc-Gal4 animals and wild-type controls. **C)** Population average sleep traces summed hourly

for indicated genotypes. n = 41-58. For all panels, animals are heterozygous for UAS transgenes.

Figure 4. The conserved Inc C-terminus is essential for Inc activity in vivo

Biochemical and behavioral analysis of *inc^1 inc-Gal4* animals expressing 3×FLAG-tagged Inc or

C-terminally truncated Inc proteins. **A)** Immunoblot analysis of whole animal lysates. **B)** Total

sleep duration. Mean ± SEM is shown. n = 27-30; *p < 0.01 for comparison to *inc¹ inc-Gal4*

animals and not significantly different from wild-type controls. **C)** Population average sleep traces

summed hourly for indicated genotypes. n = 27-30. For all panels, animals are heterozygous for

UAS transgenes.

Figure 5. Inc C-terminal point mutants do not alter Inc-Cul3 and Inc-Inc associations

Co-immunoprecipitation analysis of 3×HA-tagged Inc or Inc point mutants and 3×FLAG-Cul3 or

3×FLAG-Inc from transiently transfected S2 cells.

Figure 6. The conserved Inc C-terminal arginine is vital for Inc activity in vivo

Biochemical and behavioral analysis of *inc¹ inc-Gal4* animals expressing 3×Myc-tagged Inc or

Inc point mutants. **A)** Immunoblot analysis of whole animal lysates. **B)** Total sleep duration. Mean

± SEM is shown. n = 24-46; *p < 0.01 for comparison to *inc¹ inc-Gal4* animals and not

significantly different from wild-type controls. **C)** Population average sleep traces summed hourly

for indicated genotypes. n = 24-46. For all panels, animals are heterozygous for UAS transgenes.

Figure 7. Inc exhibits key properties of a Cul3 adaptor in neurons in vivo

Co-immunoprecipitation analysis of 3×Myc-tagged Inc and 3×FLAG-3×HA-Cul3 **(A)** or 3×HA-

Inc **(B)** from head lysates prepared from indicated genotypes. **C)** Western blots of head lysates

prepared from indicated genotypes. **D)** qRT-PCR analysis of *inc* mRNA levels of indicated

genotypes. Mean ± SEM is shown. *p < 0.05; ns, not significant (p>0.05).

Table 1. Summary of co-immunoprecipitation analysis of Inc point mutants designed to

weaken Inc-Cul3 interaction from transfected S2 cells

Table 2. Summary of co-immunoprecipitation analysis of Inc point mutants designed to

weaken Inc-Inc interaction from transfected S2 cells

Supplementary Figure 1. Sequence and crystal structure analysis of Inc and human Inc

orthologs

(A) Alignment of Inc and its human orthologs. Identical and similar residues are shaded in black

and gray, respectively. Locations of Inc truncations are indicated by arrowheads. Inc point mutants

are indicated by closed circles. **(B)** Crystal structure of the BTB domains of adjacent subunits of

human KCTD5. Residues that do not impair Inc-Cul3 association when mutated are highlighted

in cyan. Mutations that weaken Inc-Cul3 associations (F47A and F105A) are highlighted in blue.

(C) Crystal structure of the BTB domains of adjacent subunits of human KCTD5. Residues

mutated to impair Inc-Inc association are indicated. The T36A/D71A/R85E triple mutation

weakens Inc homomultimerization and decreases Inc stability. **(D)** Crystal structure of human

KCTD5. Inc C-terminal point mutants are indicated.

Supplementary Figure 2. Neuronal expression of Inc BTB-domain point mutants does not

perturb sleep

(A) Biochemical analysis of *inc-Gal4* animals expressing 3×FLAG-tagged Inc or Inc point mutants.

(B) Behavioral analysis of *elav^c155^-Gal4* animals expressing 3×FLAG-tagged Inc or Inc point

mutants. Mean ± SEM is shown. n = 5-24; ns, not significant (p>0.05**)** compared to *elav^c155^-Gal4*

control. For all panels, animals are heterozygous for UAS transgenes.

Supplementary Figure 3. Additional sleep parameters for animals expressing Inc and Inc

BTB domain point mutants

(A-D) Sleep parameters for *inc^1^ inc-Gal4* animals expressing 3×FLAG-tagged Inc or Inc point

mutants. Mean ± SEM is shown. n = 16-58 as in Fig 3B; *p < 0.01 for comparison to *inc^1^ inc-Gal4*

animals and not significantly different from wild-type controls; ‡p < 0.01 for comparisons to *inc¹*

inc-Gal4 animals and wild-type controls. **A)** Nighttime sleep. **B)** Daytime sleep. **C)** Sleep bout

duration. **D)** Sleep bout number. For all panels, animals are heterozygous for UAS transgenes.

Supplementary Figure 4. Neuronal expression of Inc C-terminal truncations does not alter

sleep

Behavioral analysis of *elav^{c155}-Gal4* animals expressing 3×FLAG-tagged Inc or C-terminally

truncated Inc proteins. Mean ± SEM is shown. n = 23-54; ns, not significant (p>0.05) compared

to *elav^{c155}-Gal4* control. For all panels, animals are heterozygous for UAS transgenes.

Supplementary Figure 5. Additional sleep parameters for animals expressing Inc C-terminal

truncations

(A-D) Sleep parameters for *inc¹ inc-Gal4* animals expressing 3×FLAG-tagged Inc or C-terminally

truncated Inc proteins. Mean ± SEM is shown. n = 27-30 as in Fig 4B; *p < 0.01 for comparison

to *inc¹ inc-Gal4* animals and not significantly different from wild-type controls; ‡p < 0.01 for

comparisons to *inc¹ inc-Gal4* animals and wild-type controls. **A)** Nighttime sleep. **B)** Daytime

sleep. **C)** Sleep bout duration. **D)** Sleep bout number. For all panels, animals are heterozygous for

UAS transgenes.

Supplementary Figure 6. Neuronal expression of Inc C-terminal point mutants does not

perturb sleep

Behavioral analysis of *elav^c155^-Gal4* animals expressing 3×Myc-tagged Inc or Inc point mutants.

Mean ± SEM is shown. n = 18-65; ns, not significant (p>0.05) compared to *elav^c155^-Gal4* control.

For all panels, animals are heterozygous for UAS transgenes.

Supplementary Figure 7. Additional sleep parameters for animals expressing Inc C-terminal

point mutants.

(A-D) Sleep parameters for *inc^1^ inc-Gal4* animals expressing 3×Myc-tagged Inc or Inc point

mutants. Mean ± SEM is shown. n = 24-46 as in Fig 4B; *p < 0.01 for comparison to *inc^1^ inc-Gal4*

animals and not significantly different from wild-type controls; ‡p < 0.01 for comparisons to *inc^1^*

inc-Gal4 animals and wild-type controls. **A)** Nighttime sleep. **B)** Daytime sleep. **C)** Sleep bout

duration. **D)** Sleep bout number. For all panels, animals are heterozygous for UAS transgenes.

References

Afonso, D.J.S., Liu, D., Machado, D.R., Pan, H., Jepson, J.E.C., Rogulja, D., Koh, K., 2015.
TARANIS Functions with Cyclin A and Cdk1 in a Novel Arousal Center to Control Sleep in
Drosophila. Curr Biol 25, 1717–1726. doi:10.1016/j.cub.2015.05.037

Balasco, N., Pirone, L., Smaldone, G., Di Gaetano, S., Esposito, L., Pedone, E.M., Vitagliano,
L., 2014. Molecular recognition of Cullin3 by KCTDs: insights from experimental and
computational investigations. Biochim. Biophys. Acta 1844, 1289–1298.
doi:10.1016/j.bbapap.2014.04.006

Brockmann, M., Blomen, V.A., Nieuwenhuis, J., Stickel, E., Raaben, M., Bleijerveld, O.B.,
Altelaar, A.F.M., Jae, L.T., Brummelkamp, T.R., 2017. Genetic wiring maps of single-cell
protein states reveal an off-switch for GPCR signalling. Nature 546, 307–311.
doi:10.1038/nature22376

Butler, A., Wei, A.G., Baker, K., Salkoff, L., 1989. A family of putative potassium channel
genes in Drosophila. Science 243, 943–947. doi:10.1126/science.2493160

Canning, P., Cooper, C.D.O., Krojer, T., Murray, J.W., Pike, A.C.W., Chaikuad, A., Keates, T.,
Thangaratnarajah, C., Hojzan, V., Ayinampudi, V., Marsden, B.D., Gileadi, O., Knapp, S.,
Delft, von, F., Bullock, A.N., 2013. Structural basis for Cul3 protein assembly with the
BTB-Kelch family of E3 ubiquitin ligases. J Biol Chem 288, 7803–7814.
doi:10.1074/jbc.M112.437996

Chiu, C.N., Rihel, J., Lee, D.A., Singh, C., Mosser, E.A., Chen, S., Sapin, V., Pham, U., Engle,
J., Niles, B.J., Montz, C.J., Chakravarthy, S., Zimmerman, S., Salehi-Ashtiani, K., Vidal, M.,
Schier, A.F., Prober, D.A., 2016. A Zebrafish Genetic Screen Identifies Neuromedin U as a
Regulator of Sleep/Wake States. Neuron 89, 842–856. doi:10.1016/j.neuron.2016.01.007

Cirelli, C., Bushey, D., Hill, S., Huber, R., Kreber, R., Ganetzky, B., Tononi, G., 2005. Reduced
sleep in Drosophila Shaker mutants. Nature 434, 1087–1092. doi:10.1038/nature03486

Dementieva, I.S., Tereshko, V., McCrossan, Z.A., Solomaha, E., Araki, D., Xu, C., Grigorieff,
N., Goldstein, S.A.N., 2009. Pentameric assembly of potassium channel tetramerization
domain-containing protein 5. J Mol Biol 387, 175–191. doi:10.1016/j.jmb.2009.01.030

DiBello, P.R., Withers, D.A., Bayer, C.A., Fristrom, J.W., Guild, G.M., 1991. The Drosophila
Broad-Complex encodes a family of related proteins containing zinc fingers. Genetics 129,
385–397.

Djagaeva, I., Doronkin, S., 2009. COP9 limits dendritic branching via Cullin3-dependent
degradation of the actin-crosslinking BTB-domain protein Kelch. PLoS ONE 4, e7598.
doi:10.1371/journal.pone.0007598

Emanuele, M.J., Elia, A.E.H., Xu, Q., Thoma, C.R., Izhar, L., Leng, Y., Guo, A., Chen, Y.-N.,
Rush, J., Hsu, P.W.-C., Yen, H.-C.S., Elledge, S.J., 2011. Global identification of modular
cullin-RING ligase substrates. Cell 147, 459–474. doi:10.1016/j.cell.2011.09.019
842

843   Errington, W.J., Khan, M.Q., Bueler, S.A., Rubinstein, J.L., Chakrabartty, A., Privé, G.G., 2012.
844   Adaptor protein self-assembly drives the control of a cullin-RING ubiquitin ligase. Structure
845   20, 1141–1153. doi:10.1016/j.str.2012.04.009
846

Funato, H., Miyoshi, C., Fujiyama, T., Kanda, T., Sato, M., Wang, Z., Ma, J., Nakane, S.,
Tomita, J., Ikkyu, A., Kakizaki, M., Hotta-Hirashima, N., Kanno, S., Komiya, H., Asano, F.,
Honda, T., Kim, S.J., Harano, K., Muramoto, H., Yonezawa, T., Mizuno, S., Miyazaki, S.,
Connor, L., Kumar, V., Miura, I., Suzuki, T., Watanabe, A., Abe, M., Sugiyama, F.,
Takahashi, S., Sakimura, K., Hayashi, Y., Liu, Q., Kume, K., Wakana, S., Takahashi, J.S.,
Yanagisawa, M., 2016. Forward-genetics analysis of sleep in randomly mutagenized mice.
Nature 539, 378–383. doi:10.1038/nature20142
854

855   Furukawa, M., He, Y.J., Borchers, C., Xiong, Y., 2003. Targeting of protein ubiquitination by
856   BTB-Cullin 3-Roc1 ubiquitin ligases. Nat Cell Biol 5, 1001–1007. doi:10.1038/ncb1056
857

858   Geyer, R., Wee, S., Anderson, S., Yates, J., Wolf, D.A., 2003. BTB/POZ domain proteins are
859   putative substrate adaptors for cullin 3 ubiquitin ligases. Mol Cell 12, 783–790.
860

861   Godt, D., Couderc, J.L., Cramton, S.E., Laski, F.A., 1993. Pattern formation in the limbs of
862   Drosophila: bric à brac is expressed in both a gradient and a wave-like pattern and is
863   required for specification and proper segmentation of the tarsus. Development 119, 799–812.
864

865   Groth, A.C., Fish, M., Nusse, R., Calos, M.P., 2004. Construction of transgenic Drosophila by
866   using the site-specific integrase from phage phiC31. Genetics 166, 1775–1782.
867

868   Harrison, S.D., Travers, A.A., 1990. The tramtrack gene encodes a Drosophila finger protein that
869   interacts with the ftz transcriptional regulatory region and shows a novel embryonic
870   expression pattern. EMBO J 9, 207–216.
871

872   Hudson, A.M., Cooley, L., 2010. Drosophila Kelch functions with Cullin-3 to organize the ring
873   canal actin cytoskeleton. J Cell Biol 188, 29–37. doi:10.1083/jcb.200909017
874

875   Iannacone, M.J., Beets, I., Lopes, L.E., Churgin, M.A., Fang-Yen, C., Nelson, M.D., Schoofs, L.,
876   Raizen, D.M., 2017. The RFamide receptor DMSR-1 regulates stress-induced sleep in C.
877   elegans. Elife 6, 377. doi:10.7554/eLife.19837
878

879   Ji, A.X., Chu, A., Nielsen, T.K., Benlekbir, S., Rubinstein, J.L., Privé, G.G., 2015. Structural
880   insights into KCTD protein assembly and Cullin3 recognition. J Mol Biol.
881   doi:10.1016/j.jmb.2015.08.019
882

Ji, A.X., Privé, G.G., 2013. Crystal Structure of KLHL3 in Complex with Cullin3. PLoS ONE 8,
e60445. doi:10.1371/journal.pone.0060445

Kasahara, K., Kawakami, Y., Kiyono, T., Yonemura, S., Kawamura, Y., Era, S., Matsuzaki, F.,
Goshima, N., Inagaki, M., 2014. Ubiquitin-proteasome system controls ciliogenesis at the
initial step of axoneme extension. Nat Comms 5, 5081. doi:10.1038/ncomms6081

Kim, E.-J., Kim, S.-H., Jin, X., Jin, X., Kim, H., 2017. KCTD2, an adaptor of Cullin3 E3
ubiquitin ligase, suppresses gliomagenesis by destabilizing c-Myc. Cell Death Differ 24,
649–659. doi:10.1038/cdd.2016.151

Koh, K., Joiner, W.J., Wu, M.N., Yue, Z., Smith, C.J., Sehgal, A., 2008. Identification of
SLEEPLESS, a sleep-promoting factor. Science 321, 372–376. doi:10.1126/science.1155942

Li, Q., Kellner, D.A., Hatch, H.A.M., Yumita, T., Sanchez, S., Machold, R.P., Frank, C.A.,
Stavropoulos, N., 2017. Conserved properties of Drosophila Insomniac link sleep regulation
and synaptic function. PLoS Genet 13, e1006815. doi:10.1371/journal.pgen.1006815

Li, Q., Stavropoulos, N., 2016. Evaluation of Ligand-Inducible Expression Systems for
Conditional Neuronal Manipulations of Sleep in Drosophila. G3: Genes | Genomes |
Genetics 6, 3351–3359. doi:10.1534/g3.116.034132

Lin, D.M., Goodman, C.S., 1994. Ectopic and increased expression of Fasciclin II alters
motoneuron growth cone guidance. Neuron 13, 507–523.

McGourty, C.A., Akopian, D., Walsh, C., Gorur, A., Werner, A., Schekman, R., Bautista, D.,
Rape, M., 2016. Regulation of the CUL3 Ubiquitin Ligase by a Calcium-Dependent Co-
adaptor. Cell 167, 525–538.e14. doi:10.1016/j.cell.2016.09.026

Mencacci, N.E., Rubio-Agusti, I., Zdebik, A., Asmus, F., Ludtmann, M.H.R., Ryten, M.,
Plagnol, V., Hauser, A.-K., Bandres-Ciga, S., Bettencourt, C., Forabosco, P., Hughes, D.,
Soutar, M.M.P., Peall, K., Morris, H.R., Trabzuni, D., Tekman, M., Stanescu, H.C., Kleta,
R., Carecchio, M., Zorzi, G., Nardocci, N., Garavaglia, B., Lohmann, E., Weissbach, A.,
Klein, C., Hardy, J., Pittman, A.M., Foltynie, T., Abramov, A.Y., Gasser, T., Bhatia, K.P.,
Wood, N.W., 2015. A missense mutation in KCTD17 causes autosomal dominant
myoclonus-dystonia. American journal of human genetics 96, 938–947.
doi:10.1016/j.ajhg.2015.04.008

Papazian, D.M., Schwarz, T.L., Tempel, B.L., Jan, Y.N., Jan, L.Y., 1987. Cloning of genomic
and complementary DNA from Shaker, a putative potassium channel gene from Drosophila.
Science 237, 749–753. doi:10.1126/science.2441470

Pfeiffenberger, C., Allada, R., 2012. Cul3 and the BTB adaptor insomniac are key regulators of
sleep homeostasis and a dopamine arousal pathway in Drosophila. PLoS Genet 8, e1003003.
doi:10.1371/journal.pgen.1003003

Pintard, L., Willis, J.H., Willems, A., Johnson, J.-L.F., Srayko, M., Kurz, T., Glaser, S., Mains,
P.E., Tyers, M., Bowerman, B., Peter, M., 2003. The BTB protein MEL-26 is a substrate-

specific adaptor of the CUL-3 ubiquitin-ligase. Nature 425, 311–316.
doi:10.1038/nature01959
933
934 Pongs, O., Kecskemethy, N., Müller, R., Krah-Jentgens, I., Baumann, A., Kiltz, H.H., Canal, I.,
935   Llamazares, S., Ferrus, A., 1988. Shaker encodes a family of putative potassium channel
936   proteins in the nervous system of Drosophila. EMBO J 7, 1087–1096.
937
938 Rogulja, D., Young, M.W., 2012. Control of sleep by cyclin A and its regulator. Science 335,
939   1617–1621. doi:10.1126/science.1212476
940
Ryder, E., Blows, F., Ashburner, M., Bautista-Llacer, R., Coulson, D., Drummond, J., Webster,
J., Gubb, D., Gunton, N., Johnson, G., O'kane, C.J., Huen, D., Sharma, P., Asztalos, Z.,
Baisch, H., Schulze, J., Kube, M., Kittlaus, K., Reuter, G., Maroy, P., Szidonya, J.,
Rasmuson-Lestander, A., Ekström, K., Dickson, B., Hugentobler, C., Stocker, H., Hafen, E.,
Lepesant, J.A., Pflugfelder, G., Heisenberg, M., Mechler, B., Serras, F., Corominas, M.,
Schneuwly, S., Preat, T., Roote, J., Russell, S., 2004. The DrosDel collection: a set of P-
element insertions for generating custom chromosomal aberrations in Drosophila
melanogaster. Genetics 167, 797–813. doi:10.1534/genetics.104.026658
949
950 Schwenk, J., Metz, M., Zolles, G., Turecek, R., Fritzius, T., Bildl, W., Tarusawa, E., Kulik, A.,
951   Unger, A., Ivankova, K., Seddik, R., Tiao, J.Y., Rajalu, M., Trojanova, J., Rohde, V.,
952   Gassmann, M., Schulte, U., Fakler, B., Bettler, B., 2010. Native GABA(B) receptors are
953   heteromultimers with a family of auxiliary subunits. Nature 465, 231–235.
954   doi:10.1038/nature08964
955
956 Singh, C., Rihel, J., Prober, D.A., 2017. Neuropeptide Y Regulates Sleep by Modulating
957   Noradrenergic Signaling. Curr Biol 27, 3796–3811.e5. doi:10.1016/j.cub.2017.11.018
958
959 Stavropoulos, N., Young, M.W., 2011. insomniac and Cullin-3 regulate sleep and wakefulness in
960   Drosophila. Neuron 72, 964–976. doi:10.1016/j.neuron.2011.12.003
961
962 Stogios, P.J., Downs, G.S., Jauhal, J.J.S., Nandra, S.K., Privé, G.G., 2005. Sequence and
963   structural analysis of BTB domain proteins. Genome Biol 6, R82. doi:10.1186/gb-2005-6-
964   10-r82
965
966 Toda, H., Williams, J.A., Gulledge, M., Sehgal, A., 2019. A sleep-inducing gene, nemuri, links
967   sleep and immune function in Drosophila. Science 363, 509–515.
968   doi:10.1126/science.aat1650
969
970 Wee, S., Geyer, R.K., Toda, T., Wolf, D.A., 2005. CSN facilitates Cullin-RING ubiquitin ligase
971   function by counteracting autocatalytic adapter instability. Nat Cell Biol 7, 387–391.
972   doi:10.1038/ncb1241
973
Wei, A., Covarrubias, M., Butler, A., Baker, K., Pak, M., Salkoff, L., 1990. K+ current diversity
is produced by an extended gene family conserved in Drosophila and mouse. Science 248,
599–603. doi:10.1126/science.2333511

Wu, M., Robinson, J.E., Joiner, W.J., 2014. SLEEPLESS is a bifunctional regulator of
excitability and cholinergic synaptic transmission. Curr Biol 24, 621–629.
doi:10.1016/j.cub.2014.02.026

Wu, M.N., Joiner, W.J., Dean, T., Yue, Z., Smith, C.J., Chen, D., Hoshi, T., Sehgal, A., Koh, K.,
2010. SLEEPLESS, a Ly-6/neurotoxin family member, regulates the levels, localization and
activity of Shaker. Nat Neurosci 13, 69–75. doi:10.1038/nn.2454

Xu, L., Wei, Y., Reboul, J., Vaglio, P., Shin, T.-H., Vidal, M., Elledge, S.J., Harper, J.W., 2003.
BTB proteins are substrate-specific adaptors in an SCF-like modular ubiquitin ligase
containing CUL-3. Nature 425, 316–321. doi:10.1038/nature01985

Zhang, D.D., Lo, S.-C., Sun, Z., Habib, G.M., Lieberman, M.W., Hannink, M., 2005.
Ubiquitination of Keap1, a BTB-Kelch substrate adaptor protein for Cul3, targets Keap1 for
degradation by a proteasome-independent pathway. J Biol Chem 280, 30091–30099.
doi:10.1074/jbc.M501279200

A

Residues

- 1-211
- 22-211
- 31-211
- 124-211
- 1-186
- 1-156
- 1-123
- 22-123

B

C

Figure 1

Figure 2

A

B

C

Figure 3

A

B

C

Figure 4

Figure 5

A

B

C

Figure 6

A

elav^{c155}-Gal4	+	+
UAS-Myc-Inc		+
UAS-3xFLAG-3xHA-Cul3		+

(kDa)

WB α-Myc — Lysate — 25
WB α-Myc — IP α-Myc — 25
WB α-FLAG — Lysate — 80
WB α-FLAG — IP α-Myc — 80

B

elav^{c155}-Gal4	+	+
UAS-Myc-Inc		+
UAS-Inc-HA		+

(kDa)

WB α-Myc — Lysate — 25
WB α-Myc — IP α-Myc — 25
WB α-HA — Lysate — 25
WB α-HA — IP α-Myc — 25

C

elav^{c155}-Gal4; UAS-Dcr2

UAS-Cul3-RNAi   -   -   +   +   *inc^l*

(kDa)
25 — Inc
50 — Tubulin

D

inc / rps3 mRNA (normalized)

* ns *

UAS-inc-RNAi	-	+		
UAS-Cul3-RNAi			-	+

elav^{c155}-Gal4; UAS-Dcr2

inc^+   *inc^l*

Figure 7

Table 1

	Inc mutant	Inc-Inc	Inc-Cul3	Inc stability
1	Inc^{F47A}	n.c.	Strongly reduced	n.c.
2	Inc^{R50E}	Reduced	Reduced	n.c.
3	Inc^{D57A}	n.c.	n.c.	n.c.
4	Inc^{D61A}	n.c.	n.c.	n.c.
5	Inc^{E104K}	n.c.	n.c.	n.c.
6	Inc^{F105A}	n.c.	Strongly reduced	n.c.
7	Inc^{Y106A}	n.c.	n.c.	n.c.
8	Inc^{N107A}	n.c.	n.c.	n.c.
9	Inc$^{F47A/F105A}$	n.c.	Strongly reduced	n.c.

n.c. (no change)

Table 2

	Inc mutant	Inc-Inc	Inc-Cul3	Inc stability
1	Inc$^{T36A/D71A/R85E}$	Strongly reduced	Reduced	Strongly reduced
2	Inc$^{T36A/D71A}$	n.c.	n.c.	n.c.
3	Inc$^{D71A/R85E}$	n.c.	Reduced	Strongly reduced
4	Inc$^{D73A/N82A}$	n.c.	n.c.	n.c.
5	Inc$^{K88D/E101K}$	n.c.	Reduced	n.c.
6	Inc^{T36A}	n.c.	n.c.	n.c.

n.c. (no change)

Supplementary Figure 1

A

| | - | Inc | Inc^{F47A} | Inc^{F105A} | Inc^{F47A/F105A} |

inc-Gal4 + + + + +

FLAG

(kDa) 25 —

Tubulin

46 —

B

ns

Sleep / day (min)

elav^{c155}-Gal4

Supplementary Figure 2

A

B

C

D