1

The Cul3 ubiquitin ligase engages Insomniac as an adaptor to impact sleep and synaptic

homeostasis

Qiuling Li[1], Kayla Y. Lim[1], Raad Altawell[3], Faith Verderose[3], Celine Dong[2], Joshua Martinez[2],

Xiling Li[2], Dion Dickman[2], and Nicholas Stavropoulos[3,*]

[1]NYU Neuroscience Institute, Department of Neuroscience and Physiology, New York

University School of Medicine, New York, NY 10016, USA

[2]Department of Neurobiology, University of Southern California, Los Angeles, CA 90089, USA

[3]Waksman Institute, Rutgers University, Piscataway, NJ 08854, USA

Keywords: Cullin-3, Inc, KCTD2, KCTD5, KCTD17, BTB, autism, schizophrenia, myoclonic

dystonia

Running Title: Cul3 impacts sleep and synapses through Insomniac

*Correspondence: stavropoulos@waksman.rutgers.edu

Abstract

Mutations of the Cullin-3 (Cul3) E3 ubiquitin ligase are associated with autism and schizophrenia, neurological disorders characterized by sleep disturbances and altered synaptic function. Cul3 engages dozens of adaptor proteins to recruit hundreds of substrates for ubiquitination, but the adaptors that impact sleep and synapses remain ill-defined. Here we implicate Insomniac (Inc), a conserved protein required for normal sleep and synaptic homeostasis in *Drosophila*, as a Cul3 adaptor. Inc binds Cul3 in vivo, and mutations within the N-terminal BTB domain of Inc that weaken Inc-Cul3 associations impair Inc activity, indicating that Inc function requires binding to the Cul3 complex. Deletion of the conserved C-terminus of Inc does not alter Cul3 binding but abolishes Inc activity in the context of sleep and synaptic homeostasis, suggesting that the C-terminal domain of Inc is a substrate recruitment domain. Mutation of a conserved, disease-associated arginine in the Inc C-terminus also abolishes Inc function, suggesting that this residue is vital for recruiting Inc targets. Inc levels are negatively regulated by Cul3 in neurons, consistent with Inc degradation by autocatalytic ubiquitination, a hallmark of Cullin adaptors. These findings link Inc and Cul3 in vivo and indicate that Cul3-Inc complexes are essential for normal sleep and synaptic function. Furthermore, these results indicate that dysregulation of conserved substrates of Cul3-Inc complexes may contribute to altered sleep and synaptic function in autism and schizophrenia associated with *Cul3* mutations.

Author Summary

Cul3 is a highly conserved gene important for brain development and function. *Cul3*

mutations are a risk factor for autism and schizophrenia, neurological disorders associated with

disturbed sleep and changes in neuronal synapses. A key challenge in understanding how *Cul3*

impacts brain function is elucidating the downstream molecular pathways. Cul3 is a ubiquitin

ligase that assembles with dozens of adaptor proteins, which in turn recruit specific protein

substrates for ubiquitination. Identifying and characterizing Cul3 adaptors in the nervous system

is thus a critical step in understanding Cul3 function. Because Cul3 and its adaptors are conserved

through evolution, simpler organisms including the fruit fly *Drosophila* provide powerful systems

for identifying and characterizing Cul3 adaptors. We found that the Insomniac (Inc) protein has

the properties of a Cul3 adaptor that impacts sleep and synaptic function in *Drosophila*. These

results suggest that human proteins related to Inc may be relevant for changes in sleep and synaptic

function in autism and schizophrenia associated with reduced Cul3 activity.

Introduction

The Cullin-3 (Cul3) E3 ubiquitin ligase is broadly expressed in the brain and is an important regulator of nervous system development and function. In humans, *Cul3* mutations are linked to autism, its associated sleep disturbances, and to schizophrenia [1–7]. Mice bearing *Cul3* mutations exhibit alterations in neurogenesis, dendritic development, synaptic transmission, and glutamate receptor levels, as well as social deficits relevant to autism spectrum disorder [8–14]. *Cul3* is highly conserved, and reduced activity of *Cul3* in worms and flies causes similar phenotypes within the nervous system, including perturbations in neuronal morphogenesis, glutamate receptor abundance, synaptic homeostasis, and sleep [15–20].

A key challenge in understanding how *Cul3* impacts brain function is elucidating the downstream molecular pathways. Cul3 assembles in a modular fashion with dozens of BTB-domain adaptor proteins that recruit substrates to Cul3 complexes for ubiquitination [21–25]. Which among the known and putative adaptors of Cul3 contribute to its phenotypes—including sleep disturbances and altered synaptic function—remains largely unknown. Though several Cul3 adaptors have been characterized in neurons [16,26–32], many known and putative Cul3 adaptors have yet to be assessed.

insomniac (*inc*) encodes a conserved BTB-domain protein that is hypothesized to serve as a Cul3 adaptor [17,18,33]. *inc* null mutations reduce sleep duration and consolidation in *Drosophila* and impair synaptic homeostasis, phenotypes that are recapitulated by reduced *Cul3* activity [17,18,20]. Inc can bind Cul3 when overexpressed in cultured cells [17,18,33], but whether these interactions occur in vivo or are required for the activity of Inc is unknown. Notably, some BTB-domain proteins associate with Cul3 but are unlikely to serve as adaptors [34]. Functional analysis has been essential to distinguish Cul3 adaptors within the BTB superfamily, which also includes GABA$_B$ receptor subunits, potassium channels, and transcriptional regulators [35,36]. In

the absence of evidence linking Inc and Cul3 in vivo, the role of Inc as a Cul3 adaptor has remained

speculative.

Here we implicate Inc as a Cul3 adaptor, using biochemical, behavioral, and

electrophysiological studies of Inc mutants. Critically, Inc binds Cul3 in vivo and Inc activity

requires the assembly of Inc-Cul3 complexes. Inc function also requires its C-terminus, a

conserved domain dispensable for Cul3 associations but vital for Inc activity, consistent with the

properties of a substrate recruitment domain. Mutation of a conserved C-terminal arginine

abolishes Inc function, suggesting that this residue is vital for recruiting Inc targets. Mutation of

the same residue in the human Inc ortholog KCTD17 is associated with myoclonic dystonia [37],

consistent with a critical and evolutionarily conserved function for this residue in Inc orthologs.

Our results reveal that Cul3-Inc complexes impact sleep and synaptic function, and provide tools

to identify targets of Cul3, Inc, and Inc orthologs that impact normal brain function and

neurological disorders.

Results

The Inc BTB domain mediates Cul3 binding and Inc multimerization

Cul3 adaptors have a modular functional organization, with BTB domains that self-

associate and bind Cul3, and distal domains that recruit substrates [21–24]. Inc has two conserved

domains, an N-terminal BTB domain and a C-terminal domain unique to Inc and its orthologs

[17,38,39] (Fig 1A and S1A Fig). To determine whether Inc has the properties of a Cul3 adaptor,

we generated progressive N- and C-terminal deletions of Inc (Fig 1A) and assessed whether these

derivatives could bind Inc and Cul3, by expressing epitope-tagged forms of these proteins in

Drosophila S2 cells and performing co-immunoprecipitations.

Deleting Inc residues preceding the BTB domain (Inc$^{22-211}$) did not alter Inc stability or

Inc-Cul3 interactions (Fig 1B and S2 Fig), indicating that the N-terminus of Inc is dispensable for

Cul3 binding. In contrast, removing part of the BTB domain (Inc$^{31-211}$) or the entire BTB domain

(Inc$^{124-211}$) significantly destabilized Inc (S2 Fig), and low steady-state levels of these proteins

precluded conclusions about their ability to bind Cul3. Inc proteins lacking 25 C-terminal residues

(Inc$^{1-186}$) or all residues following the BTB domain (Inc$^{1-123}$) were stable and associated with Cul3

similarly to full-length Inc. An Inc mutant lacking most of the Inc C-terminal domain (Inc$^{1-156}$)

was stable and interacted with Cul3 more strongly than wild-type Inc and other Inc deletions (S3A

Fig). Thus, the Inc C-terminus is dispensable for interaction with Cul3. The Inc BTB domain alone

(Inc$^{22-123}$) was sufficient to bind Cul3, albeit more weakly than wild-type Inc (S3B Fig); because

deleting the same N-terminal residues from full length Inc slightly weakened Cul3 binding (Fig

1B; Inc$^{22-211}$ versus Inc$^{1-211}$), these results suggest that Inc N-terminal residues stabilize Cul3

interactions mediated chiefly by the Inc BTB domain.

Cul3 adaptors homomultimerize through their BTB domains, and adaptor-mediated

oligomerization of Cul3 complexes can enhance substrate ubiquitination [40]. We therefore

assessed whether Inc self-association is mediated by its BTB domain. N-terminal truncations that

preserve the Inc BTB domain (Inc[22-211]) did not alter Inc-Inc associations (Fig 1C). Similarly, Inc

proteins truncated C-terminally (Inc[1-186], Inc[1-156], and Inc[1-123]) associated with Inc normally,

indicating that the Inc C-terminus is dispensable for Inc multimerization (Fig 1C). The Inc BTB

domain (Inc[22-123]) alone was sufficient to bind Inc (Fig 1C). Thus, the Inc BTB domain is sufficient

to mediate both Inc multimerization and Cul3 binding, as expected of a Cul3 adaptor.

Inc point mutants define a surface of the Inc BTB domain that binds Cul3

To determine whether Inc function in vivo requires binding to Cul3 and Inc

homomultimerization, we sought to identify Inc point mutants that selectively perturb Inc-Cul3

and Inc-Inc interactions. Our mutagenesis of Inc was informed by the structures of homodimeric

BTB-MATH and BTB-BACK-Kelch adaptors bound to Cul3 [40–42], and by the structure of

KCTD5, a human Inc ortholog that forms homopentamers [39]. Homology modeling and light

scattering measurements suggest that Inc forms a homopentamer similar to that of KCTD5 [43,44].

We selected eight conserved residues in the Inc BTB domain whose equivalents in KCTD5 reside

on a surface suggested to bind Cul3 [43,45] (S1A Fig). We mutated these residues to alanine

(F47A, D57A, D61A, F105A, N107A), to oppositely charged residues (R50E, E104K), or altered

their hydrophobicity (Y106F), and assessed the effects on Inc-Cul3 and Inc-Inc binding. Most of

these mutants did not alter Inc-Cul3 or Inc-Inc interactions, while one mutant (R50E) impaired

both associations (S1 Table). Two mutants, F47A and F105A (Fig 2A), weakened interactions

with Cul3 (Fig 2B) but did not significantly affect Inc self-association or stability (Fig 2C and S4

Fig). Similarly, an F47A/F105A double mutant was stable and selectively impaired Cul3 binding

(Fig 2B and 2C and S4 Fig). The analogous phenylalanines in KCTD5 cluster near the interface

of adjacent subunits, defining a surface for Cul3 binding (S1B Fig).

To identify Inc mutants that impair its multimerization, we selected seven conserved

residues whose equivalents in KCTD5 engage in polar and charged interactions between adjacent

subunits [39] (S1A and S1C Fig). Mutating these residues individually or in double mutant

combinations to alanine (T36A, D71A, D73A, N82A) or to oppositely charged residues (R85E,

K88D, E101K) did not significantly alter Inc self-association (S2 Table). A triple mutant

(T36A/D71A/R85E) predicted to disrupt interactions of Inc subunits with both flanking neighbors

in a putative Inc homopentamer exhibited significantly reduced stability (S4 Fig and S2 Table),

precluding conclusions about its ability to multimerize and bind Cul3. Because this triple mutant

was also unstable when expressed in vivo, we focused further analysis on Inc mutants that

selectively perturb Cul3 interactions.

Inc function in vivo requires the assembly of Inc-Cul3 complexes

To determine whether Inc impacts sleep as a Cul3 adaptor, we assessed whether Inc

mutants that weaken Cul3 binding impair Inc activity in vivo. We generated UAS transgenes

expressing 3×FLAG-tagged Inc, Inc^{F47A}, Inc^{F105A}, and Inc$^{F47A/F105A}$, integrating them at the same

genomic site to compare their activity. When expressed with *inc-Gal4*, a driver that rescues *inc*

mutants when used to restore *inc* expression [17,33], Inc and Inc mutants were similarly stable in

both *inc^1* and *inc$^+$* animals (Fig 3A and S5 Fig). Because Inc mutants deficient for Cul3 binding

might sequester substrates and function as dominant negatives, we tested whether these mutants

could alter sleep when expressed in neurons. Neuronal expression of Inc^{F47A}, Inc^{F105A}, or

Inc$^{F47A/F105A}$ did not alter sleep (S6A Fig), in contrast to neuronal *inc* RNAi [17,18]. Similarly,

expression of these mutants using *inc-Gal4* did not perturb sleep (S6D Fig). Thus, these Inc

mutants do not antagonize endogenous Inc, possibly because they retain some ability to bind Cul3

(Fig 2B). Finally, we tested whether Inc function in vivo requires the assembly of Inc-Cul3

complexes. If so, Inc mutants that weaken Cul3 binding should impair *inc* rescue. Whereas wild-

type Inc expressed with *inc-Gal4* fully rescued daily sleep duration in *inc¹* mutants, Inc^{F47A}

provided only a partial rescue, and Inc^{F105A} and $Inc^{F47A/F105A}$ failed to restore a significant amount

of sleep (Fig 3B and 3C), indicating that Inc activity requires binding to the Cul3 complex. For

other sleep parameters, rescue was more strongly impaired for Inc^{F105A} and $Inc^{F47A/F105A}$ than for

Inc^{F47A} (S7A–D Fig), suggesting that F105 is particularly critical for Cul3 binding in vivo. All

rescuing activity of UAS transgenes was Gal4-dependent, as indicated by the short sleep of *inc¹*

mutants bearing only UAS transgenes (S8 Fig). Collectively, these results indicate that Inc function

in vivo requires the assembly of Inc-Cul3 complexes and that their disruption inhibits sleep.

The Inc C-terminus is vital for Inc activity in vivo and has the properties of a substrate

recruitment domain

Although the Inc C-terminus is dispensable for Inc-Inc and Inc-Cul3 associations (Fig 1B

and 1C), its evolutionary conservation suggests a critical function. The modular structure of Cul3

adaptors suggests that if Inc serves as a Cul3 adaptor, its C-terminus would bind and recruit

substrates to Inc-Cul3 complexes. We therefore assessed the activity of C-terminally truncated Inc

mutants in vivo. 3×FLAG-tagged Inc^{1-186}, Inc^{1-156}, and Inc^{1-123} were expressed at similar levels

under *inc-Gal4* control (Fig 4A), indicating that they are stable in vivo, as in cultured cells (Fig

1B and 1C). Expression of C-terminally truncated Inc mutants neuronally with *elav-Gal4* or with

inc-Gal4 did not alter sleep, indicating that they do not inhibit endogenous *inc* function (S6B Fig

and S6D Fig). We next tested whether C-terminally truncated Inc proteins could restore sleep to

inc mutants. Wild-type Inc expressed with *inc-Gal4* fully rescued sleep in *inc¹* mutants (Fig 4B

and 4C). In contrast, C-terminal truncations of Inc curtailed or abolished Inc function. Inc^{1-186}

partially rescued *inc* mutants, while animals expressing Inc^{1-156} or Inc^{1-123} slept indistinguishably

from *inc* mutants, indicating that Inc$^{1-156}$ and Inc$^{1-123}$ lacked activity in vivo (Fig 4B and 4C and

S9A–D Fig). Thus, the Inc C-terminus is vital for Inc function. The ability of C-terminally

truncated Inc proteins to multimerize and bind Cul3 normally (Fig 1B and 1C) is consistent with

a modular function for the Inc C-terminus in recruiting targets that impact sleep.

Inc-Cul3 binding and the Inc C-terminus are vital for Inc activity at synapses

We next assessed whether Inc-Cul3 associations and the Inc C-terminus are required for

Inc function at synapses. At the larval neuromuscular junction, Inc is required for the homeostatic

upregulation of presynaptic neurotransmitter release caused by inhibition of postsynaptic

glutamate receptors [20]. Application of philanthotoxin, a glutamate receptor antagonist, reduces

the amplitude of spontaneous neurotransmission, while homeostatic upregulation of evoked

neurotransmission maintains synaptic strength in wild-type animals (Fig 5A and 5B) [46]. Synaptic

homeostasis is blocked in *inc* mutants and is restored by Inc expressed post-synaptically in muscle

with *BG57-Gal4* [20] (Fig 5A and 5B). We assessed whether Inc^{F105A}, which impairs Cul3

associations and rescue of sleep (Figs 2B, 3B, and 3C), could restore synaptic homeostasis to *inc*

mutants. In contrast to wild-type Inc, post-synaptic expression of Inc^{F105A} failed to restore synaptic

homeostasis to Inc mutants, indicating that Inc-Cul3 associations are required for Inc activity at

synapses (Fig 5B). To determine whether the activity of Cul3-Inc complexes requires the Inc C-

terminus, we similarly assessed the rescuing activity of Inc$^{1-123}$, which lacks this conserved

domain. Inc$^{1-123}$ failed to restore synaptic homeostasis to *inc* mutants (Fig 5A and 5B), indicating

that the C-terminus of Inc is essential for Inc activity at synapses. The requirement for Inc-Cul3

associations and the Inc C-terminus for both normal sleep and synaptic plasticity supports the

notion that Inc serves as a Cul3 adaptor in vivo.

Mutation of a conserved disease-associated arginine abolishes Inc function and defines a surface likely to recruit substrates

To further dissect Inc function, we mutated conserved C-terminal residues we hypothesized might bind Inc targets (Fig 6A). In particular, a missense mutation of a conserved arginine (R145H) in the human Inc ortholog KCTD17 is associated with myoclonic dystonia, a movement disorder [37]. While the mechanism of KCTD17^{R145H} pathogenesis is unknown, the analogous KCTD5 residue resides on the surface of the C-terminal domain (S1D Fig). We reasoned that this residue might have a conserved function in binding substrates, and we therefore mutated the corresponding Inc residue (Inc^{R135H}) to assess its function. We also mutated three Inc residues (N167, Y168, G169) to alanine (Inc^{AAA}); the corresponding KCTD5 residues reside in an exposed loop that might recruit substrates (S1D Fig).

In S2 cells, Inc^{R135H} and Inc^{AAA} were stable and associated with Cul3 and Inc indistinguishably from wild-type Inc (Figs 6B and 6C), indicating that these mutations do not alter Inc multimerization or Cul3 binding. In vivo, Inc^{R135H} and Inc^{AAA} were stably expressed with *inc-Gal4* (Fig 6D) and lacked dominant negative activity when expressed in neurons or with *inc-Gal4* (S6C–D Fig). Next, we tested whether Inc^{R135H} and Inc^{AAA} could rescue the sleep deficits of *inc* mutants. When expressed with *inc-Gal4*, Inc^{AAA} rescued *inc^1* mutants indistinguishably from wild-type Inc, indicating that Inc^{AAA} retains normal activity (Fig 6E and 6F and S8 Fig and S10A–D Fig). In stark contrast, Inc^{R135H} was unable to restore sleep to *inc^1* mutants, indicating that the R135H mutation abolishes Inc function (Fig 6E and 6F and S10A–D Fig). The complete loss of Inc activity caused by the R135H mutation is indistinguishable from the consequences of deleting the Inc C-terminus (Fig 4B and 4C), suggesting that R135 is essential for the function of the C-terminus and binding Inc targets. These findings furthermore suggest that the analogous

KCTD17^{R145H} mutation causes myoclonic dystonia by impairing the recruitment and

ubiquitination of KCTD17 targets.

Inc exhibits properties of a Cul3 adaptor in neurons and is negatively regulated by Cul3

Normal sleep requires the activity of both *inc* and *Cul3* in neurons [17,18]. We performed

several experiments to determine whether Inc has the properties of a Cul3 adaptor in vivo. First,

we tested whether Inc-Cul3 complexes assemble in vivo, by co-expressing epitope-tagged Inc and

Cul3 with *inc-Gal4* and performing co-immunoprecipitations. We observed that Inc and Cul3

associate in vivo (Fig 7A). Second, we tested whether Inc multimerizes in vivo, by assessing the

physical interactions of Inc-HA and Myc-Inc co-expressed with *inc-Gal4*. These proteins

associated robustly (Fig 7B), indicating that Inc forms homomultimers in vivo. Third, we tested

whether Inc is regulated by Cul3. Cul3 adaptors are often regulated by autocatalytic ubiquitination

and degradation in Cul3 complexes [22,47–49]. Neuronal RNAi against *Cul3* increased Inc protein

levels in head lysates (Fig 7C), indicating that Cul3 negatively regulates Inc abundance. qRT-PCR

confirmed that neuronal Cul3 RNAi reduced Cul3 mRNA levels in heads (Fig 7D), as reported

previously [50], but did not significantly alter *inc* transcript levels (Fig 7D). These findings

indicate that Cul3 negatively regulates Inc by a post-transcriptional mechanism. We conclude that

Inc is likely regulated by Cul3-dependent autocatalytic ubiquitination and degradation in neurons,

cells through which *inc* influences sleep [17,18]. Together, these results indicate that Inc has the

attributes of a Cul3 adaptor in vivo.

Discussion

Cul3 mutations disrupt brain development and physiology and contribute to neurological

disorders, including autism and schizophrenia. Understanding the diverse functions of *Cul3* in the

nervous system requires characterization of its relevant adaptors and substrates. Our findings

implicate Inc as a Cul3 adaptor critical for sleep and synaptic function. We have shown that Inc-

Cul3 complexes assemble in vivo, that weakening Inc-Cul3 associations impairs Inc activity, and

that Inc has modular functional domains typical of Cul3 adaptors. The conserved C-terminal

domain of Inc is vital for Inc activity but dispensable for Inc multimerization and Cul3 binding, as

expected of a substrate recruitment domain. Lastly, the negative regulation of Inc abundance by

Cul3 in vivo is characteristic of autocatalytic adaptor ubiquitination within the Cul3 complex and

subsequent proteolytic degradation.

In addition to demonstrating that Inc is regulated by autocatalytic ubiquitination, the

present findings suggest that Inc activity is controlled by additional mechanisms. Inc mutants

deficient for Cul3 binding would be expected to sequester substrates but do not perturb sleep when

overexpressed. Similarly, C-terminally truncated Inc mutants that cannot recruit substrates might

be expected to compete for Cul3 binding and inhibit ubiquitination of Inc substrates, yet these

mutants lack dominant negative activity. Consistent with these findings, overexpression of wild-

type Inc rescues *inc* mutants in a permissive manner but does not cause gain-of-function

phenotypes, for both sleep and synaptic homeostasis [17,20,33]. Inc may thus be regulated by

mechanisms that control its ability to bind Cul3, recruit substrates, or catalyze substrate

ubiquitination.

Inc and its orthologs are structurally distinct from better characterized Cul3 adaptors of the

BTB-MATH and BTB-BACK-Kelch families. In contrast to BTB-MATH and BTB-BACK-Kelch

adaptors that form homodimers and bind Cul3 independently [40–42], Inc orthologs form

homopentamers that bind Cul3 in radial assemblies [39,43,45,51–53]. Our findings indicate that

Cul3 binds to a cleft between adjacent Inc subunits, consistent with a similar homopentameric

structure for Cul3-Inc complexes. Because formation of higher order Cul3 complexes can

stimulate substrate ubiquitination [40], the pentameric structure of Inc and its orthologs may enable

efficient ubiquitination of substrates, including those that associate weakly or transiently with the

Cul3 complex.

The analysis of C-terminal Inc truncations and point mutations suggests a mechanism by

which Inc and its orthologs bind substrates. Deleting the 25 C-terminal residues of Inc reduces but

does not abolish Inc activity, indicating that these residues contribute to substrate binding. The

analogous KCTD5 residues are not resolved in crystal structures of KCTD alone [39] (Fig. S1D),

suggesting that they are flexible and may become structured upon target binding. Second, mutation

of a surface-exposed arginine (R135) abolishes Inc function and is functionally equivalent to

deleting the C-terminus. Together, these data support a model in which R135 and the surrounding

lateral surface of Inc bind substrates in conjunction with the C-terminal tail of Inc. A recent

structure of KCTD5 bound to a substrate provides direct support for this mechanism [53].

Furthermore, this mechanism may be conserved in other Inc orthologs and disrupted by

KCT17^{R145H} mutations linked to myoclonic dystonia [37]. While the simplest hypothesis is that

Inc binds substrates directly, Inc might also engage co-adaptor proteins, analogous to the

mechanism described for the BTB-Kelch adaptor KLHL12 [54].

Our findings do not identify Inc targets that mediate its impact on sleep and synaptic

homeostasis in *Drosophila*, nor do they discern whether Inc orthologs contribute to changes in

sleep or other *Cul3* phenotypes in mammals. Another important unresolved question is whether

heterozygous loss-of-function *Cul3* mutations alter sleep in rodent models, analogous to sleep

disturbances reported for human *Cul3* mutations associated with autism [4]. While Inc orthologs

are expressed in the mammalian brain [33], their characterization in vivo remains limited. In mice,

KCTD5 deletions are homozygous lethal and cause motor deficits when heterozygous [55];

possible effects on sleep have not yet been assessed. There are no existing rodent models for

mutations in *KCTD2* or *KCTD17*. Conditional genetic analysis of *inc* in flies and *Cul3* in mice

underscores developmental mechanisms of action. Inc acts in developing neurons to impact sleep

in adult flies [56], suggesting that its relevant substrates have developmental functions within the

brain. Conditional deletion of *Cul3* in mice indicates that its behavioral phenotypes, including

deficits in sociability and cognition, arise during a critical period of brain development [10],

suggesting that any sleep phenotypes of *Cul3* mutants might also arise in a developmental manner.

The molecular targets of Inc that influence sleep are likely conserved, as mammalian Inc

orthologs restore sleep to *inc* mutants [33]. Intriguing overlaps in the synaptic phenotypes of *inc*

and *Cul3* mutants in flies and mice suggest that Inc orthologs impact synaptic function through

substrates that are similarly conserved. Inc and Cul3 are required in flies for synaptic homeostasis

at glutamatergic neuromuscular synapses and are recruited to synapses during homeostatic

potentiation [20]. *Cul3* mutations in mice alter glutamatergic neurotransmission and the abundance

of glutamate receptors and other synaptic proteins [9,11]. While contributions of mammalian Inc

orthologs to *Cul3* phenotypes at glutamatergic synapses remain unexplored, Cul3 and Inc

orthologs are present in synaptosome preparations [33], suggesting that their targets include

synaptic proteins.

Characterization of Inc orthologs has identified various substrates in cultured cells. c-Myc

is a substrate for KCTD2 [57], and KCTD5 ubiquitinates ΔNp63α and RhoGDI [58,59], linking

KCTD2 and KCTD5 to pathways that regulate cell proliferation. KCTD5 has also been shown to

ubiquitinate a G-protein beta subunit and to regulate AKT and cAMP signaling, including in

cultured striatal neurons [55,60,61]. KCTD17 ubiquitinates trichoplein, a regulator of primary cilia

[62], as well as Chop and PHLPP2 [63,64], proteins that regulate lipids and adipogenesis. Whether

these targets or others mediate the impact of Inc on sleep and synapses remains unknown.

Emerging evidence from genetic studies in humans suggests that Inc orthologs have

functions in the brain that remain poorly understood. As noted above, *KCTD17* mutations are

associated with myoclonic dystonia [37]. Polymorphisms near the *KCTD2* locus are associated

with Alzheimer's [65], and another study suggests an association of *KCTD5* and daytime

sleepiness [66]. Elucidating targets of Cul3-Inc complexes within the fly brain may provide a

powerful system to understand conserved mechanisms through which Cul3 and Inc contribute to

nervous system function and neurological disorders.

Acknowledgements

We thank J. Lopez for assistance with Western blots and qRT-PCR; Z. Zuchowski for assistance in constructing plasmids; A. Arzeno and A. Gonzalez for assistance with immunoprecipitations; and N. Ringstad, J. Treisman, and members of the Stavropoulos laboratory for comments on the manuscript. This work was supported by an International Student Research Fellowship from the Howard Hughes Medical Institute (HHMI) to Q.L., by a grant from the National Institutes of Health (NS091546) to D.D., and by grants from the National Institutes of Health (NS112844 and NS111304), the Mathers Foundation, Whitehall Foundation grant 2013-05-78, fellowships from the Alfred P. Sloan and Leon Levy Foundations, a NARSAD Young Investigator Award from the Brain and Behavior Foundation, the J. Christian Gillin, M.D. Research Award from the Sleep Research Society Foundation, and a Career Scientist Award from the Irma T. Hirschl/Weill-Caulier Trust to N.S.

Author Contributions

Conceptualization: Q.L., D.D., N.S.

Investigation: Q.L., K.Y.L., R.A., F.V., C.D., J.M., X.L., and N.S.

Resources: Q.L., D.D., and N.S.

Writing - original draft: N.S. and Q.L.

Writing - review and editing: N.S., Q.L., and D.D.

Competing Interests

The authors declare that no completing interests exist.

Materials and methods

Plasmids and molecular cloning

Vectors for expression in S2 cells were as follows:

pAc5.1–Inc-HA (pNS277) encodes Inc fused to a C-terminal HA epitope (GSYPYDVPDYA) and was generated by ligating EcoRI-XhoI digested pAc5.1/V5-HisA backbone (ThermoFisher) and an EcoRI-XhoI Inc-HA fragment from pNS273 (described below).

pAc5.1–3×Myc-Inc (pNS351) was previously described (Li et al., 2017) and encodes an N-terminal 3×Myc epitope (MEQKLISEEDLGSEQKLISEEDLGSEQKLISEEDLAS) fused to Inc.

pAc5.1–3×Myc-Inc$^{22\text{-}211}$ (pNS370) was generated by ligating NheI-XhoI digested pNS309 (Stavropoulos and Young, 2011) and the NheI-XhoI fragment liberated from the PCR amplification product of pNS351 template and primers oNS315 and oNS285.

pAc5.1–3×Myc-Inc$^{31\text{-}211}$ (pNS371) was generated similarly to pNS370, substituting primers oNS316 and oNS285.

pAc5.1–3×Myc-Inc$^{124\text{-}211}$ (pNS372) was generated similarly to pNS370, substituting primers oNS317 and oNS285.

pAc5.1–3×Myc-Inc$^{1\text{-}123}$ (pNS374) was generated similarly to pNS370, substituting primers oNS277 and oNS319.

pAc5.1–3×Myc-Inc$^{1\text{-}156}$ (pNS375) was generated similarly to pNS370, substituting primers oNS277 and oNS320.

pAc5.1–3×Myc-Inc$^{1\text{-}186}$ (pNS376) was generated similarly to pNS370, substituting primers oNS277 and oNS321.

pAc5.1–3×Myc-Inc$^{22\text{-}123}$ (pNS377) was generated similarly to pNS370, substituting primers oNS315 and oNS319.

pAc5.1–3×HA-Inc   (pNS402)   encodes   an   N-terminal   3×HA   epitope

(MYPYDVPDYAGSYPYDVPDYAGSYPYDVPDYAAS) fused to Inc and was generated by

ligating NheI-XhoI digested pNS310 (Stavropoulos and Young, 2011) and an NheI-XhoI *inc*

fragment liberated from pNS351.

pAc5.1–3×FLAG-Cul3   (pNS403)   encodes   an   N-terminal   3×FLAG   epitope

(MDYKDDDDKGSDYKDDDDKGSDYKDDDDKAS) fused to *Drosophila* Cul3 and was

generated by ligating NheI-NotI digested pNS311 and an NheI-NotI Cul3 fragment liberated from

pNS314 (Stavropoulos and Young, 2011). pNS311 contains an N-terminal 3×FLAG tag and was

generated from pNS298, a pAc5.1/V5-HisA derivative that contains a C-terminal 3×FLAG tag.

To construct pNS298, oligonucleotides oNS234 and oNS235 were phosphorylated, annealed, and

cloned into XhoI-XbaI digested pAc5.1/V5-HisA. To construct pNS311, EcoRI-NotI digested

pAc5.1/V5-HisA was ligated to the EcoRI-NotI fragment liberated from the PCR amplification

product of pNS298 template and primers oNS240 and oNS241.

pAc5.1–3×FLAG-Inc   (pNS408)   encodes   an   N-terminal   3×FLAG   epitope

(MDYKDDDDKGSDYKDDDDKGSDYKDDDDKAS) fused to Inc and was generated by

ligating NheI-NotI digested pNS311 and an NheI-XhoI *inc* fragment liberated from pNS351.

pAc5.1–3×HA-Inc^{F47A} (pNS409)   encodes   an   N-terminal   3×HA   epitope

(MYPYDVPDYAGSYPYDVPDYAGSYPYDVPDYAAS) fused to Inc^{F47A} and was generated by

ligating NheI-XhoI digested pNS310 and an NheI-XhoI digested Inc^{F47A} fragment. The Inc^{F47A}

fragment was prepared by fusion PCR in two steps. First, overlapping 5' and 3' Inc^{F47A} fragments

were generated by PCR amplification of pNS408 template with primers oNS277/oNS695 and

oNS684/oNS694, respectively. An equimolar mix of these fragments was then used as template

for fusion PCR amplification with primers oNS277 and oNS684.

pAc5.1–3×HA-Inc^{R50E} (pNS410) was generated similarly to pNS409, substituting primers

oNS277/oNS697 and oNS684/oNS696 to obtain 5' and 3' Inc^{R50E} fragments, respectively.

pAc5.1–3×HA-Inc^{D57A} (pNS411) was generated similarly to pNS409, substituting primers

oNS277/oNS699 and oNS684/oNS698 to obtain 5' and 3' Inc^{D57A} fragments, respectively.

pAc5.1–3×HA-Inc^{D61A} (pNS412) was generated similarly to pNS409, substituting primers

oNS277/oNS701 and oNS684/oNS700 to obtain 5' and 3' Inc^{D61A} fragments, respectively.

pAc5.1–3×HA-Inc^{E104K} (pNS413) was generated similarly to pNS409, substituting primers

oNS277/oNS703 and oNS684/oNS702 to obtain 5' and 3' Inc^{E104K} fragments, respectively.

pAc5.1–3×HA-Inc^{F105A} (pNS419) was generated similarly to pNS409, substituting primers

oNS277/oNS1123 and oNS684/oNS1122 to obtain 5' and 3' Inc^{F105A} fragments, respectively.

pAc5.1–3×HA-Inc^{Y106F} (pNS420) was generated similarly to pNS409, substituting primers

oNS277/oNS1146 and oNS684/oNS1145 to obtain 5' and 3' Inc^{Y106F} fragments, respectively.

pAc5.1–3×HA-Inc^{N107A} (pNS421) was generated similarly to pNS409, substituting primers

oNS277/oNS1125 and oNS684/oNS1124 to obtain 5' and 3' Inc^{N107A} fragments, respectively.

pAc5.1–3×FLAG-Inc^{D73A} (pNS430) and pAc5.1–3×FLAG-Inc^{N82A} (pNS431) were

generated simultaneously, using PCR amplification of pNS408 template with primers

oNS277/oNS686 and oNS684/oNS685 to generate overlapping Inc^{N82A} and Inc^{D73A} fragments,

respectively. An equimolar mix of these fragments was used as template for fusion PCR

amplification with primers oNS277 and oNS684, and the PCR product was digested with NheI-

XhoI and ligated to NheI-XhoI digested pNS311. Sequencing of multiple clones identified both

single mutants.

pAc5.1–3×FLAG-Inc^{K88D} (pNS432) and pAc5.1–3×FLAG-Inc^{E101K} (pNS433) were

generated similarly to pNS430 and pNS431, substituting primers oNS277/oNS688 and

oNS684/oNS687 to obtain overlapping Inc^{E101K} and Inc^{K88D} fragments, respectively.

pAc5.1–3×FLAG-Inc^{T36A} (pNS434) and pAc5.1–3×FLAG-Inc^{D71A} (pNS435) were

generated similarly to pNS430 and pNS431, substituting primers oNS277/oNS693 and

oNS684/oNS689 to obtain overlapping Inc^{D71A} and Inc^{T36A} fragments, respectively.

pAc5.1–3×HA-Inc^{R85E} (pNS436) was generated similarly to pNS430 and pNS431,

substituting primers oNS277/oNS692 and oNS684/oNS691 to obtain overlapping Inc^{R85E} and

Inc^{D71A} fragments, respectively, and NheI-XhoI digested pNS310. Inc^{R85E} was identified by

sequencing multiple clones.

pAc5.1–3×HA-Inc$^{F47A/F105A}$ (pNS422) was generated similarly to pNS409, substituting

pNS409 as template and primers oNS277/oNS1123 and oNS684/oNS1122 to obtain 5' and 3'

Inc$^{F47A/F105A}$ fragments, respectively.

pAc5.1–3×HA-Inc$^{D73A/N82A}$ (pNS414) was generated similarly to pNS409, substituting

pNS430 template and primers oNS277/oNS686 to obtain the 5' Inc$^{D73A/N82A}$ fragment and pNS431

template and primers oNS684/oNS685 to obtain the 3' Inc$^{D73A/N82A}$ fragment.

pAc5.1–3×HA-Inc$^{K88D/E101K}$ (pNS415) was generated similarly to pNS409, substituting

pNS432 template and primers oNS277/oNS688 to obtain the 5' Inc$^{K88D/E101K}$ fragment and pNS433

template and primers oNS684/oNS687 to obtain the 3' Inc$^{K88D/E101K}$ fragment.

pAc5.1–3×HA-Inc$^{T36A/D71A}$ (pNS416) was generated similarly to pNS409, substituting

pNS434 template and primers oNS277/oNS693 to obtain the 5' Inc$^{T36A/D71A}$ fragment and pNS435

template and primers oNS684/oNS689 to obtain the 3' Inc$^{T36A/D71A}$ fragment.

pAc5.1–3×HA-Inc$^{D71A/R85E}$ (pNS417) was generated similarly to pNS409, substituting

pNS435 template and primers oNS277/oNS692 to obtain the 5' Inc$^{D71A/R85E}$ fragment and pNS436

template and primers oNS684/oNS691 to obtain the 3' Inc $^{D71A/R85E}$ fragment.

pAc5.1–3×HA-Inc$^{T36A/D71A/R85E}$ (pNS418) was generated by ligating EcoRV-XhoI

digested pNS434 backbone and an Inc$^{D71A/R85E}$ fragment obtained by EcoRV-XhoI digestion of

the PCR amplification product of pNS417 template and primers oNS277 and oNS684.

pAc5.1–3×HA-Inc^{R135H} (pNS426) was generated by ligating HindIII-XhoI digested

pNS402 backbone and a HindIII-XhoI Inc^{R135H} fragment. The Inc^{R135H} fragment was prepared by

fusion PCR in two steps. First, overlapping 5' and 3' Inc^{R135H} fragments were generated by PCR

amplification of pNS346 template with primers oNS1126/oNS1555 and oNS1127/oNS1554,

respectively. An equimolar mix of these fragments was then used as template for fusion PCR

amplification with primers oNS1126 and oNS1127. pUASTattB–Myc-Inc (pNS346) encodes an

N-terminal Myc epitope (MEQKLISEEDLAS) fused to Inc, as previously described (Li et al.,

2017).

pAc5.1–3×HA-Inc^{AAA} (pNS427) was generated similarly to pNS426, substituting a

HindIII-XhoI Inc^{AAA} fragment generated similarly with fusion PCR; primer pairs

oNS1126/oNS1557 and oNS1127/oNS1556 were used to obtain 5' and 3' Inc^{AAA} fragments,

respectively.

pAc5.1–3×Myc-Inc$^{T36A/D71A/R85E}$ (pNS437) was generated by ligating NheI-XhoI digested

pNS309 backbone and an NheI-XhoI Inc$^{T36A/D71A/R85E}$ fragment liberated from pNS418.

pAc5.1–3×Myc-Inc^{F47A} (pNS438) was generated by ligating NheI-XhoI digested pNS309

backbone and an NheI-XhoI Inc^{F47A} fragment liberated from pNS409.

pAc5.1–3×Myc-Inc^{F105A} (pNS439) was generated by ligating NheI-XhoI digested pNS309

backbone and an NheI-XhoI Inc^{F105A} fragment liberated from p419.

pAc5.1–3×Myc-Inc$^{F47A/F105A}$ (pNS440) was generated by ligating NheI-XhoI digested

pNS309 backbone and an NheI-XhoI Inc$^{F47A/F105A}$ fragment liberated from pNS422.

pAc5.1–3×Myc-Inc^{R135H} (pNS441) was generated by ligating HindIII-XhoI digested

pNS351 backbone and a HindIII-XhoI Inc^{R135H} fragment prepared as for pNS426.

pAc5.1–3×Myc-Inc^{AAA} (pNS442) was generated by ligating HindIII-XhoI digested

pNS351 backbone and a HindIII-XhoI Inc^{AAA} fragment prepared as for pNS427.

Vectors for *Drosophila* transgenesis were as follows:

pUASTattB–Myc-Inc^{R135H} (pNS428) was generated by ligating a BglII-XhoI digested

pUASTattB backbone (Bischof et al., 2007), prepared by liberating an unrelated BglII-XhoI insert,

and a BgII-XhoI digested Myc-Inc^{R135H} fragment, generated with fusion PCR as for pNS426.

pUASTattB–Myc-Inc^{AAA} (pNS429) was generated by ligating a BgII-XhoI pUASTattB

backbone prepared from pNS349 [33] and a BgII-XhoI digested Myc-Inc^{AAA} fragment, generated

with fusion PCR as for pNS427.

pUAST–Inc-HA (pNS273) was previously described and encodes Inc fused to a C-terminal

HA epitope (GS<u>YPYDVPDYA</u>) [33].

pUASTattB–3×FLAG-Inc (pNS404) encodes an N-terminal 3×FLAG epitope

(M<u>DYKDDDDK</u>GS<u>DYKDDDDK</u>GS<u>DYKDDDDK</u>AS) fused to Inc and was generated by three-

piece ligation of EcoRI-XhoI digested pUASTattB, an EcoRI-NheI 3×FLAG fragment liberated

from the PCR amplification product of pNS311 template and primers ACF and oNS241, and an

NheI-XhoI *inc* fragment liberated from pNS351.

pUASTattB–3×FLAG-Inc$^{1-186}$ (pNS405) was generated similarly to pNS404, substituting

an NheI-XhoI Inc$^{1-186}$ fragment liberated from pNS376.

pUASTattB–3×FLAG-Inc$^{1-156}$ (pNS406) was generated similarly to pNS404, substituting

an NheI-XhoI Inc$^{1-156}$ fragment liberated from pNS375.

pUASTattB–3×FLAG-Inc$^{1-123}$ (pNS407) was generated similarly to pNS404, substituting

an NheI-XhoI Inc$^{1-123}$ fragment liberated from pNS374.

pUASTattB–3×FLAG-Inc^{F47A} (pNS423) was generated similarly to pNS404, substituting

an NheI-XhoI Inc^{F47A} fragment liberated from pNS409.

pUASTattB–3×FLAG-Inc^{F105A} (pNS424) was generated similarly to pNS404, substituting

an NheI-XhoI Inc^{F105A} fragment liberated from pNS419.

pUASTattB–3×FLAG-Inc$^{F47A/F105A}$ (pNS425) was generated similarly to pNS404,

substituting an NheI-XhoI Inc$^{F47A/F105A}$ fragment liberated from pNS422.

Oligonucleotides

Oligonucleotides used in this work, listed 5' to 3', are as follows:

oNS180   ACGTAGATCTGAACCGCAGCAGCGGCAACACCATC

oNS184   CCAGCCATCCGACAGCGTTGAGATC

oNS234

TCGAGGCTAGCGACTACAAGGATGATGACGATAAGGGCTCCGATTACAAGGA

CGACGATGATAAGGGATCCGATTACAAGGATGATGACGACAAGTGAT

oNS235

CTAGATCACTTGTCGTCATCATCCTTGTAATCGGATCCCTTATCATCGTCGTCC

TTGTAATCGGAGCCCTTATCGTCATCATCCTTGTAGTCGCTAGCC

oNS240

ACTGGAATTCCGCGGCAACATGGACTACAAGGATGATGACGATAAGGGC

oNS241

ACTGGCGGCCGCTCCTAGGGTGCTAGCCTTGTCGTCATCATCCTTGTAATCGG

AT

524	oNS277	ACGTGCTAGCATGAGCACGGTGTTCATAAACTCGC
525	oNS285	ACGTGCTAGCTCGAGGGGTTGTGTGTGAATATATAGCGCGA
526	oNS315	ACGTGCTAGCCAGTGGGTCAAGCTGAACGTAG
527	oNS316	ACGTGCTAGCACCTACTTCCTCACCACAAAGACG
528	oNS317	ACGTGCTAGCCAGCGACCCCAAACGGACAA
529	oNS319	ACGTCTCGAGTTAATCCCTGTGCAGGATGCACTC
530	oNS320	ACGTCTCGAGTTACCTCCAGCCATCCGACAG
531	oNS321	ACGTCTCGAGTTATGTGCCACACTCTTTGGATACCA
532	oNS337	TTTTTTTTTTTTTTTTTTTTTTTTVN
533	oNS684	ACGTCTCGAGGGGTTGTGTGTGAATATATAGCGCGA
534	oNS685	CCTGATCGACAGAGCCCCAAATACTTTGC
535	oNS686	GGCGCAGGTAGGCGAGCACGGGTGCAAA
536	oNS687	TGCGCCACGGCGACCTTGTGCTCGAT
537	oNS688	TAGAACTCAGCCTCCTTCAGGACGCCTTC
538	oNS689	CTACTTCCTCACCGCCAAGACGACGCTC
539	oNS691	GCCTACCTGATCGCCAGAGACCCCAAA
540	oNS692	CAAGCTTGCCGTGTTCCAGGTAATTGAGC
541	oNS693	TTTGGGGTCTCTGGCGATCAGGTAGGC
542	oNS694	GACCCAAATTCGGCCCTCTCCCGTCTG
543	oNS695	CAGACGGGAGAGGGCCGAATTTGGGTC
544	oNS696	TCGTTCCTCTCCGAACTGATTCAGGAGG
545	oNS697	CCTCCTGAATCAGTTCGGAGAGGAACGA
546	oNS698	CAGGAGGACTGCGCCTTGATATCAGATCG
547	oNS699	CGATCTGATATCAAGGCGCAGTCCTCCTG

oNS700   CGACTTGATATCAGCCCGGGACGAGAC

oNS701   GTCTCGTCCCGGGCTGATATCAAGTCG

oNS702   CCTGGAGGAGGCTAAGTTCTACAACGTGAC

oNS703   GTCACGTTGTAGAACTTAGCCTCCTCCAGG

oNS1122   GGAGGCTGAGGCCTACAACGTGAC

oNS1123   GTCACGTTGTAGGCCTCAGCCTCC

oNS1124   GCTGAGTTCTACGCCGTGACGCAGC

oNS1125   GCTGCGTCACGGCGTAGAACTCAGC

oNS1126   CAACTGCAACTACTGAAATCTGCCAAGAAG

oNS1127   GGTAGTTTGTCCAATTATGTCACACCACAGAAG

oNS1145   GAGGCTGAGTTCTTCAACGTGACGCAGC

oNS1146   GCTGCGTCACGTTGAAGAACTCAGCCTC

oNS1554   AAGCGCGTTTATCATGTGCTGCAGTGC

oNS1555   GCACTGCAGCACATGATAAACGCGCTT

oNS1556

563   ATCAGCATGCAGTACACGGCCGCCGCGCCCTTCGAAAACAATGAGTTCCTG

oNS1557   ATTGTTTTCGAAGGGCGCGGCGGCCGTGTACTGCATGCTGATCAGCT

oNS1783   TGTCCACGTACCAGATGTGTGTGCTA

oNS1784   ACTAGCAAGTTGTGAGCCAGACGC

ACF   GACACAAAGCCGCTCCATCAG

RPS3A   CGAACCTTCCGATTTCCAAGAAACGC

RPS3B   ACGACGGACGGCCAGTCCTCC

Cell culture and biochemistry

S2 cells were cultured in S2 media containing 10% FBS, penicillin, and streptomycin, and

were transfected with Effectene (Qiagen) as described previously (Stavropoulos and Young,

2011). Transfections were performed in 6 well plates for ~24 hr, after which liposome-containing

media was replaced with fresh culture media. 400 ng of total DNA was used for each transfection.

For transfections involving two plasmids, an equal amount of each was used. For comparisons of

multiple plasmids, empty vector lacking insert was used to equalize DNA amounts as required.

Cells were harvested 36-48 hr after transfection, washed in PBS, and lysed in ice-cold NP40 lysis

buffer (50 mM Tris pH 7.6, 150mM NaCl, 0.5% NP40) containing protease inhibitors (Sigma,

P8340). Protein extracts were quantitated in duplicate (BioRad, 5000111). For measurements of

protein stability, 100 μg/ml cycloheximide (Fisher, AC357420010) was added at specified

timepoints prior to lysis.

For immunoprecipitations of truncated Inc proteins and Inc point mutants from S2 cells,

300-1000 μg of total protein was incubated overnight at 4°C with 1:100 anti-FLAG (Sigma,

F1804), anti-Myc (Sigma, C3956), or rat anti-Inc antibody [17], followed by incubation with 20

μl (50% slurry) of protein G sepharose beads (ThermoFisher, 10-1243) at 4°C on a nutator for 1

hr or overnight. Alternatively, 20 μl (50% slurry) of anti-FLAG (Sigma, F2426) or anti-Myc

affinity gel (Sigma, E6654) was used to immunoprecipitate samples. Samples were washed 4×5

min at 4°C with NP40 lysis buffer, denatured in SDS sample buffer, separated on Tris SDS-PAGE

gels, and transferred to nitrocellulose.

Protein extracts were prepared from whole flies or sieved heads by manual pestle

homogenization in ice-cold NP40 lysis buffer containing protease inhibitors. To assess in vivo

expression of Inc proteins, 30 μg of lysate was separated on Tris-SDS-PAGE gels. For

immunoprecipitations of whole fly extracts, 800 μg of protein was incubated with 30 μl (50%

slurry) of anti-Myc affinity gel for 1.5 hr at 4°C on a nutator. Samples were washed, denatured,

electrophoresed, and transferred to membranes as described above.

Western blots and antibodies

Membranes were incubated for 60-90 min at room temperature in blocking buffer (LI-

COR, 927-40000). Membranes were subsequently incubated for 1-2 hr at room temperature or 4°C

overnight in blocking buffer containing 0.1% Tween 20 and the appropriate primary antibodies:

rabbit anti-Myc (1:2,000, Sigma, C3956), mouse anti-FLAG (1:2,000, Sigma, F1804), rat anti-HA

(1:2,000, Roche, 11867431001), and rabbit anti-HA (1:2,000, Bethyl Laboratories, A190-208A),

mouse anti-tubulin (1:10,000, DSHB, 12G10), and rabbit anti-tubulin (1:60,000, VWR, 89364-

004). Membranes were washed 4×5 min in TBST (150 mM NaCl, 10mM Tris pH 7.6, and 0.1%

Tween 20) and subsequently incubated in the dark for 30-60 min at room temperature in blocking

buffer containing 0.1% Tween 20, 0.01% SDS, and appropriate secondary antibodies, all diluted

1:15,000 or 1:30,000: Alexa 680 donkey anti-rabbit (Life Technologies, A10043), Alexa 790 anti-

rabbit (Life Technologies, A11374), Alexa 680 donkey anti-mouse (Life Technologies, A10038),

Alexa 790 anti-mouse (Life Technologies, A11371), Alexa 790 anti-rat (Jackson

ImmunoResearch, 712-655-153), and Alexa 680 donkey anti-rat (Jackson ImmunoResearch, 712-

625-1533). Membranes were then washed 4×5 min in TBST, 1×5 min in TBS, and imaged on a

Li-Cor Odyssey CLx instrument.

qRT-PCR

Total RNA was isolated from adult male fly heads with TRIZOL (ThermoFisher, 15596-

026). 5 µg of RNA was reverse transcribed with $d(T)_{23}VN$ primer (oNS337) and SuperScript II

reverse transcriptase (ThermoFisher, 18064014). qPCR was performed using a BioRad DNA

Engine Opticon 2 System, SYBR Green Supermix (Bio-Rad, 1725272), and the following primers:

oNS180 and oNS184 (*inc*); oNS1783 and oNS1784 (*Cul3*); RPS3A and RPS3B (*rps3*).

Fly stocks and transgenes

elav^{c155}-Gal4 [67], *inc^1* [17], *inc^{kk3}* [20], *inc-Gal4* [17], *inc^1 inc-Gal4* [33], *BG57-Gal4*

[68], *attP2: UAS-Myc-Inc* [33], *attP2: UAS-3×FLAG-Inc* [33], and *UAS-3×FLAG-3×HA-Cul3*

[69] were previously described. *UAS-Cul3-RNAi* (11861R-2) and *UAS-inc-RNAi* (18225) were

obtained from the NIG-Fly and VDRC stock centers, respectively. Transgenic flies generated in

this study using pUASTattB-based vectors were integrated at *attP2* [70] with phiC31 recombinase

(BestGene). All transgenes were backcrossed six to eight generations to Bloomington stock 5905,

an isogenic *w^{1118}* stock described elsewhere as iso31 [71].

Sleep analysis

Crosses were set with five virgin females and three males on cornmeal, agar, and molasses

food. One- to four-day old male flies eclosing from LD-entrained cultures raised at 25°C were

loaded in glass tubes containing cornmeal, agar, and molasses food. Animals were monitored for

5-7 days at 25°C in LD cycles using DAM2 monitors (Trikinetics). The first 36-48 hours of data

were discarded and an integral number of days of data (3-5) were analyzed using custom Matlab

code. Locomotor data were collected in 1 min bins. Sleep was defined by locomotor inactivity for

5 min or more; all minutes within inactive periods exceeding 5 min were assigned as sleep. Dead

animals were excluded from analysis by a combination of automated filtering and visual inspection

of locomotor traces. Two to four independent biological replicates were performed for all sleep

assays.

Electrophysiology

Two-electrode voltage clamp (TEVC) recordings were performed in muscle 6 in abdominal segments 2 and 3, as previously described [20]. Muscles were clamped at −70 mV, with a leak current below 5 nA. mEPSCs were recorded for 1 min from each muscle cell in the absence of stimulation. Twenty EPSCs were acquired for each cell under stimulation at 0.5 Hz, using 0.5 ms stimulus duration and with stimulus intensity adjusted with an ISO-Flex Stimulus Isolator (A.M.P.I.). To acutely block postsynaptic receptors, larvae were incubated with philanthotoxin-433 (PhTx; 20 μM; Sigma) in HL-3 for 10 min. Data were analyzed using Clampfit 10.7 (Molecular Devices), MiniAnalysis (Synaptosoft), Excel (Microsoft), and GraphPad Prism (GraphPad Software).

Statistics

Normality of data and equal variance were assessed with Shapiro-Wilk and Brown-Forsythe tests, respectively. Kruskal-Wallis and Dunn's post-hoc tests were used for comparisons of total sleep, daytime sleep, nighttime sleep, sleep bout length, sleep bout number, and electrophysiological parameters. For other data, one-way ANOVA and Tukey post-hoc tests were used for data that met the assumption of normality and equal variance; for normally distributed data with unequal variance, Welch's ANOVA and Dunnet's post-hoc tests were used. Otherwise, Kruskal-Wallis and Dunn's post-hoc tests were used.

Sequence alignments

Alignments were performed with Clustal Omega 2.1 and BOXSHADE. GenBank accession numbers for proteins in Figure S1A are: Inc, NP_001284787; KCTD2, NP_056168; KCTD5, NP_061865; KCTD17.4, NP_001269615.

 Figure Legends

Fig 1. The Inc BTB domain mediates Inc-Cul3 and Inc-Inc interactions

(A) Schematic of N- and C-terminally truncated Inc proteins. Residues comprising the conserved

BTB domain (22 to 123) and C-terminal domain (CTD) (131 to 211) are indicated. **(B and C)** Co-

immunoprecipitation of 3×Myc-tagged Inc or Inc truncation mutants with 3×FLAG-Cul3 **(B)** or

Inc-HA **(C)** from transiently transfected S2 cells. LC, immunoglobulin light chain.

Fig 2. Identification of Inc point mutants that impair Inc-Cul3 interactions

(A) Schematic of Inc point mutants within the BTB domain. **(B-C)** Co-immunoprecipitation of

3×Myc-tagged Inc or Inc point mutants with 3×FLAG-Cul3 **(B)** or 3×FLAG-Inc **(C)** from

transiently transfected S2 cells.

Fig 3. Inc-Cul3 binding is required for Inc activity in vivo

In vivo analysis of *inc^1 inc-Gal4* animals expressing 3×FLAG-tagged Inc or Inc point mutants.

(A) Immunoblot of male whole animal lysates. **(B)** Total sleep duration. Mean ± SEM is shown. n

= 16-58; Kruskal-Wallis, p<0.0001 and Dunn's tests; *p < 0.05; ****p < 0.0001; ns, not significant

(p > 0.5). **(C)** Average daily sleep traces of indicated genotypes from **(B)**. Shading represents SEM.

For all panels, animals are heterozygous for UAS transgenes.

Fig 4. The conserved Inc C-terminus is required for Inc activity in vivo

In vivo analysis of *inc^1 inc-Gal4* animals expressing 3×FLAG-tagged Inc or Inc truncation

mutants. **(A)** Immunoblot of male whole animal lysates. **(B)** Total sleep duration. Mean ± SEM is

shown. n = 27-30; Kruskal-Wallis, p < 0.0001 and Dunn's tests; *p < 0.05; ****p < 0.0001; ns,

not significant (p > 0.5). **(C)** Average daily sleep traces of indicated genotypes from **(B)**. Shading

represents SEM. For all panels, animals are heterozygous for UAS transgenes.

Fig 5. Inc-Cul3 associations and the Inc C-terminus are essential for synaptic homeostasis

(A) Spontaneous and evoked electrophysiological traces at the larval NMJ in the presence and

absence of philanthotoxin. **(B)** Summary data for electrophysiological recordings in the presence

of philanthotoxin normalized to baseline values (absence of PhTx application). Homeostatic

plasticity is not expressed in animals bearing Inc^{F105A} or Inc^{1-123}. Mean ± SEM is shown. n = 8-27;

Kruskal-Wallis, p<0.0001 and Dunn's tests; ***p < 0.001; ****p < 0.0001; ns, not significant (p

> 0.5).

Fig 6. A conserved Inc C-terminal arginine is vital for Inc activity

(A) Schematic of Inc C-terminal point mutants. **(B and C)** Co-immunoprecipitation of 3×Myc-

tagged Inc or Inc point mutants with 3×FLAG-Cul3 **(B)** or 3×FLAG-Inc **(C)** from transiently

transfected S2 cells. **(D)** Immunoblot of lysates prepared from whole male animals expressing

3×Myc-tagged Inc proteins. **(E)** Total sleep duration. Mean ± SEM is shown. n = 24-45; Kruskal-

Wallis, p < 0.0001 and Dunn's tests; ns, not significant (p > 0.05). **(F)** Average daily sleep traces

of indicated genotypes from **(E)**. Shading represents SEM. For panels **(D-F)**, animals are

heterozygous for UAS transgenes.

Fig 7. Inc exhibits key properties of a Cul3 adaptor in neurons in vivo

(A) Co-immunoprecipitation of 3×FLAG-Inc and 3×FLAG-3×HA-Cul3 from male whole animal

lysates. Note immunoprecipitation (IP) was performed with anti-Inc antibody. **(B)** Co-

immunoprecipitation of Myc-Inc and Inc-HA from male whole animal lysates. **(C)** Immunoblots

of male head lysates. **(D)** qRT-PCR of *Cul3* and *inc* mRNA isolated from male heads. Neuronal

Cul3 RNAi decreases *Cul3* transcript levels but does not significantly alter those of *inc*. Each point

represents an independent biological sample measured in technical triplicate. n=3-7. Mean ± SEM

is shown. For *Cul3*, one-way ANOVA, $p < 0.05$ and Tukey tests; for *inc*, Kruskal-Wallis; *$p <$

0.05; ns, not significant ($p > 0.25$).

 Supporting Information

 S1 Fig. Mutagenesis of Inc

(A) Alignment of Inc and its human orthologs. Identical and similar residues are shaded in black

and gray respectively. Locations of Inc truncations and point mutants are indicated by arrowheads

and closed circles, respectively. **(B)** Surface rendering of adjacent BTB domains from the crystal

structure of human KCTD5 [39]. Here and in subsequent figure panels, Inc residue numbers are

superimposed on conserved equivalents in KCTD5. Inc residues whose mutation specifically

weakens Inc-Cul3 associations are highlighted in magenta; other mutated residues are highlighted

in blue. **(C)** Ribbon rendering of adjacent KCTD5 BTB domains. Side chains are shown for Inc

residues mutated to assess Inc multimerization. Note that the T36A/D71A/R85E triple mutant

alters both faces of Inc. **(D)** Surface (left) and ribbon (right) rendering of human KCTD5. Inc C-

terminal point mutants are labeled and indicated in magenta; side chains are shown in the ribbon

rendering. Note that KCTD5 residues equivalent to Inc amino acids 186-211 are not resolved in

the crystal structure, suggesting that they are disordered.

S2 Fig. Stability of Inc truncations in S2 cells

(A) Immunoblot of lysates prepared from transiently transfected S2 cells treated with

cycloheximide (CHX) for indicated durations. **(B)** Quantitation of two biological replicates. For

all timepoints, Welch's ANOVA $p<0.05$ for comparison to wild-type Inc (1-211); Dunnett's tests;

*$p < 0.05$.

S3 Fig. Interactions of Inc 1-156 and Inc 22-123 with Cul3

(A-B) Co-immunoprecipitation of 3×Myc-tagged Inc 1-156 **(A)** or Inc 22-123 **(B)** with 3×FLAG-

Cul3 from transiently transfected S2 cells.

S4 Fig. Stability of Inc point mutants in S2 cells

(A) Immunoblot of lysates prepared from transiently transfected S2 cells treated with cycloheximide (CHX) for indicated durations. **(B)** Quantitation of two biological replicates. For all timepoints, Welch's ANOVA $p < 0.05$ for comparison to wild-type Inc.

S5 Fig. Expression of Inc BTB domain point mutants in vivo

Immunoblot of male whole animal lysates expressing 3×FLAG-tagged Inc and Inc point mutants under *inc-Gal4* control. Animals are heterozygous for UAS transgenes.

S6 Fig. Inc truncations and point mutants lack dominant negative activity in vivo

Total daily sleep for animals expressing tagged Inc or Inc point mutants under the control of *elav^c155^-Gal4* **(A-C)** and *inc-Gal4* **(D)**. **(A)** *elav^c155^-Gal4* expression of 3×FLAG-tagged Inc point mutants. n = 5-24. **(B)** *elav^c155^-Gal4* expression of 3×FLAG-tagged Inc truncations. n = 23-54. **(C)** *elav^c155^-Gal4* expression of Myc-tagged Inc point mutants. Note that Inc^R135H^ is statistically indistinguishable from Inc and Inc^AAA^. n = 18-32. **(D)** *inc-Gal4* expression of 3×FLAG-tagged Myc-tagged Inc point mutants. n = 19-63; For all panels, mean ± SEM is shown. Kruskal-Wallis and Dunn's tests; *$p < 0.01$; ns, not significant ($p > 0.05$). For all panels, animals are heterozygous for UAS transgenes.

S7 Fig. Additional sleep parameters for animals expressing Inc BTB domain point mutants

(A-D) Sleep parameters for *inc^1^ inc-Gal4* animals expressing 3×FLAG-tagged Inc or Inc point mutants. **(A)** Nighttime sleep. **(B)** Daytime sleep. **(C)** Sleep bout duration. **(D)** Sleep bout number.

Mean ± SEM is shown. n = 16-58 as in Fig 3B; Kruskal-Wallis and Dunn's tests; *$p < 0.05$; ***p

< 0.001; ****$p < 0.0001$; ns, not significant ($p > 0.05$). For all panels, animals are heterozygous

for UAS transgenes.

S8 Fig. *UAS-Inc* transgenes do not alter sleep in *inc* mutants in the absence of Gal4

Total daily sleep is shown for control, *inc[1]*, and *inc[1]* animals heterozygous for indicated UAS

transgenes. Mean ± SEM is shown. n = 83-65; Kruskal-Wallis and Dunn's tests; ***$p < 0.001$; ns,

not significant ($p > 0.05$**).**

S9 Fig. Additional sleep parameters for animals expressing Inc C-terminal truncations

(A-D) Sleep parameters for *inc[1] inc-Gal4* animals expressing 3×FLAG-tagged Inc or C-terminally

truncated Inc mutants. **(A)** Nighttime sleep. **(B)** Daytime sleep. **(C)** Sleep bout duration. **(D)** Sleep

bout number. Mean ± SEM is shown. n = 27-30 as in Fig 4B; Kruskal-Wallis and Dunn's tests; *p

< 0.05; **$p < 0.01$; ****$p < 0.0001$; ns, not significant ($p > 0.05$). For all panels, animals are

heterozygous for UAS transgenes.

S10 Fig. Additional sleep parameters for animals expressing Inc C-terminal point mutants

(A-D) Sleep parameters for *inc[1] inc-Gal4* animals expressing 3×Myc-tagged Inc or Inc point

mutants. **(A)** Nighttime sleep. **(B)** Daytime sleep. **(C)** Sleep bout duration. **(D)** Sleep bout number.

Mean ± SEM is shown. n = 24-45 as in Fig 6E; Kruskal-Wallis, $p<0.0001$ and Dunn's tests; *$p <$

0.05; ***$p < 0.001$; ****$p < 0.0001$; ns, not significant ($p > 0.05$). For all panels, animals are

heterozygous for UAS transgenes.

S1 Table. Summary of Inc point mutants targeting Inc-Cul3 interactions

(PDF)

S2 Table. Summary of Inc point mutants targeting Inc-Inc interactions

(PDF)

References

References

1. Codina-Solà M, Rodríguez-Santiago B, Homs A, Santoyo J, Rigau M, Aznar-Laín G, et al.
Integrated analysis of whole-exome sequencing and transcriptome profiling in males with
autism spectrum disorders. Mol Autism. 2015;6: 21. doi:10.1186/s13229-015-0017-0

2. da Silva Montenegro EM, Costa CS, Campos G, Scliar M, de Almeida TF, Zachi EC, et al.
Meta-Analyses Support Previous and Novel Autism Candidate Genes: Outcomes of an
Unexplored Brazilian Cohort. Autism Research. 2020;13: 199–206. doi:10.1002/aur.2238

3. De Rubeis S, He X, Goldberg AP, Poultney CS, Samocha K, Cicek AE, et al. Synaptic,
transcriptional and chromatin genes disrupted in autism. Nature. 2014;515: 209–215.
doi:10.1038/nature13772

4. Iwafuchi S, Kikuchi A, Endo W, Inui T, Aihara Y, Satou K, et al. A novel stop-gain CUL3
mutation in a Japanese patient with autism spectrum disorder. Brain and Development.
2021;43: 303–307. doi:10.1016/j.braindev.2020.09.015

5. Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G, et al. Rate of de
novo mutations and the importance of father's age to disease risk. Nature. 2012;488:
471–475. doi:10.1038/nature11396

6. O'Roak BJ, Vives L, Girirajan S, Karakoc E, Krumm N, Coe BP, et al. Sporadic autism
exomes reveal a highly interconnected protein network of de novo mutations. Nature.
2012;485: 246–250. doi:10.1038/nature10989

7. Wang T, Guo H, Xiong B, Stessman HAF, Wu H, Coe BP, et al. De novo genic mutations
among a Chinese autism spectrum disorder cohort. Nat Comms. 2016;7: 13316.
doi:10.1038/ncomms13316

8. Amar M, Pramod AB, Yu N-K, Herrera VM, Qiu LR, Moran-Losada P, et al. Autism-linked
Cullin3 germline haploinsufficiency impacts cytoskeletal dynamics and cortical
neurogenesis through RhoA signaling. Mol Psychiatry. 2021;26: 3586–3613.
doi:10.1038/s41380-021-01052-x

9. Dong Z, Chen W, Chen C, Wang H, Cui W, Tan Z, et al. CUL3 Deficiency Causes Social
Deficits and Anxiety-like Behaviors by Impairing Excitation-Inhibition Balance through the
Promotion of Cap-Dependent Translation. Neuron. 2020;105: 475-490.e6.
doi:10.1016/j.neuron.2019.10.035

10. Morandell J, Schwarz LA, Basilico B, Tasciyan S, Dimchev G, Nicolas A, et al. Cul3
regulates cytoskeleton protein homeostasis and cell migration during a critical window of
brain development. Nat Comms. 2021;12: 3058–22. doi:10.1038/s41467-021-23123-x

11. Rapanelli M, Tan T, Wang W, Wang X, Wang Z-J, Zhong P, et al. Behavioral, circuitry, and
molecular aberrations by region-specific deficiency of the high-risk autism gene Cul3.
Molecular Psychiatry. 2019;22: 345–14. doi:10.1038/s41380-019-0498-x

12. Rapanelli M. Cholinergic neurons in the basal forebrain are involved in behavioral
abnormalities associated with Cul3 deficiency: Role of prefrontal cortex projections in
cognitive deficits. Translational Psychiatry. 2023.

13. Xia Q, Walker AK, Song C, Wang J, Singh A, Mobley JA, et al. Effects of heterozygous
deletion of autism-related gene Cullin-3 in mice. Li Y, editor. PLoS ONE. 2023;18:
e0283299. doi:10.1371/journal.pone.0283299

14. Gao N, Liu Z, Wang H, Shen C, Dong Z, Cui W, et al. Deficiency of Cullin 3, a Protein
Encoded by a Schizophrenia and Autism Risk Gene, Impairs Behaviors by Enhancing the
Excitability of Ventral Tegmental Area (VTA) DA Neurons. J Neurosci. 2023;43: 6249–
6267. doi:10.1523/JNEUROSCI.0247-23.2023

15. Zhu S, Perez R, Pan M, Lee T. Requirement of Cul3 for axonal arborization and dendritic
elaboration in Drosophila mushroom body neurons. Journal of Neuroscience. 2005;25:
4189–4197. doi:10.1523/JNEUROSCI.0149-05.2005

16. Schaefer H, Rongo C. KEL-8 is a substrate receptor for CUL3-dependent ubiquitin ligase
that regulates synaptic glutamate receptor turnover. Mol Biol Cell. 2006;17: 1250–1260.
doi:10.1091/mbc.E05-08-0794

17. Stavropoulos N, Young MW. insomniac and Cullin-3 regulate sleep and wakefulness in
Drosophila. Neuron. 2011;72: 964–976. doi:10.1016/j.neuron.2011.12.003

18. Pfeiffenberger C, Allada R. Cul3 and the BTB adaptor insomniac are key regulators of sleep
homeostasis and a dopamine arousal pathway in Drosophila. PLoS Genet. 2012;8:
e1003003. doi:10.1371/journal.pgen.1003003

19. Singh K, Ju JY, Walsh MB, DiIorio MA, Hart AC. Deep conservation of genes required for
both Drosphila melanogaster and Caenorhabditis elegans sleep includes a role for
dopaminergic signaling. Sleep. 2014;37: 1439–1451. doi:10.5665/sleep.3990

20. Kikuma K, Li X, Perry S, Li Q, Goel P, Chen C, et al. Cul3 and insomniac are required for
rapid ubiquitination of postsynaptic targets and retrograde homeostatic signaling. Nat
Comms. 2019;10: 2998–13. doi:10.1038/s41467-019-10992-6

21. Furukawa M, He YJ, Borchers C, Xiong Y. Targeting of protein ubiquitination by BTB-
Cullin 3-Roc1 ubiquitin ligases. Nat Cell Biol. 2003;5: 1001–1007. doi:10.1038/ncb1056

22. Geyer R, Wee S, Anderson S, Yates J, Wolf DA. BTB/POZ domain proteins are putative
substrate adaptors for cullin 3 ubiquitin ligases. Mol Cell. 2003;12: 783–790.

23. Pintard L, Willis JH, Willems A, Johnson J-LF, Srayko M, Kurz T, et al. The BTB protein
MEL-26 is a substrate-specific adaptor of the CUL-3 ubiquitin-ligase. Nature. 2003;425:
311–316. doi:10.1038/nature01959

24. Xu L, Wei Y, Reboul J, Vaglio P, Shin T-H, Vidal M, et al. BTB proteins are substrate-
specific adaptors in an SCF-like modular ubiquitin ligase containing CUL-3. Nature.
2003;425: 316–321. doi:10.1038/nature01985

25. Bennett EJ, Rush J, Gygi SP, Harper JW. Dynamics of cullin-RING ubiquitin ligase network
revealed by systematic quantitative proteomics. Cell. 2010;143: 951–965.
doi:10.1016/j.cell.2010.11.017

26. Allen E, Ding J, Wang W, Pramanik S, Chou J, Yau V, et al. Gigaxonin-controlled
degradation of MAP1B light chain is critical to neuronal survival. Nature. 2005;438: 224–
228.

27. Salinas GD, Blair LAC, Needleman LA, Gonzales JD, Chen Y, Li M, et al. Actinfilin is a
Cul3 substrate adaptor, linking GluR6 kainate receptor subunits to the ubiquitin-proteasome
pathway. J Biol Chem. 2006;281: 40164–40173. doi:10.1074/jbc.M608194200

28. Zhang Q, Zhang L, Wang B, Ou C-Y, Chien C-T, Jiang J. A hedgehog-induced BTB protein
modulates hedgehog signaling by degrading Ci/Gli transcription factor. Dev Cell. 2006;10:
719–729. doi:10.1016/j.devcel.2006.05.004

29. Sobieszczuk DF, Poliakov A, Xu Q, Wilkinson DG. A feedback loop mediated by
degradation of an inhibitor is required to initiate neuronal differentiation. Genes Dev.
2010;24: 206–218. doi:10.1101/gad.554510

30. Lin M-Y, Lin Y-M, Kao T, Chuang H-H, Chen R-H. PDZ-RhoGEF ubiquitination by
Cullin3-KLHL20 controls neurotrophin-induced neurite outgrowth. J Cell Biol. 2011;193:
985–994. doi:10.1083/jcb.201103015

31. Tseng LA, Bixby JL. Interaction of an intracellular pentraxin with a BTB-Kelch protein is
associated with ubiquitylation, aggregation and neuronal apoptosis. Mol Cell Neurosci.
2011;47: 254–264. doi:10.1016/j.mcn.2011.04.005

32. Escamilla CO, Filonova I, Walker AK, Xuan ZX, Holehonnur R, Espinosa F, et al. Kctd13
deletion reduces synaptic transmission via increased RhoA. Nature. 2017;551: 227–231.
doi:10.1038/nature24470

33. Li Q, Kellner DA, Hatch HAM, Yumita T, Sanchez S, Machold RP, et al. Conserved
properties of Drosophila Insomniac link sleep regulation and synaptic function. PLoS
Genet. 2017;13: e1006815. doi:10.1371/journal.pgen.1006815

34. Mathew R, Seiler MP, Scanlon ST, Mao A-P, Constantinides MG, Bertozzi-Villa C, et al.
BTB-ZF factors recruit the E3 ligase cullin 3 to regulate lymphoid effector programs.
Nature. 2012. doi:10.1038/nature11548

35. Schwenk J, Metz M, Zolles G, Turecek R, Fritzius T, Bildl W, et al. Native GABA(B)
receptors are heteromultimers with a family of auxiliary subunits. Nature. 2010;465: 231–
235. doi:10.1038/nature08964

36. Stogios PJ, Downs GS, Jauhal JJS, Nandra SK, Privé GG. Sequence and structural analysis
of BTB domain proteins. Genome Biol. 2005;6: R82. doi:10.1186/gb-2005-6-10-r82

37. Mencacci NE, Rubio-Agusti I, Zdebik A, Asmus F, Ludtmann MHR, Ryten M, et al. A
missense mutation in KCTD17 causes autosomal dominant myoclonus-dystonia. American
journal of human genetics. 2015;96: 938–947. doi:10.1016/j.ajhg.2015.04.008

38. Bayón Y, Trinidad AG, de la Puerta ML, Del Carmen Rodríguez M, Bogetz J, Rojas A, et al.
KCTD5, a putative substrate adaptor for cullin3 ubiquitin ligases. FEBS J. 2008;275: 3900–
3910. doi:10.1111/j.1742-4658.2008.06537.x

39. Dementieva IS, Tereshko V, McCrossan ZA, Solomaha E, Araki D, Xu C, et al. Pentameric
assembly of potassium channel tetramerization domain-containing protein 5. J Mol Biol.
2009;387: 175–191. doi:10.1016/j.jmb.2009.01.030

40. Errington WJ, Khan MQ, Bueler SA, Rubinstein JL, Chakrabartty A, Privé GG. Adaptor
protein self-assembly drives the control of a cullin-RING ubiquitin ligase. Structure.
2012;20: 1141–1153. doi:10.1016/j.str.2012.04.009

41. Canning P, Cooper CDO, Krojer T, Murray JW, Pike ACW, Chaikuad A, et al. Structural
basis for Cul3 protein assembly with the BTB-Kelch family of E3 ubiquitin ligases. J Biol
Chem. 2013;288: 7803–7814. doi:10.1074/jbc.M112.437996

42. Ji AX, Privé GG. Crystal Structure of KLHL3 in Complex with Cullin3. PLoS ONE. 2013;8:
e60445. doi:10.1371/journal.pone.0060445

43. Balasco N, Pirone L, Smaldone G, Di Gaetano S, Esposito L, Pedone EM, et al. Molecular
recognition of Cullin3 by KCTDs: insights from experimental and computational
investigations. Biochim Biophys Acta. 2014;1844: 1289–1298.
doi:10.1016/j.bbapap.2014.04.006

44. Pirone L, Smaldone G, Esposito C, Balasco N, Petoukhov MV, Spilotros A, et al. Proteins
involved in sleep homeostasis: Biophysical characterization of INC and its partners.
Biochimie. 2016;131: 106–114. doi:10.1016/j.biochi.2016.09.013

45. Ji AX, Chu A, Nielsen TK, Benlekbir S, Rubinstein JL, Privé GG. Structural insights into
KCTD protein assembly and Cullin3 recognition. J Mol Biol. 2015.
doi:10.1016/j.jmb.2015.08.019

46. Frank CA, Kennedy MJ, Goold CP, Marek KW, Davis GW. Mechanisms underlying the
rapid induction and sustained expression of synaptic homeostasis. Neuron. 2006;52: 663–
677. doi:10.1016/j.neuron.2006.09.029

47. Pintard L, Kurz T, Glaser S, Willis JH, Peter M, Bowerman B. Neddylation and
deneddylation of CUL-3 is required to target MEI-1/Katanin for degradation at the meiosis-
to-mitosis transition in C. elegans. Curr Biol. 2003;13: 911–921.

48. Wee S, Geyer RK, Toda T, Wolf DA. CSN facilitates Cullin-RING ubiquitin ligase function
by counteracting autocatalytic adapter instability. Nat Cell Biol. 2005;7: 387–391.
doi:10.1038/ncb1241

49. Zhang DD, Lo S-C, Sun Z, Habib GM, Lieberman MW, Hannink M. Ubiquitination of
Keap1, a BTB-Kelch substrate adaptor protein for Cul3, targets Keap1 for degradation by a
proteasome-independent pathway. J Biol Chem. 2005;280: 30091–30099.
doi:10.1074/jbc.M501279200

50. Tener SJ, Lin Z, Park SJ, Oraedu K, Ulgherait M, Van Beek E, et al. Neuronal knockdown of
Cullin3 as a Drosophila model of autism spectrum disorder. Sci Rep. 2024;14: 1541.
doi:10.1038/s41598-024-51657-9

51. Pinkas DM, Sanvitale CE, Bufton JC, Sorrell FJ, Solcan N, Chalk R, et al. Structural
complexity in the KCTD family of Cullin3-dependent E3 ubiquitin ligases. Biochem J.
2017;474: 3747–3761. doi:10.1042/BCJ20170527

52. Jiang W, Wang W, Kong Y, Zheng S. Structural basis for the ubiquitination of G protein βγ
subunits by KCTD5/Cullin3 E3 ligase. Sci Adv. 2023;9: eadg8369.
doi:10.1126/sciadv.adg8369

53. Nguyen DM, Rath DH, Devost D, Pétrin D, Rizk R, Ji AX, et al. Structure and dynamics of a
pentameric KCTD5/Cullin3/Gβγ E3 ubiquitin ligase complex.
http://biorxiv.org/lookup/doi/10.1101/2023.09.20.558662; 2023 Sep.
doi:10.1101/2023.09.20.558662

54. McGourty CA, Akopian D, Walsh C, Gorur A, Werner A, Schekman R, et al. Regulation of
the CUL3 Ubiquitin Ligase by a Calcium-Dependent Co-adaptor. Cell. 2016;167: 525-
538.e14. doi:10.1016/j.cell.2016.09.026

55. Muntean BS, Marwari S, Li X, Sloan DC, Young BD, Wohlschlegel JA, et al. Members of
the KCTD family are major regulators of cAMP signaling. Proceedings of the National
Academy of Sciences. 2022;119. doi:10.1073/pnas.2119237119

56. Li Q, Jang H, Lim KY, Lessing A, Stavropoulos N. insomniac links the development and
function of a sleep-regulatory circuit. Calabrese RL, Cirelli C, editors. eLife. 2021;10:
e65437. doi:10.7554/eLife.65437

57. Kim E-J, Kim S-H, Jin X, Jin X, Kim H. KCTD2, an adaptor of Cullin3 E3 ubiquitin ligase,
suppresses gliomagenesis by destabilizing c-Myc. Cell Death Differ. 2017;24: 649–659.
doi:10.1038/cdd.2016.151

58. He H, Peng Y, Fan S, Chen Y, Zheng X, Li C. Cullin3/KCTD5 induces monoubiquitination
of ΔNp63α and impairs its activity. FEBS Letters. 2018;592: 2334–2340.
doi:10.1002/1873-3468.13104

59. Cho HJ, Ryu K-J, Baek KE, Lim J, Kim T, Song CY, et al. Cullin 3/KCTD5 promotes the
ubiqutination of Rho guanine nucleotide dissociation inhibitor 1 and regulates its stability. J
Microbiol Biotechnol. 2020;30: 1–7. doi:10.4014/jmb.2007.07033

60. Brockmann M, Blomen VA, Nieuwenhuis J, Stickel E, Raaben M, Bleijerveld OB, et al.
Genetic wiring maps of single-cell protein states reveal an off-switch for GPCR signalling.
Nature. 2017;546: 307–311. doi:10.1038/nature22376

61. Young BD, Sha J, Vashisht AA, Wohlschlegel JA. Human Multisubunit E3 Ubiquitin Ligase
Required for Heterotrimeric G-Protein β-Subunit Ubiquitination and Downstream
Signaling. J Proteome Res. 2021. doi:10.1021/acs.jproteome.1c00292

62. Kasahara K, Kawakami Y, Kiyono T, Yonemura S, Kawamura Y, Era S, et al. Ubiquitin-
proteasome system controls ciliogenesis at the initial step of axoneme extension. Nat
Comms. 2014;5: 5081. doi:10.1038/ncomms6081

63. Kim K, Ryu D, Dongiovanni P, Ozcan L, Nayak S, Ueberheide B, et al. Degradation of
PHLPP2 by KCTD17, via a Glucagon-Dependent Pathway, Promotes Hepatic Steatosis.
Gastroenterology. 2017;153: 1568-1580.e10. doi:10.1053/j.gastro.2017.08.039

64. Shin MC, Jung YH, Jeong Y, Oh A-R, Lee SB, Kim K. Kctd17-mediated Chop degradation
promotes adipogenic differentiation. Biochemical and Biophysical Research
Communications. 2023;653: 126–132. doi:10.1016/j.bbrc.2023.02.068

65. Boada M, Antúnez C, Ramírez-Lorca R, DeStefano AL, González-Pérez A, Gayán J, et al.
ATP5H/KCTD2 locus is associated with Alzheimer's disease risk. Molecular
Psychiatry. 2014;19: 682–687. doi:10.1038/mp.2013.86

66. Barfield R, Wang H, Liu Y, Brody JA, Swenson B, Li R, et al. Epigenome-wide association
analysis of daytime sleepiness in the Multi-Ethnic Study of Atherosclerosis reveals African-
American-specific associations. Sleep. 2019;42: zsz101. doi:10.1093/sleep/zsz101

67. Lin DM, Goodman CS. Ectopic and increased expression of Fasciclin II alters motoneuron
growth cone guidance. Neuron. 1994;13: 507–523.

68. Budnik V, Koh Y-H, Guan B, Hartmann B, Hough C, Woods D, et al. Regulation of Synapse
Structure and Function by the Drosophila Tumor Suppressor Gene dlg. Neuron. 1996;17:
627–640. doi:10.1016/S0896-6273(00)80196-8

69. Hudson AM, Cooley L. Drosophila Kelch functions with Cullin-3 to organize the ring canal
actin cytoskeleton. J Cell Biol. 2010;188: 29–37. doi:10.1083/jcb.200909017

70. Groth AC, Fish M, Nusse R, Calos MP. Construction of transgenic Drosophila by using the
site-specific integrase from phage phiC31. Genetics. 2004;166: 1775–1782.

71. Ryder E, Blows F, Ashburner M, Bautista-Llacer R, Coulson D, Drummond J, et al. The
DrosDel collection: a set of P-element insertions for generating custom chromosomal
aberrations in Drosophila melanogaster. Genetics. 2004;167: 797–813.
doi:10.1534/genetics.104.026658

1009

A

Residues

B

C

Figure 1

A

F47A   F105A

B

	3×Myc-Inc	WT	F47A	F105A	F47A F105A

C

	3×Myc-Inc	WT	F47A	F105A	F47A F105A

Figure 2

A

B

C

Figure 3

A

B

C

Figure 4

A

Wild type | inc^{kk3}; BG57-Gal4/+ | inc^{kk3}; BG57-Gal4/+; UAS-3×FLAG-Inc/+ | inc^{kk3}; BG57-Gal4/+; UAS-3×FLAG-Inc$^{1-123}$/+

- PhTx

+ PhTx

20 nA   10 ms
200 ms   1 nA

B

mEPSC

Quantal content

% baseline (- PhTx)

ns

ns

****   ***

Wild type   -   WT   F105A   1-123

UAS-3×FLAG-Inc

inc^{kk3}; BG57-Gal4/+

Figure 5

Figure 6

Figure 7

A

Inc N-terminal truncation
Inc C-terminal truncation
Inc-Cul3 interaction point mutant
Inc-Inc interaction point mutant
Inc[R135H]
Inc[AAA]

B   **C**   **D**

Supplementary Figure 1

A

3×Myc-Inc

| 1-211 (WT) | 22-211 | 31-211 | 124-211 |

CHX (0–6h) 0 2 4 6   0 2 4 6   0 2 4 6   0 2 4 6

34 —
26 —
kDa
17 —

Myc
(Inc)

55 —

Tubulin

B

- 1-211
- 22-211
- 31-211
- 124-211

Normalized signal

Time (hr)

A

B

A

3×Myc-Inc

CHX (0–6h)

	WT	F47A	F105A	F47A F105A	T36A D71A R85E
	0 2 4 6	0 2 4 6	0 2 4 6	0 2 4 6	0 2 4 6

34 —
kDa

55 —

Myc (Inc)

Tubulin

B

WT
F47A
F105A
F47A F105A
T36A D71A R85E

Supplementary Figure 4

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 10