1 **Presynaptic mitochondria volumes and complexity of subsynaptic distribution increase**
2 **during development at a high-fidelity synapse**

4 *(Abbreviated title):* Presynaptic mitochondria volumes and distribution during development
5 Connon I. Thomas[1], Christian Keine[3], Satoko Okayama[2], Rachel Satterfield[2], Morgan Musgrove[1],
6 Debbie Guerrero-Given[1], Naomi Kamasawa[1,^], and Samuel M. Young, Jr.[3,4,^].
7
[1]Electron Microscopy Core Facility, Max Planck Florida Institute, Jupiter, FL 33458, USA
[2]Molecular Mechanisms of Synaptic Function, Max Planck Florida Institute, Jupiter, FL 33458,
USA; [3]Department of Anatomy and Cell Biology, Iowa Neuroscience Institute, University of Iowa,
Iowa City, IA 52242, USA; [4]Department of Otolaryngology, University of Iowa, Iowa City, IA
52242, USA.

^Correspondence
Naomi Kamasawa, PhD.
Electron Microscopy Core Facility,
Max Planck Florida Institute for Neuroscience
One Max Planck Way
Jupiter, FL 33458
Email: naomi.kamasawa@mpfi.org

^Correspondence
Samuel M. Young, Jr. PhD
Department of Anatomy and Cell Biology
University of Iowa
PBDB 5322
169 Newton Road
Iowa City, IA 52242
Email: samuel-m-young@uiowa.edu

Number of Pages = 48
Number of Figures = 10
Number of Tables = 2
Number of Multimedia/3D models = 7
Number of words in Abstract = 250/250
Number of words in Significance Statement = 119/120
Number of words in Introduction = 650/650
Number of words in Discussion = 1500/1500

Acknowledgements
We would like to thank the members of the Young lab for their comments on the manuscript,
Keisuke Ohta for initial discussions, and Melissa A. Ryan and Michael Morehead for help with
EM volume data analysis. This work was supported by the National Institute of Deafness and
Communication Disorders (R01 DC014093), the University of Iowa, Max Planck Society (SMY)
and a German Research Foundation postdoctoral fellowship (DFG 420075000) (CK).

Conflict of Interest: The authors declare no conflict of interests.

Abstract

The calyx of Held, a large glutamatergic presynaptic terminal in the auditory brainstem undergoes

developmental changes to support the high action-potential firing rates required for auditory

information encoding. In addition, calyx terminals are morphologically diverse which impacts

vesicle release properties and synaptic plasticity. Mitochondria influence synaptic plasticity

through calcium buffering and are crucial for providing the energy required for synaptic

transmission. Therefore, it has been postulated that mitochondrial levels increase during

development and contribute to the morphological-functional diversity in the mature calyx.

However, the developmental profile of mitochondrial volumes and subsynaptic distribution at the

calyx of Held remains unclear. To provide insight on this, we developed a helper-dependent

adenoviral vector (HdAd) that expresses the genetically encoded peroxidase marker for

mitochondria, mito-APEX2, at the mouse calyx of Held. We developed protocols to detect labeled

mitochondria for use with serial block face scanning electron microscopy to carry out semi-

automated segmentation of mitochondria, high-throughput whole terminal reconstruction and

presynaptic ultrastructure in mice of either sex. Subsequently, we measured mitochondrial

volumes and subsynaptic distributions at the immature postnatal day 7 (P7) and the mature (P21)

calyx. We found an increase of mitochondria volumes in terminals and axons from P7 to P21 but

did not observe differences between stalk and swelling subcompartments in the mature calyx.

Based on these findings, we propose that mitochondrial volumes developmentally increase to

support high firing rates but have limited contribution to morphological-functional diversity at the

calyx.

Significance Statement

Elucidating the developmental processes of auditory brainstem presynaptic terminals is critical to

understanding auditory information encoding. Additionally, morphological-functional diversity at

these terminals is proposed to enhance coding capacity. Mitochondria provide energy for synaptic

transmission and can buffer calcium, impacting synaptic plasticity; however, their developmental

profile to ultimately support the energetic demands of synapses following the onset of hearing

remains unknown. Therefore, we created a helper-dependent adenoviral vector with the

mitochondria-targeting peroxidase mito-APEX2 and expressed it at the mouse calyx of Held.

Volumetric reconstructions of serial block face electron microscopy data of immature and mature

labeled calyces reveal that mitochondrial volumes are increased to support high firing rates upon

maturity and likely have little contribution to morphological-functional diversity at the calyx.

Introduction

During development of auditory brainstem circuits, many synapses undergo morphological and physiological transformations to support the high action-potential (AP) firing rates required in the initial stages of auditory information processing (Yu and Goodrich, 2014). Synaptic transmission and developmental remodeling are energetically demanding processes (Harris et al., 2012), and mitochondria are essential organelles for supplying energy for these processes (Devine and Kittler, 2018). In addition, mitochondria play a role in calcium (Ca^{2+}) buffering which can impact synaptic plasticity (Vos et al., 2010). Therefore, understanding how presynaptic mitochondria volume and distribution change with development is integral to understanding how high-fidelity AP firing is achieved and maintained in the auditory brainstem.

The calyx of Held, a large glutamatergic presynaptic terminal in the medial nucleus of the trapezoid body (MNTB) in the auditory brainstem undergoes considerable morphological and physiological transformations around the onset of hearing at the end of the second postnatal week (Baydyuk et al., 2016). At the physiological level, the calyx of Held/MNTB synapse shifts from exhibiting frequent synaptic failures before hearing onset to reliable synaptic transmission at firing rates up to several hundred Hertz (Taschenberger and von Gersdorff, 2000). In parallel there is a reduction in release probability (*Pr*) (Taschenberger and von Gersdorff, 2000; Iwasaki and Takahashi, 2001) and a shift from micro- to nano-domain release due to tighter coupling—the distance between synaptic vesicles (SVs) and voltage gated calcium channels (VGCCs) (Fedchyshyn and Wang, 2005; Wang et al., 2008; Chen et al., 2015). At the morphological level, the calyx transforms from cup-like to a highly fenestrated structure consisting of stalks and swellings (Kandler and Friauf, 1993; Rowland et al., 2000; Wimmer et al., 2006), and undergoes ultrastructural changes such as smaller active zones (AZs) and a reduced number of docked SVs

(Taschenberger et al., 2002). Finally, the mature calyx contains mitochondria tethered to the

presynaptic plasma membrane (Rowland et al., 2000; Perkins et al., 2010) and SVs are tightly

packed around mitochondria (Wimmer et al., 2006).

In addition to these developmental changes, a morphological-functional diversity at the

mature calyx has been shown, in which calyces with many swellings have lower synaptic

depression and SV release probability (Grande and Wang, 2011). Furthermore, stalks and

swellings differ in Ca^{2+} buffering and SV coupling states, with stalks having tighter SV coupling

than swellings (Fekete et al., 2019). Nanodomain release corresponds to tight coupling with SV-

VGCC distances of ~25 nm and is considered to be energetically favorable (Eggermann et al.,

2012) Therefore, the energetic demands of AZs at stalks are predicted to be lower than swellings.

However, the developmental changes in mitochondria volumes and subsynaptic distribution

between stalks and swellings in the calyx remain unknown. To address these questions, we carried

out volumetric reconstructions at the electron microscopic (EM) level of calyces and mitochondria

at the immature postnatal day 7 (P7) and mature P21 calyx. To do so, we created a helper-

dependent adenoviral vector (HdAd) that expresses the mitochondrial EM marker mito-APEX2

(Martell et al., 2012; Lam et al., 2015; Martell et al., 2017). Since serial block face scanning EM

(SBF-SEM) makes large scale reconstructions more accessible than transmission EM (TEM)

(Lichtman and Denk, 2011), we developed protocols permitting identification of mito-APEX2

with SBF-SEM and reconstructed multiple mito-APEX2 positive terminals while resolving

presynaptic ultrastructure. We found a developmental increase in mitochondrial volumes in the

calyx and its respective axon while mitochondrial volumes were not different between stalk and

swellings of the adult calyx. These data suggest, that mitochondrial volumes developmentally

increase to support high firing rates but have limited contribution to morphological-functional

diversity at the calyx. Finally, since mito-APEX2 permits identification of calyces while

preserving ultrastructure, we optimized protocols using automated tape-collecting ultramicrotome

serial section SEM (ATUM ssSEM) to analyze single-AZ ultrastructure in partially reconstructed

calyces. We propose our viral vectors and methods will have broad applications for quantitative

analyses of subsynaptic organization of SVs, mitochondria, and their morphological continuum

using TEM, ssSEM and SBF-SEM.

Materials and Methods

Animals

All experiments were performed in accordance with animal welfare laws and approved by the

Institutional Committee for Care and Use of Animals at the Max Planck Florida Institute for

Neuroscience and the University of Iowa and complied with accepted ethical best practice.

Animals were housed at a 12-hour light/dark cycle and had access to food and water ad libitum.

Experiments were performed on C57Bl/6J mice (RRID:IMSR_JAX:000664, The Jackson

Laboratory, Bar Harbor, USA) of either sex. Virus constructs were injected at postnatal day 1 (P1)

and experiments were performed at P7 (immature calyces) or P21 (functionally mature calyces).

DNA construct and recombinant viral vector production

Mito-V5-APEX2 (Lam et al., 2015), a gift from Alice Ting (Addgene plasmid #72480,

http://n2t.net/addgene:72480, RRID:Addgene_72480), was cloned into the EcoRI and NotI sites

of the synapsin expression cassette. This cassette included the 470 bp human synapsin (hsyn)

promoter, the minute virus of mice (mvm) intron, and the bovine growth hormone polyA sequence

(Montesinos et al., 2011). HdAd was produced as previously described (Montesinos et al., 2016).

The mito V5 expression cassette was cloned into the AscI site of a modified version of pdelta

28E4, gift from Dr. Philip Ng, using InFusion (Clontech). This version of pdelta 28E4 was

modified to also contain a separate neurospecific EGFP expression cassette driven by the hsyn

promoter (Lubbert et al., 2017). HdAd was produced as previously described (Montesinos et al.,

2016). Briefly, pHdAd mito-APEX2 hsyn EGFP was digested with PmeI and into 116 producer

cells. HdAd was serially amplified in five consecutive passages. Each successive passage was

performed after cytopathic effect occurred and cell lysates were subjected to three freeze-thaw

cycles to lyse cells and thereby release the viral particles. HdAd was stored at $-80°C$ in storage

buffer containing 10 mM HEPES, 250 mM sucrose, and 1 mM $MgCl_2$ at pH 7.4.

Virus injections

Virus injections at P1 were performed as described previously (Chen et al., 2013). Briefly, mice

were anesthetized by hypothermia for 5 min in an ice bath and a stereotactic injection of ~1 μL of

HdAd (total number of virus particles <1e9) syn mito-V5-APEX2 syn EGFP into the cochlear

nucleus was performed at a rate of 1 μL/min using glass pipettes (Blaubrand, IntraMARK).

Following the injection, the glass needle was left in place for 1 min to dissipate the pressure and

then slowly retracted. Animals were then placed under an infrared heat lamp and allowed to

recover before being returned to their respective cages with their mother.

Acute brain slice preparation

Acute coronal brainstem slices (200 μm) containing the medial nucleus of the trapezoid body

(MNTB) were prepared from control and EGFP-mito-APEX2-injected animals using a Leica VT

1200S vibratome equipped with zirconia ceramic blades (EF-INZ10, Cadence Blades) in low-

calcium artificial cerebrospinal fluid (aCSF) solution, containing (in mM): 125 NaCl, 2.5 KCl,

3 $MgCl_2$, 0.1 $CaCl_2$, 10 glucose, 25 $NaHCO_3$, 1.25 NaH_2PO_4, 0.4 L-ascorbic acid, 3 myo-inositol,

and 2 Na-pyruvate, pH 7.3–7.4. The blade was advanced at a speed of 20-50 μm/s. Slices were

immediately transferred to an incubation beaker containing standard extracellular solution (same

as above but using 1 mM $MgCl_2$ and 1.2 mM $CaCl_2$ at 37°C, continuously bubbled with 95% O_2–

5% CO_2). After approximately 45 min of incubation, slices were transferred to a recording

chamber with the same saline at room temperature (~22°C).

Electrophysiology

Fiber stimulation was performed as described previously (Dong et al., 2018). Briefly, to record AP-evoked EPSCs, fibers of the trapezoid body were stimulated using a bipolar platinum-iridium electrode (FHC, Model MX214EP) positioned medially of the MNTB. Postsynaptic MNTB neurons were whole-cell voltage clamped at −73 mV using an EPC10/2 amplifier controlled by Patchmaster Software (version 2x90.2, HEKA Elektronik, RRID:SCR_000034). Slices were continuously perfused with standard aCSF solution and visualized by an upright microscope (BX51WI, Olympus) through a 60x water-immersion objective (LUMPlanFL N, Olympus) and an EMCCD camera (Andor Luca S, Oxford Instruments). To identify calyces transduced with the EGFP-mito-APEX2 virus, the slice was illuminated at an excitation wavelength of 480 nm using a Polychrome V xenon bulb monochromator (TILL Photonics). During recordings, the standard extracellular solution was supplemented with 1 mM kynurenic acid (Tocris, Cat# 0223) to avoid desensitization and saturation of postsynaptic AMPA receptors, 50 μM D-AP-5 (Tocris, Cat# 0106) to block NMDA receptors, and 20 μM bicuculline (Tocris, Cat# 0131) and 5 μM strychnine (Tocris, Cat# 2785) to block inhibitory GABA- and glycine receptors, respectively. Patch pipettes were fabricated using a PIP6 (HEKA) and had a resistance of ∼3–4 MΩ when filled with the following (in mM): 130 Cs-gluconate, 20 tetraethylammonium (TEA)-Cl, 10 HEPES, 5 Na_2-phosphocreatine, 4 MgATP, 0.3 NaGTP, 6 QX-314, and 5 EGTA, pH 7.2 (315 mOsm). Liquid junction potential was 13 mV and corrected for all reported voltages. Data were acquired at a sampling rate of 50 kHz. Series resistance (3–8 MΩ) was compensated online to <3.0 MΩ, and the remaining series resistance was further compensated offline to 0 MΩ with a time lag of 20 μs (Traynelis, 1998) using a custom-written MATLAB function (version 9.4; Mathworks, RRID:SCR_001622).

Electrophysiological data analysis

Electrophysiological data were imported to MATLAB using a custom-modified version of sigTool

(Lidierth, 2009) and analyzed offline with custom-written MATLAB functions. EPSC amplitudes

were measured as peak amplitude minus baseline preceding the EPSC. The RRP size was

calculated using the back-extrapolation method as described earlier (Neher, 2015).

EM slice preparation

Slices for EM were prepared independently of acute slices for electrophysiology experiments. P7

or P21 animals previously injected with the EGFP-mito-APEX2 viral vector were anesthetized

with an intraperitoneal injection of tribromoethanol (250 mg/ kg of body weight) and perfused

transcardially with 0.9% NaCl in MilliQ H_2O, followed by a fixative containing 2%

paraformaldehyde and 1-2% glutaraldehyde in 0.1 M cacodylate buffer (CB, pH 7.4) containing 2

mM $CaCl_2$. Brains were extracted and post fixed in the same fixative for 2 hours, then washed in

CB. The brainstem was sliced coronally at a thickness of 50 µm (TEM, ssSEM) or 100 µm (SBF-

SEM) using a Leica VT 1200S vibratome in 0.1 M CB. The EGFP expression at the injection site

and in the contralateral MNTB was confirmed under an epifluorescence microscope (CKX41,

Olympus). All EM images of calyx terminals were taken from the middle region of the MNTB.

TEM sample processing and data analysis

Slices for conventional TEM imaging were reacted in a solution containing 0.05% DAB (3,3'-

Diaminobenzidine, Sigma Aldrich) in CB (pH 7.4) for 1 hour on ice. Slices were then reacted with

0.03% H_2O_2 (v/v) in the same well for 5-10 minutes and rinsed thoroughly. Labeling was

confirmed using a standard brightfield microscope. Tissue was reacted with 1% aqueous osmium

tetroxide for 1 hour on ice, rinsed with distilled water, reacted with 1% aqueous uranyl acetate for

30 minutes in the dark at 4˚C, and rinsed again with distilled water. Tissue was then dehydrated in

a graded ethanol series (30, 50, 70, 90, 100%), 1:1 ethanol to acetone, 100% acetone, 1:1 acetone

to propylene oxide (PO), and 100% PO, each 2×5 minutes. Tissue was infiltrated using 1:3 PO

to Durcupan resin (Sigma Aldrich) for 2 hours, 1:1 PO to resin for 2 hours, and 3:1 PO to resin

overnight, then flat embedded in 100% resin on a glass slide and covered with an Aclar sheet at

60˚C for 2 days. The MNTB region contralateral to the injected cochlear nucleus was trimmed,

mounted on empty resin blocks with cyanoacrylate glue (Krazy Glue, Elmer's Products), and

sectioned at 40 nm using a Leica UC7 ultramicrotome. Sections were counterstained with uranyl

acetate and lead citrate, and examined in a Tecnai G2 Spirit BioTwin transmission electron

microscope (Thermo Fisher Science, Waltham, MA) at 100 kV acceleration voltage. Images were

taken with a Veleta CCD camera (Olympus) operated by TIA software (Thermo Fisher Scientific).

Images used for quantification were taken at 60,000x magnification. All TEM data were analyzed

using Fiji imaging analysis software (Schindelin et al., 2012), RRID:SCR_002285). Linescans

were made using the plot profile function in Fiji, which was also used for ssSEM and SBF-SEM

analyses. Active zones (AZs) were identified by the following criteria: (1) rigid opposed pre- and

post-synaptic membrane, (2) SVs clustering at the membrane, and (3) an asymmetric post synaptic

density. AZs were measured as the presynaptic membrane opposing the postsynaptic density.

Puncta adherentia (PA) were differentiated from AZs by lack of vesicles, a symmetric pre- and

post-synaptic density, and shorter length. Vesicle distance to the AZ was calculated using a 32-bit

Euclidean distance map generated from the AZ for all vesicles within 200 nm of the presynaptic

membrane. Vesicles within 5 nm of the AZ were considered to be docked (Taschenberger et al.,

2002; Yang et al., 2010).

Section based immuno-electron microscopy

The samples prepared for TEM above were used for on-section immunoelectron microscopy. The MNTB region contralateral to the injected cochlear nucleus was trimmed and sectioned at 100 nm thickness using a Leica UC7 ultramicrotome. Sections were collected onto a 100-mesh nickel grid and air dried. As an antigen unmasking procedure for Durcupan resin, grids were placed section side down on water heated on a hotplate to 93-95°C for 10 minutes (Stirling and Graff, 1995). Following heat treatment, the water was slowly cooled down for 15 minutes by mixing in room temperature water. Without drying, grids were immediately placed onto drops of blocking medium for 10 minutes composed of 0.1% Tween-20, 1% bovine serum albumin, 1% normal goat serum, 1% fish skin gelatin, and 0.005% sodium azide diluted in Tris-buffered saline (TBS) buffer (pH 7.4). Grids were then incubated on a chicken anti-GFP primary antibody solution (0.5 mg/mL, Abcam Cat# ab13970, RRID:AB_300798) diluted 1:100 in the same blocking medium, and placed in a humid chamber overnight at room temperature. They were then washed and secondary antibody incubation was performed for one hour at room temperature with a 12 nm colloidal gold donkey anti-chicken antibody (Jackson ImmunoResearch Labs Cat# 703-205-155, RRID:AB_2340368) diluted at 1:30 in blocking medium. Grids were washed again on TBS and the reaction was stabilized by placing them on 1% glutaraldehyde in PBS for 5 minutes. Grids were rinsed on water and counterstained with uranyl acetate and lead citrate, and examined in a Tecnai G2 Spirit BioTwin transmission electron microscope (Thermo Fisher Scientific) at 80 kV acceleration voltage. Images were taken with a Veleta CCD camera (Olympus) operated by TIA software. Gold probe density of transduced calyces, non-transduced calyces, and principal cells was calculated by counting the number of gold particles contained within the area of each terminal or cell and dividing by that area. Raw mitochondrial intensity was calculated using Fiji by

demarcating each mitochondrion within a terminal contained in a single section and calculating

the mean pixel intensity across all mitochondria in the terminal.

SBF-SEM sample processing and data analysis

Fixed brain slices of 100 µm thickness were trimmed to contain a single MNTB region and reacted

with DAB and H_2O_2 as above. The tissue pieces were incubated in an aqueous solution of 2%

osmium tetroxide and 1.5% potassium ferrocyanide for 30 minutes at 4˚C in the dark. Tissue was

washed with water, which was repeated between each of the following steps. Tissue was then

reacted with 0.2% tannic acid in water for 30 minutes, with 1% aqueous osmium tetroxide for 30

minutes, with 1% aqueous thiocarbohydrizide for 20 minutes, and another treatment of 1% osmium

tetroxide for 30 minutes all at room temperature, then placed in water overnight. The following

day, tissue was stained *en bloc* with 1% phosphotungstic acid in 25% ethanol at 50˚C for 30

minutes, then with 1% uranyl acetate in 25% ethanol for 30 minutes at 4˚C, and Walton's lead

aspartate for 30 minutes at 60˚C. At P7, staining of MNTB using this protocol was optimal for

imaging. At P21, however, staining was too intense due to massive axonal myelination which

readily recruits osmium. Therefore, ethanolic uranyl acetate staining was reduced to 10 minutes,

and the tannic acid and second osmification steps were omitted. Infiltration and embedding were

performed as above, taking extreme care in handling the brittle samples. Since SBF-SEM requires

the sample to be conductive to minimize charging during imaging, the tissue was trimmed to $1 \times$

0.5 mm and one side was exposed using an ultramicrotome (UC7, Leica), then turned downwards

to be remounted to a metal pin with conductive silver epoxy (CircuitWorks, CHEMTRONICS).

The new surface was then exposed and sputter-coated with a 5 nm layer of platinum. Samples were

then sectioned and imaged using 3View and Digital Micrograph (Gatan Microscopy Suite,

RRID:SCR_014492) installed on a Gemini SEM300 (Carl Zeiss Microscopy LLC.) equipped with

an OnPoint BSE detector (Gatan, Inc.). The detector magnification was calibrated prior to imaging

within SmartSEM imaging software (Carl Zeiss Microscopy LLC.) and Digital Micrograph with

a 500 nm cross line grating standard. Imaging was performed at 1.4 kV accelerating voltage,

4.45 kV extractor voltage, 30 µm aperture, working distance of ~5 mm, 0.5 µs pixel dwell time,

6.5 nm per pixel, knife speed of 0.1–0.3 mm/sec with oscillation, and 50 or 70 nm section

thickness. Images were captured at an original size of 12,000 by 12,000 pixels. The volume

analyzed for whole cell reconstruction at P7 was approximately $78 \times 78 \times 52$ µm (3.16×10^5 µm^3);

the volume analyzed for P21 was approximately $78 \times 78 \times 79$ µm, (4.81×10^5 µm^3), which was

approximately the entire thickness remaining from the 100 µm tissue slice following trimming.

Serial images were aligned using Digital Micrograph software and converted to TIFF format or

exported as TIFFs to TrakEM2 (RRID:SCR_008954) and aligned using Scale-Invariant Feature

Transform image alignment with linear feature correspondences and rigid transformation (Lowe,

2004).

For segmentation of whole terminals, images were downsampled by a factor of four to

reduce computational overhead, and only every third image was used to speed up segmentation.

Aligned images were exported in TIFF format to Microscopy Image Browser (MIB;

RRID:SCR_016560, (Belevich et al., 2016) for semi-automated segmentation of the calyx using

the watershed brush tool with interpolation. Following segmentation, interpolation was visually

verified in each section. Segmentations were saved separately and then converted to a mask for

mitochondrial segmentation, allowing extract features specific to the terminal. Dark features were

then thresholded in 3D and objects in 3D were morphologically filtered using the statistics function

until only mitochondria remained. Segmented mitochondria were smoothed and manually proofed

in each section. Volume measurements for terminals and mitochondria were calculated from

meshed surfaces of the segmented data, beginning at the base of the axon, identified as the location

where the area of the calyx cross section began to open into the main portion of the terminal. Axon

measurements were taken from the opening of the terminal to where the axon left the volume.

Only calyx terminals that were fully contained within the imaged volume were analyzed, resulting

in six calyces and eight minor terminals each at P7 and P21. Scale bars presented in figures are

extracted from the original EM micrographs.

For full resolution analysis of the ultrastructure of the subcompartments, one calyx at P7

(Fig. 5D far left) and two calyces at P21 (Fig. 5E far left and far right) were sampled. A P7 the

calyx exists largely as a cup structure, while the P21 terminal can be divided into stalk and swelling

subcompartments. We used the definition of stalk and swelling as defined by (Wimmer et al.,

2006) and (Grande and Wang, 2011). An ROI measuring $6.5 \times 6.5 \times 7$ µm was used to sample the

axons, cup, and stalks, as they are not discrete structures. Swellings could be defined as discrete

cytoplasmic compartments containing active zones, SVs, and typically mitochondria, having at

least one thin neck connecting it to another compartment, and therefore could be sampled as

distinct structures. One axon and terminal (cup) were sampled at P7. Two axons (one per calyx),

four stalks (two per calyx), and 22 swellings (16 and 6 per calyx) from two calyces were sampled

at P21.

For the partial volume at P7, a series of 300 sections at native XYZ resolution (6.5 nm per

pixel, 50 nm thick) was used to semi-automatically segment the calyx terminal and mitochondria.

A complete stack of aligned TIFF images was cropped to the size of the terminal using Fiji, then

exported to MIB. Gaussian smoothing was performed on the images and the terminal was

segmented using the semi-automatic graphcut segmentation function. Segmentations were proofed

manually in each section.

For all reconstructions, Amira (version 6.5, Amira 3D analysis, RRID:SCR_014305) was

used as a measurement tool, visualization tool, and video creation tool. Blender (version 2.79, 2.8;

RRID:SCR_008606) was used to create 3D renderings of terminals.

ssSEM sample processing and data analysis

Tissue slices were processed with the same method as for TEM above. Once embedded, samples

were trimmed to a roughly 2×3 mm rectangle, sectioned at 50 nm using the ATUMtome (RMC

Boeckeler, Tuscon, AZ), and collected on a reel of Kapton tape. Tape containing sections was

attached to a 4-inch silicon wafer in rows using double sided carbon tape (EMS, Cat# 77817-50)

and sections were post-stained with uranyl acetate and lead citrate. Wafers were coated with a

5 nm layer of carbon before imaging to limit charging (Leica, Wetzlar, Germany, ACE600). Image

acquisition was performed semi-automatically using a Merlin VP Compact SEM equipped with

Atlas 5 AT software (Carl Zeiss Microscopy LLC., Oberkochen, Germany). Overview images of

sections were acquired and registered using SE2 signal at 2 µm per pixel. Medium magnification

images of the general region of interest were acquired to identify potential synaptic terminals at

30 nm per pixel using a BSE detector. High magnification images of targets were captured using

an Inlens Duo detector at an accelerating voltage of 1.9 kV, 4.45 kV extractor voltage, 60 µm

aperture, ~6.8 mm working distance, ~12.8 µs pixel dwell time, and 5 nm per pixel. The detector

was calibrated prior to imaging using a 500 nm cross line grating replica. Images were batch

exported as TIFF files and aligned automatically using the TrakEM2 plugin using Scale-Invariant

Feature Transform image alignment with linear feature correspondences and rigid transformation.

Microscopy Image Browser (MIB) was used to semi-automatically segment calyx of Held

terminals using the watershed brush tool. Amira was used for analysis of the 3D vesicle pool,

vesicle docking, and extraction of AZ surface areas; as a visualization and video creation tool. All

vesicles within a 200 nm distance from AZ were manually selected from the image stack, and a

custom script was used to calculate the nearest distance from each vesicle to the surface of the AZ.

AZ surface areas were extracted from the presynaptic side of the triangular mesh formed following

reconstruction. Estimates of the total number of AZs and PA were quantified using MIB.

Controls for EM analysis

Internal controls were taken from the non-transfected MNTB ipsilateral to the injection site, or, if

expression was not ubiquitous in the contralateral MNTB, individual cells identified as being

negative for EGFP expression and by the lack of electron dense mitochondria (compared to nearby

infected cells) were used. For post-embedded samples, lack of gold particles labeling EGFP was

also used to confirm a terminal was uninfected. We did not observe ipsilateral expression of EGFP

or mito-APEX2 in any processed tissue by fluorescence, DAB reaction product, or EM dense

staining at the mitochondrion. For linescan analyses, principal neuron mitochondria were used as

a staining control. For the immuno-EM study, antibody specificity was examined by omitting

primary antibody from the diluent solution during the staining protocol.

Experimental Design and Statistical analysis

For the electrophysiological data analysis, individual neurons were considered independent

samples. For immuno-TEM analysis, different cell compartments from one MNTB region (i.e.

calyx terminals or principal cell bodies) were considered independent samples. For thin-section

TEM analysis, AZs were considered independent samples 120 total AZs, (3 animals, 40 AZs each).

Data were analyzed in MATLAB (version 9.4, RRID:SCR_001622), GraphPad Prism

(RRID:SCR_002798), R Project for Statistical Computing (version 3.5.2, RRID:SCR_001905),

and JASP (version 0.9.2.0, RRID:SCR_015823). Data distributions were tested for Gaussianity

using the Shapiro-Wilk test. To compare two groups, a two-tailed unpaired Student's t test with

Welch's correction (normal distribution) or two-tailed Mann-Whitney U test (non-normal

distribution) were used. Immuno-TEM data groups were compared using Mood's median test.

Linear regressions of total mitochondria volume to terminal volume were fit through the origin of

the axes. For mitochondria volume ratio measurements of calyx terminals, a two-way repeated

measures ANOVA was conducted to compare the effects of age and compartment on volume ratio.

Mitochondria diameter was compared using one-way ANOVA. P-values from multiple

comparisons were adjusted using the Bonferroni method. Data are reported as mean ± standard

error of the mean, unless otherwise stated. For interpretation of all data, a P-value of 0.05 was

deemed significant. P-values in figures are denoted *P<0.05; **P<0.01; ***P<0.001;

****P<0.0001. Since non-significant P-values do not provide evidence for the lack of effect or to

draw conclusions about the probability of the null hypothesis (H0) over the alternative hypothesis

(H1), we additionally calculated the Bayes Factor using a Cauchy distribution with scale factor 1

as the prior distribution implemented as described previously (Rouder et al., 2009). The Bayes

Factor is reported as BF_{10} for immuno-TEM indicating the likelihood of the data under H1

(treatment has an effect) relative to H0 (treatment has no effect), where values larger than 1 are in

favor of H1. For thin-section-TEM and electrophysiology, BF_{01} is reported, indicating the

likelihood of the data under H0 (treatment has no effect) relative to H1 (treatment has an effect),

where values larger than 1 are in favor of H0. Effect sizes were calculated using the MES toolbox

in MATLAB (Hentschke and Stuttgen, 2011) and are reported as Cohen's U_1 (proportion of non-

overlap of the two distributions) for two-sample comparisons and eta-squared (η^2, the proportion

of the total variation explained by the variation between groups) for multiple-sample comparisons.

Confidence intervals of effect sizes were calculated using bootstrapping with 10,000 repetitions.

No statistical test was performed to pre-determine sample sizes. Exact *P*-values, test statistics,

Bayes Factors, and effect sizes for all statistical comparisons are summarized in Table 1.

Data and Software Accessibility

All experimental data, movies, 3D reconstructions and custom-written software central to the

conclusion of this study are available at http://dx.doi.org/10.17632/v88r5t5myz under the terms of

the Creative Commons Attribution 4.0 License (CC BY 4.0).

Results

HdAd EGFP mito-APEX2 transduction leads to efficient labeling of mitochondria at the calyx of Held

To specifically label mitochondria in the calyx we created an HdAd which expresses mito-APEX2 and EGFP under the control of the hsyn promoter (HdAd mito-APEX2) (Fig. 1A1). This dual expression allowed for identification of transduced calyces with both electron and fluorescence microscopy. Stereotactic injection of HdAd mito-APEX2 into the anteroventral cochlear nucleus (aVCN) of C57Bl/6J mice was performed at postnatal day 1 (P1) as previously described (Chen et al., 2013) and experiments were carried out on P7 or P21 calyces representing the immature and mature calyx, respectively (Fig. 1A2) (Borst and Soria van Hoeve, 2012). HdAd injection into the aVCN resulted in the transduction of calyces which were identified by EGFP expression (Fig. 1B). We confirmed that these EGFP positive terminals contained APEX2-tagged mitochondria by carrying out DAB staining which resulted in dark reaction products in resin-embedded tissue under brightfield illumination. As expected, only the MNTB contralateral to the injection site contained DAB reaction products (Fig. 1C, D). At high magnification, individual calyx terminals were visible as areas of dark deposition surrounding the somata of principal cells (Fig. 1E, F). To confirm that these electron dense precipitates were sufficient to identify mitochondria in the transduced calyx terminals, we imaged ultrathin sections using TEM. We found that presynaptic calyx terminals in both P7 and P21 samples contained electron-dense mitochondria with DAB precipitates accumulated in the mitochondria matrix while the postsynaptic cell bodies did not (Fig. 1G-I). Additionally, labeled calyx mitochondria were easily detectable and distinguishable from unlabeled calyx mitochondria (Fig. 1J, K). This indicates that expression of mito-APEX2 using HdAd is sufficient to label mitochondria for EM applications.

EGFP and mito-APEX2 dual expression does not alter presynaptic ultrastructure

To confirm and characterize the dual expression of EGFP and mito-APEX2 we performed post-

embedding immunogold labeling of EGFP on ultrathin sections from P21 calyces labeled for mito-

APEX2 (Fig. 2). The cytoplasm of all calyx terminals containing electron dense mitochondria also

contained gold particle labeling of EGFP (Fig. 2A, B). To quantify the expression of EGFP and

mito-APEX2, we analyzed gold particle density in calyces containing labeled mitochondria

compared to non-transduced calyces and principal cells (Fig. 2C). Immunogold particle density

was highest in terminals containing mito-APEX2 labeled mitochondria (terminal

control $= 2.70 \pm 0.37$ particles/μm^2, principal cell control $= 3.55 \pm 0.34$ particles/μm^2 vs. EGFP-

mito-APEX2 $= 14.48 \pm 1.10$ particles/μm^2, $P = 0.00015$; Fig. 2C, Table 1). Although mito-

APEX2 was constrained to mitochondria, we tested if its expression affected presynaptic

ultrastructure. Using ultrathin sections from P21 animals, we analyzed active zone (AZ) length,

SV distribution, and docked SV number (within 5 nm of the presynaptic membrane)

(Taschenberger et al., 2002) for both mito-APEX2 expressing calyces and their respective in-slice

control. We observed no significant difference in any measured parameter between control and

mito-APEX2 expressing terminals (Fig. 2D-F and Table 1).

Mito-APEX2 expression does not alter synaptic transmission

HdAd mito-APEX2 lead to efficient mitochondria labeling in the calyx and ability to visualize

presynaptic ultrastructure. However, it has been shown that extreme levels of APEX expression

can impact mitochondria structure (Martell et al., 2017). Since proper mitochondria function is

integral for synaptic transmission and plasticity (Billups and Forsythe, 2002; Cserep et al., 2018;

Lee et al., 2018), we tested if mito-APEX2 expression impacts synaptic transmission. First, we

performed afferent fiber stimulation of transduced calyces at P7 and recorded whole-cell AMPAR

currents of MNTB principal cells using 50 Hz stimulation. We did not find evidence for altered

synaptic transmission compared to age-matched WT control (Fig. 3A, Table 1). Finally, we tested

if chronic expression of HdAd mito-APEX2 impacted synaptic transmission and repeated the

experiments with P21 animals. Using afferent fiber stimulation at 300 Hz, we found that chronic

mito-APEX2 expression did not alter synaptic transmission (Fig. 3B, Table 1). These data

combined with the ultrastructural measurements indicate that chronic expression of mito-APEX2

does not alter presynaptic ultrastructure or synaptic transmission.

Large scale volumetric analysis of calyces and mitochondria with SBF-SEM

Due to the large size of the calyx, volumetric analysis at fine-scale EM resolution of mitochondria

has been restricted to partial reconstructions of single calyces using TEM (Rowland et al., 2000;

Wimmer et al., 2006), ssSEM (Horstmann et al., 2012), or electron tomography (Perkins et al.,

2010). Furthermore, ultrathin section-based imaging platforms are constrained by alignment issues

of single sections which hampers large scale volumetric analysis (Lichtman and Denk, 2011). In

contrast, SBF-SEM requires minimal image alignment and is therefore ideal to efficiently image

large EM volumes. Although SBF-SEM has been successfully used for calyx reconstructions

(Holcomb et al., 2013; Xiao et al., 2013) it requires liberal application of heavy metals, which may

prevent detection of mito-APEX2 labeling. To test this, we used the 3View with OnPoint BSE

detector and optimized the imaging conditions (see Methods) to confirm that mito-APEX2 labeled

mitochondria could be distinguished in both P7 and P21 (Fig. 4). We found calyx terminals

containing mito-APEX2 positive mitochondria were clearly identifiable in both P7 and P21

samples as the electron density of labeled mitochondria in the calyx terminal was considerably

higher than that of unlabeled mitochondria in the principal cell (Fig. 4E, F). Although we could

visualize SVs at this resolution, identification of AZs (using the postsynaptic density as a

reference) and PAs was difficult due to the lower signal-to-noise ratio of images acquired with

short dwell times (Fig 4B, D).

Subsequently, we acquired and analyzed volumes of $78 \times 78 \times 52$ µm (P7, Fig. 5A) and 78

$\times 78 \times 79$ µm (P21, Fig. 5B) at an XY pixel size of 6.5 nm and section thickness of 70 nm. We

found that a pixel dwell time of 0.5 µs provided sufficient image quality for tracing membranes

while reducing imaging time, which allowed us to collect the full volume in ~30 hours per sample.

We segmented individual positive calyx terminals that were within the bounds of the datasets and

carried out whole terminal reconstructions. To increase segmentation throughput and minimize

dataset size, images were downsampled and every third image was used. In the acquired image

volume, all six calyx terminals containing mito-APEX2 positive mitochondria plus eight labeled

minor inputs were segmented at P7 in ~40 working hours. Due to the complex calyx structure in

P21 animals, segmentation of six terminals and eight labeled minor inputs required ~100 working

hours. In those volume data sets, the reconstructed terminals at P7 and P21 showed clear

morphological differences as previously described (Satzler et al., 2002; Taschenberger et al., 2002;

Wimmer et al., 2006; Grande and Wang, 2011; Horstmann et al., 2012; Holcomb et al., 2013) (Fig.

5A, B, and D, E, Movie 1 and 2). At P7, several filopodial extensions branched off from the cup-

like terminal (Fig. 5D). These cups were thin and sheet-like, while terminals at P21 were composed

of mainly thick, bulbous compartments without filopodial extensions (Fig. 5E). We measured the

volume of individual calyx terminals containing mito-APEX2 positive mitochondria and found

that calyx volumes at P21 were larger than at P7 (1307 ± 124 µm^3 vs. 841.2 ± 136.1 µm^3, $P = 0.03$;

Table 2, Fig. 5C). The terminals at P7 had volumes between 9 and 1171 µm^3 and 5 of the 6 MNTB

principal neurons were contacted by a large calyx presynaptic terminal (>700 µm^3) with minor

terminals, while one principal neuron had a dominant, but smaller calyx terminal of ~250 µm^3 and

multiple minor terminals ($<100\ \mu m^3$) (Fig. 5A, D, yellow terminals, Movie 1). Consistent with our

data, principal cells with small dominant calyx and multiple minor competing terminals are rare at

P7, comprising only roughly 20% of cells (Holcomb et al., 2013).

At P21, terminals ranged from 10 to 1620 μm^3. We found that all (6/6) principal cells at

this age were contacted by a single dominant calyx similar to previous reports (Holcomb et al.,

2013) and exhibited a markedly different structure with variable degrees of fenestration as

described previously (Grande and Wang, 2011; Wang et al., 2015); Fig. 5E, Movie 2). In addition,

half of those principal cells (3/6) had mito-APEX2 labeled minor terminals (Fig. 5E, Table 2),

with morphologies similar to "migratory calycigenic axons" in the young opossum (Morest, 1968)

and calyceal collaterals in rodents (Rodriguez-Contreras et al., 2006). Notably, one principal cell

had a moderately sized terminal (272 μm^3) in addition to minor terminals (Fig 5B, E, green

terminals). Therefore, we conclude that even though principal cells are innervated by a dominant

calyx, minor terminal still exist in the functionally mature stage of development, which might be

comprised by calyceal collaterals. Finally, in addition to the labeled minor terminals, numerous

small unlabeled terminals contacted the same principal neurons at both ages but were not

segmented or analyzed. These terminals may compose additional excitatory or inhibitory inputs to

the MNTB principal cell that do not originate from the cochlear nucleus (Hamann et al., 2003;

Awatramani et al., 2004; Albrecht et al., 2014).

Mitochondrial volume ratios are higher in the mature calyx and axon than the immature

calyx

Since the high-contrast mitochondria could be easily extracted using thresholding techniques, we

carried out high resolution whole terminal mitochondria reconstructions at P7 and P21 using the

same volume data sets acquired with SBF-SEM. After downsampling the volume data and using

every third image, whole terminal mitochondrial volume reconstructions could be completed and

proofed in several hours per terminal (Fig. 6A-D, Table 2, Movies 3 and 4). Since at both

developmental stages, principal cells had labeled minor terminals, we separately analyzed

mitochondrial volume in the calyx terminal and the minor terminals. Mitochondrial volumes scaled

linearly with terminal volume at both developmental time points. However, we found that the

scaling of mitochondria to calyx volume is steeper at P21 than at P7 (Slope = 0.2022, R^2 = 0.95

vs. 0.0737, R^2 = 0.82; Fig 6E). Comparison of the mitochondria to volume ratio of the terminal

revealed that 20% of the P21 calyx volume is composed of mitochondria, while at P7 mitochondria

only take up 7.8% of the terminal—a ~2.6-fold increase throughout development ($P < 0.0001$;

Fig. 6E, Table 2). Mitochondrial volumes in the minor terminals at both P7 and P21 showed a

similar linear scaling of mitochondrial volumes (Slope = 0.1883, R^2 = 0.99 vs. 0.1066, R^2 = 0.95)

and revealed a ~40% increase in the mitochondrial volumes at P21 ($P = 0.0245$; Fig. 6F).

During maturation, the calyx transforms into a high-fidelity synapse that can sustain action

potential (AP) firing upwards to a kilohertz (Baydyuk et al., 2016). Since AP firing is an

energetically expensive process (Harris et al., 2012), this suggested that mitochondria levels are

increased in the axon also. To determine if mitochondria volumes are increased in the axon to

support the increased firing rates during development, we analyzed the mitochondrial volumes in

the calyx axon. We found that at P7 mitochondria composed 6.4% of the axon volume, while at

P21 the mitochondria volume was increased 2-fold, comprising 12.5% of axon volume

($P = 0.0003$). When comparing the relative mitochondrial volumes in the calyx terminal vs. axon,

we found that the calyx had significantly higher relative mitochondria volumes than the axon at

P21 ($P = 0.0009$), while there was no difference between the P7 terminal and axon ($P = 0.7116$;

Fig. 6E, Table 2).

Taken together, our results revealed that mitochondrial volumes increase in both the calyx

and its axon during maturation. Furthermore, mitochondrial volumes in the mature calyx are

selectively increased compared to its respective axon, resulting in an asymmetry not found in the

immature calyx. Therefore, we conclude that increases in mitochondrial volumes support the

energetic demands required for high fidelity synaptic transmission.

Stalks and swellings of the mature calyx have similar relative mitochondrial volumes

In the mature calyx, Ca^{2+} transients are higher in swellings compared to stalks, and stalks have

higher *Pr* than swellings (Fekete et al., 2019). Since mitochondria can act as Ca^{2+} buffers, this

suggests mitochondria volumes may be lower in swellings compared to stalks. In addition,

swellings have more loosely coupled SVs than stalks which are proposed to be in a less

energetically favorable configuration (Eggermann et al., 2012) which may impact mitochondrial

volumes between the two subcompartments. To determine mitochondrial volumes in these

subcompartments we segmented small subvolumes at full resolution and reconstructed

mitochondria in 4 stalks and 22 swellings from two calyces (P21, far left and far right calyces in

Fig. 5E; 7C-E). Subsequent analysis revealed that stalks and swellings contain similar relative

volumes of mitochondria (stalks = 17.5% vs. swellings = 21.4%, Fig. 7D, E, Table 2).

Since we found that mitochondrial volumes in P21 calyces differ between axons and

terminals (Fig. 6E), we further analyzed mitochondrial morphology by measuring the

mitochondria diameter in axons (P7, P21), cup (P7), stalks and swellings (P21) from the same full

resolution reconstruction of subcompartment data (Table 2). At P7, we found no difference

between the mitochondrial diameters between axon and cup (0.4 ± 5 nm larger in the axon,

$P > 0.9999$; Fig. 7F). Within subcompartments at P21 we found that stalks and swellings contained

larger mitochondria diameters compared to the axon (44 ± 2 nm larger in stalks, $P < 0.0001$; 40 ± 2

nm larger in swellings, $P < 0.0001$), but found no difference between stalk and swellings (4 ± 2

nm larger in stalks, $P = 0.1173$). Across developmental stages, we observed an increase in the

mitochondria diameter between P7 and P21 axons (21 ± 4 nm larger at P21, $P < 0.0001$) (Fig. 7F).

Taken together, our findings reveal that stalks and swellings have similar mitochondrial volumes

and diameters, and mitochondrial diameters are larger in P21 than in P7 calyces. Therefore, we

conclude that differences in Ca^{2+} buffering between stalks and swellings are not caused by

differences in mitochondrial volumes. Furthermore, the developmental increase in mitochondrial

volumes is largely due to an increase in mitochondria size.

At P7, calyces had multiple filopodial extensions with various morphologies as shown in

Figures 5A and 5D, some of which sparsely contained mitochondria (Fig. 6A, C). To further

analyze the mitochondria distribution in these extensions we performed another full resolution

partial reconstruction of a P7 calyx and overlaid its segmented mitochondria (Fig. 8, Movie 5). In

this terminal, extensions from the cup-like calyx rarely contained mitochondria (Fig. 8B, D). In

addition, we found multiple invaginations into the calyx from the principal cell, including large

excrescences with complex membrane topologies (white arrows, Fig. 8A, C). These excrescences

were largely composed of principal cell processes, though some processes could be traced back to

the calyx terminal (Movie 5). Similar to filopodial extensions, these structures lacked

mitochondria. This indicates that at P7 mitochondria are mostly restricted to the cup.

Mito-APEX2 in conjunction with ATUM ssSEM allows for fine scale 3D analysis of

presynaptic ultrastructure

In addition to using SBF-SEM in combination with mito-APEX2 to analyze mitochondrial

volumes, one of our goals was to use mito-APEX2 as a genetic tool to analyze presynaptic

ultrastructure in a defined population of genetically labeled neurons. However, we were unable to

reliably detect the AZ with SBF-SEM. Therefore, we used ATUM-ssSEM to obtain SEM images

with higher signal to noise ratio. We first established protocols to reliably detect mito-APEX2-

labeled mitochondria with ATUM ssSEM and then used mito-APEX2 to carry out partial

reconstructions of multiple AZs in the calyx.

Using our protocols, we could clearly identify labeled mitochondria in calyx terminals with

detailed synaptic structure after post-staining sections to improve signal-to-noise ratio (SNR) for

better membrane visualization in both P7 (Fig. 9A, B) and P21 (Fig. 9C, D, see Methods). Imaging

with ssSEM was performed with several sessions at progressively higher resolutions. Beginning

with a resolution of 2 µm/pixel, we collected images covering each section on a wafer, which

required roughly a few hours. Following this, we imaged a broad region of interest in the middle

of the MNTB containing dozens of principal cells at 30 nm/pixel resolution and captured ~500

images to discriminate mito-APEX2-labeled terminals and to identify potential sites containing

the fewest section wrinkles. Finally, we captured high resolution images at 5 nm/pixel resolution

and a pixel dwell time of ~13 µs, which was sufficient for analysis of synaptic specializations (Fig.

9B, D top). We morphologically differentiated AZs (Fig. 9B, D, top) from puncta adherentia (PA)

(Fig. 9B, D, bottom) and separately quantified them for each calyx. Similar to SBF-SEM, we

confirmed that calyx terminals containing mito-APEX2 positive mitochondria could be

differentiated in both P7 and P21 ATUM-ssSEM samples (Fig. 9E, F).

Next, using 100 high-resolution serial sections captured from both P7 and P21 samples

expressing mito-APEX2, we created partial reconstructions and carried out volumetric analysis for

two calyces at each age (Fig. 10). The estimated surface areas of the partially reconstructed calyces

facing the soma ranged from 117 to 269 µm². (Fig. 10A, B, Table 2, and Movie 6 and 7). At P7,

the number of PAs was approximately two-fold higher than AZs (37% AZ vs. 63% PA; Fig. 10A,

Table 2), similar to previously reported values in P9 rat (Satzler et al., 2002). At P21 we observed

a similar proportion of 40% AZ to 60% PA (Fig. 10B, Table 2).

Since both AZ size and SV distributions change during early development (Taschenberger

et al., 2002), we analyzed AZ area and SV distributions of 10 individual AZs (5/calyx) from 3D

reconstructed models at P7 and P21. We found larger AZ at P7 compared to P21 (P7:

$0.203 \pm 0.040 \, \mu m^2$, n = 10 vs. P21: $0.066 \pm 0.008 \, \mu m^2$, n = 10; Fig 10C, D, E, and Table 2).

Comparing SV distribution within 200 nm of the AZ we found that AZ at P7 contained more SVs

than P21 (P7: 128.2 ± 18.55, n = 10; vs. P21: 92.9 ± 5.363, n = 10; Fig. 10F, G, Table 2). However,

when normalizing the SV number to AZ area to correct for the difference in AZ size, we found

that the P21 AZ contained more SVs within 200 nm per AZ unit area (Fig. 10G). We observed no

difference in overall SV distribution between P7 and P21 in a 200 nm distance from AZs in 3D

space (Fig. 10H). At both ages, docked SVs (<5 nm to AZ) per AZ were similar (Table 2) and

comparable to previous reports (Satzler et al., 2002; Taschenberger et al., 2002; Dondzillo et al.,

2010), however we did not observe a developmental decrease in contrast to a previous study

(Taschenberger et al., 2002).

Discussion

To interrogate how mitochondria volumes are regulated during the development of the calyx of

Held presynaptic terminal we developed a new tool, HdAd mito-APEX2, and optimized protocols

and visualization workflows for use with SBF-SEM. We found that during development

mitochondrial volumes increase in both the calyx and its respective axon, but that the mature calyx

contains a larger relative mitochondrial volume than the axon. In addition, there was no difference

in the relative mitochondrial volumes in the stalk and swellings of the mature calyx. Therefore, we

propose that mitochondria levels are developmentally increased at the calyx of Held and its

respective axon to support the energetic demands of high-frequency firing required to encode

auditory information but do not significantly contribute to the calyx's morphological-functional

diversity. Finally, we believe as a technical advancement, that our HdAd vector in combination

with optimized protocols for ATUM-SEM and SBF-SEM will be of broad use in combination with

molecular perturbations studies using LM and EM inside and outside of the neuroscience

community.

A tubular network of mitochondria throughout development

Whole-terminal volumetric reconstructions of mitochondria at the two developmental stages

revealed a complex tubular network in the P7 and P21 calyx. These findings of many long finger-

like mitochondria are in line with previous reports from partial calyx reconstructions from P8 and

P22 rats (Wimmer et al., 2006) and adult cat (Rowland et al., 2000) using ultra-thin section TEM.

Long tubular mitochondria have been observed in multiple neuron types and their respective axons

and across species using different imaging modalities, EM or fluorescence microscopy in both live

cell and fixed tissues (Brodin et al., 1999; Misgeld et al., 2007; Saxton and Hollenbeck, 2012;

Rossi and Pekkurnaz, 2019). However, the extensive tubular network might be specific to large

synapses with high energy demands since many conventional presynaptic terminals in the CNS

have a mitochondrion at or near the presynaptic terminal (Devine and Kittler, 2018). Since

mitochondria can be functionally coupled (Amchenkova et al., 1988; Saxton and Hollenbeck,

2012), the tight packing and interconnected mitochondria observed in this and previous studies

(Wimmer et al., 2006) suggest that mitochondria in the calyx and many other large presynaptic

terminals are functionally coupled.

Mitochondria become larger during calyx development

Based on our finding that mitochondria are more densely packed and mitochondria volumes are higher in the mature calyx compared to immature calyx, we propose that a developmental increase in mitochondrial volumes is critical for high-frequency transmission. Qualitative analysis of P7 calyces revealed large patches of the calyx without mitochondria while this was not found at the P21 calyx. Quantitative analysis of mitochondria diameters revealed that mitochondrial diameters were larger at the P21 calyces than those at P7. Though we found larger mitochondria at the mature stage, an increase in the total number of mitochondria may also contribute to the increase in relative volume. Obtaining absolute counts of mitochondria is difficult due to their ability to fuse and divide. However, qualitatively it appears that P21 calyces also contain more mitochondria. Although spontaneous activity in the auditory brainstem increases during early development (Tritsch and Bergles, 2010), the increase in mitochondrial volumes was larger at the calyx (~3-fold), compared to the secondary inputs (~1.4-fold) on the MNTB principal neuron. However, at P7 minor terminals had higher relative mitochondrial volumes (~2.5-fold) compared to the calyx.

What are the signals that promote this transformation? During calyx development, synaptic activity increases ~10-fold which lead to high rates of SV release and replenishment (Baydyuk et al., 2016). Thus, a probable signal is synaptic activity which is closely linked to mitochondria function (Nguyen et al., 1997; Rossi and Pekkurnaz, 2019). Total mitochondrial volume correlates with levels of exocytosis (Ivannikov et al., 2013), while individual mitochondrial volumes are larger at high-activity synapses (Cserep et al., 2018). In addition, mitochondria are more likely to pause during high activity levels and reside in the same position over a time course of days (Obashi and Okabe, 2013). Therefore, it possible that the increase in calyx activity increases the residency time of mitochondria in the terminal.

681   Comparison of axonal mitochondria volumes revealed that mitochondria were also more

682 densely packed in P21 compared to P7. This indicates that increases in axonal excitability during

683 development (Leao et al., 2005) are supported by increased mitochondria number to meet the

684 energy demands of AP firing (Harris et al., 2012). Unlike the P7 axon, the mature calyx axon is

685 myelinated except for the nodes of Ranvier and the heminode. Na^+ channels are highly clustered

686 at the heminode and excluded from the terminal (Leao et al., 2005). Although the last heminode

687 in the mature calyx axon is well-defined using confocal imaging (Berret et al., 2016; Xu et al.,

688 2017), we had difficulty defining the heminode from the myelination morphology only. Despite

689 this, we were unable to detect any local increases in mitochondrial densities along the last 5-50 µm

690 of the axons in the reconstructed volume. Therefore, our data indicate that mitochondria levels in

691 the axon are globally increased without higher densities at the heminode. However, since

692 mitochondrial movement is on the order of seconds it is possible that mitochondria pause

693 movement at these heminodes during AP firing to create temporary increases in density which

694 may not be detectable due to the timescale of aldehyde fixation.

695   We also found that mitochondrial volumes in the mature calyx were greater than its

696 respective axon, while this was not found in the P7 calyx. This finding suggests SV release and

697 replenishment in the mature calyx has higher energetic demands than AP firing, while the energetic

698 demands at P7 maybe equivalent between the calyx and its respective axon. The mitochondrial

699 adherens complex (MAC) is located within the mature calyx but has not been reported in the axon

700 (Rowland et al., 2000). Therefore, it is possible that the MAC contributes to the increase in

701 mitochondrial volume at the mature calyx due to its tethering of mitochondria. However, due to

702 the limitations of SBF-SEM we were unable to identify the MAC and determine if the complex

appearance of mitochondria in the calyx is developmentally regulated. Future studies will need to

be carried out to resolve this issue.

Mitochondria and their contribution to the calyx morphological-functional continuum

Although the calyx is considered a reliable relay synapse (Englitz et al., 2009; Lorteije et al., 2009;

Stasiak et al., 2018) a morphological-functional continuum of calyces that differ in the number of

stalks and swellings in the hearing animal has been observed (Grande and Wang, 2011; Wang et

al., 2015). Differences in Ca^{2+} buffering and *Pr* between stalks and swellings has been proposed

to expand the coding capacity of the calyx/MNTB synapse, (Fekete et al., 2019). Mitochondria

contribute to Ca^{2+} buffering (Billups and Forsythe, 2002; Guo et al., 2005; Verstreken et al., 2005)

which can impact SV release (Devine and Kittler, 2018). In particular at the prehearing calyx,

mitochondria slow the clearance of Ca^{2+} which can impact sustained synaptic transmission

(Billups and Forsythe, 2002). However, at the prehearing calyx, mitochondria contribution to

synaptic transmission is only prominent when the Na^+/Ca^{2+} exchanger is saturated (Kim et al.,

2005). Although swellings contain larger and faster Ca^{2+} transients and more Cav2.1 channel

clusters (Fekete et al., 2019), we found no difference in the mitochondria volumes between stalks

and swellings. Thus, we propose that the contribution of mitochondria to the morphological-

functional continuum is most likely negligible. However, it is possible that only under long periods

of high firing rates would mitochondria contribute to difference in Ca^{2+} clearance in these

subcompartments.

Identification of multiple inputs at the principal cells of the MNTB

Consistent with a previous report (Hoffpauir et al., 2006), we found multiple inputs innervating

the MNTB principal neuron at P7. Strikingly, at P21 we also found labeled minor terminals in

addition to a single calyx terminal. Although injection of HdAd with the human synapsin promoter

labels other cell types in the CN in addition to the globular bushy cells (GBCs, Montesinos et al.,

2011), none of these other cell types are reported to make synapses on MNTB principal cells (Cant

and Benson, 2003). Therefore, these labeled terminals are likely of GBC origin and distinct from

local excitatory and inhibitory inputs (Hamann et al., 2003; Albrecht et al., 2014). Due to limited

volumes imaged in this study we were unable to traces these axons to their origin. Since GBC

axons give rise to multiple collaterals which form other small terminals (coincident) onto MNTB

cells (Morest, 1968; Rodriguez-Contreras et al., 2006; Rodriguez-Contreras et al., 2008) it is

possible that they are the source of the minor terminals that we observed. Future studies will need

to be carried out to determine their role in regulating AP firing in the MNTB principal neuron.

HdAd Mito-APEX2 as presynaptic marker for morphological studies

HdAds are a dsDNA viruses and have faster onset of transgene expression than single-stranded

rAAVs which require second-strand synthesis (Hastie and Samulski, 2015; Montesinos et al.,

2016). We took advantage of this fast onset of expression and carried out the EM analysis as early

as P7, six days after HdAd injection, to analyze immature calyx ultrastructure. Although we did

not test earlier time points, HdAd vectors have led to successful transduction of calyces as early

as 72 hours (Montesinos et al., 2011). Since HdAd expression allows for long-term expression of

transgenes for years (Kim et al., 2001) and our data demonstrate lack of toxicity of synapsin

promoter driven expression of EGFP and mito-APEX2, this suggest that use of HdAd EGFP-mito-

APEX2 will enable long term morphological studies on the scales of months to possibly years.

More importantly, our HdAd vectors allows for the labeling of specific populations of neurons.

When using mitochondria for volumetric reconstruction, the cellular compartments of interest

should have a high mitochondria density for reliable distinction from unlabeled cells or

compartments.

ATUM ssSEM vs. SBF-SEM for presynaptic ultrastructure analysis

Use of HdAd mito-APEX2 with ATUM ssSEM allowed us to analyze presynaptic ultrastructure in addition to mitochondrial volumes. Both SBF-SEM and ATUM ssSEM have their strengths and weaknesses, therefore the different visualization platforms should be selected based on the target ultrastructure. Although SBF-SEM permitted large scale volumetric analysis, we were unable to segment fine AZ structures, such as the MAC or the puncta adherentia due to the low SNR required for consistent sectioning. Additionally, the low SNR prevented analysis of SV distribution. The low SNR was a result of the minimum acceptable beam dosage required to avoid sample damage and achieve consistent cutting of the block. Future improvements will need to be developed to enable better SNR in combination with thinner cutting thickness for fine structure analysis. Since ATUM ssSEM allows longer dwell times and therefore better SNR than SBF-SEM it was ideal for measuring SV distributions and AZ areas in a relatively large terminal. However, the tradeoff is the inherent issue with section alignment and possible compression artifacts, thus ATUM ssSEM is ideal for small scale volumetric reconstructions. Using ATUM ssSEM we found that the SV density per AZ area increases during development. While a previous study found a decrease in docked SV number from P5 to P14 (Taschenberger et al., 2002), we observed no change between P7 and P21. However, we only analyzed 10 AZs at each stage and differences in fixation protocols may lead to this discrepancy. Taken together, we believe that our viral vector and methods will be broadly applicable to quantitative analyses of subsynaptic organization of SVs, mitochondria, and their morphological continuum using TEM, ssSEM and SBF-SEM.

References

Albrecht O, Dondzillo A, Mayer F, Thompson JA, Klug A (2014) Inhibitory projections from the ventral
nucleus of the trapezoid body to the medial nucleus of the trapezoid body in the mouse. Front
Neural Circuits 8:83.

Amchenkova AA, Bakeeva LE, Chentsov YS, Skulachev VP, Zorov DB (1988) Coupling membranes as
energy-transmitting cables. I. Filamentous mitochondria in fibroblasts and mitochondrial clusters
in cardiomyocytes. The Journal of cell biology 107:481-495.

Awatramani GB, Turecek R, Trussell LO (2004) Inhibitory control at a synaptic relay. The Journal of
neuroscience : the official journal of the Society for Neuroscience 24:2643-2647.

Baydyuk M, Xu J, Wu LG (2016) The calyx of Held in the auditory system: Structure, function, and
development. Hear Res 338:22-31.

Belevich I, Joensuu M, Kumar D, Vihinen H, Jokitalo E (2016) Microscopy Image Browser: A Platform
for Segmentation and Analysis of Multidimensional Datasets. PLoS Biol 14:e1002340.

Berret E, Kim SE, Lee SY, Kushmerick C, Kim JH (2016) Functional and structural properties of ion
channels at the nerve terminal depends on compact myelin. The Journal of physiology 594:5593-
5609.

Billups B, Forsythe ID (2002) Presynaptic mitochondrial calcium sequestration influences transmission at
mammalian central synapses. J Neurosci 22:5840-5847.

Borst JG, Soria van Hoeve J (2012) The calyx of held synapse: from model synapse to auditory relay. Annu
Rev Physiol 74:199-224.

Brodin L, Bakeeva L, Shupliakov O (1999) Presynaptic mitochondria and the temporal pattern of
neurotransmitter release. Philosophical transactions of the Royal Society of London Series B,
Biological sciences 354:365-372.

Cant NB, Benson CG (2003) Parallel auditory pathways: projection patterns of the different neuronal
populations in the dorsal and ventral cochlear nuclei. Brain Res Bull 60:457-474.

Chen Z, Cooper B, Kalla S, Varoqueaux F, Young SM, Jr. (2013) The Munc13 proteins differentially
regulate readily releasable pool dynamics and calcium-dependent recovery at a central synapse.
The Journal of neuroscience : the official journal of the Society for Neuroscience 33:8336-8351.

Chen Z, Das B, Nakamura Y, DiGregorio DA, Young SM, Jr. (2015) Ca2+ channel to synaptic vesicle
distance accounts for the readily releasable pool kinetics at a functionally mature auditory synapse.
J Neurosci 35:2083-2100.

Cserep C, Posfai B, Schwarcz AD, Denes A (2018) Mitochondrial Ultrastructure Is Coupled to Synaptic
Performance at Axonal Release Sites. eNeuro 5.

Devine MJ, Kittler JT (2018) Mitochondria at the neuronal presynapse in health and disease. Nat Rev
Neurosci 19:63-80.

Dondzillo A, Satzler K, Horstmann H, Altrock WD, Gundelfinger ED, Kuner T (2010) Targeted three-
dimensional immunohistochemistry reveals localization of presynaptic proteins Bassoon and
Piccolo in the rat calyx of Held before and after the onset of hearing. The Journal of comparative
neurology 518:1008-1029.

Dong W, Radulovic T, Goral RO, Thomas C, Suarez Montesinos M, Guerrero-Given D, Hagiwara A,
Putzke T, Hida Y, Abe M, Sakimura K, Kamasawa N, Ohtsuka T, Young SM, Jr. (2018)

CAST/ELKS Proteins Control Voltage-Gated Ca(2+) Channel Density and Synaptic Release
Probability at a Mammalian Central Synapse. Cell reports 24:284-293 e286.

Eggermann E, Bucurenciu I, Goswami SP, Jonas P (2012) Nanodomain coupling between Ca(2)(+)
channels and sensors of exocytosis at fast mammalian synapses. Nat Rev Neurosci 13:7-21.

Englitz B, Tolnai S, Typlt M, Jost J, Rubsamen R (2009) Reliability of synaptic transmission at the synapses
of Held in vivo under acoustic stimulation. PloS one 4:e7014.

Fedchyshyn MJ, Wang LY (2005) Developmental transformation of the release modality at the calyx of
Held synapse. The Journal of neuroscience : the official journal of the Society for Neuroscience
25:4131-4140.

Fekete A, Nakamura Y, Yang YM, Herlitze S, Mark MD, DiGregorio DA, Wang LY (2019) Underpinning
heterogeneity in synaptic transmission by presynaptic ensembles of distinct morphological
modules. Nat Commun 10:826.

Grande G, Wang LY (2011) Morphological and functional continuum underlying heterogeneity in the
spiking fidelity at the calyx of Held synapse in vitro. The Journal of neuroscience : the official
journal of the Society for Neuroscience 31:13386-13399.

Guo X, Macleod GT, Wellington A, Hu F, Panchumarthi S, Schoenfield M, Marin L, Charlton MP, Atwood
HL, Zinsmaier KE (2005) The GTPase dMiro is required for axonal transport of mitochondria to
Drosophila synapses. Neuron 47:379-393.

Hamann M, Billups B, Forsythe ID (2003) Non-calyceal excitatory inputs mediate low fidelity synaptic
transmission in rat auditory brainstem slices. The European journal of neuroscience 18:2899-2902.

Harris JJ, Jolivet R, Attwell D (2012) Synaptic energy use and supply. Neuron 75:762-777.

Hastie E, Samulski RJ (2015) Adeno-associated virus at 50: a golden anniversary of discovery, research,
and gene therapy success--a personal perspective. Human gene therapy 26:257-265.

Hentschke H, Stuttgen MC (2011) Computation of measures of effect size for neuroscience data sets. The
European journal of neuroscience 34:1887-1894.

Hoffpauir BK, Grimes JL, Mathers PH, Spirou GA (2006) Synaptogenesis of the calyx of Held: rapid onset
of function and one-to-one morphological innervation. J Neurosci 26:5511-5523.

Holcomb PS, Hoffpauir BK, Hoyson MC, Jackson DR, Deerinck TJ, Marrs GS, Dehoff M, Wu J, Ellisman
MH, Spirou GA (2013) Synaptic inputs compete during rapid formation of the calyx of Held: a
new model system for neural development. J Neurosci 33:12954-12969.

Horstmann H, Korber C, Satzler K, Aydin D, Kuner T (2012) Serial section scanning electron microscopy
(S3EM) on silicon wafers for ultra-structural volume imaging of cells and tissues. PloS one
7:e35172.

Ivannikov MV, Sugimori M, Llinas RR (2013) Synaptic vesicle exocytosis in hippocampal synaptosomes
correlates directly with total mitochondrial volume. J Mol Neurosci 49:223-230.

Iwasaki S, Takahashi T (2001) Developmental regulation of transmitter release at the calyx of Held in rat
auditory brainstem. The Journal of physiology 534:861-871.

Kandler K, Friauf E (1993) Pre- and postnatal development of efferent connections of the cochlear nucleus
in the rat. The Journal of comparative neurology 328:161-184.

Kim IH, Jozkowicz A, Piedra PA, Oka K, Chan L (2001) Lifetime correction of genetic deficiency in mice
with a single injection of helper-dependent adenoviral vector. Proc Natl Acad Sci U S A 98:13282-
13287.

Kim MH, Korogod N, Schneggenburger R, Ho WK, Lee SH (2005) Interplay between Na+/Ca2+
exchangers and mitochondria in Ca2+ clearance at the calyx of Held. J Neurosci 25:6057-6065.

Lam SS, Martell JD, Kamer KJ, Deerinck TJ, Ellisman MH, Mootha VK, Ting AY (2015) Directed
evolution of APEX2 for electron microscopy and proximity labeling. Nature methods 12:51-54.

Leao RM, Kushmerick C, Pinaud R, Renden R, Li GL, Taschenberger H, Spirou G, Levinson SR, von
Gersdorff H (2005) Presynaptic Na+ channels: locus, development, and recovery from inactivation
at a high-fidelity synapse. J Neurosci 25:3724-3738.

Lee A, Hirabayashi Y, Kwon SK, Lewis TL, Jr., Polleux F (2018) Emerging roles of mitochondria in
synaptic transmission and neurodegeneration. Curr Opin Physiol 3:82-93.

Lichtman JW, Denk W (2011) The big and the small: challenges of imaging the brain's circuits. Science
334:618-623.

Lidierth M (2009) sigTOOL: A MATLAB-based environment for sharing laboratory-developed software
to analyze biological signals. J Neurosci Methods 178:188-196.

Lorteije JA, Rusu SI, Kushmerick C, Borst JG (2009) Reliability and precision of the mouse calyx of Held
synapse. The Journal of neuroscience : the official journal of the Society for Neuroscience
29:13770-13784.

Lowe DG (2004) Distinctive Image Features from Scale-Invariant Keypoints. International Journal of
Computer Vision 60:91-110.

Lubbert M, Goral RO, Satterfield R, Putzke T, van den Maagdenberg AM, Kamasawa N, Young SM, Jr.
(2017) A novel region in the CaV2.1 alpha1 subunit C-terminus regulates fast synaptic vesicle
fusion and vesicle docking at the mammalian presynaptic active zone. eLife 6.

Martell JD, Deerinck TJ, Lam SS, Ellisman MH, Ting AY (2017) Electron microscopy using the genetically
encoded APEX2 tag in cultured mammalian cells. Nat Protoc 12:1792-1816.

Martell JD, Deerinck TJ, Sancak Y, Poulos TL, Mootha VK, Sosinsky GE, Ellisman MH, Ting AY (2012)
Engineered ascorbate peroxidase as a genetically encoded reporter for electron microscopy. Nature
biotechnology 30:1143-1148.

Misgeld T, Kerschensteiner M, Bareyre FM, Burgess RW, Lichtman JW (2007) Imaging axonal transport
of mitochondria in vivo. Nature methods 4:559-561.

Montesinos MS, Chen Z, Young SM, Jr. (2011) pUNISHER: a high-level expression cassette for use with
recombinant viral vectors for rapid and long term in vivo neuronal expression in the CNS. J
Neurophysiol 106:3230-3244.

Montesinos MS, Satterfield R, Young SM, Jr. (2016) Helper-Dependent Adenoviral Vectors and Their Use
for Neuroscience Applications. Methods Mol Biol 1474:73-90.

Morest DK (1968) The growth of synaptic endings in the mammalian brain: a study of the calyces of the
trapezoid body. Z Anat Entwicklungsgesch 127:201-220.

Neher E (2015) Merits and Limitations of Vesicle Pool Models in View of Heterogeneous Populations of
Synaptic Vesicles. Neuron 87:1131-1142.

Nguyen PV, Marin L, Atwood HL (1997) Synaptic physiology and mitochondrial function in crayfish tonic
and phasic motor neurons. J Neurophysiol 78:281-294.

Obashi K, Okabe S (2013) Regulation of mitochondrial dynamics and distribution by synapse position and
neuronal activity in the axon. The European journal of neuroscience 38:2350-2363.

Perkins GA, Tjong J, Brown JM, Poquiz PH, Scott RT, Kolson DR, Ellisman MH, Spirou GA (2010) The
micro-architecture of mitochondria at active zones: electron tomography reveals novel anchoring
scaffolds and cristae structured for high-rate metabolism. The Journal of neuroscience : the official
journal of the Society for Neuroscience 30:1015-1026.

Rodriguez-Contreras A, de Lange RP, Lucassen PJ, Borst JG (2006) Branching of calyceal afferents during
postnatal development in the rat auditory brainstem. The Journal of comparative neurology
496:214-228.

Rodriguez-Contreras A, van Hoeve JS, Habets RL, Locher H, Borst JG (2008) Dynamic development of
the calyx of Held synapse. Proc Natl Acad Sci U S A 105:5603-5608.

Rossi MJ, Pekkurnaz G (2019) Powerhouse of the mind: mitochondrial plasticity at the synapse. Curr Opin
Neurobiol 57:149-155.

Rouder JN, Speckman PL, Sun D, Morey RD, Iverson G (2009) Bayesian t tests for accepting and rejecting
the null hypothesis. Psychon Bull Rev 16:225-237.

Rowland KC, Irby NK, Spirou GA (2000) Specialized synapse-associated structures within the calyx of
Held. J Neurosci 20:9135-9144.

Satzler K, Sohl LF, Bollmann JH, Borst JG, Frotscher M, Sakmann B, Lubke JH (2002) Three-dimensional
reconstruction of a calyx of Held and its postsynaptic principal neuron in the medial nucleus of the
trapezoid body. The Journal of neuroscience : the official journal of the Society for Neuroscience
22:10567-10579.

Saxton WM, Hollenbeck PJ (2012) The axonal transport of mitochondria. Journal of cell science 125:2095-
2104.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,
Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A
(2012) Fiji: an open-source platform for biological-image analysis. Nature methods 9:676-682.

Stasiak A, Sayles M, Winter IM (2018) Perfidious synaptic transmission in the guinea-pig auditory
brainstem. PloS one 13:e0203712.

Stirling JW, Graff PS (1995) Antigen unmasking for immunoelectron microscopy: labeling is improved by
treating with sodium ethoxide or sodium metaperiodate, then heating on retrieval medium. The
journal of histochemistry and cytochemistry : official journal of the Histochemistry Society 43:115-
123.

Taschenberger H, von Gersdorff H (2000) Fine-tuning an auditory synapse for speed and fidelity:
developmental changes in presynaptic waveform, EPSC kinetics, and synaptic plasticity. The
Journal of neuroscience : the official journal of the Society for Neuroscience 20:9162-9173.

Taschenberger H, Leao RM, Rowland KC, Spirou GA, von Gersdorff H (2002) Optimizing synaptic
architecture and efficiency for high-frequency transmission. Neuron 36:1127-1143.

Traynelis SF (1998) Software-based correction of single compartment series resistance errors. J Neurosci
Methods 86:25-34.

Tritsch NX, Bergles DE (2010) Developmental regulation of spontaneous activity in the Mammalian
cochlea. J Neurosci 30:1539-1550.

Verstreken P, Ly CV, Venken KJ, Koh TW, Zhou Y, Bellen HJ (2005) Synaptic mitochondria are critical
for mobilization of reserve pool vesicles at Drosophila neuromuscular junctions. Neuron 47:365-
378.

Vos M, Lauwers E, Verstreken P (2010) Synaptic mitochondria in synaptic transmission and organization
of vesicle pools in health and disease. Frontiers in synaptic neuroscience 2:139.

Wang LY, Neher E, Taschenberger H (2008) Synaptic vesicles in mature calyx of Held synapses sense
higher nanodomain calcium concentrations during action potential-evoked glutamate release. The
Journal of neuroscience : the official journal of the Society for Neuroscience 28:14450-14458.

Wang T, de Kok L, Willemsen R, Elgersma Y, Borst JG (2015) In vivo synaptic transmission and
morphology in mouse models of Tuberous sclerosis, Fragile X syndrome, Neurofibromatosis type
1, and Costello syndrome. Frontiers in cellular neuroscience 9:234.

Wimmer VC, Horstmann H, Groh A, Kuner T (2006) Donut-like topology of synaptic vesicles with a
central cluster of mitochondria wrapped into membrane protrusions: a novel structure-function
module of the adult calyx of Held. The Journal of neuroscience : the official journal of the Society
for Neuroscience 26:109-116.

Xiao L, Michalski N, Kronander E, Gjoni E, Genoud C, Knott G, Schneggenburger R (2013) BMP signaling
specifies the development of a large and fast CNS synapse. Nat Neurosci 16:856-864.

Xu J, Berret E, Kim JH (2017) Activity-dependent formation and location of voltage-gated sodium channel
clusters at a CNS nerve terminal during postnatal development. J Neurophysiol 117:582-593.

Yang YM, Fedchyshyn MJ, Grande G, Aitoubah J, Tsang CW, Xie H, Ackerley CA, Trimble WS, Wang
LY (2010) Septins regulate developmental switching from microdomain to nanodomain coupling
of Ca(2+) influx to neurotransmitter release at a central synapse. Neuron 67:100-115.

Yu WM, Goodrich LV (2014) Morphological and physiological development of auditory synapses. Hear
Res 311:3-16.

Figure legends

Figure 1. Expression of EGFP-mito-APEX2 at the calyx of Held presynaptic terminal. *A1*, Schematic

of the DNA construct encoding EGFP (green) and mitochondrial matrix-targeted APEX2 (red). *A2*,

Developmental (top) and experimental (bottom) timeline of the calyx around the onset of hearing (P12).

Mito-APEX2 expressing HdAds were injected at P1 and experiments performed at either P7 or P21. *B*,

Expression of EGFP following HdAd injection into the anteroventral cochlear nucleus resulted in green

fluorescence in both the cochlear nucleus (left) and the calyx of Held terminals in the contralateral MNTB

(right). Images were acquired at P21. *C*, *D*, Following DAB reaction, mito-APEX2 labeling of the

contralateral MNTB was visible in embedded P7 (*C*) and P21 (*D*) tissue slices under brightfield

microscope. Dashed line represents the slice midline, dotted lines demarcate the MNTB region. Yellow

dotted line represents the transduced MNTB region contralateral to the injection site. *E*, *F*, High

magnification brightfield images of transduced calyx terminals from *C* and *D*. Mito-APEX2 expressing

calyces where visible by the dark area around the MNTB principal cell soma. *G*, *H*, Transmission

electron micrographs at P7 (*G*) and P21 (*H*) of transduced calyx of Held terminals showing electron

dense mitochondria in juxtaposition to mitochondria in uninfected MNTB principal cells. *I*, High

magnification image of labeled mitochondria in a transfected calyx terminal showing labeling is

restricted to the matrix of the mitochondrion, while the intermembrane space remains unstained. *J1*,

Transduced terminals at P21 (red) contain darker mitochondria (having higher electron density)

compared to mitochondria of the opposing principal cell (aqua). *K1*, Non-transduced terminal

(blue) mitochondria compared to the principal cell. *J2*, *K2*, Intensity line profiles of mitochondrial

intensity in transduced and non-transduced terminals, $\Delta\mu$ = difference in mean pixel intensity

between calyx and PC mitochondria. coH, calyx of Held; PC, principal cell. Scale bars: *B*, 200 µm, 50

µm; *C*, *D*, 300 µm; *E*, *F*, 25 µm; *G*, *H*, 1 µm, *I*, 200 nm. *J*, *K*, 500 nm.

Figure 2. Double labeling of EGFP and mito-APEX2 using a combination of pre-embedding DAB-reaction labeling and post-embedding immunogold labeling. Expression of EGFP and mito-APEX2 has no effect on ultrastructure. *A*, Transduced calyx of Held terminal (red) at P21 with electron-dense mitochondria makes synaptic contact with a single principal cell (aqua). *B*, Comparison of a transduced calyx terminal expressing mito-APEX2 (red) versus a non-transduced control terminal (blue), both contacting separate principal cells (aqua). 12 nm gold particles labeling EGFP are visible within the presynaptic terminal of transduced cells. *C*, Gold particle density is significantly higher in calyx terminals expressing EGFP and mito-APEX2 compared to non-transduced negative terminals and principal cells. *D*, Two docked vesicles (orange) convene at a synaptic release site. Active zone is demarcated in yellow. Vesicles less than 5 nm from the active zone were consider docked. *E*, Active zone length is unchanged between control and mito-APEX2 expressing terminals. A total of 120 AZs from control and mito-APEX2 expressing terminals of three different animals (P21) were analyzed. *F*, Single-section distribution of SVs up to 200 nm from the AZ. Box plot whiskers extend to the minimum/maximum within 1.5 interquartile range, open circles indicate individual observations. Scale bars: *A*, 5 µm; *B*, 500 nm, *D*, 200 nm, 200 nm

Figure 3. Expression of mito-APEX2 does not affect synaptic transmission at the calyx of Held-MNTB synapse. Synaptic transmission at the calyx of Held after expression of mito-APEX2 (red) was investigated at P7 (*A*) and P21 (*B*) and compared to age-matched wildtype control (blue). *A1*, Representative traces of evoked EPSCs during afferent fiber stimulation (50 Hz) at P7 for control and mito-APEX2-transduced calyces (injected at P1). Stimulus artifacts are blanked. Vertical black lines

above the traces indicate time points of fiber stimulation. *A2*, Averaged EPSC amplitudes normalized to

the first EPSC showed comparable short-term plasticity between control and mito-APEX2-transduced

calyces. *A3*, Cumulative EPSC amplitudes to determine the readily releasable pool (RRP) by back-

extrapolating the linear fit of the last 10 data points; the y-intercept is an estimate of the RRP size. *A4-*

A7, No differences between control and mito-APEX2-transduced calyces were detected regarding EPSC

amplitude (*A4*), RRP size (*A5*), release probability (*A6*), or replenishment rate (*A7*). *B,* Same layout as

in *A*. Afferent fiber stimulation was performed at 300 Hz. Following chronic expression (20 days) of

mito-APEX2, the calyx of Held-MNTB synapse showed normal synaptic transmission and no change in

EPSC amplitude (*B4*), RRP (*B5*) release probability (*B6*), or replenishment rate (*B7*). Box plot whiskers

extend to the minimum/maximum within 1.5 interquartile range, open circles indicate individual cells.

1013

Figure 4. Mito-APEX2 labeling visualized by SBF-SEM at P7 (*A*, *B*) and P21 (*C*, *D*, *E1*) allows for

identification of transduced calyx of Held terminals. *A*, *C*, Block face images show the calyx of Held

(red) contacting to an MNTB principal cell (aqua). *B*, *D*, High magnification image of the white squared

area in *A* and *C*, respectively, showing electron dense mitochondria in the calyx of Held terminal. *E1*,

Transduced terminals contain darker mitochondria compared to mitochondria of the opposing

principal cell. *F1*, Non-transduced terminal mitochondria compared to the principal cell

mitochondria. *E2*, *F2*, Intensity line profiles of mitochondrial intensity in transduced and non-

transduced terminals, $\Delta\mu$ = difference in mean pixel intensity between calyx and PC mitochondria

respectively. coH, calyx of Held; PC, principal cell. Scale bars: *A*, *C*, 5 µm; *B*, *D*, 1 µm; *E1*, *F1*,

500 nm

1024

Figure 5. Large volume reconstructions of mito-APEX2-positive calyx of Held terminals using

SBF-SEM at P7 (*A*, *D*) and P21 (*B*, *E*). *A*, Volumetric positions of six transduced calyx terminals

and eight minor inputs contacting six different MNTB principal cells (not shown), reconstructed

from a down-sampled volume of $78 \times 78 \times 52$ µm at P7. *B*, Volumetric positions of six transduced

terminals with eight minor inputs contacting six principal cells (not shown) reconstructed from a down-

sampled volume of $78 \times 78 \times 79$ µm at P21. *C*, Volumes of calyx terminal (left) and minor terminal

(right) at P7 and P21. *D*, Individual renderings of mito-APEX2-positive calyx terminals together

with 1-3 minor labeled terminals contacting each principal cell (not shown). Several filopodia

extend from the cup-like calyx terminals (examples are pointed with white arrowheads). *E*,

Individual renderings of mito-APEX2 positive calyx terminal inputs contacting principal cells (not

shown). Thick digitated terminals at P21 had few filopodial extensions, but 3 out of 6 principal

cells had minor labeled inputs (migratory calycigenic axons; second, fourth, and fifth from left).

Box plot whiskers extend to the minimum/maximum within 1.5 interquartile range, open circles

indicate individual terminals. Scale bars: *A*, *B*, 20 µm; *D*, *E*, 10 µm.

1039

Figure 6. Volumetric reconstructions of mitochondria from the same whole terminals and axons

shown in Figure 5 using SBF-SEM at P7 and P21. *A*, *B*, Volumetric positions of mito-APEX2-

labeled mitochondria of the six calyx of Held terminals at P7 (*A*) and P21 (*B*). *C*, *D*, individual

renderings of mito-APEX2-labeled mitochondria with the membrane surface view of the whole

terminals in transparent gray from one terminal at P7 (*C*) and P21 (*D*). *E*, Mitochondrial volume

scales linearly with the calyx terminal volume at both developmental stages. Mitochondrial volume

ratio is not different between axons and terminals at P7, but volume ratio does increase in axons

and terminals across developmental stages. Ratios are also higher in P21 terminals compared to

the P21 axons. *F.* Mitochondrial volume scales linearly with terminal volume in minor terminals

at both developmental stages. Mitochondria volume ratios are slightly higher in P21 minor

terminals compared to P7. Box plot whiskers extend to the minimum/maximum within 1.5

interquartile range, open circles indicate individual axons or terminals. Scale bars: *A*, *B*, 20 µm; *C, D*

10 µm

1053

Figure 7. Full resolution reconstructions of subcompartments and mitochondria of calyx of Held

terminals at P7 and P21 using SBF-SEM. *A-E*, Reconstruction of a portion of P7 calyx of Held

axon (*A*) and terminal (cup) (*B*); and example reconstructions of a portion of P21 axon (*C*), stalk

(*D*), and swellings (*E*). Left images show the membrane surface views of the subcompartments

and right images show the mitochondria inside. Black arrowheads (*E*) point to several individual

swellings of different sizes. White arrowhead indicates a small swelling not containing

mitochondria. *F*, Mitochondrial diameter comparison among subcompartments at P7 and P21.

Scale bars: *A-E*, 5 µm.

1062

Figure 8. Mitochondria and membrane surface details of a segment of mito-APEX2-positive calyx of

Held terminal reconstructed at full resolution from an SBF-SEM dataset at P7. *A*, Part of a block face

image of the terminal (blue) showing the organization of electron dense mitochondria. The principal cell-

contacting face of the calyx terminal revealed a complex membrane topology. Protrusions of the principal

cell invade the calyx, visualized as indentations in the calyx membrane. White arrows highlight

excrescences of the principal cell (also shown in *C*). *B*, Top down view of the reconstructed terminal

(blue) with mitochondria (green). Mitochondria are generally excluded from filopodial extensions of the

calyx (black arrows in **B** and **D**). **C**, **D**, Face-on view of terminal surface (**C**) and mitochondria (green)

within the terminal (**D**), which form a complex network. Scale bars: **A**-**D** 5μm.

1072

Figure 9. Mito-APEX2 labeling visualized by ssSEM at P7 (**A**, **B**) and P21 (**C, D, E1**) allows for

identification of transduced calyx of Held terminals. **A**, **C**, Ultrathin sections show calyx of Held input

(red) to an MNTB principal cell (aqua) at P7 and P21, respectively. **B, D**, Zoomed view of the terminals

(top) depicted in **A** or **C**, respectively, showing electron dense mitochondria and an active zone (top,

thick arrowheads). Puncta adherentia (PA; bottom, thin arrowheads) are also located at the calyx-

principal cell interface. High signal-to-noise ratio by ssSEM imaging allows for discrimination of

individual SVs as well as pre- and post-synaptic densities. **E1**, Transduced terminals contain darker

mitochondria compared to mitochondria of the opposing principal cell. **F1**, Non-transduced

terminal mitochondria compared to the principal cell mitochondria. **E2**, **F2**, Intensity line profiles

of mitochondrial intensity in transduced and non-transduced terminals, Δμ = difference in mean

pixel intensity between calyx and PC mitochondria respectively. coH, calyx of Held; PC, principal cell;

Scale bars: **A**, **C**, 5 μm; **B**, **D**, **E**, **F**, 500 nm.

1085

Figure 10. Identification of transduced terminals by ssSEM for 3D synaptic analysis. **A**, **B**, Partial

3D reconstructions from 100 serial sections of the calyx of Held terminals at P7 (**A**) and P21 (**B**)

respectively, showing active zones (pink) and puncta adherentia (yellow). **C**, **D**, Ball models of an

example active zone (pink) at the calyx of Held-MNTB synapse. SVs within 200 nm of the presynaptic

membrane are modeled in 3D as 46 nm spheres (orange). Distances to the AZ from 0 to 200 nm are

represented as colored lines. **E**, AZ surface area at P7 and P21. **F**, Total number of vesicles within 200 nm

of the AZ. **G**, Vesicle density as a function of AZ area, binned from 0 to 200 nm from the AZ. **H**, Fraction

of SVs within 5 nm bins up to 200 nm from the AZ. Box plot whiskers extend to the minimum/maximum

within 1.5 interquartile range, open circles indicate individual observations. coH, calyx of Held; PC,

principal cell; Scale bars: *A*, *B*, 5 μm; *C D*, colorimetric scale bar 0-200 nm.

Movie legends

Movie 1. Whole reconstructions of calyx of Held terminals contacting six MNTB principal cells at P7 in a 78 by 78 by 52 µm volume using SBF-SEM. Related to Figure 5A.

Movie 2. Whole reconstructions of calyx of Held terminals contacting six MNTB principal cells at P21 in a 78 by 78 by 79 µm volume using SBF-SEM. Related to Figure 5B.

Movie 3. Mitochondria reconstructions of calyx of Held terminals contacting six MNTB principal cells at P7 in the same volume as Movie 3. Related to Figure 6A.

Movie 4. Mitochondria reconstructions of calyx of Held terminals contacting six MNTB principal cells at P21 in the same volume as Movie 4. Related to Figure 6B.

Movie 5. Semi-automated partial reconstruction of a calyx of Held terminal at P7 in a 23 by 22 by 15 µm volume using SBF-SEM. Mitochondria (orange) were also semi-automatically segmented from the volume. Related to Figure 8.

Movie 6. Partial reconstruction of a labeled calyx of Held terminal at P7 in a 26 by 20 by 5 µm volume using ssSEM. Active zones (pink) and puncta adherentia (yellow) are shown on the calyx face in contact with the MNTB principal cell. Related to Figure 10A.

Movie 7. Partial reconstruction of a labeled calyx of Held terminal at P21 in a 27 by 21 by 5 µm volume using ssSEM. Active zones (pink) and puncta adherentia (yellow) are shown on the calyx face in contact with the MNTB principal cell. Related to Figure 10B.

Figure 1

A1
SV40 polyA | pCI-neo intron | EGFP | hsyn promoter | HdAd | hsyn promoter | mito matrix | V5 | APEX2 | mvm intron | BGH polyA

Ψ

8.1 Kb

18.2 Kb

500 bp

B Cochlear nucleus

MNTB

D / M

D / L

A2
immature | hearing onset | adult-like

P1 | P7 | P12 | P21

HdAd injection | experiments | experiments

C P7

E

G PC coH

I

D P21

F

H PC coH PC

J1 P21

J2

Pixel intensity (8 bit)

Calyx | PC

Δμ = 64.88

Distance (nm) | 1778

K1 P21

K2

Pixel intensity (8 bit)

Calyx | PC

Δμ = 8.60

Distance (nm) | 1544

Figure 2

Figure 3

P7 (50 Hz stimulation)

Figure 4

Figure 5

Figure 6

Figure 7

A   P7 axon

B   P7 cup

C   P21 axon

D   P21 stalk

E   P21 swellings

F

Figure 8

Figure 9

Figure 10