METABOLIC RECOVERY AND COMPENSATORY SHELL GROWTH OF JUVENILE PACIFIC GEODUCK *PANOPEA GENEROSA* FOLLOWING SHORT-TERM EXPOSURE TO ACIDIFIED SEAWATER

Samuel J. Gurr[1*], Brent Vadopalas[2], Steven B. Roberts[3], Hollie M. Putnam[1]

[1] University of Rhode Island, College of the Environment and Life Sciences, 120 Flagg Rd, Kingston, RI 02881 USA
[2] University of Washington, Washington Sea Grant, 3716 Brooklyn Ave NE, Seattle, WA 98105 USA
[3] University of Washington, School of Aquatic and Fishery Sciences, 1122 NE Boat St, Seattle, WA 98105 USA

*Corresponding author: Fax: Phone:1-401-874-9510 Email: samuel_gurr@uri.edu

Lay summary: (50-word summary of the paper)
Commercial shellfish hatcheries provide essential food security, but often production can be hampered by sensitivity of shellfish at early life stages. Repeated short-term exposures can increase tolerance and performance of the geoduck clam with implications for sustainable aquaculture.

Author contributions:
SJG, BV, SBR, and HMP designed the experiments, SJG conducted the experiments, SJG, BV, SBR, and HMP drafted, revised, read and approved the final version of the manuscript.

Abstract

While acute stressors can be detrimental, environmental stress conditioning can improve performance. To test the hypothesis that physiological status is altered by stress conditioning, we subjected juvenile Pacific geoduck, *Panopea generosa,* to repeated exposures of elevated pCO_2 in a commercial hatchery setting followed by a period in ambient common garden. Respiration rate and shell length were measured for juvenile geoduck periodically throughout short-term repeated reciprocal exposure periods in ambient (~550 µatm) or elevated (~2400 µatm) pCO_2 treatments and in common, ambient conditions, five months after exposure. Short-term exposure periods comprised an initial 10-day exposure followed by 14 days in ambient before a secondary 6-day reciprocal exposure. The initial exposure to elevated pCO_2 significantly reduced respiration rate by 25% relative to ambient conditions, but no effect on shell growth was detected. Following 14 days in common garden, ambient conditions, reciprocal exposure to elevated or ambient pCO_2 did not alter juvenile respiration rates, indicating ability for metabolic recovery under subsequent conditions. Shell growth was negatively affected during the reciprocal treatment in both exposure histories, however clams exposed to the initial elevated pCO_2 showed compensatory growth with 5.8% greater shell length (on average between the two secondary exposures) after five months in ambient conditions. Additionally, clams exposed to the secondary elevated pCO_2 showed 52.4% increase in respiration rate after five months in ambient conditions. Early exposure to low pH appears to trigger carry-over effects suggesting bioenergetic re-allocation facilitates growth compensation. Life stage-specific exposures to stress can determine when it may be especially detrimental, or advantageous, to apply stress conditioning for commercial production of this long-lived burrowing clam.

1. Introduction

Sustainable food production minimizes overexploitation of wild populations and degradation of ecological health (Campbell *et al.*, 1998; Shumway *et al.*, 2003; Orensanz *et al.*, 2004; Zhang and Hand, 2006). Shellfish aquaculture has expanded worldwide in recent decades to satisfy international trade (FAO 2018). However, early larval and juvenile rearing pose a production bottleneck. For example, early life histories are highly sensitive to biotic (e.g. harmful algae, pathogens; Prado *et al.*, 2005; Rojas *et al.*, 2015) and abiotic stressors (e.g. pH, salinity, thermal, and hypoxic stress; Baker and Mann 1992; Przeslawski et al. 2015; Kroeker *et al.*, 2010; Gimenez *et al.*, 2018). These stressors are known to intensify in coastal marine systems (Cloern, 2001; Diaz and Rosenberg, 2001; Cai *et al.*, 2011; Wallace *et al.*, 2014) causing mass mortality for early-stage bivalves in wild or hatchery settings (Elston *et al.*, 2008; Barton *et al.*, 2015). Local and global anthropogenic stressors such as CO_2-induced changes in pH and carbonate mineral saturation states can reduce performance and normal shell development (White *et al.*, 2013; Waldbusser *et al.*, 2015; Kapsenberg *et al.*, 2018).

Ocean acidification, or the decrease of oceanic pH due to elevated atmospheric partial pressures (µatm pCO_2), poses a threat to aquaculture (Barton *et al.*, 2012; Froehlich *et al.*, 2018; Mangi *et al.*, 2018). Elevated pCO_2 and aragonite undersaturation ($\Omega_{aragonite} < 1$) generally have detrimental consequences for aerobic performance (Pörtner *et al.*, 2004; Portner and Farrell, 2008) and shell biomineralization in marine calcifiers (Shirayama, 2005; Talmage and Gobler, 2010; Waldbusser *et al.*, 2010, 2015; Gazeau *et al.*, 2013). Responses to acidification can be species (Ries *et al.*, 2009) and population specific (Lemasson *et al.*, 2018), but it is widely established to be impactful during early life stages for bivalves (Dupont and Thorndyke, 2009; Gazeau *et al.*, 2010; Kroeker *et al.*, 2010; Gimenez *et al.*, 2018). Experimental research is commonly focused on species with short generational times, (Parker *et al.*, 2011, 2015; Lohbeck *et al.*, 2012) limiting

evidence for effects of acidification on long-lived mollusks important for food and economic

security (Melzner *et al.*, 2009).

The Pacific geoduck *Panopea generosa* is a large and long-lived infaunal clam of cultural

and ecological importance (Dethier, 2006) with an increasing presence in sustainable shellfish

industry (Cubillo *et al.*, 2018). Geoduck production in Washington (USA) provides ~90% of

global supply (Shamshak and King, 2015) and alone constitutes 27% of the overall shellfish

revenue in the state valued at >$24 million yr^{-1} and >$14 $pound^{-1}$ as of 2015 (Washington Sea

Grant, 2015). Geoduck are known to live in dynamic CO_2-enriched low pH waters such as Hood

Canal in Puget Sound, WA where conditions in summer can reach $\Omega_{aragonite}$ 0.4 and pH 7.4 (Feely

et al., 2010). Although *P. generosa* may be adapted and able to acclimatize to local stressors

(Putnam *et al.*, 2017; Spencer *et al.*, 2018), acidification has caused massive losses of larval

bivalves in hatcheries (Barton *et al.*, 2015), identifying a critical need for assessment of

physiological stress tolerance during early life stages.

Evidence of acclimatory mechanisms in response to acidification (Goncalves *et al.*, 2018)

and enhanced performance within and across generations (Parker *et al.*, 2011, 2015; Putnam and

Gates, 2015; Ross *et al.*, 2016; Thomsen *et al.*, 2017; Zhao *et al.*, 2017) support conditioning as a

viable strategy to mitigate the negative effects of stress exposure and enhance organismal

performance under high pCO_2 (Parker *et al.*, 2011; Dupont *et al.*, 2012; Suckling *et al.*, 2015; Foo

and Byrne, 2016). Hormesis is a biphasic low-dose-stimulatory response, as identified in

toxicological studies (Calabrese, 2008), and suggests beneficial carry-over effects of moderate

stress exposure (Calabrese *et al.*, 2007; Costantini *et al.*, 2010; Costantini, 2014; Putnam *et al.*,

2018). Conditioning-hormesis can explain patterns of intra- and transgenerational plasticity for

organisms under environmental change (Calabrese and Mattson, 2011; Costantini *et al.*, 2012;

López-Martínez and Hahn, 2012; Putnam *et al.*, 2018; Visser *et al.*, 2018), but is understudied for

stress resilience in bivalves likely due to generally negative physiological implications of

acidification (Gazeau *et al.*, 2013). In one example of early-life stage conditioning in bivalves,

Putnam et al. (2017) found *P. generosa* exhibit compensatory shell growth after an acute exposure

under elevated pCO_2. This finding suggests acute exposures may present a strategy for stress-

hardening and enhancement of sustainable geoduck production. We therefore tested the hypothesis

that repeated stress exposure under elevated pCO_2 can enhance intragenerational performance for

Pacific geoduck. To this end, we measured the respiration rate and shell growth of juvenile

geoduck in a commercial hatchery under repeated acute periods (~6-10 days) of elevated pCO_2

and aragonite undersaturation, and the longer term (~5 months) carry-over effects.

2. Methods

2.1. Exposure of juveniles

Juvenile geoduck (n = 640; mean ± SEM initial size, 5.08 ± 0.66 mm shell length [measured

parallel to hinge]) were reared in trays (Heath/Tecna water tray) with rinsed sediment for ~16

weeks (pediveliger to juvenile stage) by Jamestown Point Whitney Shellfish Hatchery before

allocated into eight trays for the experiment (Fig. 1; n = 80 clams per tray). During typical hatchery

practice, geoduck are reared from 'settlers' (pediveliger stage; 30 days old) to 'seed' (juvenile

stage; 4-6 months old) in either downwellers or stacked trays; juveniles are then planted *in-situ* to

grow for several years until market size. Following aquaculture practice, trays were filled with 5

mm depth of rinsed sand (35-45 µm grain size) that allowed juvenile geoduck to burrow and

siphons could clearly be seen extended above the sediment throughout the experiments. To enable

measurements of metabolic activity and shell growth, 30 geoduck were placed in an open circular

dish (6.5 cm diameter and 3 cm height) with equal mesh size and sand depth submerged in each

tray, the remaining 50 geoduck in each tray burrowed in the surrounding sediment. Seawater at

the Jamestown Point Whitney Shellfish Hatchery (Brinnon, WA, USA) was pumped from offshore

(100 m) in Quilcene Bay (WA, USA), bag-filtered (5 µm), and UV sterilized before fed to 250-L

conical tanks at rate of 1 L min^{-1}. Four conical tanks were used as replicates for two treatments:

elevated pCO_2 level of ~2300-2500 µatm and ~7.3 pH (total scale); and ambient hatchery

conditions of ~500-600 µatm and ~7.8-7.9 pH (total scale). The elevated pCO_2 level was set with

a pH-stat system (Neptune Apex Controller System; Putnam *et al.*, 2016) and gas solenoid valves

for a target pH of 7.2 (NBS scale) and pH and temperature (°C) were measured every 10 seconds

in conicals (Neptune Systems; accuracy: ± 0.01 pH units and ± 0.1°C, resolution: ± 0.1 pH units

and ± 0.1°C). These treatments were delivered to replicate exposure trays, which were gravity fed

seawater from conicals (Fig. 1; n = 4 per treatment). The experiment began with an initial exposure

period of 10 days under elevated pCO_2 (2345 µatm) and ambient treatments (608 µatm; Table 1).

Preliminary exposure was followed by 14 days in ambient common garden (557 ± 17 µatm; $pH_{t.s.}$

7.9 ± 0.01; $\Omega_{aragonite}$ 1.46 ± 0.04, mean ± SEM) before secondary exposure for 6 days to reciprocal

treatments of elevated pCO_2 (2552 µatm) and ambient treatments (506 µatm; Table 2). For the

secondary exposure period, one tray was crossed to the opposite treatment to address both repeated

and reciprocal exposure (n = 2 trays per initial×secondary pCO_2 treatment; Fig. 1). Following this

the juveniles were exposed to ambient conditions for 157 days within the replicate trays.

Figure 1. Schematic of the repeated exposure experimental design for two exposure trials, initial (10-day) and secondary (6-day), in ambient and elevated pCO_2 treatments. Timeline displays respiration and growth measurements as solid white circles.

Juvenile geoduck were fed semi-continuously with a mixed algae diet (30% *Isocrysis galbana*, 30% *Pavlova lutheri*, and 40% *Tetraselmis suecica*) throughout the 30-d experiment with a programmable dosing pump (Jebao DP-4 auto dosing pump). Large algae batch cultures were counted daily via bright-field image-based analysis (Nexcelom T4 Cellometer; Gurr *et al.*, 2018) to calculate a daily ration of 5×10^7 live algae cells d^{-1} individual^{-1}. Diet was calculated with an equation in Utting & Spencer (1991) catered for 5-mm clams:

$V = (S \times 0.4) \div (7 \times W \times C)$; this equation accounts for a feed ration of 0.4 mg dried algae mg live animal weight^{-1} week^{-1}, the live animal weight (mg) of spat (S; estimated from regression of shell length and weight of Manilla clams in Utting and Spencer 1991), weight (mg) of one million algal cells (W), and cell concentration of the culture (cells µl^{-1}) to calculate the total volume (V) of each

species in a mixed-algae diet. Tray flow rates (mean flow rate, approx. 480 ± 9 ml^{-1} min^{-1}) and

food delivery were measured and adjusted daily.

All geoduck survived the exposure periods. Half of the remaining juveniles burrowed in

each tray were maintained at the hatchery, positioned in the same replicate trays and stacked for

continuous and high flow of ambient seawater (~8-10 L minute^{-1}). Stacked trays, commonly used

for incubation of finfish, present a promising innovation for geoduck aquaculture; the experiment

stack occurred alongside prototype stacked growing trays stocked by Jamestown Point Whitney

Shellfish. The juveniles were fed cultured algae *ad libitum* daily for 157 days before shell length

and respiration rates were measured.

2.2. Respirometry and shell length measurements

Juvenile geoduck were measured on days 2, 5, 8, and 10 of initial exposure, days 0, 2, 4,

and 6 (cumulatively as day 24, 26, 28, and 30, respectively) of secondary exposure, and 157 days

after the exposure period (cumulatively as day 187) to assess rates of oxygen consumption

normalized to shell length. Calibrated optical sensor vials (PreSens, SensorVial SV-PSt5-4ml)

were used to measure oxygen consumption in 4 ml vials on a 24-well plate sensor system (Presens

SDR SensorDish). Juveniles in each treatment dish were divided into three sensor vials (10

individuals vial^{-1} for exposure periods; 1 individual vial^{-1} at 157-d post-exposure), each filled with

0.2 µm-filtered seawater from corresponding trays. Three blank vials per tray, filled only with 0.2

µm-filtered seawater, were used to account for potential microbial oxygen consumption.

Respiratory runs occurred within an incubator at 15°C, with the vials and sensor placed on a rotator

for mixing. Each set of measurements lasted ~30 minutes and trials ceased when oxygen

concentration declined ~70-80% saturation to avoid hypoxic stress and isolate the effect of pCO$_2$

treatment on respiration rate. Siphons were observed pre and post-respirometry and were fully

extended (~1-2 times shell length). Geoduck were subsequently photographed and shell length

(parallel to hinge) was measured using Image J with a size standard (1 mm stage micrometer).

Rates of respiration (oxygen consumption) were calculated from repeated local linear

regressions using the R package LoLinR (Olito *et al.*, 2017). An initial criterion of fixed constants

(from the LoLin R package) for weighting method ($L_{\%}$) and observations (alpha = 0.2) was run

individually for each respirometry measurement over the full 30-minute record as a "reference"

dataset. These are considered to be the most robust parameters as suggested by the R package

authors (Olito *et al.*, 2017). Diagnostic plots (from the LoLin R package) were individually

observed and $L_{\%}$ and alpha were altered as necessary to best approximate the peak empirical

distribution of local linear regressions (see https://doi.org/10.5281/zenodo.3588326 for full

details). To determine the optimal set of parameters, respiration data was calculated using three

alpha values and data truncations (alpha = 0.2, 0.4, and 0.6; truncation = 10-20 minutes, 10-25

minutes, and no truncation; weighting method = $L_{\%}$) and each was compared to the initial reference

dataset with two curve fitting steps (local polynomial regressions) to calculate unbiased and

reproducible rates of oxygen consumption similar to the reference (10-day exposure, r^2=0.88; 6-

day exposure, r^2=0.95). Final metabolic rates of juvenile geoduck were corrected for vial volume,

rates of oxygen change in the blank vials, and standardized by mean shell length ($\mu g\ O_2\ hr^{-1}\ mm^{-1}$

).

2.3. Seawater carbonate chemistry

Total alkalinity (TA; $\mu mol\ kg^{-1}$ seawater) water samples were collected from trays once

daily during treatment periods, in combination with measurements of pH by handheld probe

(Mettler Toledo pH probe; resolution: 1 mV, 0.01 pH ; accuracy: ± 1 mV, ± 0.01 pH; Thermo

Scientific Orion Star A series A325), salinity (Orion 013010MD Conductivity Cell; range 1 µS/cm

- 200 mS/cm; accuracy: $\pm$ 0.01 psu), and temperature (Fisherbrand Traceable Platinum Ultra-

Accurate Digital Thermometer; resolution; 0.001°C; accuracy: $\pm$ 0.05 °C). Seawater chemistry

was measured for three consecutive days during the 14 days of ambient common garden between

initial and secondary treatment periods. Quality control for pH data was assessed daily with Tris

standard (Dickson Lab Tris Standard Batch T27) and handheld conductivity probes used for

discrete measurements were calibrated every three days. TA was measured using an open cell

titration (SOP 3b; Dickson *et al.*, 2007) with certified HCl titrant ($\sim$0.1 mol kg^{-1}, $\sim$0.6 mol kg^{-1}

NaCl; Dickson Lab) and TA measurements identified <1% error when compared against certified

reference materials (Dickson Lab CO_2 CRM Batches 137 and 168). Seawater chemistry was

completed following Guide to Best Practices (Dickson *et al.*, 2007); daily measurements were used

to calculate carbonate chemistry, CO_2, pCO_2, HCO_3^{3-}, CO_3, and $\Omega_{aragonite}$, using the SEACARB

package (Gattuso *et al.*, 2018) in R v3.5.1 (R Core Team, 2018).

2.4. Data Analysis

A two-way Analysis of Variance (ANOVA) was used to analyze the effect of time (fixed),

pCO_2 treatment (fixed), and time$\times pCO_2$ interaction for respiration and shell length during initial

exposure. A t-test was used to test the effect of initial pCO_2 treatment on respiration rate and shell

length prior to the secondary exposure (last day of ambient common garden, cumulatively day 24,

day 0). For the secondary exposure period, a three-way ANOVA was used to test the effects of

time (fixed), initial pCO_2 treatment (fixed), secondary pCO_2 treatment (fixed), and their

interactions on respiration rate and shell length. No significant differences in seawater chemistry

were detected between trays of the same treatment (pH, pCO_2, TA, salinity, and temperature;

https://doi.org/10.5281/zenodo.3588326), thus tray effects were assumed negligible. Significant

model effects were followed with pairwise comparisons with a Tukey's *a posteriori* HSD. We

used a two-way ANOVA to analyze the effects of initial (fixed) and secondary (fixed) pCO_2

treatments on respiration and shell length after 157 days in ambient conditions. In all cases, model

residuals were tested for normality assumptions with visual inspection of diagnostic plots (residual

vs. fitted and normal Q-Q; Kozak and Piepho, 2018) and homogeneity of variance was tested with

Levene's test. Model effects using raw data were robust to transformation(s) that resolved

normality assumptions via Shapiro-Wilk test. Statistical tests were completed using R (v3.5.1; R

Core Team, 2018) and all code is available (https://doi.org/10.5281/zenodo.3588326).

3. Results

3.1. Exposure 1

The respiration rate of juvenile clams (4.26 ± 0.85 mm shell length; mean $\pm$ SD) prior to

exposure was $0.29 \pm 0.16\ \mu g\ O_2\ hr^{-1}\ mm^{-1}$ (mean $\pm$ SD). Elevated pCO_2 had a significant effect on

respiration rate over the initial 10-day exposure (pCO_2 treatment, $F_{1,88} = 7.512$; P < 0.01) with a

25% reduction (averaged across all days) in respiration rate in elevated pCO_2 treatment relative to

ambient (Fig. 2A). Juvenile geoduck grew significantly with time under the initial 10-d exposure

(time, $F_{3,949} = 3.392$; P = 0.018) with a 3.6% increase in shell length between days 2 and 10 (Fig.

2B), but there was no effect of pCO_2 treatment on shell length (Table 2). Significant differences in

respiration rate from the initial pCO_2 treatment were still apparent after 14 days in ambient

common garden and before the onset of the secondary exposure (Table 2 and Fig. 3A). In contrast,

there was no significant change in shell length due to initial pCO_2 treatment after 14 days in

ambient common garden (Table 2).

3.2. Exposure 2

There was no interaction between initial and secondary pCO_2 treatments nor between

treatments and time on respiration rate or shell length (Table 2). There was a marginal effect of

time on respiration rate (Table 2; time, $F_{2,60} = 3.137$; P = 0.0506) with a 31% increase in average

respiration rate between days 2 and 6. Initial pCO_2 treatment had a significant effect on shell length,

with on average a ~4% reduction in shell size under high pCO_2 relative to ambient initial exposure

(Fig. 3B; $pCO_{2\_initial}$, $F_{1,709} = 15.821$; P < 0.001). This same trend was present under the secondary

high pCO_2 exposure, (Fig. 3B; $pCO_{2\_secondary}$, $F_{1,709} = 9.917$; P = 0.002) with 3.20% smaller shells

for individuals exposed to elevated pCO_2 treatments. There were pairwise differences in shell size

between animals only exposed to ambient and animals repeatedly exposed to elevated pCO_2 (Fig.

3B; day 6, P = 0.0415; day 6 ambient - day 4 elevated, P = 0.0406).

3.3. Common garden after exposure periods

There was no interaction between initial and secondary pCO_2 treatments on respiration rate

or shell length (Table 2). The initial exposure period had a significant effect on shell length of

juveniles previously exposed to high pCO_2, after 157 days in ambient common garden (Fig. 4A;

$pCO_{2\_initial}$, $F_{1,170} = 5.228$; P = 0.023), where average shell lengths were 5.8% larger in juveniles

exposed to initial elevated pCO_2. Secondary 6-day exposure had a significant effect on respiration

rates after 157 days in ambient common garden (Fig. 4B; $pCO_{2\_seccondary}$, $F_{1,31} = 13.008$; P = 0.001)

with an average of 52.4% greater respiration rates in juveniles secondarily exposed to elevated

pCO_2. Visual examination during screening indicated low mortality (1-4 tray^{-1}) over the ~5-month

grow-out period. Shell lengths of dead animals (as empty shells) were similar to the size of juvenile

geoduck during the 30-d exposure period suggesting low mortality occurred at the start of the

grow-out period possibly due to handling stress.

4. Discussion

Metabolic recovery and compensatory shell growth by juvenile *P. generosa* present a novel

application of hormetic framework for resilience of a mollusc to acidification. To date, within-

generation carry-over effects remain poorly understood for marine molluscs (Ross *et al.*, 2016)

with few examples of either positive and negative responses after stress challenges (Hettinger *et*

al., 2012; Gobler and Talmage, 2013; Putnam *et al.*, 2017). Further study on conditioning-hormesis

in response to pCO$_2$ stress must address cellular-level energy allocation, in addition to whole

organism physiology, to account for essential functions with more holistic implications for stress

resilience (Pan *et al.* 2015).

4.1. Metabolic depression and compensatory response

Metabolic depression, such that was found under initial exposure of geoduck to elevated

pCO$_2$, has been suggested as an adaptive mechanism to extend survival (Guppy and Withers,

1999). Stress-induced metabolic depression has been documented for a variety of marine

invertebrates in response to environmental stress. For example, in the New Zealand geoduck,

Panopea zelandica, there was a 2-fold reduction in respiration rate under hypoxia (Le *et al.*, 2016).

Prior work has shown metabolic reductions up to 60-95% of basal performance at rest for marine

molluscs (Guppy and Withers, 1999). Here, depression of oxygen consumption rate by juvenile

geoduck to ~25% in comparison with rates under ambient conditions suggests *P. generosa* are

relatively tolerant to short-term acidification and may have adaptive physiology to cope with

environmental acidification and high pCO$_2$. Responsiveness to acidification is critical for pH-

tolerant taxa to maintain buffering capacity and cope with acidosis (high intracellular pCO$_2$;

(Melzner *et al.*, 2009). However, pH-induced metabolic depression to a similar degree found in

this study has caused a permanent decrease in extracellular pH and increase in protein degradation

and ammonia excretion in the Mediterranean mussel (*Mytilus galloprovincialis*) (Michaelidis *et*

al., 2005). Conversely, metabolic elevation is relatively common for early-life stage bivalves

exposed to low pH and $\Omega_{aragonite}$ undersaturation and typically coincides with consequences for

performance and survival (Michaelidis *et al.*, 2005; Beniash *et al.*, 2010; Thomsen and Melzner,

2010; Fernández-Reiriz *et al.*, 2011; Waldbusser *et al.*, 2015; Lemasson *et al.*, 2018). Whether

depressed or elevated, stress-induced metabolic alterations are known to contribute to biochemical

outcomes such as intracellular hypercapnia and hemolymph acidosis (Pörtner *et al.*, 2004; Spicer

et al., 2011) and increased ammonia excretion and reduced growth for invertebrate fauna

(Michaelidis *et al.*, 2005; Beniash *et al.*, 2010; Lannig *et al.*, 2010; Thomsen and Melzner, 2010;

Gazeau *et al.*, 2013). However, pCO_2 did not impair shell growth during the initial period further

demonstrative of the pH/hypercapnia-tolerance of *P. generosa.*

Juvenile geoduck repeatedly exposed to elevated pCO_2 showed possible stress "memory"

with rebound from metabolic depression under subsequent stress and higher respiration rate and

compensatory shell growth after long-term recovery. Metabolic rebound supports a hormetic-like

response by *P. generosa* (Calabrese *et al.*, 2007; Costantini, 2014) and prompts further

investigation of energy budget, cellular, and -omic measures under repeated reciprocal stress

encounters to improve our understanding of the mechanism underpinning hormesis. Use of

hormesis to conceptualize carry-over effects of mild stress exposure is largely confined to model

insects, plants, and microorganisms (Lee *et al.*, 1987; Calabrese and Blain, 2009; López-Martínez

and Hahn, 2012; Visser *et al.*, 2018). For example, Visser et al. (2018) found the Caribbean fruit

fly, *Anastrepha suspensa*, exposed to oxidative stress early in life enhanced survivorship and

investment in fertility and lipid synthesis under subsequent stress during adulthood. Mechanistic

molecular and biochemical assessments under different and repeated stress intensities (i.e.

magnitude, duration, and frequency) are planned to determine the threshold between low-dose

stimulation and high-dose inhibition from stress-conditioning.

4.2. Age and intensity dependence of shell growth

Metabolic recovery was coupled with reduced shell growth under a repeated stress

encounter (Fig. 3) and compensatory shell growth after approximately five months in ambient

conditions (Fig. 4). This could be explained by several hypotheses such as: carry-over effect from

metabolic depression under initial exposure to elevated pCO_2 (Fig. 2A), differing sensitivity to

stress intensity (Table 1), and/or age dependence for environmental hardening, or the interaction

with increasing temperature through the season (see Supplementary Figure 1.). Bivalves known to

exhibit metabolic suppression under acute and long-term acidification are often attributed with

increased ammonia excretion rates and decreased ingestion and clearance rates as possible

contributors to protein degradation and reduced growth (Michaelidis *et al.*, 2005; Thomsen and

Melzner, 2010; Fernández-Reiriz *et al.*, 2011; Navarro *et al.*, 2013). Therefore, decreased shell

length under secondary exposure may be a latent effect of metabolic depression during initial

exposure. However, shell length was also reduced for clams initially exposed to the elevated

treatment in the second exposure period (Table 2, Fig. 3B), indicating potential age-dependence

of calcification and bioenergetic effects for juvenile *P. generosa.* This reduction however, could

also be explained by the fact the secondary elevated pCO_2 treatment was on average ~0.04 pH

units lower than the initial exposure (Table 1), suggesting possible sensitivity to increased stress

intensity. It is likely that both temporal dynamics and stress thresholds influence intragenerational

carry-over effects and further experimental efforts with repeated reciprocal design are needed.

Respiration rates and shell growth five months post-exposure show a latent enhancement

for animals repeatedly stressed or exposed to a stress event earlier in life, emphasizing the

importance of the severity, duration, and timing of intragenerational stress-conditioning. These

specific findings present a window in their life history where it may be advantageous to condition

Pacific geoduck for enhancement of sustainable aquaculture.

4.3. Commercial and environmental applications of experimental findings

Our findings infer both positive and negative implications for aquaculture. Although

advantageous to elicit carry-over effects exhibited by stress-conditioned animals, results imply

greater feed (ingestion rate) to sustain enhanced aerobic metabolism and compensatory shell

growth; this can heighten labor and financial costs for industry, likely not incentivized by a

marginal 5.8% increase in shell size. However, typical protocols for geoduck aquaculture yield 5-

month-old juvenile clams in the hatchery before grown *in-situ* for 5-7 years. Consequently, latency

of enhanced performance in this study (~9-month-old juveniles), overlaid with the standard

timeline for geoduck industry, does not present additional expenses. Further related tests on stress

conditioning and production of resilient strains (i.e. phenotypes and/or epigenotypes) must account

for distinct life-stages and species-specific attributes in aquaculture practice.

Shellfish farming has adapted in recent years to implement "climate-proofing" technology

to maintain production and combat both coastal and climate-related stressors (e.g. ocean

acidification, sea-level rise, coastal development; Allison et al. 2011). For example, chemical

buffering systems (e.g. mixing sodium bicarbonate) are increasingly common in shellfish industry

to elevate aragonite saturation levels and reduce deleterious effects of ocean acidification;

hatcheries report increases in productivity by 30-50%, offsetting the cost to maintain optimal

carbonate chemistry year-round (Barton et al. 2015). Although buffering systems are advantageous

to yield juvenile 'seed', alleviation of aragonite undersaturation in the short-term may leave

juveniles and adults unprepared to cope with the heterogeneity of environmental chemistry during

long growing periods *in-situ*. As conditions in coastal bays report deteriorating water quality

(Feely et al. 2008; Wallace et al. 2014; Cloern 2001; Melzner et al. 2013), acclimatization and

selective breeding posit alternate and more robust solutions to generate stress-resilience (Barton et

al. 2015). Implementation and tests of effectiveness of stress conditioning remain uncommon for

scientists and aquaculture; our novel findings collected in a hatchery setting provide incentive to

fine-tune stress exposures and build a mechanistic understanding of physiological, cellular, and

molecular responses. Critical questions to test the practical application of stress conditioning are:

(1) what are the effects of repeated stress exposures on energy budget? (2) what life-stages and/or

pCO_2 stress intensity (i.e. magnitude and duration) optimizes establishment of resilient phenotypes

and genotypes during hatchery-rearing? (3) does stress history under elevated pCO_2 affect the

stability and longevity of carry-over effects later in life? Answers to these challenges will result in

effective implementation of conditioning to both reduce pressure on wild stocks and sustain food

security under environmental change.

Although this study was primarily focused on production enhancement in a hatchery

setting, effects on shell growth and metabolism have important applications to natural systems.

Seawater carbonate chemistry targeted for stress treatments was more severe than levels

commonly used in experimental research (Gazeau *et al.*, 2010; Navarro *et al.*, 2013; Diaz *et al.*,

2018), but relevant to summer subsurface conditions within the natural range of *P. generosa* (pH

7.4 and $\Omega_{aragonite}$ 0.4 in Hood Canal, WA; Feely *et al.*, 2010). Thus, survival, metabolic recovery,

and compensatory growth in *P. generosa* in this study demonstrates a resilience to short-term

acidification in the water column. Enhanced growth rates during juvenile development can present

benefits for burrowing behavior (Green *et al.*, 2009; Clements *et al.*, 2016; Meseck *et al.*, 2018)

and survival due to decreased risk of predation and susceptibility to environmental stress

(Przeslawski and Webb, 2009; Johnson and Smee, 2012). Specific to juvenile *P. generosa*, time to

metamorphosis (to dissoconch), pre-burrowing time (time elapsed to anchor into substrate and

obtain upright position), and burrowing depth are directly related to growth and survival (Goodwin

and Pease, 1989; Tapia-Morales *et al.*, 2015). Thus, stress conditioning under CO_2-enrichment and

low pH may enhance survivorship of juvenile geoduck in natural systems. Water column carbonate

chemistry may be critical for sustainable production of infaunal clams, such as *P. generosa*, that

are out-planted for several years *in-situ* on mudflats known to exhibit dynamic abiotic gradients

(Green *et al.*, 1993; Burdige *et al.*, 2008) adjacent to seasonally acidified and undersaturated water

bodies (Feely *et al.*, 2010; Reum *et al.*, 2014).

Conclusion

Data in this present study provides evidence of capacity to cope with short-term

acidification for an understudied infaunal clam of high economic importance. Survival of all

individuals over the 30-d experiment demonstrates the resilience of this species to low pH and

reduced carbonate saturation. Juvenile geoduck exposed to low pH for 10 days recovered from

metabolic depression under subsequent stress exposure and conditioned animals showed a

significant increase in both shell length and metabolic rate compared to controls after five months

under ambient conditions, suggesting stress "memory" and compensatory growth as possible

indicators of enhanced performance from intragenerational stress-conditioning. Our focus on

industry enhancement must expand to test developmental morphology, physiology, and genetic

and non-genetic markers over larval and juvenile stages in a multi-generational experiment to

generate a more holistic assessment of stress hardening and the effects of exposure on cellular

stress response (Costantini *et al.*, 2010; Foo and Byrne, 2016; Eirin-Lopez and Putnam, 2018) for

advancement of sustainable aquaculture (Branch *et al.*, 2013). Advancements in genome

sequencing will facilitate further research to synthesize -omic profiling (i.e global DNA

methylation and differential expression) with physiological responses throughout reproductive and

offspring development under environmental stress (Gavery and Roberts, 2014; Li *et al.*, 2019) to

determine if these mechanisms are transferable among species. Stress conditioning within a generation at critical life stages may yield beneficial responses for food production and provide a baseline for other long-lived burrowing bivalves of ecological and economic importance.

Supplementary Material:

Supplementary Figure 1. Continuous temp and pH data from APEX system

Funding:

This work was funded in part through a grant from the Foundation for Food and Agriculture research; Grant ID: 554012, Development of Environmental Conditioning Practices to Decrease Impacts of Climate Change on Shellfish Aquaculture. The content of this publication is solely the responsibility of the authors and does not necessarily represent the official views of the Foundation for Food and Agriculture Research

Acknowledgements:

We thank the Jamestown S'Klallam Tribe and Kurt Grinnell for providing the animals and facilities for this research. We also thank the management staff and technicians at the Jamestown Point Whitney Shellfish Hatchery, Matt Henderson, Josh Valley, and Clara Duncan, for their assistance, and Maddie Sherman, Emma Strand, and Kaitlyn Mitchell for analytical support.

References:

Allison EH, Badjeck M-C, Meinhold K (2011) The Implications of Global Climate Change for Molluscan Aquaculture. *Shellfish Aquaculture and the Environment*.

Barton A, Hales B, Waldbusser GG, Langdon C, Feely RA (2012) The Pacific oyster, Crassostrea
gigas, shows negative correlation to naturally elevated carbon dioxide levels: Implications for
near-term ocean acidification effects. *Limnol Oceanogr* 57: 698–710.

Barton A, Hatchery WCS, Waldbusser G, Feely R, Weisberg S, Newton J, Hales B, Cudd S, Eudeline
B, Langdon C, *et al.* (2015) Impacts of coastal acidification on the Pacific Northwest shellfish
industry and adaptation strategies implemented in response. *Oceanography* 25: 146–159.

Beniash E, Ivanina A, Lieb NS, Kurochkin I, Sokolova IM (2010) Elevated level of carbon dioxide
affects metabolism and shell formation in oysters Crassostrea virginica (Gmelin). *Mar Ecol Prog*
Ser 419: 95–108.

Branch TA, DeJoseph BM, Ray LJ, Wagner CA (2013) Impacts of ocean acidification on marine
seafood. *Trends Ecol Evol* 28: 178–186.

Burdige DJ, Zimmerman RC, Hu X (2008) Rates of carbonate dissolution in permeable sediments
estimated from pore-water profiles: The role of sea grasses. *Limnol Oceanogr* 53: 549–565.

Cai W-J, Hu X, Huang W-J, Murrell MC, Lehrter JC, Lohrenz SE, Chou W-C, Zhai W, Hollibaugh
JT, Wang Y, *et al.* (2011) Acidification of subsurface coastal waters enhanced by eutrophication.
Nat Geosci 4: 766–770.

Calabrese EJ (2008) Hormesis: why it is important to toxicology and toxicologists. *Environ Toxicol*
Chem 27: 1451–1474.

Calabrese EJ, Bachmann KA, Bailer AJ, Bolger PM, Borak J, Cai L, Cedergreen N, Cherian MG,
Chiueh CC, Clarkson TW, *et al.* (2007) Biological stress response terminology: Integrating the
concepts of adaptive response and preconditioning stress within a hormetic dose-response
framework. *Toxicol Appl Pharmacol* 222: 122–128.

Calabrese EJ, Blain RB (2009) Hormesis and plant biology. *Environmental Pollution*.

Calabrese EJ, Mattson MP (2011) Hormesis provides a generalized quantitative estimate of biological
plasticity. *J Cell Commun Signal* 5: 25–38.

Campbell A, Harbo RM, Hand CM (1998) Harvesting and distribution of Pacific geoduck clams,
Panopea abrupta, in Brtitish Columbia. *Can Spec Publ Fish Aquat Sci/Publ Spec Can Sci Halieut*
Aquat 125: 349–358.

Cao R, Liu Y, Wang Q, Yang D, Liu H, Ran W, Qu Y, Zhao J (2018) Seawater acidification reduced
the resistance of Crassostrea gigas to Vibrio splendidus challenge: an energy metabolism
perspective. *Frontiers in Physiology*.

Clements JC, Woodard KD, Hunt HL (2016) Porewater acidification alters the burrowing behavior

and post-settlement dispersal of juvenile soft-shell clams (Mya arenaria). *J Exp Mar Bio Ecol*

477: 103–111.

Cloern JE (2001) Our evolving conceptual model of the coastal eutrophication problem. *Marine*

Ecology Progress Series.

Cole VJ, Parker LM, O'Connor SJ, O'Connor WA, Scanes E, Byrne M, Ross PM (2016) Effects of

multiple climate change stressors: ocean acidification interacts with warming, hyposalinity, and

low food supply on the larvae of the brooding flat oyster Ostrea angasi. *Mar Biol* 163.

doi:10.1007/s00227-016-2880-4

Costantini D (2014) Does hormesis foster organism resistance to extreme events? *Frontiers in*

Ecology and the Environment.

Costantini D (2019) Hormesis promotes evolutionary change. *Dose Response* 17: 1–4.

Costantini D, Metcalfe NB, Monaghan P (2010) Ecological processes in a hormetic framework. *Ecol*

Lett 13: 1435–1447.

Costantini D, Monaghan P, Metcalfe NB (2012) Early life experience primes resistance to oxidative

stress. *J Exp Biol* 215: 2820–2826.

Cubillo AM, Ferreira JG, Pearce CM, Marshall R, Cheney D, Hudson B (2018) Ecosystem services of

geoduck farming in South Puget Sound, USA: a modeling analysis. *Aquac Int* 26: 1427–1443.

Dethier M (2006) Native Shellfish in Nearshore Ecosystems of Puget Sound. Puget Sound Nearshore

Partnership Report No. 2006-04. Seattle District, U.S. Army Corps of Engineers, Seattle,

Washington.

Diaz RJ, Rosenberg R (2001) Overview of anthropogenically-induced hypoxic effects on marine

benthic fauna. *Coastal and Estuarine Studies*.

Diaz R, Lardies MA, Tapia FJ, Tarifeño E, Vargas CA (2018) Transgenerational effects of pCO_2-

driven ocean acidification on adult mussels Mytilus chilensis modulate physiological response to

multiple stressors in larvae. *Front Physiol* 9. doi:10.3389/fphys.2018.01349

Dickson AG, Sabine CL, Christian JR, eds. (2007) Guide to Best Practices for Ocean CO_2

Measurements. PICES.

Dupont S, Dorey N, Stumpp M, Melzner F, Thorndyke M (2012) Long-term and trans-life-cycle

effects of exposure to ocean acidification in the green sea urchin Strongylocentrotus

droebachiensis. *Mar Biol* 160: 1835–1843.

Dupont S, Thorndyke MC (2009) Impact of CO_2-driven ocean acidification on invertebrates early

life-history – What we know, what we need to know and what we can do. *Biogeosciences*

Discussions.

Eirin-Lopez JM, Putnam HM (2018) Marine environmental epigenetics. *Ann Rev Mar Sci*.

doi:10.1146/annurev-marine-010318-095114

Elston RA, Hasegawa H, Humphrey KL, Polyak IK, Häse CC (2008) Re-emergence of Vibrio

tubiashii in bivalve shellfish aquaculture: severity, environmental drivers, geographic extent and

management. *Dis Aquat Organ* 82: 119–134.

FAO (2018) The State of World Fisheries and Aquaculture 2018 - Meeting the Sustainable

Development Goals.

Feely RA, Alin SR, Newton J, Sabine CL, Warner M, Devol A, Krembs C, Maloy C (2010) The

combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation

in an urbanized estuary. *Estuar Coast Shelf Sci* 88: 442–449.

Fernández-Reiriz MJ, Range P, Álvarez-Salgado XA, Labarta U (2011) Physiological energetics of

juvenile clams Ruditapes decussatus in a high CO_2 coastal ocean. *Mar Ecol Prog Ser* 433: 97–

105.

Foo SA, Byrne M (2016) Acclimatization and adaptive capacity of marine species in a changing

ocean. In: Advances in Marine Biology. pp 69–116.

Froehlich HE, Gentry RR, Halpern BS (2018) Global change in marine aquaculture production

potential under climate change. *Nat Ecol Evol* 2: 1745–1750.

García E, Clemente S, Hernández JC (2018) Effects of natural current pH variability on the sea urchin

Paracentrotus lividus larvae development and settlement. *Mar Environ Res* 139: 11–18.

Gattuso JP, Epitalon JM, Lavine H (2018) Seacarb: Seawater Carbonate Chemistry.

Gavery MR, Roberts SB (2014) A context dependent role for DNA methylation in bivalves. *Brief*

Funct Genomics 13: 217–222.

Gazeau F, Gattuso JP, Dawber C, Pronker AE, Peene F, Peene J, Heip CHR, Middelburg JJ (2010)

Effect of ocean acidification on the early life stages of the blue mussel (Mytilus edulis).

Biogeosci Discuss 7: 2927–2947.

Gazeau F, Parker LM, Comeau S, Gattuso J-P, O'Connor WA, Martin S, Pörtner H-O, Ross PM

(2013) Impacts of ocean acidification on marine shelled molluscs. *Mar Biol* 160: 2207–2245.

Gimenez I, Waldbusser GG, Hales B (2018) Ocean acidification stress index for shellfish (OASIS):

Linking Pacific oyster larval survival and exposure to variable carbonate chemistry regimes.

Elem Sci Anth 6: 51.

Gobler CJ, Talmage SC (2013) Short- and long-term consequences of larval stage exposure to

constantly and ephemerally elevated carbon dioxide for marine bivalve populations.

Biogeosciences.

Goncalves P, Anderson K, Raftos DA, Thompson EL (2018) The capacity of oysters to regulate

energy metabolism-related processes may be key to their resilience against ocean acidification.

Aquac Res 49: 2059–2071.

Goodwin L, Pease B (1989) Species Profiles: Life Histories and Environmental Requirements of

Coastal Fish and Invertebrates (Pacific Northwest): Pacific Geoduck Clam. US Fish and

Wildlife.

Green MA, Aller RC, Aller JY (1993) Carbonate dissolution and temporal abundances of

Foraminifera in Long Island Sound sediments. *Limnol Oceanogr* 38: 331–345.

Green MA, Waldbusser GG, Reilly SL, Emerson K, O'Donnell S (2009) Death by dissolution:

Sediment saturation state as a mortality factor for juvenile bivalves. *Limnol Oceanogr* 54: 1037–

1047.

Gunderson AR, Armstrong EJ, Stillman JH (2016) Multiple stressors in a changing world: the need

for an improved perspective on physiological responses to the dynamic marine environment. *Ann*

Rev Mar Sci 8: 357–378.

Guppy M, Withers P (1999) Metabolic depression in animals: physiological perspectives and

biochemical generalizations. *Biol Rev Camb Philos Soc* 74: 1–40.

Gurr SJ, Rollando C, Chan LL, Vadopalas B, Putnam HM, Roberts SB (2018) Alternative Image-

Based Technique for Phytoplankton Cell Counts in Shellfish Aquaculture (No. 1001481).

Nexcelom.

Hettinger A, Sanford E, Hill TM, Russell AD, Sato KNS, Hoey J, Forsch M, Page HN, Gaylord B

(2012) Persistent carry-over effects of planktonic exposure to ocean acidification in the Olympia

oyster. *Ecology* 93: 2758–2768.

Johnson KD, Smee DL (2012) Size matters for risk assessment and resource allocation in bivalves.

Marine Ecology Progress Series.

Kapsenberg L, Miglioli A, Bitter MC, Tambutté E, Dumollard R, Gattuso JP (2018) Ocean pH

fluctuations affect mussel larvae at key developmental transitions. *Proceedings of the Royal*

Society B: Biological Sciences 285: 20182381.

Kozak M, Piepho HP (2018) What's normal anyway? Residual plots are more telling than

significance tests when checking ANOVA assumptions. *Journal of Agronomy and Crop Science*.

Kroeker KJ, Kordas RL, Crim RN, Singh GG (2010) Meta-analysis reveals negative yet variable

effects of ocean acidification on marine organisms. *Ecol Lett* 13: 1419–1434.

Lannig G, Eilers S, Pörtner HO, Sokolova IM, Bock C (2010) Impact of ocean acidification on energy

metabolism of oyster, Crassostrea gigas—changes in metabolic pathways and thermal response.

Marine Drugs.

Le DV, Alfaro AC, Ragg NLC, Hilton Z, King N (2016) Aerobic scope and oxygen regulation of

New Zealand geoduck (Panopea zelandica) in response to progressive hypoxia. *Aquaculture* 463:

28–36.

Lee RE Jr, Chen CP, Denlinger DL (1987) A rapid cold-hardening process in insects. *Science* 238:

1415–1417.

Lemasson AJ, Hall-Spencer JM, Fletcher S, Provstgaard-Morys S, Knights AM (2018) Indications of

future performance of native and non-native adult oysters under acidification and warming. *Mar*

Environ Res 142: 178–189.

Li Y, Zhang L, Li Y, Li W, Guo Z, Li R, Hu X, Bao Z, Wang S (2019) Dynamics of DNA

methylation and DNMT expression during gametogenesis and early development of scallop

Patinopecten yessoensis. *Mar Biotechnol*. doi:10.1007/s10126-018-09871-w

Lohbeck KT, Riebesell U, Reusch TBH (2012) Adaptive evolution of a key phytoplankton species to

ocean acidification. *Nat Geosci* 5: 346–351.

López-Martínez G, Hahn DA (2012) Short-term anoxic conditioning hormesis boosts antioxidant

defenses, lowers oxidative damage following irradiation and enhances male sexual performance

in the Caribbean fruit fly, Anastrepha suspensa. *J Exp Biol* 215: 2150–2161.

Mackenzie CL, Ormondroyd GA, Curling SF, Ball RJ, Whiteley NM, Malham SK (2014) Ocean

warming, more than acidification, reduces shell strength in a commercial shellfish species during

food limitation. *PLoS ONE*.

Mangi SC, Lee J, Pinnegar JK, Law RJ, Tyllianakis E, Birchenough SNR (2018) The economic

impacts of ocean acidification on shellfish fisheries and aquaculture in the United Kingdom.

Environ Sci Policy 86: 95–105.

Melzner F, Gutowska MA, Langenbuch M, Dupont S, Lucassen M, Thorndyke MC, Bleich M,

Pörtner HO (2009) Physiological basis for high CO_2 tolerance in marine ectothermic animals:

pre-adaptation through lifestyle and ontogeny? *Biogeosci Discuss* 6: 4693–4738.

Meseck SL, Mercaldo-Allen R, Kuropat C, Clark P, Goldberg R (2018) Variability in sediment-water

carbonate chemistry and bivalve abundance after bivalve settlement in Long Island Sound,

Milford, Connecticut. *Mar Pollut Bull* 135: 165–175.

Michaelidis B, Ouzounis C, Paleras A, Pörtner HO (2005) Effects of long-term moderate hypercapnia

on acid-base balance and growth rate in marine mussels Mytilus galloprovincialis. *Mar Ecol*

Prog Ser 293: 109–118.

Navarro JM, Torres R, Acuña K, Duarte C, Manriquez PH, Lardies M, Lagos NA, Vargas C, Aguilera

V (2013) Impact of medium-term exposure to elevated pCO_2 levels on the physiological

energetics of the mussel Mytilus chilensis. *Chemosphere* 90: 1242–1248.

Olito C, White CR, Marshall DJ, Barneche DR (2017) Estimating monotonic rates from biological

data using local linear regression. *J Exp Biol* 220: 759–764.

Orensanz JM (lobo), Hand CM, Parma AM, Valero J, Hilborn R (2004) Precaution in the harvest of

Methuselah's clams the difficulty of getting timely feedback from slow-paced dynamics. *Can J*

Fish Aquat Sci 61: 1355–1372.

Pan T-CF, Applebaum SL, Manahan DT (2015) Experimental ocean acidification alters the allocation

of metabolic energy. *Proc Natl Acad Sci USA* 112: 4696–4701.

Parker LM, O'Connor WA, Byrne M, Coleman RA, Virtue P, Dove M, Gibbs M, Spohr L, Scanes E,

Ross PM (2017) Adult exposure to ocean acidification is maladaptive for larvae of the Sydney

rock oyster in the presence of multiple stressors. *Biol Lett* 13. doi:10.1098/rsbl.2016.0798

Parker LM, O'Connor WA, Raftos DA, Pörtner H-O, Ross PM (2015) Persistence of positive

carryover effects in the oyster, Saccostrea glomerata, following transgenerational exposure to

ocean acidification. *PLoS One* 10: e0132276.

Parker LM, Ross PM, O'Connor WA, Borysko L, Raftos DA, Pörtner H-O (2011) Adult exposure

influences offspring response to ocean acidification in oysters. *Glob Chang Biol* 18: 82–92.

Portner HO, Farrell AP (2008) Physiology and climate change. *Science* 322: 690–692.

Pörtner HO, Langenbuch M, Reipschläger A (2004) Biological impact of elevated ocean CO_2

concentrations: lessons from animal physiology and earth history. *J Oceanogr* 60: 705–718.

Prado S, Romalde JL, Montes J, Barja JL (2005) Pathogenic bacteria isolated from disease outbreaks

in shellfish hatcheries. First description of Vibrio neptunius as an oyster pathogen. *Dis Aquat*

Organ 67: 209–215.

Przeslawski R, Webb AR (2009) Natural variation in larval size and developmental rate of the

northern quahog Mercenaria mercenaria and associated effects on larval and juvenile fitness.

Journal of Shellfish Research.

Putnam HM, Davidson JM, Gates RD (2016) Ocean acidification influences host DNA methylation

and phenotypic plasticity in environmentally susceptible corals. *Evol Appl* 9: 1165–1178.

Putnam HM, Gates RD (2015) Preconditioning in the reef-building coral Pocillopora damicornis and

the potential for trans-generational acclimatization in coral larvae under future climate change

conditions. *J Exp Biol* 218: 2365–2372.

Putnam HM, Ritson-Williams R, Cruz JA, Davidson JM, Gates RD (2018) Nurtured by nature:

considering the role of environmental and parental legacies in coral ecological performance.

bioRxiv. doi:10.1101/317453

Putnam H, Roberts S, Spencer LH (2017) Capacity or adaptation and acclimatization to ocean

acidification in geoduck through epigenetic mechanisms.

R Core Team (2018) A Language and Environment for Statistical Computing.

Reum JCP, Alin SR, Feely RA, Newton J, Warner M, McElhany P (2014) Seasonal carbonate

chemistry covariation with temperature, oxygen, and salinity in a fjord estuary: implications for

the design of ocean acidification experiments. *PLoS One* 9: e89619.

Ries JB, Cohen AL, McCorkle DC (2009) Marine calcifiers exhibit mixed responses to CO_2-induced

ocean acidification. *Geology* 37: 1131–1134.

Rojas R, Miranda CD, Opazo R, Romero J (2015) Characterization and pathogenicity of Vibrio

splendidus strains associated with massive mortalities of commercial hatchery-reared larvae of

scallop Argopecten purpuratus (Lamarck, 1819). *J Invertebr Pathol* 124: 61–69.

Ross PM, Parker L, Byrne M (2016) Transgenerational responses of molluscs and echinoderms to

changing ocean conditions. *ICES Journal of Marine Science: Journal du Conseil*.

Sanders MB, Bean TP, Hutchinson TH, Le Quesne WJF (2013) Juvenile king scallop, Pecten

maximus, is potentially tolerant to low levels of ocean acidification when food is unrestricted.

PLoS One 8: e74118.

Scanes E, Parker LM, O'Connor WA, Stapp LS, Ross PM (2017) Intertidal oysters reach their

physiological limit in a future high-CO_2 world. *J Exp Biol* 220: 765–774.

Shamshak GL, King JR (2015) From cannery to culinary luxury: the evolution of the global geoduck

market. *Mar Policy* 55: 81–89.

Shirayama Y (2005) Effect of increased atmospheric CO_2 on shallow water marine benthos. *J*

Geophys Res 110. doi:10.1029/2004jc002618

Shumway SE, Davis C, Downey R, Karney R, Kraeuter J, Parsons J, Rheault R, Wikfors G (2003)

Shellfish aquaculture–in praise of sustainable economies and environments. *World Aquacult* 34:

8–10.

Spencer LH, Horwith M, Lowe AT, Venkataraman YR, Timmins-Schiffman E, Nunn BL, Roberts SB

(2018) Pacific geoduck (Panopea generosa) resilience to natural pH variation.

Spicer JI, Widdicombe S, Needham HR, Berge JA (2011) Impact of CO_2-acidified seawater on the

extracellular acid–base balance of the northern sea urchin Strongylocentrotus dröebachiensis.

Journal of Experimental Marine Biology and Ecology.

Stevens AM, Gobler CJ (2018) Interactive effects of acidification, hypoxia, and thermal stress on

growth, respiration, and survival of four North Atlantic bivalves. *Mar Ecol Prog Ser* 604: 143–

161.

Suckling CC, Clark MS, Richard J, Morley SA, Thorne MAS, Harper EM, Peck LS (2015) Adult

acclimation to combined temperature and pH stressors significantly enhances reproductive

outcomes compared to short-term exposures. *J Anim Ecol* 84: 773–784.

Talmage SC, Gobler CJ (2010) Effects of past, present, and future ocean carbon dioxide

concentrations on the growth and survival of larval shellfish. *Proc Natl Acad Sci U S A* 107:

17246–17251.

Tapia-Morales S, García-Esquivel Z, Vadopalas B, Davis J (2015) Growth and burrowing rates of

juvenile geoducks Panopea generosa and Panopea globosa under laboratory conditions. *Journal*

of Shellfish Research.

Thomsen J, Casties I, Pansch C, Körtzinger A, Melzner F (2013) Food availability outweighs ocean

acidification effects in juvenile Mytilus edulis: laboratory and field experiments. *Glob Chang*

Biol 19: 1017–1027.

Thomsen J, Melzner F (2010) Moderate seawater acidification does not elicit long-term metabolic

depression in the blue mussel Mytilus edulis. *Mar Biol* 157: 2667–2676.

Thomsen J, Stapp LS, Haynert K, Schade H, Danelli M, Lannig G, Mathias Wegner K, Melzner F

(2017) Naturally acidified habitat selects for ocean acidification–tolerant mussels. *Science*

Advances 3: e1602411.

Utting SD, Millican PF (1997) Techniques for the hatchery conditioning of bivalve broodstocks and

the subsequent effect on egg quality and larval viability. *Aquaculture* 155: 45–54.

Visser B, Williams CM, Hahn DA, Short CA, López-Martínez G (2018) Hormetic benefits of prior

anoxia exposure in buffering anoxia stress in a soil-pupating insect. *The Journal of Experimental*

Biology.

Waldbusser GG, Hales B, Langdon CJ, Haley BA, Schrader P, Brunner EL, Gray MW, Miller CA,

Gimenez I, Hutchinson G (2015) Ocean acidification has multiple modes of action on bivalve

larvae. *PLoS One* 10: e0128376.

Waldbusser GG, Voigt EP, Bergschneider H, Green MA, Newell RIE (2010) Biocalcification in the

eastern oyster (Crassostrea virginica) in relation to long-term trends in Chesapeake Bay pH.

Estuaries Coasts 34: 221–231.

Wallace RB, Baumann H, Grear JS, Aller RC, Gobler CJ (2014) Coastal ocean acidification: The

other eutrophication problem. *Estuarine, Coastal and Shelf Science*.

Washington Sea Grant (2015) Shellfish Aquaculture in Washington State. Final report to the

Washington State Legislature. Washington Sea Grant.

White MM, McCorkle DC, Mullineaux LS, Cohen AL (2013) Early exposure of bay scallops

(Argopecten irradians) to high CO_2 causes a decrease in larval shell growth. *PLoS One* 8:

e61065.

Zhang Z, Hand C (2006) Recruitment patterns and precautionary exploitation rates for geoduck

(Panopea abrupta) populations in British Columbia. *J Shellfish Res* 25: 445–453.

Zhao L, Schöne BR, Mertz-Kraus R, Yang F (2017) Sodium provides unique insights into

transgenerational effects of ocean acidification on bivalve shell formation. *Sci Total Environ*

577: 360–366.

Table 1. pH, salinity, and temperature measured with handheld probes and total alkalinity
measured daily with 60 mL from each heath tray via Gran titration (n = 4 per treatment) during
initial (10-d) and secondary (6-d) exposure trials. Seawater carbonate chemistry (CO_2, pCO_2,
HCO_3^-, CO_3^{2-}, DIC, aragonite saturation state) was calculated with the seacarb R package
(Gattuso *et al.*, 2018).

Treatment	Temperature	Salinity	Flow rate mL min^{-1}	pH, Total Scale	CO_2 µmol kg^{-1}	pCO_2 µatm	HCO_3 µmol kg^{-1}	CO_3 µmol kg^{-1}	DIC µmol kg^{-1}	Total Alkalinity µmol kg^{-1}	Aragonite Saturation state
Initial exposure											
Ambient	14.82 ± 0.12	29 ± 0.03	504 ± 21	7.86 ± 0.007	24 ± 0.5	608 ± 11	1842 ± 4	86 ± 1.4	1952 ± 3	2056 ± 1	1.35 ± 0.02
Low	14.91 ± 0.12	29 ± 0.04	484 ± 17	7.31 ± 0.004	91 ± 0.7	2345 ± 20	1992 ± 1	26 ± 0.20	2108 ± 1	2056 ± 1	0.41 ± 0.003
Ambient common garden											
Ambient	15.01 ± 0.22	29 ± 0.05	449 ± 18	7.89 ± 0.012	21 ± 0.7	561 ± 17	1821 ± 7	93 ± 2.6	1936 ± 5	2051 ± 1	1.45 ± 0.04
Secondary exposure											
Ambient	16.33 ± 0.22	28.67 ± 0.03	494 ± 29	7.93 ± 0.004	19 ± 0.3	506 ± 5	1781 ± 5	102 ± 1.4	1902 ± 4	2033 ± 2	1.60 ± 0.02
Low	16.40 ± 0.22	28.67 ± 0.04	471 ± 18	7.27 ± 0.007	95 ± 1.3	2551 ± 42	1972 ± 3	25 ± 0.3	2091 ± 3	2033 ± 3	0.39 ± 0.005

Table 2. Two-way and three-way ANOVA tests for metabolic rate and shell length during initial and secondary exposures, respectively. A Welch's t-test was used on day zero of secondary exposure to test for differences in mean respiration rate and shell length from initial treatments and a two-way ANOVA tested for treatment effects after 157 days. Significant effects are bolded for $P < 0.05$.

		df	SS	MS	F	P
Initial exposure	*Two-way ANOVA*					
Respiration rate	time	3	0.0323	0.011	0.822	0.485
	$p\,CO_2$	1	0.0983	0.098	7.512	**0.007**
	$p\,CO_2 \times$ time	3	0.0475	0.016	1.210	0.311
Shell length	time	3	4.250	1.415	3.392	**0.018**
	$p\,CO_2$	1	0	0.0005	0.0012	0.973
	$p\,CO_2 \times$ time	3	0.170	0.058	0.138	0.937
Ambient common garden	*Welch Two Sample t-test*	df	t	P		
Respiration rate	$p\,CO_2$	19.833	2.673	**0.015**	-	-
Shell length	$p\,CO_2$	1.146	236.680	0.253	-	-
Secondary exposure	*Three-way ANOVA*					
Respiration rate	time	2	0.068	0.034	3.137	0.051
	$p\,CO_{2\ initial}$	1	0.021	0.021	1.916	0.171
	$p\,CO_{2\ secondary}$	1	0.032	0.032	2.926	0.092
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary}$	1	0.023	0.023	2.080	0.154
	$p\,CO_{2\ initial} \times$ time	2	0.016	0.008	0.724	0.489
	$p\,CO_{2\ secondary} \times$ time	2	0.002	0.001	0.103	0.903
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary} \times$ time	2	0.035	0.017	1.608	0.209
Shell length	time	2	0.190	0.095	0.152	0.859
	$p\,CO_{2\ initial}$	1	9.910	9.910	15.821	**<0.001**
	$p\,CO_{2\ secondary}$	1	6.210	6.212	9.917	**0.002**
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary}$	1	0.060	0.063	0.100	0.752
	$p\,CO_{2\ initial} \times$ time	2	0	0.001	0.002	0.998
	$p\,CO_{2\ secondary} \times$ time	2	0.460	0.231	0.368	0.692
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary} \times$ time	2	0.100	0.048	0.076	0.927
157 days post	*Two-way ANOVA*					
Respiration rate	$p\,CO_{2\ initial}$	1	0.003	0.002	0.011	0.919
	$p\,CO_{2\ secondary}$	1	3.037	3.037	13.008	**0.001**
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary}$	1	0.050	0.050	0.212	0.648
Shell length	$p\,CO_{2\ initial}$	1	10.600	10.597	5.228	**0.023**
	$p\,CO_{2\ secondary}$	1	0.210	0.214	0.105	0.746
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary}$	1	3.510	3.507	1.730	0.190

760 Significant P-values (< 0.05) are bolded.

761
762 **Figure 1.** Schematic of the repeated exposure experimental design for two exposure trials, initial
763 (10-day) and secondary (6-day), in ambient and elevated pCO_2 treatments. Timeline displays
764 respiration and growth measurements as solid white circles.

765
766 **Figure 2.** Respiration rates (A) and shell length (B) of juvenile geoduck under the initial 10-day
767 exposure displayed as box whisker plots with mean (dashed line), median (solid line), 25-50%
768 range (box), and interquartile range (whiskers). Small solid circles represent all data points.

769
Figure 3. Respiration rates (A) and shell length (B) of juvenile geoduck under the secondary 6-day exposure displayed as box whisker plots with mean (dashed line), median (solid line), 25-50% range (box), and interquartile range (whiskers). Small solid circles represent all data points. Letters display significant post hoc effects and an asterisk shows difference via t-test (P < 0.05).

774
775 **Figure 4**. Shell length (A) and metabolic rates (B) of juvenile geoduck after 157 days in ambient
776 common garden conditions post-exposure. Data is displayed as box whisker plots with mean
777 (dashed line), median (solid line), 25-50% range (box), and interquartile range (whiskers). Small
778 solid circles represent all data points.

779
780 **Supplementary Figure 1**. Hourly averages (mean ± SEM) of temperature (°C) and pH (NBS
781 scale) data from the Neptune Apex system. Data were collected every 10 minutes.
782

Treatment	Temperature	Salinity	Flow rate mL min^{-1}	pH, Total Scale	CO_2 µmol kg^{-1}	pCO_2 µatm	HCO_3 µmol kg^{-1}	CO_3 µmol kg^{-1}	DIC µmol kg^{-1}	Total Alkalinity µmol kg^{-1}	Aragonite Saturation state
Initial exposure											
Ambient	14.82 ± 0.12	29 ± 0.03	504 ± 21	7.86 ± 0.007	24 ± 0.5	608 ± 11	1842 ± 4	86 ± 1.4	1952 ± 3	2056 ± 1	1.35 ± 0.02
Low	14.91 ± 0.12	29 ± 0.04	484 ± 17	7.31 ± 0.004	91 ± 0.7	2345 ± 20	1992 ± 1	26 ± 0.20	2108 ± 1	2056 ± 1	0.41 ± 0.003
Ambient common garden											
Ambient	15.01 ± 0.22	29 ± 0.05	449 ± 18	7.89 ± 0.012	21 ± 0.7	561 ± 17	1821 ± 7	93 ± 2.6	1936 ± 5	2051 ± 1	1.45 ± 0.04
Secondary exposure											
Ambient	16.33 ± 0.22	28.67 ± 0.03	494 ± 29	7.93 ± 0.004	19 ± 0.3	506 ± 5	1781 ± 5	102 ± 1.4	1902 ± 4	2033 ± 2	1.60 ± 0.02
Low	16.40 ± 0.22	28.67 ± 0.04	471 ± 18	7.27 ± 0.007	95 ± 1.3	2551 ± 42	1972 ± 3	25 ± 0.3	2091 ± 3	2033 ± 3	0.39 ± 0.005

		df	SS	MS	F	P
Initial exposure	*Two-way ANOVA*					
Respiration rate	time	3	0.0323	0.011	0.822	0.485
	$p\,CO_2$	1	0.0983	0.098	7.512	**0.007**
	$p\,CO_2 \times$ time	3	0.0475	0.016	1.210	0.311
Shell length	time	3	4.250	1.415	3.392	**0.018**
	$p\,CO_2$	1	0	0.0005	0.0012	0.973
	$p\,CO_2 \times$ time	3	0.170	0.058	0.138	0.937
Ambient common garden	*Welch Two Sample t-test*	df	t	P		
Respiration rate	$p\,CO_2$	19.833	2.673	**0.015**	-	-
Shell length	$p\,CO_2$	1.146	236.680	0.253	-	-
Secondary exposure	*Three-way ANOVA*					
Respiration rate	time	2	0.068	0.034	3.137	0.051
	$p\,CO_{2\ initial}$	1	0.021	0.021	1.916	0.171
	$p\,CO_{2\ secondary}$	1	0.032	0.032	2.926	0.092
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary}$	1	0.023	0.023	2.080	0.154
	$p\,CO_{2\ initial} \times$ time	2	0.016	0.008	0.724	0.489
	$p\,CO_{2\ secondary} \times$ time	2	0.002	0.001	0.103	0.903
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary} \times$ time	2	0.035	0.017	1.608	0.209
Shell length	time	2	0.190	0.095	0.152	0.859
	$p\,CO_{2\ initial}$	1	9.910	9.910	15.821	**<0.001**
	$p\,CO_{2\ secondary}$	1	6.210	6.212	9.917	**0.002**
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary}$	1	0.060	0.063	0.100	0.752
	$p\,CO_{2\ initial} \times$ time	2	0	0.001	0.002	0.998
	$p\,CO_{2\ secondary} \times$ time	2	0.460	0.231	0.368	0.692
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary} \times$ time	2	0.100	0.048	0.076	0.927
157 days post	*Two-way ANOVA*					
Respiration rate	$p\,CO_{2\ initial}$	1	0.003	0.002	0.011	0.919
	$p\,CO_{2\ secondary}$	1	3.037	3.037	13.008	**0.001**
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary}$	1	0.050	0.050	0.212	0.648
Shell length	$p\,CO_{2\ initial}$	1	10.600	10.597	5.228	**0.023**
	$p\,CO_{2\ secondary}$	1	0.210	0.214	0.105	0.746
	$p\,CO_{2\ initial} \times p\,CO_{2\ secondary}$	1	3.510	3.507	1.730	0.190

Significant P-values (< 0.05) are bolded.