Effects of unilateral cortical resection of the visual cortex on bilateral human white matter

Anne Margarette S. Maallo[1,2], Erez Freud[3,4], Tina Tong Liu[1,5], Christina Patterson[6],

Marlene Behrmann[1,2]

[1] Department of Psychology, Carnegie Mellon University, Pittsburgh, PA, USA

[2] Center for the Neural Basis of Cognition, Pittsburgh, PA, USA

[3] Department of Psychology, York University, Toronto, ON, Canada

[4] Center for Vision Research, ON, Canada

[5] Laboratory of Brain and Cognition, National Institutes of Health, Bethesda, MD

[6] Department of Pediatrics, University of Pittsburgh, Pittsburgh, PA

Correspondence: Anne Margarette S. Maallo, Ph.D.

Email: maallo@cmu.edu

Department of Psychology

Baker Hall 331G

Carnegie Mellon University

5000 Forbes Ave., Pittsburgh PA, 15213

Highlights

- There is well-circumscribed white matter damage in pediatric epilepsy after surgery
- White matter pathways are normal distal as well as contralesional to the resection
- Contralesional network properties differ after left or right hemisphere resection
- Preserved cortex and white matter may be sufficient for normal perception

Abstract

Children with unilateral resections of ventral occipito-temporal cortex (VOTC) typically do not evince visual perceptual impairments, even when relatively large swathes of VOTC are resected. In search of possible explanations for this behavioral competence, we evaluated white matter microstructure and connectivity in eight pediatric epilepsy patients following unilateral cortical resection and 15 age-matched controls. To uncover both local and broader resection-induced effects, we analyzed tractography data using two complementary approaches. First, the microstructural properties were measured in the inferior longitudinal and the inferior fronto-occipital fasciculi, the major VOTC association tracts. Group differences were only evident in the ipsilesional, and not in the contralesional, hemisphere, and single-subject analyses revealed that these differences were limited to the site of the resection. Second, graph theory was used to characterize the connectivity of the contralesional occipito-temporal regions. There were no changes to the network properties in patients with left VOTC resections nor in patients with resections outside the VOTC, but altered network efficiency was observed in two cases with right VOTC resections. These results suggest that, in many, although perhaps not all, cases of unilateral VOTC resections in childhood, the white matter integrity in the preserved contralesional hemisphere along with residual neural activity might be sufficient for normal visual perception.

45 **Keywords**
46
47 white matter, tractography, graph theory, lobectomy, visual cortex, epilepsy
48

Abbreviations

AD axial diffusivity

DNET dysembryoplastic neuroepithelial tumor

FA fractional anisotropy

fMRI functional magnetic resonance imaging

IFOF inferior fronto-occipital fasciculus

ILF inferior longitudinal fasciculus

LH left hemisphere

MRI magnetic resonance imaging

RD radial diffusivity

RH right hemisphere

ROI region of interest

VOTC ventral occipito-temporal cortex

WM white matter

1 Introduction

Recent studies have revealed that visual perception, for example contour integration and face and word recognition, is typically intact in children who have undergone unilateral visual cortical resection to manage pharmacoresistant epilepsy. One such case, UD, with surgical removal of the ventral occipito-temporal cortex (VOTC) of the right hemisphere (RH), showed normal intermediate and higher-order visual perceptual behavior despite the persistent hemianopia (Liu *et al*., 2018). This behavior was especially striking given that, as revealed through a longitudinal functional magnetic resonance imaging (fMRI) study, the resection impacted some category-selective regions in the RH (e.g. right fusiform face area) that are crucial for perceptual function. The findings of normal visual perception were corroborated in a study with a larger group of children, with the exception of two children with neurological comorbidities (Liu, Freud *et al*., 2019). Also, as was true for UD, too, normal category-selectivity in response to visual stimuli (e.g. images of faces and words) was observed in the remaining cortex (primarily in the VOTC of the structurally intact hemisphere) using fMRI.

A compelling question concerns the neural basis of the intact perceptual abilities in these patients. Given that surgery has been shown to be successful in managing the seizure disorder in the majority of the cases (Helmstaedter *et al*., 2019), it is important to understand the consequences and functional outcomes of this surgical procedure. One candidate mechanism that might impact behavioral outcome is the status of cortical connectivity postoperatively. Several studies have examined functional connectivity in pediatric epilepsy patients using graph-theoretic measures. For example, resting state fMRI studies have revealed that network modularity, path length, and global efficiency were independently associated with epilepsy duration (Paldino *et al*., 2017c). More relevant for establishing cognitive outcome after surgery, these same network measures have been associated with full

scale IQ in children (Paldino *et al.*, 2017b, 2019; Zhang *et al.*, 2018). Of note is that network

properties not only provide biologically and psychologically relevant measures in the

pediatric epileptic brain cross-sectionally, but are also highly replicable longitudinally

(Paldino *et al.*, 2017a).

An alternative way to examine cortical connectivity is to focus on structure and

quantify the integrity of the underlying white matter (WM) obtained from diffusion MRI.

Some diffusion neuroimaging studies have revealed abnormalities even beyond the

epileptogenic zone, including in regions that are typically associated with the default mode

network (DeSalvo *et al.*, 2014) and in the limbic system (Bonilha *et al.*, 2012). In addition to

insights that can be gleaned about the structural connectivity, diffusion MRI can also be used

to assess the microstructural properties of the WM. For example, Pustina et al. (2014)

revealed a decrease in fractional anisotropy (FA), which is a measure of WM integrity, in

numerous tracts of patients with cortical resections, but especially in the ipsilesional tracts

including the uncinate and fornix. However, these structural neuroimaging studies were

conducted mostly with adults where the opportunity for plasticity is less than in younger

patients.

Here, we used diffusion MRI with children who had undergone unilateral cortical

resection. To our knowledge, there has not been a systematic assessment of WM in the

VOTC postoperatively, and this is especially warranted in light of the previous findings of

normal visual perception as described above. Moreover, because a reduction in structural

connectivity is associated with reduced regional grey matter volume (Bonilha *et al.*, 2010), a

precise characterization of WM changes might be especially revelatory.

We adopted two different approaches to characterize the integrity of WM and its

organization: first, we examined the microstructural properties of specific WM tracts that

traverse the VOTC, and second, we evaluated cortical connectivity of the intact VOTC. In the

first approach, we examined the local effects of resection on two VOTC WM tracts: the

inferior longitudinal (ILF) and inferior fronto-occipital fasciculi (IFOF). We measured the

FA, as well as the axial diffusivity (AD), a measure of diffusion parallel to the length of a

tract, and the radial diffusivity (RD), a measure of the diffusion perpendicular to the tract. We

selected these dependent measures as they generally offer clearer insight into the nature of

injury than other measures, say, the mean diffusivity (Aung *et al*., 2013). We expected to

observe hemispheric asymmetries in microstructural WM properties that could result from

either an enhancement in the contralesional tracts, indicative of a compensatory mechanism,

or an asymmetry in which compromised ipsilesional tracts are evident while the

microstructural indices of contralesional tracts do not differ from those of the controls. In the

second approach, to assess any possible WM changes on a broader scale, we used graph

theory to characterize the network organization of the structurally intact contralesional VOTC

and compared the network properties in the patients versus the control children.

The findings from this study have the potential to uncover important alterations in

neural mechanisms following an assault on cortical development during childhood. Major

changes to the WM might point to adaptive compensatory mechanisms, thus enabling normal

perception. Conversely, the absence of significant alterations supports the hypothesis that the

residual existing structure may be sufficient to support perception.

2 Materials and Methods

The procedures were reviewed and approved by the Institutional Review Boards of

Carnegie Mellon University and the University of Pittsburgh. Parents provided informed

consent and minor participants gave assent prior to the scanning session.

2.1 Participants

Participants included six pediatric patients with resections to the VOTC and two

patients with resections outside the VOTC, and 15 age-matched typically-developing controls

(three female, 12 male, mean age 14.5 ± 3.1 years; no age difference across groups,

Wilcoxon rank sum test p>0.67). In total, three patients had resections to the RH and five had

resections to the left hemisphere (LH). In seven of the eight patients, surgery was performed

to manage pharmacoresistant epilepsy and, in the remaining case, surgery was performed for

an emergent evacuation of cerebral hematoma at day one of life (Table 1).

In six of these eight patients, the functional category-selective neural responses have

previously been documented (Liu *et al*., 2018, Liu, Freud *et al*., 2019), but additional data

from patients not included in those studies (SN and DX) are included in the Supplementary

Materials (Figs. A.1 and A.2). Briefly, each participant completed three runs of an fMRI scan

in which neural responses to images (faces, words, house and objects, as well as scrambled

images) were measured, and used to define category-selective regions, which preferentially

responded to one of the visual categories. In two patients with LH resection (SN and TC), the

visual word form area, which is necessary for processing of words and typically localized in

the LH, was instead localized in the RH.

The behavioral profiles for six patients have also been previously documented (Liu *et*

al., 2018, Liu, Freud *et al*., 2019), and the data from the two additional patients (SN and DX)

are presented here. Briefly, all participants completed several tasks designed to assess their

visual perception (see Supplementary Materials Section C for more details). In these tasks,

images were presented centrally, and patients were free to make saccades. Participants

completed two tasks assessing their global perception: one that evaluates contour integration

and one that measures their ability to detect "swirl" in glass patterns. Two further tasks

examined their pattern recognition: one designed to assess object perception and one to

measure face recognition.

We compared patients' behaviors (Table 2), in terms of collinearity thresholds,

accuracy, and response times, to age-matched controls. Although it is difficult to know

exactly what the patients' phenomenological experience was like and self-report is often

imprecise (as is true for other cases of field defects such as hemianopia or quadrantanopia),

the patients' performance was nevertheless comparable to that of matched typically

developing children on a wide range of dependent measures, indicating objective similarities

in perception. An impairment was revealed both for the global perception and object

recognition tasks in patient NN (who had polymicrogyria and low IQ as a complicating

comorbidity), and for the global perception task in patient DX who had non-VOTC resection.

2.2 MRI Protocols

Images were acquired on a Siemens Verio 3T scanner with a 32-channel head coil at

Carnegie Mellon University. A T_1-weighted image with $1\times1\times1mm^3$ resolution (MPRAGE,

TE=1.97ms, TR=2300ms, acquisition time=5m21s) and one set of diffusion-weighted images

with $2\times2\times2mm^3$ resolution (2D EPI, 113 directions, maximum b=4000s/mm^2, TE=121ms,

TR=3981ms, acquisition time=7m46s) were acquired for all participants. The neuroimaging

data presented are being uploaded to KiltHub, Carnegie Mellon's online data repository

(reserved DOI: 10.1184/R1/9856205). However, the DOI will appear empty as the process is

pending institutional approval. We trust that this will be available online after the review

process.

2.3 Microstructural properties of VOTC white matter pathways

This first approach allowed us to determine whether there were local microstructural

alterations to either of the two major WM tracts that traverse the VOTC (Catani *et al*., 2003,

Catani, 2006). The ILF connects the extrastriate occipital lobe and the anterior temporal lobe,

running along the lateral and inferior wall of the lateral ventricles. The IFOF is an indirect

pathway that runs from the occipital to the temporal lobes and then passes through the

uncinate and corona radiata along the lateral border of the caudate nucleus, finally

terminating in the frontal lobes.

The diffusion data were reconstructed in DSI Studio (http://dsi-studio.labsolver.org)

using generalized q-sampling imaging (Yeh *et al*., 2010) with 1.25 diffusion sampling ratio.

We used a deterministic fiber tracking algorithm (Yeh *et al*., 2013) with whole-brain seeding,

0.20 quantitative anisotropy threshold, 0.20mm step size, 20mm minimum length, and

150mm maximum length to generate four sets of tractograms with one million streamlines

each, at angular thresholds of 30°, 40°, 50°, and 60° that were ultimately combined to create

one effective whole-brain tractogram. These thresholds were consistent with values that

allow for robust fiber tract reconstructions as used in the literature (Dennis *et al*., 2015).

Next, we delineated the ILF and IFOF in the patient group, in both the ipsilesional

and contralesional hemispheres, as well as in both hemispheres of the control group. We

filtered the global tractogram described above to extract the tracts using the following

standard procedure, which has been previously used successfully to generate reproducible

results (Wakana *et al*., 2007). The ILF was defined using two regions of interest (ROIs) as

inclusion masks. One ROI included the entire occipital cortex and was located coronally at

the most anterior slice of the parieto-occipital sulcus. The other ROI included the entire

temporal lobe and was located coronally at the most posterior slice in which the temporal

lobe was detached from the frontal lobe. The IFOF was defined similarly: one ROI included

the entire occipital cortex and was located coronally halfway between the anterior and

posterior slices of the parieto-occipital sulcus. The other ROI included the entire frontal lobe

and was located coronally at the anterior edge of the genu of the corpus callosum. For both

tracts, streamlines that projected across the midline were discarded.

We measured the FA, AD, and RD of the tracts also in DSI Studio. These indices

were only measured along the tracts, which were external to the resection areas (i.e. the tracts

were located superior or inferior to, or more medially/laterally with respect to the resection).

A lesion mask was manually drawn based on the anatomical image to enable us to determine

the extent of the resection in the anteroposterior direction and, combined with anteroposterior

profiles of the microstructural indices, we were able to determine the locations of any

abnormalities relative to the resection.

We computed the mean microstructural indices in two regimes. In the first regime, we

computed the mean indices over the entire length of each tract and used these values for

between-group comparisons in patients and controls. In the second regime, we computed the

mean indices over segments of each tract and used these values in single-subject analyses.

Given the heterogeneity in the resection site in the patients and the generally large extent of

the resections, we deemed it sufficient to bisect the tracts into two segments representing the

anterior or the posterior segment of each tract. For instance, if a tract was of length L, we

bisected it into segments, each of length $L/2$ and this was done independently for the left and

right ILF and IFOF in each individual.

2.4 Network properties in the contralesional hemisphere

This second approach allowed us to determine whether there were broader resection-

induced changes in the connectivity of regions in the occipito-temporal cortex. However,

because of the heterogeneity in the location and extent of the resection and the variability in

the preserved ipsilesional tissue, it was difficult to study the ipsilesional hemisphere across

the patient group. Additionally, automated parcellation of the ipsilesional hemisphere was

likely to be unreliable given the structural abnormalities and manual demarcation of regions

may not have been sufficiently systematic across cases. Therefore, we studied only the

contralesional hemisphere and its network properties.

For each participant, we used only a single tractogram to define the network

connectivity, with an angular threshold of $60°$. All other thresholds were the same as in

Section 2.3, except for the number of streamlines. Instead of terminating tractography once a

specific number of streamlines was reached, here, we controlled the seed density (20/voxel)

by placing seeds only in the mask encompassing the preserved hemisphere in each individual

to control for the variability in the participants' head sizes.

Next, we used cortical ROIs based on the parcellation of the anatomical images using

the Destrieux atlas (Destrieux *et al.*, 2000) in FreeSurfer (Dale *et al.*, 1999). The parcellation

output was visually inspected and confirmed to be reliable in the structurally preserved

hemisphere of the patients (see Supplementary Materials Section B for list of the regions

comprising the VOTC that were used as network nodes). We generated a connectivity matrix

in DSI Studio, such that, for any ROI pair, the connectivity value was the mean FA of all

tracts that pass through both ROIs. We binarized the connectivity matrix for each individual

by using a threshold of one standard deviation below the mean FA (excluding FA values of 0

– i.e. ROI pairs that were not connected by any streamline was not used in the computation of

the threshold). Any value higher than the threshold was assigned a value of 1, and every other

value was assigned a value of 0. With this binary connectivity matrix, we computed four

standard network measures: transitivity, modularity, characteristic path length, and global

network efficiency using the Brain Connectivity Toolbox (Rubinov and Sporns, 2010) for

MatLab. We then compared these graph-theoretic measures between patients and controls

using between-group permutation and single-subject analyses. Additionally, we also

explored, on an individual level, the effects of using a different threshold in binarizing the

connectivity matrices.

2.5 Statistical analyses

2.5.1 Group comparisons

In the between-group comparisons, given the small number of VOTC resection patients, we generated 100,000 permutations of data from the two groups (six VOTC resection patients and 15 controls) and used a two-sample t-test, thus creating a null distribution of t-scores. We did not include the non-VOTC resection patients for this analysis. Then, p-values were computed as the ratio of values that were more extreme than the actual patient versus control group t-score relative to the number of permutations (the number of $|t_{permutation}| > t_{actual}$ divided by 100,000).

2.5.2 Single-subject level comparisons

In the single subject analyses, we used established statistical procedures (Crawford and Howell, 1998) to compare each individual patient's data to the data from the controls. With a normative sample size (degree of freedom=14, t_c=2.145), a modified t-test was used in which each patient was treated as a sample of n=1, thereby eliminating the contribution of the single subject to the estimate of within-group variance.

3 Results

3.1 ILF and IFOF in VOTC resection patients

We defined the ILF and IFOF in six children with unilateral resections to their VOTC, as well as in the controls. We were able to reconstruct the ILF and the IFOF in most patients, even in the ipsilesional hemisphere, as long as the streamlines were outside the resection (Fig. 1), with the exception of the ipsilesional ILF and IFOF in KQ and the ipsilesional IFOF in SN.

3.1.1 Between-group differences in microstructural properties of the tracts

To evaluate the differences between the patients and controls, we computed the mean FA, AD, and RD in both of the identified tracts (Fig. 2). Perhaps surprisingly, there were multiple peaks in the null distribution histograms, a result of having extreme values from patients in the pooled data set from both groups that was resampled with no replacement. Relative to this derived null distribution, we found that, for the ipsilesional ILF, FA was lower and both AD and RD were higher in patients than controls (actual between group $|t|>2.5$, all $p<0.01$). Meanwhile, for the ipsilesional IFOF, there were only significant differences between patients and controls on RD (higher in patients than controls, between group $t=2.5$, $p<0.05$). In contrast, for both the contralesional ILF and IFOF, the findings did not differ significantly between the patients and the controls on any of the microstructural indices (between group $|t|<1.69$), all $p>0.11$). Moreover, the results were the same irrespective of whether the patient data were compared to data from either the LH or the RH of the controls, or to the mean of the bilateral tracts in controls.

3.1.2 Microstructural properties of the ILF

Next, we wished to determine whether the observed ipsilesional group differences were present along the entire length of the tracts or restricted to the proximate site of the resection. We plotted the three different microstructural indices along the anteroposterior axis of the ILF (Fig. 3A). In an exemplar control participant (top rows, Fig. 3A), the respective microstructural indices of the ILF were each comparable bilaterally along the entire length of the tract. In patients in whom we recovered bilateral ILF, unlike in controls, there appeared to be differences in the microstructural indices of the two hemispheres: visual inspection suggests that FA was lower, and AD and RD were higher in the ipsilesional than in the

contralesional ILF (consistent with the finding from the between-group analysis above).

Furthermore, these differences appeared to be confined to the extent of the resection of each

patient (preserved hemisphere serves as within-subject comparison, Fig. 3A). Thus, in

patients SN and OT who have anterior temporal lobe resections, the within-subject

hemispheric asymmetry was evident mostly in the anterior segment of the ILF. While in

patients UD, TC, and NN, who had more posterior resections, the within-subject hemispheric

asymmetries were evident in the posterior segment of the ILF.

We quantified the microstructural damage by comparing the mean indices in both the

anterior and posterior segments of the ILF in patients to the corresponding values from

controls (Fig. 3B), and found that only patient OT exhibited normal microstructural indices

across the board. Critically, for the other patients, the significant differences to the

microstructural indices of the ILF was confined to the extent of the resection (Fig. 3B, all

$|t|>2.4$, all $p<0.05$). That is, compared to controls, patients with posterior resections had

compromised microstructural indices only in the posterior segment of the ILF and patient SN

(the only patient who had an anterior resection with bilaterally defined ILF) exhibited

compromised microstructural indices only in anterior microstructural indices of the ILF.

Moreover, both segments of the contralesional ILF exhibited normal microstructural indices

compared to controls.

3.1.3 Microstructural properties of the IFOF

As with the ILF, the anteroposterior profiles of microstructural indices of the IFOF in

patients are also different from those of controls, as well as from the profiles of the same

measures in the ILF in patients. In an exemplar control participant (top rows, Fig. 4A), the

respective microstructural indices of the IFOF were each comparable bilaterally along the

entire length of the tract, similar to the ILF. And in patients in whom we recovered bilateral

IFOF, visual inspection indicates that within-subject hemispheric differences in the IFOF

were present only in patients with more posterior VOTC resections (preserved hemisphere

serves as within-subject comparison, Fig. 4A), wherein the following patterns, similar to

those from the ILF, were seen: FA was lower, AD and RD were higher in the ipsilesional

than in the contralesional IFOF. However, the IFOF could not be defined in two of the three

patients with anterior resection and so this finding ought to be interpreted with caution. Thus,

in patients with posterior resections (UD, TC, and NN), the within-subject hemispheric

asymmetries were evident in the posterior segment of the IFOF.

We also quantified the microstructural damage by comparing the mean indices in both

the anterior and posterior segments of the IFOF in patients to the corresponding values from

controls (Fig. 4B), and found that the significant differences to the microstructural indices of

the IFOF was confined to the extent of the resection (Fig. 4B, all $|t| > 2.4$, all $p < 0.05$). That is,

the patients with posterior resections had compromised microstructural indices only in the

posterior segment of the IFOF. And similar to the ILF, the contralesional IFOF exhibited

normal microstructural indices compared to controls.

3.1.4 Normal microstructural properties of the ILF and IFOF in non-VOTC resection

We next characterized the ILF and IFOF (Fig. 5A) in the two patients with

pharmacoresistant epilepsy who had resections outside their VOTC (EK and DX). Using

identical procedures as in the children with VOTC resections, we found no qualitative within-

subject hemispheric asymmetry in the anteroposterior profiles of the microstructural indices

of either tract in these two patients (Fig. 5B-C) and visual inspection suggests that these plots

resemble the profiles seen in controls (Figs. 3A and 4A, top rows). Furthermore, single-

subject analysis revealed no significant differences in either case compared to the controls

across any of the indices (all |t|<1.23, all p>0.24), be it in the ipsilesional or in the

contralesional tracts.

3.2 Network properties of the contralesional occipito-temporal cortex

Thus far, we have uncovered local changes to specific WM tracts by revealing

microstructural damage restricted to the extent of resection. In a complementary approach,

we wished to examine whether there were also broader resection-induced changes to the

structural connectivity of the visual cortex. We used the same between-group and single-

subject approaches to characterize the network properties in the intact contralesional

hemisphere in the VOTC resection patients as in previous sections.

3.2.1 Between-group comparisons of network properties

First, we combined the data from patients and controls in order to perform

permutation testing on the network properties including transitivity, modularity, path length,

and efficiency. On a group level (Fig. 6A), relative to the null distribution, there were no

difference between patients and controls, and this was true for all network properties (all

|t|<0.4568, all p>0.65). Note that now, there is only a single peak, in the histograms, as the

patient group values fall within the same normal distribution as the controls.

3.2.2 Single-subject comparison of network properties

Next, we looked at the network properties on a single-subject level compared to the

normative group and we observed a difference in the network properties of a subset of the

patients with VOTC resections compared with those of the controls. The four patients with

LH VOTC resections had normal graph-theoretic values on all four dependent measures. The

two patients with RH VOTC resections (KQ and UD), however, exhibited altered network

properties, relative to the controls (Fig. 6B). KQ had higher modularity and characteristic

path length and lower network efficiency (all |t|>2.4, all p<0.05), and UD had higher

characteristic path length and lower network efficiency (both |t|>2.2, both p<0.05).

Importantly, consistent with the results in Section 3.1.4, there were no changes to the network

properties in the two patients with resections outside their VOTC (all |t|<1.94, all p>0.11).

3.2.3 Effects of using different thresholds on the network properties

Network analysis is contingent on using robust connectivity matrices. Thus, we also

explored the effects of using different thresholds in binarizing the connectivity matrices (see

supplementary materials, Fig. D.1). At a higher threshold, there were no significant

differences between any of the patients and controls in any of the dependent measures. At

lower thresholds, only one patient with left resection, OT, had a significantly different path

length and efficiency (all |t|>2.14, all p<0.05), and in the two patients with right resection,

KQ had significantly different properties to controls, while these effects in UD disappeared.

4 Discussion

Surprisingly, in contrast with the visual perceptual impairments that ensue even after

a fairly circumscribed lesion to VOTC in adulthood, visual perception in pediatric epilepsy

patients with relatively large resections appears remarkably normal (as shown in Liu *et al*.,

2018, Liu, Freud *et al*., 2019). The key question addressed here is whether alterations in WM

underlie the positive visuoperceptual outcomes. As has already been documented (Taylor *et*

al., 2015), behavioral impairments are not always correlated with the size and site of affected

cortex, and aberrant cortico-cortical interactions might positively or negatively affect the

expected brain-behavior correspondence. To elucidate possible neural mechanisms

supporting the normal perceptual competence in the patients with unilateral resections

presented here, we characterized the integrity of the two major WM association tracts that

traverse the VOTC, the ILF and the IFOF, as well as the WM network properties.

4.1 Ipsilesional tracts were present in some patients

We first reconstructed the ILF and IFOF, which have previously been shown to

underlie object recognition (Thomas *et al*., 2009; Tavor *et al*., 2014; Behrmann and Plaut,

2013; Decramer *et al*., 2019; Ortibus *et al*., 2011) and mediate visual perception (Mishkin *et*

al., 1983). We were able to delineate all contralesional tracts and even the ipsilesional tracts

in most patients, except in those with extremely large resections of the entire anterior

temporal lobe. That we could not demarcate all ipsilesional tracts was not surprising given

the extent of resection that affected the anterior temporal lobe in some of these patients.

Moreover, the large resections caused automated tractography algorithms to fail in producing

streamlines that survived the tractography thresholds, as voxels containing cerebrospinal

fluid, which now filled the resection site, have different diffusion values to voxels containing

WM. Nevertheless, the mere presence of the ipsilesional ILF and IFOF in most of the patients

suggests that any damage caused by the removal of cortical tissue does not result in an

obvious cascade of axonal injury that affects the entire ipsilesional hemisphere beyond the

resection site.

4.1.1 Microstructural changes were specific to site of resection

To provide a more quantitative analysis of the WM, we characterized the tracts in

terms of their FA, AD, and RD. These indices were chosen because they offer a mechanistic

relationship to the nature of injury (Aung *et al*., 2013); while FA has typically been used to

characterize microstructural integrity, AD and RD have been shown to be more sensitive

biomarkers for WM defects (Arfanakis *et al*., 2002; Xu *et al*., 2007; Moen *et al*., 2016),

though the associated mechanisms might differ. Our results showed that, in children with a

VOTC resection, there was damage to the ipsilesional ILF and IFOF that manifested in one

or all of the following neuroimaging markers: there was within-subject hemispheric

asymmetry in the microstructural indices – FA was lower, while both AD and RD were

higher in the ipsilesional than in the contralesional tract. Importantly, the microstructural

deficits in the pediatric epilepsy resection cases were only observed in regions proximal to

the resection. In the ILF, abnormalities were evident in patients with either anterior or

posterior resection, while in the IFOF, abnormalities were seen only in patients with posterior

resections. This is unsurprising as the trajectory of the IFOF was more medial (Makris *et al*.,

2007), while the anterior resections in the patients here were more lateral, thereby sparing the

IFOF. Distal to the resection (either antero- or retrograde) in the ipsilesional tracts, the

microstructural indices were normal compared to controls. Moreover, the microstructural

indices in the contralesional ILF and IFOF in VOTC resection patients were also normal.

And in the non-VOTC resection patients, both ipsilesional and contralesional ILF and IFOF

also exhibited normal microstructural properties. Together, these results speak to the

specificity of the damage as a result of surgery.

Abnormalities in the microstructural indices in circumscribed regions of the

ipsilesional tracts are informative. AD has been shown to either increase (de Ruiter *et al*.,

2011; Della Nave *et al*., 2011; Zhu *et al*., 2013; Counsell, 2006; Acosta-Cabronero *et al*.,

2010) or decrease (Budde *et al*., 2009; Kubicki *et al*., 2013) as a result of WM pathology.

Meanwhile, RD has been more consistently shown to increase due to demyelination (Song *et*

al., 2002, 2005), and, specifically, myelin sheath degradation (Seal, *et al*., 2008, Klawiter *et*

al., 2011). The results here showed increased AD values that might indicate axonal

fragmentation, resulting in restricted diffusion along the principal tract direction, as well as

increased RD values that might indicate myelin degradation, resulting in diffusion out of the

WM bundle.

As noted previously, these circumscribed findings ought to be interpreted with

caution in that there is a limited number of patients in the sample, and in some patients, the

ipsilesional tracts could not be defined. Nevertheless, the results from the VOTC and the non-

VOTC patients together reveal that these effects are likely a specific outcome of the resection

without further degeneration of the WM as a result of surgery, and not due to a more

pervasive WM deficits linked to epilepsy or other comorbidities.

4.1.2 Normal microstructural indices in VOTC resection

Among the cases presented here, surprisingly, a single patient, OT, exhibited normal

microstructural indices, even in the ipsilesional ILF and IFOF. This cannot be simply due to

the age at surgery, as other patients were either younger (SN, UD) or older (KQ, NN) at the

time of resection and none showed entirely normal microstructural indices. Similarly, the

normal microstructure of the tracts cannot obviously be accounted for in terms of the time

when the data were acquired postoperatively. One plausible explanation for the integrity of

OT's tracts is etiological — OT's surgery was prompted by the presence of a

dysembryoplastic neuroepithelial tumor (DNET). We speculate that the chronic growth of the

DNET triggered reorganization in OT's ipsilesional hemisphere such that the WM tracts were

slowly displaced over time, and consequently, spared from damage due to the resection.

However, we did not see this preservation in the only other patient with a DNET, UD, who

evinced microstructural deficits ipsilesionally and network changes contralesionally. Of note,

however, is that UD's surgery occurred when he was 6 years old, but OT's surgery took place

when he was 13 years old. OT's extended preoperative development may account for the

difference between him and UD, but these speculative comments await additional

experimental confirmation. We also consider UD in further detail below (Section 4.2.1).

4.2 Network properties of the contralesional hemisphere

In search for broader resection-induced changes, we characterized the connectivity of

the contralesional occipito-temporal cortex by measuring transitivity and modularity (as these

reflect segregation of the nodes into sub-networks) and characteristic path length and

efficiency (as these reflect the ease of information flow in the network) (Bullmore and

Sporns, 2009; Rubinov and Sporns, 2010; Paldino *et al*., 2019). There were no between-

group differences across any of these dependent measures. However, on an individual level,

while there were no changes to the network in the contralesional hemisphere of the LH

VOTC resection patients or in the two non-VOTC resection patients, we did observe changes

to the network in the contralesional hemisphere in the RH VOTC resection patients.

A similar pattern of network differences was reported in patients with LH lesions,

previously (Crofts *et al*., 2011). While the etiology of the lesions in those patients – stroke –

is different to the cases presented here, the hemispheric differences bear some resemblance.

Also, Crofts et al. (2012) measured network communicability, a measure of the ease with

which information can travel across a network. This is somewhat similar to the network

efficiency that we measured here and for which we saw significant differences in the RH

VOTC resection. The authors postulated that such alterations possibly arise from secondary

degradation of long-range WM pathways that may or may not be directly connected to the

damaged area.

4.2.1 Differential hemispheric effects of resection on network properties

Given the small number of patients, we cannot definitively attribute the different

profiles of network configuration to differences in hemispheric specialization.

Notwithstanding the recommendation of caution in interpretation, the findings reported here

are provocative enough to warrant some speculation, but additional exploration of this issue

is obviously warranted.

There is well-known hemispheric lateralization of language to the LH and visual

function to the RH in the majority of the population (Corballis, 1983). That we only see

adverse changes to network properties in the two patients with RH VOTC resections might

indicate that, in such cases, WM is reorganized in the contralesional LH to accommodate

functions that are typically lateralized to, and dominant in, the RH. That is, the visual

functions from the RH must be localized elsewhere postoperatively, and a potential location

would be in the homologous LH regions. This transfer of function might be made possible

via the corpus callosum, which plays a key role in plasticity after brain injury (Restani and

Caleo, 2016) and whose maturation depends on coordinated activation across the two

hemispheres (Pietrasanta *et al.*, 2012).

One explanation of why there is no commensurate reorganization of the contralesional

RH following LH VOTC resections is that the RH may already be finely tuned to subserve

visual function. Remember that here, only the occipito-temporal regions were considered in

the network topology. The cases of SN and TC support this view in that we do not see altered

network properties in the contralesional RH when we know from fMRI studies that some

visual functions of the resected regions have indeed emerged in the contralesional RH (as in

the right VWFA in patients SN and TC, Table 1). The idea then is that, for visual function,

the RH is already well tuned and can easily absorb the LH visual function, but the converse

does not hold.

4.2.2 Use of graph theory on structural data from pediatric epilepsy cases

Graph theory measures, applied to structural data, have previously been used in medial temporal lobe epilepsy in adults (Bonilha *et al.*, 2012), where the authors showed normal network efficiency in patients prior to surgery. The absence of network changes pre-operatively suggests that epilepsy itself may not always alter the network efficiency in the epileptic brain, and that any changes are likely resection-induced. However, to our knowledge, aside from the present study, there has not been a systematic study of graph-theoretic measures applied to WM postoperative connectivity in children.

Because network properties rely on robust connectivity matrices, we studied the effects of using different threshold levels on the stability of the measures derived from tractography (see Supplementary Materials, Fig. D1). On the one hand, without thresholding (i.e. using data that potentially include spurious connections with low FA), only patients KQ and OT (right and left resection, respectively) had altered network properties. On the other hand, at an extremely high threshold, there were no alterations to the network properties, likely resulting in false negatives. Notably, in EK and DX (non-VOTC resection patients), the network properties of the contralesional temporo-occipital regions were consistently normal independent of threshold, suggesting that the alterations were likely a result of the resection to the VOTC and not due to the epilepsy per se.

Ultimately, using a reasonable threshold that was used to minimize noise, we found results from the RH VOTC resection cases that are consistent with results from a previous study that showed a moderate decrease in the network efficiency in epileptic patients (Taylor *et al.*, 2018). It must be noted, however, that the result of Taylor et al. (2018) was based on an artificial resection of the (whole-brain) connectome data acquired preoperatively, whereas we provide an explicit characterization of the properties of the contralesional temporo-occipital network using in vivo data acquired postoperatively.

All that being said, we recognize that cognitive impairments in domains that are not

directly associated with visual function (e.g. IQ below 70) might already be present

premorbidly (Veersema *et al.*, 2019) and might impact the network topography

postoperatively. In our patients, only NN (with neurological comorbidities: extant

polymicrogyria and low IQ) and DX showed some intermediate and high-level visual

behavioral deficits (Liu, Freud *et al.*, 2019). The other LH resection patients had no

impairments and alterations of their contralesional network properties, while the RH VOTC

resection patients had no impairments with alterations to their network properties. This

heterogeneity confounds determination of the relationship between network properties vis-a-

vis possible compensatory behavior. Nevertheless, network properties based on resting state

data have already been shown to be reliable biomarkers for cognitive ability in children with

epilepsy (Paldino *et al.*, 2017b). Thus, it may be a worthwhile endeavor to determine whether

network properties, as defined from structural connectivity, can be correlated with behavior

in a larger sample of postoperative participants.

4.3 Normal visual perception despite persistent degradation in ipsilesional hemisphere

The main goal of this study was to determine the possible role of WM changes in

relation to the preserved visual competence in pediatric patients with unilateral resections. A

recent longitudinal study following individuals with anterior temporal lobectomy has

documented changes over time to the FA of various WM tracts, including the bilateral ILF

(Li *et al.*, 2019). However, the authors did not measure behavioral outcomes in these patients.

In Decramer et al (2019), recovery of behavioral function was seen in a single patient who

had persistently damaged WM. As we only have cross-sectional diffusion data, albeit coupled

with several behavior measures, we cannot comment on the longitudinal variations in the

microstructural damage in our patients. However, the data from each patient used in the

analyses were obtained at various times postoperatively, ranging from only a few months to

more than a decade, and the patterns in microstructural damage are, nevertheless, largely

similar across all the cases. It is, therefore, likely that the postoperative focal damage is

persistent in the patients presented here and this is in the face of their largely normal higher-

order perceptual abilities.

In sum, the current findings reveal an apparent dissociation between the postoperative

structural damage and the intact behavioral competence in these patients. This finding

contrasts with several studies reporting an association between compromised VOTC WM

tracts and visual behavioral deficits, such as in prosopagnosia (Gomez *et al*., 2015, Thomas *et*

al., 2009; Geskin and Behrmann, 2018). For instance, in the absence of obvious neurological

damage and ample visual experience, individuals with prosopagnosia do not acquire normal

face recognition skills. In such individuals, both the ILF and IFOF were abnormal and,

interestingly, the severity of the face recognition deficit was correlated with the extent to

which the ILF was compromised (Gomez *et al*., 2015, Thomas *et al*., 2009; Geskin and

Behrmann, 2018). Along similar lines, there is also an association between perception of

facial emotion and damage to the right IFOF (Philippi *et al*., 2009). Together, these findings

demonstrate that WM atypicalities can have clear behavioral consequences. This is just not

the case in many of the patients described here.

One explanation for the largely normal perceptual abilities, that we favor

speculatively, is that the remaining intact cortex – the entire contralesional hemisphere

together with the varying degree of intact tissue in the ipsilesional hemisphere in patients – is

sufficient to support normal function and behavior. That is, the normal WM integrity outside

the resection, without need for any enhancement either ipsi- or contralesionally, already

allows for effective neural activity. The cases of SN and TC with LH VOTC resections offer

strong support for this view. Whereas a typical visual word form area is localized in the LH

in majority of the population, in SN and TC, this region emerged in the RH, and yet, we see

no enhancement of the right WM pathways relative to controls. Additionally, the WM

damage here is well-circumscribed even in the ipsilesional hemisphere. Whereas the

degradation of WM associated with face processing deficits in neurodevelopmental cases is

diffuse across a tract, the damage as a result of surgery in our patients is confined to the site

of resection.

5 Conclusions

The goal of this study was to determine the neural basis of the largely normal

perceptual abilities in children with unilateral resections of visual cortical regions as a result

of pharmacoresistant epilepsy (save for patients NN with neurological comorbidities and

DX). To do so, we adopted two complementary approaches to probe local and broader effects

on the underlying WM. First, the microstructural indices, used to characterize specific fiber

tracts, revealed focal damage only to the ipsilesional ILF and IFOF, without degradation of

tracts distal to the resection. Moreover, the contralesional tracts exhibited normal

microstructural indices indicating preserved WM integrity without enhancement. Second, a

network approach uncovered changes to the contralesional network properties of only the

patients with RH VOTC resection, but not in patients with LH VOTC resection nor in

patients with non-VOTC resection. Because of the small number of participants, namely only

two/four patients with RH/LH VOTC resection, respectively, this finding, although

provocative, requires further confirmation. Taken together, we do not observe obvious

patterns of reorganization of WM. Instead, our findings suggest that, in cases of unilateral

resections to the VOTC in childhood, the category selectivity in the preserved contralesional

hemisphere and the intact tissue in the ipsilesional hemisphere, together with structurally

preserved ILF and IFOF outside the resection, might suffice to subserve normal visual

perception.

Acknowledgments

We thank the patients, the controls, and their families for their time and cooperation. We also

thank Drs John Pyles and Tim Verstynen for assistance in determining the optimal scan

protocols, as well as the MRI technologists Scott Kurdilla and Mark Vignone for help with

the acquisition of the imaging data, and Dr Frank Yeh and the VisCog group at CMU for the

fruitful discussions.

Funding

This work was supported by the National Institutes of Health grant RO1 EY027018 to M.B.

Competing Interests

The authors report no competing interests.

References
649
1. Acosta-Cabronero, J., Williams, G. B., Pengas, G., and Nestor, P. J. Absolute

diffusivities define the landscape of white matter degeneration in Alzheimer's disease.

Brain 2010; 133:529–539.

2. Arfanakis, K., Haughton, V. M., Carew, J. D., Rogers, B. P., Dempsey, R. J., and

Meyerand, M. E. Diffusion tensor MR Imaging in diffuse axonal injury. American

Journal of Neuroradiology 2002; 23:794–802.

3. Aung, W. Y., Mar, S., and Benzinger, T. L. Diffusion tensor MRI as a biomarker in

axonal and myelin damage. Imaging in Medicine 2013; 5:427–440.

4. Behrmann, M. and Plaut, D. C. Distributed circuits, not circumscribed centers,

mediate visual recognition. Trends in Cognitive Sciences 2013; 17:210–219.

5. Bonilha, L., Edwards, J. C., Kinsman, S. L., Morgan, P. S., Fridriksson, J., Rorden,

C., Rumboldt, Z., Roberts, D. R., Eckert, M. A., and Halfor, J. J. Extrahippocampal

gray matter loss and hippocampal deafferentation in patients with temporal lobe

epilepsy. Epilepsia 2010; 51: 519 – 528.

6. Bonilha, L., Nesland, T., Martz, G. U., Joseph, J. E., Spampinato, M. V., Edwards, J.

C., and Tabesh, A. Medial temporal lobe epilepsy is associated with neuronal fibre

loss and paradoxical increase in structural connectivity of limbic structures. Journal of

Neurology, Neurosurgery and Psychiatry 2012; 83:903–909.

7. Budde, M. D., Xie, M., Cross, A. H., and Song, S.K. Axial Diffusivity Is the Primary

Correlate of Axonal Injury in the Experimental Autoimmune Encephalomyelitis

Spinal Cord: A Quantitative Pixelwise Analysis. Journal of Neuroscience 2009;

29:2805–2813.

8. Bullmore, E. and Sporns, O. Complex brain networks: graph theoretical analysis of

structural and functional systems. Nature Reviews Neuroscience 2009; 10:186–98.

9. Catani, M. Diffusion tensor magnetic resonance imaging tractography in cognitive

disorders. Current Opinion in Neurology 2006; 19:599 –606.

10. Catani, M., Jones, D. K., Donato, R., ffytche, D. H. Occipito-temporal connections in

the human brain. Brain 2003; 126:2093 –2107.

11. Corballis, M. C. (1983) *Human Laterality*. Auckland, New Zealand: Academic Press.

12. Counsell, S. J. Axial and Radial Diffusivity in Preterm Infants Who Have Diffuse

White Matter Changes on Magnetic Resonance Imaging at Term-Equivalent Age.

Pediatrics 2006; 117:376–386.

13. Crawford, J. R. and Howell, D. C. Comparing an individual's test score against norms

derived from small Samples. The Clinical Neurophsychologist 1998; 12:482–486.

14. Crofts, J. J., Higham, D. J., Bosnell, R., Jbabdi, S., Matthews, P. M., Behrens, T. E. J.

Johansen-Berg, H. Network analysis detects changes in the contralesional hemisphere

following stroke. NeuroImage 2011; 54:161–169.

15. Dale, A. M., Fischl, B., and Sereno, M. I. Cortical surface-based analysis: I.

Segmentation and surface reconstruction. NeuroImage 1999; 9:179–194.

16. de Ruiter, M. B., Reneman, L., Boogerd, W., Veltman, D. J., Caan, M., Douaud, G.,

Lavini, C., Linn, S. C., Boven, E., van Dam, F. S. A. M., Schagen, S. B. Late effects

of high-dose adjuvant chemotherapy on white and gray matter in breast cancer

survivors: Converging results from multimodal magnetic resonance imaging. Human

Brain Mapping 2011; 33:2971– 2983.

17. Decramer, T., Premereur, E., Lagae, L., van Loon, J., Janssen, P., Sunaert, S., Theys,

T. Patient MW: transient visual hemi-agnosia. Journal of Neurology 2019; 266:691–

698.

697 18. Della Nave, R., Ginestroni, A., Diciotti, S., Salvatore, E., Soricelli, A., and Mascalchi,

698    M. Axial diffusivity is increased in the degenerating superior cerebellar peduncles of

699    Friedreich's ataxia. Neuroradiology 2011; 53:367–372.

700 19. Dennis, E. L., Jin, Y., Kernan, C., Babikian, T., Mink, R., Babbitt, C., Johnson, J.,

701    Giza, C. C., Asarnow, R., Thompson, P. M. White matter integrity in traumatic brain

702    injury: Effects of permissible fiber turning angle. In 2015 IEEE 12th International

703    Symposium on Biomedical Imaging; pages 930–933. IEEE.

704 20. DeSalvo, M. N., Douw, L., Tanaka, N., Reisenberger, C., and Stufflebeam, S. M.

705    Altered Structural Connectome in Temporal Lobe Epilepsy. Neuroradiology 2014;

706    270:842–848.

707 21. Destrieux, C., Fischl, B., Dale, A., and Halgren, E. Automatic parcellation of human

708    cortical gyri and sulci using standard anatomical nomenclature. NeuroImage 2000;

709    53:1–15.

710 22. Fang-Cheng Yeh, Wedeen, V. J., and Tseng, W.-Y. I. Generalized q-sampling

711    imaging. IEEE Transactions on Medical Imaging 2010; 29:1626–1635.

712 23. Geskin, J. and Behrmann, M. Congenital prosopagnosia without object agnosia? A

713    literature review. Cognitive Neuropsychology 2018; 35:4–54.

714 24. Gomez, J., Pestilli, F., Witthoft, N., Golarai, G., Liberman, A., Poltoratski, S., Yoon,

715    J., Grill-Spector, K. Functionally Defined White Matter Reveals Segregated Pathways

716    in Human Ventral Temporal Cortex Associated with Category-Specific Processing.

717    Neuron 2015; 85:216–227.

718 25. Helmstaedter, C., Beeres, K., Elger, C.E., Kuczaty, S., Schramm, J., Hoppe, C. (in

719    press). Cognitive outcome of pediatric epilepsy surgery across ages and different

720    types of surgeries: A monocentric 1-year follow-up study in 306 patients of school

721    age. Seizure. doi: 10.1016/j.seizure.2019.07.021

722 26. Klawiter, E. C., Schmidt, R. E., Trinkause, K., Liang, H. F., Budde, M. D., Naismith,
723   R. T., Song, S. K., Cross, A. H., Benzinger, T. L. Radial diffusivity predicts
724   demyelination in ex vivo multiple sclerosis spinal cords. NeuroImage 2011; 55: 1454
725   −1460.

726 27. Kubicki, M., Shenton, M. E., Maciejewski, P. K., Pelavin, P. E., Hawley, K. J.,
727   Ballinger, T., Swisher, T., Jabbar, G. A. Theremenos, H.W., Keshavan, M.S.,
728   Seidman, L.J., DeLisi, L.E. Decreased axial diffusivity within language connections:
729   A possible biomarker of schizophrenia risk. Schizophrenia Research 2013; 148:67–
730   73.

731 28. Li, W., An, D., Tong, X., Liu, W., Xiao, F., Ren, J., Niu, R., Tang, Y., Zhou, B., Lei,
732   D., Jiang, Y., Luo, C., Yao, D., Gong, Q., Zhou, D. Different patterns of white matter
733   changes after successful surgery of mesial temporal lobe epilepsy. NeuroImage:
734   Clinical 2019; 21:10163.

735 29. Liu, T. T., Freud, E., Patterson, C., and Behrmann, M. Perceptual function and
736   category-selective neural organization in children with resections of visual cortex.
737   The Journal of Neuroscience 2019; 39: 6299 – 6314.

738 30. Liu, T. T., Nestor, A., Vida, M. D., Pyles, J. A., Patterson, C., Yang, Y., Yang, F. N.,
739   Freud, E., Behrmann, M. Successful reorganization of category-selective visual cortex
740   following occipito-temporal lobectomy in childhood. Cell Reports 2018; 24:1113–
741   1122.

742 31. Makris, N., Papadimitriou, G. M., Sorg, S., Kennedy, D. N., Caviness, V. S., and
743   Pandya, D. N. The occipitofrontal fascicle in humans: A quantitative, in vivo, DT-
744   MRI study. NeuroImage 2007; 37:1100-1111.

745 32. Mishkin, M., Ungerleider, L. G., and Macko, K. A. Object vision and spatial vision:
746   two cortical pathways. Trends in Neurosciences 1983; 6:414–417.

747   33. Moen, K. G., Vik, A., Olsen, A., Skandsen, T., Haberg, A. K., Evensen, K. A. I.,

748   Eikenes, L. Traumatic Axonal Injury: Relationships Between Lesions in the Early

749   Phase and Diffusion Tensor Imaging Parameters in the Chronic Phase of Traumatic

750   Brain Injury. Journal of Neuroscience Research 2016; 635:623–635.

751   34. Ortibus, E., Verhoeven, J., Sunaert, S., Casteels, I., de Cock, P., and Lagae, L.

752   Integrity of the inferior longitudinal fasciculus and impaired object recognition in

753   children: A diffusion tensor imaging study. Developmental Medicine and Child

754   Neurology 2011; 54:38–43.

755   35. Paldino, M. J., Chu, Z. D., Chapieski, M. L., Golriz, F., and Zhang, W. Repeatability

756   of graph theoretical metrics derived from resting-state functional networks in

757   pediatric epilepsy patients. British Journal of Radiology 2017a; 90:20160656.

758   36. Paldino, M. J., Golriz, F., Chapieski, M. L., Zhang, W., and Chu, Z. D. Brain network

759   architecture and global intelligence in children with focal epilepsy. American Journal

760   of Neuroradiology 2017b; 38:349–356.

761   37. Paldino, M. J., Golriz, F., Zhang, W., and Chu, Z. Normalization enhances brain

762   network features that predict individual intelligence in children with epilepsy. PLoS

763   ONE 2019; 14:e0212901.

764   38. Paldino, M. J., Hedges, K., Rodrigues, K. M., and Barboriak, D. P. Repeatability of

765   quantitative metrics derived from MR diffusion tractography in pediatric patients with

766   epilepsy. British Journal of Radiology 2014a; 87: 20140095.

767   39. Paldino, M. J., Zhang, W., Chu, Z. D., and Golriz, F. Metrics of brain network

768   architecture capture the impact of disease in children with epilepsy. NeuroImage:

769   Clinical 2017c; 13:201–208.

40. Philippi, C. L., Mehta, S., Grabowski, T., Adolphs, R., and Rudrauf, D. Damage to
Association Fiber Tracts Impairs Recognition of the Facial Expression of Emotion.
The Journal of Neuroscience 2009; 29:15089–15099.

41. Pietrasanta, M., Restani, L., and Caleo, M. The corpus callosum and the visual cortex:
Plasticity is a game for two. Neural Plasticity 2012; 2012:838672.

42. Pustina, D., Doucet, G., Evans, J., Sharan, A., Sperling, M., Skidmore, C., Tracy, J.
Distinct types of white matter changes are observed after anterior temporal lobectomy
in epilepsy. PLoS ONE 2014; 9:e104211.

43. Restani, L. and Caleo, M. Reorganization of Visual Callosal Connections Following
Alterations of Retinal Input and Brain Damage. Frontiers in Systems Neuroscience
2016; 10:86:1– 17.

44. Rubinov, M. and Sporns, O. Complex network measures of brain connectivity: Uses
and interpretations. NeuroImage 2010; 52:1059–1069.

45. Seal, M., Yucel, M., Fornito, A., Wood, S. J., Harrison, B. J., Walterfang, M., Pell, G.
S., Pantelis, C. Abnormal white matter microstructure in schizophrenia: A voxelwise
analysis of axial and radial diffusivity. Schizophrenia Research 2008; 1–3; 106–110.

46. Slinger, G., Sinke, M. R. T., Braun, K. P. J., Otte, W. M. White matter abnormalities
at a regional and voxel level in focal and generalized epilepsy: A systematic review
and meta-analysis. NeuroImage: Clinical 2016; 12: 902 – 909.

47. Song, S. K., Sun, S.W., Ramsbottom, M. J., Chang, C., Russell, J., and Cross, A. H.
Dysmyelination revealed through MRI as increased radial (but unchanged axial)
diffusion of water. NeuroImage 2002; 17:1429–1436.

48. Song, S.K., Yoshino, J., Le, T. Q., Lin, S.J., Sun, S.-W., Cross, A. H., Armstrong, R.
C. Demyelination increases radial diffusivity in corpus callosum of mouse brain.
NeuroImage 2005; 26:132–140.

49. Tavor, I., Yablonski, M., Mezer, A., Rom, S., Assaf, Y., and Yovel, G. Separate parts

of occipito-temporal white matter fibers are associated with recognition of faces and

places. NeuroImage 2014; 86:123–130.

50. Taylor, P. N., Han, C. E., Schoene-Bake, J.C., Weber, B., and Kaiser, M. Structural

connectivity changes in temporal lobe epilepsy: Spatial features contribute more than

topological measures. NeuroImage: Clinical 2015; 8:322–328.

51. Taylor, P. N., Sinha, N., Wang, Y., Vos, S. B., de Tisi, J., Miserocchi, A., McEvoy,

A. W., Winston, G. P., Duncan, J. S. The impact of epilepsy surgery on the structural

connectome and its relation to outcome. NeuroImage: Clinical 2018; 18:202–214.

52. Thomas, C., Avidan, G., Humphreys, K., Jung, K. J., Gao, F., and Behrmann, M.

Reduced structural connectivity in ventral visual cortex in congenital prosopagnosia.

Nature Neuroscience 2009; 12:29–31.

53. Veersema, T. J., van Schooneveld, M. M., Ferrier, C. H., van Eijsden, P., Gosselaar,

P. H., van Rijen, P. C., Spliet, W. G. M., Muhlebner, A., Aronica, E. Braun, K. P. J.

Cognitive functioning after epilepsy surgery in children with mild malformation of

cortical development and focal cortical dysplasia. Epilepsy and Behavior 2019;

94:209–215.

54. Wakana, S., Caprihan, A., Panzenboeck, M. M., Fallon, J. H., Perry, M., Gollub, R.

L., Hua, K., Zhang, J., Jiang, H., Dubey, P., Blitz, A., van Zijl, P., Mori, S.

Reproducibility of quantitative tractography methods applied to cerebral white matter.

NeuroImage, 2007; 36:630– 644.

55. Wheeler-Kingshott, C. A. and Cercignani, M. About "axial" and "radial" diffusivities.

Magnetic Resonance in Medicine 2009; 61:1255–1260.

56. Xu, J., Rasmussen, I.-A. J., Lagopoulos, J., and Haberg, A. Diffuse axonal injury in

severe traumatic brain injury visualized using high-resolution diffusion tensor

imaging. Journal of Neurotrauma 2007; 24:753–765.

57. Yeh, F.C., Verstynen, T. D., Wang, Y., Fernandez-Miranda, J. C., and Tseng, W.Y. I.

Deterministic diffusion fiber tracking improved by quantitative anisotropy. PLoS

ONE 2013; 8:e80713.

58. Zhang, W., Muravina, V., Azencott, R., Chu, Z. D., and Paldino, M. J. Mutual

Information Better Quantifies Brain Network Architecture in Children with Epilepsy.

Computational and Mathematical Methods in Medicine 2018; 2018:6142898.

59. Zhu, T., Zhong, J., Hu, R., Tivarus, M., Ekholm, S., Harezlak, J., Ombao, H., Navia,

B., Cohen, R., Schifitto, G. Patterns of white matter injury in HIV infection after

partial immune reconstitution: A DTI tract-based spatial statistics study. Journal of

NeuroVirology 2013; 19:10–23.

 Figure Legends

Figure 1: ILF and IFOF in patients with VOTC resections. ILF and IFOF in exemplar

control and patients with VOTC resections to their right hemisphere (RH) and left

hemisphere (LH). Anatomical images show the resection site in each patient. Both

contralesional and ipsilesional tracts were present in most patients, except for right

(ipsilesional) ILF and IFOF in KQ, and left (ipsilesional) IFOF in SN. Ages shown for

patients are at time of data acquisition.

Figure 2: Group comparison of microstructural indices between VOTC resection

patients and controls. Results of permutation testing of mean microstructural indices of the

(A) ILF and (B) IFOF. There were significant group differences at $\alpha = 0.05$ between controls

and patients (red dots) for all measures in ipsilesional ILF but only in RD for the IFOF. Both

contralesional tracts had group differences of $p > 0.11$ across all microstructural indices.

Patient data were compared to the mean of the indices of both hemispheres in controls (top

rows) or to indices from either hemisphere (2nd and 3rd rows). Black vertical lines indicate

upper and lower 2.5th percentile of the null distribution (gray histogram).

Figure 3: Microstructural indices of the ILF. (A). Anteroposterior profiles of FA, AD, and

RD in an exemplar control and in patients with resections to their VOTC. Shaded regions

indicate extent of resection. (B). Single-subject comparisons against normative group values

(n=15) showed significant differences between ipsilesional and corresponding ILF in controls

only for values in the segment corresponding to the resection in each patient. Shaded areas

indicate patients with posterior resection, white areas indicate patients with anterior resection.

p-values uncorrected for multiple comparisons: $*p < 0.05, ** p < 0.01, *** p < 0.001$.

Figure 4: Microstructural indices of the IFOF. (A). Anteroposterior profiles of FA, AD,

and RD in an exemplar control and in patients with resections to their VOTC. Shaded regions

indicate extent of resection. (B). Single-subject comparisons against normative group values

showed significant differences between ipsilesional and corresponding IFOF in controls only

for values in patients with posterior resection. Shaded areas indicate patients with posterior

resection, white areas indicate patients with anterior resection. p-values uncorrected for

multiple comparisons: $*p < 0.05, ** p < 0.01, *** p < 0.001$.

Figure 5: ILF and IFOF in children with resections outside VOTC. Microstructural

indices of the (B) ILF and (C) IFOF in patients with resections outside their VOTC. There

were no single-subject level differences to control values for any of the indices of bisected

tracts, all $|t|<1.23$, all $p>0.24$. Same colors and orientation as in Fig. 1.

869
Figure 6: Comparison of graph-theoretic measures between VOTC resection patients

and controls. (A) Results of permutation testing of network properties reveal normal

measures in patients compared to controls on a group level (all $p>0.65$) and are consistent

whether patient data were compared to the mean of both hemispheres in controls (top rows)

or to either hemisphere (2nd and 3rd rows). (B) Patients with LH VOTC resections exhibit

normal network properties while patients with RH VOTC resections (KQ and UD) exhibit

significantly different network properties compared to controls. Shaded areas indicate

patients with posterior resection, white areas indicate patients with anterior resection. p-

values uncorrected for multiple comparisons: $*p < 0.05, ** p < 0.01$.

883 **Figure 2.**

885
886

887   **Figure 3.**

889
890

891   **Figure 4.**

893
894

895   **Figure 5.**

898
899

900   **Figure 6.**
901

902
903

904   Table 1. Surgery and data acquisition history of patients.

Patient	Surgical Procedure	Age at resection	Time between resection & MRI scan	Category selective ROI*
KQ	Right anterior temporal lobectomy and hippocampectomy	15y, 10m	5m	None missing
SN***	Evacuation of left temporal hematoma	1d	12y, 6m	Missing, L: VWFA**
OT	Left temporal lobectomy with preservation of mesial structures, gross total resection of left mesial temporal dysembryoplastic neuroepithelial tumor (DNET)	13y, 4m	4y, 3m	None missing
UD	Right occipital and posterior temporal lobectomy with resection of inferomesial temporal DNET	6y, 9m	4y, 3m	Missing, R: EVC, FFA, PPA, LOC
TC	Left parietal and occipital lobectomy	13y, 3m	8m	Missing, L: EVC, FFA, PPA, LOC, VWFA**
NN	Left occipital lobectomy, left posterior temporal and left parietal corticectomy	15y, 8m	3y, 7m	Missing, L: EVC, PPA, LOC, VWFA
EK	Gross total resection of right frontal neuroglial/gliotic lesion	17y, 1m	8m	None missing
DX***	Left frontal corticectomy with resection of focal cortical dysplastic lesion	15y, 4m	3m	None missing

∗ Category selectivity was defined in (Liu, Freud *et al.*, 2019) and, in each individual, nine

regions were identified where possible: bilateral early visual cortex (EVC), bilateral fusiform

face area (FFA), bilateral parahippocampal place area (PPA), bilateral lateral occipital cortex

(LOC), and left visual word form area (VWFA).

∗∗ SN's and TC's VWFA were localized in the right hemisphere.

∗∗∗ Category selectivity in patients SN and DX has not been documented previously, but

their regions of interest are presented in the supplementary materials (Figs. A.1 and A.2).

913   Table 2. Behavioral measures in patients with unilateral cortical resections

| Patient | Intermediate-level vision | | | High-level vision | | | |
| | Contour integration | | Glass pattern | CFMT-C | | Object matching | |
	Threshold (±0 collinearity)	Threshold (±20 collinearity)	Threshold	Upright faces	Inverted faces	Accuracy	RT
KQ	68.45	74.21	35	90.00%	78.33%	99.00%	1018.14
SN	67.27	80	33.33	95%	73.30%	95.00%	825.7
OT	54.01	65.73	31.67	62.50%	55.56%	99.00%	929.73
UD	51.96	76.88	25.83	83.33%	68.33%	91.00%	1366.96
TC	66.12	77.27	45.83	83.33%	46.67%	89.00%	1047.66
NN	**75.95**	78.63	25.83	80.56%	45.83%	93.00%	**1726.1**
EK	54.01	75.51	54.17	88.33%	81.67%	100.00%	1192.12
DX	**77.27**	80	36.67	100.00%	93.00%	96.00%	1208.9
Control mean	56.45±5.05	74.04±3.53	39.4±7.74	88.22±11.55%	70.36±11.79%	94.86±3.08%	869.03±225.65

915   Only NN and DX showed significant differences in performance relative to controls. Values

916   in bold are significantly different to controls at p<0.05.