Integrating NMR and Simulations Reveals Motions in the UUCG Tetraloop

Sandro Bottaro[1,2]*, **Parker J. Nichols**[3], **Beat Vögeli**[3], **Michele Parrinello**[1], **Kresten Lindorff-Larsen**[2]*

*For correspondence:
sandro.bottaro@iit.it (SB);
lindorff@bio.ku.dk (KLL)

[1]Atomistic Simulations Laboratory, Istituto Italiano di Tecnologia, Genova, Italy; [2]Structural Biology and NMR Laboratory, Department of Biology, University of Copenhagen, Copenhagen, Denmark; [3]Department of Biochemistry and Molecular Genetics, University of Colorado Denver, Anschutz Medical Campus, Aurora, Colorado, USA.

Abstract We provide an atomic-level description of the structure and dynamics of the UUCG RNA stem-loop by combining molecular dynamics simulations with experimental data. The integration of simulations with exact nuclear Overhauser enhancements data allowed us to characterize two distinct states of this molecule. The most stable conformation corresponds to the consensus three-dimensional structure. The second state is characterized by the absence of the peculiar non-Watson-Crick interactions in the loop region. By using machine learning techniques we identify a set of experimental measurements that are most sensitive to the presence of non-native states. We find that although our MD ensemble, as well as the consensus UUCG tetraloop structures, are in good agreement with experiments, there are remaining discrepancies. Together, our results show that i) the structural interpretation of experimental data for dynamic RNAs is highly complex, even for a simple model system such as the UUCG tetraloop ii) the MD simulation overstabilize a non-native loop conformation, and iii) eNOE data support its presence with a population of $\approx 10\%$.

INTRODUCTION

RNA loops are structural elements that cap A-form double helices, and as such are fundamental structural units in RNA molecules. The great majority of known RNA loops contain four nucleotides *Wolters* (*1992*), and these so-called tetraloops are one of the most common and well-studied three-dimensional RNA motifs. The great majority of known RNA tetraloops have the sequence GNRA or UNCG, where

Figure 1. Consensus secondary structure (left) and three dimensional structure (right) of the UUCG tetraloop *Nozinovic et al.* (*2010*). The stem is formed by 5 consecutive Watson-Crick base-pairs capped by the loop U6-U7-C8-G9. One of the most distinctive feature of this structure is the trans-Sugar-Watson interaction between U6 and G9 (bottom). Extended secondary structure annotation follows the Leontis-Westhof nomenclature *Leontis and Westhof* (*2001*)

N is any nucleotide and R is guanine or adenine *Bottaro and Lindorff-Larsen* (*2017*). Their small size, together with their biological relevance, has made these systems primary targets for nuclear magnetic resonance (NMR) spectroscopy, X-ray-crystallography, and atomistic molecular dynamics (MD) simulation studies *Cheong et al.* (*1990*); *Woese et al.* (*1990*); *Ferner et al.* (*2008*).

The UUCG tetraloop has been long known to be highly stable, and both crystallographic and NMR studies suggest that this tetraloop adopts a well-defined three dimensional structure including a characteristic trans-Sugar-Watson (tSW) interaction between U6 and G9 *Ennifar et al.* (*2000*); *Nozinovic et al.* (*2010*) (Fig. 1). Experimentally, the UUCG tetraloop is used to stabilize the secondary structure of larger RNA molecules without interacting with other RNAs or proteins *Hall* (*2015*).

Despite its stability, the UUCG tetraloop is not rigid. In particular, three recent studies by independent groups indicate the presence of alternative loop conformations *Borkar et al.* (*2017*); *Hartlmüller et al.* (*2017*); *Nichols et al.* (*2018b*). Earlier NMR studies *Nozinovic et al.* (*2010*); *Duchardt and Schwalbe* (*2005*) also suggested the presence of loop dynamics, without providing a detailed structural interpretation of the data. More generally, the atomic-detailed characterization

of RNA structure and dynamics requires specialized techniques and substantial experimental effort, including NMR measurements of nuclear Overhauser effects (NOE), scalar couplings, chemical shifts, residual dipolar couplings, cross-correlated relaxation rates as well as a wide range of relaxation-dispersion type NMR experiments *Salmon et al.* (*2014*); *Marušič et al.* (*2019*).

While NOEs are typically used to determine RNA and protein structures, they also contain dynamic information. Because ensemble-averaged NOEs are highly sensitive to the underlying distance fluctuations, they may contain contributions even from minor populations. Normally, such information is difficult to extract because standard NOE measurements are relatively inaccurate. It has, however, been demonstrated that a substantial part of the information content inherent to these probes can be obtained from exact NOE measurements (eNOEs) *Nichols et al.* (*2018b*). As opposed to conventional NOEs, eNOEs can be converted into tight upper and lower distance limit restraints *Vögeli* (*2014*); *Nichols et al.* (*2017*, 2018a).

Previous computational studies of the UUCG tetraloops focused either on the dynamics around the near-native state *Giambaşu et al.* (*2015*) or on the difficulty in separating force-field inaccuracies from insufficient sampling *Banás et al.* (*2010*); *Bergonzo et al.* (*2015*). In a previous study we reported converged free-energy landscape for RNA 8-mer and 6-mer loops, and we have shown that native-like states are not the global free-energy minimum using the current AMBER RNA force-field *Bottaro et al.* (*2016*). This problem has been addressed in a new parameterization of the AMBER force-field, that improves the description of the UUCG 14-mer and other RNA systems *Tan et al.* (*2018*). Nevertheless, it remains difficult to assess the accuracy of these simulations, because experiments alone do not provide an atomic-detailed description of structure and dynamics that serve as a benchmark.

Here, we use extensive atomistic MD simulations to map the conformational landscape of the UUCG tetraloop using enhanced sampling techniques and a recent force-field parameterization. To improve the description of this system further, we perform an a posteriori refinement of the MD simulation using experimental data via a Bayesian/maximum entropy procedure *Hummer and Köfinger* (*2015*); *Bottaro et al.* (*2018a*). We validate the eNOE-refined ensemble against independent NMR measurements and find an agreement that is on average comparable with NMR structures of the UUCG tetraloop deposited in the Protein Data Bank (PDB).

Our experimentally-refined ensemble reveals the presence of two conformational states. The dominant, major state (here called state A) is the consensus UUCG structure shown in Fig. 1. The second, previously unreported lowly-populated state (state B) is characterized by the absence of the signature U6-G9 non-Watson-Crick base pair, with the C8 and G9 bases exposed into solution. We

employ a random forest classifier to identify the structural properties that discriminate state A from state B. Furthermore, we use the same method in the space of experimental data to identify specific measurements that are most sensitive to the presence of state B. By construction our refined ensemble better agrees with eNOE compared to the original MD simulation and to the consensus PBD structures.

The paper is organized as follows: we first compare the predictions obtained from MD simulation against different experimental datasets. We then discuss the effect of the refinement procedure, showing how it improves the agreement with experiments and how it affects the population of different conformations. We proceed by identifying the relevant degrees of freedom and contacts that characterize the two states. Finally, we identify experiments sensitive to the presence of state B. We accompany this paper with the commented code, in form of Jupyter notebooks, to reproduce step-by-step the complete analysis, including all figures and supplementary results presented in the manuscript.

Results

MD simulations and comparison with experimental data

We simulate the RNA 14-mer with sequence `GGCACUUCGGUGCC` starting from a completely extended conformation. Studying the folding free-energy landscape of this system is computationally expensive: for this reason previous attempts required μs-long simulations in combination with tempering protocols *Tan et al.* (*2018*); *Kuhrova et al.* (*2013*); *Chen and García* (*2013*).

Here, we combine two enhanced sampling techniques: solute tempering in the REST2 formulation *Wang et al.* (*2011*) and well-tempered metadynamics *Barducci et al.* (*2008*). We used a nucleic-acid specific metric, called eRMSD, *Bottaro et al.* (*2014*) as a collective variable for enhanced sampling. The MD simulation setup and convergence analysis are presented in supporting information 1 (SI1).

Before describing the conformational ensemble provided by MD, we compare the computational prediction with available NMR spectroscopy data. More precisely, we consider the following experimental datasets:

- **Dataset A**. Exact eNOEs *Nichols et al.* (*2018b*), consisting in 62 bidirectional exact NOE, 177 unidirectional eNOE and 77 generic normalized eNOE (gn-eNOE). This dataset alone was used to determine the structure of the UUCG tetraloop with PDB accession codes 6BY4 and 6BY5. In addition to the original dataset, we added 1 new eNOE and 6 new gn-eNOEs, as described in SI2.
- **Dataset B**. 97 ^{3}J scalar couplings, 31 RDCs and 250 NOE distances. This data, among other NMR measurements, was used to calculate the consensus UUCG tetraloop structure (PDB 2KOC *Nozinovic et al.* (*2010*)).

• **Dataset C**. 38 (RDC1) plus 13 (RDC2) residual dipolar couplings. These RDCs
have been used in conjunction with MD simulations to obtain a dynamic
ensemble of the UUCG tetraloop. *Borkar et al.* (*2017*).
• **Dataset D**. 91 solvent paramagnetic resonance enhancement (sPRE) mea-
surements *Hartlmüller et al.* (*2017*).

The orange bars in the four panels of Fig. 2 show the agreement between
simulation and the different experimental datasets. The agreement between ex-
periment and simulations is expressed using the reduced χ^2 statistics, defined as
the average square difference between the experimental measurement (F^{exp}) and
the back-calculated ensemble average ($< F(\mathbf{x}) >$) normalized by the experimental
error σ:

$$\chi^2 = \frac{1}{m} \sum_i^m \frac{(< F(\mathbf{x}) >_i - F_i^{EXP})^2}{\sigma_i^2} \tag{1}$$

143 Hence, the lower the χ^2, the better the agreement. As a rule of thumb, $\chi^2 < 1$
144 can be considered small, as the difference between experiment and prediction
145 is within experimental error. See SI2 and SI3 for additional details on this com-
parison. As a reference, we report in Fig. 2 the agreement calculated on the

Figure 2. Comparison between experiment and simulations. **a)** χ^2 between experimental dataset A against the MD ensemble (MD) and against the refined ensembles (MD+set A, MD+set B, MD+set C, MD+set D). As a reference, values calculated from all NMR models from PDB structures 2KOC and 6BY5 are shown as dashed lines. The agreement between the same ensembles and datasets B, C, D, are shown in panels **b**,**c**, and **d**, respectively. Error bars show the standard error estimated using four blocks.

147 PDB ensembles 6BY5 *Nichols et al.* (*2018b*) and 2KOC *Nozinovic et al.* (*2010*). For

---

set A, the agreement of the MD with experiment is considerably poorer than the
one calculated on 6BY5. We recall that this latter ensemble was determined by
fitting dataset A, we thus expect χ^2 to be small in this case. On datasets B, D the
MD better agrees with experiments compared to 2KOC, 6BY5, while differences
on set C are smaller. When considering other statistics (e.g. root mean square
error, Pearson correlation, number of violations), the same conclusions apply.
Note that χ^2 for sPRE and for scalar couplings (SI 3) is very large. This discrepancy
may arise both from the imperfect ensembles, from the underestimation of the
experimental error, as well as from the limitation of the function used to calculate
the experimental quantity from the atomic positions (i.e. the forward model). As
an example, the parameters in the Karplus equation for HCOP couplings critically
depend on a single experimental data point measured in 1969 *Bentrude and*
Hargis (*1969*).

Bayesian/Maximum entropy refinement of the MD ensemble

As described above, our MD simulation provides a conformational ensemble
consisting of a rich and diverse set of conformations, that, however, as a whole
does not match all experimental data perfectly.

In order to improve the description provided by the MD simulation, we calcu-
late a refined conformational ensemble by a posteriori including experimental
information into simulations. In brief, the refinement is obtained by assigning a
new weight to each MD snapshot, in such a way that the averages calculated with
these new weights match a set of input (or "training") experimental data within a
given error. Among all the possible solutions to this underdetermined problem,
we use the one that maximize the Shannon cross-entropy *Pitera and Chodera*
(*2012*); *Boomsma et al.* (*2014*); *Orioli et al.* (*2019*).

Since we have four independent experimental datasets, we can refine the
simulation by using one of them, using the other three as test set. We choose the
free hyper-parameter of the algorithm (θ) by performing a k-fold cross-validation
procedure (see SI 4). By construction, the refinement procedure improves the
agreement on the training data, but it does not guarantee improved agreement
on the test sets. Figure 2 shows the effect of the refinement for different com-
binations of training/test set. For example, the blue bar in panel **a** shows the
agreement between experiment and the MD simulations refined against set A
(MD+set A) evaluated on set A itself (training). The blue bars in panels **b,c,d** show
the χ^2 of MD+set A on datasets B,C,D (test). The MD+set B is trained on set B and
tested on datasets A,C,D, and so on.

We observe that including into simulations dataset A has a small, yet detrimen-
tal effect on the agreement with sets B and D. The MD+set B ensemble shows
improved agreement with dataset D, but performs worse than the original MD on
set A and C. We observe a similar behavior for MD+set D. Among the four refined

ensembles the MD+set C is the one that behaves better, as it shows a smaller or equal χ^2 (relative to MD) on the three tests sets A,B,D.

Taken together, our results show that the MD and the refined ensembles fit available experimental data to a degree that is comparable to the one calculated from PDB structures 2KOC and 6BY5 (Fig. 2). However, there exists substantial discrepancies with experimental data, and none of the considered ensembles clearly outperforms the others.

Free energy landscape

In this section we analyse in detail the refined MD ensembles, and discuss the differences with respect to the original simulation and previously determined structures. We consider the histogram of the distance (structural dissimilarity) from the consensus structure (PDB 2KOC). Distances are measured using the eRMSD, a nucleic-acid specific metric that takes into account both position and orientations between nucleobases *Bottaro et al.* (*2014*), although a similar picture is obtained using the standard RMSD metric.

The distribution obtained from the original MD ensemble is shown in orange in Fig. 3**a**, and repeated in the four panels. The two peaks correspond to structures where the stem is fully formed, but with different loop conformations. Structures in the left peak (eRMSD<0.7, state A) display the signature interactions present in the consensus structure, while in the second peak (state B) the loop is disordered. The population of state A in our MD is $60\% \pm 4$, larger than the one calculated from the 180μs simulated tempering simulation by Tan et al *Tan et al.* (*2018*) ($\approx 40\%$).

The inclusion of experimental data in simulations affects the histogram in different ways. When including eNOE data from dataset A (Fig 3a, blue), we observe an increase in the population of state A to $83\% \pm 16$. The refinement obtained with datasets B and D (Fig. 3 panels **b**,**d**) reduces the population of state A down to $44\% \pm 6$ and $44\% \pm 10$, respectively. In the MD+set C ensemble, the population of state A is unaffected (65 ± 5). Interestingly, we observe one new peak appearing at eRMSD > 1.5, also present in the MD+set B ensemble. This peak corresponds to fully extended conformations, and it is present when including RDC measurements. If we trust our simulation, the forward model including the complex issue of estimating the alignment tensor, and the refinement procedure, we can conclude that RDC are sensitive to a small population of either dimers or unfolded structures. Unfortunately, we do not have the possibility to test this hypothesis, which is left for future investigations. Here, instead, we focus on the results obtained from MD+set A. First, because this ensemble has the highest population of state A, and it is thus less "surprising". Second, because we better understand and control the experimental data, and lastly because we can take advantage of our previous experience in using NOE for RNA ensemble refinement *Bottaro et al.* (*2018a*).

Figure 3. Histograms of the eRMSD from native. The original MD simulation (orange) is compared with the four refined ensembles: MD+set A in panel **a**, MD+set B in panel **b**, MD+set C in panel **c**, and MD+set D in **d**. Shades show the standard error estimated using four blocks. The vertical dashed line in panel **a** shows the separation between state A (eRMSD < 0.7) and state B (eRMSD ≥ 0.7).

Structural differences between state A and B

Having discovered this new B-state, we proceed to analyse its structural features. While state A is known and structurally well-defined (Fig. 4a), it is not trivial from a simple visual inspection to identify which are the main features of state B (Fig. 4b). Here, we address this question by using a random forest classifier. In practice, we first extract samples from the MD+set A ensemble using a bootstrapping procedure. Second, we assign a label to each sample depending on the distance from native (state A if eRMSD<0.7, and state B otherwise). Last, we calculate structural properties (e.g. torsion angles, distances) for each of the the labelled sample, that are used to train a random forest classifier. In this context we are not interested in the decision tree per se, but rather in its ability to rank the importance of the input features in the classification problem. Thus, we can find the most relevant degrees of freedom that discriminate the two states. The result of such analysis on all dihedral angles $\alpha, \beta, \gamma, \delta, \epsilon, \zeta, \chi$ in the 14-mer reveals that the two highest-ranked angles are ζ in C8 and G9. Fig. 4d,e show the histograms for these two angles: the angle in C8 is a good classifier, as all samples from state A (empty curve) are in $gauche+$ conformation, while all samples from state B (filled curve) are in $gauche-$. Information on state A/B differences are also contained in G9 ζ, although the separation of the two states is not as striking as in C8. The

importance of C8 and G9 is further confirmed when using the distance between the center of the six-membered rings in the nucleobases as input features. In this case, the distances between U6-C8 and U6-G9 are the two most important degrees of freedom that distinguish state A from state B. The distribution of these two distances is shown in Fig. 4c. In the consensus structure U6 and G9 interact through a trans sugar-Watson base-pair and U6 and C8 are stacked. State B is characterized instead by the absence of the stacking interaction (large U6-C8 distance), and of the non-Watson-Crick base-pair (large U6-G9 distance).

Figure 4. a) Representative (random) conformations sampled from state A. **b**) Representative (random) conformations sampled from state B. The color code is identical to Fig. 1: U6 in ochre, U7 in purple, C8 in green and G9 in red. **c**) Free energy surface projected onto the the U6-C8/U6-G9 distance between ring centres. The units of the colorbar are in $k_B T$. **d**) Histogram of ζ dihedral angle in C8. The open filled area indicates conformations belonging to state A, and the filled area indicates conformations belonging to state B. **e**) Histogram of ζ dihedral angle in G9.

255 **Experimental measurements sensitive to the presence of state B.**

In the previous section we have described the structural differences between
state A and state B, and we now seek for additional experimental validation. In
particular, we would like to answer the following question: does the MD+set A
ensemble provide a better loop description compared to the consensus NMR
structure 2KOC?

The four experimental sets contain hundreds of datapoints in total, and it is
therefore not trivial to identify specific measurements that probe directly the pres-
ence of state B. In machine learning this is called a feature importance problem,
that we solve using a random forest classifier, as we did in the previous section.
Here, however, the features are not structural properties, but back-calculated
experimental data. We construct a random forest classifier for each experimental
dataset, and we obtain four sub-datasets with the features (measurements) that
are most important for classifying (i.e. separating) state A and B. Consistently with
Fig. 4, we find that these measurements involve nucleotides in the loop region.

For each measurement in the sub-datasets, we calculate the difference be-
tween the ensemble average and experiments divided by the error σ (i.e. the
Z-score).

$$|\text{Z-score}|_i = \frac{|<F(\mathbf{x})>_i - F_i^{EXP}|}{\sigma_i} \tag{2}$$

274 Figure 5 shows the Z-score calculated on the MD+set A ensemble (x-axis)
275 against the same quantity calculated on the 2KOC structures. Points above the
276 diagonal (such as 1 in panel a, 6 in panel d) indicate measurements for which
277 the MD+set A ensemble better agrees with experimental data ($|\text{Z-score}|_{MD+set A} <$
278 $|\text{Z-score}|_{2KOC}$), while points below the diagonal (e.g. 3,4,7,8) indicate the opposite.
279

Overall, there are more points below the diagonal (42) than above (36), sug-
gesting that the original 2KOC ensemble provide a better description of the loop
region. For example, the RDC in set C 8C3'-8H3 is within experimental error for
2KOC, but not for our MD+set A ensemble (point 4, Fig. 6). Conversely, the eNOE
C8 H5' to G9 H8 in dataset A is significantly closer to the experiment in the MD+set
A ensemble (point 1, Fig. 6). Note that major discrepancies are present in both
ensembles, such as point 2 (^{3}J in G9 H5''-H4), point 3(RDC C8 C1'-H1'), point 5 (RDC
G10 C8-H8), and point 7 (sPRE G10 1H5'). Again, we stress that these discrepan-
cies can be ascribed to errors in the ensembles, but also to inaccuracies in the
empirical model employed to calculate experimental data from structures, or to
errors in the data.

Figure 5. Comparison between 2KOC and MD+set A ensemble on experimental data that are most sensitive to the presence of state B. Scatter plots show the Z-score calculated on the MD+set A ensemble (x-axis) versus the same quantity calculated on the PDB ensemble 2KOC (y-axis). The four panels show data belonging to the four datasets: set A in panel **a**, set B in panel **b**, set C in panel **c**, and set D in panel **d**. Points discussed in the main text and shown in Fig. 6 are labeled in red.

Figure 6. Comparison between calculated and experimental data for selected measurements discussed in the text and labelled as in Fig.5.

Conclusions

Based on our extensive MD simulations and integrating them with exact NOE data, we report the free energy landscape of a prototype stem-loop RNA 14-mer

known as the UUCG tetraloop. By combining a recently refined force field for RNA with enhanced sampling MD we were able to fold the tetraloop to its native conformation(s) as judged by very good agreement with several sets of experiments. The main finding of the present study is the presence of a low populated, non-native conformation (state B). The low-populated state differs from the consensus structure (state A) only in the loop region, and it is characterized by the absence of the tSW base-pair between U6 and G9, with C8 and G9 partially exposed into solution (Fig. 4). This result has been obtained by using atomistic MD simulations and eNOE, without the need of additional data.

The free-energy surfaces and estimated population provided here are based on the available experimental data, on the employed model, and the extent of our sampling. Therefore, they are subject to inaccuracies. However, both simulations and eNOE data are consistent with the presence of the B state as described in this paper. This interpretation is qualitatively consistent with several NMR studies, that also suggested the presence of dynamics in G9 *Duchardt and Schwalbe* (*2005*); *Nozinovic et al.* (*2010*); *Hartlmüller et al.* (*2017*). Conversely, on-resonance 13C R1ρ relaxation dispersion experiments on a UUCG tetraloop with a different stem sequence showed no significant exchange contributions, indicating the absence of motions with substantial chemical shift variation in the μ-ms timescale *Salmon et al.* (*2015*). Note also that G9-exposed structures were reported in previous MD simulations *Kuhrova et al.* (*2013*); *Bergonzo et al.* (*2015*); *Cesari et al.* (*2019*), suggesting our finding to be robust with respect to the choice of the force-field and water model.

In this work we have used eNOEs to reweight a posteriori the ensemble generated via enhanced sampling MD simulations. This refinement procedure is a computationally cheap post-processing *Graf et al.* (*2007*); *Hummer and Köfinger* (*2015*); *Bottaro et al.* (*2018a*) that allows one to try different combinations of training and test sets, as we did in this work *Orioli et al.* (*2019*). Note that refinement is in principle less powerful compared to on-the-fly methods that samples directly from the target probability distribution *Bonomi et al.* (*2017*); *Reißer et al.* (*2019*).

In our study we refine the simulation by matching RDC data (set B, C) or solvent PRE (set D) as well. Only when we use set C for training we obtain an improved or equal agreement on the test sets relative to the original MD simulation (Fig. 2). Additionally, different data affect the MD conformational ensemble in different ways (Fig. 3). Several reasons can contribute to this behaviour. First, we do not expect all experimental data to be perfectly compatible one with the other, because measurements were conducted in similar, but not identical conditions. Second, the forward models might not be accurate for arbitrary molecular conformations. For example, if the forward model accurately predicts the RDC given the native structure, but fails on unfolded/misfolded conformations, we obtain artefacts that cannot be easily accounted for in our refinement procedure. Note that this

problem is typically less relevant when using experimental RDC, sPRE or chemical
shift data for scoring structures *Sripakdeevong et al.* (*2014*); *Salmon et al.* (*2015*);
Hartlmüller et al. (*2017*).

Based on the above observations, and considering our previous experience
with eNOE data, we here analyse in detail the results obtained using MD refined
using set A (Fig. 3a). The structural features that are most important to discrimi-
nate between state A and state B are identified using a random forest classifier.
The problem of concisely interpret differences in biomolecular conformations
has been recently pursued using a variety of machine learning methods, includ-
ing linear discriminant analysis *Piccini et al.* (*2018*), decision trees *Brandt et al.*
(*2018*), and others *Fleetwood et al.* (*2019*). In this work we extend this idea, and
use back-calculated experimental data as input features for the random forest
classifier. In this way we identify individual (available) measurements that are
most sensitive to the presence of state B. Note that this approach can also be
used in a generative fashion to design experiments that probe the existence of
specific conformational states.

We closely inspect the selected set of measurements that are sensitive to state
B (Fig. 5). In the majority of the cases we find the presence of the additional state
to provide a worse agreement with experiment compared to the consensus NMR
structure (PDB code 2KOC) (see e.g. Fig. 6, points 3,8). In other cases (Fig. 6,
points 1,6), instead, the MD+set A performs better than 2KOC. Several other data
significantly deviate from experiments in both ensembles (Fig. 6, points 2,5,7).
This suggest the possibility that conformations that are different from state A are
indeed present, but do not correspond to the state B as described in Fig.4b.

Finally, we note that the approach taken here is general and it is applicable
to other RNA or protein systems *Escobedo et al.* (*2019*); *Crehuet et al.* (*2019*).
Previous characterization of slow, larger motions in RNA molecules have mostly
relied on relaxation-dispersion, chemical exchange saturation transfer or related
NMR experiments that probe chemical shift differences between different con-
formational states. We hope that the integration of MD simulations and eNOE
measurements provides further opportunities for characterizing the free energy
landscapes of RNA molecules.

Acknowledgements

The research is funded by a grant from The Velux Foundations (S.B. and K.L.-L.),
the Lundbeck Foundation BRAINSTRUC initiative (K.L.-L.), a start-up package from
the University of Colorado (B.V.), and by NSF grant 1917254 for Infrastructure
Innovation for Biological Research (B.V.). The authors thank Giovanni Bussi for
helpful comments. We also thank Tobias Madl, Harald Schwalbe, and coworkers
for useful discussions and for sharing their data.

Data Availability

Jupyter notebooks to reproduce the analysis and all figures are included as supporting information. The MD trajectory, together with the experimental data are hosted on github (https://github.com/KULL-Centre/papers/tree/master/2020/UUCG-dynamics-Bottaro-et-al). The Plumed input file to reproduce the simulation is hosted on the Plumed nest *Bonomi et al.* (*2019*) under the accession code 19.070.

Methods

Integrating MD simulation and experimental data

We combine the MD simulation with experimental data using a maximum entropy/Bayesian procedure *Boomsma et al.* (*2014*); *Beauchamp et al.* (*2014*); *Hummer and Köfinger* (*2015*). In our previous work, we have described this reweighting procedure as Bayesian/MaxEnt (BME) *Bottaro et al.* (*2018b*,a) . In BME we use the experimental data to modify a posteriori the simulation so that the new conformational ensemble has the following properties: (i) the calculated averages are close to the experimental values taking uncertainty into account and (ii) it maximizes the relative Shannon entropy with respect to the original simulation ensemble. The modification comes in the form of a new set of weights w_j^*, one for each simulation frame.

It can be shown that this problem can be cast as a minimization problem, in which one seeks the minimum of the function Γ with respect to the set of Lagrange multipliers $\bar{\lambda} = \lambda_1 \cdots \lambda_m$, with m being the number of experimental constraints.

$$\Gamma(\bar{\lambda}) = \log(Z(\bar{\lambda})) + \sum_i^m \lambda_i F_i^{\text{exp}} + \frac{\theta}{2} \sum_i^m \lambda_i^2 \sigma_i^2 \qquad (3)$$

Here, σ_i are the uncertainties on the experimental measurements F_i^{exp} and include experimental errors and inaccuracies introduced by the calculation of the experimental quantity from the atomic positions ($F(\mathbf{x})$). θ is a free parameter, while the partition function Z is defined as

$$Z(\bar{\lambda}) = \sum_{j=1}^n w_j^0 \exp[-\sum_i^m \lambda_i F_i(\mathbf{x}_j)] \qquad (4)$$

The sum over the index j runs over the n frames in the simulation, and w_j^0 are the original weights. $w^0 = 1/n$ when using plain MD simulations or enhanced sampling techniques that sample directly from the target distribution (e.g. parallel tempering). In this paper we use WT-METAD, and the original weights w^0 are estimated using the final bias potential *Branduardi et al.* (*2012*). The minimization of Eq. 3 yields a set of Lagrange multipliers $\bar{\lambda}^*$ that are used to calculate the

optimal weights

$$w_j^* = \frac{1}{Z(\bar{\lambda}^*)} w_j^0 \exp[-\sum_i^m \lambda_i^* F_i(\mathbf{x}_j)] \tag{5}$$

In the context of the UUCG tetraloop, we use the dataset A described in the previous section to refine the simulation ensemble, and cross-validate the results against datasets B, C, and D. Details on the comparison between simulations and experiments, on the BME procedure and on the choice of the regularization parameter θ can be found in SI 4.

Random Forest Classifier

The random forest analysis is set up according to the following procedure:

- Bootstrap $n = 50000$ samples from the MD simulation trajectories. The weight of each sample, w_j^*, is calculated as described above.
- Samples are assigned to state A if the eRMSD from the first model of the NMR structure 2KOC is less than 0.7, and state B otherwise.
- Calculate a set of m features (torsion angles, distances between ring centres, experimental data) for each sample **Bottaro et al.** (**2019**).
- Construct a random forest classifier using n samples, m features and 2 classes (state A and state B). In our work, we use the implementation of the random forest algorithm **Breiman** (**2001**) available in sklearn 0.22 using a maximum tree depth of 3. 80% of the samples are used for training, while the remaining 20% are used to evaluate the accuracy of the classifier (>97% in all cases).
- Rank the m features by their importance. In the analysis of the experimental data in Fig. 5, only features with importance greater than 0.2 are shown.

References

Banás P, Hollas D, Zgarbová M, Jurecka P, Orozco M, Cheatham III TE, Sponer J, Otyepka M. Performance of molecular mechanics force fields for RNA simulations: stability of UUCG and GNRA hairpins. J Chem Theory Comput. 2010; 6(12):3836–3849.

Barducci A, Bussi G, Parrinello M. Well-tempered metadynamics: a smoothly converging and tunable free-energy method. Phys Rev Lett. 2008; 100(2):020603.

Beauchamp KA, Pande VS, Das R. Bayesian energy landscape tilting: towards concordant models of molecular ensembles. Biophys J. 2014; 106(6):1381–1390.

Bentrude WG, Hargis JH. Conformations of 6-membered-ring phosphorus heterocycles: the 5-t-butyl-2-oxo-1, 3, 2-dioxaphosphorinans. J Chem Soc D. 1969; (19):1113b–1114.

Bergonzo C, Henriksen NM, Roe DR, Cheatham TE. Highly sampled tetranucleotide and tetraloop motifs enable evaluation of common RNA force fields. RNA. 2015; 21(9):1578–1590.

Bonomi M, Bussi G, Camilloni C, Tribello G, Bonas P, Barducci A, Bernetti M, Bolhuis PG,
Bottaro S, Branduardi D, et al. Promoting transparency and reproducibility in enhanced
molecular simulations. Nature methods. 2019; 16(8):670–673.

Bonomi M, Heller GT, Camilloni C, Vendruscolo M. Principles of protein structural ensem-
ble determination. Curr Opin Struct Biol. 2017; 42:106–116.

Boomsma W, Ferkinghoff-Borg J, Lindorff-Larsen K. Combining experiments and simula-
tions using the maximum entropy principle. PLoS Comput Biol. 2014; 10(2):e1003406.

Borkar AN, Vallurupalli P, Camilloni C, Kay LE, Vendruscolo M. Simultaneous NMR charac-
terisation of multiple minima in the free energy landscape of an RNA UUCG tetraloop.
Phys Chem Chem Phys. 2017; 19(4):2797–2804.

Bottaro S, Banas P, Sponer J, Bussi G. Free energy landscape of GAGA and UUCG RNA
tetraloops. J Phys Chem Lett. 2016; 7(20):4032–4038.

Bottaro S, Bengtsen T, Lindorff-Larsen K. Integrating Molecular Simulation and Experi-
mental Data: A Bayesian/Maximum Entropy Reweighting Approach. bioRxiv. 2018; p.
457952.

Bottaro S, Bussi G, Kennedy SD, Turner DH, Lindorff-Larsen K. Conformational ensembles
of RNA oligonucleotides from integrating NMR and molecular simulations. Sci Adv.
2018; 4(5):eaar8521.

Bottaro S, Bussi G, Pinamonti G, Reisser S, Boomsma W, Lindorff-Larsen K. Barnaba:
software for analysis of nucleic acid structures and trajectories. RNA. 2019; 25(2):219–
231.

Bottaro S, Di Palma F, Bussi G. The role of nucleobase interactions in RNA structure and
dynamics. Nucl Acids Res. 2014; 42(21):13306–13314.

Bottaro S, Lindorff-Larsen K. Mapping the universe of RNA tetraloop folds. Biophys J.
2017; 113(2):257–267.

Brandt S, Sittel F, Ernst M, Stock G. Machine learning of biomolecular reaction coordinates.
The journal of physical chemistry letters. 2018; 9(9):2144–2150.

Branduardi D, Bussi G, Parrinello M. Metadynamics with adaptive Gaussians. Journal of
chemical theory and computation. 2012; 8(7):2247–2254.

Breiman L. Random forests. Machine learning. 2001; 45(1):5–32.

Cesari A, Bottaro S, Lindorff-Larsen K, Banáš P, Sponer J, Bussi G. Fitting corrections to an
RNA force field using experimental data. Journal of chemical theory and computation.
2019; .

Chen AA, García AE. High-resolution reversible folding of hyperstable RNA tetraloops
using molecular dynamics simulations. Proc Natl Acad Sci USA. 2013; 110(42):16820–
16825.

Cheong C, Varani G, Tinoco Jr I. Solution structure of an unusually stable RNA hairpin,
5GGAC (UUCG) GUCC. Nature. 1990; 346(6285):680.

Crehuet R, Jorro PJB, Lindorff-Larsen K, Salvatella X. Bayesian-Maximum-Entropy reweight-
ing of IDPs ensembles based on NMR chemical shifts. BioRxiv. 2019; p. 689083.

Duchardt E, Schwalbe H. Residue specific ribose and nucleobase dynamics of the cUUCGg
RNA tetraloop motif by MNMR 13 C relaxation. J Biomol NMR. 2005; 32(4):295–308.

Ennifar E, Nikulin A, Tishchenko S, Serganov A, Nevskaya N, Garber M, Ehresmann B,
Ehresmann C, Nikonov S, Dumas P. The crystal structure of UUCG tetraloop1. J Mol Biol.
2000; 304(1):35–42.

Escobedo A, Topal B, Kunze MBA, Aranda J, Chiesa G, Mungianu D, Bernardo-Seisedos G,
Eftekharzadeh B, Gairi M, Pieratelli R, et al. Side chain to main chain hydrogen bonds
stabilize polyglutamine helices in transcription factors. Nat Comm. 2019; 10(2034).

Ferner J, Villa A, Duchardt E, Widjajakusuma E, Wöhnert J, Stock G, Schwalbe H. NMR and
MD studies of the temperature-dependent dynamics of RNA YNMG-tetraloops. Nucleic
acids research. 2008; 36(6):1928–1940.

Fleetwood O, Kasimova MA, Westerlund AM, Delemotte L. Extracting molecular insights
from conformational ensembles using Machine Learning. BioRxiv. 2019; p. 695254.

Giambaşu GM, York DM, Case DA. Structural fidelity and NMR relaxation analysis in a
prototype RNA hairpin. RNA. 2015; 21(5):963–974.

Graf J, Nguyen PH, Stock G, Schwalbe H. Structure and dynamics of the homologous
series of alanine peptides: a joint molecular dynamics/NMR study. Journal of the
American Chemical Society. 2007; 129(5):1179–1189.

Hall KB. Mighty tiny. RNA. 2015; 21(4):630–631.

Hartlmüller C, Günther JC, Wolter AC, Wöhnert J, Sattler M, Madl T. RNA structure
refinement using NMR solvent accessibility data. Sci Rep. 2017; 7(1):5393.

Hummer G, Köfinger J. Bayesian ensemble refinement by replica simulations and
reweighting. J Chem Phys. 2015; 143(24):12B634_1.

Kuhrova P, Banas P, Best RB, Sponer J, Otyepka M. Computer folding of RNA tetraloops?
Are we there yet? J Chem Theory Comput. 2013; 9(4):2115–2125.

Leontis NB, Westhof E. Geometric nomenclature and classification of RNA base pairs.
Rna. 2001; 7(4):499–512.

Marušič M, Schlagnitweit J, Petzold K. RNA dynamics by NMR. ChemBioChem. 2019; .

Nichols P, Born A, Henen M, Strotz D, Orts J, Olsson S, Güntert P, Chi C, Vögeli B. The exact
nuclear overhauser enhancement: recent advances. Molecules. 2017; 22(7):1176.

Nichols PJ, Born A, Henen MA, Strotz D, Celestine CN, Güntert P, Vögeli B. Extending the
Applicability of Exact Nuclear Overhauser Enhancements to Large Proteins and RNA.
ChemBioChem. 2018; 19(16):1695–1701.

Nichols PJ, Henen MA, Born A, Strotz D, Güntert P, Vögeli B. High-resolution small RNA structures from exact nuclear Overhauser enhancement measurements without additional restraints. Communications Biology. 2018; 1(1):61.

Nozinovic S, Fürtig B, Jonker HR, Richter C, Schwalbe H. High-resolution NMR structure of an RNA model system: the 14-mer cUUCGg tetraloop hairpin RNA. Nucleic Acids Res. 2010; 38(2):683–694.

Orioli S, Larsen AH, Bottaro S, Lindorff-Larsen K. How to learn from inconsistencies: Integrating molecular simulations with experimental data. arXiv preprint arXiv:190906780. 2019; .

Piccini G, Mendels D, Parrinello M. Metadynamics with discriminants: A tool for understanding chemistry. J Chem Theory Comput. 2018; 14(10):5040–5044.

Pitera JW, Chodera JD. On the use of experimental observations to bias simulated ensembles. Journal of chemical theory and computation. 2012; 8(10):3445–3451.

Reißer S, Zucchelli S, Gustincich S, Bussi G. Conformational ensembles of an RNA hairpin using molecular dynamics and sparse NMR data. Nucleic Acids Research. 2019; .

Salmon L, Giambasu GM, Nikolova EN, Petzold K, Bhattacharya A, Case DA, Al-Hashimi HM. Modulating RNA alignment using directional dynamic kinks: application in determining an atomic-resolution ensemble for a hairpin using NMR residual dipolar couplings. Journal of the American Chemical Society. 2015; 137(40):12954–12965.

Salmon L, Yang S, Al-Hashimi HM. Advances in the determination of nucleic acid conformational ensembles. Annu Rev Phys Chem. 2014; 65:293–316.

Sripakdeevong P, Cevec M, Chang AT, Erat MC, Ziegeler M, Zhao Q, Fox GE, Gao X, Kennedy SD, Kierzek R, et al. Structure determination of noncanonical RNA motifs guided by 1 H NMR chemical shifts. Nature Methods. 2014; 11(4):413.

Tan D, Piana S, Dirks RM, Shaw DE. RNA force field with accuracy comparable to state-of-the-art protein force fields. Proc Natl Acad Sci USA. 2018; p. 201713027.

Vögeli B. The nuclear Overhauser effect from a quantitative perspective. Prog Nucl Mag Res Sp. 2014; 78:1–46.

Wang L, Friesner RA, Berne B. Replica exchange with solute scaling: a more efficient version of replica exchange with solute tempering (REST2). J Phys Chem B. 2011; 115(30):9431–9438.

Woese C, Winker S, Gutell R. Architecture of ribosomal RNA: constraints on the sequence of" tetra-loops". Proc Natl Acad Sci USA. 1990; 87(21):8467–8471.

Wolters J. The nature of preferred hairpin structures in 16S-like rRNA variable regions. Nucleic acids research. 1992; 20(8):1843–1850.