Steady-state responses in the somatosensory system interact with endogenous beta activity

Michel J. Wälti[1,2*], Marc Bächinger[1,2], Kathy L. Ruddy[1,3], Nicole Wenderoth[1,2]

[1]Neural Control of Movement Lab, Department of Health Sciences and Technology, ETH Zurich,

Switzerland

[2]Neuroscience Center Zurich (ZNZ), University and ETH Zurich, Zurich, Switzerland

[3]Institute of Neuroscience, Trinity College Dublin, Ireland

* Correspondence:

Michel J. Wälti

michel.waelti@hest.ethz.ch

Keywords: steady-state evoked potentials, entrainment, somatosensory system, neural

oscillations, vibrotactile stimulation

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

Author Contributions

MJW, MB and KLR designed research; MJW performed research; MJW and MB analyzed data; MJW,

MB, KLR and NW wrote the paper.

Funding

This research was funded by the Swiss National Science Foundation (No. 320030_175616).

Acknowledgments

We thank Pearl Saldanha and Marionna Münger for their help with data acquisition, and the Statistical

Counseling Service of ETH Zurich, especially Janine Burren, for methodological support. We also

acknowledge the support of the Neuroscience Center Zurich (ZNZ).

Abstract

Brain oscillations have been related to many aspects of human behavior. To understand a potential

causal relationship, it is of great importance to develop methods for modulating ongoing neural activity.

It has been shown that external rhythmic stimulation leads to an oscillatory brain response that follows

the temporal structure of the presented stimulus and is assumed to reflect the synchronization of

ongoing neural oscillations with the stimulation rhythm. This interaction between individual brain

activity and so called steady-state evoked potentials (SSEPs) is the fundamental requirement of neural

entrainment. Here, we investigate whether neural responses to rhythmic vibrotactile stimulation,

measured with EEG, are dependent on ongoing individual brain oscillations, and therefore reflect

entrained oscillatory activity. For this, we measured phase synchronization in response to rhythmic

stimulation across various frequencies in the alpha and beta band. Three different stimulation

intensities were applied for each frequency relative to the individual sensory threshold. We found that

a higher stimulation intensity, compared to lower intensities, resulted in a more pronounced phase

synchronization with the stimulation signal. Moreover, EEG responses to low stimulation frequencies

closer to individual beta peak frequencies revealed a higher degree of entrainment, compared to

stimulation conditions with frequencies that were more distant to endogenous oscillations. Our findings

provide evidence that the efficacy of vibrotactile rhythmic beta stimulation to evoke a SSEPs is

dependent on ongoing brain oscillations.

Introduction

During cognitive tasks brain rhythms of different frequencies can be recorded at the scalp using

electroencephalography (EEG), that are correlated with cognitive and behavioral processes. In recent

years, it has been suggested that various non-invasive methods, such as repetitive transcranial magnetic

stimulation (rTMS) or transcranial alternating current stimulation (tACS) are capable of frequency-

specific modulation of brain rhythms. Hence, they have been used to test the causal relationship

between the strength of neuronal oscillations and behavior (for a review, see Thut, Schyns, & Gross,

2011). However, the interaction of these externally applied magnetic or electric fields with ongoing

brain rhythms is still poorly understood. An alternative form of frequency-specific modulation of brain

rhythms can be achieved with rhythmic sensory stimulation which induces steady-state evoked

potentials (SSEPs) following the temporal frequency of the driving stimulus (Regan, 1977). SSEPs are

usually measured with electroencephalography (EEG) and have been documented in the visual (steady-
state visually evoked potentials, SSVEPs), the auditory (auditory steady-state responses, ASSRs) and
the somatosensory (steady-state somatosensory evoked potentials, SSSEPs) systems (for a review, see
Vialatte, Maurice, Dauwels, & Cichocki, 2010). However, the underlying mechanism of SSEPs, is still
debated (see Zoefel, Ten Oever, & Sack, 2018). While some studies report evidence for entrainment
of ongoing brain oscillations by showing an interaction between stimulation and endogenous activity
(Notbohm, Kurths, & Herrmann, 2016; Schwab et al., 2006), others found responses to be nothing
more than regular repetitions of evoked neural potentials (Capilla, Pazo-Alvarez, Darriba, Campo, &
Gross, 2011; C. Keitel, Quigley, & Ruhnau, 2014). In contrast to simply overlaying evoked potentials,
entrainment modulates brain oscillations. To test whether brain responses to rhythmic sensory
stimulation reflect entrainment, it is therefore necessary to provide evidence of a dependency on
ongoing brain oscillations. This dependency arises from theoretical considerations and has been the
main research interest in studies supporting the entrainment hypothesis (see Zoefel et al., 2018). The
so-called Arnold tongue describes a classical model of entrainment between two oscillators depending
on stimulation rhythm and intensity (Pikovsky, Rosenblum, Zaks, & Kurths, 1999). The model predicts
that a high stimulation intensity evokes a larger response than a low intensity but also entrainment to
be most pronounced when the stimulation frequency matches the frequency of the endogenous system
(e.g. Fröhlich, 2015). In a recent study, Notbohm et al. (2016) reported that the coupling of EEG
oscillations with rhythmic visual flicker follows the characteristics of the Arnold tongue. They showed
that the EEG response to the stimulation depends on both the stimulation intensity and the distance
between the flicker frequency and the participants' individual alpha frequency (IAF; Notbohm et al.,
2016). Other studies found similar effects for tACS over occipital cortex which either increases EEG
power at participants' visual alpha frequency (Helfrich et al., 2014; Zaehle, Rach, & Herrmann, 2010),
or biases ongoing alpha oscillations in visual cortex towards the exogenous stimulation frequency
(Minami & Amano, 2017). Most previous studies have tested entrainment in visual cortex targeting
the alpha rhythm. Here we test whether it is also possible to entrain endogenous brain rhythms of
primary sensorimotor cortex via rhythmic tactile stimuli. It is well known that the sensorimotor cortex
exhibits strong activity in the alpha (8 – 12 Hz) and beta band (15 – 30 Hz) when subjects are at rest
(Crone et al., 1998; Fransen, Dimitriadis, van Ede, & Maris, 2016). During sensorimotor processing
and, particularly, during motor execution, both rhythms desynchronize (Bauer, Oostenveld, Peeters, &
Fries, 2006; Kilavik, Zaepffel, Brovelli, MacKay, & Riehle, 2013; Spitzer & Haegens, 2017) but re-
synchronize quickly once the movement is completed, implicating both rhythms in sensorimotor
processing (Kilavik et al., 2013; Pfurtscheller & Lopes da Silva, 1999; Ploner, Gross, Timmermann,

Pollok, & Schnitzler, 2006; Rajagovindan & Ding, 2011). Here we ask whether the alpha and the beta

rhythms in primary sensorimotor cortex are equally susceptible to entrainment via tactile stimuli. We

hypothesized that repetitive tactile stimulation at or close to the intrinsic alpha and beta frequency

peaks of the sensorimotor system should be more effective than stimulation at other frequencies. To

that end, we probed whether characteristics of the Arnold tongue can be found when measuring phase

synchronization in response to vibrotactile stimulation at different frequencies in the alpha and beta

bands. Three different stimulation intensities were applied for each frequency which were defined

relative to individual sensory threshold. We expected steady-state EEG responses to be more

pronounced for (i) high-intensity stimulations compared to low intensities, and (ii) at stimulation

frequencies closer to individual alpha (IAF) and/or beta frequencies (IBF) compared to frequencies

more distant from endogenous oscillations.

Materials and methods

Participants

25 participants took part in the experiment in exchange for monetary compensation (20 Swiss francs

per hour). The participants had no reports of psychiatric disorders and were all right-handed according

to the Edinburgh Handedness Inventory (Oldfield, 1971). The study protocol was approved by the local

ethics committee and was conducted in accordance with the Declaration of Helsinki. All participants

provided written consent. 3 participants were excluded from the analysis due to technical problems

during EEG recording or because they did not follow the instructions (e.g. falling asleep during the

EEG recording), resulting in a final sample of 22 participants (female: 8; age: range $= 18 - 29$, M $\pm$

SD $= 22.8 \pm 2.6$).

Design and procedure

The experiment was conducted in two parts. In a first part, we determined participants' IAF and IBF

in a simple finger tapping task. The second part consisted of the main experiment with different

conditions of vibrotactile stimulation to participants' right index finger. Throughout both parts, EEG

was recorded (see below).

Part I: Detection of IAF and IBF

Participants were comfortably seated in a dark sound-attenuated room, approximately 100 cm away
from a computer screen (27-inch) and were instructed to execute a self-paced finger tapping task with
their right hand, in which they tapped and gently squeezed their thumb against each of the other fingers
successively. One run consisted of an alternating green and red fixation cross appearing in the center
of the screen (each 9 times for 10 seconds). Green indicated that the participant should execute the
finger tapping, and red indicated that they should rest their hand. Each participant performed two runs.

Part II: Vibrotactile stimulation

During the main experiment, participants' right index finger was stimulated with a rhythmic
vibrotactile stimulation. For the stimulation, an earphone (YVE-01B-03, Yeil Electronics Co., South
Korea) with a diameter of 17 mm was taped to the participants finger. The earphone was modified to
produce vibrations via a coil actuator with a mass attached which vibrates when current is passed
through. The stimulation signals were generated in MATLAB (The MathWorks Inc., USA) and sent
to the stimulation devices via a National Instruments Card (NI-USB 6343) and an amplifier (Dancer
Design, St. Helens, UK). The stimulation was elicited by driving the stimulator with a carrier frequency
of 200 Hz which was frequency-modulated by a square wave corresponding to the stimulation
frequency (Breitwieser, Kaiser, Neuper, & Müller-Putz, 2012). This time course of the driving voltage
had a resolution of 0.1 ms. For each participant, the individual stimulation threshold was determined
prior to the main experiment by applying a 20 Hz stimulation stimulus which started at sub-threshold
intensity and increased in steps of 10% increments of output voltage until it was perceived. This
procedure was repeated three times. The mean values of the detection thresholds indicated the
individual stimulation threshold for each participant. Based on the individual stimulation threshold
three stimulation intensities were set: (i) sub-threshold intensity corresponding to 0.5 x threshold
intensity (sub-threshold), (ii) a low supra-threshold intensity corresponding to 20 x threshold intensity
(low), and (iii) a high supra-threshold intensity corresponding to 200 x threshold intensity (high).
During the main experiment (Figure 2.1), all stimulation frequencies (i.e. 6 – 24 Hz) were applied with
the same sub-threshold, low and high intensities, respectively. Note that the range of our experimental
frequencies was rather narrow such that all activate the same tactile mechanoreceptors (Meissner's
corpuscles, 5 – 50 Hz; see Johansson & Flanagan, 2009). After threshold detection, the main
experiment began which consisted of 3 blocks each consisting of 10 stimulation conditions which were
carried out in random order (resulting in 30 stimulation conditions in total). These 30 conditions were
composed of 10 different stimulation frequencies ranging from 6 to 24 Hz in steps of 2 Hz, each
administered with the three stimulation intensities (sub-threshold, low or high). Each stimulation lasted

for 40 seconds and was preceded by a 10 second period without stimulation. Participants were
instructed to sit as still as possible and direct their gaze at a white fixation cross centered on a black
screen.

Figure 1. Experimental procedure. Participants' right index finger was stimulated with frequencies ranging
from 6 to 24 Hz at 3 different stimulation intensities. Each stimulation lasted for 40 s, preceded by a 10 s period
without stimulation (rest).

Data acquisition

EEG data were acquired at 1000 Hz using a 64-channel Hydrocel Geodesic EEG System (Electrical
Geodesic Inc., USA), referenced to Cz (vertex), with an online Notch filter (50 Hz) and high-pass filter
at 0.3 Hz. Impedances were kept below 50 kΩ.

Preprocessing of EEG data

EEG data for both parts of the experiment were preprocessed and analyzed offline. Data were cleaned
(detection and interpolation of bad electrodes), band-pass filtered (5 – 25 Hz) and further processed
using independent component analysis (ICA). Artifact components (ICs) were automatically detected
and removed from the data using a custom built toolbox (see Liu, Ganzetti, Wenderoth, & Mantini,
2018).

EEG data analysis

Part I: IAF and IBF

To detect IAF and IBF, preprocessed data were epoched regarding the two conditions (movement and

rest; each from +1 s to +9 s after start of the condition) and analyzed using fast Fourier transform (FFT).

For each subject, frequency-band specific power differences in the alpha (8 – 12 Hz) and beta band (15

– 25 Hz) were calculated between movement and rest to detect event-related de-synchronization (ERD)

that typically appears during movement as well as event-related synchronization (ERS) immediately

after movement. Power changes were determined as percentage change from rest versus movement

(Pfurtscheller & Lopes da Silva, 1999):

173
$$\Delta\text{Power} = \frac{\text{power(rest)} - \text{power(movement)}}{\text{power(movement)}} * 100.$$

The electrode(s) exhibiting the strongest ΔPower over the contralateral hemisphere (right hand, left

hemisphere) for the alpha and beta band, respectively, was selected and we determined (i) IAF as the

frequency with the biggest power increase between 6 and 14 Hz (range of the stimulation frequencies);

and (ii) IBF as the frequency with the most pronounced power increase between 16 and 24 Hz. Similar

to before, we defined power increases as percentage change in power in the rest condition compared

to the move condition (ΔPower) and normalized the resulting power spectrum to the power at the alpha

or beta peak respectively.

Part II: Steady-state evoked potentials

In order to evaluate general effects of vibrotactile stimulation on EEG signals, we measured frequency-

specific power and phase-based consistency of EEG time-series during stimulation. For this,

preprocessed EEG data from the main experiment were epoched to each stimulation condition (+1 s to

+39 s relative to stimulation onset). Further, time-series data were cropped in 1 s segments without

overlap which were later treated as separate trials. In order to analyze steady-state responses to the

stimulation frequencies we pre-defined a region of interest (ROI) consisting of nine electrodes over the

somatosensory area. For each participant, we evaluated the electrode that showed the strongest SSEPs

across alpha (6 Hz, 8 Hz, 10 Hz, 12 Hz, 14 Hz) and beta (16 Hz, 18 Hz, 20 Hz, 22 Hz, 24 Hz)

stimulation conditions separately. For this, data averaged across trials of each high-intensity

stimulation condition were transformed into its frequency components using a fast Fourier transform

(FFT) and power at the corresponding stimulation frequency as well as neighboring frequencies (-2

Hz, -1Hz, +1Hz, +2Hz) were determined for each electrode within the ROI. We then calculated

percentage increase of power at the stimulation frequency compared to power at neighboring

frequencies for each electrode. To evaluate the electrode showing the strongest response across the

alpha stimulation conditions, we averaged the calculated power changes across the corresponding

conditions. The same procedure was carried out for the beta stimulation conditions. To determine

stimulation effects on frequency-specific power, FFT was performed on data that had been

preprocessed and averaged across trials, derived from these electrodes (strongest alpha and beta for

each participant). To determine frequency-specific phase consistency of the EEG signals, we further

calculated intertrial phase clustering (ITPC), which measures the extent of phase angle distributions at

each time-frequency point across trials (Cohen, 2014).

Part II: Phase-based synchronization between EEG and stimulation signal (entrainment)

To determine entrainment, we used data from the electrodes which were earlier defined as detecting

the strongest alpha and beta signals. We then measured changes in intersite phase clustering (ISPC)

between the preprocessed and averaged time-series data of each stimulation condition and the

corresponding stimulation signal. ISPC is a measure of phase-based connectivity between two time-

series signals (Cohen, 2014). Similar to ITPC, ISPC measures phase angle distributions, with the key

distinction that phase angle *differences* between two signals are measured. Similarly, to the EEG data,

the stimulation signals were band-pass filtered in the range from 5 to 25 Hz and down-sampled to 1000

Hz. The single-channel EEG time-series were further band-pass filtered in the range of ±2 Hz regarding

the corresponding stimulation frequency. To detect changes in phase-locked activity in response to the

stimulation, we compared ISPC between the stimulation signal and the EEG signal during stimulation

with ISPC prior to each condition (i.e. preceding rest period). Similar to the previously described

procedure, EEG data prior to stimulation onset (-9 s to -1 s) was cropped in 1 s segments and then

averaged across trials. ΔISPC was calculated by subtracting the mean pre-stimulation ISPC from the

mean ISPC during stimulation for each of the 30 conditions. All scripts to execute the experiment and

to preprocess and analyze EEG data were written in MATLAB (The MathWorks Inc., USA), and used

functions from EEGLAB (Delorme & Makeig, 2004). Scripts using FFT, ITPC and ISPC analysis were

adapted from M. X. Cohen (Cohen, 2014). For the experiment presentation code, PsychToolbox was

used (Brainard, 1997).

Statistical analysis

All statistical analyzes were performed separately for conditions targeting alpha oscillations

(stimulation frequencies: 6 Hz, 8 Hz, 10 Hz, 12 Hz, 14 Hz) and beta oscillations (stimulation

frequencies: 16 Hz, 18 Hz, 20 Hz, 22 Hz, 24 Hz). General effects of the stimulation signals on

frequency-specific power and ITPC were analyzed using repeated-measures ANOVA models in SPSS

Version 25 (IBM, USA). To test the hypothesized effects regarding phase-based connectivity between

stimulation and EEG signals, we used R (R Core Team, 2018) and the lme4 package (Bates, Mächler,

Bolker, & Walker, 2015) to perform linear mixed effects analyzes. We used linear mixed effects

models because they account for variance arising from individual differences (Baayen, Davidson, &

Bates, 2008). To ensure that observed effects are not driven by individual variability, subjects were

included in the models as random factors. Fixed effects represent the influence of experimental factors

on the dependent variable (here: ΔISPC). In our models, the fixed effects include stimulation intensity

(sub-threshold intensity as baseline) and distance of the stimulation frequency to individual

endogenous frequencies in the alpha (ΔIAF), or the beta (ΔIBF) band. Validation of our models was

done by plotting residuals against fitted values.

Results

239 The current study was designed to reveal variations in the effect of vibrotactile stimulation on the

240 individual neuronal response in sensorimotor cortex. Participants were rhythmically stimulated at their

241 right index finger (dominant hand) using three stimulation intensities and ten stimulation frequencies

242 across the alpha and beta bands. Each stimulation lasted for 40 seconds and was preceded by a 10

243 second period without any stimulation.

Detection of IAF and IBF

To detect IAF and IBF, frequency-band specific power changes due to event-related de-

/synchronization (ERD/ERS) in a simple finger tapping task were determined for each subject in the

alpha and beta bands. Figure 2.2 A/B show topographical plots and power spectra from an example

subject, revealing strong power increases in the left primary motor areas (rest vs. movement) and clear

peaks in the power spectra in both alpha and beta bands. Across subjects, average IAF was 11.0 Hz (±

2.1) and average IBF was 20.3 Hz (± 1.9) (see Figures 2.2 C/D). To evaluate whether the two spectral

peaks in alpha and beta present two independent rhythms, we calculated linear correlation with the

Pearson correlation coefficient between IAFs and IBFs across subjects. We found only a weak negative

correlation (r = -0.22, p = 0.319) and conclude that the power changes related to the finger tapping task

found in alpha and beta represent two distinct endogenous brain rhythms rather than harmonics.

Figure 2. Participants' IAF and IBF. Detection of participants' endogenous peak frequencies in the alpha (IAF) and beta (IBF) bands by calculating frequency-band specific power changes due to event-related de-/synchronization (ERD/ERS) in a simple finger tapping task. A: Example subject's topographical plot of ΔPower in the alpha band and power spectrum of the electrode showing the highest value (red dot). IAF was determined as frequency with the highest ΔPower (here: 9.875 Hz, marked with an asterisk). B: Same example subject's topographical plot and power spectrum as in A, but for the beta band (IBF: 22 Hz, marked with an asterisk). C: Ordered IAF frequencies across all subjects. D: Ordered IBF frequencies across all subjects. E: Depiction of correlation with regression line between IAF and IBF values. Pearson correlation coefficient reveals only a weak negative linear dependency (r = -0.22).

Stimulation intensity differences on steady-state evoked potentials in alpha and beta

In order to evaluate entrainment effects, we first need to determine whether our stimulation protocol reveals typical characteristics of steady-state evoked potentials over the somatosensory area across

alpha and beta frequencies. Further, we are interested in general differences between the two frequency

bands regarding effects of stimulation intensity. Therefore, a two-way repeated measures ANOVA

with factors frequency band (alpha and beta) and stimulation intensity (sub-threshold, low, high) was

used to detect general effects of different stimulation conditions on frequency-specific power and

phase-clustering (ITPC) values. Figures 2.3 and 2.4 depict stimulation effects averaged across all

subjects. Figure 2.3 shows changes in the power spectra across all stimulation conditions and Figure

2.4 reveals effects on ITPC. Our first analysis used power as dependent variable (derived from FFT).

The results from the ANOVA reveal a main effect of stimulation intensity ($F_{(2,20)} = 50.7$, $p < 0.001$),

but not of frequency band ($F_{(1,21)} = 0.7$, $p < 0.411$). Bonferroni adjusted post-hoc comparisons

between intensities revealed a significant difference between all three levels (sub-threshold vs. low: p

= 0.030; sub-threshold vs. high: $p < 0.001$; low vs. high: $p < 0.001$). Further, the interaction between

frequency band and stimulation intensity show a significant effect, meaning that vibrotactile

stimulations in alpha and beta have different effects on frequency-specific power across the different

intensities ($F_{(2,20)} = 3.7$, $p = 0.041$). This steeper increase in power across the stimulation intensities

at beta compared to alpha (see Figure 2.5 A) is in line with previous research suggesting that the

somatosensory system has a temporal resonance frequency in the beta band (Müller, Neuper, &

Pfurtscheller, 2001; Snyder, 1992; Tobimatsu, Zhang, & Kato, 1999), thus, responding with greater

amplitudes to stimulation frequencies in beta. The same approach was used to detect effects on ITPC

across the two frequency bands and three stimulation intensities. Again, we found a main effect of

stimulation intensity ($F_{(2,20)} = 24.8$, $p < 0.001$), but not of frequency band ($F_{(1,21)} = 1.0$, $p = 0.341$).

Post-hoc comparisons between intensities revealed a significant difference between high intensity

compared to the other two, but not between sub-threshold and low levels (sub-threshold vs. low: p =

0.293; sub-threshold vs. high: $p < 0.001$; low vs. high: $p < 0.001$). Further, no significant interaction

between frequency band and stimulation intensity on ITPC was observed ($F_{(2,20)} = 2.7$, $p = 0.093$).

We conclude that rhythmic vibrotactile stimulation has similar effects on alpha and beta band

frequencies regarding phase-angle distributions across trials (ITPC).

Figure 3. Power-based analysis of each stimulation condition averaged across subjects. A: Topographical plot of changes to electrode power across high-intensity alpha stimulations (red dots: ROI electrodes in the somatosensory area). B: Power spectra (FFT) of time-series of the selected electrode for stimulation frequencies in the alpha range. C: Closer depiction of power spectra at stimulation frequencies. D-F: Equivalent to A-C, across beta stimulation frequencies.

Figure 4. Phase-based analysis of each stimulation condition averaged across subjects. Time-series data from chosen electrodes (see Figure 2.3 A&D) were cropped in 1 s segments. A: Time-frequency plots of ITPC for stimulation frequencies delivered with high intensity in the alpha band with indication of the stimulation frequency (white transparent area). B: Time course of ITPC for each stimulation across alpha conditions. Dotted

lines represent mean ITPC for each condition over the time course of the 1 second segment (averaged across

trials). C&D: Equivalent to A&B, across beta stimulation frequencies.

Figure 5. Differences of vibrotactile stimulation on frequency-specific power and ITPC across both

frequency bands. A: Higher stimulation intensities reveal higher power values across the frequency bands. This

main effect of intensity seems to be similar for alpha and beta (no interaction effect of frequency band and

intensity). B: Also for ITPC we found a main effect of stimulation intensity. However, a significant interaction

effect between frequency band and intensity suggests that the intensity effect differs between alpha and beta.

ISPC in alpha and beta depend on stimulation intensity

To investigate entrainment effects of the stimulation signals, we computed changes to intersite phase

clustering (ΔISPC) for each condition and used linear mixed effects models for the statistical analysis.

Our model for the alpha band conditions included stimulation intensity and absolute distance from

stimulation frequency to IAF (ΔIAF) as fixed factors. Any variations in slope and intercept of the data

regarding individual variability was accounted for with the inclusion of subject as random variable. To

test for an effect of stimulation intensity across all stimulation frequencies, we used mean-shifted ΔIAF

values. Using the mean-shifted values allowed us to determine stimulation frequency-independent

differences across the three intensity levels by directly comparing the resulting intercepts of the factor

stimulation intensity. In line with our results from the power and ITPC analysis, we found the ΔISPC

between stimulation and EEG signal to be more pronounced with increasing stimulation intensity. A

post-hoc Tukey test revealed higher ISPC for the high intensity compared to sub-threshold and low

(sub-threshold vs. high: $p < 0.001$; low vs. high: $p = 0.004$). However, the increase in ΔISPC from sub-

threshold to low intensity showed no difference ($p = 0.936$). Testing the effect of stimulation intensity

in the beta band, revealed similar results as seen for alpha. Again, linear mixed effects models were

used with ΔISPC as dependent variable, stimulation intensity and mean-shifted distance from

stimulation frequency to IBF (ΔIBF) as fixed variables and subject as random variable. Positive and

statistically significant estimates for the post-hoc comparison between all intensity levels (sub-

threshold vs. low: $p < 0.001$; sub-threshold vs. high: $p < 0.001$; low vs. high: $p < 0.001$), confirm the

effect of stimulation intensity on steady-state neuronal response as hypothesized.

Phase-coupling in beta depends on the interaction of intensity and distance to IBF

In order to test the main requirement regarding entrainment effects, and therefore the most important

feature of the Arnold tongue, the dependency of stimulation effects on endogenous neuronal activity

was investigated. We started with the analysis of the data obtained from stimulation conditions

targeting alpha oscillations (stimulation frequencies from 6 to 14 Hz). The shape of the Arnold tongue

suggests a dependency of entrainment effects (here: ΔISPC) on the distance between stimulation

frequency to endogenous peak frequency (ΔIAF/IBF). In our statistical model this dependency can be

depicted by plotting ΔISPC values according to their ΔIAF/IBF (see Figure 2.6 A/B). Negative slopes

(derived for each stimulation intensity) which significantly differ from 0 can be expected, if the

hypothesized dependency is true (i.e. stimulation frequencies closer to IAF/IBF lead to higher ΔISPC

values). Although the slopes in the alpha conditions for low and high intensities show a negative

direction (see Figure 2.6 A), statistically, they were not significantly different from 0 (ΔIAF:low: $p =$

0.998; ΔIAF:high: $p = 0.915$). Our next analysis focused on the beta band conditions (stimulation

frequencies from 16 to 24 Hz). For beta, the slope of the low intensity differed significantly from 0 (p

$= 0.013$), while sub-threshold and high intensities showed no such effect (ΔIBF:sub-threshold: $p =$

0.998; ΔIBF:high: $p = 0.999$). Our results from the beta band reveal a relationship between endogenous

oscillations, stimulation intensity and phase coupling as would be assumed by the Arnold tongue. In

our low intensity conditions, the distance of the stimulation frequency to IBF plays a crucial role in

predicting coupling between stimulation and the underlying oscillation. The Arnold tongue widens by

increasing the stimulation intensity, inferring a decreased effect of ΔIBF. This is because the system

experiences such a strong enforcement that the ongoing oscillatory activity has no effect on ISPC.

Figure 6. Depiction of the effects of different stimulation intensities on changes in ISPC in relation to the distance to endogenous frequencies (IAF and IBF). A: Stimulation conditions in the alpha band reveal no difference from 0 for any of the slopes and therefore no evidence for entrainment. B: In the beta band, low stimulation intensity reveals a negative slope which differs significantly from 0. This dependency of entrainment effect (ΔISPC) is in line with the hypothesized shape of the Arnold tongue.

Discussion

In the present study, we investigated the effects of rhythmic vibrotactile stimulation across alpha and
beta frequencies on neuronal activity in the somatosensory cortex. Our main results reveal a clear effect
of stimulation intensity on frequency-specific power and intertrial phase clustering (ITPC) across both
frequency bands. A significant interaction effect revealed a difference between alpha and beta
stimulation frequencies with regard to their effect on frequency-specific power. The steeper power
increase across the three stimulation intensities in beta compared to alpha, as well as the higher power
values attained from high-intensity stimulation (Figure 2.5 A) is in line with previous findings
suggesting that the resonance frequency of the somatosensory system is in the beta band (Müller et al.,
2001; Snyder, 1992; Tobimatsu et al., 1999). Further, we provide evidence for the hypothesized

entrainment effect by revealing a dependency of phase-locked activity between stimulation signal and

endogenous beta brain oscillations on the interaction of stimulation intensity and frequency regarding

its distance to individual beta peaks. This dependency is in line with the hypothesized Arnold tongue

which predicts that entrainment via external signals of low intensity is stronger when the stimulation

frequency matches the frequency of the endogenous system (Fröhlich, 2015; Notbohm et al., 2016;

Pikovsky et al., 1999). Stimulation conditions in the alpha band revealed no such entrainment effect

although effects of stimulation intensity on ISPC were found as well.

Rhythmic sensory stimulation reveals online entrainment effects

Entrainment effects of sensory and non-invasive transcranial stimulation have been reported in various

studies (for reviews see: Thut et al., 2011; Zoefel et al., 2018), yet our study is the first to demonstrate

this effect for a wider range of endogenous brain oscillations in the somatosensory cortex. Our main

finding that external rhythmic stimulation interacts with ongoing endogenous brain rhythms of the

sensorimotor cortex is in line with previous reports using either sensory stimulation (Gulbinaite, van

Viegen, Wieling, Cohen, & VanRullen, 2017; Notbohm et al., 2016) or various forms of rhythmic,

non-invasive brain stimulation methods (Helfrich et al., 2014; Minami & Amano, 2017; Zaehle et al.,

2010). However, until now entrainment effects have been mainly studied by targeting the visual alpha

rhythm and its behavioral correlates. Previous studies revealed that steady-state evoked potentials

depend on endogenous brain oscillations (Gulbinaite et al., 2017; Notbohm et al., 2016) and that

modulating these endogenous rhythms of the visual cortex affects visual processing as measured at the

behavioral level (Cecere, Rees, & Romei, 2015; Ronconi, Busch, & Melcher, 2018; Ronconi &

Melcher, 2017). Together, these findings provide evidence for alpha entrainment effects in the visual

system and also demonstrate the underlying causal relationship between brain oscillations and human

behavior.

Few studies have investigated the modulation of neuronal oscillations in human somatosensory cortex.

Two recent studies used transcranial alternating current stimulation (tACS) of the mu-rhythm (i.e. the

alpha rhythm of sensorimotor cortex) to modulate sensorimotor perception threshold (Gundlach,

Müller, Nierhaus, Villringer, & Sehm, 2016) and to reveal entrainment effects specific to endogenous

mu-frequencies (Gundlach, Muller, Nierhaus, Villringer, & Sehm, 2017). In a first study, Gundlach et

al. (2016) applied tACS stimulation at individual mu-alpha frequencies and revealed phase-dependent

modulations of human somatosensory perception. However, in a tACS-EEG follow-up study,

investigating the electrophysiological correlates of their tACS stimulation on endogenous mu-alpha

oscillations, their findings reveal a counter-intuitive effect of a decrease of mu-alpha amplitude after

stimulation (Gundlach et al., 2017). While the authors argue that homeostatic neuroplastic processes,

as well as the electrode placement (bilateral) might have led to the decrease in mu-alpha, the underlying

neural mechanism of their first study remains unclear. Even though their approach is well-suited to

reveal functional modulations with tACS, it does not allow to investigate online entrainment effects

due to large electrical artifacts. Further, due to the lack of different stimulation intensities, as well as

variations in stimulation frequencies away from endogenous frequencies, the Arnold tongue cannot be

derived from these studies. In addition, only mu-alpha frequency stimulation was applied, preventing

conclusions in regard of other frequency bands. Here, we used tactile stimulation across various

frequencies covering alpha and beta frequencies, which was provided mechanically and, therefore, did

not interfere with the EEG recordings, thus, enabling the analysis of potential entrainment effects

during stimulation. Our results suggest that rhythmic activity in somatosensory cortex can be

influenced, but that potential entrainment effects were only found in the beta band.

Stronger response to rhythmic beta stimulation in the somatosensory system

Although studies investigating spontaneous (resting-state) EEG activity, found both beta (mu-beta) and

alpha (mu-alpha) oscillations in sensorimotor areas (Hillebrand, Barnes, Bosboom, Berendse, & Stam,

2012; A. Keitel & Gross, 2016; Ramkumar, Parkkonen, & Hyvarinen, 2014), the steady-state evoked

potentials in the two frequency bands differ clearly (see Figure 2.3). This finding is in line with previous

studies using rhythmic sensory stimulation, revealing that each sensory system responds maximally to

a specific stimulation frequency (for a review: Vialatte et al., 2010). Arguably, this represents a

mechanism by which each of the sensory systems processes information optimally (Hutcheon &

Yarom, 2000). Such a temporal tuning function appears in the visual system around 10 Hz (Regan,

1989), in the auditory system around 40 Hz (Stapells, Makeig, & Galambos, 1987) and in the

somatosensory system around 21 Hz (Müller et al., 2001; Snyder, 1992; Tobimatsu et al., 1999;

Tobimatsu, Zhang, Suga, & Kato, 2000). Such resonance mechanisms could be responsible for our

mixed results for alpha and beta stimulation frequencies.

True entrainment or rhythmically evoked potentials?

Our results are in line with previously stated dependencies of EEG responses to rhythmic stimulation

on endogenous brain oscillations. However, it is difficult to draw a final conclusion on whether steady-

state somatosensory evoked potentials (SSSEPs), as produced in our study with vibrotactile

stimulation, represent entrained neural oscillations. As Haegens and Zion-Golumbic (2018) point out

in a recent review, the interpretation of neuronal responses to rhythmic stimulation as neural

entrainment might lack the fundamental distinction from a rhythmic series of evoked potentials.

Although some studies have shown that non-rhythmic (= jittered) stimulations do not reveal a non-

linear response with dependency on endogenous oscillations (Kayser, Ince, Gross, & Kayser, 2015;

Notbohm et al., 2016), others found the opposite effect (Capilla et al., 2011; C. Keitel, Thut, & Gross,

2017; see also: Zoefel et al., 2018). Our findings can be seen in agreement with an entrainment

mechanism, as we find dependencies of SSSEPs on endogenous oscillations only with a low

stimulation intensity and only in beta. Arguably, if SSSEPs would in fact represent only regular

repetitions of evoked potentials, we should have seen the same effects for alpha and beta stimulation.

However, we do not evaluate our finding in comparison to non-rhythmic stimulation. Regarding the

ongoing debate on whether neural entrainment occurs in response to rhythmic sensory stimulation,

modulating functional correlates of neural oscillations (e.g. perception, memory) would serve as a

convincing outcome measure (Haegens & Zion Golumbic, 2018). Further, in the present study, we

used sensory data from selected electrodes that revealed strongest frequency-specific power changes

in a finger tapping task prior to the main experiment. Although our data reveal clear peaks in the alpha

and beta bands for most of the participants, our analyses have to be interpreted with caution. For

example, a decline in attention and alertness throughout the experiment could have led to variations in

IAF and/or IBF (Mierau, Klimesch, & Lefebvre, 2017), reducing potential entrainment effects as

analyzed here. Nevertheless, our findings provide valuable insight in the neural processing of rhythmic

sensory stimulation and should be considered in future studies using somatosensory entrainment.

Conclusion

The concept of modulating brain oscillations by rhythmic stimulation has become increasingly popular

in research on brain rhythms and their underlying functions (Thut et al., 2011). Despite the limitations

of the present study, we show that there is a significant frequency-specific dependency of brain

responses to rhythmic vibrotactile stimulation on individual neuronal activity in the beta band. We

conclude that our findings support the notion of entrainment of endogenous brain oscillations in

somatosensory cortex. This opens up possibilities to causally test interactions between oscillatory

activity and behavior by utilizing entrainment via somatosensory stimulation.

References

Baayen, R. H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed random effects for subjects and items. *Journal of Memory and Language, 59*(4), 390-412. doi:10.1016/j.jml.2007.12.005

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Usinglme4. *Journal of Statistical Software, 67*(1). doi:10.18637/jss.v067.i01

Bauer, M., Oostenveld, R., Peeters, M., & Fries, P. (2006). Tactile spatial attention enhances gamma-band activity in somatosensory cortex and reduces low-frequency activity in parieto-occipital areas. *J Neurosci, 26*(2), 490-501. doi:10.1523/JNEUROSCI.5228-04.2006

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision, 10*(4), 433-436.

Breitwieser, C., Kaiser, V., Neuper, C., & Müller-Putz, G. R. (2012). Stability and distribution of steady-state somatosensory evoked potentials elicited by vibro-tactile stimulation. *Med Biol Eng Comput, 50*(4), 347-357. doi:10.1007/s11517-012-0877-9

Capilla, A., Pazo-Alvarez, P., Darriba, A., Campo, P., & Gross, J. (2011). Steady-state visual evoked potentials can be explained by temporal superposition of transient event-related responses. *PLoS One, 6*(1), e14543. doi:10.1371/journal.pone.0014543

Cecere, R., Rees, G., & Romei, V. (2015). Individual differences in alpha frequency drive crossmodal illusory perception. *Curr Biol, 25*(2), 231-235. doi:10.1016/j.cub.2014.11.034

Cohen, M. X. (2014). *Analyzing neural time series data: theory and practice*: MIT press.

Crone, N. E., Miglioretti, D. L., Gordon, B., Sieracki, J. M., Wilson, M. T., Uematsu, S., & Lesser, R. P. (1998). Functional mapping of human sensorimotor cortex with electrocorticographic spectral analysis I. Alpha and beta event-related desynchronization. *Brain, 121*, 2271-2299.

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J Neurosci Methods, 134*(1), 9-21. doi:10.1016/j.jneumeth.2003.10.009

Fransen, A. M., Dimitriadis, G., van Ede, F., & Maris, E. (2016). Distinct alpha- and beta-band rhythms over rat somatosensory cortex with similar properties as in humans. *J Neurophysiol, 115*(6), 3030-3044. doi:10.1152/jn.00507.2015

Fröhlich, F. (2015). Experiments and models of cortical oscillations as a target for noninvasive brain stimulation. *Progress in Brain Research, 222*, 41-73. doi:10.1016/bs.pbr.2015.07.025

Gulbinaite, R., van Viegen, T., Wieling, M., Cohen, M. X., & VanRullen, R. (2017). Individual Alpha Peak Frequency Predicts 10 Hz Flicker Effects on Selective Attention. *J Neurosci, 37*(42), 10173-10184. doi:10.1523/JNEUROSCI.1163-17.2017

Gundlach, C., Muller, M. M., Nierhaus, T., Villringer, A., & Sehm, B. (2017). Modulation of Somatosensory Alpha Rhythm by Transcranial Alternating Current Stimulation at Mu-Frequency. *Front Hum Neurosci, 11*, 432. doi:10.3389/fnhum.2017.00432

Gundlach, C., Müller, M. M., Nierhaus, T., Villringer, A., & Sehm, B. (2016). Phasic Modulation of Human Somatosensory Perception by Transcranially Applied Oscillating Currents. *Brain Stimulation, 9*(5), 712-719. doi:10.1016/j.brs.2016.04.014

Haegens, S., & Zion Golumbic, E. (2018). Rhythmic facilitation of sensory processing: A critical review. *Neurosci Biobehav Rev, 86*, 150-165. doi:10.1016/j.neubiorev.2017.12.002

Helfrich, R. F., Schneider, T. R., Rach, S., Trautmann-Lengsfeld, S. A., Engel, A. K., & Herrmann, C. S. (2014). Entrainment of brain oscillations by transcranial alternating current stimulation. *Curr Biol, 24*(3), 333-339. doi:10.1016/j.cub.2013.12.041

Hillebrand, A., Barnes, G. R., Bosboom, J. L., Berendse, H. W., & Stam, C. J. (2012). Frequency-dependent functional connectivity within resting-state networks: an atlas-based MEG beamformer solution. *Neuroimage, 59*(4), 3909-3921. doi:10.1016/j.neuroimage.2011.11.005

Hutcheon, B., & Yarom, Y. (2000). Resonance, oscillation and the intrinsic frequency preferences of neurons. *Trends in Neuroscience, 23*, 216–222.

Johansson, R. S., & Flanagan, J. R. (2009). Coding and use of tactile signals from the fingertips in object manipulation tasks. *Nat Rev Neurosci, 10*(5), 345-359. doi:10.1038/nrn2621

Kayser, S. J., Ince, R. A., Gross, J., & Kayser, C. (2015). Irregular Speech Rate Dissociates Auditory Cortical Entrainment, Evoked Responses, and Frontal Alpha. *J Neurosci, 35*(44), 14691-14701. doi:10.1523/JNEUROSCI.2243-15.2015

Keitel, A., & Gross, J. (2016). Individual Human Brain Areas Can Be Identified from Their Characteristic Spectral Activation Fingerprints. *PLoS Biol, 14*(6), e1002498. doi:10.1371/journal.pbio.1002498

Keitel, C., Quigley, C., & Ruhnau, P. (2014). Stimulus-driven brain oscillations in the alpha range: entrainment of intrinsic rhythms or frequency-following response? *J Neurosci, 34*(31), 10137-10140. doi:10.1523/JNEUROSCI.1904-14.2014

Keitel, C., Thut, G., & Gross, J. (2017). Visual cortex responses reflect temporal structure of continuous quasi-rhythmic sensory stimulation. *Neuroimage, 146*, 58-70. doi:10.1016/j.neuroimage.2016.11.043

Kilavik, B. E., Zaepffel, M., Brovelli, A., MacKay, W. A., & Riehle, A. (2013). The ups and downs of beta oscillations in sensorimotor cortex. *Exp Neurol, 245*, 15-26. doi:10.1016/j.expneurol.2012.09.014

Liu, Q., Ganzetti, M., Wenderoth, N., & Mantini, D. (2018). Detecting Large-Scale Brain Networks Using EEG: Impact of Electrode Density, Head Modeling and Source Localization. *Front Neuroinform, 12*, 4. doi:10.3389/fninf.2018.00004

Mierau, A., Klimesch, W., & Lefebvre, J. (2017). State-dependent alpha peak frequency shifts: Experimental evidence, potential mechanisms and functional implications. *Neuroscience, 360*, 146-154. doi:10.1016/j.neuroscience.2017.07.037

Minami, S., & Amano, K. (2017). Illusory Jitter Perceived at the Frequency of Alpha Oscillations. *Curr Biol, 27*(15), 2344-2351 e2344. doi:10.1016/j.cub.2017.06.033

Müller, G. R., Neuper, C., & Pfurtscheller, G. (2001). „Resonance-like" frequencies of sensorimotor areas evoked by repetitive tactile stimulation. *Biomedical Engineering, 46*(7-8), 186-190.

Notbohm, A., Kurths, J., & Herrmann, C. S. (2016). Modification of Brain Oscillations via Rhythmic Light Stimulation Provides Evidence for Entrainment but Not for Superposition of Event-Related Responses. *Front Hum Neurosci, 10*, 10. doi:10.3389/fnhum.2016.00010

Oldfield, R. C. (1971). The Assessment and Analysis of Handedness: The Edinburgh Inventory. *Neuropsychologia, 9*, 97-113.

Pfurtscheller, G., & Lopes da Silva, F. H. (1999). Event-related EEG/MEG synchronization and desynchronization: basic principles. *Clinical Neurophysiology, 110*, 1842-1857.

Pikovsky, A. S., Rosenblum, M. G., Zaks, M. A., & Kurths, J. (1999). Phase Synchronization of
Regular and Chaotic oscillators. In H. G. Schuster (Ed.), *Handbook of Chaos Control*.
Weinheim: Wiley-VCH.

Ploner, M., Gross, J., Timmermann, L., Pollok, B., & Schnitzler, A. (2006). Oscillatory activity reflects
the excitability of the human somatosensory system. *Neuroimage, 32*(3), 1231-1236.
doi:10.1016/j.neuroimage.2006.06.004

R Core Team. (2018). R: A language and environment for statistical computing. Vienna, Austria.
Retrieved from https://www.r-project.org/

Rajagovindan, R., & Ding, M. (2011). From Prestimulus Alpha Oscillation to Visual-evoked
Response: An Inverted-U Function and Its Attentional Modulation. *Journal of Cognitive
Neuroscience, 23*(6), 1379-1394.

Ramkumar, P., Parkkonen, L., & Hyvarinen, A. (2014). Group-level spatial independent component
analysis of Fourier envelopes of resting-state MEG data. *Neuroimage, 86*, 480-491.
doi:10.1016/j.neuroimage.2013.10.032

Regan, D. (1977). Steady-state evoked potentials. *JOSA, 67*(11), 1475-1489.

Regan, D. (1989). *Human brain electrophysiology: evoked poten- tials and evoked magnetic fields in
science and medicine*. New York: Elsevier.

Ronconi, L., Busch, N. A., & Melcher, D. (2018). Alpha-band sensory entrainment alters the duration
of temporal windows in visual perception. *Sci Rep, 8*(1), 11810. doi:10.1038/s41598-018-
29671-5

Ronconi, L., & Melcher, D. (2017). The Role of Oscillatory Phase in Determining the Temporal
Organization of Perception: Evidence from Sensory Entrainment. *J Neurosci, 37*(44), 10636-
10644. doi:10.1523/JNEUROSCI.1704-17.2017

Schwab, K., Ligges, C., Jungmann, T., Hilgenfeld, B., Haueisen, J., & Witte, H. (2006). Alpha
entrainment in human electroencephalogram and magnetoencephalogram recordings.
Neuroreport, 17(17), 1829-1833.

Snyder, A. Z. (1992). Steady-state vibration evoked potentials: description of technique and
characterization of responses. *Electroencephalography and Clinical Neurophysiology/Evoked
potentials Section, 84*(3), 257-268.

Spitzer, B., & Haegens, S. (2017). Beyond the Status Quo: A Role for Beta Oscillations in Endogenous
Content (Re)Activation. *eNeuro, 4*(4). doi:10.1523/ENEURO.0170-17.2017

Stapells, D. R., Makeig, S., & Galambos, R. (1987). Auditory steady-state responses: threshold
prediction using phase coherence. *Electroencephalography and Clinical Neurophysiology,
67*(3), 260-270.

Thut, G., Schyns, P. G., & Gross, J. (2011). Entrainment of perceptually relevant brain oscillations by
non-invasive rhythmic stimulation of the human brain. *Front Psychol, 2*, 170.
doi:10.3389/fpsyg.2011.00170

Tobimatsu, S., Zhang, Y. M., & Kato, M. (1999). Steady-state vibration somatosensory evoked
potentials: physiological characteristics and tuning function. *Clinical Neurophysiology,
110*(11), 1953-1958.

Tobimatsu, S., Zhang, Y. M., Suga, R., & Kato, M. (2000). Differential temporal coding of the
vibratory sense in the hand and foot in man. *Clinical Neurophysiology, 111*(3), 398-404.

Vialatte, F. B., Maurice, M., Dauwels, J., & Cichocki, A. (2010). Steady-state visually evoked
potentials: focus on essential paradigms and future perspectives. *Prog Neurobiol, 90*(4), 418-
438. doi:10.1016/j.pneurobio.2009.11.005

Zaehle, T., Rach, S., & Herrmann, C. S. (2010). Transcranial alternating current stimulation enhances
individual alpha activity in human EEG. *PLoS One, 5*(11), e13766.
doi:10.1371/journal.pone.0013766

Zoefel, B., Ten Oever, S., & Sack, A. T. (2018). The Involvement of Endogenous Neural Oscillations
in the Processing of Rhythmic Input: More Than a Regular Repetition of Evoked Neural
Responses. *Front Neurosci, 12*, 95. doi:10.3389/fnins.2018.00095