High-throughput transposon sequencing highlights the cell wall as an important barrier for salt stress in methicillin resistant *Staphylococcus aureus*

Short Title: The global salt stress response in S. aureus

Christopher F. Schuster[a], David M. Wiedemann[a], Freja C. M. Kirsebom[a,1], Marina Santiago[b],

Suzanne Walker[b] and Angelika Gründling[a,2]

ORCID IDs: 0000-0002-4566-7226 (CFS), 0000-0001-6961-0980 (DMW), 0000-0003-4585-319X (FCK), 0000-0002-5446-3452 (MS), 0000-0002-0545-914X (SW), 0000-0002-6235-8687 (AG)

[a]Section of Molecular Microbiology and MRC Centre for Molecular Bacteriology and Infection, Imperial College London, London, SW7 2AZ, U.K.

[b]Department of Microbiology and Immunobiology, Harvard Medical School, Boston, MA 02115, U.S.A.

[1]Current address: Section of Respiratory Infections, National Heart and Lung Institute, Imperial College London, London, W2 1PG, U.K.

[2]To whom correspondence should be addressed. Email: a.grundling@imperial.ac.uk, Phone: 0044 20 7594 5256

Keywords: osmotic stress, TN-seq, Tradis, NaCl, *S. aureus*, DUF25380, Pbp2

Summary

Staphylococcus aureus is an opportunistic pathogen that can cause soft tissue infections but is

also a frequent cause of foodborne illnesses. One contributing factor for this food association is its

high salt tolerance allowing this organism to survive the commonly used food preservation method

of salting. How this resistance is mediated is poorly understood, particularly during long-term

exposure. In this study, we used TN-seq based screens and identified a number of previously

uncharacterized genes that are involved in the salt stress response. Amongst these, we identified

the DUF2538 domain containing gene *SAUSA300_0957* (gene *957*) as essential under salt stress.

Interestingly, a *957* mutant was less susceptible to oxacillin and showed increased peptidoglycan

crosslinking. The salt sensitivity phenotype could be suppressed by amino acid substitutions in the

transglycosylase domain of the penicillin binding protein Pbp2, and these changes restored the

peptidoglycan crosslinking to WT levels. These results indicate that increased crosslinking of the

peptidoglycan polymer can be detrimental and highlight a critical role of the bacterial cell wall for

osmotic stress resistance. This study will serve as a starting point for future research on osmotic

stress response and help develop better strategies to tackle foodborne staphylococcal infections.

Introduction

Staphylococcus aureus is a Gram-positive bacterium that is carried by about 20 % of healthy

individuals (Wertheim *et al.*, 2005). It is also an opportunistic pathogen causing a variety of diseases,

including bacteremia, endocarditis, soft tissue infections and food poisoning (Tong *et al.*, 2015). One

characteristic of *S. aureus* is its ability to grow in the presence of a very high salt concentration (up

to 2.5 M). Many other bacteria are unable to grow under these conditions (Measures, 1975), and

this has been used extensively for the isolation of staphylococci. This salt tolerance also allows *S.*

aureus to thrive in its niches such as the human nares, skin (Wertheim *et al.*, 2005) and certain food

products (Adams & Moss, 2008).

Bacterial cells are highly pressurized (up to 1.9 MPa for *Bacillus subtilis* (Whatmore & Reed,

1990)) and contain a strongly crowded cytoplasm that is hyperosmotic to the environment. Water

is not transported into the cell actively but rather influxes passively into the cell due to the high

internal osmolality. Whenever the extracellular osmolality increases, water exits the cell which can

cause problems to multiple cellular processes, in part due to molecular crowding (Wood, 2011, van

den Berg *et al.*, 2017). In the case of salt, an increase in the intracellular sodium concentration can

lead to mis- and unfolding of proteins and inhibit enzymatic reactions due to the strong ionic

interactions produced. Hence, *S. aureus* must possess strategies to counter the effects of salt

damage, but these are poorly understood, as most work has been performed in *Escherichia coli* and

B. subtilis, which possess much lower salt tolerances than Staphylococci.

General mechanisms of hyper-osmotic stress mitigation in bacteria are the rapid import of

potassium and the accumulation of small osmotically active compounds, named compatible solutes

(reviewed in depth by (Bremer & Krämer, 2019, Gunde-Cimerman *et al.*, 2018, Booth, 2014, Cox *et

al.*, 2018)). Potassium levels quickly subside in favor of an increase in the concentration of

compatible solutes.

In *S. aureus,* results consistent with the uptake potassium (Csonka, 1989, Csonka & Hanson,

1991, Sleator & Hill, 2002) followed by the accumulation of compatible solutes have been reported

(Measures, 1975, Graham & Wilkinson, 1992, Christian & Waltho, 1964). The most effective

compatible solutes in *S. aureus* are glycine betaine and proline (Miller *et al.*, 1991, Measures, 1975,

Townsend & Wilkinson, 1992, Anderson & Witter, 1982, Bae & Miller, 1992, Pourkomailian & Booth,

1992), which are imported by specific osmolyte transporters (Bae & Miller, 1992, Pourkomailian &

Booth, 1992). These preferred osmolytes can accumulate in the cell to very high concentrations

without negatively affecting cellular processes (Csonka & Hanson, 1991). In addition, *S. aureus* can

also synthesize compatible solutes *de novo*, as in the case of glutamine (Anderson & Witter, 1982),

but this process is much slower than the uptake of osmolytes and requires more energy.

Previous research investigating the underlying genetic factors of the *S. aureus* osmotic stress

response implicated several osmolyte transporters in this process (Price-Whelan *et al.*, 2013,

Scybert *et al.*, 2003, Schwan *et al.*, 2006, Vijaranakul *et al.*, 1998, Wengender & Miller, 1995). These

include the potassium transport systems Kdp and Ktr (Price-Whelan *et al.*, 2013), the proline

transporter PutP (Wengender & Miller, 1995, Schwan *et al.*, 2006), the arsenic transport system Ars

(Scybert *et al.*, 2003) and the branched chain amino acid uptake system BrnQ (Vijaranakul *et al.*,

1998). In addition, although currently not experimentally verified in *S. aureus*, it can be assumed

that the main glycine betaine uptake system OpuD (Wetzel *et al.*, 2011, Zeden *et al.*, 2018) is an

important factor in *S. aureus* for osmotic adaptation as glycine betaine uptake reduces sensitivity to

NaCl exposure (Miller *et al.*, 1991). Of note, in recent time the second messenger cyclic-

dinucleotide-AMP (c-di-AMP) (Corrigan & Gründling, 2013) has been tied to the regulation of

osmotic stress in *S. aureus* and other Gram-positive bacteria (reviewed by Commichau *et al.*, 2018).

This is through the interaction and regulation of a number of potassium (Corrigan *et al.*, 2013) and

compatible solute transporters by c-di-AMP (Schuster *et al.*, 2016).

Despite these multiple countermeasures, exposure of *S. aureus* to osmotic stress leads to

morphological changes, including the formation of larger cells (Vijaranakul *et al.*, 1995) and a

thickening of the cell wall (Onyango *et al.*, 2013). However, if and which genetic factors are required

for those processes are currently not well understood. The cell wall of Gram-positive bacteria is an

important barrier and acts as a counterpart to the pressurized cytoplasm. A major component of

the cell wall is peptidoglycan, a polymer consisting of chains of repeating N-acetylglucosamine

(GlcNAc) and N-acetylmuramic acid (MurNAc) units that are crosslinked with neighboring chains via

short peptides and, in the case of *S. aureus,* pentaglycine cross-bridges (Reviewed by (Vollmer *et al.*,

2008)). In the past, the contribution of the cell wall towards osmotic stress resistance was often not

considered in the bacterial osmotic stress resistance field. Partly, because the integrity of the cell

wall is often assumed to be a prerequisite rather than an active adaptation to survive osmotic

stresses and partly because most cell envelope stress sensing systems are unable to directly sense

osmotic stresses (Jordan *et al.*, 2008).

Nevertheless, several possible changes to the cell wall have been observed *in S. aureus* during

NaCl stress, such as shortened interpeptide bridges in the peptidoglycan (Vijaranakul *et al.*, 1995),

an increase in resistance to methicillin (Madiraju *et al.*, 1987) and increased autolysis activity

(Stapleton *et al.*, 2007, Ochiai, 2000). In addition to peptidoglycan, other components within the cell

envelope, such as teichoic acids and more specifically their modifications with D-alanine (Koprivnjak

et al., 2006) and sugar residues (Mistretta *et al.*, 2019, Kho & Meredith, 2018) are affected by

osmotic stress. Combined, these findings indicate that the integrity of the cell wall is an important

requirement for osmotic resistance.

Using a transposon insertion sequencing (TN-seq) method (Santiago *et al.*, 2015), we

determined on a whole genome level genes that are essential or dispensable during long-term salt

exposure. Amongst other novel candidate genes, we identified gene *SAUSA300_0957* as an

important salt resistance gene in *S. aureus* and show that the encoded protein is important for

peptidoglycan homeostasis in *S. aureus*. Another protein, which we identified as important during

salt stress, is the penicillin binding protein Pbp2, further emphasizing a key function of

peptidoglycan during NaCl-induced osmotic stress. Taken together, these experiments highlight the

bacterial cell wall as a key player in the salt stress response of *S. aureus*.

Results

TN-seq is a robust method to identify *S. aureus* genes involved in NaCl stress

To identify *S. aureus* genes that are needed during salt stress, we conducted transposon

sequencing (TN-seq) screens in the presence and absence of salt (Fig. 1A). These experiments were

repeated in two independent TN-seq screens using a previously published library (Santiago *et al.*,

2015, Coe *et al.*, 2019) with six transposon variants (Replicate A) and a library containing only one

transposon variant (Replicate B) (Supplementary Table S1 for overview). The libraries were

propagated for 17 generations in either LBN or LB medium with 0.5 M NaCl and the expected growth

reduction was observed in the presence of 0.5 M NaCl (Fig. 1B). The Tn insertions were found to be

distributed evenly throughout the genome in the libraries grown in LBN medium. When the libraries

were grown in high salt medium, clustering of transposon insertions in certain regions was observed

due to the enrichment of better salt-adapted TN-strains (Supplementary Fig. S1). The fold-changes

of transposon insertions per gene under the salt stress conditions compared to the LBN condition

(input library) as well as q-values were determined for each gene (Supplementary Table S2-S3). The

data were plotted as volcano plots (Fig. 1C), revealing similar numbers of conditional salt essential

and dispensable genes in both replicates. Furthermore, a good overlap of the top essential and

dispensable genes was observed between the replicates confirming the robustness of the TN-Seq

screens (Supplementary Tables S3-7).

The TN-seq screens highlight the essentiality of transporters and cell envelope related genes and

the dispensability of respiration genes under high salt conditions

In both replicates, genes coding for transporters and cell envelope related genes were the most

prominent conditionally salt essential genes (Supplementary Tables S4-S7). More specifically, the

TN-seq data indicated that inactivation of the putative magnesium transporter MgtE (Schuster *et*

al., 2019) and the putative D-serine/D-alanine/glycine transporter AapA is detrimental under salt

stress (Supplementary Tables S4 and S6). Inactivation of AapA has previously been shown to

suppress the temperature sensitivity phenotype seen in an *S. aureus fmtC/mprF* mutant (Ishibashi

et al., 2007) and to lead to increased amoxacillin and daptomycin sensitivity (Mlynek *et al.*, 2016,

Coe *et al.*, 2019). In addition, the *dlt* operon, coding for enzymes required for the D-alanylation of

teichoic acids, an important factor for cationic peptide resistance (Brown *et al.*, 2013), as well as the

stationary and stress sigma factor gene *sigB* were found to be essential.

An even larger number of genes was identified as dispensable during salt stress (Supplementary

Tables S5 and S7). Inactivating or reducing the expression of such genes is expected to help *S. aureus*

survive under salt stress. Among these dispensable genes were genes coding for the sodium

antiporter Mnh2, which has previously been shown to transport Na^+/H^+ and K^+/H^+ in high pH

medium and to play an important role in cytoplasmic pH maintenance (Vaish *et al.*, 2018). Strikingly,

a large number of genes involved in respiration, such as the *cta/qox/men/hem* genes as well as

genes involved indirectly in respiration through quinone synthesis (shikimate pathway, *aro* operon)

were also identified as dispensable during salt stress. The data also reproducibly indicated that

genes coding for the phosphodiesterases GdpP and Pde2 involved in degradation of the signaling

molecule c-di-AMP are dispensable under high salt conditions. Using the TN-seq approach, we were

able to identify a large number of *S. aureus* genes as potentially dispensable or essential when this

organism is exposed to high salt.

Tn-seq reveals previously unidentified salt resistance genes during long-term salt exposure

The TN-seq method relies on the enrichment of strains adapted to certain growth conditions,

and is therefore ideal for exploring the long-term effects on growth in response to salt stress. In a

previous study (Price-Whelan *et al.*, 2013), the adaptation of *S. aureus* to 2 M NaCl stress after 6

generations of growth was determined on a transcriptome level. This work highlighted the

importance of K$^+$ transporters for *S. aureus* to survive the high salt exposure. To evaluate how our results compare, we determined the overlaps between our datasets and the microarray data. To this end, the *S. aureus* strain COL locus tags used in the microarray study were converted to TCH1516 locus tags where possible (Supplementary Table S8) and compared to the two TN-seq datasets. The overlap between the downregulated (microarray) and dispensable (TN-seq) genes was minimal in both replicates (1 and 7) and also very small for the upregulated and essential genes (both 14). Three operons were consistently essential or upregulated between all datasets: the *cap5*, *SAUSA300_0771-2*, and the *sda* operon. The *cap5* genes are involved in capsule production (Chan *et al.*, 2014) and this type of capsule has been previously reported to be induced by the addition of salt (Pöhlmann-Dietze *et al.*, 2000). Genes *SAUSA300_0771-2* are hypothetical membrane proteins and presumably part of a threonine/serine exporter (Interpro database IDs: IPR010619 and IPR024528) and the *sda* genes are annotated as L-serine dehydratase components and a regulator protein. The low overlap between the microarray and TN-seq data (Supplementary Table S8) suggests that in our screen we have identified a number of previously unknown salt tolerance genes.

Confirmation of genes essential during salt stress using defined mutants

To determine to what extent different genes identified in the TN-seq screen contribute to the salt resistance of *S. aureus*, a set of potentially salt essential genes from dataset B were chosen for further investigation (Supplementary Table S9). For this analysis we used defined transposon mutants available from the *S. aureus* NTML transposon mutant library (Fey *et al.*, 2013). The growth of the different NTML transposon mutant strains was assessed in LBN and LB 0.5 M NaCl medium. In LBN medium, most strains grew similarly to the WT strain JE2, indicating that inactivation of the respective gene does not greatly affect bacterial growth in normal LB medium (Supplementary Figs. S2A-D, left panels). Exceptions were strains NE1109 (*sigB*), NE1778 (*lcpB*) and NE188 (*mfd*), cultures of which reached slightly lower final optical densities (Supplementary Figs. S2A-D, left panels). At

high osmolality conditions (LB 0.5 M NaCl), all strains showed a growth defect compared to the WT

strain (Supplementary Figs. S2A-D, right panels), validating the TN-seq as a method to identify *S.*

aureus genes that are important during salt stress. Several strains exhibited strong growth defects

in the presence of 0.5 M NaCl with strain NE1384, containing a transposon insertion in

SAUSA300_0957 showing extremely reduced growth (Supplementary Figs. S2C and D). To confirm

that the salt sensitivity was mediated by the inactivation of the genes in questions, we constructed

complementation plasmids for seven of the most promising candidates by either expressing the

gene of interest from its native promoter or an anhydrotetracycline (Atet) inducible promoter in the

transposon mutant strains. While no complementation was observed for four strains

(Supplementary Figs. S3A and S3B), the salt-dependent growth defect could be complemented for

strains carrying mutations in *SAUSA300_0694* (Fig. 2A), *SAUSA300_0910* (*mgtE*) and

SAUSA300_0957 (Fig. 2B). *SAUSA300_0694* encodes a hypothetical protein with 6 predicted

transmembrane helices but no other identifiable domain motif. MgtE (SAUSA300_0910) is a

predicted magnesium transporter, which we have recently shown can also contribute to cobalt

toxicity in *S. aureus* (Schuster *et al.*, 2019). *SAUSA300_0957* (from here on out referred to as gene

957) codes for a cytoplasmic protein with a DUF2538 domain of unknown function that is conserved

in Actinobacteria and Firmicutes. We further investigated this gene in this study because the *957*

mutant exhibited the strongest salt-sensitivity phenotype (Fig. 2B).

Deletion of gene *957* causes physiological changes leading to a smaller cell size

As all previous experiments were performed with transposon mutants, we first constructed

strain LAC*Δ*957* with a marker-less in-frame deletion of gene *957*. Strain LAC*Δ*957* had the

expected growth defect in the presence of 0.5 M NaCl and this phenotype could be complemented

by expressing a functional copy of *957* (Fig. 3A). To gain further insight into the molecular function

of gene *957*, we analyzed the neighboring genomic area. Transcriptome data (Mäder *et al.*, 2016)

suggested that gene *957* is in an operon with *lcpB* (Fig. 3B), which is one of three wall teichoic acid

(WTA) ligases present in *S. aureus* (Chan *et al.*, 2013, Over *et al.*, 2011, Schaefer *et al.*, 2017). *fmtA*

is located upstream of *957* and has been proposed to code for an esterase that can remove D-

alanine modifications from teichoic acids (Rahman *et al.*, 2016) and is involved in methicillin

resistance (Berger-Bächi *et al.*, 1989). Genes coding for a predicted acetyltransferase and *atl* coding

for the major *S. aureus* autolysin are found immediately downstream of the *957* operon (Oshida *et*

al., 1995). Due to the presence of many cell envelope related genes in the vicinity of gene *957*, we

proceeded to test if the *957* mutant displays other phenotypes connected to cell envelope

homeostasis. First, WT LAC*, the *957* mutant and the complementation strain were grown in LBN

and the ratio of colony forming units to optical density (CFU/OD) was determined. The CFU/OD ratio

was significantly larger for the *957* mutant ($60.8\pm6.8 \times 10^7$ CFU/OD) mutant as compared to strain

LAC* ($37.5\pm9.3 \times 10^7$ CFU/OD) when grown in LBN, suggesting a change in cell shape or size (Fig. 3C).

This phenotype was restored in the complementation strain ($37.3\pm3.6 \times 10^7$ CFU/OD) (Fig. 3C). When

the average cell diameter was determined by microscopy, a small but significant reduction in cell

diameter was seen for the *957* mutant (1.25 ± 0.02 µm) as compared to the WT (1.36 ± 0.03 µm) and

the cell size was restored to wild type levels in the complementation strain (1.35 ± 0.05 nm) (Fig. 3D).

These results support our hypothesis that the envelope might be a target of *gene 957* and prompted

us to look into these processes in more detail.

Salt sensitivity of *957* is not due to interaction with LcpB but likely due to changes in peptidoglycan

crosslinking

Since the gene *957* is on the same transcript as *lcpB* coding for a WTA ligase (Fig. 3A), the

contribution of WTA to salt stress resistance as well as the contribution of *957* to the attachment of

WTA to the cell wall was assessed. Initially the growth of a *tagO* (*tarO/llm*) mutant, a strain unable

to produce WTA, was tested in LB and LB 0.5 M NaCl medium. When cells were grown in LBN, they

grew normally (Supplementary Fig. S4A), but under high salt conditions, the *tagO* mutant showed reduced growth, similar to that of the *957* mutant (Fig. 4A). This is consistent with what has been reported previously for a *Staphylococcus epidermidis tagO* mutant (Holland *et al.*, 2011) and indicates an important role for WTA during salt stress. When the *lcpB* mutant was propagated in LBN medium, its growth was slightly inhibited (Supplementary Fig. S4A). However, in high salt medium (Fig. 4A) it grew unimpeded and like WT. Since the *lcpB* mutant did not show a growth defect in high salt medium, this made it unlikely that *957* is a regulator of LcpB activity, and hence is involved in the attachment of WTA to the peptidoglycan. To specifically investigate the requirement of 957 in WTA attachment, we isolated WTA from strains grown in either LBN or LB containing 0.4 M NaCl and separated the polymer on native polyacrylamide gels. The slightly lower concentration of salt was chosen because growth of the *957* and the *tagO* mutants was too strongly inhibited at 0.5 M NaCl. As expected the *tagO* mutant did not produce any WTA (Supplementary Fig. S4B, Fig. 4B) and consistent with previous studies (Over *et al.*, 2011, Chan *et al.*, 2013), the *lcpB* mutant exhibited slightly reduced WTA levels compared to the other strains (Supplementary Fig. S4B, Fig. 4B). No reduction in WTA was observable in the *957* mutant compared to the WT strain in either LBN (Supplementary Fig. S4B & S4C) or LB 0.4 M NaCl (Fig. 4B & 4C) but rather a small increase, which could be complemented. Taken together, these data indicate that 957 neither directly nor indirectly (e.g. through regulating the activity of LcpB) plays a major role in WTA attachment.

Gene *957* is found upstream of the bi-functional autolysin gene *atl* and therefore we next investigated potential changes in autolytic activity using zymograms. A small increase in autolytic activity was detected in the *957* mutant (Fig. 4D) for bands around 55 kDa and 70 kDa compared to the WT and complementation strain, indicating a possible change in Atl availability. We also tested the susceptibility of the *957* mutant towards the cell wall active beta-lactam antibiotic oxacillin. The *957* mutant exhibited a small but significant increase in MIC towards this antibiotic compared to the WT and the complementation strain (Fig. 4E), indicating potential changes to the peptidoglycan

structure. To test this, we next analyzed the muropeptide profile of mutanolysin-digested

peptidoglycan isolated from the WT, the *957* mutant and complementation strain after growth in

LBN or LB 0.4 M NaCl medium. Upon first inspection, no peaks were absent in any condition or strain

(Supplementary Fig. S4D and S4E). Probing further, mono- and multimers of muropeptides up to 7-

mers were quantified but no significant changes were found when strains were cultured in LBN

medium (Fig. 5A). At the 0.4 M NaCl condition however, the muropeptide profiles of the *957* mutant

strain exhibited a significant decrease in the total amount of di- and trimers and a significant

increase in higher multimers compared to the WT and complementation strain (Fig. 5B). This

underlines the involvement of gene *957* in the cell wall homeostasis and suggests that the sensitivity

to NaCl could potentially be caused by higher crosslinking and potentially rigidity of the

peptidoglycan in the mutant.

Salt-resistant suppressors possess variations in the transglycosylase domain of Pbp2

Next, we attempted to generate *957* suppressor strains that showed improved growth in the

presence of 0.5 M NaCl, with the idea that by mapping and investigating the compensatory

mutations, further insight into the cellular functions of protein 957 could be gained. We readily

obtained suppressor strains by growth if the *957* mutant strain in NaCl containing liquid medium.

These strains grew as well as or even better than the WT LAC* strain in high salt conditions (Fig. 6A)

and also grew similar to the WT in LBN (Supplementary Fig. S5A). The genome sequences of 10

independent suppressor strains were determined and single nucleotide polymorphisms (SNPs) in

one or two genes could be identified for each strain (Supplementary Table S10), except for one

strain where the coverage was insufficient, and this strain was omitted from further analysis. A

common denominator for all suppressor strains were SNPs in the *pbp2* gene, coding for the Penicillin

binding protein 2 (Pbp2) (Fig. 6B). *S. aureus* Pbp2 is a bifunctional enzyme (Wyke, 1984, Murakami

et al., 1994), which possesses transglycosylase and transpeptidase activity and is involved in

peptidoglycan synthesis. All nine discovered SNPs were unique and led to mutations in the amino

acid sequence in the transglycosylase domain but neither to stop codons, nor frameshifts nor

mutations in the transpeptidase domain. The mutations were mapped (Fig. 6B) onto an available

Pbp2 structure (3DWK) (Lovering *et al.*, 2008) to see if a specific area of the transglycosylase domain

had been targeted, but the amino acid substitutions were found throughout the molecule and not

only in the active site. Interestingly, also from the Tn-Seq data, *pbp2* appeared to be considerably

less important for growth at 0.5 M NaCl (Supplementary Fig. S5B and S5C), although transposon

insertions were mainly located at the beginning and end of the gene or within the promoter region.

To confirm that the *pbp2* SNPs were indeed responsible for the suppression of the salt

sensitivity of the *957* mutant, we transferred the *pbp2* SNPs from two strains into a fresh LAC*Δ*957*

background by co-transduction of a transposon in a nearby gene (Supplementary Fig. S5D,

schematic). As expected, these "recreated" suppressor strains showed also increased salt resistance

(Fig. 6C, Supplementary Fig. S6A). In addition, we repaired the *pbp2* gene in two suppressors to the

WT *pbp2* allele and this reduced their ability to cope with NaCl stress, all consistent with the

hypothesis that the SNPs in *pbp2* are responsible for the suppression phenotype (Fig. 6D,

Supplementary Fig. S6B). When the same *pbp2* SNPs were transferred into a WT LAC* strain, a

reproducible growth improvement of the SNP-bearing strains was observed in high salt medium, an

indication that the *pbp2* mutations lead to a general growth improvement of *S. aureus* under NaCl

stress conditions (Fig. 6E, Supplementary Fig. S6C). Overall these results show that several

independent mutations in the *pbp2* gene can suppress the salt sensitivity of the *957* mutant and can

improve growth of the WT strain.

Mutations in the *pbp2* gene alter moenomycin susceptibility and result in reduced peptidoglycan crosslinking

Moenomycin is a phosphoglycolipid antibiotic that inhibits the transglycosylase activity of the

S. aureus Pbp2 enzyme (reviewed in (Ostash & Walker, 2010)). To test if a decrease in Pbp2

transglycosylase activity results in improved growth of the *957* mutant in high salt conditions, cells

were grown on LB agar containing 0.5 M NaCl with or without moenomycin. The growth behaviors

of the WT strain and the *957* mutant were similar when grown on solid LB agar containing 0.5 M

NaCl but lacking moenomycin (Fig. 7A, top panel). This discrepancy in the growth behavior of the

957 mutant in high salt liquid versus solid medium was already noted when we attempted to

generate suppressors on agar plates. On the other hand, in medium supplemented with 0.02 µg/ml

moenomycin, we observed better growth of the *957* mutant compared to the WT strain (Fig. 7A,

lower panel). In addition, the *957* mutant also showed improved growth compared to the two

suppressor strains S2 and S4 when moenomycin was added (Fig. 7A, lower panel). These findings

are consistent with the idea that a partial inhibition of the glycosyltransferase activity of Pbp2 can

improve the growth of the *957* mutant in high salt medium.

Because the LAC*Δ*957* strain showed increased peptidoglycan crosslinking when compared to

a WT strain, we hypothesized that the peptidoglycan of the suppressor strains would be again less

crosslinked. To test this, the peptidoglycan from two of the suppressor strains grown in LB 0.4 M

NaCl was isolated, the muropeptide profiles determined and compared to that of the original *957*

mutant (Fig. 7B, Supplementary Fig. S6D). As before, the muropeptide profiles looked similar but

quantification revealed a significant reduction in crosslinked peptidoglycan fragments and a

significantly overrepresentation of monomeric and dimeric fragments in the suppressor strains as

compared to the original *957* mutant (Fig. 7B). These results indicate that the amount of crosslinking

of the peptidoglycan polymer is an important factor in the salt resistance of *S. aureus* and that

dysregulation of peptidoglycan crosslinking can lead to destabilizing effects.

Discussion

Discussion

Using TN-seq, we successfully identified several unidentified salt tolerance genes in the USA300

S. aureus LAC* strain and further characterized gene *957,* a gene without previously assigned

function. Our results indicate that the gene product is involved in cell envelope homeostasis but

likely not WTA attachment and we show that its absence leads to increased crosslinking of the

peptidoglycan in the presence of NaCl. Suppressors of a *957* mutant acquired mutations in *pbp2* and

the increase in peptidoglycan crosslinking was reversed in such stains. This demonstrates that the

synthesis of the peptidoglycan layer is a tightly regulated process that plays an essential role during

osmotic stress.

TN-seq experiments track the growth ability of single mutants in a mixed population and can

therefore quickly and accurately determine genes that are required under certain growth condition.

It is however always a concern that a single TN-seq experiment might not be reproducible due to

stochastic extinction of individual mutants or certain growth dynamics. In this study, we therefore

performed two independent NaCl TN-seq experiments using two different libraries and slightly

different experimental setups. Although we detected differences between experiments, overall the

data proved to be rather consistent suggesting that a single experiment would be sufficient in many

cases.

Most of the mutants we identified as NaCl essential in the TN-seq screen, proved to be sensitive

to 0.5 M NaCl, confirming the effectiveness of the TN-seq screen (Supplementary Fig. S2). In

particular, previously uncharacterized genes such as *SAUSA300_0694*, *mgtE,* and the *957* gene could

be identified as essential during NaCl stress and their phenotypes could be complemented (Fig. 2).

In our study, we opted not to experimentally validate any of the dispensable genes, as the

verification of a growth advantage is more difficult to prove than a growth defect, but these genes

will provide interesting starting points for future research. In addition, it will be interesting to

investigate higher concentrations of NaCl to see if the same or different genes are flagged as

essential and dispensable. Somewhat contradictory, our data indicated, that the c-di-AMP

phosphodiesterase enzymes GdpP and Pde2 become dispensable when cells are exposed to high

salt conditions. This is in contrast to the reports that construction and propagation of a *dacA* mutant

(coding for the cognate c-di-AMP cyclase DacA) is only possible at high salt concentrations (or in

defined chemical medium) (Zeden *et al.*, 2018). These findings do however point to an involvement

of c-di-AMP in salt mediated osmotic stress adaption in *S. aureus*.

Gene *957* was selected for further characterization as part of this study, because it had

previously not been linked to NaCl stress and had not been investigated before. Initial tests

supported our proposed link to the cell envelope. The differences in CFU/OD correlations of the WT

and *957* mutant could be explained by the reduced cell size of the *957* mutant, as this would produce

more cells per OD unit. The underlying mechanism as to why *957* mutant cells are smaller in size is

still unclear but could be related to changes in peptidoglycan homeostasis.

As gene *957* is co-transcribed with *lcpB,* we initially assumed the encoded protein is involved in

WTA attachment via LcpB. However, we were unable to demonstrate such a link. Instead the *957*

mutant exhibited increased muropeptide crosslinking in the presence of 0.4 M NaCl, which could be

suppressed by *pbp2* SNPs suggesting an involvement in peptidoglycan synthesis. We can only

speculate if the SNPs identified in *pbp2* increase or inhibit the activity of Pbp2. However, judging by

the number of independent SNPs, it seems more likely that the mutations lead to a decrease rather

than an increase in transglycosylase activity. In addition, the results from the moenomycin

sensitivity experiments using a sublethal antibiotic concentration highlights that partial inactivation

of the transglycosylase activity of Pbp2 improves the growth of the *957* strain in high salt conditions.

This is consistent with the idea that the obtained *pbp2* SNPs lead to a decrease in transglycosylase

activity. The absence of frameshift or non-sense mutations can be explained by the importance of

the C-terminal transpeptidase domain of *pbp2* (Pinho *et al.*, 2001) and this is also reflected in the

transposon insertion distribution in *pbp2*. Intriguingly, the suppressor mutants show reduced

growth compared to the *957* mutant on moenomycin plates, presumably due to transglycosylation

activity being reduced to levels that are detrimental to the cell. At this point it is unclear how

mutations in the transglycosylase domain alter the muropeptide pattern, since the peptide bonds

are made by the transpeptidase, not the transglycosylase domain. We hypothesize that a decrease

in the efficiency of the glycosylation activity and slowing down the glycan chain synthesis process

will reduce the efficiency of the subsequent transpeptidation process, resulting in decreased

peptidoglycan crosslinking in the *pbp2* SNP strains. It is also noteworthy that the *pbp2* gene becomes

dispensable under NaCl conditions, an additional indicator that Pbp2 activity needs to be avoided

under high salt conditions.

Interestingly, the *957* mutant exhibited a slightly higher oxacillin MIC than the WT. This is in

contrast to what we expected as MICs are determined in 2% (0.342 M) salt medium and this should

lead to a growth disadvantage of the mutant. However, when seen in light of the increase in

peptidoglycan crosslinking, this could explain the growth advantage of the *957* mutant in the

presence of a beta-lactam antibiotic, as this could counter the inhibition of transpeptidases by

oxacillin.

Although we were unable to determine the exact molecular mechanism, it is clear from our

data that gene *957* is involved in cell wall synthesis either directly or indirectly. Based on these data,

we hypothesize that the rigidity of the peptidoglycan cell wall could be an important factor to

counter osmotic stresses but further experimental work is needed to explore this in depth.

In conclusion, we have identified a number of previously uncharacterized factors involved in

the osmotic stress response of *S. aureus,* which in particular highlighted the importance of the cell

envelope. The generated data, will also provide a great resource for further studies on the

staphylococcal NaCl stress response.

Experimental procedures

Growth of bacteria

E. coli and *S. aureus* strains were streaked from frozen stock onto Lysogenic Broth (LB) or Tryptic Soya agar (TSA) plates, respectively. For all *E. coli* and most *S. aureus* experiments, the bacteria were grown in LB medium (LB normal, LBN, 10 g Tryptone, 5 g Yeast extract, 5 g NaCl) pH 7.5 or LB medium with extra 0.5 M NaCl added. For molecular cloning and some other experiments (as indicated in the text), *S. aureus* strains were grown in TSB medium. Growth curves were done in 125 ml flasks with 20 ml of medium without antibiotics. Cultures were inoculated from overnight cultures to a starting OD_{600} of 0.01 and growth was followed by determining OD_{600} readings every one or two hours.

TN-seq experiments

A previously published highly-saturated *S. aureus* USA300 library with a mix of six different transposons was used (Santiago *et al.*, 2015, Coe *et al.*, 2019) for replicate A. For replicate B, a new transposon library with only the blunt transposon was produced using the same techniques as described in Santiago *et al.* (Santiago *et al.*, 2015). This library contained approximately 1.2 million individual clones.

For each experiment, one tube of the transposon library was thawed on ice and used to set up a 20 ml pre-culture in LBN/Erm 5 µg/ml with a starting OD_{600} of 0.1. In the case of replicate A, Erm was added throughout all following cultivation steps, whereas Erm was not added to replicate B after the pre-culture. The resulting starter culture was grown for 1 hour and subsequently used to inoculate 25 ml of either LBN or LB 0.5 M to a starting OD_{600} of 0.00125. When the cultures reached an OD_{600} of approximately 0.3, the cultures were back-diluted 1:100-fold into 40 ml fresh medium and 10-12 ml harvested by centrifugation when the back-diluted culture reached an OD_{600} of approx.

1.4 (17 generations). The bacterial pellets were washed once with their respective growth medium

and stored at -20°C for further processing.

The library preparation for sequencing was done as described previously (Santiago *et al.*, 2015).

Briefly, genomic DNA was isolated, cut with *Not*I, biotinylated adapters ligated, the DNA cut again

with *Mme*I, and fragments ligated to Illumina adapters and the products PCR amplified,

incorporating bar codes and the Illumina adaptor sequences P5 and P7. The samples were

sequenced on an Illumina HiSeq machine after spiking with 40% PhiX DNA. The data analysis was

done using the Tufts Galaxy Server and custom python scripts as described earlier (Santiago *et al.*,

2015). To this end, the reads were trimmed up to the barcode and de-multiplexed by strain barcode.

Due to the variability in the DNA cleavage by the *Mme*I restriction enzyme, the reads that could not

be mapped to a barcode were trimmed by an additional base and the process repeated. Reads were

then mapped to the USA300_TCH1516 (CP000730.1) genome and a hop table was generated.

Statistical analysis (Mann-Whitney tests and Benjamini-Hochberg) was done using custom python

scripts (Santiago *et al.*, 2015). Further exploration of the data was done using Excel and custom R

scripts.

S. aureus cell diameter measurements

For cell diameter measurements, *S. aureus* strains ANG4054, ANG4340, ANG4341 were grown

overnight (14-20 hours) at 37°C with 180 rpm shaking in 5 ml LBN in the presence of 100 ng/ml Atet.

100 µl of these overnight cultures was transferred to a 1.5 ml reaction tube and 1.5 µl of BODIPY-

FL-vancomycin (100 µg/ml in phosphate buffered saline) was added to each tube to stain the cell

wall. After a 30 min static incubation step at 37°C, 1.5 µl of the suspension was pipetted onto a slide

covered with a 1% agarose pad and analyzed by microscopy using an Axio Imager.A2 microscope

with an EC Plan-Neofluar 100x/1.30 Oil M27 objective and images recorded using an AxioCam MR

R3. Native CZI files were opened in FIJI (Schindelin *et al.*, 2012) and cell diameters of 200 cells were

measured using the line and measure tool. Only cells without any visible septa were measured.

Experiment was conducted five times with independent biological cultures (n=5, 200 cells each).

CFU/OD correlations

The different *S. aureus* strains were grown as described for the cell diameter measurements.

Optical densities of cultures were measured as well as 100 µl of 10^{-6} dilutions (made in their

respective growth medium) plated onto TSA plates. The next day, the number of colonies were

counted and the ratios of OD_{600} to CFU calculated. The experiment was conducted six time with

independent biological samples (n=6).

Zymogram assays

The different *S. aureus* strains were grown over night in TSB. The next day, the bacteria from an

OD_{600} equivalent of 20 were pelleted by centrifugation for 3 min at 17000 x *g*. The cells were washed

twice with 600 µl PBS and subsequently suspended in 50 µl SDS sample buffer. The suspension was

boiled for 20 min with interspersed shaking, centrifuged for 5 min at 17000 x *g* and 20 µl of each

sample was loaded onto Zymogram gels, which where were prepared as described previously

(Atilano *et al.*, 2014). Gels were stained and de-stained using methylene blue and water. The

experiment was performed twice, and a representative result is shown.

WTA isolation and analysis

Overnight cultures of *S. aureus* strains ANG4054, ANG4340, ANG4341, ANG1575, ANG4290,

ANG4748, ANG4749, ANG4759 were prepared in LBN and used to inoculate 50 ml LBN or LB 0.4 M

NaCl to an OD_{600} of 0.01. The cultures were incubated at 37°C with shaking until they reached an

OD_{600} of 5 to 6. Where appropriate, 10 µg/ml chloramphenicol and 100 ng/ml Atet was also added

to the medium. The bacteria from 20-24 ml culture were harvested by centrifugation for 10 min at

7000 x g. The bacterial pellets were washed with 50 mM 2-(N-morpholino) ethanesulfonic acid

(MES) pH 6.5 buffer and stored at -20°C until further use. Pellets were then processed and WTA

separated on a 20% native polyacrylamide gel by electrophoresis on a Biorad Protein XL ii cell as

described previously (Covas *et al.*, 2016) and WTA visualized by silver staining according to the

manufacturer protocol. Experiments were done with three biological replicates (n=3) and a

representative result is shown.

Peptidoglycan isolation and analysis

Peptidoglycan was prepared as described previously (Corrigan *et al.*, 2011) with the following

modifications: 0.5 L (ANG4290, ANG4382, ANG4384) or 1.0 L (ANG5054, ANG4340, ANG4341) of LB

0.4 M NaCl medium (all strains) or LBN (ANG5054, ANG4340, ANG4341) medium were inoculated

to an OD_{600} of 0.01. For strains carrying derivatives of plasmid piTET, the medium was supplemented

with 100 ng/ml Atet starting from the pre-cultures. The cultures were grown at 37°C with shaking

at 180 rpm until they reached an OD_{600} of 2-3 and bacteria were subsequently harvested by

centrifugation. Chromatography of mutanolysin digested peptidoglycan was performed as

described previously using an Agilent 1260 infinity system (Corrigan *et al.*, 2011) and muropeptide

peaks assigned according to de Jonge *et al.* (de Jonge *et al.*, 1992). For the muropeptide

quantification, a baseline was drawn, the total peak area determined as well as the peak areas for

mono- and the different multimer peaks and calculated as percentage of the total peak area. The

peak area quantification was done three times for each HPLC chromatogram and average values

were calculated. Experiments were done with three biological replicates (n=3).

Data analysis, software and statistics

Data were processed with a combination of Python 3.6 (https://www.python.org), R 3.3 & 3.5 (https://www.r-project.org/), RStudio 1.1 & 1.2 (https://www.rstudio.com/), Prism 7 and 8 (https://www.graphpad.com), ChemStation OpenLab C.01.05 (https://www.agilent.com/) and Microsoft Excel 15 and 16 (https://www.office.com/). Microscopy images were analyzed using FIJI 1.0 (https://fiji.sc/). Statistical analysis was performed with Prism using appropriate tests as described in the figure legends.

Raising of Δ*957* suppressor strains

Multiple independent overnight cultures of strain LAC*Δ*957* (ANG4290) were prepared in 5 ml LBN. These cultures were back diluted the next day into 20 ml LB 0.5 M NaCl to a starting OD_{600} of 0.05 and grown for 8-10 hours at 37°C with shaking until the cultures were slightly turbid. 50 μl of each culture was passed into 5 ml of LBN and grown overnight. The next day, appropriate culture dilutions were plated onto LBN agar plates and incubated overnight at 37°C. For each independent culture, multiple single colonies were picked and used to inoculate individual wells of a 96-well microtiter plates containing 100 μl LBN and the plates were subsequently incubated at 37°C in a 96-well plate incubator with shaking at 650 rpm. The next day, a culture aliquot was stored at -80°C, and the cultures were also diluted 1:50 in LB 0.5 M NaCl medium and 20 μl of these diluted cultures used to inoculate 180 μl of LB 0.5 M NaCl medium. The plates were incubated overnight with shaking at 37°C and the next morning the growth of the potential *957* suppressor strains compared to that of the original LAC*Δ*957* deletion strain (ANG4290), which showed low growth and the WT LAC* (ANG1575) strain, which showed good growth. For each lineage, four suppressors that showed good growth were streaked out from the previously frozen stocks and subsequently single colonies selected to set up overnight cultures. The deletion of gene *SAUSA300_0957* was confirmed by PCR

and after performing growth curves in culture flasks, one strain that showed significant growth

improvement compared to the original LAC*Δ*957* (ANG4290) strain was selected from each lineage.

These strains were streaked out again for single colonies and used to inoculate an overnight culture,

giving rise to the independently raised suppressor strains ANG4381 through ANG4394. The

compensatory mutations for several of these suppressor strains were subsequently determined by

whole genome sequencing.

Transfer of *pbp2* SNPs by phage transduction

In order to demonstrate that the SNPs in *pbp2* are responsible for the growth rescue of strain

LAC*Δ*957* (ANG4290), we transferred two different *pbp2* SNPs by phage transduction into the

original LAC*Δ*957* strain as well as repaired these SNPs in the suppressor strains. This was done by

placing an Erm-marked transposon in proximity of the *pbp*2 gene allowing at a certain frequency for

the co-transduction of the Erm marker and the *pbp2* SNP. To this end, the NTML transposon mutant

library strain NE789 (Fey *et al.*, 2013) was used, which harbors a transposon insertion in

SAUSA300_1332 (putative exonuclease) about 10 kbp upstream of the *pbp2* gene. *SAUSA300_1332*

is expected to be unrelated to salt stress and the cell wall synthesis machinery but close enough to

lead to an intermediate rate of co-transference with the *pbp2* SNPs. The transposon from NE789

was transduced into WT LAC* (Boles *et al.*, 2010) and strain LAC*Δ*957* (ANG4290) using phage Φ85

yielding strains ANG4561 and ANG452, respectively. The transposon was also transduced into the

suppressor strains ANG4382 and ANG4384 transferring either *SAUSA300_1332::Tn* only, yielding

strains ANG4527 and 4528 or transferring *SAUSA300_1332::Tn* as well as replacing the *pbp2* SNP

with a WT *pbp2* allele yielding *pbp2* repaired strains ANG4557 and ANG4568. Lysates from strains

ANG4527 and 4528 (suppressors containing *pbp2* SNPs and *SAUSA300_1332::Tn*) were used to

transfer the *pbp2* SNPs back into a clean LAC*Δ*957* strain background yielding strains ANG4624 and

ANG4625 and into WT LAC*, yielding strains ANG4563 and ANG4564. Successful repair or transfer

of the SNPs was checked by PCR and subsequent restriction digest of the product choosing enzymes

that recognize sites present in either the WT or SNP allele (ANG4382 *pbp2* SNP: BsrI. Recognition

site CCWGG, additional site introduced by SNP; ANG4384 *pbp2* SNP: SspI. Recognition site: AATATT,

site missing in SNP).

Determination of oxacillin MICs using Etest strips

S. aureus strains LAC* piTET (ANG4054), LAC*Δ*957* piTET (ANG4340) and the complementation

strain LAC*Δ*957* piTET-*957* (ANG4341) were grown overnight (22 hours) in 5 ml TSB containing 10

µg/ml chloramphenicol and 100 ng/ml Atet. The next day, the cultures were diluted to an OD_{600} of

0.1 in sterile PBS buffer and streaked with cotton swabs onto cation adjusted Mueller-Hinton agar

plates supplemented with 2% NaCl and 100 ng/ml Atet. One M.I.C.Evaluator strip was placed on

each plate and the plates were incubated for 24 hours at 35°C. MICs were then read directly from

the strips. The experiment was done with 4 biological replicates (n=4).

Moenomycin growth improvement test

S. aureus strains LAC* (ANG1575), LAC* Δ*957* (ANG4290), LAC* Δ*957* S2 (ANG4382) and LAC*

Δ*957* S4 (ANG4384) were grown overnight (18 hours) in 5 ml LBN. The next day, the cells were

diluted to an OD_{600} of 0.01 in LBN, grown until mid-exponential phase (OD_{600} 0.4-0.6) and normalized

to an OD_{600} of 0.1. The cells were 10-fold serially diluted and 5 µl of each dilution spotted onto LB

agar containing 0.5 M NaCl with either 0.02 µg/ml moenomycin (mix of moenomycin A, A12, C1, C3

and C4, Santa Cruz Biotechnology) or no moenomycin. Plates were incubated at 37°C overnight (18-

22 hours) and photographed. The experiment was done with 3 biological replicates (n=3) and one

representative result is shown.

Whole genome sequencing

Genomic DNA was either isolated using the Promega Gene Wizard kit according to the manufacturer instructions or using chloroform-isoamylalcohol as described previously (Schuster & Bertram, 2014). DNA was sent off for whole genome sequencing to MicrobesNG, Birmingham, U.K. or libraries prepared using the Illumina Nextera DNA kit and sequenced at the London Institute of Medical Sciences. Short reads were trimmed in CLC workbench genomics (Qiagen), then mapped to a custom *S. aureus* USA300 LAC* reference genome generated in a previous study (Bowman *et al.*, 2016) and single nucleotide polymorphisms called based on at least 80% frequency of occurrence. This list was compared to a manually curated list of well non false-positives and these entries were removed.

Nebraska Transposon Mutant Library (NTML) strains and complementation strains

All Nebraska transposon mutant library (NTML) strains and primers used in this study are listed in Supplementary Tables S9, S11 and S12. Transposon insertions in the respective genes were confirmed by PCRs. For complementation analysis, NTML strains NE535 (JE2 *SAUSA300_0867*::*Tn*), NE867 (JE2 *SAUSA300_0483*::*Tn*) were transformed as controls with the empty plasmid pCL55 (Lee *et al.*, 1991), yielding strains ANG4326 and ANG4328 and strains NE188 (JE2 *SAUSA300_0481*::*Tn*), NE251 (JE2 SAUSA300_0482::*Tn*), NE526 (JE2 *SAUSA300_0694*::*Tn*), NE736 (JE2 *SAUSA300_0910*::*Tn*), NE1384 (JE2 *SAUSA300_0957*::*Tn*) with plasmid piTET (Gründling & Schneewind, 2007), yielding ANG4308, ANG4310, ANG4377, ANG4336, ANG4338. The WT strain JE2 (Fey *et al.*, 2013) was also transformed with both plasmids yielding JE2 pCL55 (ANG4325) and JE2 piTET (ANG4307).

The complementation plasmid piTET-*481* for strain NE188 (JE2 *SAUSA300_0481::Tn*) was constructed by amplifying the *SAUSA300_0481* gene using primers P2378/P2379, digesting the

product with AvrII/BglII and ligating the fragment with plasmid piTET cut with the same enzymes.

The plasmid was then introduced into *E. coli* strain XL1-Blue, yielding ANG4139. After shuttling the

plasmid through *E. coli* strain IM08B (Monk *et al.*, 2012), creating strain ANG4150, the plasmid was

integrated into the *geh* locus of NE188 (JE2 *SAUSA300_0481*::*Tn*) yielding strain ANG4309. The

complementation plasmid piTET-*482* for strain NE251 (JE2 SAUSA300_0482::*Tn*), piTET-*694* for

strain NE526 (JE2 *SAUSA300_0694*::*Tn*), piTET-*910* for strain NE736 (JE2 *SAUSA300_0910*::*Tn*), and

piTET-*957* for strain NE1384 (JE2 *SAUSA300_0957*::*Tn*) were constructed in a similar manner using

primer pairs P2380/P2381 (*SAUSA300_0482*), P2388/P2389 (*SAUSA300_0694*), P2384/P2385

(*SAUSA300_0910*) and P2386/P2387 (*SAUSA300_0957*), respectively. The complementation

plasmids were recovered in *E. coli* XL1-Blue yielding strains ANG4140 (SAUSA300_0482), ANG4142

(*SAUSA300_0694*), ANG4144 (*SAUSA300_0910*) and ANG4145 (*SAUSA300_0957*), shuttled through

E. coli IM08B giving strains ANG4151 (*SAUSA300_0482*), ANG4153 (*SAUSA300_0694*), ANG4155

(*SAUSA300_0910*) and ANG4156 (*SAUSA300_0957*) and finally introduced in the respective NTML

strain yielding the complementation strains ANG4311 (*SAUSA300_0482*), ANG4378

(*SAUSA300_0694*), ANG4337 (*SAUSA300_0910*) and ANG4339 (*SAUSA300_0957*).

Complementation plasmid pCL55-*483* for strain NE867 (JE2 *SAUSA300_0483*::*Tn*) was made by

fusing PCR products of the operon promoter (in front of *SAUSA300_0481*) amplified with primers

P2588/P2553 and the *SAUSA300_0483* gene, amplified with primers P2554/P2589 together using

primers P2588/P2589. The resulting fragment was cloned into pCL55 using EcoRI/BamHI restriction

sites. The plasmid was recovered in XL1-Blue, creating strain ANG4291, shuttled through *E. coli*

IM08B (ANG4292) and subsequently introduced into NE867 (JE2 *SAUSA300_0483*::*Tn*) to create

strain ANG4327. The complementation plasmid pCL55-*867* for strain NE535 (JE2

SAUSA300_0867::*Tn*) was constructed in a similar manner, using primers P2590/P2556 and

P2557/P2591 to amplify the promoter and *SAUSA300_0867* gene, which were subsequently fused

in a second PCR using primers P2590/P2591. The resulting fragment was cloning into pCL55 using

EcoRI/BamHI restriction sites and the resulting plasmid recovered in *E. coli* XL1-Blue (ANG4293),

shuttling through IM08B (ANG4294) and finally introduced into NE535 (JE2 *SAUSA300_0867*::*Tn*),

creating the complementation strain ANG4329.

Construction of *S. aureus* gene deletion and complementation strains

S. aureus strains with in-frame gene deletions were constructed by allelic exchange using

plasmids pIMAY (Monk *et al.*, 2012) and pIMAY* (Schuster *et al.*, 2019). The gene deletion plasmids

were designed to contain approximately 1000 bp up- and downstream regions around the deletion

site, amplified from LAC* genomic DNA (Boles *et al.*, 2010), and the first and last 30 bp of the open

reading frame to be deleted.

For construction of plasmid pIMAY-Δ*957*, the up- and downstream regions of *SAUSA300_0957*

were amplified using primers P2370/P2371 and P2372/P2373, spliced together in a second PCR

using primers P2370/P2373 and cloned into pIMAY using XmaI/EcoRI. Plasmid pIMAY-Δ957 was

recovered in *E. coli* XL1-Blue, creating strain ANG4147, shuttled through *E. coli* IM08B (ANG4159)

and subsequently introduced into S*. aureus* LAC*Δ*957*. The allelic exchange to delete

SAUSA300_0957 and to create strain LAC*Δ*957* (ANG4290) was performed as previously described

(Monk *et al.*, 2012). For complementation analysis, plasmid piTET-*957* was integrated into the

chromosome of strain LAC*Δ*957*, giving rise to the complementation strain LAC*Δ*957* piTET-*957*

(ANG4341). As control, the empty plasmid piTET from ANG4163 was also integrated into the

chromosome of LAC*Δ*957* yielding strain LAC*Δ*957* piTET (ANG4340). The *lcpB* (SAUSA300_0958)

gene was deleted in a similar manner using primers P2844/P2845 and P2846/P2847 for the first and

primers P2844/P2847 for the second PCR and cloning the fragment XmaI/EcoRI into pIMAY. Plasmid

pIMAY-Δ*lcpB* was recovered in *E. coli* XL1blue (ANG4740), shuttled through *E. coli* IM08B (ANG4742)

and introduced into the *S. aureus* LAC* (ANG4744) and finally yielding the *lcpB* deletion strain

LAC*Δ*lcpB* (ANG4748).

Plasmid pIMAY*-Δ*957-958* was constructed for the production of an *SAUSA300_0957-0958*

(*lcpB*) double mutant strain in which the first 30 bases of *SAUSA300_0957* were fused to the last 30

bp of *SAUSA300_0958* (*lcpB*). This was done using primers P2844/P2848 and P2849/P2373 in the

first and primers P2844/P2373 in a second PCR. The fragment was cloned using XmaI/EcoRI into

pIMAY* (Schuster *et al.*, 2019) and the resulting plasmid pIMAY*-Δ*957-958* recovered in *E. coli* XL1-

Blue (ANG4741). The plasmid was shuttled through *E. coli* IM08B (ANG4743), introduced into *S.*

aureus LAC* (ANG4745) and following the allelic exchange procedure yielding the *S. aureus* strain

LAC*Δ*957-958* (*lcpB*) (ANG4749).

The plasmid pIMAY-Δ*mgtE* was designed to contain up- and downstream regions of

SAUSA300_0910 (*mgtE*) fused together by the first and last 30 bp of the gene in order to create a

LAC*Δ*mgtE* mutant. For this, primer pairs P2374/P2375 and P2376/P2377 were used to amplify up-

and downstream regions and subsequently spliced together by PCR using primers P2374/P2377.

The PCR fragment was digested with XmaI/EcoRI and ligated into pIMAY cut with the same enzymes.

The plasmid pIMAY-Δ*mgtE* was recovered in *E. coli* XL1-Blue yielding strain ANG4146 and shuttled

through *E. coli* IM08B yielding strain ANG4158. The plasmid pIMAY-Δ*mgtE* was then introduced into

LAC* yielding strain LAC*Δ*SAUSA300_0910* (*mgtE*) (ANG4422) after the knockout procedure. To

create an empty vector-containing control strain and a plasmid-based complementation strain, the

plasmid piTET and piTET-*910* from strains ANG4163 and ANG4155 were isolated and used to

transform strain LAC*Δ*SAUSA300_0910* (*mgtE*) (ANG4422), yielding strains ANG4445 (LAC*Δ*mgtE*

piTET) and ANG4446 (LAC*Δ*mgtE* piTET-*mgtE).*

The *tagO* gene (*SAUSA300_0731*) was inactivated using the targetron intron homing system

(Yao *et al.*, 2006) or by allelic exchange using pIMAY* (Schuster *et al.,* 2019). For the targetron

mutagenesis, plasmid pNL9164-*tagO* was produced by amplifying the upstream targetron fragment

with primers P2887 and the EBS universal primer (P2890) and the downstream fragment using

primers P2888/P2889, the fragments were fused using the IBS primer (P2887) and the EBS1d primer

(P2888). The final fragment was cloned using HindIII/BsrGI into pNL9164 (Yao *et al.*, 2006). Plasmid

pNL9164-*tagO* was recovered in *E. coli* XL1-Blue, yielding strain ANG4703, introduced into *E. coli*

IM08B to give rise to strain ANG4704 and finally introduced into *S. aureus* LAC* (Boles *et al.*, 2010)

to create strain LAC* pNL9164-*tagO* (ANG4751). The final *tagO* targetron mutant strain

LAC**tagO*::targetron (ANG4753) was generated using the method described by Yao *et al.* (Yao *et*

al., 2006). For the construction of *S. aureus* strain LAC*Δ*tagO* with an in-frame deletion in *tagO,* the

allelic exchange plasmid pIMAY*-Δ*tagO* was used. To this end, up- and downstream *tagO* fragments

were produced using primer pairs P2883/P2884 and P2885/P2886 and introduced by Gibson cloning

into XmaI/EcoRI pre-cut pIMAY*. Plasmid pIMAY*-Δ*tagO* was recovered in *E. coli* XL1-Blue,

generating strain ANG4755. After shuttling through *E. coli* strain IM08B (ANG4756), the plasmid was

introduced into *S. aureus* LAC* to create strain ANG4757. Finally, the *tagO* locus was deleted by

allelic exchange, generating strain LAC*Δ*tagO* (ANG4759). Because initial generation of the mutant

failed, cells were plated unselectively onto TSA plates and incubated at room temperature (18-25°C)

for about two weeks to enable colony differentiation between WT and Δ*tagO* mutant strains. Upon

prolonged incubation, *tagO* mutant strain colonies have an opaque appearance.

Data access

The whole genome sequencing data were deposited at the European Nucleotide Archive (ENA)

under accession id ERP115099 and TN-seq data in the short read archive (SRA) at the National

Center for Biotechnology Information (NCBI) under BioProject ID PRJNA544248. Analyzed tables can

be found in the supplementary files. For other data, please contact the corresponding author.

Acknowledgements

This research was funded by the Wellcome Trust through grants 100289 and 210671/Z/18/Z to

A.G. and the NIH grant P01AI083214 to S.W.. C.F.S. was supported by the German Research

Foundation [Deutsche Forschungsgemeinschaft (DFG)] through grant SCHU 3159/1–1 and F.C.M.K.

by a Wellcome trust Ph.D. studentship. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript. We thank Microbes NG and the

London Institute of Medical Sciences for providing whole genome sequencing services and the Tufts

sequencing facility for the sequencing services for the TN-seq experiments. We also would like to

thank Ute Bertsche for kind advice on HPLC analyses.

Author contributions

C.F.S., D.M.W., F.C.K. and A.G. performed the research; C.F.S., D.M.W., F.C.M.K., M.S., S.W. and

A.G. analyzed the data; C.F.S. and A.G. designed the research; C.F.S. and A.G. wrote the paper. All

authors approved the final version of the manuscript.

Disclosure declaration

The authors declare not conflict of interest.

Graphical abstract

Abbreviated summary

Staphylococcus aureus is able to grow in the presence of high concentrations of NaCl but the

exact genetic factors contributing to this are unknown. Using a high-throughput TN-seq approach,

we identified gene *957* as an important factor for the salt stress resistance in *S. aureus*. A *957-*

mutant was not only salt sensitive but also showed increased peptidoglycan crosslinking, altogether

highlighting the cell wall as an important barrier against osmotic stress.

References

Adams, M.R., and Moss, M.O., (2008) 7.14.2 The Organism and its Characteristics. In: Food
Microbiology (3rd Edition). Royal Society of Chemistry, pp.
Anderson, C.B., and Witter, L.D. (1982) Glutamine and proline accumulation by *Staphylococcus*
aureus with reduction in water activity. *Appl Environ Microbiol* **43**: 1501-1503.
Atilano, M.L., Pereira, P.M., Vaz, F., Catalão, M.J., Reed, P., Grilo, I.R., Sobral, R.G., Ligoxygakis, P.,
Pinho, M.G., and Filipe, S.R. (2014) Bacterial autolysins trim cell surface peptidoglycan to
prevent detection by the *Drosophila* innate immune system. *Elife* **3**: e02277.
Bae, J.H., and Miller, K.J. (1992) Identification of two proline transport systems in *Staphylococcus*
aureus and their possible roles in osmoregulation. *Appl Environ Microbiol* **58**: 471-475.
Berger-Bächi, B., Barberis-Maino, L., Strässle, A., and Kayser, F.H. (1989) FemA, a host-mediated
factor essential for methicillin resistance in *Staphylococcus aureus*: molecular cloning and
characterization. *Mol Gen Genet* **219**: 263-269.
Boles, B.R., Thoendel, M., Roth, A.J., and Horswill, A.R. (2010) Identification of genes involved in
polysaccharide-independent *Staphylococcus aureus* biofilm formation. *PLoS One* **5**: e10146.
Booth, I.R. (2014) Bacterial mechanosensitive channels: progress towards an understanding of their
roles in cell physiology. *Curr Opin Microbiol* **18**: 16-22.
Bowman, L., Zeden, M.S., Schuster, C.F., Kaever, V., and Gründling, A. (2016) New Insights into the
Cyclic Di-adenosine Monophosphate (c-di-AMP) Degradation Pathway and the Requirement
of the Cyclic Dinucleotide for Acid Stress Resistance in *Staphylococcus aureus*. *J Biol Chem*
291: 26970-26986.
Bremer, E., and Krämer, R. (2019) Responses of Microorganisms to Osmotic Stress. *Annu Rev*
Microbiol **73**: 313-334.
Brown, S., Santa Maria, J.P., Jr., and Walker, S. (2013) Wall teichoic acids of gram-positive bacteria.
Annu Rev Microbiol **67**: 313-336.
Chan, Y.G., Frankel, M.B., Dengler, V., Schneewind, O., and Missiakas, D. (2013) *Staphylococcus*
aureus mutants lacking the LytR-CpsA-Psr family of enzymes release cell wall teichoic acids
into the extracellular medium. *J Bacteriol* **195**: 4650-4659.
Chan, Y.G., Kim, H.K., Schneewind, O., and Missiakas, D. (2014) The capsular polysaccharide of
Staphylococcus aureus is attached to peptidoglycan by the LytR-CpsA-Psr (LCP) family of
enzymes. *J Biol Chem* **289**: 15680-15690.
Christian, J.H., and Waltho, J.A. (1964) The Composition of *Staphylococcus aureus* in Relation to the
Water Activity of the Growth Medium. *J Gen Microbiol* **35**: 205-213.
Coe, K.A., Lee, W., Komazin-Meredith, G., Meredith, T.C., Grad, Y.H., and Walker, S. (2019)
Comparative Tn-Seq reveals common daptomycin resistance determinants in
Staphylococcus aureus despite strain-dependent differences in essentiality of shared cell
envelope genes. *bioRxiv*: 648246.
Commichau, F.M., Gibhardt, J., Halbedel, S., Gundlach, J., and Stülke, J. (2018) A Delicate
Connection: c-di-AMP Affects Cell Integrity by Controlling Osmolyte Transport. *Trends*
Microbiol **26**: 175-185.
Corrigan, R.M., Abbott, J.C., Burhenne, H., Kaever, V., and Gründling, A. (2011) c-di-AMP is a new
second messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope
stress. *PLoS Pathog* **7**: e1002217.
Corrigan, R.M., Campeotto, I., Jeganathan, T., Roelofs, K.G., Lee, V.T., and Gründling, A. (2013)
Systematic identification of conserved bacterial c-di-AMP receptor proteins. *Proc Natl Acad*
Sci U S A **110**: 9084-9089.
Corrigan, R.M., and Gründling, A. (2013) Cyclic di-AMP: another second messenger enters the fray.
Nat Rev Microbiol **11**: 513-524.

Covas, G., Vaz, F., Henriques, G., Pinho, M.G., and Filipe, S.R. (2016) Analysis of Cell Wall Teichoic
Acids in *Staphylococcus aureus*. *Methods Mol Biol* **1440**: 201-213.
Cox, C.D., Bavi, N., and Martinac, B. (2018) Bacterial Mechanosensors. *Annu Rev Physiol* **80**: 71-93.
Csonka, L.N. (1989) Physiological and genetic responses of bacteria to osmotic stress. *Microbiol Rev*
53: 121-147.
Csonka, L.N., and Hanson, A.D. (1991) Prokaryotic osmoregulation: genetics and physiology. *Annu*
Rev Microbiol **45**: 569-606.
de Jonge, B.L., Chang, Y.S., Gage, D., and Tomasz, A. (1992) Peptidoglycan composition of a highly
methicillin-resistant *Staphylococcus aureus* strain. The role of penicillin binding protein 2A. *J*
Biol Chem **267**: 11248-11254.
Fey, P.D., Endres, J.L., Yajjala, V.K., Widhelm, T.J., Boissy, R.J., Bose, J.L., and Bayles, K.W. (2013) A
genetic resource for rapid and comprehensive phenotype screening of nonessential
Staphylococcus aureus genes. *MBio* **4**: e00537-00512.
Graham, J.E., and Wilkinson, B.J. (1992) *Staphylococcus aureus* osmoregulation: roles for choline,
glycine betaine, proline, and taurine. *J Bacteriol* **174**: 2711-2716.
Gründling, A., and Schneewind, O. (2007) Genes required for glycolipid synthesis and lipoteichoic
acid anchoring in *Staphylococcus aureus*. *J Bacteriol* **189**: 2521-2530.
Gunde-Cimerman, N., Plemenitas, A., and Oren, A. (2018) Strategies of adaptation of
microorganisms of the three domains of life to high salt concentrations. *FEMS Microbiol Rev*
42: 353-375.
Holland, L.M., Conlon, B., and O'Gara, J.P. (2011) Mutation of *tagO* reveals an essential role for wall
teichoic acids in *Staphylococcus epidermidis* biofilm development. *Microbiology* **157**: 408-
418.
Ishibashi, M., Kurokawa, K., Nishida, S., Ueno, K., Matsuo, M., and Sekimizu, K. (2007) Isolation of
temperature-sensitive mutations in *murC* of *Staphylococcus aureus*. *FEMS Microbiology*
Letters **274**: 204-209.
Jordan, S., Hutchings, M.I., and Mascher, T. (2008) Cell envelope stress response in Gram-positive
bacteria. *FEMS Microbiol Rev* **32**: 107-146.
Kho, K., and Meredith, T.C. (2018) Salt-Induced Stress Stimulates a Lipoteichoic Acid-Specific Three-
Component Glycosylation System in *Staphylococcus aureus*. *J Bacteriol* **200**.
Koprivnjak, T., Mlakar, V., Swanson, L., Fournier, B., Peschel, A., and Weiss, J.P. (2006) Cation-
induced transcriptional regulation of the dlt operon of *Staphylococcus aureus*. *J Bacteriol*
188: 3622-3630.
Lee, C.Y., Buranen, S.L., and Ye, Z.H. (1991) Construction of single-copy integration vectors for
Staphylococcus aureus. *Gene* **103**: 101-105.
Lovering, A.L., De Castro, L., and Strynadka, N.C. (2008) Identification of dynamic structural motifs
involved in peptidoglycan glycosyltransfer. *J Mol Biol* **383**: 167-177.
Mäder, U., Nicolas, P., Depke, M., Pané-Farré, J., Debarbouille, M., van der Kooi-Pol, M.M., Guérin,
C., Dérozier, S., Hiron, A., Jarmer, H., Leduc, A., Michalik, S., Reilman, E., Schaffer, M.,
Schmidt, F., Bessières, P., Noirot, P., Hecker, M., Msadek, T., Völker, U., and van Dijl, J.M.
(2016) *Staphylococcus aureus* Transcriptome Architecture: From Laboratory to Infection-
Mimicking Conditions. *PLoS Genet* **12**: e1005962.
Madiraju, M.V., Brunner, D.P., and Wilkinson, B.J. (1987) Effects of temperature, NaCl, and
methicillin on penicillin-binding proteins, growth, peptidoglycan synthesis, and autolysis in
methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* **31**: 1727-1733.
Measures, J.C. (1975) Role of amino acids in osmoregulation of non-halophilic bacteria. *Nature* **257**:
398-400.
Miller, K.J., Zelt, S.C., and Bae, J.-H. (1991) Glycine betaine and proline are the principal compatible
solutes of *Staphylococcus aureus*. *Current Microbiology* **23**: 131-137.

Mistretta, N., Brossaud, M., Telles, F., Sanchez, V., Talaga, P., and Rokbi, B. (2019) Glycosylation of
Staphylococcus aureus cell wall teichoic acid is influenced by environmental conditions. *Sci*
Rep **9**: 3212.
Mlynek, K.D., Callahan, M.T., Shimkevitch, A.V., Farmer, J.T., Endres, J.L., Marchand, M., Bayles,
K.W., Horswill, A.R., and Kaplan, J.B. (2016) Effects of Low-Dose Amoxicillin on
Staphylococcus aureus USA300 Biofilms. *Antimicrob Agents Chemother* **60**: 2639-2651.
Monk, I.R., Shah, I.M., Xu, M., Tan, M.W., and Foster, T.J. (2012) Transforming the untransformable:
application of direct transformation to manipulate genetically *Staphylococcus aureus* and
Staphylococcus epidermidis. *MBio* **3**.
Murakami, K., Fujimura, T., and Doi, M. (1994) Nucleotide sequence of the structural gene for the
penicillin-binding protein 2 of *Staphylococcus aureus* and the presence of a homologous
gene in other staphylococci. *FEMS Microbiol Lett* **117**: 131-136.
Ochiai, T. (2000) *Staphylococcus aureus* produces autolysin-susceptible cell walls during growth in a
high-NaCl and low-Ca2+ concentration medium. *Microbiol Immunol* **44**: 97-104.
Onyango, L.A., Hugh Dunstan, R., Roberts, T.K., Macdonald, M.M., and Gottfries, J. (2013)
Phenotypic variants of staphylococci and their underlying population distributions following
exposure to stress. *PLoS One* **8**: e77614.
Oshida, T., Sugai, M., Komatsuzawa, H., Hong, Y.M., Suginaka, H., and Tomasz, A. (1995) A
Staphylococcus aureus autolysin that has an N-acetylmuramoyl-L-alanine amidase domain
and an endo-beta-N-acetylglucosaminidase domain: cloning, sequence analysis, and
characterization. *Proc Natl Acad Sci U S A* **92**: 285-289.
Ostash, B., and Walker, S. (2010) Moenomycin family antibiotics: chemical synthesis, biosynthesis,
and biological activity. *Nat Prod Rep* **27**: 1594-1617.
Over, B., Heusser, R., McCallum, N., Schulthess, B., Kupferschmied, P., Gaiani, J.M., Sifri, C.D., Berger-
Bächi, B., and Stutzmann Meier, P. (2011) LytR-CpsA-Psr proteins in *Staphylococcus aureus*
display partial functional redundancy and the deletion of all three severely impairs septum
placement and cell separation. *FEMS Microbiol Lett* **320**: 142-151.
Pinho, M.G., Filipe, S.R., de Lencastre, H., and Tomasz, A. (2001) Complementation of the essential
peptidoglycan transpeptidase function of penicillin-binding protein 2 (PBP2) by the drug
resistance protein PBP2A in *Staphylococcus aureus*. *J Bacteriol* **183**: 6525-6531.
Pöhlmann-Dietze, P., Ulrich, M., Kiser, K.B., Döring, G., Lee, J.C., Fournier, J.M., Botzenhart, K., and
Wolz, C. (2000) Adherence of *Staphylococcus aureus* to endothelial cells: influence of
capsular polysaccharide, global regulator agr, and bacterial growth phase. *Infect Immun* **68**:
4865-4871.
Pourkomailian, B., and Booth, I.R. (1992) Glycine betaine transport by *Staphylococcus aureus*:
evidence for feedback regulation of the activity of the two transport systems. *Microbiology*
140 (Pt 11): 3131-3138.
Price-Whelan, A., Poon, C.K., Benson, M.A., Eidem, T.T., Roux, C.M., Boyd, J.M., Dunman, P.M.,
Torres, V.J., and Krulwich, T.A. (2013) Transcriptional profiling of *Staphylococcus aureus*
during growth in 2 M NaCl leads to clarification of physiological roles for Kdp and Ktr K+
uptake systems. *MBio* **4**.
Rahman, M.M., Hunter, H.N., Prova, S., Verma, V., Qamar, A., and Golemi-Kotra, D. (2016) The
Staphylococcus aureus Methicillin Resistance Factor FmtA Is a d-Amino Esterase That Acts
on Teichoic Acids. *MBio* **7**: e02070-02015.
Santiago, M., Matano, L.M., Moussa, S.H., Gilmore, M.S., Walker, S., and Meredith, T.C. (2015) A
new platform for ultra-high density *Staphylococcus aureus* transposon libraries. *BMC*
Genomics **16**: 252.
Schaefer, K., Matano, L.M., Qiao, Y., Kahne, D., and Walker, S. (2017) In vitro reconstitution
demonstrates the cell wall ligase activity of LCP proteins. *Nat Chem Biol* **13**: 396-401.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V., Eliceiri, K.,
Tomancak, P., and Cardona, A. (2012) Fiji: an open-source platform for biological-image
analysis. *Nat Methods* **9**: 676-682.
Schuster, C.F., Bellows, L.E., Tosi, T., Campeotto, I., Corrigan, R.M., Freemont, P., and Gründling, A.
(2016) The second messenger c-di-AMP inhibits the osmolyte uptake system OpuC in
Staphylococcus aureus. *Sci Signal* **9**: ra81.
Schuster, C.F., and Bertram, R. (2014) Fluorescence based primer extension technique to determine
transcriptional starting points and cleavage sites of RNases *in vivo*. *J Vis Exp*: e52134.
Schuster, C.F., Howard, S.A., and Gründling, A. (2019) Use of the counter selectable marker PheS*
for genome engineering in *Staphylococcus aureus*. *Microbiology*.
Schwan, W.R., Lehmann, L., and McCormick, J. (2006) Transcriptional activation of the
Staphylococcus aureus putP gene by low-proline-high osmotic conditions and during
infection of murine and human tissues. *Infect Immun* **74**: 399-409.
Scybert, S., Pechous, R., Sitthisak, S., Nadakavukaren, M.J., Wilkinson, B.J., and Jayaswal, R.K. (2003)
NaCl-sensitive mutant of *Staphylococcus aureus* has a Tn917-lacZ insertion in its ars operon.
FEMS Microbiol Lett **222**: 171-176.
Sleator, R.D., and Hill, C. (2002) Bacterial osmoadaptation: the role of osmolytes in bacterial stress
and virulence. *FEMS Microbiol Rev* **26**: 49-71.
Stapleton, M.R., Horsburgh, M.J., Hayhurst, E.J., Wright, L., Jonsson, I.M., Tarkowski, A., Kokai-Kun,
J.F., Mond, J.J., and Foster, S.J. (2007) Characterization of IsaA and SceD, two putative lytic
transglycosylases of *Staphylococcus aureus*. *J Bacteriol* **189**: 7316-7325.
Tong, S.Y., Davis, J.S., Eichenberger, E., Holland, T.L., and Fowler, V.G., Jr. (2015) *Staphylococcus
aureus* infections: epidemiology, pathophysiology, clinical manifestations, and
management. *Clin Microbiol Rev* **28**: 603-661.
Townsend, D.E., and Wilkinson, B.J. (1992) Proline transport in *Staphylococcus aureus*: a high-
affinity system and a low-affinity system involved in osmoregulation. *J Bacteriol* **174**: 2702-
2710.
Vaish, M., Price-Whelan, A., Reyes-Robles, T., Liu, J., Jereen, A., Christie, S., Alonzo, F., 3rd, Benson,
M.A., Torres, V.J., and Krulwich, T.A. (2018) Roles of *Staphylococcus aureus* Mnh1 and Mnh2
Antiporters in Salt Tolerance, Alkali Tolerance, and Pathogenesis. *J Bacteriol* **200**.
van den Berg, J., Boersma, A.J., and Poolman, B. (2017) Microorganisms maintain crowding
homeostasis. *Nat Rev Microbiol* **15**: 309-318.
Vijaranakul, U., Nadakavukaren, M.J., de Jonge, B.L., Wilkinson, B.J., and Jayaswal, R.K. (1995)
Increased cell size and shortened peptidoglycan interpeptide bridge of NaCl-stressed
Staphylococcus aureus and their reversal by glycine betaine. *J Bacteriol* **177**: 5116-5121.
Vijaranakul, U., Xiong, A., Lockwood, K., and Jayaswal, R.K. (1998) Cloning and nucleotide
sequencing of a *Staphylococcus aureus* gene encoding a branched-chain-amino-acid
transporter. *Appl Environ Microbiol* **64**: 763-767.
Vollmer, W., Blanot, D., and de Pedro, M.A. (2008) Peptidoglycan structure and architecture. *FEMS
Microbiol Rev* **32**: 149-167.
Wengender, P.A., and Miller, K.J. (1995) Identification of a PutP proline permease gene homolog
from *Staphylococcus aureus* by expression cloning of the high-affinity proline transport
system in Escherichia coli. *Appl Environ Microbiol* **61**: 252-259.
Wertheim, H.F., Melles, D.C., Vos, M.C., van Leeuwen, W., van Belkum, A., Verbrugh, H.A., and
Nouwen, J.L. (2005) The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet
Infect Dis* **5**: 751-762.

Wetzel, K.J., Bjorge, D., and Schwan, W.R. (2011) Mutational and transcriptional analyses of the
Staphylococcus aureus low-affinity proline transporter OpuD during *in vitro* growth and
infection of murine tissues. *FEMS Immunol Med Microbiol* **61**: 346-355.

Whatmore, A.M., and Reed, R.H. (1990) Determination of turgor pressure in Bacillus subtilis: a
possible role for K+ in turgor regulation. *J Gen Microbiol* **136**: 2521-2526.

Wood, J.M. (2011) Bacterial osmoregulation: a paradigm for the study of cellular homeostasis. *Annu*
Rev Microbiol **65**: 215-238.

Wyke, A.W. (1984) Isolation of five penicillin-binding proteins from *Staphylococcus aureus*. *FEMS*
Microbiology Letters **22**: 133-138.

Yao, J., Zhong, J., Fang, Y., Geisinger, E., Novick, R.P., and Lambowitz, A.M. (2006) Use of targetrons
to disrupt essential and nonessential genes in *Staphylococcus aureus* reveals temperature
sensitivity of Ll.LtrB group II intron splicing. *RNA* **12**: 1271-1281.

Zeden, M.S., Schuster, C.F., Bowman, L., Zhong, Q., Williams, H.D., and Gründling, A. (2018) Cyclic
di-adenosine monophosphate (c-di-AMP) is required for osmotic regulation in
Staphylococcus aureus but dispensable for viability in anaerobic conditions. *J Biol Chem* **293**:
3180-3200.

Figure legends

Fig. 1. TN-Seq screen reveals essential and dispensable genes during prolonged NaCl stress in *S. aureus*.

A. Workflow of TN-seq experiments. A saturated transposon library was pre-cultured for one hour,

then used to inoculate LB medium containing either normal levels (0.086 M) or 0.5 M of NaCl. After

17 generations, cells were harvested, and transposon insertion sites determined by high throughput

sequencing.

B. Growth curve. Growth of the *S. aureus* library culture in LBN and LB 0.5 M medium was followed

by determining OD_{600} readings. Dotted line indicates the 17 generation threshold. Cultures were

back-diluted once when they reached an OD of approximately 0.3 and the optical densities shown

are calculated from measured ODs time dilution factor. Culture from replicate B (n=1) is shown and

a similar growth profile was seen for replicate A.

C. Volcano plots of essential and dispensable genes in NaCl conditions compared to LBN. Negative

$\log_2$ fold-changes indicate essential and positive $\log_2$ fold-changes dispensable genes. q-value stands

for Benjamini-Hochberg false discovery rate. Black horizontal line indicates q-value of 0.05 (cut-off)

and red, orange and green vertical lines 10-, 5- and 2-fold differences in either direction. Colored

circles indicate genes for which a significant change in the number of transposon insertions was

observed above q-value cutoff and follow the same color code as the vertical lines. Very small q-

values were truncated to fit onto the graph. Data from replicate A are shown in the left panel and

from replicate B in the right panel (n=1 for each plot).

Fig. 2. Growth curves and complementation analysis of *S. aureus* strains with transposon insertions in salt essential genes.

A-B. *S. aureus* strains JE2 (WT), *SAUSA300_0694::Tn* mutant with empty plasmid and complementation strain (A) or JE2 (WT), *SAUSA300_0910* (*mgtE*), *SAUSA300_0957::Tn* with empty plasmid (B) and respective complementation strains were grown in LBN (left panels) or LB 0.5 M NaCl medium (right panels) supplemented with 100 ng/ml Atet and OD_{600} readings determined at timed intervals. Growth curves were performed in triplicates (n=3) and means and SDs of OD_{600} readings were plotted.

923
924

Fig. 3. Gene _SAUSA300_0957_ is located in a genomic region with cell envelope genes and its

deletion leads to salt sensitivity and other phenotypic changes.

A. Growth curves. _S. aureus_ strains LAC* piTET (WT), LAC*Δ_957_ piTET (Δ_957_) and the

complementation strain LAC*Δ_957_ piTET-_957_ (compl.) were grown in LBN or LB containing 0.5 M

NaCl medium and 100 ng/ml Atet. The growth was monitored by determining OD_{600} readings the

means and SDs from three independent experiments were plotted.

B. Schematic of the _S. aureus_ USA300 FPR3757 chromosomal region with gene _SAUSA300_0957_

(gene _957_). Putative promoters are shown as angled arrows (adapted from (Mäder _et al._, 2016)) and

black bars indicate the corresponding transcripts.

C. Determination of CFU/OD ratios. OD_{600} values as well as CFU/ml were determined for overnight

cultures of the *S. aureus* strains described in panel B and the means and SDs of the CFU per ml OD_{600}

of 1 from five independent experiments were plotted.

D. Cell size measurements. The cell walls of the *S. aureus* strains described in panel B were stained

with fluorescently labelled vancomycin and the cells subsequently observed under a microscope.

The diameters of 200 cells were measured and the means calculated. This average of the means and

SDs from five independent experiments were plotted.

For statistical analysis, a Kruskal-Wallis one-way ANOVA test was performed followed by a Dunn's

post-hoc test to determine p-values. Asterisks (*) indicate $p \leq 0.05$ and two asterisks (**) $p \leq 0.01$ and

ns=not significant.

Fig. 4. Inactivation of *gene 957* does not significantly alter WTA levels or autolysis but leads to

changes in oxacillin resistance

A. Growth curves. *S. aureus* strains LAC* (WT), LAC*Δ*957* (Δ*957*), LAC*Δ*lcpB*, (Δ*957*), and LAC*Δ*tagO*

(Δ*tagO*) were grown in LB 0.5 M NaCl medium and OD$_{600}$ readings determined at timed intervals.

The means and SDs from three biological replicates were plotted.

B. Detection of WTA on silver stained gels. WTA was isolated from *S. aureus* strains described in

panel A following growth in LB 0.4 M NaCl medium. The WTA was separated by electrophoresis and

visualized by silver stain. The experiment was performed three times and one representative gel

image is shown.

C. Detection of WTA on silver stained gels. Same as panel B but using the strains LAC* piTET (WT),

LAC*Δ*957* piTET (Δ*957*), and the complementation strain LAC*Δ*957* piTET-*957* (compl.) grown in LB

0.4 M NaCl medium also containing 100 ng/ml Atet.

D. Zymogram gel. Cell extracts prepared from *S. aureus* strains described in panel C were separated

on a gel containing heat killed *Micrococcus luteus* cells. Autolysins were renatured and the zones of

lysis visualized by methylene blue staining. The experiment was performed twice and one

experiment is shown.

E. Determination of oxacillin MICs. Oxacillin MICs were determined for *S. aureus* strains described

in panel C using Etest strips. The median and SDs from four biological replicates were plotted.

For statistical analysis, a Kruskal-Wallis one-way ANOVA test was performed followed by a Dunn's

post-hoc test to determine p-values. Asterisks (*) indicate p≤0.05 and ns=not significant.

Fig. 5. Inactivation of *gene 957* leads to an increase in peptidoglycan crosslinking.

A-B. Muropeptide quantification. *S. aureus* strains LAC* piTET (WT), LAC*Δ*957* piTET (Δ*957*), and the complementation strain LAC*Δ*957* piTET-*957* (compl.) were grown in either LBN (A) or LB 0.4 M NaCl (B) containing 100 ng/ml Atet. After reaching late-exponential phase, the peptidoglycan was isolated, digested and the muropeptides separated by HPLC. The areas under the peaks from the chromatograms (representative chromatograms are shown in Supplementary Fig. S5) were quantified for each mono-, di- or multimer. Bars represent means, and error bars standard deviations from three independent experiments. For statistical analyses, two-way ANOVAs were performed, followed by Dunnett's post-hoc tests using the WT as a comparison. The differences in (A) were not significant. For (B), one asterisk (*) indicates $p \leq 0.05$, two asterisks (**) $p \leq 0.01$ and four asterisks (****) $p \leq 0.0001$.

Fig. 6. The growth and peptidoglycan defect observed for the *957* mutant can be rescued by

compensatory mutations in *pbp2*.

A. Growth curves. *S. aureus* strains LAC* (WT), LAC*Δ*957* (Δ*957*) and the LAC*Δ*957* suppressors S1-

10 (S1 to 10) were grown in LB 0.5 M NaCl medium, OD$_{600}$ readings determined and the means and

SDs of four biological replicates plotted.

B. Schematic of Pbp2 with amino acid substitutions identified. Top: Structure of the *S. aureus*

penicillin binding protein Pbp2 (PDB: 3DWK, (Lovering *et al.*, 2008)) with amino acids that are altered

in the obtained suppressor strains shown in red. Bottom: Schematic of the Pbp2 enzyme with the

transglycosylase and transpeptidase domains as well as the observed amino acid changes in the

transglycosylase domain indicated.

C. Growth curves using *957* mutant strains containing *pbp2* alleles from suppressors S2 and S4.

Growth curves were performed in LB 0.5 M NaCl medium and plotted as described in panel A but

using *S. aureus* strains LAC* *1332*::Tn (WT), LAC*Δ*957 1332*::Tn (Δ*957*), LAC*Δ*957 1332*::Tn *pbp2*

SNP S2 (Δ*957 pbp2* SNP S2), LAC*Δ*957* suppressor S4 *1332*::Tn (Δ*957 pbp2* SNP S4).

D. Growth curves using *957* suppressors with their *pbp2* gene repaired to WT. Growth curves were

performed in LB 0.5 M NaCl medium and plotted as described in panel A but using *S. aureus* strains

LAC* *1332*::Tn (WT), LAC*Δ*957 1332*::Tn (Δ*957*), LAC*Δ*957* suppressor S2 *1332*::Tn (S2), LAC*Δ*957*

suppressor S4 *1332*::Tn (S4), LAC*Δ*957* suppressor S2 *1332*::Tn repaired WT *pbp2* (S2 WT *pbp2*),

and LAC*Δ*957* suppressor S4 *1332*::Tn repaired WT *pbp2* (S4 WT *pbp2*).

E. Growth curves using WT strains carrying *pbp2* SNPs. *S. aureus* strains LAC* *1332*::Tn (WT), LAC*

1332::Tn *pbp2* SNP S2 (WT *pbp2* SNP S2) and LAC* *1332*::Tn *pbp2* SNP S4 (WT *pbp2* SNP S4) were

grown in LB 0.5 M NaCl medium and OD_{600} readings determined and the means and SDs of three

biological replicates plotted.

1003

1004

Fig. 7. The Δ*957* suppressor strains show decreased peptidoglycan crosslinking compared the Δ*957*

mutant.

A. Bacterial growth on moenomycin supplemented agar plates. *S. aureus* strains LAC* (WT),

LAC*Δ*957* (Δ*957*), LAC*Δ*957* Suppressor S2 (S2) and S4 LAC*Δ*957* Suppressor S4 (S4) were grown

to exponential phase, normalized to an OD_{600} of 0.1 and plated either neat (n) or in 10-fold dilutions

onto LB agar containing 0.5 M NaCl without (top) or with (bottom) moenomycin. Shown is one

representative image of three biological replicates.

B. Muropeptide analysis. *S. aureus* strains LAC*Δ*957* (Δ*957*), LAC*Δ*957* Suppressor S2 (S2) and S4

LAC*Δ*957* Suppressor S4 (S4) were grown in 0.4 M NaCl LB medium and muropeptides quantified

as described in Fig. 5. Representative chromatograms are shown in Supplementary Fig. S6D. The

means and SDs from three biological replicates were plotted. For statistical analysis a two-way

ANOVA and Dunnett's post-hoc test was performed. One asterisk (*) indicates p≤0.05, two asterisks

(**) p≤0.01 and four asterisks (****) p≤0.0001.

1018