High-throughput transposon sequencing highlights the cell wall as an important barrier for osmotic stress in methicillin resistant *Staphylococcus aureus* and underlines a tailored response to different osmotic stressors

Short Title: Global analysis of osmotic stress responses in Staphylococcus aureus

Christopher F. Schuster[a], David M. Wiedemann[a], Freja C. M. Kirsebom[a,1], Marina Santiago[b], Suzanne Walker[b] and Angelika Gründling[a,2]

ORCID IDs: 0000-0002-4566-7226 (CFS), 0000-0001-6961-0980 (DMW), 0000-0003-4585-319X (FCK), 0000-0002-5446-3452 (MS), 0000-0002-0545-914X (SW), 0000-0002-6235-8687 (AG)

[a]Section of Molecular Microbiology and MRC Centre for Molecular Bacteriology and Infection, Imperial College London, London, SW7 2AZ, U.K.

[b]Department of Microbiology and Immunobiology, Harvard Medical School, Boston, MA 02115, U.S.A.

[1]Current address: Section of Respiratory Infections, National Heart and Lung Institute, Imperial College London, London, W2 1PG, U.K.

[2]To whom correspondence should be addressed. Email: a.grundling@imperial.ac.uk, Phone: 0044 20 7594 5256

Keywords: osmotic stress, NaCl, KCl, Sucrose, *S. aureus*, DUF25380, PBP2

Summary

Staphylococcus aureus is an opportunistic pathogen that can cause soft tissue infections but is

also a frequent cause of foodborne illnesses. One contributing factor for this food association is its

high salt tolerance allowing this organism to survive commonly used food preservation methods.

How this resistance is mediated is poorly understood, particularly during long-term exposure. In

this study, we used TN-seq to understand how the responses to osmotic stressors differ. Our

results revealed distinctly different long-term responses to NaCl, KCl and sucrose stresses. In

addition, we identified the DUF2538 domain containing gene *SAUSA300_0957* (gene *957*) as

essential under salt stress. Interestingly, a *957* mutant was less susceptible to oxacillin and showed

increased peptidoglycan crosslinking. The salt sensitivity phenotype could be suppressed by amino

acid substitutions in the transglycosylase domain of the penicillin binding protein Pbp2, and these

changes restored the peptidoglycan crosslinking to WT levels. These results indicate that increased

crosslinking of the peptidoglycan polymer can be detrimental and highlight a critical role of the

bacterial cell wall for osmotic stress resistance. This study will serve as a starting point for future

research on osmotic stress response and help develop better strategies to tackle foodborne

staphylococcal infections.

Introduction

Staphylococcus aureus is a Gram-positive bacterium that is carried by about 20 % of healthy

individuals (Wertheim *et al.*, 2005). It is also an opportunistic pathogen causing a variety of

diseases, including bacteremia, endocarditis, soft tissue infections and food poisoning (Tong *et al.*,

2015). One characteristic of *S. aureus* is its ability to grow in the presence of a high salt

concentration (up to 2.5 M). Many other bacteria are unable to grow under these conditions

(Measures, 1975), and this has been used extensively for the isolation of staphylococci. This salt

tolerance also allows *S. aureus* to thrive in its niches such as the human nares, skin (Wertheim *et*

al., 2005) and certain food products (Adams & Moss, 2008).

Bacterial cells are highly pressurized (up to 1.9 MPa for *Bacillus subtilis* (Whatmore & Reed,

1990)) and contain a strongly crowded cytoplasm that is hyperosmotic to the environment. Water

is not transported into the cell actively but rather influxes passively into the cell due to the high

internal osmolality. Whenever the extracellular osmolality increases, water exits the cell which can

cause problems to multiple cellular processes, in part due to molecular crowding (Wood, 2011,

van den Berg *et al.,* 2017). Hence, *S. aureus* must possess strategies to counter the effects of salt

damage, but these are not entirely understood, as most work has been performed in *Escherichia*

coli and *B. subtilis.*

General mechanisms of hyper-osmotic stress mitigation in bacteria are the rapid import of

potassium and the accumulation of small osmotically active compounds, named compatible

solutes (reviewed in depth by (Bremer & Krämer, 2019, Gunde-Cimerman *et al.,* 2018, Booth,

2014, Cox *et al.,* 2018)). Potassium levels quickly subside in favor of an increase in the

concentration of compatible solutes.

In *S. aureus,* results consistent with the uptake of potassium (Csonka, 1989, Csonka & Hanson,

1991, Sleator & Hill, 2002) followed by the accumulation of compatible solutes have been

reported (Measures, 1975, Graham & Wilkinson, 1992, Christian & Waltho, 1964). The most

effective compatible solutes in *S. aureus* are glycine betaine and proline (Miller *et al.,* 1991,

Measures, 1975, Townsend & Wilkinson, 1992, Anderson & Witter, 1982, Bae & Miller, 1992,

Pourkomailian & Booth, 1992), which are imported by specific osmolyte transporters (Bae &

Miller, 1992, Pourkomailian & Booth, 1992). These preferred osmolytes can accumulate in the cell

to very high concentrations without negatively affecting cellular processes (Csonka & Hanson,

1991). In addition, *S. aureus* can also synthesize compatible solutes *de novo,* as in the case of

glutamine (Anderson & Witter, 1982), but this process is much slower than the uptake of

osmolytes and requires more energy.

Previous research investigating the underlying genetic factors of the *S. aureus* osmotic stress

response implicated several osmolyte transporters in this process (Price-Whelan *et al.*, 2013,

Scybert *et al.*, 2003, Schwan *et al.*, 2006, Vijaranakul *et al.*, 1998, Wengender & Miller, 1995).

These include the potassium transport systems Kdp and Ktr (Price-Whelan *et al.*, 2013), the proline

transporter PutP (Wengender & Miller, 1995, Schwan *et al.*, 2006), the arsenic transport system

Ars (Scybert *et al.*, 2003) and the branched chain amino acid uptake system BrnQ (Vijaranakul *et*

al., 1998). In addition, although currently not experimentally verified in *S. aureus*, it can be

assumed that the main glycine betaine uptake system OpuD (Wetzel *et al.*, 2011, Zeden *et al.*,

2018) is an important factor in *S. aureus* for osmotic adaptation as glycine betaine uptake reduces

sensitivity to NaCl exposure (Miller *et al.*, 1991). Of note, in recent time the second messenger

cyclic-dinucleotide-AMP (c-di-AMP) (Corrigan & Gründling, 2013) has been tied to the regulation

of osmotic stress in *S. aureus* and other Gram-positive bacteria (reviewed by Commichau *et al.*,

2018). This is through the interaction and regulation of a number of potassium (Corrigan *et al.*,

2013) and compatible solute transporters by c-di-AMP (Schuster *et al.*, 2016).

Despite these multiple countermeasures, exposure of *S. aureus* to osmotic stress leads to

morphological changes, including the formation of larger cells (Vijaranakul *et al.*, 1995) and a

thickening of the cell wall (Onyango *et al.*, 2013). However, if and which genetic factors are

required for those processes are currently not well understood. The cell wall of Gram-positive

bacteria is an important barrier and acts as a counterpart to the pressurized cytoplasm. A major

component of the cell wall is peptidoglycan, a polymer consisting of chains of repeating N-

acetylglucosamine (GlcNAc) and N-acetylmuramic acid (MurNAc) units that are crosslinked with

neighboring chains via short peptides and, in the case of *S. aureus,* pentaglycine cross-bridges

(Reviewed by (Vollmer *et al.*, 2008)). In the past, the contribution of the cell wall towards osmotic

stress resistance was often not considered in the bacterial osmotic stress resistance field. Partly,

because the integrity of the cell wall is often assumed to be a prerequisite rather than an active

adaptation to survive osmotic stresses and partly because most cell envelope stress sensing

systems are unable to directly sense osmotic stresses (Jordan *et al.*, 2008).

Nevertheless, several possible changes to the cell wall have been observed *in S. aureus* during

NaCl stress, such as shortened interpeptide bridges in the peptidoglycan (Vijaranakul *et al.*, 1995),

an increase in resistance to methicillin (Madiraju *et al.*, 1987) and increased autolysis activity

(Stapleton *et al.*, 2007, Ochiai, 2000). In addition to peptidoglycan, other components within the

cell envelope, such as teichoic acids and more specifically their modifications with D-alanine

(Koprivnjak *et al.*, 2006) and sugar residues (Mistretta *et al.*, 2019, Kho & Meredith, 2018) are

affected by osmotic stress. Combined, these findings indicate that the integrity of the cell wall is

an important requirement for osmotic resistance.

Most studies on osmotic stress in *S. aureus* have been conducted using NaCl salt as the

osmotic stressor. The ionic properties of NaCl have important implications on protein stability and

activity as its accumulation can lead to denaturation of proteins. A few studies in *S. aureus* have

also been performed with non-ionic osmotic stressors such as sucrose, sorbitol glycerol and amino

acids (Stewart *et al.*, 2005, Stewart *et al.*, 2002, Mitchell & Moyle, 1959, Schwan *et al.*, 2006). All

of these different stresses are often used synonymously with osmotic stress, in part influenced by

the findings that the initial mitigation of osmotic stress by either salts or sugars can be prevented

by the accumulation of potassium and compatible solutes. However, there are potential

differences in the long-term adaptation to ionic and non-ionic osmotic stressors, which up to date

have not been investigated in detail. It is currently also not known if the adaptations besides the

accumulation of potassium and osmolytes are similar or differ depending on the osmotic stressor,

and this was addressed as part of this study.

Using a transposon insertion sequencing (TN-seq) method (Santiago *et al.*, 2015), we

determined on a whole genome level genes that are essential or dispensable during long-term

osmotic stress caused by exposure to NaCl, KCl and sucrose. This yielded very different sets of

essential and dispensable candidate genes, providing experimental evidence of the distinct nature

of these stresses and linking a number of previously unknown factors to the osmotic stress

response in *S. aureus*. Amongst other novel candidate genes, we identified gene *SAUSA300_0957*

as an important NaCl resistance gene in *S. aureus* and show that the encoded protein is important

for peptidoglycan homeostasis in *S. aureus*. Another protein, which we identified as important

during salt stress, is the penicillin binding protein Pbp2, further emphasizing a key function of

peptidoglycan during NaCl-induced osmotic stress. Taken together, these experiments highlight

the bacterial cell wall as a key player in the salt stress response of *S. aureus*.

Results

Different types of osmotic stresses target different sets of genes

In the literature, osmotic stress is a loosely used term, however the type of ion or osmolyte can

potentially have a great impact on how bacteria respond. To address this issue, we determined

how the stress responses to different commonly used osmolytes compare. A highly saturated *S.*

aureus transposon library was generated with a promoter-less transposon (Santiago *et al.*, 2015)

and subsequently used for transposon sequencing (TN-seq) experiments (Supplementary Table 1

for details on all TN-seq experiments). The library was propagated for 16 generations in either LB

medium (Lennox) or in LB with extra 0.5 M NaCl, 0.5 M KCl or 1.0 M sucrose added (Fig. 1A). The

molarity of sucrose was doubled to accommodate for the dissociation of the salts into two ions

and hence leading to roughly the same osmolality. The culture challenged with 0.5 M NaCl grew

the slowest, followed by 1.0 M sucrose, whereas the cells grown in 0.5 M KCl grew similarly to the

cells grown in LB (Fig. 1B). As expected for a good quality transposon library, transposon insertions

were found under all conditions in genes throughout the whole genome (Fig. 1C). Next, the

number of TN-insertions per gene following growth under the NaCl, KCl or sucrose stress condition

was compared to the number of TN-insertions per gene after growth in LB medium

(Supplementary Table 2) and in this manner conditionally essential (Supplementary Table S3) and

dispensable (Supplementary Table S4) genes identified. Fold-changes in transposon insertions per

gene and q-values were plotted in volcano plots (Fig. 1D). From this, it was evident that under KCl

stress the number of essential and dispensable genes was much lower than for NaCl or sucrose

stress, indicating a much less severe effect of KCl on *S. aureus* cells. This was also reflected when

inspecting the gene lists of the top 30 essential or dispensable genes (Supplementary Table S3 and

S4) as in the KCl condition only 15 and 2 genes respectively met the requirements of q-value ≤ 0.05

and a fold-change of 5. To identify common genes between conditions, the fold-change stringency

was relaxed to 2-fold and the overlap of genes was visualized in Venn diagrams (Fig. 1E, individual

genes in Supplementary Table S5). Three genes, namely *SAUSA300_0425* (*USA300HOU_0457*,

mpsA (*nuoF*)) coding for a cation transporter of the respiratory chain (Mayer *et al.*, 2015),

SAUSA300_0750 (*USA300HOU_0796*, *whiA*) coding for a protein of unknown function and

SAUSA300_0846 (USA300HOU_0903) encoding a possible sodium:proton antiporter were

essential in all conditions. In the case of dispensable genes, only one gene, *SAUSA300_1255*

(*USA300HOU_1294*, *mprF/fmtC*), coding for a phosphatidylglycerol lysyltransferase involved in the

defense against cationic microbial peptides, was identified in all conditions (Fig. 1E,

Supplementary Table S5). To assess similarities and differences on a gene function and cellular

pathway level, functional Voronoi maps were generated from the conditionally essential

(Supplementary Fig. S1) and dispensable (Supplementary Fig. S2) genes regardless of their p-

values. These images highlighted the essentiality of the wall teichoic acid through *tagO* and the *dlt*

operon (Supplementary Fig. S1A) and the transporters AapA and MgtE in both NaCl

(Supplementary Fig. S1A) and KCl (Supplementary Fig. S1B) but not in the sucrose stress condition

(Supplementary Fig. S1C). When the dispensable genes were investigated (Supplementary Fig. S2),

the number and type of genes in the NaCl condition differed considerably from KCl and sucrose

stress. Under NaCl stress (Supplementary Fig. S2A), in addition to the penicillin binding protein

pbp2, a large number of genes involved in respiration were found to be dispensable

(Supplementary Fig. S2A). The number of dispensable genes in the KCl (Supplementary Fig. S2B)

and sucrose (Supplementary Fig. S2C) conditions were considerably lower. Taken together, the

overlaps between conditions were small, indicating distinct modes of actions for each osmotic

stress.

TN-seq is a robust method to identify *S. aureus* genes involved in NaCl stress

In our experiments, NaCl stress elicited the strongest response. We therefore focused our

further analysis on the *S. aureus* salt stress response and conducted two additional TN-seq screens

to confirm the robustness of this approach (Figure 2A). TN-seq screens were performed using a

previously published library (Santiago *et al.*, 2015, Coe *et al.*, 2019) with six transposon variants

and the library containing only one transposon variant used in the first TN-seq experiment

(Supplementary Table S1 for overview). The libraries were propagated for 17 generations in either

LB or in medium containing 0.5 M NaCl (Fig. 2B). When the libraries were grown in high salt

medium, clustering of transposon insertions in certain regions was observed due to the

enrichment of better salt-adapted TN-strains (Supplementary Fig. S3). The fold-changes of

transposon insertions per gene under the salt stress conditions compared to the LB condition

(input library) as well as q-values were determined for each gene (Supplementary Table S6-S7).

The data were plotted as volcano plots (Fig. 2C), revealing similar numbers of conditional salt

essential and dispensable genes in both replicates. A good overlap of the top essential and

dispensable genes was observed between the replicates, confirming the robustness of the TN-Seq

screens (Supplementary Tables S8-11).

The TN-seq screens highlight the essentiality of transporters and cell envelope related genes and

the dispensability of respiration genes under high salt conditions

In both replicates, genes coding for transporters and cell envelope related genes were the

most prominent conditionally salt essential genes (Voronoi maps in Supplementary Fig. S4-S5, Top

essential and dispensable genes in Supplementary Tables S8-S11). More specifically, the TN-seq

data indicated that inactivation of the putative magnesium transporter MgtE (Schuster *et al.*,

2019) and the putative D-serine/D-alanine/glycine transporter AapA is detrimental under salt

stress (Voronoi maps in Supplementary Fig. S4, Top essential genes in Supplementary Tables S8

and S10). In addition, the *dlt* operon, coding for enzymes required for the D-alanylation of teichoic

acids, an important factor for cationic peptide resistance (Brown *et al.*, 2013), as well as the

stationary and stress sigma factor gene *sigB* were found to be essential.

An even larger number of genes was identified as dispensable during salt stress (Voronoi

maps in Supplementary Fig. S5, Top dispensable genes in Supplementary Tables S9 and S11).

Inactivating or reducing the expression of such genes is expected to help *S. aureus* survive under

salt stress. Among these dispensable genes were genes coding for the sodium antiporter Mnh2,

which has previously been shown to transport Na^+/H^+ and K^+/H^+ in high pH medium and to play an

important role in cytoplasmic pH maintenance (Vaish *et al.*, 2018). As noted earlier, a large

number of genes involved in respiration, such as the *cta/qox/men/hem* genes as well as genes

involved indirectly in respiration through quinone synthesis (shikimate pathway, *aro* operon) were

also identified as dispensable during salt stress. The data also reproducibly indicated that genes

coding for the phosphodiesterases GdpP and Pde2 involved in degradation of the signaling

molecule c-di-AMP are dispensable under high salt conditions. Using the TN-seq approach, we

were able to identify a large number of *S. aureus* genes as potentially dispensable or essential

when this organism is exposed to high salt.

Tn-seq reveals previously unidentified salt resistance genes during long-term salt exposure

The TN-seq method relies on the enrichment of strains adapted to certain growth conditions

and is therefore ideal for exploring the long-term effects on growth in response to salt stress. In a

previous study (Price-Whelan *et al.*, 2013), the adaptation of *S. aureus* to 2 M NaCl stress after 6

generations of growth was determined on a transcriptome level. This work highlighted the

importance of K$^+$ transporters for *S. aureus* to survive a high salt exposure. To evaluate how our

results compare, we determined the overlaps between our datasets and the microarray data. To

this end, the *S. aureus* strain COL locus tags used in the microarray study were converted to

TCH1516 locus tags where possible (Supplementary Table S12) and compared to the two TN-seq

datasets. The overlap between the downregulated (microarray) and dispensable (TN-seq) genes

was minimal in both replicates (1 and 7) and also very small for the upregulated and essential

genes (both 14). Three operons were consistently essential or upregulated between all datasets:

the *cap5*, *SAUSA300_0771-2*, and the *sda* operon. The *cap5* genes are involved in capsule

production (Chan *et al.*, 2014) and have been previously reported to be induced by the addition of

salt (Pöhlmann-Dietze *et al.*, 2000). Genes *SAUSA300_0771-2* encode for membrane proteins that

are predicted to form a threonine/serine exporter (Interpro database IDs: IPR010619 and

IPR024528) and the *sda* genes are annotated as L-serine dehydratase components and a regulator

protein. The low overlap between the microarray and TN-seq data (Supplementary Table S12)

suggests that in our screen we have identified a number of previously unknown salt tolerance

genes.

Confirmation of genes essential during salt stress using defined mutants

To determine to what extent different genes identified in the TN-seq screen contribute to the salt resistance of *S. aureus*, a set of potentially salt essential genes from dataset B were chosen for further investigation (Supplementary Table S13). For this analysis we used defined transposon mutants available from the *S. aureus* NTML transposon mutant library (Fey *et al.*, 2013). The growth of the different NTML transposon mutant strains was assessed in LB and LB 0.5 M NaCl medium. In LB medium, most strains grew similarly to the WT strain JE2, indicating that inactivation of the respective gene does not greatly affect bacterial growth in normal LB medium (Supplementary Figs. S6A-D, left panels). Exceptions were strains NE1109 (*sigB*), NE1778 (*lcpB*) and NE188 (*mfd*), cultures of which reached slightly lower final optical densities (Supplementary Figs. S6A-D, left panels). At high osmolality conditions (LB 0.5 M NaCl), all strains showed a growth defect compared to the WT strain (Supplementary Figs. S6A-D, right panels), validating the TN-seq as a method to identify *S. aureus* genes that are important during salt stress. Several strains exhibited strong growth defects in the presence of 0.5 M NaCl with strain NE1384, containing a transposon insertion in *SAUSA300_0957* showing extremely reduced growth (Supplementary Figs. S6C and D). To confirm that the salt sensitivity was mediated by the inactivation of the genes in questions, we constructed complementation plasmids for seven of the most promising candidates by either expressing the gene of interest from its native promoter or an anhydrotetracycline (Atet) inducible promoter in the transposon mutant strains. While no complementation was observed for four strains (Supplementary Figs. S7A and S7B), the salt-dependent growth defect could be complemented for strains carrying mutations in *SAUSA300_0694* (Fig. 3A), *SAUSA300_0910* (*mgtE*) and *SAUSA300_0957* (Fig. 3B). *SAUSA300_0694* encodes a hypothetical protein with 6 predicted transmembrane helices but no other identifiable domain motif. MgtE (SAUSA300_0910) is a predicted magnesium transporter, which we have recently shown can also contribute to cobalt toxicity in *S. aureus* (Schuster *et al.*, 2019). *SAUSA300_0957* (from here on out referred to as gene

242   *957*) codes for a cytoplasmic protein with a DUF2538 domain of unknown function that is

243   conserved in Actinobacteria and Firmicutes. We further investigated this gene in this study

244   because the *957* mutant exhibited the strongest salt-sensitivity phenotype (Fig. 3B).

Deletion of gene *957* causes physiological changes leading to a smaller cell size

As all previous experiments were performed with transposon mutants, we first constructed

strain LAC*Δ*957* with a marker-less in-frame deletion of gene *957*. Strain LAC*Δ*957* had the

expected growth defect in the presence of 0.5 M NaCl and this phenotype could be complemented

by expressing a functional copy of *957* (Fig. 4A). To gain further insight into the molecular function

of gene *957*, we analyzed the neighboring genomic area. Transcriptome data (Mäder *et al.*, 2016)

suggested that gene *957* is in an operon with *lcpB* (Fig. 4B), which is one of three wall teichoic acid

(WTA) ligases present in *S. aureus* (Chan *et al.*, 2013, Over *et al.*, 2011, Schaefer *et al.*, 2017). *fmtA*

is located upstream of *957* and has been proposed to code for an esterase that can remove D-

alanine modifications from teichoic acids (Rahman *et al.*, 2016) and is involved in methicillin

resistance (Berger-Bächi *et al.*, 1989). Genes coding for a predicted acetyltransferase and *atl*

coding for the major *S. aureus* autolysin are found immediately downstream of the *957* operon

(Oshida *et al.*, 1995). Due to the presence of many cell envelope related genes in the vicinity of

gene *957*, we proceeded to test if the *957* mutant displays other phenotypes connected to cell

envelope homeostasis. First, WT LAC*, the *957* mutant and the complementation strain were

grown in LB and the ratio of colony forming units to optical density (CFU/OD) was determined. The

CFU/OD ratio was significantly larger for the *957* mutant (60.8 ± 6.8 x10^7 CFU/OD) mutant as

compared to strain LAC* (37.5 ± 9.3 x10^7 CFU/OD) when grown in LB, suggesting a change in cell

shape or size (Fig. 4C). This phenotype was restored in the complementation strain (37.3 ± 3.6 x10^7

CFU/OD) (Fig. 4C). When the average cell diameter was determined by microscopy, a small but

significant reduction in cell diameter was seen for the *957* mutant (1.25 ± 0.02 μm) as compared to

the WT (1.36±0.03 µm) and the cell size was restored to wild type levels in the complementation

strain (1.35±0.05 nm) (Fig. 4D). These results support our hypothesis that the envelope might be a

target of *gene 957* and prompted us to look into these processes in more detail.

Salt sensitivity of *957* is not due to interaction with LcpB but likely due to changes in

peptidoglycan crosslinking

Since the gene *957* is on the same transcript as *lcpB* coding for a WTA ligase (Fig. 4B), the

contribution of WTA to salt stress resistance as well as the contribution of *957* to the attachment

of WTA to the cell wall was assessed. Initially the growth of a *tagO* (*tarO/llm*) mutant, a strain

unable to produce WTA, was tested in LB and LB 0.5 M NaCl medium. When cells were grown in

LB, they grew normally (Supplementary Fig. S8A), but under high salt conditions, the *tagO* mutant

showed reduced growth, similar to that of the *957* mutant (Fig. 5A). This is consistent with what

has been reported previously for a *Staphylococcus epidermidis tagO* mutant (Holland *et al.*, 2011)

and indicates an important role for WTA during salt stress. When the *lcpB* mutant was propagated

in LB medium, its growth was slightly inhibited (Supplementary Fig. S8A). However, in high salt

medium (Fig. 5A) it grew unimpeded and like WT. Since the *lcpB* mutant did not show a growth

defect in high salt medium, this made it unlikely that *957* is a regulator of LcpB activity, and hence

is involved in the attachment of WTA to the peptidoglycan. To specifically investigate the

requirement of 957 in WTA attachment, we isolated WTA from strains grown in either LB or LB

containing 0.4 M NaCl and separated the polymer on native polyacrylamide gels. The slightly lower

concentration of salt was chosen because growth of the *957* and the *tagO* mutants was too

strongly inhibited at 0.5 M NaCl. As expected the *tagO* mutant did not produce any WTA

(Supplementary Fig. S8B, Fig. 5B) and consistent with previous studies (Over *et al.*, 2011, Chan *et*

al., 2013), the *lcpB* mutant exhibited slightly reduced WTA levels compared to the other strains

(Supplementary Fig. S8B, Fig. 5B). No reduction in WTA was observable in the *957* mutant

compared to the WT strain in either LB (Supplementary Fig. S8B & S8C) or LB 0.4 M NaCl (Fig. 5B &

5C) but rather a small increase, which could be complemented. Taken together, these data

indicate that 957 neither directly nor indirectly (e.g. through regulating the activity of LcpB) plays a

major role in WTA attachment.

Gene *957* is found upstream of the bi-functional autolysin gene *atl* and therefore we next

investigated potential changes in autolytic activity using zymograms. A small increase in autolytic

activity was detected in the *957* mutant (Fig. 5D) for bands around 55 kDa and 70 kDa compared

to the WT and complementation strain, indicating a possible change in Atl availability. We also

tested the susceptibility of the *957* mutant towards the cell wall active beta-lactam antibiotic

oxacillin. The *957* mutant exhibited a small but significant increase in MIC towards this antibiotic

compared to the WT and the complementation strain (Fig. 5E), indicating potential changes to the

peptidoglycan structure. To test this, we next analyzed the muropeptide profile of mutanolysin-

digested peptidoglycan isolated from the WT, the *957* mutant and complementation strain after

growth in LB or LB 0.4 M NaCl medium. Upon first inspection, no peaks were absent in any

condition or strain (Supplementary Fig. S8D and S8E). Probing further, mono- and multimers of

muropeptides up to 7-mers were quantified but no significant changes were found when strains

were cultured in LB medium (Fig. 6A). At the 0.4 M NaCl condition however, the muropeptide

profiles of the *957* mutant strain exhibited a significant decrease in the total amount of di- and

trimers and a significant increase in higher multimers compared to the WT and complementation

strain (Fig. 6B). This underlines the involvement of gene *957* in the cell wall homeostasis and

suggests that the sensitivity to NaCl could potentially be caused by higher crosslinking and possibly

rigidity of the peptidoglycan in the mutant.

Salt-resistant suppressors possess variations in the transglycosylase domain of Pbp2

Next, we attempted to generate *957* suppressor strains that showed improved growth in the

presence of 0.5 M NaCl, with the idea that by mapping and investigating the compensatory

mutations, further insight into the cellular functions of protein 957 could be gained. We readily

obtained suppressor strains by growth of the *957* mutant strain in NaCl containing liquid medium.

These strains grew as well as or even better than the WT LAC* strain in high salt conditions (Fig.

7A) and also grew similar to the WT in LB (Supplementary Fig. S9A). The genome sequences of 10

independent suppressor strains were determined and single nucleotide polymorphisms (SNPs) in

one or two genes could be identified for each strain (Supplementary Table S14), except for one

strain where the coverage was insufficient, and this strain was omitted from further analysis. A

common denominator for all suppressor strains were SNPs in the *pbp2* gene, coding for the

Penicillin binding protein 2 (Pbp2) (Fig. 7B). *S. aureus* Pbp2 is a bifunctional enzyme (Wyke, 1984,

Murakami *et al.*, 1994), which possesses transglycosylase and transpeptidase activity and is

involved in peptidoglycan synthesis. All nine discovered SNPs were unique and led to mutations in

the amino acid sequence in the transglycosylase domain but neither to stop codons, nor

frameshifts nor mutations in the transpeptidase domain. The mutations were mapped (Fig. 7B)

onto an available Pbp2 structure (3DWK) (Lovering *et al.*, 2008) to see if a specific area of the

transglycosylase domain had been targeted, but the amino acid substitutions were found

throughout the molecule and not only in the active site. Interestingly, also from the Tn-Seq data,

pbp2 appeared to be considerably less important for growth at 0.5 M NaCl (Supplementary Fig.

S9B and S9C), although transposon insertions were mainly located at the beginning and end of the

gene or within the promoter region.

To confirm that the *pbp2* SNPs were indeed responsible for the suppression of the salt

sensitivity of the *957* mutant, we transferred the *pbp2* SNPs from two strains into a fresh

LAC*Δ*957* background by co-transduction of a transposon in a nearby gene (Supplementary Fig.

S9D, schematic). As expected, these "recreated" suppressor strains showed also increased salt

resistance (Fig. 7C, Supplementary Fig. S10A). In addition, we repaired the *pbp2* gene in two

suppressors to the WT *pbp2* allele and this reduced their ability to cope with NaCl stress, all

consistent with the hypothesis that the SNPs in *pbp2* are responsible for the suppression

phenotype (Fig. 7D, Supplementary Fig. S10B). When the same *pbp2* SNPs were transferred into a

WT LAC* strain, a reproducible growth improvement of the SNP-bearing strains was observed in

high salt medium, an indication that the *pbp2* mutations lead to a general growth improvement of

S. aureus under NaCl stress conditions (Fig. 7E, Supplementary Fig. S10C). Overall these results

show that several independent mutations in the *pbp2* gene can suppress the salt sensitivity of the

957 mutant and can improve growth of the WT strain.

Mutations in the *pbp2* gene alter moenomycin susceptibility and result in reduced peptidoglycan

crosslinking

Moenomycin is a phosphoglycolipid antibiotic that inhibits the transglycosylase activity of the

S. aureus Pbp2 enzyme (reviewed in (Ostash & Walker, 2010)). To test if a decrease in Pbp2

transglycosylase activity results in improved growth of the *957* mutant in high salt conditions, cells

were grown on LB agar containing 0.5 M NaCl with or without moenomycin. The growth behaviors

of the WT strain and the *957* mutant were similar when grown on solid LB agar containing 0.5 M

NaCl but lacking moenomycin (Fig. 8A, top panel). This discrepancy in the growth behavior of the

957 mutant in high salt liquid versus solid medium was already noted when we attempted to

generate suppressors on agar plates. On the other hand, in medium supplemented with 0.02

µg/ml moenomycin, we observed better growth of the *957* mutant compared to the WT strain

(Fig. 8A, lower panel). In addition, the *957* mutant also showed improved growth compared to the

two suppressor strains S2 and S4 when moenomycin was added (Fig. 8A, lower panel). These

findings are consistent with the idea that a partial inhibition of the glycosyltransferase activity of

Pbp2 can improve the growth of the *957* mutant in high salt medium.

Because the LAC*Δ*957* strain showed increased peptidoglycan crosslinking when compared to

a WT strain, we hypothesized that the peptidoglycan of the suppressor strains would be again less

crosslinked. To test this, the peptidoglycan from two of the suppressor strains grown in LB 0.4 M

NaCl was isolated, the muropeptide profiles determined and compared to that of the original *957*

mutant (Fig. 8B, Supplementary Fig. S10D). As before, the muropeptide profiles looked similar but

quantification revealed a significant reduction in crosslinked peptidoglycan fragments and a

significantly overrepresentation of monomeric and dimeric fragments in the suppressor strains as

compared to the original *957* mutant (Fig. 8B). These results indicate that the amount of

crosslinking of the peptidoglycan polymer is an important factor in the salt resistance of *S. aureus*

and that dysregulation of peptidoglycan crosslinking can lead to destabilizing effects.

Discussion

In this work, we performed TN-seq screens with *S. aureus* exposed to different osmotic

stresses and could show that each osmotic stressor affects a defined but different set of genes. It

is therefore important to use the term osmotic stress carefully, as our results highlight that there

is not one general osmotic stress response but rather responses tailored to the individual

osmolyte. Using the TN-seq method, we successfully identified several unknown salt tolerance

genes in *S. aureus* and further characterized gene *957,* a gene without previously assigned

function. Our results indicate that this gene product is involved in cell envelope homeostasis but

likely not WTA attachment and we show that its absence leads to increased crosslinking of the

peptidoglycan in the presence of NaCl. Suppressors of a *957* mutant acquired mutations in *pbp2*

and the increase in peptidoglycan crosslinking was reversed in such stains. This demonstrates that

the synthesis of the peptidoglycan layer is a tightly regulated process that plays an essential role

during osmotic stress.

TN-seq experiments track the growth ability of single mutants in a mixed population and can

therefore quickly and accurately determine genes that are required under certain growth

condition. It is however always a concern that a single TN-seq experiment might not be

reproducible due to stochastic extinction of individual mutants or certain growth dynamics. In this

study, we therefore performed two independent NaCl TN-seq experiments using two different

libraries and slightly different experimental setups. Although we detected differences between

experiments, overall the data proved to be rather consistent suggesting that a single experiment

would be sufficient in many cases.

From our TN-seq data and the use of different osmotic stressors (NaCl, KCl and sucrose), it is

evident that although all these compounds exert similar osmotic stresses, the responses differs

greatly between them (Figure 1 and Supplementary Figure S1-S2). NaCl and KCl conditions share a

set of genes that become essential such as *aapA* and *mgtE* but differ in the set of dispensable

genes. It is intriguing that a whole set of respiration related genes become dispensable under NaCl

stress but not under KCl stress. This could indicate that respiration should be inhibited or slowed

down in high NaCl conditions, possibly by interference with sodium pumps a process less likely to

be inhibited by potassium ions. The set of essential genes under sucrose conditions was vastly

different from that of the NaCl and KCl conditions whereas the set of sucrose dispensable genes

looked similar to the KCl set. This highlights the differences in osmotic stress adaptations and

serves as a reminder that the type of stressor remains important.

Most of the strains with mutations in genes we identified as NaCl essential in the TN-seq

screens, proved to be sensitive to 0.5 M NaCl, confirming the effectiveness of the TN-seq screens

(Supplementary Fig. S6). In particular, previously uncharacterized genes such as *SAUSA300_0694*,

mgtE, and the *957* gene could be identified as essential during NaCl stress and their phenotypes

could be complemented (Fig. 3). In our study, we opted not to experimentally validate any of the

dispensable genes, as the verification of a growth advantage is more difficult to prove than a

growth defect, but these genes will provide interesting starting points for future research. In

addition, it will be interesting to investigate higher concentrations of NaCl to see if the same or

different genes are flagged as essential and dispensable. Somewhat contradictory, our data

indicated, that the c-di-AMP phosphodiesterase enzymes GdpP and Pde2 are dispensable when

cells are exposed to high salt conditions. This is in contrast to the reports that construction and

propagation of a *dacA* mutant (coding for the cognate c-di-AMP cyclase DacA) is only possible at

high salt concentrations (or in defined chemical medium) (Zeden *et al.*, 2018). These findings do

however point to an involvement of c-di-AMP in salt mediated osmotic stress adaption in *S.*

aureus.

Gene *957* was selected for further characterization as part of this study, because it had

previously not been linked to NaCl stress and had not been investigated before. Initial tests

supported our proposed link to the cell envelope. The differences in CFU/OD correlations of the

WT and *957* mutant could be explained by the reduced cell size of the *957* mutant, as this would

produce more cells per OD unit. The underlying mechanism as to why *957* mutant cells are smaller

in size is still unclear but could be related to changes in peptidoglycan homeostasis.

As gene *957* is co-transcribed with *lcpB,* we initially assumed the encoded protein is involved

in WTA attachment via LcpB. However, we were unable to demonstrate such a link. Instead the

957 mutant exhibited increased muropeptide crosslinking in the presence of 0.4 M NaCl, which

could be suppressed by *pbp2* SNPs suggesting an involvement in peptidoglycan synthesis. We can

only speculate if the SNPs identified in *pbp2* increase or inhibit the activity of Pbp2. However,

judging by the number of independent SNPs, it seems more likely that the mutations lead to a

decrease rather than an increase in transglycosylase activity. In addition, the results from the

moenomycin sensitivity experiments using a sublethal antibiotic concentration highlights that

partial inactivation of the transglycosylase activity of Pbp2 improves the growth of the *957* strain

in high salt conditions. This is consistent with the idea that the obtained *pbp2* SNPs lead to a

decrease in transglycosylase activity. The absence of frameshift or non-sense mutations can be

explained by the importance of the C-terminal transpeptidase domain of *pbp2* (Pinho *et al.*, 2001)

and this is also reflected in the transposon insertion distribution in *pbp2*. Intriguingly, the

suppressor mutants show reduced growth compared to the *957* mutant on moenomycin plates,

presumably due to transglycosylation activity being reduced to levels that are detrimental to the

cell. At this point it is unclear how mutations in the transglycosylase domain alter the muropeptide

pattern, since the peptide bonds are made by the transpeptidase, not the transglycosylase

domain. We hypothesize that a decrease in the efficiency of the glycosylation activity and slowing

down the glycan chain synthesis process will reduce the efficiency of the subsequent

transpeptidation process, resulting in decreased peptidoglycan crosslinking in the *pbp2* SNP

strains. It is also noteworthy that the *pbp2* gene becomes dispensable under NaCl conditions, an

additional indicator that Pbp2 activity needs to be avoided under high salt conditions.

Interestingly, the *957* mutant exhibited a slightly higher oxacillin MIC than the WT. This is in

contrast to what we expected as MICs are determined in 2% (0.342 M) salt medium and this

should lead to a growth disadvantage of the mutant. However, when seen in light of the increase

in peptidoglycan crosslinking, this could explain the growth advantage of the *957* mutant in the

presence of a beta-lactam antibiotic, as this could counter the inhibition of transpeptidases by

oxacillin.

Although we were unable to determine the exact molecular mechanism, it is clear from our

data that gene *957* is involved in cell wall synthesis either directly or indirectly. Based on these

data, we hypothesize that the rigidity of the peptidoglycan cell wall could be an important factor

to counter osmotic stresses, but further experimental work is needed to explore this in depth.

In conclusion, we have demonstrated differences between the osmotic stress response to

NaCl, KCl and sucrose in *S. aureus*. In addition, we identified a number of previously

uncharacterized factors involved in the osmotic stress response of *S. aureus,* which in particular

highlighted the importance of the cell envelope. The generated data will also provide a great

resource for further studies on the staphylococcal NaCl stress response.

Experimental procedures

Growth of bacteria

E. coli and *S. aureus* strains (see Supplementary Table S15 for details) were streaked from

frozen stocks onto Lysogenic Broth (LB) or Tryptic Soya agar (TSA) plates, respectively. For all *E.*

coli and most *S. aureus* experiments, the bacteria were grown in LB medium (Lennox Recipe: 10 g

Tryptone, 5 g Yeast extract, 5 g NaCl) pH 7.5 or LB medium with extra 0.5 M NaCl, 0.5 M KCl or 1.0

M Sucrose added. For molecular cloning and some other experiments (as indicated in the text), *S.*

aureus strains were grown in TSB medium. Growth curves were performed in 125 ml flasks with 20

ml of medium without antibiotics. Cultures were inoculated from overnight cultures to a starting

OD_{600} of 0.01 and growth was followed by determining OD_{600} readings every one or two hours.

TN-seq experiments

A previously published highly-saturated *S. aureus* USA300 library with a mix of six different

transposons was used (Santiago *et al.*, 2015, Coe *et al.*, 2019). In addition, a new transposon

library with only the blunt transposon, was produced using the same techniques as described

(Santiago *et al.*, 2015) containing approximately 1.2 million individual clones.

In the first experiment (NaCl, KCl and sucrose stress), a tube of the blunt library was thawed on ice

and used to set up a 20 ml pre-culture in LB/Erm 10 µg/ml with a starting OD_{600} of 0.1. This starter

culture was grown for 1 hour and subsequently used to inoculate 25 ml of either LB (Lennox

Recipe: 10 g Tryptone, 5 g Yeast extract, 5 g NaCl) pH 7.5 or LB with extra 0.5 M NaCl, extra 0.5 M

KCl or extra 1 M sucrose without Erm to a starting OD_{600} of 0.00125. When the cultures reached an

OD_{600} of approximately 0.3, the cultures were back-diluted 1:100-fold into 40 ml fresh medium

and 10-12 ml harvested by centrifugation when the back-diluted culture reached an OD_{600} of

approx. 1.0 (16 generations). The bacterial pellets were washed once with their respective growth

medium and stored at -20°C for further processing.

In the second experiment (NaCl, Replicate A), the multiplexed promoter transposon library was

used and Erm 5 µg/ml was added to the pre-culture and subsequent cultures in either LB (Lennox

Recipe: 10 g Tryptone, 5 g Yeast extract, 5 g NaCl) pH 7.5 or LB medium containing a final

concentration of 0.5 M NaCl. Samples were collected at an OD_{600} of about 1.4 (17 generations).

The third experiment (NaCl, replicate B) was set up similar to replicate A, however the blunt library

was used in this case and Erm 5 µg/ml was only added to the pre-culture.

The library preparation for sequencing was done as described previously (Santiago *et al.,*

2015). Briefly, genomic DNA was isolated, cut with *Not*I, biotinylated adapters ligated, the DNA cut

again with *Mme*I, and fragments ligated to Illumina adapters and the products PCR amplified,

incorporating bar codes and the Illumina adaptor sequences P5 and P7. The samples were

sequenced on an Illumina HiSeq machine after spiking with 40% PhiX DNA. The data analysis was

done using the Tufts Galaxy Server and custom python scripts as described earlier (Santiago *et al.,*

2015). To this end, the reads were trimmed up to the barcode and de-multiplexed by strain

barcode. Due to the variability in the DNA cleavage by the *Mme*I restriction enzyme, the reads that

could not be mapped to a barcode were trimmed by an additional base and the process repeated.

Reads were then mapped to the USA300_TCH1516 (CP000730.1) genome and a hop table was

generated. Statistical analysis (Mann-Whitney tests and Benjamini-Hochberg) was done using

custom python scripts (Santiago *et al.*, 2015). Further exploration of the data was done using Excel

and custom R scripts.

***S. aureus* cell diameter measurements**

For cell diameter measurements, *S. aureus* strains ANG4054, ANG4340, ANG4341 were grown

overnight (14-20 hours) at 37°C with 180 rpm shaking in 5 ml LB in the presence of 100 ng/ml Atet.

100 µl of these overnight cultures was transferred to a 1.5 ml reaction tube and 1.5 µl of BODIPY-

FL-vancomycin (100 µg/ml in phosphate buffered saline) was added to each tube to stain the cell

wall. After a 30 min static incubation step at 37°C, 1.5 µl of the suspension was pipetted onto a

slide covered with a 1% agarose pad and analyzed by microscopy using an Axio Imager.A2

microscope with an EC Plan-Neofluar 100x/1.30 Oil M27 objective and images recorded using an

AxioCam MR R3. Native CZI files were opened in FIJI (Schindelin *et al.*, 2012) and cell diameters of

200 cells were measured using the line and measure tool. Only cells without any visible septa were

measured. Experiment was conducted five times with independent biological cultures (n=5, 200

cells each).

CFU/OD correlations

The different *S. aureus* strains were grown as described for the cell diameter measurements.

Optical densities of cultures were measured as well as 100 µl of 10^{-6} dilutions (made in their

respective growth medium) plated onto TSA plates. The next day, the number of colonies were

counted and the ratios of OD_{600} to CFU calculated. The experiment was conducted six time with

independent biological samples (n=6).

Zymogram assays

The different *S. aureus* strains were grown over night in TSB. The next day, the bacteria from

an OD_{600} equivalent of 20 were pelleted by centrifugation for 3 min at 17000 x *g*. The cells were

washed twice with 600 µl PBS and subsequently suspended in 50 µl SDS sample buffer. The

suspension was boiled for 20 min with interspersed shaking, centrifuged for 5 min at 17000 x *g*

and 20 µl of each sample was loaded onto Zymogram gels, which where were prepared as

described previously (Atilano *et al.*, 2014). Gels were stained and de-stained using methylene blue

and water. The experiment was performed twice, and a representative result is shown.

WTA isolation and analysis

Overnight cultures of *S. aureus* strains ANG4054, ANG4340, ANG4341, ANG1575, ANG4290,

ANG4748, ANG4749, ANG4759 were prepared in LB and used to inoculate 50 ml LB or LB 0.4 M

NaCl to an OD_{600} of 0.01. The cultures were incubated at 37°C with shaking until they reached an

OD_{600} of 5 to 6. Where appropriate, 10 µg/ml chloramphenicol and 100 ng/ml Atet was also added

to the medium. The bacteria from 20-24 ml culture were harvested by centrifugation for 10 min at

7000 x *g*. The bacterial pellets were washed with 50 mM 2-(N-morpholino) ethanesulfonic acid

(MES) pH 6.5 buffer and stored at -20°C until further use. Pellets were then processed and WTA

separated on a 20% native polyacrylamide gel by electrophoresis on a Biorad Protein XL ii cell as

described previously (Covas *et al.*, 2016) and WTA visualized by silver staining according to the

manufacturer protocol. Experiments were done with three biological replicates (n=3) and a

representative result is shown.

Peptidoglycan isolation and analysis

Peptidoglycan was prepared as described previously (Corrigan *et al.*, 2011) with the following modifications: 0.5 L (ANG4290, ANG4382, ANG4384) or 1.0 L (ANG5054, ANG4340, ANG4341) of LB 0.4 M NaCl medium (all strains) or LB (ANG5054, ANG4340, ANG4341) medium were inoculated to an OD_{600} of 0.01. For strains carrying derivatives of plasmid piTET, the medium was supplemented with 100 ng/ml Atet starting from the pre-cultures. The cultures were grown at 37°C with shaking at 180 rpm until they reached an OD_{600} of 2-3 and bacteria were subsequently harvested by centrifugation. Chromatography of mutanolysin digested peptidoglycan was performed as described previously using an Agilent 1260 infinity system (Corrigan *et al.*, 2011) and muropeptide peaks assigned according to de Jonge *et al.* (de Jonge *et al.*, 1992). For the muropeptide quantification, a baseline was drawn, the total peak area determined as well as the peak areas for mono- and the different multimer peaks and calculated as percentage of the total peak area. The peak area quantification was done three times for each HPLC chromatogram and average values were calculated. Experiments were done with three biological replicates (n=3).

Data analysis, software and statistics

Data were processed with a combination of Python 3.6 (https://www.python.org), R 3.3 & 3.5 (https://www.r-project.org/), RStudio 1.1 & 1.2 (https://www.rstudio.com/), Prism 7 and 8 (https://www.graphpad.com), ChemStation OpenLab C.01.05 (https://www.agilent.com/) and Microsoft Excel 15 and 16 (https://www.office.com/). Microscopy images were analyzed using FIJI 1.0 (https://fiji.sc/). Statistical analysis was performed with Prism using appropriate tests as described in the figure legends.

Raising of Δ957 suppressor strains

Multiple independent overnight cultures of strain LAC*Δ957 (ANG4290) were prepared in 5 ml

LB. These cultures were back diluted the next day into 20 ml LB 0.5 M NaCl to a starting OD_{600} of

0.05 and grown for 8-10 hours at 37°C with shaking until the cultures were slightly turbid. 50 μl of

each culture was passed into 5 ml of LB and grown overnight. The next day, appropriate culture

dilutions were plated onto LB agar plates and incubated overnight at 37°C. For each independent

culture, multiple single colonies were picked and used to inoculate individual wells of a 96-well

microtiter plates containing 100 μl LB and the plates were subsequently incubated at 37°C in a 96-

well plate incubator with shaking at 650 rpm. The next day, a culture aliquot was stored at -80°C,

and the cultures were also diluted 1:50 in LB 0.5 M NaCl medium and 20 μl of these diluted

cultures used to inoculate 180 μl of LB 0.5 M NaCl medium. The plates were incubated overnight

with shaking at 37°C and the next morning the growth of the potential 957 suppressor strains

compared to that of the original LAC*Δ957 deletion strain (ANG4290), which showed low growth

and the WT LAC* (ANG1575) strain, which showed good growth. For each lineage, four

suppressors that showed good growth were streaked out from the previously frozen stocks and

subsequently single colonies selected to set up overnight cultures. The deletion of gene

SAUSA300_0957 was confirmed by PCR and after performing growth curves in culture flasks, one

strain that showed significant growth improvement compared to the original LAC*Δ957

(ANG4290) strain was selected from each lineage. These strains were streaked out again for single

colonies and used to inoculate an overnight culture, giving rise to the independently raised

suppressor strains ANG4381 through ANG4394. The compensatory mutations for several of these

suppressor strains were subsequently determined by whole genome sequencing.

Transfer of *pbp2* SNPs by phage transduction

In order to demonstrate that the SNPs in *pbp2* are responsible for the growth rescue of strain

LAC*Δ*957* (ANG4290), we transferred two different *pbp2* SNPs by phage transduction into the

original LAC*Δ*957* strain as well as repaired these SNPs in the suppressor strains. This was done by

placing an Erm-marked transposon in proximity of the *pbp*2 gene allowing at a certain frequency

for the co-transduction of the Erm marker and the *pbp2* SNP. To this end, the NTML transposon

mutant library strain NE789 (Fey *et al.*, 2013) was used, which harbors a transposon insertion in

SAUSA300_1332 (putative exonuclease) about 10 kbp upstream of the *pbp2* gene.

SAUSA300_1332 is expected to be unrelated to salt stress and the cell wall synthesis machinery

but close enough to lead to an intermediate rate of co-transference with the *pbp2* SNPs. The

transposon from NE789 was transduced into WT LAC* (Boles *et al.*, 2010) and strain LAC*Δ*957*

(ANG4290) using phage Φ85 yielding strains ANG4561 and ANG452, respectively. The transposon

was also transduced into the suppressor strains ANG4382 and ANG4384 transferring either

SAUSA300_1332::Tn only, yielding strains ANG4527 and 4528 or transferring *SAUSA300_1332::Tn*

as well as replacing the *pbp2* SNP with a WT *pbp2* allele yielding *pbp2* repaired strains ANG4557

and ANG4568. Lysates from strains ANG4527 and 4528 (suppressors containing *pbp2* SNPs and

SAUSA300_1332::Tn) were used to transfer the *pbp2* SNPs back into a clean LAC*Δ*957* strain

background yielding strains ANG4624 and ANG4625 and into WT LAC*, yielding strains ANG4563

and ANG4564. Successful repair or transfer of the SNPs was checked by PCR and subsequent

restriction digest of the product choosing enzymes that recognize sites present in either the WT or

SNP allele (ANG4382 *pbp2* SNP: BsrI. Recognition site CCWGG, additional site introduced by SNP;

ANG4384 *pbp2* SNP: SspI. Recognition site: AATATT, site missing in SNP).

Determination of oxacillin MICs using Etest strips

S. aureus strains LAC* piTET (ANG4054), LAC*Δ957 piTET (ANG4340) and the complementation strain LAC*Δ957 piTET-957 (ANG4341) were grown overnight (22 hours) in 5 ml TSB containing 10 µg/ml chloramphenicol and 100 ng/ml Atet. The next day, the cultures were diluted to an OD_{600} of 0.1 in sterile PBS buffer and streaked with cotton swabs onto cation adjusted Mueller-Hinton agar plates supplemented with 2% NaCl and 100 ng/ml Atet. One M.I.C.Evaluator strip was placed on each plate and the plates were incubated for 24 hours at 35°C. MICs were then read directly from the strips. The experiment was done with 4 biological replicates (n=4).

Moenomycin growth improvement test

S. aureus strains LAC* (ANG1575), LAC* Δ957 (ANG4290), LAC* Δ957 S2 (ANG4382) and LAC* Δ957 S4 (ANG4384) were grown overnight (18 hours) in 5 ml LB. The next day, the cells were diluted to an OD_{600} of 0.01 in LB, grown until mid-exponential phase (OD_{600} 0.4-0.6) and normalized to an OD_{600} of 0.1. The cells were 10-fold serially diluted and 5 µl of each dilution spotted onto LB agar containing 0.5 M NaCl with either 0.02 µg/ml moenomycin (mix of moenomycin A, A12, C1, C3 and C4, Santa Cruz Biotechnology) or no moenomycin. Plates were incubated at 37°C overnight (18-22 hours) and photographed. The experiment was done with 3 biological replicates (n=3) and one representative result is shown.

Whole genome sequencing

Genomic DNA was either isolated using the Promega Gene Wizard kit according to the manufacturer instructions or using chloroform-isoamylalcohol as described previously (Schuster & Bertram, 2014). DNA was sent off for whole genome sequencing to MicrobesNG, Birmingham, U.K.

or libraries prepared using the Illumina Nextera DNA kit and sequenced at the London Institute of

Medical Sciences. Short reads were trimmed in CLC workbench genomics (Qiagen), then mapped

to a custom *S. aureus* USA300 LAC* reference genome generated in a previous study (Bowman *et*

al., 2016) and single nucleotide polymorphisms called based on at least 80% frequency of

occurrence. This list was compared to a manually curated list of well non false-positives and these

entries were removed.

Nebraska Transposon Mutant Library (NTML) strains and complementation strains

All Nebraska transposon mutant library (NTML) strains and primers used in this study are

listed in Supplementary Tables S13, S15 and S16. Transposon insertions in the respective genes

were confirmed by PCRs. For complementation analysis, NTML strains NE535 (JE2

SAUSA300_0867::*Tn*), NE867 (JE2 *SAUSA300_0483*::*Tn*) were transformed as controls with the

empty plasmid pCL55 (Lee *et al.*, 1991), yielding strains ANG4326 and ANG4328 and strains NE188

(JE2 *SAUSA300_0481*::*Tn*), NE251 (JE2 SAUSA300_0482::*Tn*), NE526 (JE2 *SAUSA300_0694*::*Tn*),

NE736 (JE2 *SAUSA300_0910*::*Tn*), NE1384 (JE2 *SAUSA300_0957*::*Tn*) with plasmid piTET (Gründling

& Schneewind, 2007), yielding ANG4308, ANG4310, ANG4377, ANG4336, ANG4338. The WT strain

JE2 (Fey *et al.*, 2013) was also transformed with both plasmids yielding JE2 pCL55 (ANG4325) and

JE2 piTET (ANG4307).

The complementation plasmid piTET-*481* for strain NE188 (JE2 *SAUSA300_0481::Tn*) was

constructed by amplifying the *SAUSA300_0481* gene using primers P2378/P2379, digesting the

product with AvrII/BglII and ligating the fragment with plasmid piTET cut with the same enzymes.

The plasmid was then introduced into *E. coli* strain XL1-Blue, yielding ANG4139. After shuttling the

plasmid through *E. coli* strain IM08B (Monk *et al.*, 2012), creating strain ANG4150, the plasmid was

integrated into the *geh* locus of NE188 (JE2 *SAUSA300_0481*::*Tn*) yielding strain ANG4309. The

complementation plasmid piTET-*482* for strain NE251 (JE2 SAUSA300_0482::*Tn*), piTET-*694* for

strain NE526 (JE2 *SAUSA300_0694*::*Tn*), piTET-*910* for strain NE736 (JE2 *SAUSA300_0910*::*Tn*), and

piTET-*957* for strain NE1384 (JE2 *SAUSA300_0957*::*Tn*) were constructed in a similar manner using

primer pairs P2380/P2381 (*SAUSA300_0482*), P2388/P2389 (*SAUSA300_0694*), P2384/P2385

(*SAUSA300_0910*) and P2386/P2387 (*SAUSA300_0957*), respectively. The complementation

plasmids were recovered in *E. coli* XL1-Blue yielding strains ANG4140 (SAUSA300_0482), ANG4142

(*SAUSA300_0694*), ANG4144 (*SAUSA300_0910*) and ANG4145 (*SAUSA300_0957*), shuttled through

E. coli IM08B giving strains ANG4151 (*SAUSA300_0482*), ANG4153 (*SAUSA300_0694*), ANG4155

(*SAUSA300_0910*) and ANG4156 (*SAUSA300_0957*) and finally introduced in the respective NTML

strain yielding the complementation strains ANG4311 (*SAUSA300_0482*), ANG4378

(*SAUSA300_0694*), ANG4337 (*SAUSA300_0910*) and ANG4339 (*SAUSA300_0957*).

Complementation plasmid pCL55-*483* for strain NE867 (JE2 *SAUSA300_0483*::*Tn*) was made by

fusing PCR products of the operon promoter (in front of *SAUSA300_0481*) amplified with primers

P2588/P2553 and the *SAUSA300_0483* gene, amplified with primers P2554/P2589 together using

primers P2588/P2589. The resulting fragment was cloned into pCL55 using EcoRI/BamHI

restriction sites. The plasmid was recovered in XL1-Blue, creating strain ANG4291, shuttled

through *E. coli* IM08B (ANG4292) and subsequently introduced into NE867 (JE2

SAUSA300_0483::*Tn*) to create strain ANG4327. The complementation plasmid pCL55-*867* for

strain NE535 (JE2 *SAUSA300_0867*::*Tn*) was constructed in a similar manner, using primers

P2590/P2556 and P2557/P2591 to amplify the promoter and *SAUSA300_0867* gene, which were

subsequently fused in a second PCR using primers P2590/P2591. The resulting fragment was

cloning into pCL55 using EcoRI/BamHI restriction sites and the resulting plasmid recovered in *E.*

coli XL1-Blue (ANG4293), shuttling through IM08B (ANG4294) and finally introduced into NE535

(JE2 *SAUSA300_0867*::*Tn*), creating the complementation strain ANG4329.

Construction of *S. aureus* gene deletion and complementation strains

S. aureus strains with in-frame gene deletions were constructed by allelic exchange using plasmids pIMAY (Monk *et al.*, 2012) and pIMAY* (Schuster *et al.*, 2019). The gene deletion plasmids were designed to contain approximately 1000 bp up- and downstream regions around the deletion site, amplified from LAC* genomic DNA (Boles *et al.*, 2010), and the first and last 30 bp of the open reading frame to be deleted.

For construction of plasmid pIMAY-Δ*957*, the up- and downstream regions of *SAUSA300_0957* were amplified using primers P2370/P2371 and P2372/P2373, spliced together in a second PCR using primers P2370/P2373 and cloned into pIMAY using XmaI/EcoRI. Plasmid pIMAY-Δ957 was recovered in *E. coli* XL1-Blue, creating strain ANG4147, shuttled through *E. coli* IM08B (ANG4159) and subsequently introduced into S. *aureus* LAC*Δ*957*. The allelic exchange to delete *SAUSA300_0957* and to create strain LAC*Δ*957* (ANG4290) was performed as previously described (Monk *et al.*, 2012). For complementation analysis, plasmid piTET-*957* was integrated into the chromosome of strain LAC*Δ*957*, giving rise to the complementation strain LAC*Δ*957* piTET-*957* (ANG4341). As control, the empty plasmid piTET from ANG4163 was also integrated into the chromosome of LAC*Δ*957* yielding strain LAC*Δ*957* piTET (ANG4340). The *lcpB* (SAUSA300_0958) gene was deleted in a similar manner using primers P2844/P2845 and P2846/P2847 for the first and primers P2844/P2847 for the second PCR and cloning the fragment XmaI/EcoRI into pIMAY. Plasmid pIMAY-Δ*lcpB* was recovered in *E. coli* XL1blue (ANG4740), shuttled through *E. coli* IM08B (ANG4742) and introduced into the *S. aureus* LAC* (ANG4744) and finally yielding the *lcpB* deletion strain LAC*Δ*lcpB* (ANG4748).

Plasmid pIMAY*-Δ*957-958* was constructed for the production of an *SAUSA300_0957-0958* (*lcpB*) double mutant strain in which the first 30 bases of *SAUSA300_0957* were fused to the last 30 bp of *SAUSA300_0958* (*lcpB*). This was done using primers P2844/P2848 and P2849/P2373 in the first and primers P2844/P2373 in a second PCR. The fragment was cloned using XmaI/EcoRI

into pIMAY* (Schuster *et al.*, 2019) and the resulting plasmid pIMAY*-Δ*957-958* recovered in *E.*

coli XL1-Blue (ANG4741). The plasmid was shuttled through *E. coli* IM08B (ANG4743), introduced

into *S. aureus* LAC* (ANG4745) and following the allelic exchange procedure yielding the *S. aureus*

strain LAC*Δ*957-958* (*lcpB*) (ANG4749).

The plasmid pIMAY-Δ*mgtE* was designed to contain up- and downstream regions of

SAUSA300_0910 (*mgtE*) fused together by the first and last 30 bp of the gene in order to create a

LAC*Δ*mgtE* mutant. For this, primer pairs P2374/P2375 and P2376/P2377 were used to amplify

up- and downstream regions and subsequently spliced together by PCR using primers

P2374/P2377. The PCR fragment was digested with XmaI/EcoRI and ligated into pIMAY cut with

the same enzymes. The plasmid pIMAY-Δ*mgtE* was recovered in *E. coli* XL1-Blue yielding strain

ANG4146 and shuttled through *E. coli* IM08B yielding strain ANG4158. The plasmid pIMAY-Δ*mgtE*

was then introduced into LAC* yielding strain LAC*Δ*SAUSA300_0910* (*mgtE*) (ANG4422) after the

knockout procedure. To create an empty vector-containing control strain and a plasmid-based

complementation strain, the plasmid piTET and piTET-*910* from strains ANG4163 and ANG4155

were isolated and used to transform strain LAC*Δ*SAUSA300_0910* (*mgtE*) (ANG4422), yielding

strains ANG4445 (LAC*Δ*mgtE* piTET) and ANG4446 (LAC*Δ*mgtE* piTET-*mgtE)*.

The *tagO* gene (*SAUSA300_0731*) was inactivated using the targetron intron homing system

(Yao *et al.*, 2006) or by allelic exchange using pIMAY* (Schuster *et al.*, 2019). For the targetron

mutagenesis, plasmid pNL9164-*tagO* was produced by amplifying the upstream targetron

fragment with primers P2887 and the EBS universal primer (P2890) and the downstream fragment

using primers P2888/P2889, the fragments were fused using the IBS primer (P2887) and the EBS1d

primer (P2888). The final fragment was cloned using HindIII/BsrGI into pNL9164 (Yao *et al.*, 2006).

Plasmid pNL9164-*tagO* was recovered in *E. coli* XL1-Blue, yielding strain ANG4703, introduced into

E. coli IM08B to give rise to strain ANG4704 and finally introduced into *S. aureus* LAC* (Boles *et al.*,

2010) to create strain LAC* pNL9164-*tagO* (ANG4751). The final *tagO* targetron mutant strain

LAC*tagO::targetron (ANG4753) was generated using the method described by Yao *et al.* (Yao *et*

al., 2006). For the construction of *S. aureus* strain LAC*ΔtagO with an in-frame deletion in *tagO,*

the allelic exchange plasmid pIMAY*-ΔtagO was used. To this end, up- and downstream *tagO*

fragments were produced using primer pairs P2883/P2884 and P2885/P2886 and introduced by

Gibson cloning into XmaI/EcoRI pre-cut pIMAY*. Plasmid pIMAY*-ΔtagO was recovered in *E. coli*

XL1-Blue, generating strain ANG4755. After shuttling through *E. coli* strain IM08B (ANG4756), the

plasmid was introduced into *S. aureus* LAC* to create strain ANG4757. Finally, the *tagO* locus was

deleted by allelic exchange, generating strain LAC*ΔtagO (ANG4759). Because initial generation of

the mutant failed, cells were plated unselectively onto TSA plates and incubated at room

temperature (18-25°C) for about two weeks to enable colony differentiation between WT and

ΔtagO mutant strains. Upon prolonged incubation, *tagO* mutant strain colonies have an opaque

appearance.

Data access

The whole genome sequencing data were deposited at the European Nucleotide Archive

(ENA) under accession id ERP115099 and TN-seq data in the short read archive (SRA) at the

National Center for Biotechnology Information (NCBI) under BioProject ID PRJNA544248. Analyzed

tables can be found in the supplementary files. For other data, please contact the corresponding

author.

Acknowledgements

This research was funded by the Wellcome Trust through grants 100289 and 210671/Z/18/Z

to A.G. and the NIH grant P01AI083214 to S.W.. C.F.S. was supported by the German Research

Foundation [Deutsche Forschungsgemeinschaft (DFG)] through grant SCHU 3159/1–1 and F.C.M.K.

by a Wellcome trust Ph.D. studentship. The funders had no role in study design, data collection

and analysis, decision to publish, or preparation of the manuscript. We thank Microbes NG and the

London Institute of Medical Sciences for providing whole genome sequencing services and the

Tufts sequencing facility for the sequencing services for the TN-seq experiments. We also would

like to thank Ute Bertsche for kind advice on HPLC analyses.

Author contributions

C.F.S., D.M.W., F.C.K. and A.G. performed the research; C.F.S., D.M.W., F.C.M.K., M.S., S.W.

and A.G. analyzed the data; C.F.S. and A.G. designed the research; C.F.S. and A.G. wrote the paper.

All authors approved the final version of the manuscript.

Disclosure declaration

The authors declare not conflict of interest.

Graphical abstract

Abbreviated summary

Staphylococcus aureus is able to grow in the presence of high concentrations of NaCl but the

exact genetic factors contributing to this are unknown. Using a high-throughput TN-seq approach,

we identified gene *957* as an important factor for the salt stress resistance in *S. aureus*. A *957*-

mutant was not only salt sensitive but also showed increased peptidoglycan crosslinking,

altogether highlighting the cell wall as an important barrier against osmotic stress.

References

Adams, M.R., and Moss, M.O., (2008) 7.14.2 The Organism and its Characteristics. In: Food
Microbiology (3rd Edition). Royal Society of Chemistry, pp.
Anderson, C.B., and Witter, L.D. (1982) Glutamine and proline accumulation by *Staphylococcus*
aureus with reduction in water activity. *Appl Environ Microbiol* **43**: 1501-1503.
Atilano, M.L., Pereira, P.M., Vaz, F., Catalão, M.J., Reed, P., Grilo, I.R., Sobral, R.G., Ligoxygakis, P.,
Pinho, M.G., and Filipe, S.R. (2014) Bacterial autolysins trim cell surface peptidoglycan to
prevent detection by the *Drosophila* innate immune system. *Elife* **3**: e02277.
Bae, J.H., and Miller, K.J. (1992) Identification of two proline transport systems in *Staphylococcus*
aureus and their possible roles in osmoregulation. *Appl Environ Microbiol* **58**: 471-475.
Berger-Bächi, B., Barberis-Maino, L., Strässle, A., and Kayser, F.H. (1989) FemA, a host-mediated
factor essential for methicillin resistance in *Staphylococcus aureus*: molecular cloning and
characterization. *Mol Gen Genet* **219**: 263-269.
Boles, B.R., Thoendel, M., Roth, A.J., and Horswill, A.R. (2010) Identification of genes involved in
polysaccharide-independent *Staphylococcus aureus* biofilm formation. *PLoS One* **5**: e10146.
Booth, I.R. (2014) Bacterial mechanosensitive channels: progress towards an understanding of
their roles in cell physiology. *Curr Opin Microbiol* **18**: 16-22.
Bowman, L., Zeden, M.S., Schuster, C.F., Kaever, V., and Grundling, A. (2016) New Insights into the
Cyclic Di-adenosine Monophosphate (c-di-AMP) Degradation Pathway and the
Requirement of the Cyclic Dinucleotide for Acid Stress Resistance in *Staphylococcus aureus*.
J Biol Chem **291**: 26970-26986.
Bremer, E., and Krämer, R. (2019) Responses of Microorganisms to Osmotic Stress. *Annu Rev*
Microbiol **73**: 313-334.
Brown, S., Santa Maria, J.P., Jr., and Walker, S. (2013) Wall teichoic acids of gram-positive bacteria.
Annu Rev Microbiol **67**: 313-336.
Chan, Y.G., Frankel, M.B., Dengler, V., Schneewind, O., and Missiakas, D. (2013) *Staphylococcus*
aureus mutants lacking the LytR-CpsA-Psr family of enzymes release cell wall teichoic acids
into the extracellular medium. *J Bacteriol* **195**: 4650-4659.
Chan, Y.G., Kim, H.K., Schneewind, O., and Missiakas, D. (2014) The capsular polysaccharide of
Staphylococcus aureus is attached to peptidoglycan by the LytR-CpsA-Psr (LCP) family of
enzymes. *J Biol Chem* **289**: 15680-15690.
Christian, J.H., and Waltho, J.A. (1964) The Composition of *Staphylococcus aureus* in Relation to
the Water Activity of the Growth Medium. *J Gen Microbiol* **35**: 205-213.
Coe, K.A., Lee, W., Komazin-Meredith, G., Meredith, T.C., Grad, Y.H., and Walker, S. (2019)
Comparative Tn-Seq reveals common daptomycin resistance determinants in
Staphylococcus aureus despite strain-dependent differences in essentiality of shared cell
envelope genes. *bioRxiv*: 648246.
Commichau, F.M., Gibhardt, J., Halbedel, S., Gundlach, J., and Stülke, J. (2018) A Delicate
Connection: c-di-AMP Affects Cell Integrity by Controlling Osmolyte Transport. *Trends*
Microbiol **26**: 175-185.
Corrigan, R.M., Abbott, J.C., Burhenne, H., Kaever, V., and Gründling, A. (2011) c-di-AMP is a new
second messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope
stress. *PLoS Pathog* **7**: e1002217.
Corrigan, R.M., Campeotto, I., Jeganathan, T., Roelofs, K.G., Lee, V.T., and Gründling, A. (2013)
Systematic identification of conserved bacterial c-di-AMP receptor proteins. *Proc Natl Acad*
Sci U S A **110**: 9084-9089.
Corrigan, R.M., and Gründling, A. (2013) Cyclic di-AMP: another second messenger enters the fray.
Nat Rev Microbiol **11**: 513-524.

Covas, G., Vaz, F., Henriques, G., Pinho, M.G., and Filipe, S.R. (2016) Analysis of Cell Wall Teichoic
Acids in *Staphylococcus aureus*. *Methods Mol Biol* **1440**: 201-213.
Cox, C.D., Bavi, N., and Martinac, B. (2018) Bacterial Mechanosensors. *Annu Rev Physiol* **80**: 71-93.
Csonka, L.N. (1989) Physiological and genetic responses of bacteria to osmotic stress. *Microbiol*
Rev **53**: 121-147.
Csonka, L.N., and Hanson, A.D. (1991) Prokaryotic osmoregulation: genetics and physiology. *Annu*
Rev Microbiol **45**: 569-606.
de Jonge, B.L., Chang, Y.S., Gage, D., and Tomasz, A. (1992) Peptidoglycan composition of a highly
methicillin-resistant *Staphylococcus aureus* strain. The role of penicillin binding protein 2A.
J Biol Chem **267**: 11248-11254.
Fey, P.D., Endres, J.L., Yajjala, V.K., Widhelm, T.J., Boissy, R.J., Bose, J.L., and Bayles, K.W. (2013) A
genetic resource for rapid and comprehensive phenotype screening of nonessential
Staphylococcus aureus genes. *MBio* **4**: e00537-00512.
Graham, J.E., and Wilkinson, B.J. (1992) *Staphylococcus aureus* osmoregulation: roles for choline,
glycine betaine, proline, and taurine. *J Bacteriol* **174**: 2711-2716.
Gründling, A., and Schneewind, O. (2007) Genes required for glycolipid synthesis and lipoteichoic
acid anchoring in *Staphylococcus aureus*. *J Bacteriol* **189**: 2521-2530.
Gunde-Cimerman, N., Plemenitas, A., and Oren, A. (2018) Strategies of adaptation of
microorganisms of the three domains of life to high salt concentrations. *FEMS Microbiol*
Rev **42**: 353-375.
Holland, L.M., Conlon, B., and O'Gara, J.P. (2011) Mutation of *tagO* reveals an essential role for
wall teichoic acids in *Staphylococcus epidermidis* biofilm development. *Microbiology* **157**:
408-418.
Jordan, S., Hutchings, M.I., and Mascher, T. (2008) Cell envelope stress response in Gram-positive
bacteria. *FEMS Microbiol Rev* **32**: 107-146.
Kho, K., and Meredith, T.C. (2018) Salt-Induced Stress Stimulates a Lipoteichoic Acid-Specific
Three-Component Glycosylation System in *Staphylococcus aureus*. *J Bacteriol* **200**.
Koprivnjak, T., Mlakar, V., Swanson, L., Fournier, B., Peschel, A., and Weiss, J.P. (2006) Cation-
induced transcriptional regulation of the dlt operon of *Staphylococcus aureus*. *J Bacteriol*
188: 3622-3630.
Lee, C.Y., Buranen, S.L., and Ye, Z.H. (1991) Construction of single-copy integration vectors for
Staphylococcus aureus. *Gene* **103**: 101-105.
Lovering, A.L., De Castro, L., and Strynadka, N.C. (2008) Identification of dynamic structural motifs
involved in peptidoglycan glycosyltransfer. *J Mol Biol* **383**: 167-177.
Mäder, U., Nicolas, P., Depke, M., Pané-Farré, J., Debarbouille, M., van der Kooi-Pol, M.M., Guérin,
C., Dérozier, S., Hiron, A., Jarmer, H., Leduc, A., Michalik, S., Reilman, E., Schaffer, M.,
Schmidt, F., Bessières, P., Noirot, P., Hecker, M., Msadek, T., Völker, U., and van Dijl, J.M.
(2016) *Staphylococcus aureus* Transcriptome Architecture: From Laboratory to Infection-
Mimicking Conditions. *PLoS Genet* **12**: e1005962.
Madiraju, M.V., Brunner, D.P., and Wilkinson, B.J. (1987) Effects of temperature, NaCl, and
methicillin on penicillin-binding proteins, growth, peptidoglycan synthesis, and autolysis in
methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* **31**: 1727-1733.
Mayer, S., Steffen, W., Steuber, J., and Götz, F. (2015) The *Staphylococcus aureus* NuoL-like protein
MpsA contributes to the generation of membrane potential. *J Bacteriol* **197**: 794-806.
Measures, J.C. (1975) Role of amino acids in osmoregulation of non-halophilic bacteria. *Nature*
257: 398-400.
Miller, K.J., Zelt, S.C., and Bae, J.-H. (1991) Glycine betaine and proline are the principal compatible
solutes of *Staphylococcus aureus*. *Current Microbiology* **23**: 131-137.

Mistretta, N., Brossaud, M., Telles, F., Sanchez, V., Talaga, P., and Rokbi, B. (2019) Glycosylation of
Staphylococcus aureus cell wall teichoic acid is influenced by environmental conditions. *Sci*
Rep **9**: 3212.
Mitchell, P., and Moyle, J. (1959) Permeability of the envelopes of *Staphylococcus aureus* to some
salts, amino acids, and non-electrolytes. *J Gen Microbiol* **20**: 434-441.
Monk, I.R., Shah, I.M., Xu, M., Tan, M.W., and Foster, T.J. (2012) Transforming the
untransformable: application of direct transformation to manipulate genetically
Staphylococcus aureus and *Staphylococcus epidermidis. MBio* **3**.
Murakami, K., Fujimura, T., and Doi, M. (1994) Nucleotide sequence of the structural gene for the
penicillin-binding protein 2 of *Staphylococcus aureus* and the presence of a homologous
gene in other staphylococci. *FEMS Microbiol Lett* **117**: 131-136.
Ochiai, T. (2000) *Staphylococcus aureus* produces autolysin-susceptible cell walls during growth in
a high-NaCl and low-Ca2+ concentration medium. *Microbiol Immunol* **44**: 97-104.
Onyango, L.A., Hugh Dunstan, R., Roberts, T.K., Macdonald, M.M., and Gottfries, J. (2013)
Phenotypic variants of staphylococci and their underlying population distributions
following exposure to stress. *PLoS One* **8**: e77614.
Oshida, T., Sugai, M., Komatsuzawa, H., Hong, Y.M., Suginaka, H., and Tomasz, A. (1995) A
Staphylococcus aureus autolysin that has an N-acetylmuramoyl-L-alanine amidase domain
and an endo-beta-N-acetylglucosaminidase domain: cloning, sequence analysis, and
characterization. *Proc Natl Acad Sci U S A* **92**: 285-289.
Ostash, B., and Walker, S. (2010) Moenomycin family antibiotics: chemical synthesis, biosynthesis,
and biological activity. *Nat Prod Rep* **27**: 1594-1617.
Over, B., Heusser, R., McCallum, N., Schulthess, B., Kupferschmied, P., Gaiani, J.M., Sifri, C.D.,
Berger-Bächi, B., and Stutzmann Meier, P. (2011) LytR-CpsA-Psr proteins in *Staphylococcus*
aureus display partial functional redundancy and the deletion of all three severely impairs
septum placement and cell separation. *FEMS Microbiol Lett* **320**: 142-151.
Pinho, M.G., Filipe, S.R., de Lencastre, H., and Tomasz, A. (2001) Complementation of the essential
peptidoglycan transpeptidase function of penicillin-binding protein 2 (PBP2) by the drug
resistance protein PBP2A in *Staphylococcus aureus. J Bacteriol* **183**: 6525-6531.
Pöhlmann-Dietze, P., Ulrich, M., Kiser, K.B., Döring, G., Lee, J.C., Fournier, J.M., Botzenhart, K., and
Wolz, C. (2000) Adherence of *Staphylococcus aureus* to endothelial cells: influence of
capsular polysaccharide, global regulator agr, and bacterial growth phase. *Infect Immun* **68**:
4865-4871.
Pourkomailian, B., and Booth, I.R. (1992) Glycine betaine transport by *Staphylococcus aureus*:
evidence for feedback regulation of the activity of the two transport systems. *Microbiology*
140 (Pt 11): 3131-3138.
Price-Whelan, A., Poon, C.K., Benson, M.A., Eidem, T.T., Roux, C.M., Boyd, J.M., Dunman, P.M.,
Torres, V.J., and Krulwich, T.A. (2013) Transcriptional profiling of *Staphylococcus aureus*
during growth in 2 M NaCl leads to clarification of physiological roles for Kdp and Ktr K+
uptake systems. *MBio* **4**.
Rahman, M.M., Hunter, H.N., Prova, S., Verma, V., Qamar, A., and Golemi-Kotra, D. (2016) The
Staphylococcus aureus Methicillin Resistance Factor FmtA Is a d-Amino Esterase That Acts
on Teichoic Acids. *MBio* **7**: e02070-02015.
Santiago, M., Matano, L.M., Moussa, S.H., Gilmore, M.S., Walker, S., and Meredith, T.C. (2015) A
new platform for ultra-high density *Staphylococcus aureus* transposon libraries. *BMC*
Genomics **16**: 252.
Schaefer, K., Matano, L.M., Qiao, Y., Kahne, D., and Walker, S. (2017) In vitro reconstitution
demonstrates the cell wall ligase activity of LCP proteins. *Nat Chem Biol* **13**: 396-401.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V., Eliceiri, K.,
Tomancak, P., and Cardona, A. (2012) Fiji: an open-source platform for biological-image
analysis. *Nat Methods* **9**: 676-682.
Schuster, C.F., Bellows, L.E., Tosi, T., Campeotto, I., Corrigan, R.M., Freemont, P., and Gründling, A.
(2016) The second messenger c-di-AMP inhibits the osmolyte uptake system OpuC in
Staphylococcus aureus. *Sci Signal* **9**: ra81.
Schuster, C.F., and Bertram, R. (2014) Fluorescence based primer extension technique to
determine transcriptional starting points and cleavage sites of RNases *in vivo*. *J Vis Exp*:
e52134.
Schuster, C.F., Howard, S.A., and Gründling, A. (2019) Use of the counter selectable marker PheS*
for genome engineering in *Staphylococcus aureus*. *Microbiology*.
Schwan, W.R., Lehmann, L., and McCormick, J. (2006) Transcriptional activation of the
Staphylococcus aureus putP gene by low-proline-high osmotic conditions and during
infection of murine and human tissues. *Infect Immun* **74**: 399-409.
Scybert, S., Pechous, R., Sitthisak, S., Nadakavukaren, M.J., Wilkinson, B.J., and Jayaswal, R.K.
(2003) NaCl-sensitive mutant of *Staphylococcus aureus* has a Tn917-lacZ insertion in its ars
operon. *FEMS Microbiol Lett* **222**: 171-176.
Sleator, R.D., and Hill, C. (2002) Bacterial osmoadaptation: the role of osmolytes in bacterial stress
and virulence. *FEMS Microbiol Rev* **26**: 49-71.
Stapleton, M.R., Horsburgh, M.J., Hayhurst, E.J., Wright, L., Jonsson, I.M., Tarkowski, A., Kokai-Kun,
J.F., Mond, J.J., and Foster, S.J. (2007) Characterization of IsaA and SceD, two putative lytic
transglycosylases of *Staphylococcus aureus*. *J Bacteriol* **189**: 7316-7325.
Stewart, C.M., Cole, M.B., Legan, J.D., Slade, L., and Schaffner, D.W. (2005) Solute-specific effects
of osmotic stress on *Staphylococcus aureus*. *J Appl Microbiol* **98**: 193-202.
Stewart, C.M., Cole, M.B., Legan, J.D., Slade, L., Vandeven, M.H., and Schaffner, D.W. (2002)
Staphylococcus aureus growth boundaries: moving towards mechanistic predictive models
based on solute-specific effects. *Appl Environ Microbiol* **68**: 1864-1871.
Tong, S.Y., Davis, J.S., Eichenberger, E., Holland, T.L., and Fowler, V.G., Jr. (2015) *Staphylococcus
aureus* infections: epidemiology, pathophysiology, clinical manifestations, and
management. *Clin Microbiol Rev* **28**: 603-661.
Townsend, D.E., and Wilkinson, B.J. (1992) Proline transport in *Staphylococcus aureus*: a high-
affinity system and a low-affinity system involved in osmoregulation. *J Bacteriol* **174**: 2702-
2710.
Vaish, M., Price-Whelan, A., Reyes-Robles, T., Liu, J., Jereen, A., Christie, S., Alonzo, F., 3rd, Benson,
M.A., Torres, V.J., and Krulwich, T.A. (2018) Roles of *Staphylococcus aureus* Mnh1 and
Mnh2 Antiporters in Salt Tolerance, Alkali Tolerance, and Pathogenesis. *J Bacteriol* **200**.
van den Berg, J., Boersma, A.J., and Poolman, B. (2017) Microorganisms maintain crowding
homeostasis. *Nat Rev Microbiol* **15**: 309-318.
Vijaranakul, U., Nadakavukaren, M.J., de Jonge, B.L., Wilkinson, B.J., and Jayaswal, R.K. (1995)
Increased cell size and shortened peptidoglycan interpeptide bridge of NaCl-stressed
Staphylococcus aureus and their reversal by glycine betaine. *J Bacteriol* **177**: 5116-5121.
Vijaranakul, U., Xiong, A., Lockwood, K., and Jayaswal, R.K. (1998) Cloning and nucleotide
sequencing of a *Staphylococcus aureus* gene encoding a branched-chain-amino-acid
transporter. *Appl Environ Microbiol* **64**: 763-767.
Vollmer, W., Blanot, D., and de Pedro, M.A. (2008) Peptidoglycan structure and architecture. *FEMS
Microbiol Rev* **32**: 149-167.

Wengender, P.A., and Miller, K.J. (1995) Identification of a PutP proline permease gene homolog
from *Staphylococcus aureus* by expression cloning of the high-affinity proline transport
system in Escherichia coli. *Appl Environ Microbiol* **61**: 252-259.
Wertheim, H.F., Melles, D.C., Vos, M.C., van Leeuwen, W., van Belkum, A., Verbrugh, H.A., and
Nouwen, J.L. (2005) The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet*
Infect Dis **5**: 751-762.
Wetzel, K.J., Bjorge, D., and Schwan, W.R. (2011) Mutational and transcriptional analyses of the
Staphylococcus aureus low-affinity proline transporter OpuD during *in vitro* growth and
infection of murine tissues. *FEMS Immunol Med Microbiol* **61**: 346-355.
Whatmore, A.M., and Reed, R.H. (1990) Determination of turgor pressure in Bacillus subtilis: a
possible role for K+ in turgor regulation. *J Gen Microbiol* **136**: 2521-2526.
Wood, J.M. (2011) Bacterial osmoregulation: a paradigm for the study of cellular homeostasis.
Annu Rev Microbiol **65**: 215-238.
Wyke, A.W. (1984) Isolation of five penicillin-binding proteins from *Staphylococcus aureus*. *FEMS*
Microbiology Letters **22**: 133-138.
Yao, J., Zhong, J., Fang, Y., Geisinger, E., Novick, R.P., and Lambowitz, A.M. (2006) Use of
targetrons to disrupt essential and nonessential genes in *Staphylococcus aureus* reveals
temperature sensitivity of Ll.LtrB group II intron splicing. *RNA* **12**: 1271-1281.
Zeden, M.S., Schuster, C.F., Bowman, L., Zhong, Q., Williams, H.D., and Gründling, A. (2018) Cyclic
di-adenosine monophosphate (c-di-AMP) is required for osmotic regulation in
Staphylococcus aureus but dispensable for viability in anaerobic conditions. *J Biol Chem*
293: 3180-3200.
984
985

987
988

Fig. 1. Different sets of genes are conditionally essential for the growth of *S. aureus* under NaCl,

KCl or sucrose stress conditions.

A. Workflow of the TN-seq experiment. A TN-seq library was pre-cultured and then used to

inoculate LB, LB containing 0.5 M NaCl, 0.5 M KCl or 1 M sucrose. Cells were kept in exponential

phase for 16 generations, harvested and transposon insertion sites determined by high

throughput sequencing.

B. Growth curves. For the TN-seq experiment (n=1), the growth of an *S. aureus* library cultured in

LB, LB 0.5M NaCl, LB 0.5 M KCl or LB 1.0 M sucrose medium was followed by taking OD$_{600}$

measurements at timed intervals. The cultures were back-diluted once when they reached an

OD_{600} of approximately 0.3. The plotted OD_{600} values were calculated by multiplying the measured

OD_{600} with the dilution factor. The dotted line indicates the 16 generation cut-off at which cultures

were harvested.

C. Circular plot showing the transposon insertion density along the *S. aureus* genome under

different osmotic stress conditions. The two outer rings depict genes located on the (+) or (-)

strand in *S. aureus* strain USA300 FPR3757. The inner four rings show the histograms of

transposon insertions on a per gene basis after growth of the library in LB (orange), 0.5 M NaCl

(brown), 0.5 M KCl (blue) and 1 M sucrose (pink) medium for 16 generations.

D. Volcano plots of conditionally essential and dispensable genes following growth of *S. aureus* in

0.5 M NaCl, 0.5 M KCl or 1 M sucrose medium and compared to LB conditions. q-values represent

Benjamini-Hochberg corrected p-values, and log_2 (fold changes) indicate essential genes (negative

values) or dispensable genes (positive values). The black horizontal line indicates a q-value of 0.05,

which was deemed the significance level. Vertical lines indicate 2- (green), 5- (orange) or 10- (red)

fold changes in either direction. Each dot represents one gene and coloring follows the fold-

change scheme whenever the q-value threshold was met. Very small q-values were truncated to

fit onto the graph. The TN-seq experiments were conducted once (n=1).

E. Venn diagrams showing overlaps of conditionally essential and dispensable genes during

prolonged NaCl, KCl or sucrose stress. Genes with a 2-fold decrease (essential genes) or a 2-fold

increase (dispensable genes) in transposon insertions under each stress condition compared to the

LB condition and a q-value of ≤0.05 were determined and the overlap between these gene lists

displayed in Venn diagrams.

1019

Fig. 2. TN-Seq screen reveals essential and dispensable genes during prolonged NaCl stress in *S.*
***aureus*.**

A. Workflow of TN-seq experiments. A saturated transposon library was pre-cultured for one hour,

then used to inoculate LB medium containing either normal levels (0.086 M) or 0.5 M of NaCl.

After 17 generations, cells were harvested, and transposon insertion sites determined by high

throughput sequencing.

B. Growth curve. Growth of the *S. aureus* library culture in LB and LB 0.5 M medium was followed

by determining OD_{600} readings. Dotted line indicates the 17 generation threshold. Cultures were

back-diluted once when they reached an OD of approximately 0.3 and the optical densities shown

are calculated from measured ODs time dilution factor. Culture from replicate B (n=1) is shown

and a similar growth profile was seen for replicate A.

C. Volcano plots of essential and dispensable genes in NaCl conditions compared to LB. Negative

$\log_2$ fold-changes indicate essential and positive $\log_2$ fold-changes dispensable genes. q-value

stands for Benjamini-Hochberg false discovery rate. Black horizontal line indicates q-value of 0.05

(cut-off) and red, orange and green vertical lines 10-, 5- and 2-fold differences in either direction.

Colored circles indicate genes for which a significant change in the number of transposon

insertions was observed above q-value cutoff and follow the same color code as the vertical lines.

Very small q-values were truncated to fit onto the graph. Data from replicate A are shown in the

left panel and from replicate B in the right panel (n=1 for each plot).

1041

Fig. 3. Growth curves and complementation analysis of *S. aureus* strains with transposon

insertions in salt essential genes.

A-B. *S. aureus* strains JE2 (WT), *SAUSA300_0694::Tn* mutant with empty plasmid and

complementation strain (A) or JE2 (WT), *SAUSA300_0910* (*mgtE*), *SAUSA300_0957::Tn* with empty

plasmid (B) and respective complementation strains were grown in LB (left panels) or LB 0.5 M

NaCl medium (right panels) supplemented with 100 ng/ml Atet and OD_{600} readings determined at

timed intervals. Growth curves were performed in triplicates (n=3) and means and SDs of OD_{600}

readings were plotted.

1050

1051
1052

Fig. 4. Gene *SAUSA300_0957* is located in a genomic region with cell envelope genes and its

deletion leads to salt sensitivity and other phenotypic changes.

A. Growth curves. *S. aureus* strains LAC* piTET (WT), LAC*Δ*957* piTET (Δ*957*) and the

complementation strain LAC*Δ*957* piTET-*957* (compl.) were grown in LB or LB containing 0.5 M

NaCl medium and 100 ng/ml Atet. The growth was monitored by determining OD_{600} readings the

means and SDs from three independent experiments were plotted.

B. Schematic of the *S. aureus* USA300 FPR3757 chromosomal region with gene *SAUSA300_0957*

(gene *957*). Putative promoters are shown as angled arrows (adapted from (Mäder *et al.*, 2016))

and black bars indicate the corresponding transcripts.

C. Determination of CFU/OD ratios. OD_{600} values as well as CFU/ml were determined for overnight

cultures of the *S. aureus* strains described in panel B and the means and SDs of the CFU per ml

OD_{600} of 1 from five independent experiments were plotted.

D. Cell size measurements. The cell walls of the *S. aureus* strains described in panel B were stained

with fluorescently labelled vancomycin and the cells subsequently observed under a microscope.

The diameters of 200 cells were measured and the means calculated. This average of the means

and SDs from five independent experiments were plotted.

For statistical analysis, a Kruskal-Wallis one-way ANOVA test was performed followed by a Dunn's

post-hoc test to determine p-values. Asterisks (*) indicate $p \leq 0.05$ and two asterisks (**) $p \leq 0.01$

and ns=not significant.

1072

1073

Fig. 5. Inactivation of *gene 957* does not significantly alter WTA levels or autolysis but leads to changes in oxacillin resistance.

A. Growth curves. *S. aureus* strains LAC* (WT), LAC*Δ957 (Δ957), LAC*ΔlcpB, (Δ957), and LAC*ΔtagO (ΔtagO) were grown in LB 0.5 M NaCl medium and OD_{600} readings determined at timed intervals. The means and SDs from three biological replicates were plotted.

B. Detection of WTA on silver stained gels. WTA was isolated from *S. aureus* strains described in panel A following growth in LB 0.4 M NaCl medium. The WTA was separated by electrophoresis and visualized by silver stain. The experiment was performed three times and one representative gel image is shown.

C. Detection of WTA on silver stained gels. Same as panel B but using the strains LAC* piTET (WT), LAC*Δ957 piTET (Δ957), and the complementation strain LAC*Δ957 piTET-957 (compl.) grown in LB 0.4 M NaCl medium also containing 100 ng/ml Atet.

D. Zymogram gel. Cell extracts prepared from *S. aureus* strains described in panel C were separated on a gel containing heat killed *Micrococcus luteus* cells. Autolysins were renatured and

the zones of lysis visualized by methylene blue staining. The experiment was performed twice and

one experiment is shown.

E. Determination of oxacillin MICs. Oxacillin MICs were determined for *S. aureus* strains described

in panel C using Etest strips. The median and SDs from four biological replicates were plotted.

For statistical analysis, a Kruskal-Wallis one-way ANOVA test was performed followed by a Dunn's

post-hoc test to determine p-values. Asterisks (*) indicate $p \leq 0.05$ and ns=not significant.

1094

1095

Fig. 6. Inactivation of *gene 957* leads to an increase in peptidoglycan crosslinking.

A-B. Muropeptide quantification. *S. aureus* strains LAC* piTET (WT), LAC*Δ957 piTET (Δ957), and

the complementation strain LAC*Δ957 piTET-957 (compl.) were grown in either LB (A) or LB 0.4 M

NaCl (B) containing 100 ng/ml Atet. After reaching late-exponential phase, the peptidoglycan was

isolated, digested and the muropeptides separated by HPLC. The areas under the peaks from the

chromatograms (representative chromatograms are shown in Supplementary Fig. S8) were

quantified for each mono-, di- or multimer. Bars represent means, and error bars standard

deviations from three independent experiments. For statistical analyses, two-way ANOVAs were

performed, followed by Dunnett's post-hoc tests using the WT as a comparison. The differences in

(A) were not significant. For (B), one asterisk (*) indicates p≤0.05, two asterisks (**) p≤0.01 and

four asterisks (****) p≤0.0001.

1107

Fig. 7. The growth and peptidoglycan defect observed for the *957* mutant can be rescued by

compensatory mutations in *pbp2*.

A. Growth curves. *S. aureus* strains LAC* (WT), LAC*Δ*957* (Δ*957*) and the LAC*Δ*957* suppressors

S1-10 (S1 to 10) were grown in LB 0.5 M NaCl medium, OD_{600} readings determined and the means

and SDs of four biological replicates plotted.

B. Schematic of Pbp2 with amino acid substitutions identified. Top: Structure of the *S. aureus*

penicillin binding protein Pbp2 (PDB: 3DWK, (Lovering *et al.*, 2008)) with amino acids that are

altered in the obtained suppressor strains shown in red. Bottom: Schematic of the Pbp2 enzyme

with the transglycosylase and transpeptidase domains as well as the observed amino acid changes

in the transglycosylase domain indicated.

C. Growth curves using *957* mutant strains containing *pbp2* alleles from suppressors S2 and S4.

Growth curves were performed in LB 0.5 M NaCl medium and plotted as described in panel A but

using *S. aureus* strains LAC* *1332*::Tn (WT), LAC*Δ*957 1332*::Tn (Δ*957*), LAC*Δ*957 1332*::Tn *pbp2*

SNP S2 (Δ*957 pbp2* SNP S2), LAC*Δ*957* suppressor S4 *1332*::Tn (Δ*957 pbp2* SNP S4).

D. Growth curves using *957* suppressors with their *pbp2* gene repaired to WT. Growth curves were

performed in LB 0.5 M NaCl medium and plotted as described in panel A but using *S. aureus*

strains LAC* *1332*::Tn (WT), LAC*Δ*957 1332*::Tn (Δ*957*), LAC*Δ*957* suppressor S2 *1332*::Tn (S2),

LAC*Δ*957* suppressor S4 *1332*::Tn (S4), LAC*Δ*957* suppressor S2 *1332*::Tn repaired WT *pbp2* (S2

WT *pbp2*), and LAC*Δ*957* suppressor S4 *1332*::Tn repaired WT *pbp2* (S4 WT *pbp2*).

E. Growth curves using WT strains carrying *pbp2* SNPs. *S. aureus* strains LAC* *1332*::Tn (WT), LAC*

1332::Tn *pbp2* SNP S2 (WT *pbp2* SNP S2) and LAC* *1332*::Tn *pbp2* SNP S4 (WT *pbp2* SNP S4) were

grown in LB 0.5 M NaCl medium and OD_{600} readings determined and the means and SDs of three

biological replicates plotted.

1131

1132

Fig. 8. The Δ957 suppressor strains show decreased peptidoglycan crosslinking compared the

Δ957 mutant.

A. Bacterial growth on moenomycin supplemented agar plates. *S. aureus* strains LAC* (WT),

LAC*Δ957 (Δ957), LAC*Δ957 Suppressor S2 (S2) and S4 LAC*Δ957 Suppressor S4 (S4) were grown

to exponential phase, normalized to an OD_{600} of 0.1 and plated either neat (n) or in 10-fold

dilutions onto LB agar containing 0.5 M NaCl without (top) or with (bottom) moenomycin. Shown

is one representative image of three biological replicates.

B. Muropeptide analysis. *S. aureus* strains LAC*Δ957 (Δ957), LAC*Δ957 Suppressor S2 (S2) and S4

LAC*Δ957 Suppressor S4 (S4) were grown in 0.4 M NaCl LB medium and muropeptides quantified

as described in Fig. 5. Representative chromatograms are shown in Supplementary Fig. S10D. The

means and SDs from three biological replicates were plotted. For statistical analysis a two-way

ANOVA and Dunnett's post-hoc test was performed. One asterisk (*) indicates $p \leq 0.05$, two

asterisks (**) $p \leq 0.01$ and four asterisks (****) $p \leq 0.0001$.

1146