High-throughput transposon sequencing highlights cell wall as an important barrier for osmotic stress in methicillin resistant *Staphylococcus aureus* and underlines a tailored response to different osmotic stressors

Short Title: Global analysis of osmotic stress responses in Staphylococcus aureus

Christopher F. Schuster[a], David M. Wiedemann[a], Freja C. M. Kirsebom[a,1], Marina Santiago[b,2], Suzanne Walker[b] and Angelika Gründling[a,3]

ORCID IDs: 0000-0002-4566-7226 (CFS), 0000-0001-6961-0980 (DMW), 0000-0003-4585-319X (FCK), 0000-0002-5446-3452 (MS), 0000-0002-0545-914X (SW), 0000-0002-6235-8687 (AG)

[a]Section of Microbiology and MRC Centre for Molecular Bacteriology and Infection, Imperial College London, London, SW7 2AZ, U.K.

[b]Department of Microbiology and Immunobiology, Harvard Medical School, Boston, MA 02115, U.S.A.

[1]Current address: Section of Respiratory Infections, National Heart and Lung Institute, Imperial College London, London, W2 1PG, U.K.

[3]To whom correspondence should be addressed. Email: a.grundling@imperial.ac.uk

Keywords: osmotic stress, TN-seq, Tradis, NaCl, KCl, Sucrose, *S. aureus*, DUF25380, Pbp2

Abstract

Staphylococcus aureus is an opportunistic pathogen that causes a variety of diseases. It presents a

problem in hospitals as well as communities partly due to the acquisition of multiple antibiotic

resistances, which make infections difficult to treat. *S. aureus* is also a frequent cause of foodborne

illnesses due to its ability to produce heat stabile toxins that cause nausea, vomiting and diarrhoea

even in the absence of living cells. One contributing factor for the food association is its high salt

tolerance, which allows this organism to survive commonly used methods of food preservation. How

this resistance is mediated is poorly understood. In this study, we used TN-seq-based high

throughput screens to find genes that are involved in the salt tolerance of *S. aureus* and identified

the previously uncharacterized DUF2538 domain containing gene *SAUSA300_0957* (gene *957*) as

essential under salt stress. Further experiments revealed that a *957* mutant strain is less susceptible

to oxacillin and shows increased peptidoglycan crosslinking. The salt sensitivity phenotype could be

suppressed by point mutations in the transglycosylase domain of the penicillin binding protein gene

pbp2, and these mutations also restored the peptidoglycan crosslinking to WT levels. These results

indicate that increased crosslinking of the peptidoglycan can be detrimental and highlight the role

of the bacterial cell wall for osmotic stress resistance. To gain more information on the general

osmotic stress response of *S. aureus* and how responses differ between different osmotic stressors,

additional TN-seq studies were performed with KCl and sucrose. Although it is generally assumed

that a generic osmotic stress response exists, our results revealed distinctly different long-term

responses to NaCl, KCl and sucrose stress. Using a global and genome-wide TN-seq approach, we

were able to link numerous previously unknown factors to the osmotic stress response in *S. aureus*.

This study will also serve as a starting point for future research in osmotic stress and might help us

develop strategies to tackle foodborne staphylococcal infections.

Introduction

Staphylococcus aureus is a Gram-positive bacterium that is carried by about 20% of healthy individuals [1]. It is also an opportunistic pathogen causing a variety of diseases, including bacteremia, endocarditis, soft tissue infections and food poisoning [2]. One characteristic of *S. aureus* is its ability to grow in the presence of a very high salt concentration up to 2.5 M, conditions under which other bacteria are unable to survive [3], and this has been used extensively for the isolation of staphylococci. This salt tolerance also allows *S. aureus* to thrive in its niches such as the human nares, skin [1] and certain food products [4].

In general, the bacterial cytoplasm is hyperosmotic to the environment and can therefore tolerate a certain level of external osmotic stress. However, an increase in the intracellular sodium concentration can lead to mis- and unfolding of proteins and inhibit enzymatic reactions. Hence, *S. aureus* must possess strategies to counter the effects of osmotic damage, but these are poorly understood, as most work has been performed in *Escherichia coli* and *Bacillus subtilis*, which possess much lower salt tolerances than Staphylococci.

General mechanisms of osmotic stress mitigation are the rapid import of potassium into the cell to reduce the influx of sodium and efflux of water [5-7]. However, during prolonged NaCl stress, potassium levels decrease again after its initial accumulation in the cytoplasm of *S. aureus* [8, 9]. Instead, *S. aureus* accumulates small osmotically active compounds, named compatible solutes [3]. The most effective compatible solutes in *S. aureus* are glycine betaine and proline [3, 10-14], which are imported by specific osmolyte transporters [13, 14]. These preferred osmolytes can accumulate in the cell to very high concentrations without negatively affecting cellular processes [6]. In addition, *S. aureus* can also synthesize compatible solutes *de novo*, as in the case of glutamine [12], but this process is much slower than the uptake of osmolytes and requires more energy.

Consistent with these observations, previous research studying the underlying genetic factors

of the *S. aureus* osmotic stress response implicated several osmolyte transporters in this process

[15-19]. These include the potassium transport systems Kdp and Ktr [15], the proline transporter

PutP [17, 19], the arsenic transport system Ars [16] and the branched chain amino acid uptake

system BrnQ [18]. In addition, although currently not experimentally verified, it can be assumed that

the main glycine betaine uptake system OpuD [20, 21] is an important factor for osmotic adaptation

as glycine betaine uptake reduces sensitivity to NaCl exposure significantly [10].

Despite these multiple countermeasures, exposure of *S. aureus* to high salt concentrations also

leads to morphological changes, including the formation of larger cells [22] and a thickening of the

cell wall [23]. However, if and which genetic factors are required for those processes are currently

not well understood. The cell wall of Gram-positive bacteria is an important barrier against osmotic

stresses and acts as a counterpart to the pressurized cytoplasm. A major component of the cell wall

is peptidoglycan, a polymer consisting of chains of repeating N-acetylglucosamine (GlcNAc) and N-

acetylmuramic acid (MurNAc) units that are crosslinked with neighboring chains via short peptides

and, in the case of *S. aureus,* pentaglycine cross-bridges (Reviewed by [24]). In the past, the

contribution of the cell wall towards osmotic stress resistance was often not considered in the

bacterial osmotic stress resistance field, presumably because the integrity of the cell wall was

assumed to be a prerequisite rather than an active adaptation to survive osmotic stresses.

Nevertheless, several possible changes to the cell wall have been observed *in S. aureus* during NaCl

stress, such as shortened interpeptide bridges in the peptidoglycan [22], an increase in resistance

to methicillin [25] and increased autolysis activity [26, 27]. In addition to peptidoglycan, other

components within the cell envelope, such as teichoic acids and more specifically their modifications

with D-alanine [28] and sugar residues [29, 30] are affected by osmotic stress. Combined, these

findings indicate that the integrity of the cell wall is an important requirement for osmotic

resistance.

Most studies on osmotic stress in *S. aureus* have been conducted with NaCl as osmotic stressor.

The ionic properties of NaCl have important implications on protein stability and activity as its

accumulation can lead to denaturation of proteins. A few studies in *S. aureus* have also been

performed with non-ionic osmotic stressors such as sucrose, sorbitol glycerol and amino acids [17,

31-33]. These different stresses are often used synonymously with osmotic stress, in part influenced

by the findings that the initial mitigation of osmotic stress by either salts or sugars can be prevented

by the accumulation of potassium and compatible solutes. However, there are potential differences

in the long-term adaptation to ionic and non-ionic osmotic stressors, which have not been

investigated in detail up to date. It is currently also not known if adaptations besides the

accumulation of potassium and osmolytes are similar or differ depending on the osmotic stressor,

and this was addressed as part of this study.

Using a transposon insertion sequencing (TN-seq) method [34], we determined on a whole

genome level genes that are essential or dispensable during long-term osmotic stress caused by the

exposure to NaCl. Several unknown and understudied genes were identified as essential under salt

stress including *SAUSA300_0957,* coding for a cytoplasmic protein, which we show here is important

for peptidoglycan homeostasis in *S. aureus*. Another protein, which we identified as important

during salt stress and essential for peptidoglycan synthesis, is the penicillin binding protein Pbp2,

again highlighting a key function of peptidoglycan during NaCl-induced salt stress. To determine

how the response varies depending on the osmolyte used, TN-seq screens were also performed in

the presence of KCl and sucrose. This yielded very different sets of essential and dispensable

candidate genes, providing experimental evidence of the distinct nature of these stresses and linking

a number of previously unknown factors to the osmotic stress response in *S. aureus*.

Results

TN-seq analysis identifies *S. aureus* genes involved in NaCl stress

To identify genes involved in the *S. aureus* salt stress response, two independent transposon sequencing (TN-seq) screens were conducted at different salt concentrations and generation times (Figure 1A). A multiplexed transposon library, generated using 6 different transposons possessing different or no promoters [34, 35], was used in replicate A and an independent Tn-library using a transposon without an outward facing promoter was used in replicate B. In both experiments, LB medium with either the normal salt concentration of 85 mM NaCl (LBN), LB medium with extra 0.25M NaCl or LB medium with extra 0.5 M NaCl was used (See Supplementary Table S1 for a list of all TN-seq experiments). Monitoring bacterial growth over seventeen generations, revealed that growth in LB 0.25M NaCl was comparable to that of cells grown in LBN (Figure 1B), whereas a strong reduction in growth was seen in the presence of 0.5M NaCl (Figure 1B). For the TN-seq analysis, samples were taken after 10 and 17 doublings. As expected, the Tn insertions were distributed evenly throughout the genome when the bacteria were propagated in LBN but became more unbalanced at higher salt concentrations and longer generation times due to the enrichment of better salt-adapted TN-strains (Supplementary Figure 1). The fold-change of transposon insertions per gene under the salt stress conditions compared to the LBN condition (input library) as well as q-values (false discovery rate according to Benjamini-Hochberg) were determined for each gene (Supplementary Table S2 & 3). The data were also plotted as volcano plots (Figure 1C), in which negative values on the x-axis indicate genes that are essential under high salt conditions (fewer TN insertions in high salt), whereas positive values indicate dispensable genes, for which an inactivation might be beneficial under high salt conditions. For replicate A, after 10 generations in the presence of 0.25M NaCl the number of TN insertions per gene was not drastically altered compared to LBN (Figure 1C, top row). An increase in the salt concentration with the same number of generations

slightly increased the number of essential and dispensable genes, but the number was still quite low

(Figure 1C, top row). Extending the growth to 17 generations (Figure 1C, second row) increased the

number of genes with significant differences in the number of TN-insertions during growth at 0.25

M NaCl and even more so at 0.5 M NaCl, indicating a much more stringent depletion or enrichment

of strains with TN-insertions in specific genes from the input library. Therefore, the TN-seq

experiment for replicate B was only performed with 17 generation samples (Figure 1C, bottom row;

Supplementary Table 3). Lists of the top common shared genes between conditions were created

(Supplementary Tables S4-7) and the reproducibility between experiments was also assessed by a

principal component analysis (PCA) (Figure 1D). For the 17 generation data sets, a clear difference

between cells grown in either 0.25 M (orange ellipse) or 0.5 M NaCl (red ellipse) was observable and

experimental data points clustered together for each condition. The 0.25 and 0.5 M NaCl data from

the 10 generation experiment, which was only conducted once, clustered both closely with the 17

generation 0.25 M NaCl samples, consistent with considerably fewer changes in depleted or

enriched TN-strains under these conditions as compared to growth for 17 generations in LB 0.5 M

NaCl. From this analysis we concluded that growing *S. aureus* for 17 generations in 0.5 M NaCl

medium provides sufficiently stringent conditions to identify genes that are either essential or

dispensable in high salt conditions and our further analysis focused on these data sets.

137   The TN-seq screens highlight the essentiality of transporter and cell envelope related genes and

138   the dispensability of respiration genes under high salt conditions

To provide a visual and genome wide guide to the cellular functions of salt essential or dispensable

genes, Voronoi maps were generated for the 17 generation 0.5 M NaCl datasets [36, 37]. For this

global visualization, the observed decrease (essential genes) or increase (dispensable genes) in Tn

insertions per gene compared to LBN medium were plotted regardless of their p-value

(Supplementary Figures 2 and 3). In both replicates, genes coding for transporters and cell envelope

related genes were the most prominent conditionally salt essential genes (Supplementary Figure 2).

More specifically, the TN-Seq data indicated that inactivation of the putative magnesium

transporter MgtE [38] and the putative D-serine/D-alanine/glycine transporter AapA is detrimental

under salt stress. Inactivation of AapA has previously been shown to suppress the temperature

sensitivity phenotype seen in an *S. aureus fmtC/mprF* mutant [39] and to lead to increased

amoxacillin and daptomycin sensitivity [35, 40]. In addition, the *dlt* operon, coding for enzymes

required for the D-alanylation of teichoic acids, an important factor for cationic peptide resistance

[41], as well as the stationary and stress sigma factor gene *sigB* were found to be essential.

An even larger number of genes was identified as dispensable during salt stress (Supplementary

Figure 3). Inactivating or reducing the expression of such genes is expected to help *S. aureus* survive

under salt stress. Among these dispensable genes were genes coding for the sodium antiporter

Mnh2, which has previously been shown to transport Na^+/H^+ and K^+/H^+ in high pH medium and to

play an important role in cytoplasmic pH maintenance [42]. Strikingly, a large number of genes

involved in respiration, such as the *cta/qox/men/hem* operons as well as genes involved indirectly

in respiration through quinone synthesis (shikimate pathway, *aro* operon) were also identified as

dispensable during salt stress. Curiously, the data also reproducibly indicated that genes coding for

the phosphodiesterases GdpP and Pde2 involved in degradation of the signaling molecule c-di-AMP

are dispensable under high salt conditions. Somewhat contradictory, the construction of a *dacA*

mutant coding for the cognate c-di-AMP cyclase DacA was only possible at high salt concentrations

(or in defined chemical medium) [21]; but altogether these findings confirm the involvement of c-

di-AMP in salt mediated osmotic stress adaption in *S. aureus*.

Confirmation of genes essential during salt stress using individual mutants

To determine to what extent different genes identified in the TN-seq screen contribute to the salt resistance of *S. aureus*, a set of candidate genes from the top hits of the 17 generations, 0.5 M NaCl LB dataset B were chosen for further analysis. For this analysis we used defined transposon mutants available from the *S. aureus* NTML transposon mutant library [43] (Supplementary Table S8). Most of the chosen genes showed similar trends in dataset A, but intriguingly two of the tested genes, *rsbV* and *rsbW*, were flagged as highly dispensable in dataset A but as essential in dataset B. Since no strains with transposon insertions in the *dlt* operon were available in the NTLM transposon mutant library the contribution of these genes to the salt stress resistance was not further investigated.

Next, the growth of the different NTML transposon mutant strains was assessed in LBN and LB 0.5 M NaCl medium. In LBN medium, most strains grew similarly to the WT strain JE2, indicating that inactivation of the respective gene does not greatly affect bacterial growth in normal LB medium (Figures 2A-D, left panels). Exceptions were strains NE1109 (*sigB*) (Figure 2A), NE1778 (*lcpB*) (Figure 2B) and NE188 (*mfd*) (Figure 2C), cultures of which reached slightly lower final optical densities. At high osmolarity conditions (LB 0.5 M NaCl), all strains showed a growth defect compared to the WT strain (Figures 2A-D), validating TN-seq as a method to identify genes important in *S. aureus* during salt stress. Several strains exhibited strong growth defects in the presence of 0.5 M NaCl (Figures 2C and D) with strain NE1384 (*SAUSA300_0957*) showing extremely reduced growth (Figure 2D). To confirm that the salt sensitivity was mediated by the inactivation of the genes in questions, we constructed complementation plasmids for seven of the most promising candidates by either expressing the gene of interest from its native promoter or an anhydrotetracycline (Atet) inducible promoter in the transposon mutant strains. While no complementation was observed for four strains (Supplementary Figures 4A and B), the salt-dependent growth defect could be complemented for strains carrying mutations in *SAUSA300_0694* (Figure 3A), *SAUSA300_0910*

(*mgtE*) and *SAUSA300_0957* (Figure 3B). *SAUSA300_0694* encodes a hypothetical protein with 6

predicted transmembrane helices but no other identifiable domain motif. MgtE (SAUSA300_0910)

is a predicted magnesium transporter. *SAUSA300_0957* (from here on also referred to as gene *957*)

codes for a cytoplasmic protein of unknown function and was further investigated in this study

because a *957* mutant exhibited the strongest salt-sensitivity phenotype.

Genomic organization of *957* suggests a potential function in cell envelope homeostasis

SAUSA300_0957 encodes a protein with a DUF2538 domain of unknown function. Proteins with this

domain are found mainly in Gram-positive bacteria (Actinobacteria and Firmicutes) (based on an

Interpro search, https://www.ebi.ac.uk/interpro/entry/IPR024469). Although a crystal structure of

the 957 protein has been deposited in the PDB (3KBY), to our knowledge nothing is known about

the actual cellular function of this protein. Transcriptome data [44] suggest that gene *957* is in an

operon with *lcpB* (Figure 4A). LcpB is one of three wall teichoic acid (WTA) ligases present in *S.*

aureus that attaches WTA to the peptidoglycan polymer [45-47]. Inactivation of LcpB leads to slightly

lower WTA levels in the cell wall, whereas deletion of all three WT ligases completely abolishes WTA

anchoring to the cell wall [45]. In addition, LcpB cell wall ligase activity was demonstrated *in vitro*

[47]. *fmtA* is located upstream of the *957* operon and has been proposed to code for an esterase

that can remove D-alanine modifications from teichoic acids [48] and is involved in methicillin

resistance [49]. Genes coding for a predicted acetyltransferase and *atl* coding for the major *S. aureus*

autolysin are found immediately downstream of the *957* operon [50]. In this early work, gene *957*

was first described as an ORF of "unknown function". Since functionally related genes often cluster

together in bacteria, this prompted us to next look at cell-envelope related phenotypes for a *957*

deletion strain.

Deletion of gene *957* causes physiological changes leading to a smaller cell size

As all previous experiments were performed with transposon mutants, we first constructed strain LAC*Δ*957* with a marker-less in-frame deletion of gene *957*. Strain LAC*Δ*957* had the expected growth defect in the presence of 0.5 M NaCl, which could be complemented by expressing a functional copy of *957* (Figure 4B). We then proceeded to test if the *957* mutant displays other phenotypes and to investigate if gene *957* is involved in cell envelope homeostasis. First, WT LAC*, the *957* mutant and the complementation strain were grown in LBN and the ratio of colony forming units to optical density (CDU/OD) was determined. The CFU/OD ratio was significantly larger for the *957* mutant ($60.8 \pm 6.8 \times 10^7$ CFU/OD) mutant as compared to strain LAC* ($37.5 \pm 9.3 \times 10^7$ CFU/OD) when grown in LBN, suggesting a change in cell shape or size (Figure 4C). This phenotype was restored in the complementation strain ($37.3 \pm 3.6 \times 10^7$ CFU/OD) (Figure 4C). When the average cell diameter was determined by microscopy, a small but significant reduction in cell diameter was seen for the *957* mutant (1.25 ± 0.02 µm) as compared to the WT (1.36 ± 0.03 µm) and the cell size was restored to wild type levels in the complementation strain (1.35 ± 0.05 nm) (Figure 4D).

Salt sensitivity of *957* is not due to interaction with LcpB but likely due to changes in peptidoglycan crosslinking

Since the gene *957* is on the same transcript as *lcpB* coding for a WTA ligase (Figure 4A), the contribution of WTA to salt stress resistance as well as the contribution of *957* to the attachment of WTA to the cell wall was assessed. Initially the growth of a *tagO* (*tarO/llm*) mutant, a strain unable to produce WTA, was tested in LB and LB 0.5 M NaCl medium. When cells were grown in LBN, they grew normally (Supplementary Figure 5A), but under high salt conditions, the *tagO* mutant showed reduced growth, similar to that of the *957* mutant (Figure 5A). This is consistent with what has been reported previously for a *Staphylococcus epidermidis tagO* mutant [51] and indicates an important

role for WTA during salt stress. When the *lcpB* mutant was propagated in LBN medium, its growth was slightly inhibited (Supplementary Figure 5A). However, in high salt medium (Figure 5A) it grew unimpeded and like WT. Since the *lcpB* mutant did not show a growth defect in high salt medium, this made it unlikely that *957* is a regulator of LcpB activity. To further exclude that gene *957* could be somehow involved in the attachment of WTA to the peptidoglycan, we isolated WTA from strains grown in either LB or LB containing 0.4 M NaCl and separated the polymer on native polyacrylamide gels. The slightly lower concentration of salt was chosen because growth of the *957* and the *tagO* mutants was too strongly inhibited at 0.5 M NaCl. As expected the *tagO* mutant did not produce any WTA (Supplementary Figure 5B, Figure 5B) and consistent with previous studies [45, 46], the *lcpB* mutant exhibited slightly reduced WTA levels compared to all other strains (Supplementary Figure 5B, Figure 5B). There was no reduction of WTA observable in the *957* mutant compared to the WT strain in either LBN (Supplementary Figure 5B & 5C) or LB 0.4 M NaCl (Figure 5B & 5C) but rather a small, complementable increase, suggesting no influence of *957* over LcpB on WTA attachment.

Gene *957* is found upstream of the bi-functional autolysin gene *atl* and therefore we next investigated potential changes in autolytic activity using zymograms. Cell extracts from overnight cultures of the WT, the *957* mutant and the complementation strain were separated on polyacrylamide gels with embedded *Micrococcus luteus* cells and incubated overnight in buffer to re-fold the autolysins and enable the digestion of the peptidoglycan. A small increase in autolytic activity was detected in the *957* mutant (Figure 5D) for bands around 55 kDa and 70 kDa compared to the WT and complementation strain, indicating a possible change in Atl availability. We also tested the susceptibility of the *957* mutant towards the cell wall active beta-lactam antibiotic oxacillin. In contrast to our expectations, the *957* mutant exhibited a small but significant increase in MIC towards this antibiotic than the WT or the complementation strain (Figure 5E), indicating potential changes to the peptidoglycan structure. To test this, we next analyzed the muropeptide

profile of mutanolysin-digested peptidoglycan isolated from the WT, the *957* mutant and

complementation strain after growth in LBN or LB 0.4 M NaCl medium. Upon first inspection, no

peaks were absent in any condition or strain (Supplementary Figure 5D, Supplementary Figure 5E).

Probing further, mono- and multimers of muropeptides up to 7-mers (see Supplementary Figure 5D

for how binning was performed) were quantified but no significant changes were found when

strains were cultured in LBN medium (Supplementary Figure 5F). However, when the muropeptide

profiles from cells cultured in 0.4 M NaCl were compared, a significant decrease in the total amount

of di- and trimers and a significant increase in higher multimers was observed for the peptidoglycan

isolated from the *957* mutant (Figure 5F), suggesting that sensitivity to NaCl could potentially be

caused by higher crosslinking and rigidity of the peptidoglycan in the mutant.

Salt-resistant suppressors possess variations in the transglycosylase domain of Pbp2 and show

reduced peptidoglycan crosslinking

Next, we attempted to generate *957* suppressor strains that showed improved growth in the

presence of 0.5 M NaCl, with the idea that by mapping and investigating the compensatory

mutations, further insight into the cellular functions of protein 957 could be gained. We initially

attempted to raise suppressor strains on solid medium but failed to do so, because the growth

defect of the *957* mutant was not as prevalent on agar plates. We therefore used liquid medium

instead, where we readily obtained suppressor strains. These strains grew as well as or even better

than the WT LAC* strain in the high salt conditions (Figure 6A) and also grew similar to the WT in

LBN (Supplementary Figure 6A). The genome sequences of 10 independent suppressor strains were

determined and single nucleotide polymorphisms (SNPs) in one or two genes could be identified for

each strain (Supplementary table S9), except for one strain where the coverage was insufficient, and

this strain was omitted from further analysis. A common denominator for all suppressor strains were

SNPs in the *pbp2* gene, coding for the Penicillin binding protein 2 (Pbp2) (Figure 6B). *S. aureus* Pbp2

is a bifunctional enzyme [52, 53], which possesses transglycosylase and transpeptidase activity and

is involved in peptidoglycan synthesis. All nine discovered SNPs were unique and led to mutations

in the amino acid sequence in the transglycosylase domain but neither to stop codons, nor

frameshifts nor mutations in the transpeptidase domain. The mutations were mapped (Figure 6B)

onto an available Pbp2 structure (3DWK) [54] to see if a specific area of the transglycosylase domain

had been targeted, but the amino acid substitutions were found throughout the molecule and not

only in the active site. Interestingly, *pbp2* also appeared to be considerably less important for

growth at 0.5 M NaCl (Supplementary Figure 6B and 6C), although transposons were mainly

localized at the beginning and end of the gene or within the promoter region.

To confirm that the *pbp2* SNPs were indeed responsible for the suppression of the salt sensitivity of

the *957* mutant, we transferred the *pbp2* SNPs from two strains into a fresh LAC*Δ*957* background

by co-transduction of a transposon in a nearby gene (Supplementary Figure 6D, schematic). As

expected, these "recreated" suppressor strains showed also increased salt resistance (Figure 6C,

Supplementary Figure 7A). In addition, we repaired the *pbp2* gene in two suppressors to the WT

pbp2 allele and this reduced their ability to cope with NaCl stress, all consistent with the hypothesis

that the SNPs in *pbp2* are responsible for the suppression phenotype (Figure 6D, Supplementary

Figure 7B). When the same *pbp2* SNPs were transferred into a WT LAC* strain, a reproducible

growth improvement of the SNP-bearing strains was observed in high salt medium, an indication

that the *pbp2* mutations lead to a general growth improvement of *S. aureus* under NaCl stress

conditions (Figure 6E, Supplementary Figure 7C).

Moenomycin is a phosphoglycolipid antibiotic that inhibits the transglycosylase activity of the *S.*

aureus Pbp2 enzyme (reviewed in [55]). To test if a decrease in Pbp2 transglycosylase activity results

in the improved growth of the *957* mutant in high salt conditions, cells were grown on LB agar

containing 0.5 M NaCl with or without moenomycin. The growth behaviors of the WT strain and the

957 mutant were similar when grown on solid LB agar containing 0.5 M NaCl but lacking

moenomycin (Figure 6F, top panel). As stated above, this discrepancy in the growth behavior of the

957 mutant in high salt liquid versus solid medium was already noted when we attempted to

generate suppressors on agar plates. On the other hand, in medium supplemented with 0.02 µg/ml

moenomycin, we observed better growth of the *957* mutant compared to the WT strain (Figure 6F,

lower panel). In addition, the *957* mutant also showed improved growth compared to the two

suppressor strains S2 and S4 when moenomycin was added (Figure 6F, lower panel). These findings

are consistant with the idea that a partial inhibition of the glycosyltransferase activity of Pbp2 can

improve the growth of the *957* mutant in high salt medium.

Because the LAC*Δ*957* strain showed increased peptidoglycan crosslinking when compared

to a WT strain, we hypothesized that the peptidoglycan of the suppressor strains would be again

less crosslinked. To test this, the peptidoglycan from two of the suppressor strains grown in LB 0.4

M NaCl was isolated, the muropeptide profiles determined and compared to that of the original *957*

mutant (Figure 6G, Supplementary Figure 7D). As before, the muropeptide profiles looked similar

(Supplementary Figure 7D) but quantification revealed a significant reduction in crosslinked

peptidoglycan fragments and a significantly overrepresentation of monomeric and dimeric

fragments in the suppressor strains as compared to the original *957* mutant (Figure 6G). These

results indicate that the amount of crosslinking of the peptidoglycan polymer is an important factor

in the salt resistance of *S. aureus* and that dysregulation of peptidoglycan crosslinking can lead to

destabilizing effects.

Different types of osmotic stresses target different sets of genes

In the literature, osmotic stress is a loosely used term for any accumulation of osmolytes, however

the type of ion or osmolyte can potentially have a great impact how bacteria respond. To address

this issue, we then determined how the stress response between different commonly used

osmolytes compare. To this end, the highly saturated *S. aureus* transposon library with the

promoter-less transposon was propagated for 16 generations while challenged with either 0.5 M

NaCl, 0.5 M KCl or 1.0 M sucrose (Figure 7A). The molarity for sucrose was doubled, as the salts

dissociate into two molecules whereas the sucrose does not. The culture challenged with 0.5 M NaCl

grew the slowest, followed by 1.0 M sucrose, whereas the cells grown in 0.5 M KCl grew similarly to

the cells grown in LBN (Figure 7B). Of note, the optical density measurements of *S. aureus* in 1.0 M

sucrose proved to be difficult because the OD_{600} values dropped dramatically upon diluting cells into

this medium. In all likelihood, the exposure leads to shrinking of the cells and therefore reduced

absorbance, resulting in values too low to detect in the first two hours.

As expected by using a large and good quality transposon library, transposon insertions were found

under all conditions in genes throughout the whole genome (Figure 7C). Next, the number of TN-

insertions per gene following growth under the NaCl, KCl or sucrose stress condition was compared

to the number of TN-insertions per gene after growth in LBN medium (Supplementary table S10)

and in this manner conditionally essential (Supplementary table S11) and dispensable

(Supplementary table S12) genes identified. Fold-changes in transposon insertions per gene and q-

values were plotted in volcano plots (Figure 7D). From this, it was evident that under KCl stress the

number of essential and dispensable genes was much lower than for NaCl or sucrose stress,

indicating a much less severe effect of KCl on *S. aureus* cells. Sucrose stress also showed a reduced

set of essential and dispensable genes compared to NaCl stress, but considerably higher than under

the KCl condition. This was also reflected when inspecting the gene lists of the top 30 essential or

dispensable genes (Supplementary table S11-12) as in the KCl condition only 15 and 2 genes

respectively met the requirements of q-value ≤ 0.05 and a fold-change of 5. To identify common

genes between conditions, the fold-change stringency was relaxed to 2-fold and the overlap of

genes was visualized in Venn diagrams (Figure 7E, individual genes in Supplementary Table S13).

Three genes, namely *SAUSA300_0425* (*USA300HOU_0457*, *mpsA (nuoF)*, a cation transporter of the

respiratory chain [56]) *SAUSA300_0750* (*USA300HOU_0796*, *whiA*, a hypothetical protein, possibly

involved in cell wall synthesis) and SAUSA300_0846 (USA300HOU_0903, a possible sodium:proton antiporter) were essential in all conditions. In the case of dispensable genes, only one gene, *SAUSA300_1255* (*USA300HOU_1294*, *mprF/fmtC*), coding for a phosphatidylglycerol lysyltransferase involved in the defense against cationic microbial peptides, was identified in all conditions (Figure 7E, (Supplementary Table S12). Overall, the overlaps between conditions were small, indicating distinct modes of actions for each osmotic stress.

Next, in order to assess similarities and differences on a gene function and cellular pathway level, functional Voronoi maps were generated from the conditionally essential (Figure 8) and dispensable (Supplementary Figure 8) genes regardless of their p-values. These images highlighted the essentiality of the wall teichoic acid through *tagO* and the *dlt* operon (Figure 8A) as reported in the first two experiments for the 0.5 M NaCl condition and the essentiality of the transporters AapA and MgtE in both NaCl (Figure 8A) and KCl (Figure 8B) stress conditions. In the high sucrose condition, these transporters were much less prominent (Figure 8C). The patterns of essential genes suggested a similar stress response for the ionic stressors NaCl and KCl that was distinct from sucrose stress. When the dispensable genes were investigated (Supplementary Figure 8), the number and type of genes in the NaCl condition differed considerably from KCl and sucrose stress. Under NaCl stress (Supplementary Figure 8A), in addition to the penicillin binding protein *pbp2*, a large number of genes involved either directly in respiration or indirectly by their role in ubiquinone synthesis were found to be dispensable (Supplementary Figures 3A, 3B and 8A). The number of dispensable genes in the KCl (Supplementary Figure 8B) and sucrose (Supplementary Figure 8C) conditions were considerably less, indicating differences in the osmotic stress response between NaCl and KCl.

Verification of KCl sensitivity of conditionally essential genes identified by TN-seq

Next, from the pool of genes postulated to be essential under KCl stress, and also taking into account genes that overlapped between NaCl and KCl conditions (Figure 8A and B), mutants were selected

for growth analysis in medium containing 0.5 M KCl. This included strains NE736 (*SAUSA300_0910*,

mgtE), NE867 (*SAUSA300_0483*, *mazG*) and NE810 (*SAUSA300_1642*, *aapA*) (Figure 9). In addition,

NE1384 (SAUSA300_0957, *gene 957*) was also included. When grown in LB medium supplemented

with 0.5 M KCl, all strains showed a strong growth defect compared to the WT strain JE2 (Figure 9A,

LBN curves in Figure 2). The strongest growth defect was observed for the *mgtE* mutant strain

NE736. This is in line with our TN-seq data, as the *mgtE* gene was identified as the top candidate of

essential genes in the KCl condition (Supplementary table S11). MgtE is a putative

magnesium/cobalt transporter with a cystathionine-beta-synthase (CBS) domain that is thought to

regulate ion translocation [57] but not much is known in *S. aureus*. In previous studies, we

determined that the CBS domain of MgtE is not a c-di-AMP target [58] and we recently

demonstrated that MgtE is an important contributor to cobalt toxicity in *S. aureus* [38]. To verify the

KCl sensitivity, growth curves were repeated with a clean Δ*mgtE* mutant and complementation

strain. In LBN medium a slight, but complementable, growth defect was detectable for the *mgtE*

mutant but an almost complete growth arrest was seen in LB medium supplemented with 0.5 M KCl

(Figure 9B). Taken together, we present robust data on different osmotic stresses and our results

indicate that the osmotic stresses caused by NaCl, KCl and sucrose are distinct from each other.

Nevertheless, some genes such as *mgtE* play an important role in more than one condition and are

therefore valuable targets for upcoming studies.

Discussion

In this work, we successfully identified several salt tolerance genes in the USA300 *S. aureus* strain

and further characterized gene *957,* a gene without previously assigned function. Our results

indicate that the gene product is involved in cell envelope homeostasis but likely not WTA

attachment and we show that its absence leads to more crosslinked peptidoglycan. Suppressors of

a *957* mutant acquired mutations in *pbp2* and the increase in peptidoglycan crosslinking was

reversed in such stains. This demonstrates that the synthesis of the peptidoglycan layer is a tightly

regulated process that plays an essential role during osmotic stress. In addition, we tested other

osmotic stresses and performed TN-seq screens and could show that each osmotic stressor affects

a defined but different set of genes. It is therefore important to use the term osmotic stress

carefully, as our results highlight that there is not a general osmotic stress response but rather

responses tailored to the individual osmolyte.

TN-seq experiments provide a wealth of information about the growth ability of single mutants in a

mixed population and can therefore quickly and accurately determine genes that are required under

certain growth condition. It is however always a concern that a single TN-seq experiment might not

be reproducible due to stochastic extinction of individual mutants or certain growth dynamics. In

this study, we performed three independent NaCl TN-seq experiments using two different libraries

and slightly different experimental setups. Although there were differences between experiments,

overall the data (0.5 M NaCl LB, 17 generations) proved to be rather consistent between different

runs (Figure 1D) suggesting that a single experiment would be sufficient in most cases.

Because TN-seq relies on the selection of strains adapted to certain growth conditions, competition

between mutants and subsequent depletion of the input library it is to be expected that this method

is ideal for exploring the long-term effects on growth instead of quick, short-term response to

stresses. To evaluate how our TN-seq results compared to previous results from high-throughput

studies of *S. aureus* cells grown in high salt conditions, we determined the overlaps between our 0.5 M NaCl, 17 generation datasets and a microarray data set acquired from cells grown for ~6 generations in 2 M NaCl [15]. To compare the datasets, the *S. aureus* strain COL locus tags used in the microarray study were first converted to TCH1516 locus tags where possible (Supplementary Table S14) and afterwards compared with the first two 0.5 M NaCl, 17 generation TN-seq datasets. The overlap between the downregulated (microarray) and dispensable (TN-seq) genes was minimal in both replicates (1 and 7) and also very small for the upregulated and essential genes (both 14). Three operons were consistently essential or upregulated between all datasets: the *cap5*, *SAUSA300_0771-2*, and the *sda* operon. The *cap5* genes are involved in capsule production [59] and this type of capsule has been previously reported to be induced by the addition of salt [60]. Genes *SAUSA300_0771-2* are hypothetical membrane proteins and presumably part of a threonine/serine exporter (Interpro database IDs: IPR010619 and IPR024528) and the *sda* genes are annotated as L-serine dehydratase components and a regulator protein. Taken together, the low overlap between the microarray and TN-seq data (Supplementary Table S14) further highlights that in our screen we have identified a number of previously unknown salt tolerance genes.

Most mutants of potential NaCl essential genes identified in the TN-seq screen proved to be sensitive to 0.5 M NaCl (Figure 2A-D), confirming the effectiveness of the TN-seq screen. In particular, previously uncharacterized genes such as *SAUSA300_0694*, *mgtE,* and the *957* gene could be identified as essential during NaCl stress and their phenotypes could be complemented (Figure 3). In our study, we opted not to experimentally validate any of the dispensable genes, as the verification of a growth advantage is more difficult to prove than a growth defect, but these genes will provide interesting starting points for future research. In addition, it will be interesting to investigate higher concentrations of NaCl to see if the same or different genes are flagged as essential and dispensable.

Gene *957* was selected for further characterization as part of this study, because it had previously not been linked to NaCl stress and had not been investigated before. As gene *957* and *lcpB* are co-transcribed we initially assumed that protein 957 is involved in WTA attachment via LcpB but we were unable to demonstrate such a link. Instead the *957* mutant exhibited increased muropeptide crosslinking, which could be suppressed by certain *pbp2* SNPs suggesting involvement in peptidoglycan synthesis. We can only speculate if the SNPs identified in *pbp2* increase or inhibit the activity of Pbp2. However, judging by the number of independent SNPs, it seems more likely that the mutations lead to a decrease rather than an increase in transglycosylase activity. In addition, the results from the moenomycin sensitivity experiments using a sublethal antibiotic concentration (Figure 6F) highlights that partial inactivation of the transglycosylase activity of Pbp2 improves the growth of the *957* strain in high salt conditions. This is consistent with the idea that the obtained *pbp2* SNPs lead to a decrease in transglycolylase activity. The absence of frameshift or non-sense mutations can be explained by the importance of the C-terminal transpeptidase domain of *pbp2* [61] and this is reflected in the transposon insertion distributions (Supplementary Figure 6B & C). Intriguingly, the suppressor mutants show reduced growth compared to the *957* mutant on moenomycin plates, presumably due to transglycosylation activity reduced to levels that are detrimental to the cell. At this point it is also unclear how mutations in the transglycosylase domain alter the muropeptide pattern, since the peptide bonds are made by the transpeptidase, not the transglycosylase domain. We hypothesize that a decrease in the efficiency of the glycosylation activity and slowing down the glycan chain synthesis process, will allow reduce the efficiency of the subsequent transpeptidation process, resulting in decreased peptidoglycan crosslinking in the *pbp2* SNP strains. It is also noteworthy that the *pbp2* gene becomes dispensable under NaCl conditions (Supplementary Figure 8A), an additional indicator that Pbp2 activity needs to be avoided under high salt conditions.

Interestingly, the *957* mutant exhibited a slightly higher oxacillin MIC than the WT. This is in contrast

to what we expected as MICs are determined in 2% (0.342 M) salt medium and this should lead to

a growth disadvantage of the mutant. However, when seen in light of the increase in peptidoglycan

crosslinking, this could explain the growth advantage of the *957* mutant in the presence of this beta-

lactam antibiotic, as this could counter the inhibition of transpeptidases by oxacillin.

Although we were unable to determine the exact molecular mechanism, it is clear from our data

that gene *957* is involved in cell wall synthesis either directly or indirectly. More generally, our work

suggests that the rigidity of the peptidoglycan cell wall is an important factor to counter osmotic

stresses and it is conceivable that the more crosslinked cell wall of the *957* mutant is less elastic and

therefore easier to break under strong osmotic stress caused by NaCl exposure (Summarized in

Figure 10).

From our TN-seq data and the use of different osmotic stressors (NaCl, KCl and sucrose), it is evident

that although all these compounds exert osmotic stress, the response differs greatly between them

(Figure 7, Figure 8 and Supplementary Figure 8). NaCl and KCl conditions share a set of genes that

become essential such as the transporters *aapA* and *mgtE* (Figure 8A and B) but differ in the set of

dispensable genes (Supplementary Figure 8A and B). It is intriguing that a whole set of respiration

related genes become dispensable under NaCl stress but not under KCl stress. This could indicate

that respiration should be inhibited or slowed down in high NaCl conditions, possibly by interference

with sodium pumps a process less likely to be inhibited by potassium ions. The set of essential genes

under sucrose conditions (Figure 8C) was vastly different from that of the NaCl and KCl conditions

whereas the set of sucrose dispensable genes (Supplementary Figure 8C) looked similar to the KCl

set. This highlights the differences in osmotic stress adaptations and serves as a reminder that the

type of stressor remains important. Therefore, although the term osmotic pressure is useful in some

contexts, it often needs to be taken with a grain of salt.

Using our TN-seq approach, we were able to identify another important factor, the predicted

magnesium transporter MgtE, as essential during KCl, as well as NaCl stress. From experiments

performed in *E. coli*, it can be inferred that MgtE transports Mg^{2+} ions from the outside of the cell

into the cytoplasm [57]. In our previous work, we have shown that the addition of Co^{2+} to the growth

medium of a MgtE-carrying *S. aureus* strain leads to a growth defect that can be rescued by deletion

of *mgtE* [38], which is in line with cation import by MgtE. Using *in vitro* assays, it was shown for the

E. coli MgtE transporter that the addition of K^+ or Na^+ does not inhibit Mg^{2+} transport, which means

that MgtE is unable to import sodium or potassium. This raises the question why NaCl or KCl would

inhibit the growth of an *mgtE* mutant. It is possible that an abundance of KCl or NaCl in the growth

medium leads to an increase in these ions in the cell envelope. This could displace magnesium ions,

which are an important factor for the stability of the cell wall and a higher requirement of

magnesium under NaCl stress has been demonstrated [62]. Deletion of *mgtE* presumably reduces

the uptake of magnesium and could reduce magnesium levels in the cell envelope leading to a

destabilization of the cell.

In conclusion, we have identified a number of previously uncharacterized factors involved in the

osmotic stress response of *S. aureus,* which in particular highlighted the importance of the cell

envelope and the data generated will provide a great resource for further studies of the

staphylococcal osmotic stress response.

Acknowledgements

This research was funded by the Wellcome Trust through grants 100289 and 210671/Z/18/Z to A.G.

and the NIH grant P01AI083214 to S.W.. C.F.S. is funded by the German Research Foundation

[Deutsche Forschungsgemeinschaft (DFG)] through grant SCHU 3159/1–1 and F.C.M.K. was

supported by a Wellcome trust Ph.D. studentship. We thank Microbes NG and the London Institute

of Medical Sciences for providing whole genome sequencing services and the Tufts sequencing

facility for the sequencing services for the TN-seq experiments. We also would like to thank Ute

Bertsche for kind advice on HPLC analyses.

Author contributions

523 C.F.S., D.M.W., F.C.K. and A.G. performed the research; C.F.S., D.M.W., F.C.M.K., M.S., S.W. and A.G.

524 analyzed the data; C.F.S. and A.G. designed the research; C.F.S. and A.G. wrote the paper. All authors

525 approved the final version of the manuscript.

Conflict of interests

527 The authors declare not conflict of interest.

Data availability

529 The whole genome sequencing data were deposited at the European Nucleotide Archive (ENA)

530 under accession id ERP115099 and TN-seq data in the short read archive (SRA) at the National

531 Center for Biotechnology Information (NCBI) under BioProject ID PRJNA544248. Analyzed tables can

532 be found in the supplementary files. For other data, please contact the corresponding author.

Material and Methods

Growth of bacteria

535 *E. coli* and *S. aureus* strains were streaked from frozen stock onto Lysogenic Broth (LB) or Tryptic

536 Soya agar (TSA) plates, respectively. For all *E. coli* and most *S. aureus* experiments, the bacteria were

537 grown in LB medium (LB normal, LBN, 10 g Tryptone, 5 g Yeast extract, 5 g NaCl) pH 7.5 or LB medium

538 with extra 0.25 M, 0.5 M NaCl, 0.5 M KCl or 1 M sucrose added. For molecular cloning and some

539 other experiments (as indicated in the text), *S. aureus* strains were grown in TSB medium. Growth

curves were done in 125 ml flasks with 20 ml of medium without antibiotics. Cultures were inoculated from overnight cultures to a starting OD_{600} of 0.01 and growth was followed by determining OD_{600} readings every one or two hours.

TN-seq experiments

A previously published highly-saturated *S. aureus* USA300 library with a mix of six different transposons was used [34, 35]. In addition, a new transposon library with only the blunt transposon, containing approximately 1.2 million individual clones, was produced using the same techniques as described in Santiago *et al.* [34].

In the first experiment (replicate A), a tube of the mixed library was thawed on ice and used to set up a 20 ml pre-culture in LBN/Erm 5 µg/ml with a starting OD_{600} of 0.1. This starter culture was grown for 1 hour and subsequently used to inoculate 25 ml of either LBN, LB 0.25 M NaCl or LB 0.5 M NaCl containing Erm 5 µg/ml to a starting OD_{600} of 0.00125. When the cultures reached an OD_{600} of approximately 1.4, bacteria from 10-12 ml culture were harvested by centrifugation (10 generation sample). For, the 17 generation sample, the first cultures were back-diluted 1:100-fold into 25 ml fresh medium when they reached an OD_{600} of 0.3 and bacteria from 10-12 ml harvested by centrifugation when the back-diluted culture reached an OD_{600} of 1.3 to 1.6 (17 generations). The bacterial pellets were washed once with their respective growth medium and stored at -20°C for further processing. In the second experiment (replicate B), the transposon library containing only the blunt transposon was used, Erm was omitted after the pre-culture step and samples were only collected at around 16 generations. In the third experiment, the blunt library was used, Erm 10 µg/ml was only added to the pre-culture, and the bacteria grown in either LBN, LB 0.5 M NaCl, LB 0.5 M KCl or LB 1.0 M sucrose medium and cells collected after 16 generations.

The library preparation for sequencing was done as described previously [34]. Briefly, genomic DNA was isolated, cut with *Not*I, biotinylated adapters ligated, the DNA cut again with *Mme*I, and

fragments ligated to Illumina adapters and the products PCR amplified, incorporating bar codes and

the Illumina adaptor sequences P5 and P7. The samples were sequenced on an Illumina HiSeq

machine after spiking with 40% PhiX DNA. The data analysis was done using the Tufts Galaxy Server

and custom python scripts as described earlier [34]. To this end, the reads were trimmed up to the

barcode and de-multiplexed by strain barcode. Due to the variability in the DNA cleavage by the

*Mme*I restriction enzyme, the reads that could not be mapped to a barcode were trimmed by an

additional base and the process repeated. Reads were then mapped to the USA300_TCH1516

(CP000730.1) genome and a hop table was generated. Statistical analysis (Mann-Whitney tests and

Benjamini-Hochberg) was done using custom python scripts [34]. Further exploration of the data

was done using Excel and custom R scripts.

S. aureus cell diameter measurements

For cell diameter measurements, *S. aureus* strains ANG4054, ANG4340, ANG4341 were grown

overnight (14-20 hours) at 37°C with 180 rpm shaking in 5 ml LBN in the presence of 100 ng/ml Atet.

100 µl of these overnight cultures was transferred to a 1.5 ml reaction tube and 1.5 µl of BODIPY-

FL-vancomycin (100 µg/ml in phosphate buffered saline) was added to each tube to stain the cell

wall. After a 30 min static incubation step at 37°C, 1.5 µl of the suspension was pipetted onto a slide

covered with a 1% agarose pad and analyzed by microscopy using an Axio Imager.A2 microscope

with an EC Plan-Neofluar 100x/1.30 Oil M27 objective and images recorded using an AxioCam MR

R3. Native CZI files were opened in FIJI [63] and cell diameters of 200 cells were measured using the

line and measure tool. Only cells without any visible septa were measured. Experiment was

conducted five times with independent biological cultures (n=5, 200 cells each).

CFU/OD correlations

The different *S. aureus* strains were grown as described for the cell diameter measurements. Optical densities of cultures were measured as well as 100 µl of 10^{-6} dilutions (made in their respective growth medium) plated onto TSA plates. The next day, the number of colonies were counted and the ratios of OD_{600} to CFU calculated. The experiment was conducted six time with independent biological samples (n=6).

Zymogram assays

The different *S. aureus* strains were grown over night in TSB. The next day, the bacteria from an OD_{600} equivalent of 20 were pelleted by centrifugation for 3 min at 17000 x *g*. The cells were washed twice with 600 µl PBS and subsequently suspended in 50 µl SDS sample buffer. The suspension was boiled for 20 min with interspersed shaking, centrifuged for 5 min at 17000 x *g* and 20 µl of each sample was loaded onto Zymogram gels, which where were prepared as described previously [64]. Gels were stained and de-stained using methylene blue and water. The experiment was performed twice, and a representative result is shown.

WTA isolation and analysis

Overnight cultures of *S. aureus* strains ANG4054, ANG4340, ANG4341, ANG1575, ANG4290, ANG4748, ANG4749, ANG4759 were prepared in LBN and used to inoculate 50 ml LBN or LB 0.4 M NaCl to an OD_{600} of 0.01. The cultures were incubated at 37°C with shaking until they reached an OD_{600} of 5 to 6. Where appropriate, 10 µg/ml chloramphenicol and 100 ng/ml Atet was also added to the medium. The bacteria from 20-24 ml culture were harvested by centrifugation for 10 min at 7000 x *g*. The bacterial pellets were washed with 50 mM 2-(N-morpholino) ethanesulfonic acid (MES) pH 6.5 buffer and stored at -20°C until further use. Pellets were then processed and WTA

607 separated on a 20% native polyacrylamide gel by electrophoresis on a Biorad Protein XL ii cell as

608 described previously [65] and WTA visualized by silver staining according to the manufacturer

609 protocol. Experiments were done with three biological replicates (n=3) and a representative result

610 is shown.

Peptidoglycan isolation and analysis

Peptidoglycan was prepared as described previously [66] with the following modifications: 0.5 L

(ANG4290, ANG4382, ANG4384) or 1.0 L (ANG5054, ANG4340, ANG4341) of LB 0.4 M NaCl medium

(all strains) or LBN (ANG5054, ANG4340, ANG4341) medium were inoculated to an OD_{600} of 0.01.

For strains carrying derivatives of plasmid piTET, the medium was supplemented with 100 ng/ml

Atet starting from the pre-cultures. The cultures were grown at 37°C with shaking at 180 rpm until

they reached an OD_{600} of 2-3 and bacteria were subsequently harvested by centrifugation.

Chromatography of mutanolysin digested peptidoglycan was performed as described previously

using an Agilent 1260 infinity system [66] and muropeptide peaks assigned according to de Jonge *et*

al. [67]. For the muropeptide quantification, a baseline was drawn, the total peak area determined

as well as the peak areas for mono- and the different multimer peaks and calculated as percentage

of the total peak area. The peak area quantification was done three times for each HPLC

chromatogram and average values were calculated. Experiments were done with three biological

replicates (n=3).

Data analysis, software and statistics

Data were processed with a combination of Python 3.6 (https://www.python.org), R 3.3 & 3.5

(https://www.r-project.org/), RStudio 1.1 & 1.2 (https://www.rstudio.com/), Prism 7 and 8

(https://www.graphpad.com), ChemStation OpenLab C.01.05 (https://www.agilent.com/) and

Microsoft Excel 15 and 16 (https://www.office.com/). Voronoi images were drawn using

Proteomaps V2 [36, 37] using custom TMD files. Microscopy images were analyzed using FIJI 1.0

(https://fiji.sc/). Statistical analysis was performed with Prism using appropriate tests as described

in the figure legends.

Raising of Δ*957* suppressor strains

Multiple independent overnight cultures of strain LAC*Δ*957* (ANG4290) were prepared in 5 ml LBN.

These cultures were back diluted the next day into 20 ml LB 0.5 M NaCl to a starting OD_{600} of 0.05

and grown for 8-10 hours at 37°C with shaking until the cultures were slightly turbid. 50 µl of each

culture was passed into 5 ml of LBN and grown overnight. The next day, appropriate culture dilutions

were plated onto LBN agar plates and incubated overnight at 37°C. For each independent culture,

multiple single colonies were picked and used to inoculate individual wells of a 96-well microtiter

plates containing 100 µl LBN and the plates were subsequently incubated at 37°C in a 96-well plate

incubator with shaking at 650 rpm. The next day, a culture aliquot was stored at -80°C, and the

cultures were also diluted 1:50 in LB 0.5 M NaCl medium and 20 µl of these diluted cultures used to

inoculate 180 µl of LB 0.5 M NaCl medium. The plates were incubated overnight with shaking at

37°C and the next morning the growth of the potential *957* suppressor strains compared to that of

the original LAC*Δ*957* deletion strain (ANG4290), which showed low growth and the WT LAC*

(ANG1575) strain, which showed good growth. For each lineage, four suppressors that showed good

growth were streaked out from the previously frozen stocks and subsequently single colonies

selected to set up overnight cultures. The deletion of gene *SAUSA300_0957* was confirmed by PCR

and after performing growth curves in culture flasks, one strain that showed significant growth

improvement compared to the original LAC*Δ*957* (ANG4290) strain was selected from each lineage.

These strains were streaked out again for single colonies and used to inoculate an overnight culture,

giving rise to the independently raised suppressor strains ANG4381 through ANG4394. The

compensatory mutations for several of these suppressor strains were subsequently determined by

whole genome sequencing.

Transfer of *pbp2* SNPs by phage transduction

In order to demonstrate that the SNPs in *pbp2* are responsible for the growth rescue of strain

LAC*Δ*957* (ANG4290), we transferred two different *pbp2* SNPs by phage transduction into the

original LAC*Δ*957* strain as well as repaired these SNPs in the suppressor strains. This was done by

placing an Erm-marked transposon in proximity of the *pbp*2 gene allowing at a certain frequency for

the co-transduction of the Erm marker and the *pbp2* SNP. To this end, the NTML transposon mutant

library strain NE789 [43] was used, which harbors a transposon insertion in *SAUSA300_1332*

(putative exonuclease) about 10 kbp upstream of the *pbp2* gene. *SAUSA300_1332* is expected to be

unrelated to salt stress and the cell wall synthesis machinery but close enough to lead to an

intermediate rate of co-transference with the *pbp2* SNPs. The transposon from NE789 was

transduced into WT LAC* [68] and strain LAC*Δ*957* (ANG4290) using phage Φ85 yielding strains

ANG4561 and ANG452, respectively. The transposon was also transduced into the suppressor

strains ANG4382 and ANG4384 transferring either *SAUSA300_1332::Tn* only, yielding strains

ANG4527 and 4528 or transferring *SAUSA300_1332::Tn* as well as replacing the *pbp2* SNP with a WT

pbp2 allele yielding *pbp2* repaired strains ANG4557 and ANG4568. Lysates from strains ANG4527

and 4528 (suppressors containing *pbp2* SNPs and *SAUSA300_1332::Tn*) were used to transfer the

pbp2 SNPs back into a clean LAC*Δ*957* strain background yielding strains ANG4624 and ANG4625

and into WT LAC*, yielding strains ANG4563 and ANG4564. Successful repair or transfer of the SNPs

was checked by PCR and subsequent restriction digest of the product choosing enzymes that

recognize sites present in either the WT or SNP allele (ANG4382 *pbp2* SNP: BsrI. Recognition site

CCWGG, additional site introduced by SNP; ANG4384 *pbp2* SNP: SspI. Recognition site: AATATT, site

missing in SNP).

Determination of oxacillin MICs using Etest strips

S. aureus strains LAC* piTET (ANG4054), LAC*Δ*957* piTET (ANG4340) and the complementation strain LAC*Δ*957* piTET-*957* (ANG4341) were grown overnight (22 hours) in 5 ml TSB containing 10 µg/ml chloramphenicol and 100 ng/ml Atet. The next day, the cultures were diluted to an OD_{600} of 0.1 in sterile PBS buffer and streaked with cotton swabs onto cation adjusted Mueller-Hinton agar plates supplemented with 2% NaCl and 100 ng/ml Atet. One M.I.C.Evaluator strip was placed on each plate and the plates were incubated for 24 hours at 35°C. MICs were then read directly from the strips. The experiment was done with 4 biological replicates (n=4).

Moenomomycin growth improvement test

S. aureus strains LAC* (ANG1575), LAC* Δ*957* (ANG4290), LAC* Δ*957* S2 (ANG4382) and LAC* Δ*957* S4 (ANG4384) were grown overnight (18 hours) in 5 ml LBN. The next day, the cells were diluted to an OD_{600} of 0.01 in LBN, grown until mid-exponential phase (OD_{600} 0.4-0.6) and normalized to an OD_{600} of 0.1. The cells were 10-fold serially diluted and 5 µl of each dilution spotted onto LB agar containing 0.5 M NaCl with either 0.02 µg/ml moenomycin (mix of moenomycin A, A12, C1, C3 and C4, Santa Cruz Biotechnology) or no moenomycin. Plates were incubated at 37°C overnight (18-22 hours) and photographed. The experiment was done with 3 biological replicates (n=3) and one representative result is shown.

Whole genome sequencing

Genomic DNA was either isolated using the Promega Gene Wizard kit according to the manufacturer instructions or using chloroform-isoamylalcohol as described previously [69]. DNA was sent off for whole genome sequencing to MicrobesNG, Birmingham, U.K. or libraries prepared using the Illumina Nextera DNA kit and sequenced at the London Institute of Medical Sciences. Short reads

699 were trimmed in CLC workbench genomics (Qiagen), then mapped to a custom *S. aureus* USA300

700 LAC* reference genome generated in a previous study [70] and single nucleotide polymorphisms

701 called based on at least 80% frequency of occurrence. This list was compared to a manually curated

702 list of well non false-positives and these entries were removed.

Nebraska Transposon Mutant Library (NTML) strains and complementation strains

All Nebraska transposon mutant library (NTML) strains and primers used in this study are listed in

supplementary tables S8, S15 and S16. Transposon insertions in the respective genes were

confirmed by PCRs. For complementation analysis, NTML strains NE535 (JE2 *SAUSA300_0867*::*Tn*),

NE867 (JE2 *SAUSA300_0483*::*Tn*) were transformed as controls with the empty plasmid pCL55 [71],

yielding strains ANG4326 and ANG4328 and strains NE188 (JE2 *SAUSA300_0481*::*Tn*), NE251 (JE2

SAUSA300_0482::*Tn*), NE526 (JE2 *SAUSA300_0694*::*Tn*), NE736 (JE2 *SAUSA300_0910*::*Tn*), NE1384

(JE2 *SAUSA300_0957*::*Tn*) with plasmid piTET [72], yielding ANG4308, ANG4310, ANG4377,

ANG4336, ANG4338. The WT strain JE2 [43] was also transformed with both plasmids yielding JE2

pCL55 (ANG4325) and JE2 piTET (ANG4307).

The complementation plasmid piTET-*481* for strain NE188 (JE2 *SAUSA300_0481::Tn*) was

constructed by amplifying the *SAUSA300_0481* gene using primers P2378/P2379, digesting the

product with AvrII/BglII and ligating the fragment with plasmid piTET cut with the same enzymes.

The plasmid was then introduced into *E. coli* strain XL1-Blue, yielding ANG4139. After shuttling the

plasmid through *E. coli* strain IM08B [73], creating strain ANG4150, the plasmid was integrated into

the *geh* locus of NE188 (JE2 *SAUSA300_0481*::*Tn*) yielding strain ANG4309. The complementation

plasmid piTET-*482* for strain NE251 (JE2 SAUSA300_0482::*Tn*), piTET-*694* for strain NE526 (JE2

SAUSA300_0694::*Tn*), piTET-*910* for strain NE736 (JE2 *SAUSA300_0910*::*Tn*), and piTET-*957* for

strain NE1384 (JE2 *SAUSA300_0957*::*Tn*) were constructed in a similar manner using primer pairs

P2380/P2381 (*SAUSA300_0482*), P2388/P2389 (*SAUSA300_0694*), P2384/P2385 (*SAUSA300_0910*)

and P2386/P2387 (*SAUSA300_0957*), respectively. The complementation plasmids were recovered

in *E. coli* XL1-Blue yielding strains ANG4140 (SAUSA300_0482), ANG4142 (*SAUSA300_0694*),

ANG4144 (*SAUSA300_0910*) and ANG4145 (*SAUSA300_0957*), shuttled through *E. coli* IM08B giving

strains ANG4151 (*SAUSA300_0482*), ANG4153 (*SAUSA300_0694*), ANG4155 (*SAUSA300_0910*) and

ANG4156 (*SAUSA300_0957*) and finally introduced in the respective NTML strain yielding the

complementation strains ANG4311 (*SAUSA300_0482*), ANG4378 (*SAUSA300_0694*), ANG4337

(*SAUSA300_0910*) and ANG4339 (*SAUSA300_0957*).

Complementation plasmid pCL55-*483* for strain NE867 (JE2 *SAUSA300_0483*::*Tn*) was made by

fusing PCR products of the operon promoter (in front of *SAUSA300_0481*) amplified with primers

P2588/P2553 and the *SAUSA300_0483* gene, amplified with primers P2554/P2589 together using

primers P2588/P2589. The resulting fragment was cloned into pCL55 using EcoRI/BamHI restriction

sites. The plasmid was recovered in XL1-Blue, creating strain ANG4291, shuttled through *E. coli*

IM08B (ANG4292) and subsequently introduced into NE867 (JE2 *SAUSA300_0483*::*Tn*) to create

strain ANG4327. The complementation plasmid pCL55-*867* for strain NE535 (JE2

SAUSA300_0867::*Tn*) was constructed in a similar manner, using primers P2590/P2556 and

P2557/P2591 to amplify the promoter and *SAUSA300_0867* gene, which were subsequently fused

in a second PCR using primers P2590/P2591. The resulting fragment was cloning into pCL55 using

EcoRI/BamHI restriction sites and the resulting plasmid recovered in *E. coli* XL1-Blue (ANG4293),

shuttling through IM08B (ANG4294) and finally introduced into NE535 (JE2 *SAUSA300_0867*::*Tn*),

creating the complementation strain ANG4329.

Construction of *S. aureus* gene deletion and complementation strains

S. aureus strains with in-frame gene deletions were constructed by allelic exchange using plasmids

pIMAY [73] and pIMAY* [38]. The gene deletion plasmids were designed to contain approximately

1000 bp up- and downstream regions around the deletion site, amplified from LAC* genomic DNA

[68], and the first and last 30 bp of the open reading frame to be deleted.

For construction of plasmid pIMAY-Δ*957*, the up- and downstream regions of *SAUSA300_0957* were

amplified using primers P2370/P2371 and P2372/P2373, spliced together in a second PCR using

primers P2370/P2373 and cloned into pIMAY using XmaI/EcoRI. Plasmid pIMAY-Δ957 was recovered

in *E. coli* XL1-Blue, creating strain ANG4147, shuttled through *E. coli* IM08B (ANG4159) and

subsequently introduced into S*. aureus* LAC*Δ*957*. The allelic exchange to delete *SAUSA300_0957*

and to create strain LAC*Δ*957* (ANG4290) was performed as previously described [73]. For

complementation analysis, the above described plasmid piTET-*957* was integrated into the

chromosome of strain LAC*Δ*957*, giving rise to the complementation strain LAC*Δ*957* piTET-*957*

(ANG4341). As control, the empty plasmid piTET from ANG4163 was also integrated into the

chromosome of LAC*Δ*957* yielding strain LAC*Δ*957* piTET (ANG4340). The *lcpB* (SAUSA300_0958)

gene was deleted in a similar manner using primers P2844/P2845 and P2846/P2847 for the first and

primers P2844/P2847 for the second PCR and cloning the fragment XmaI/EcoRI into pIMAY. Plasmid

pIMAY-Δ*lcpB* was recovered in *E. coli* XL1blue (ANG4740), shuttled through *E. coli* IM08B (ANG4742)

and introduced into the *S. aureus* LAC* (ANG4744) and finally yielding the *lcpB* deletion strain

LAC*Δ*lcpB* (ANG4748).

Plasmid pIMAY*-Δ*957-958* was constructed for the production of an *SAUSA300_0957-0958* (*lcpB*)

double mutant strain in which the first 30 bases of *SAUSA300_0957* are fused to the last 30 bp of

SAUSA300_0958 (*lcpB*). This was done using primers P2844/P2848 and P2849/P2373 in the first and

primers P2844/P2373 in a second PCR. The fragment was cloned using XmaI/EcoRI into pIMAY* [38]

and the resulting plasmid pIMAY*-Δ*957-958* recovered in *E. coli* XL1-Blue (ANG4741). The plasmid

was shuttled through *E. coli* IM08B (ANG4743), introduced into *S. aureus* LAC* (ANG4745) and

following the allelic exchange procedure yielding the *S. aureus* strain LAC*Δ*957-958* (*lcpB*)

(ANG4749).

The plasmid pIMAY-Δ*mgtE* was designed to contain up- and downstream regions of

SAUSA300_0910 (*mgtE*) fused together by the first and last 30 bp of the gene in order to create a

LAC*Δ*mgtE* mutant. For this, primer pairs P2374/P2375 and P2376/P2377 were used to amplify up-

and downstream regions and subsequently spliced together by PCR using primers P2374/P2377.

The PCR fragment was digested with XmaI/EcoRI and ligated into pIMAY cut with the same enzymes.

The plasmid pIMAY-Δ*mgtE* was recovered in *E. coli* XL1-Blue yielding strain ANG4146 and shuttled

through *E. coli* IM08B yielding strain ANG4158. The plasmid pIMAY-Δ*mgtE* was then introduced into

LAC* yielding strain LAC*Δ*SAUSA300_0910* (*mgtE*) (ANG4422) after the knockout procedure. To

create an empty vector-containing control strain and a plasmid-based complementation strain, the

plasmid piTET and piTET-*910* from strains ANG4163 and ANG4155 were isolated and used to

transform strain LAC*Δ*SAUSA300_0910* (*mgtE*) (ANG4422), yielding strains ANG4445 (LAC*Δ*mgtE*

piTET) and ANG4446 (LAC*Δ*mgtE* piTET-*mgtE)*.

The *tagO* gene (*SAUSA300_0731*) was inactivated using the targetron intron homing system [74] or

by allelic exchange using pIMAY* [38]. For the targetron mutagenesis, plasmid pNL9164-*tagO* was

produced by amplifying the upstream targetron fragment with primers P2887 and the EBS universal

primer (P2890) and the downstream fragment using primers P2888/P2889, the fragments were

fused using the IBS primer (P2887) and the EBS1d primer (P2888). The final fragment was cloned

using HindIII/BsrGI into pNL9164 [74]. Plasmid pNL9164-*tagO* was recovered in *E. coli* XL1-Blue,

yielding strain ANG4703, introduced into *E. coli* IM08B to give rise to strain ANG4704 and finally

introduced into *S. aureus* LAC* [68] to create strain LAC* pNL9164-*tagO* (ANG4751). The final *tagO*

targetron mutant strain LAC*tagO*::targetron (ANG4753) was generated using the method

described by Yao *et al.* [74]. For the construction of *S. aureus* strain LAC*Δ*tagO* with an in-frame

deletion in *tagO,* the allelic exchange plasmid pIMAY*-Δ*tagO* was used. To this end, up- and

downstream *tagO* fragments were produced using primer pairs P2883/P2884 and P2885/P2886 and

introduced by Gibson cloning into XmaI/EcoRI pre-cut pIMAY*. Plasmid pIMAY*-Δ*tagO* was

recovered in *E. coli* XL1-Blue, generating strain ANG4755. After shuttling through *E. coli* strain IM08B

(ANG4756), the plasmid was introduced into *S. aureus* LAC* to create strain ANG4757. Finally, the

tagO locus was deleted by allelic exchange, generating strain LAC*Δ*tagO* (ANG4759). Because initial

generation of the mutant failed, cells were plated unselectively onto TSA plates and incubated at

room temperature (18-25°C) for about two weeks to enable colony differentiation between WT and

Δ*tagO* mutant strains. Upon prolonged incubation, *tagO* mutant strain colonies have an opaque

appearance.

References

1. Wertheim HF, Melles DC, Vos MC, van Leeuwen W, van Belkum A, Verbrugh HA, et al. The

role of nasal carriage in *Staphylococcus aureus* infections. Lancet Infect Dis. 2005;5(12):751-62. Epub

2005/11/29. doi: 10.1016/S1473-3099(05)70295-4. PubMed PMID: 16310147.

2. Tong SY, Davis JS, Eichenberger E, Holland TL, Fowler VG, Jr. *Staphylococcus aureus*

infections: epidemiology, pathophysiology, clinical manifestations, and management. Clin Microbiol

Rev. 2015;28(3):603-61. Epub 2015/05/29. doi: 10.1128/CMR.00134-14. PubMed PMID: 26016486;

PubMed Central PMCID: PMCPMC4451395.

3. Measures JC. Role of amino acids in osmoregulation of non-halophilic bacteria. Nature.

1975;257(5525):398-400. Epub 1975/10/02. PubMed PMID: 241020.

4. Adams MR, Moss MO. 7.14.2 The Organism and its Characteristics. Food Microbiology (3rd

Edition): Royal Society of Chemistry; 2008.

5. Csonka LN. Physiological and genetic responses of bacteria to osmotic stress. Microbiol Rev.

1989;53(1):121-47. Epub 1989/03/01. PubMed PMID: 2651863; PubMed Central PMCID:

PMCPMC372720.

6.   Csonka LN, Hanson AD. Prokaryotic osmoregulation: genetics and physiology. Annu Rev

Microbiol. 1991;45:569-606. Epub 1991/01/01. doi: 10.1146/annurev.mi.45.100191.003033.

PubMed PMID: 1741624.

7.   Sleator RD, Hill C. Bacterial osmoadaptation: the role of osmolytes in bacterial stress and

virulence. FEMS Microbiol Rev. 2002;26(1):49-71. Epub 2002/05/15. doi: 10.1111/j.1574-

6976.2002.tb00598.x. PubMed PMID: 12007642.

8.   Graham JE, Wilkinson BJ. *Staphylococcus aureus* osmoregulation: roles for choline, glycine

betaine, proline, and taurine. J Bacteriol. 1992;174(8):2711-6. Epub 1992/04/01. PubMed PMID:

1556089; PubMed Central PMCID: PMCPMC205912.

9.   Christian JH, Waltho JA. The Composition of *Staphylococcus aureus* in Relation to the Water

Activity of the Growth Medium. J Gen Microbiol. 1964;35:205-13. Epub 1964/05/01. doi:

10.1099/00221287-35-2-205. PubMed PMID: 14179669.

10.   Miller KJ, Zelt SC, Bae J-H. Glycine betaine and proline are the principal compatible solutes

of *Staphylococcus aureus*. Current Microbiology. 1991;23(3):131-7. doi: 10.1007/bf02091971.

11.   Townsend DE, Wilkinson BJ. Proline transport in *Staphylococcus aureus*: a high-affinity

system and a low-affinity system involved in osmoregulation. J Bacteriol. 1992;174(8):2702-10. Epub

1992/04/01. PubMed PMID: 1556088; PubMed Central PMCID: PMCPMC205911.

12.   Anderson CB, Witter LD. Glutamine and proline accumulation by *Staphylococcus aureus* with

reduction in water activity. Appl Environ Microbiol. 1982;43(6):1501-3. Epub 1982/06/01. PubMed

PMID: 7103493; PubMed Central PMCID: PMCPMC244260.

13.   Bae JH, Miller KJ. Identification of two proline transport systems in *Staphylococcus aureus*

and their possible roles in osmoregulation. Appl Environ Microbiol. 1992;58(2):471-5. Epub

1992/02/01. PubMed PMID: 1610171; PubMed Central PMCID: PMCPMC195271.

14.   Pourkomailian B, Booth IR. Glycine betaine transport by *Staphylococcus aureus*: evidence for

feedback regulation of the activity of the two transport systems. Microbiology. 1992;140 (Pt

11):3131-8. Epub 1994/11/01. doi: 10.1099/13500872-140-11-3131. PubMed PMID: 7812452.

15.   Price-Whelan A, Poon CK, Benson MA, Eidem TT, Roux CM, Boyd JM, et al. Transcriptional

profiling of *Staphylococcus aureus* during growth in 2 M NaCl leads to clarification of physiological

roles for Kdp and Ktr K+ uptake systems. MBio. 2013;4(4). Epub 2013/08/22. doi:

10.1128/mBio.00407-13. PubMed PMID: 23963175; PubMed Central PMCID: PMCPMC3747578.

16.   Scybert S, Pechous R, Sitthisak S, Nadakavukaren MJ, Wilkinson BJ, Jayaswal RK. NaCl-

sensitive mutant of *Staphylococcus aureus* has a Tn917-lacZ insertion in its ars operon. FEMS

Microbiol Lett. 2003;222(2):171-6. Epub 2003/05/29. doi: 10.1016/S0378-1097(03)00312-4.

PubMed PMID: 12770703.

17.   Schwan WR, Lehmann L, McCormick J. Transcriptional activation of the *Staphylococcus

aureus* putP gene by low-proline-high osmotic conditions and during infection of murine and human

tissues. Infect Immun. 2006;74(1):399-409. Epub 2005/12/22. doi: 10.1128/IAI.74.1.399-409.2006.

PubMed PMID: 16368996; PubMed Central PMCID: PMCPMC1346636.

18.   Vijaranakul U, Xiong A, Lockwood K, Jayaswal RK. Cloning and nucleotide sequencing of a

Staphylococcus aureus gene encoding a branched-chain-amino-acid transporter. Appl Environ

Microbiol. 1998;64(2):763-7. Epub 1998/02/17. PubMed PMID: 9464420; PubMed Central PMCID:

PMCPMC106115.

19.   Wengender PA, Miller KJ. Identification of a PutP proline permease gene homolog from

Staphylococcus aureus by expression cloning of the high-affinity proline transport system in

Escherichia coli. Appl Environ Microbiol. 1995;61(1):252-9. Epub 1995/01/01. PubMed PMID:

7887605; PubMed Central PMCID: PMCPMC167280.

20.   Wetzel KJ, Bjorge D, Schwan WR. Mutational and transcriptional analyses of the

Staphylococcus aureus low-affinity proline transporter OpuD during *in vitro* growth and infection of

murine tissues. FEMS Immunol Med Microbiol. 2011;61(3):346-55. Epub 2011/01/15. doi:

10.1111/j.1574-695X.2011.00781.x. PubMed PMID: 21231964; PubMed Central PMCID:

PMCPMC3055929.

21.   Zeden MS, Schuster CF, Bowman L, Zhong Q, Williams HD, Gründling A. Cyclic di-adenosine

monophosphate (c-di-AMP) is required for osmotic regulation in *Staphylococcus aureus* but

dispensable for viability in anaerobic conditions. J Biol Chem. 2018;293(9):3180-200. Epub

2018/01/13. doi: 10.1074/jbc.M117.818716. PubMed PMID: 29326168; PubMed Central PMCID:

PMCPMC5836111.

22.   Vijaranakul U, Nadakavukaren MJ, de Jonge BL, Wilkinson BJ, Jayaswal RK. Increased cell size

and shortened peptidoglycan interpeptide bridge of NaCl-stressed *Staphylococcus aureus* and their

reversal by glycine betaine. J Bacteriol. 1995;177(17):5116-21. Epub 1995/09/01. PubMed PMID:

7665491; PubMed Central PMCID: PMCPMC177291.

23.   Onyango LA, Hugh Dunstan R, Roberts TK, Macdonald MM, Gottfries J. Phenotypic variants

of staphylococci and their underlying population distributions following exposure to stress. PLoS

One. 2013;8(10):e77614. Epub 2013/11/10. doi: 10.1371/journal.pone.0077614. PubMed PMID:

24204894; PubMed Central PMCID: PMCPMC3799968.

24.   Vollmer W, Blanot D, de Pedro MA. Peptidoglycan structure and architecture. FEMS

Microbiol Rev. 2008;32(2):149-67. Epub 2008/01/16. doi: 10.1111/j.1574-6976.2007.00094.x.

PubMed PMID: 18194336.

25.   Madiraju MV, Brunner DP, Wilkinson BJ. Effects of temperature, NaCl, and methicillin on

penicillin-binding proteins, growth, peptidoglycan synthesis, and autolysis in methicillin-resistant

Staphylococcus aureus. Antimicrob Agents Chemother. 1987;31(11):1727-33. Epub 1987/11/01.

PubMed PMID: 3435121; PubMed Central PMCID: PMCPMC175029.

26.   Stapleton MR, Horsburgh MJ, Hayhurst EJ, Wright L, Jonsson IM, Tarkowski A, et al.

Characterization of IsaA and SceD, two putative lytic transglycosylases of *Staphylococcus aureus*. J

Bacteriol. 2007;189(20):7316-25. Epub 2007/08/07. doi: 10.1128/JB.00734-07. PubMed PMID:

17675373; PubMed Central PMCID: PMCPMC2168438.

27. Ochiai T. Staphylococcus aureus produces autolysin-susceptible cell walls during growth in a

high-NaCl and low-Ca2+ concentration medium. Microbiol Immunol. 2000;44(2):97-104. Epub

2000/05/10. PubMed PMID: 10803496.

28. Koprivnjak T, Mlakar V, Swanson L, Fournier B, Peschel A, Weiss JP. Cation-induced

transcriptional regulation of the dlt operon of *Staphylococcus aureus*. J Bacteriol.

2006;188(10):3622-30. Epub 2006/05/05. doi: 10.1128/JB.188.10.3622-3630.2006. PubMed PMID:

16672616; PubMed Central PMCID: PMCPMC1482844.

29. Mistretta N, Brossaud M, Telles F, Sanchez V, Talaga P, Rokbi B. Glycosylation of

Staphylococcus aureus cell wall teichoic acid is influenced by environmental conditions. Sci Rep.

2019;9(1):3212. Epub 2019/03/03. doi: 10.1038/s41598-019-39929-1. PubMed PMID: 30824758;

PubMed Central PMCID: PMCPMC6397182.

30. Kho K, Meredith TC. Salt-Induced Stress Stimulates a Lipoteichoic Acid-Specific Three-

Component Glycosylation System in *Staphylococcus aureus*. J Bacteriol. 2018;200(12). Epub

2018/04/11. doi: 10.1128/JB.00017-18. PubMed PMID: 29632092; PubMed Central PMCID:

PMCPMC5971477.

31. Stewart CM, Cole MB, Legan JD, Slade L, Schaffner DW. Solute-specific effects of osmotic

stress on *Staphylococcus aureus*. J Appl Microbiol. 2005;98(1):193-202. Epub 2004/12/22. doi:

10.1111/j.1365-2672.2004.02445.x. PubMed PMID: 15610432.

32. Stewart CM, Cole MB, Legan JD, Slade L, Vandeven MH, Schaffner DW. *Staphylococcus*

aureus growth boundaries: moving towards mechanistic predictive models based on solute-specific

effects. Appl Environ Microbiol. 2002;68(4):1864-71. Epub 2002/03/28. doi:

10.1128/aem.68.4.1864-1871.2002. PubMed PMID: 11916706; PubMed Central PMCID:

PMCPMC123857.

33.   Mitchell P, Moyle J. Permeability of the envelopes of *Staphylococcus aureus* to some salts,

amino acids, and non-electrolytes. J Gen Microbiol. 1959;20(2):434-41. Epub 1959/04/01. doi:

10.1099/00221287-20-2-434. PubMed PMID: 13654736.

34.   Santiago M, Matano LM, Moussa SH, Gilmore MS, Walker S, Meredith TC. A new platform

for ultra-high density *Staphylococcus aureus* transposon libraries. BMC Genomics. 2015;16:252. doi:

10.1186/s12864-015-1361-3.   PubMed   PMID: 25888466;   PubMed   Central   PMCID:

PMCPMC4389836.

35.   Coe KA, Lee W, Komazin-Meredith G, Meredith TC, Grad YH, Walker S. Comparative Tn-Seq

reveals common daptomycin resistance determinants in *Staphylococcus aureus* despite strain-

dependent differences in essentiality of shared cell envelope genes. bioRxiv. 2019:648246. doi:

10.1101/648246.

36.   Liebermeister W, Noor E, Flamholz A, Davidi D, Bernhardt J, Milo R. Visual account of protein

investment in cellular functions. Proc Natl Acad Sci U S A. 2014;111(23):8488-93. Epub 2014/06/04.

doi: 10.1073/pnas.1314810111.   PubMed   PMID: 24889604;   PubMed   Central   PMCID:

PMCPMC4060655.

37.   Otto A, Bernhardt J, Meyer H, Schaffer M, Herbst FA, Siebourg J, et al. Systems-wide

temporal proteomic profiling in glucose-starved Bacillus subtilis. Nat Commun. 2010;1:137. Epub

2011/01/27. doi: 10.1038/ncomms1137. PubMed PMID: 21266987; PubMed Central PMCID:

PMCPMC3105300.

38.   Schuster CF, Howard SA, Gründling A. Use of the counter selectable marker PheS* for

genome engineering in *Staphylococcus aureus*. Microbiology. 2019. Epub 2019/04/04. doi:

10.1099/mic.0.000791. PubMed PMID: 30942689.

39.   Ishibashi M, Kurokawa K, Nishida S, Ueno K, Matsuo M, Sekimizu K. Isolation of temperature-

sensitive mutations in *murC* of *Staphylococcus aureus*. FEMS Microbiology Letters. 2007;274(2):204-

9. doi: 10.1111/j.1574-6968.2007.00829.x.

40. Mlynek KD, Callahan MT, Shimkevitch AV, Farmer JT, Endres JL, Marchand M, et al. Effects

of Low-Dose Amoxicillin on Staphylococcus aureus USA300 Biofilms. Antimicrob Agents Chemother.

2016;60(5):2639-51. Epub 2016/02/10. doi: 10.1128/AAC.02070-15. PubMed PMID: 26856828;

PubMed Central PMCID: PMCPMC4862544.

41. Brown S, Santa Maria JP, Jr., Walker S. Wall teichoic acids of gram-positive bacteria. Annu

Rev Microbiol. 2013;67:313-36. doi: 10.1146/annurev-micro-092412-155620. PubMed PMID:

24024634; PubMed Central PMCID: PMCPMC3883102.

42. Vaish M, Price-Whelan A, Reyes-Robles T, Liu J, Jereen A, Christie S, et al. Roles of

Staphylococcus aureus Mnh1 and Mnh2 Antiporters in Salt Tolerance, Alkali Tolerance, and

Pathogenesis. J Bacteriol. 2018;200(5). Epub 2017/12/22. doi: 10.1128/JB.00611-17. PubMed PMID:

29263099; PubMed Central PMCID: PMCPMC5809693.

43. Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, et al. A genetic resource for

rapid and comprehensive phenotype screening of nonessential *Staphylococcus aureus* genes. MBio.

2013;4(1):e00537-12. doi: 10.1128/mBio.00537-12. PubMed PMID: 23404398; PubMed Central

PMCID: PMCPMC3573662.

44. Mäder U, Nicolas P, Depke M, Pané-Farré J, Debarbouille M, van der Kooi-Pol MM, et al.

Staphylococcus aureus Transcriptome Architecture: From Laboratory to Infection-Mimicking

Conditions. PLoS Genet. 2016;12(4):e1005962. Epub 2016/04/02. doi:

10.1371/journal.pgen.1005962. PubMed PMID: 27035918; PubMed Central PMCID:

PMCPMC4818034.

45. Chan YG, Frankel MB, Dengler V, Schneewind O, Missiakas D. *Staphylococcus aureus* mutants

lacking the LytR-CpsA-Psr family of enzymes release cell wall teichoic acids into the extracellular

medium. J Bacteriol. 2013;195(20):4650-9. doi: 10.1128/JB.00544-13. PubMed PMID: 23935043;

PubMed Central PMCID: PMCPMC3807444.

46.   Over B, Heusser R, McCallum N, Schulthess B, Kupferschmied P, Gaiani JM, et al. LytR-CpsA-

Psr proteins in *Staphylococcus aureus* display partial functional redundancy and the deletion of all

three severely impairs septum placement and cell separation. FEMS Microbiol Lett.

2011;320(2):142-51. doi: 10.1111/j.1574-6968.2011.02303.x. PubMed PMID: 21554381.

47.   Schaefer K, Matano LM, Qiao Y, Kahne D, Walker S. In vitro reconstitution demonstrates the

cell wall ligase activity of LCP proteins. Nat Chem Biol. 2017;13(4):396-401. Epub 2017/02/07. doi:

10.1038/nchembio.2302. PubMed PMID: 28166208; PubMed Central PMCID: PMCPMC5362317.

48.   Rahman MM, Hunter HN, Prova S, Verma V, Qamar A, Golemi-Kotra D. The *Staphylococcus*

aureus Methicillin Resistance Factor FmtA Is a d-Amino Esterase That Acts on Teichoic Acids. MBio.

2016;7(1):e02070-15. Epub 2016/02/11. doi: 10.1128/mBio.02070-15. PubMed PMID: 26861022;

PubMed Central PMCID: PMCPMC4752606.

49.   Berger-Bächi B, Barberis-Maino L, Strässle A, Kayser FH. FemA, a host-mediated factor

essential for methicillin resistance in *Staphylococcus aureus*: molecular cloning and characterization.

Mol Gen Genet. 1989;219(1-2):263-9. PubMed PMID: 2559314.

50.   Oshida T, Sugai M, Komatsuzawa H, Hong YM, Suginaka H, Tomasz A. A *Staphylococcus*

aureus autolysin that has an N-acetylmuramoyl-L-alanine amidase domain and an endo-beta-N-

acetylglucosaminidase domain: cloning, sequence analysis, and characterization. Proc Natl Acad Sci

U S A. 1995;92(1):285-9. PubMed PMID: 7816834; PubMed Central PMCID: PMCPMC42863.

51.   Holland LM, Conlon B, O'Gara JP. Mutation of *tagO* reveals an essential role for wall teichoic

acids in *Staphylococcus epidermidis* biofilm development. Microbiology. 2011;157(Pt 2):408-18.

Epub 2010/11/06. doi: 10.1099/mic.0.042234-0. PubMed PMID: 21051486.

52.   Wyke AW. Isolation of five penicillin-binding proteins from *Staphylococcus aureus*. FEMS

Microbiology Letters. 1984;22(2):133-8. doi: doi:10.1111/j.1574-6968.1984.tb00711.x.

53.   Murakami K, Fujimura T, Doi M. Nucleotide sequence of the structural gene for the penicillin-

binding protein 2 of *Staphylococcus aureus* and the presence of a homologous gene in other

staphylococci. FEMS Microbiol Lett. 1994;117(2):131-6. doi: 10.1111/j.1574-6968.1994.tb06754.x.

PubMed PMID: 8181715.

54.   Lovering AL, De Castro L, Strynadka NC. Identification of dynamic structural motifs involved

in peptidoglycan glycosyltransfer. J Mol Biol. 2008;383(1):167-77. doi: 10.1016/j.jmb.2008.08.020.

PubMed PMID: 18760285.

55.   Ostash B, Walker S. Moenomycin family antibiotics: chemical synthesis, biosynthesis, and

biological activity. Nat Prod Rep. 2010;27(11):1594-617. Epub 2010/08/24. doi: 10.1039/c001461n.

PubMed PMID: 20730219; PubMed Central PMCID: PMCPMC2987538.

56.   Mayer S, Steffen W, Steuber J, Götz F. The *Staphylococcus aureus* NuoL-like protein MpsA

contributes to the generation of membrane potential. J Bacteriol. 2015;197(5):794-806. Epub

2014/12/03. doi: 10.1128/JB.02127-14. PubMed PMID: 25448817; PubMed Central PMCID:

PMCPMC4325100.

57.   Tomita A, Zhang M, Jin F, Zhuang W, Takeda H, Maruyama T, et al. ATP-dependent

modulation of MgtE in Mg(2+) homeostasis. Nat Commun. 2017;8(1):148. Epub 2017/07/28. doi:

10.1038/s41467-017-00082-w. PubMed PMID: 28747715; PubMed Central PMCID:

PMCPMC5529423.

58.   Schuster CF, Bellows LE, Tosi T, Campeotto I, Corrigan RM, Freemont P, et al. The second

messenger c-di-AMP inhibits the osmolyte uptake system OpuC in *Staphylococcus aureus*. Sci Signal.

2016;9(441):ra81. Epub 2016/08/18. doi: 10.1126/scisignal.aaf7279. PubMed PMID: 27531650;

PubMed Central PMCID: PMCPMC5248971.

59.   Chan YG, Kim HK, Schneewind O, Missiakas D. The capsular polysaccharide of *Staphylococcus*

aureus is attached to peptidoglycan by the LytR-CpsA-Psr (LCP) family of enzymes. J Biol Chem.

2014;289(22):15680-90. Epub 2014/04/23. doi: 10.1074/jbc.M114.567669. PubMed PMID:

24753256; PubMed Central PMCID: PMCPMC4140922.

60.   Pöhlmann-Dietze P, Ulrich M, Kiser KB, Döring G, Lee JC, Fournier JM, et al. Adherence of

Staphylococcus aureus to endothelial cells: influence of capsular polysaccharide, global regulator

agr, and bacterial growth phase. Infect Immun. 2000;68(9):4865-71. Epub 2000/08/19. PubMed

PMID: 10948098; PubMed Central PMCID: PMCPMC101683.

61.   Pinho MG, Filipe SR, de Lencastre H, Tomasz A. Complementation of the essential

peptidoglycan transpeptidase function of penicillin-binding protein 2 (PBP2) by the drug resistance

protein PBP2A in *Staphylococcus aureus*. J Bacteriol. 2001;183(22):6525-31. Epub 2001/10/24. doi:

10.1128/JB.183.22.6525-6531.2001.   PubMed   PMID: 11673420;   PubMed   Central   PMCID:

PMCPMC95481.

62.   Ochiai T. *Staphylococcus aureus* requires increased level of Ca(2+) or Mn(2+) to grow

normally in a high-NaCl/low-Mg(2+) medium. Microbiol Immunol. 2001;45(11):769-76. Epub

2002/01/17. PubMed PMID: 11791670.

63.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-

source   platform   for   biological-image   analysis.   Nat   Methods.   2012;9(7):676-82.   doi:

10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID: PMCPMC3855844.

64.   Atilano ML, Pereira PM, Vaz F, Catalão MJ, Reed P, Grilo IR, et al. Bacterial autolysins trim

cell surface peptidoglycan to prevent detection by the *Drosophila* innate immune system. Elife.

2014;3:e02277. doi: 10.7554/eLife.02277. PubMed PMID: 24692449; PubMed Central PMCID:

PMCPMC3971415.

65.   Covas G, Vaz F, Henriques G, Pinho MG, Filipe SR. Analysis of Cell Wall Teichoic Acids in

Staphylococcus aureus. Methods Mol Biol. 2016;1440(1440):201-13. doi: 10.1007/978-1-4939-

3676-2_15. PubMed PMID: 27311674.

66.   Corrigan RM, Abbott JC, Burhenne H, Kaever V, Gründling A. c-di-AMP is a new second

messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope stress. PLoS

Pathog. 2011;7(9):e1002217. doi: 10.1371/journal.ppat.1002217. PubMed PMID: 21909268;

PubMed Central PMCID: PMCPMC3164647.

67.   de Jonge BL, Chang YS, Gage D, Tomasz A. Peptidoglycan composition of a highly methicillin-

resistant *Staphylococcus aureus* strain. The role of penicillin binding protein 2A. J Biol Chem.

1992;267(16):11248-54. PubMed PMID: 1597460.

68.   Boles BR, Thoendel M, Roth AJ, Horswill AR. Identification of genes involved in

polysaccharide-independent *Staphylococcus aureus* biofilm formation. PLoS One.

2010;5(4):e10146. doi: 10.1371/journal.pone.0010146. PubMed PMID: 20418950; PubMed Central

PMCID: PMCPMC2854687.

69.   Schuster CF, Bertram R. Fluorescence based primer extension technique to determine

transcriptional starting points and cleavage sites of RNases *in vivo*. J Vis Exp. 2014;(92):e52134. doi:

10.3791/52134. PubMed PMID: 25406941; PubMed Central PMCID: PMCPMC4353401.

70.   Bowman L, Zeden MS, Schuster CF, Kaever V, Grundling A. New Insights into the Cyclic Di-

adenosine Monophosphate (c-di-AMP) Degradation Pathway and the Requirement of the Cyclic

Dinucleotide for Acid Stress Resistance in *Staphylococcus aureus*. J Biol Chem. 2016;291(53):26970-

86. Epub 2016/11/12. doi: 10.1074/jbc.M116.747709. PubMed PMID: 27834680; PubMed Central

PMCID: PMCPMC5207132.

71.   Lee CY, Buranen SL, Ye ZH. Construction of single-copy integration vectors for *Staphylococcus*

aureus. Gene. 1991;103(1):101-5. PubMed PMID: 1652539.

72.   Gründling A, Schneewind O. Genes required for glycolipid synthesis and lipoteichoic acid

anchoring in *Staphylococcus aureus*. J Bacteriol. 2007;189(6):2521-30. doi: 10.1128/JB.01683-06.

PubMed PMID: 17209021; PubMed Central PMCID: PMCPMC1899383.

73.   Monk IR, Shah IM, Xu M, Tan MW, Foster TJ. Transforming the untransformable: application

of direct transformation to manipulate genetically *Staphylococcus aureus* and *Staphylococcus*

epidermidis. MBio. 2012;3(2). doi: 10.1128/mBio.00277-11. PubMed PMID: 22434850; PubMed

Central PMCID: PMCPMC3312211.

74.   Yao J, Zhong J, Fang Y, Geisinger E, Novick RP, Lambowitz AM. Use of targetrons to disrupt

essential and nonessential genes in *Staphylococcus aureus* reveals temperature sensitivity of Ll.LtrB

group II intron splicing. RNA. 2006;12(7):1271-81. doi: 10.1261/rna.68706. PubMed PMID:

16741231; PubMed Central PMCID: PMCPMC1484445.

1069

1070

Figure legends

Fig. 1. TN-Seq screen reveals essential and dispensable genes during prolonged NaCl stress in *S. aureus*.

(A) Workflow of TN-seq experiments. A saturated transposon library was pre-cultured for one hour,

then used to inoculate LB medium containing either normal levels (0.086 M), 0.25 M or 0.5 M of

NaCl. After 10 and 17 generations, cells were harvested, and transposon insertion sites determined

by high throughput sequencing.

(B) Growth curve. Growth of the *S. aureus* library culture in LBN, LB 0.25 M NaCl and LB 0.5 M

medium was followed by determining OD$_{600}$ readings. Dotted lines indicate 10 and 17 generation

thresholds. Cultures were back-diluted once when they reached an OD of approximately 0.3 and the

optical densities shown are calculated from measured ODs time dilution factor. Culture from

replicate B (n=1) is shown and a similar growth profile was seen for replicate A.

(C) Volcano plots of essential and dispensable genes in NaCl conditions compared to LBN. Negative

$\log_2$ fold-changes indicate essential and positive $\log_2$ fold-changes dispensable genes. q-value stands

for Benjamini-Hochberg false discovery rate. Black horizontal line indicates q-value of 0.05 (cut-off)

and red, orange and green vertical lines 10-, 5- and 2-fold differences in either direction. Colored

circles indicate genes for which a significant change in the number of transposon insertions was

observed above q-value cutoff and follow the same color code as the vertical lines. Bacteria in

replicate A (top) were grown in 0.25 M and 0.5 M NaCl for 10 and 17 generations, whereas bacteria

in replicate B (bottom) were only grown for 17 generations (n=1 for each plot).

(D) Principal component analysis (PCA). A PCA was performed on the TN-seq data for replicates A

and B shown in panel (C). The data for the 17 generation 0.25 M (orange) and 0.5 M NaCl (red)

growth conditions clustered together, indicating good reproducibility between the two replicates.

1095

1096

Fig. 2. Growth curves of *S. aureus* strains with transposon insertions in potential salt essential genes.

S. aureus strain JE2 (WT) and strains from the Nebraska Transposon Mutant Library (NTML) containing transposon insertions in potential salt essential genes were grown in LBN (left column) or 0.5 M NaCl LB medium (right column) and their growth monitored over 8 hours. Transposon mutant strains shown in panels (A) and (B) had a mild growth defect while strains shown in panels (C) and (D) had a stronger growth defect when grown in the high salt medium. NE numbers

1104   correspond to the NTML mutant number. Locus tag number and where available gene names with

1105   transposon insertion site are given in parenthesis. Growth curves were performed in triplicates

1106   (n=3) and means and SDs were plotted.

Fig. 3. Growth curves and complementation analysis of *S. aureus* strains with transposon insertions in salt essential genes.

S. aureus strains (A) JE2 (WT), *SAUSA300_0694::Tn* mutant with empty plasmid and complementation strain or (B) JE2 (WT), *SAUSA300_0910* (*mgtE*), *SAUSA300_0957::Tn* with empty plasmid and respective complementation strains were grown in LBN (left panels) or LB 0.5 M NaCl medium (right panels) supplemented with 100 ng/ml Atet and OD_{600} readings determined at timed intervals. Growth curves were performed in triplicates (n=3) and means and SDs of OD_{600} readings were plotted.

Fig. 4. Gene *SAUSA300_0957* is located in a genomic region with cell envelope genes and its

deletion leads to salt sensitivity and other phenotypic changes.

(A) Schematic of the *S. aureus* USA300 FPR3757 chromosomal region with gene *SAUSA300_0957*

(gene *957*). Gene *957* is encoded in an operon with *lcpB*, which codes for a wall teichoic acid ligase.

fmtA, a proposed D-amino esterase acting on teichoic acid and genes coding for a predicted

acyltransferase and the major autolysin Atl are located up and downstream of this operon. Putative

promoters are shown as angled arrows (adapted from [44]) and black bars indicate the

corresponding transcripts.

(B) Growth curves. *S. aureus* strains LAC* piTET (WT), LAC*Δ*957* piTET (Δ*957*) and the

complementation strain LAC*Δ*957* piTET-*957* (compl.) were grown in LBN or LB containing 0.5 M

NaCl medium and 100 ng/ml Atet. The growth was monitored by determining OD_{600} readings the

means and SDs from three independent experiments were plotted.

(C) Determination of CFU/OD ratios. OD_{600} values as well as CFU/ml were determined for overnight

cultures of the *S. aureus* strains described in panel (B) and the means and SDs of the CFU per ml

OD_{600} of 1 from five independent experiments were plotted.

(D) Cell size measurements. The cell walls of the *S. aureus* strains described in panel (B) were stained

with fluorescently labelled vancomycin and the cells subsequently observed under a microscope.

The diameters of 200 cells were measured and the means calculated. This average of the means and

SDs from five independent experiments were plotted.

For statistical analysis, a Kruskal-Wallis one-way ANOVA test was performed followed by a Dunn's

post-hoc test to determine p-values. Asterisks (*) indicate p≤0.05 and two asterisks (**) p≤0.01 and

ns=not significant.

1141

Fig. 5. Inactivation of gene 957 leads to an increase in peptidoglycan crosslinking.

(A) Growth curves. *S. aureus* strains LAC* (WT), LAC*Δ*957* (Δ*957*), LAC*Δ*lcpB*, (Δ*957*), and

LAC*Δ*tagO* (Δ*tagO*) were grown in LB 0.5 M NaCl medium and OD_{600} readings determined at timed

intervals. The means and SDs from three biological replicates were plotted.

(B) Detection of WTA on silver stained gels. WTA was isolated from *S. aureus* strains described in

panel (A) following growth in LB 0.4 M NaCl medium. The WTA was separated by electrophoresis

and visualized by silver stain. The experiment was performed three times and one representative

gel image is shown.

(C) Detection of WTA on silver stained gels. Same as (B) but using the strains LAC* piTET (WT),

LAC*Δ*957* piTET (Δ*957*), and the complementation strain LAC*Δ*957* piTET-*957* (compl.) grown in LB

0.4 M NaCl medium also containing 100 ng/ml Atet.

(D) Zymogram gel. Cell extracts prepared from *S. aureus* strains described in panel (C) were

separated on a gel containing heat killed *Micrococcus luteus* cells. Autolysins were renatured and

the zones of lysis visualized by methylene blue staining. The experiment was performed twice and

one experiment is shown.

(E) Determination of oxacillin MICs. Oxacillin MICs were determined for *S. aureus* strains described

in panel (C) using Etest strips. The median and SDs from four biological replicates were plotted.

(F) Muropeptide analysis. The same *S. aureus* strains as described in (C) were grown in LB 0.4 M

NaCl medium also containing 100 ng/ml Atet, peptidoglycan isolated and digested with mutanolysin

and the resulting muropeptide fragments separated by HPLC. Representative chromatograms are

shown in Supplementary Figures S5E. The peak area of individual peaks (corresponding to monomer

up to 8-mers and above) were quantified and the means and SDs from three biological replicates

plotted.

For statistical analysis of the oxacillin MICs in (E) a Kruskal-Wallis one-way ANOVA and a Dunn's

post-hoc test was performed and for the muropeptide analysis in (F) a two-way ANOVA and a

Dunnett's post-hoc test. One asterisk (*) indicates p≤0.05, two asterisks (**) p≤0.01 and four

asterisks (****) p≤0.0001. ns=not significant.

1170

1171

Fig. 6. The growth and peptidoglycan defect observed for the *957* mutant can be rescued by

compensatory mutations in *pbp2*.

(A) Growth curves. *S. aureus* strains LAC* (WT), LAC*Δ*957* (Δ*957*) and the LAC*Δ*957* suppressors

S1-10 (S1 to 10) were grown in LB 0.5 M NaCl medium, OD_{600} readings determined and the means

and SDs of four biological replicates plotted.

(B) Schematic of Pbp2 with amino acid substitutions identified. Top: Structure of the *S. aureus*

penicillin binding protein Pbp2 (PDB: 3DWK, [54]) with amino acids that are altered in the obtained

suppressor strains shown in red. Bottom: Schematic of the Pbp2 enzyme with the transglycosylase

and transpeptidase domains as well as the observed amino acid changes in the transglycosylase

domain indicated.

(C) Growth curves using *957* mutant strains containing *pbp2* alleles from suppressors S2 and S4.

Growth curves were performed in LB 0.5 M NaCl medium and plotted as described in panel (A) but

using *S. aureus* strains LAC* *1332*::Tn (WT), LAC*Δ*957* *1332*::Tn (Δ*957*), LAC*Δ*957* *1332*::Tn *pbp2*

SNP S2 (Δ*957 pbp2* SNP S2), LAC*Δ*957* suppressor S4 *1332*::Tn (Δ*957 pbp2* SNP S4).

(D) Growth curves using *957* suppressors with their *pbp2* gene repaired to WT. Growth curves were

performed in LB 0.5 M NaCl medium and plotted as described in panel (A) but using *S. aureus* strains

LAC* *1332*::Tn (WT), LAC*Δ*957* *1332*::Tn (Δ*957*), LAC*Δ*957* suppressor S2 *1332*::Tn (S2), LAC*Δ*957*

suppressor S4 *1332*::Tn (S4), LAC*Δ*957* suppressor S2 *1332*::Tn repaired WT *pbp2* (S2 WT *pbp2*),

and LAC*Δ*957* suppressor S4 *1332*::Tn repaired WT *pbp2* (S4 WT *pbp2*).

(E) Growth curves using WT strains carrying *pbp2* SNPs. *S. aureus* strains LAC* *1332*::Tn (WT), LAC*

1332::Tn *pbp2* SNP S2 (WT *pbp2* SNP S2) and LAC* *1332*::Tn *pbp2* SNP S4 (WT *pbp2* SNP S4) were

grown in LB 0.5 M NaCl medium and OD_{600} readings determined and the means and SDs of three

biological replicates plotted.

(F) Bacterial growth on moenomycin supplemented agar plates. *S. aureus* strains LAC* (WT),

LAC*Δ*957* (Δ*957*), LAC*Δ*957* Suppressor S2 (S2) and S4 LAC*Δ*957* Suppressor S4 (S4) were grown

to exponential phase, normalized to an OD_{600} of 0.1 and plated either neat (n) or in 10-fold dilutions

onto LB agar containing 0.5 M NaCl without (top) or with (bottom) moenomycin. Shown is one

representative image of three biological replicates.

(G) Muropeptide analysis. *S. aureus* strains LAC*Δ*957* (Δ*957*), LAC*Δ*957* Suppressor S2 (S2) and S4

LAC*Δ*957* Suppressor S4 (S4) were grown in 0.4 M NaCl LB medium and muropeptides quantified

as described in Fig 5F. Representative chromatograms are shown in Supplementary Figure S7D. The

means and SDs from three biological replicates were plotted. For statistical analysis a two-way

ANOVA and Dunnett's post-hoc test was performed. One asterisk (*) indicates p≤0.05, two asterisks

(**) p≤0.01 and four asterisks (****) p≤0.0001.

1206

Fig. 7. Different sets of genes are conditionally essential for the growth of *S. aureus* under NaCl,

KCl or sucrose stress conditions.

(A) Workflow of the TN-seq experiment. A TN-seq library was pre-cultured and then used to

inoculate LBN, LB containing 0.5 M NaCl, 0.5 M KCl or 1 M sucrose. Cells were kept in exponential

phase for 16 generations, harvested and transposon insertion sites determined by high throughput

sequencing.

(B) Growth curves. For the TN-seq experiment (n=1), the growth of an *S. aureus* library cultured in

LBN, LB 0.5M NaCl, LB 0.5 M KCl or LB 1.0 M sucrose medium was followed by taking OD_{600}

measurements at timed intervals. The cultures were back-diluted once when they reached an OD_{600}

of approximately 0.3. The plotted OD_{600} values were calculated by multiplying the measured OD_{600}

with the dilution factor. The dotted line indicates the 16 generation cut-off at which cultures were

harvested.

(C) Circular plot showing the transposon insertion density along the *S. aureus* genome under

different osmotic stress conditions. The two outer rings depict genes located on the (+) or (-) strand

in *S. aureus* strain USA300 FPR3757. The inner four rings show the histograms of transposon

insertions on a per gene basis after growth of the library in LBN (orange), 0.5 M NaCl (brown), 0.5

M KCl (blue) and 1 M sucrose (pink) medium for 16 generations.

(D) Volcano plots of conditionally essential and dispensable genes following growth of *S. aureus* in

0.5 M NaCl, 0.5 M KCl or 1 M sucrose medium and compared to LBN conditions. q-values represent

Benjamini-Hochberg corrected p-values, and $\log_2$ (fold changes) indicate essential genes (negative

values) or dispensable genes (positive values). The black horizontal line indicates a q-value of 0.05,

which was deemed the significance level. Vertical lines indicate 2- (green), 5- (orange) or 10- (red)

fold changes in either direction. Each dot represents one gene and coloring follows the fold-change

scheme whenever the q-value threshold was met. The TN-seq experiments were conducted once

(n=1).

(E) Venn diagrams showing overlaps of conditionally essential and dispensable genes during

prolonged NaCl, KCl or sucrose stress. Genes with a 2-fold decrease (essential genes) or a 2-fold

increase (dispensable genes) in transposon insertions under each stress condition compared to the

LBN condition and a q-value of ≤0.05 were determined and the overlap between these gene lists

displayed in Venn diagrams.

1237

Conditionally essential genes

Fig. 8. Voronoi diagram-based visualization of conditionally essential *S. aureus* genes when exposed to different osmotic stressors underlines differences.

Conditionally essential genes of *S. aureus* strains grown either in 0.5 M NaCl (A), 0.5 M KCl (B) or 1.0 M sucrose (C) were mapped to cellular functions. Area sizes were adjusted to their essentiality regardless of p- or q-value to visualize differences on a genome wide level. The larger the area, the

1244   higher the number of transposon insertions in the LBN control condition as compared to the stress

1245   conditions. Colors of polygons indicate cellular functions as detailed in (D).

1246

1247

Fig. 9. Growth curves of putative potassium sensitive mutants in LB containing 0.5 M KCl.

(A) Growth curves using putative KCl sensitive mutants. *S. aureus* strain JE2 (WT) and strains from

the Nebraska Transposon Mutant Library (NTML) containing transposon insertions in the indicated

and potential KCl essential genes were grown in LB 0.5 M KCl medium and the mean and SDs of the

OD_{600} readings from three biological replicates plotted.

(B) Growth curves using a clean Δ*mgtE* deletion and complementation strain. *S. aureus* strains LAC*

piTET (WT), LAC*Δ*mgtE* piTET (Δ*mgtE*) and the complementation strain LAC*Δ*mgtE* piTET-*mgtE*

(compl.) were grown in LB 0.5 M KCl also containing 100 ng/ml Atet and the mean and SDs of the

OD_{600} readings from three biological replicates were plotted.

1257

1258

Fig. 10. Proposed model of 957 activity during NaCl stress.

S. aureus is normally able to grow in medium containing up to 2.5 M NaCl. The deletion of gene *957*

leads to a strong growth defect in high salt conditions and changes in the peptidoglycan structure.

We speculate that the observed increase in peptidoglycan crosslinking will lead to an increased

rigidity of the cell wall and disadvantage and increased cell wall damage when these cells are

exposed to osmotic stress conditions.

1265

Supplementary Figures

10 generations
Dataset A

17 generations
Dataset A

17 generations
Dataset B

Genes (+) strand
Genes (-) strand
Histograms
LBN
LB 0.25 M NaCl
LB 0.5 M NaCl

1267

Supplementary Fig. 1. Circos plots showing transposon insertions in the *S. aureus* genome

following growth under different conditions.

The outer two bands represent genes on the forward (blue) and reverse (green) strand, respectively.

The inner three rings represent transposon insertions per gene following the growth of *S. aureus* in

in either LBN (red), LB 0.25 M NaCl (purple) or LB 0.5 M NaCl (blue). The top two panels are for

dataset A and growth for 10 or 17 generations and the lower panel is for dataset B and growth for

17 generations.

1275

Conditionally essential genes at 0.5 M NaCl and 17 generations

Supplementary Fig. 2. Voronoi diagram-based visualization of *S. aureus* conditionally essential genes when exposed to 0.5 M NaCl.

Conditionally essential *S. aureus* genes from replicate A (A) and replicate B (B) grown in 0.5 M NaCl for 17 generations were mapped to cellular functions with area sizes adjusted to their essentiality regardless of p- or q-value. The larger the area, the lower the number of transposon insertions in stress condition as compared to the LBN control condition. Colors of polygons indicate cellular functions as detailed in (C).

Conditionally dispensable genes at 0.5 M NaCl and 17 generations

1285

Supplementary Fig. 3. Voronoi diagram-based visualization of *S. aureus* conditionally dispensable

genes when exposed to 0.5 M NaCl.

Conditionally dispensable *S. aureus* genes from replicate A (A) and replicate B (B) grown in 0.5 M

NaCl for 17 generations were mapped to cellular functions with area sizes adjusted to their

dispensability regardless of p- or q-value. The larger the area, the higher the number of transposon

insertions in the stress condition as compared to the LBN control. *recU* was scaled to 1/5 of the

original ratio to preserve space. Colors of polygons indicate cellular functions as detailed in (C).

1293

Supplementary Fig. 4. Complementation analysis using *S. aureus* mutants with transposon insertions in putative salt resistance genes.

1294

1295

(A) Growth curves. *S. aureus* strains JE2 (WT), NE867 and NE535 with integrated pCL55 and *S. aureus*

strains NE867 and NE535 with respective complementation plasmids were grown in either LBN or

LB 0.5 M NaCl and their growth monitored by determining OD_{600} readings.

(B) As in (A) but using strains JE2 piTET (WT) or NE188 and NE251 containing plasmid piTET or the

respective complementation plasmid. Strains were grown in LBN or 0.5 M NaCl LB medium

supplemented with 100 ng/ml Atet. All experiments were conducted three times and the means

and standard deviations were plotted.

1303

1304

Supplementary Fig. 5. Investigating the involvement of gene *957* in cell wall homeostasis.

(A) Growth curves using strains with mutations in WTA synthesis genes. *S. aureus* strains LAC* (WT),

LAC*Δ*957* (Δ*957*), LAC*Δ*lcpB* (Δ*lcpB*) and LAC*Δ*tagO* (Δ*tagO*) were grown in LBN and growth

monitored by measuring OD_{600} readings. The experiment was performed three times and means

and standard deviations were plotted.

(B) Detection of WTA. WTA was isolated from the strains described in (A) following growth in LBN

medium, then subjected to electrophoresis on polyacrylamide gels and visualized by silver staining.

One representative result is shown out of three independent experiments.

(C) Detection of WTA. Experimental setup is the same as in (B) but using the strains LAC* piTET (WT),

LAC*Δ*957* piTET (Δ*957*) and LAC* Δ*957* piTET-*957* (compl.).

(D) Muropeptide profile. The same strains as described in panel (C) were grown in LBN medium and

the peptidoglycan isolated, digestion with mutanolysin and the muropeptides subsequently

separated by HPLC. Shown are representative chromatograms from three independent

experiments. Retention time ranges of mono-, di- and higher multi-mers are indicated by vertical

lines.

(E) Muropeptide profiles. As in (D) but bacteria were grown in LB 0.4 M NaCl.

(F) Muropeptide quantification. The areas under the peaks from the chromatograms shown in panel

(D) and two additional replicates were quantified for each mono-, di- or multimer area. Bars

represent mean and error bars standard deviations from three independent experiments. For

statistical analyses, a two-way ANOVA was performed, followed by a Dunnett's post-hoc test but no

significant changes were present between either the wild type and the mutant or the wild type and

the deletion strain. Where induction was necessary, the medium was supplemented with 100 ng/µl

Atet.

1328

Supplementary Fig. 6. Growth of Δ957 suppressor strains and schematic of the co-transduction

strategy for complementation analysis.

(A) Growth curves of Δ957 suppressor strains. Strains LAC* (WT), LAC*Δ957 (Δ957) and LAC*Δ957

suppressors S1 trough S10 (S1-S10) were grown in LBN medium and their growth monitored by

taking OD_{600} readings. The experiment was performed three times and means and standard

deviations were plotted.

(B) TN insertion map. Transposon insertions from replicate A (ANG dataset) were mapped to the

recU/*pbp2* operon from a library grown in LBN (red) or LB medium containing 0.5 M NaCl salt (blue).

(C) TN insertion map. As in (B) but using replicate B (FCK dataset).

(D) Schematic of phage co-transduction experiment to either repair the *pbp2* SNP in the suppressor

strains (left branch) or to transduce the *pbp2* suppressor mutations into a clean *957* mutant (right

branch). A phage lysate was prepared using an NTML library strain with a transposon insertion in an

unrelated gene (*SAUSA300_1332*) located approximately 10 kb from the *pbp2* SNPs present in the

suppressor strains. This lysate was used for transductions using different LAC*Δ*957* suppressor

strains as recipient strains. With about 50% frequency, the *pbp2* SNPs were exchanged to the wild

type allele by transducing *SAUSA300_1332*::*Tn* and the WT *pbp2* allele producing strain LAC*Δ*957*

suppressor *1332*::*Tn pbp2* repaired (left branch). In the other cases, the *SAUSA300_1332*::*Tn* region

was transferred to a suppressor strain without replacing the *pbp2* SNP (right branch). A phage lysate

prepared on this intermediate strain (LAC*Δ*957* suppressor *1332*::*Tn*) was then used to transduce

the *pbp2* suppressor SNP into a clean LAC*Δ*957* strain, resulting in the clean suppressor strain

LAC*Δ*957 1332*::Tn *pbp2* SNP transduced.

1350

Supplementary Fig. 7. Growth and peptidoglycan analysis of *957* mutant and suppressor strains.

(A) Growth curves using *957* mutants with transduced *pbp2* SNPs. Strains LAC* *1332*::Tn (WT),

LAC*Δ*957* *1332*::Tn (Δ*957*), LAC*Δ*957* *1332*::Tn *pbp2* SNP S2 (Δ*957* *pbp2* SNP S2), LAC*Δ*957*

suppressor S4 *1332*::Tn (Δ*957* *pbp2* SNP S4) were grown in LBN medium and growth monitored by

determining OD$_{600}$ readings. The means and standard deviations from three independent

experiments were plotted.

(B) Growth curves using *957* suppressor strains carrying a repaired WT *pbp2* gene. Same as in (A)

but using strains LAC* *1332*::Tn (WT), LAC*Δ*957 1332*::Tn (Δ*957*), LAC*Δ*957* suppressor S2 *1332*::Tn

(S2), LAC*Δ*957* suppressor S4 *1332*::Tn (S4), LAC*Δ*957* suppressor S2 *1332*::Tn repaired WT *pbp2*

(S2 WT *pbp2*), and LAC*Δ*957* suppressor S4 *1332*::Tn repaired WT *pbp2* (S4 WT *pbp2*).

(C) Growth curves using WT strains carrying *pbp2* SNP mutations. Same as in (A) but using strains

LAC* *1332*::Tn (WT), LAC* *1332*::Tn *pbp2* SNP S2 (WT *pbp2* SNP S2) and LAC* *1332*::Tn *pbp2* SNP S4

(WT *pbp2* SNP S4).

(D) Muropeptide profiles. Strains LAC*Δ*957* (Δ*957*), LAC*Δ*957* Suppressor S2 (S2) and S4 LAC*Δ*957*

Suppressor S4 (S4) were grown in LB medium with 0.4 M NaCl, the peptidoglycan was extracted,

digested with mutanolysin, and the muropeptide separated by HPLC. Retention ranges for

monomers, di- and higher oligomers are indicated by vertical lines. The experiment was conducted

with three times and one representative result is shown.

Supplementary Fig. 8. Voronoi diagram-based visualization of conditionally dispensable *S. aureus* genes when exposed to different osmotic stressors.

Conditionally dispensable genes of *S. aureus* strains grown either in 0.5 M NaCl (A), 0.5 M KCl (B) or 1.0 M sucrose (C) were mapped to cellular functions with area sizes adjusted to their essentiality regardless of p- or q-value. The larger the area, the higher the number of transposon insertions in

1375   relation to the LBN control. *recU* was scaled to 1/10 of the original ratio to preserve space. Colors

1376   of polygons indicate cellular functions as detailed in (D).

1377