High-throughput transposon sequencing highlights the cell wall as an important barrier for osmotic stress in methicillin resistant *Staphylococcus aureus* and underlines a tailored response to different osmotic stressors

Short Title: Global analysis of osmotic stress responses in Staphylococcus aureus

Christopher F. Schuster[a], David M. Wiedemann[a], Freja C. M. Kirsebom[a,1], Marina Santiago[b,2], Suzanne Walker[b] and Angelika Gründling[a,3]

ORCID IDs: 0000-0002-4566-7226 (CFS), 0000-0001-6961-0980 (DMW), 0000-0003-4585-319X (FCK), 0000-0002-5446-3452 (MS), 0000-0002-0545-914X (SW), 0000-0002-6235-8687 (AG)

[a]Section of Microbiology and MRC Centre for Molecular Bacteriology and Infection, Imperial College London, London, SW7 2AZ, U.K.

[b]Department of Microbiology and Immunobiology, Harvard Medical School, Boston, MA 02115, U.S.A.

[1]Current address: Section of Respiratory Infections, National Heart and Lung Institute, Imperial College London, London, W2 1PG, U.K.

[3]To whom correspondence should be addressed. Email: a.grundling@imperial.ac.uk

Keywords: osmotic stress, TN-seq, Tradis, NaCl, KCl, Sucrose, *S. aureus*, DUF25380, Pbp2

Abstract

Staphylococcus aureus is an opportunistic pathogen that apart from causing soft tissue infections,

is also a frequent cause of foodborne illnesses. One contributing factor for this food association is

its high osmotic tolerance allowing this organism to survive commonly used methods of food

preservation. How this resistance is mediated is poorly understood, particularly during long-term

exposure. In this study, we used TN-seq based screens to identify genes that are involved in the

osmotic stress response of *S. aureus* and to determine how responses differ between different

osmotic stressors. Although it is generally assumed that a generic osmotic stress response exists,

our results revealed distinctly different long-term responses to NaCl, KCl and sucrose stresses. We

were able to link numerous previously unknown factors to the osmotic stress response in *S. aureus*

and identified the DUF2538 domain containing gene *SAUSA300_0957* (gene *957*) as essential under

salt stress. Interestingly, a *957* mutant was less susceptible to oxacillin and showed increased

peptidoglycan crosslinking. The salt sensitivity phenotype could be suppressed by mutations in the

transglycosylase domain of the penicillin binding protein gene *pbp2*, and these mutations also

restored the peptidoglycan crosslinking to WT levels. These results indicate that increased

crosslinking of the peptidoglycan can be detrimental and highlight a critical role of the bacterial cell

wall for osmotic stress resistance. This study will serve as a starting point for future research on

osmotic stress response and help develop better strategies to tackle foodborne staphylococcal

infections.

Introduction

Staphylococcus aureus is a Gram-positive bacterium that is carried by about 20% of healthy individuals (Wertheim *et al.* 2005). It is also an opportunistic pathogen causing a variety of diseases, including bacteremia, endocarditis, soft tissue infections and food poisoning (Tong *et al.* 2015). One characteristic of *S. aureus* is its ability to grow in the presence of a very high salt concentration of up to 2.5 M, conditions under which other bacteria are unable to survive (Measures 1975), and this has been used extensively for the isolation of staphylococci. This salt tolerance also allows *S. aureus* to thrive in its niches such as the human nares, skin (Wertheim *et al.* 2005) and certain food products (Adams and Moss 2008).

In general, the bacterial cytoplasm is hyperosmotic to the environment and can therefore tolerate a certain level of external osmotic stress. However, an increase in the intracellular sodium concentration can lead to mis- and unfolding of proteins and inhibit enzymatic reactions. Hence, *S. aureus* must possess strategies to counter the effects of osmotic damage, but these are poorly understood, as most work has been performed in *Escherichia coli* and *Bacillus subtilis*, which possess much lower salt tolerances than Staphylococci.

General short term mechanisms of osmotic stress mitigation are the rapid import of potassium into the cell to reduce the influx of sodium and efflux of water (Csonka 1989; Csonka and Hanson 1991; Sleator and Hill 2002). However, during prolonged NaCl stress, potassium levels decrease again after its initial accumulation in the cytoplasm of *S. aureus* (Christian and Waltho 1964; Graham and Wilkinson 1992). Subsequently, *S. aureus* accumulates small osmotically active compounds, named compatible solutes (Measures 1975). The most effective compatible solutes in *S. aureus* are glycine betaine and proline (Measures 1975; Anderson and Witter 1982; Miller *et al.* 1991; Bae and Miller 1992; Pourkomailian and Booth 1992; Townsend and Wilkinson 1992), which are imported by specific osmolyte transporters (Bae and Miller 1992; Pourkomailian and Booth 1992). These

preferred osmolytes can accumulate in the cell to very high concentrations without negatively

affecting cellular processes (Csonka and Hanson 1991). In addition, *S. aureus* can also synthesize

compatible solutes *de novo*, as in the case of glutamine (Anderson and Witter 1982), but this process

is much slower than the uptake of osmolytes and requires more energy.

Consistent with these observations, previous research studying the underlying genetic factors

of the *S. aureus* osmotic stress response implicated several osmolyte transporters in this process

(Wengender and Miller 1995; Vijaranakul *et al.* 1998; Scybert *et al.* 2003; Schwan *et al.* 2006; Price-

Whelan *et al.* 2013). These include the potassium transport systems Kdp and Ktr (Price-Whelan *et*

al. 2013), the proline transporter PutP (Wengender and Miller 1995; Schwan *et al.* 2006), the arsenic

transport system Ars (Scybert *et al.* 2003) and the branched chain amino acid uptake system BrnQ

(Vijaranakul *et al.* 1998). In addition, although currently not experimentally verified, it can be

assumed that the main glycine betaine uptake system OpuD (Wetzel *et al.* 2011; Zeden *et al.* 2018)

is an important factor for osmotic adaptation as glycine betaine uptake reduces sensitivity to NaCl

exposure significantly (Miller *et al.* 1991).

Despite these multiple countermeasures, exposure of *S. aureus* to osmotic stress such as high

salt concentrations also leads to morphological changes, including the formation of larger cells

(Vijaranakul *et al.* 1995) and a thickening of the cell wall (Onyango *et al.* 2013). However, if and

which genetic factors are required for those processes are currently not well understood. The cell

wall of Gram-positive bacteria is an important barrier against osmotic stresses and acts as a

counterpart to the pressurized cytoplasm. A major component of the cell wall is peptidoglycan, a

polymer consisting of chains of repeating N-acetylglucosamine (GlcNAc) and N-acetylmuramic acid

(MurNAc) units that are crosslinked with neighboring chains via short peptides and, in the case of *S.*

aureus, pentaglycine cross-bridges (Reviewed by (Vollmer *et al.* 2008)). In the past, the contribution

of the cell wall towards osmotic stress resistance was often not considered in the bacterial osmotic

stress resistance field, presumably because the integrity of the cell wall was assumed to be a

prerequisite rather than an active adaptation to survive osmotic stresses. Nevertheless, several

possible changes to the cell wall have been observed *in S. aureus* during NaCl stress, such as

shortened interpeptide bridges in the peptidoglycan (Vijaranakul *et al.* 1995), an increase in

resistance to methicillin (Madiraju *et al.* 1987) and increased autolysis activity (Ochiai 2000;

Stapleton *et al.* 2007). In addition to peptidoglycan, other components within the cell envelope,

such as teichoic acids and more specifically their modifications with D-alanine (Koprivnjak *et al.*

2006) and sugar residues (Kho and Meredith 2018; Mistretta *et al.* 2019) are affected by osmotic

stress. Combined, these findings indicate that the integrity of the cell wall is an important

requirement for osmotic resistance.

Most studies on osmotic stress in *S. aureus* have been conducted using NaCl as the osmotic

stressor. The ionic properties of NaCl have important implications on protein stability and activity

as its accumulation can lead to denaturation of proteins. A few studies in *S. aureus* have also been

performed with non-ionic osmotic stressors such as sucrose, sorbitol glycerol and amino acids

(Mitchell and Moyle 1959; Stewart *et al.* 2002; Stewart *et al.* 2005; Schwan *et al.* 2006). All of these

different stresses are often used synonymously with osmotic stress, in part influenced by the

findings that the initial mitigation of osmotic stress by either salts or sugars can be prevented by the

accumulation of potassium and compatible solutes. However, there are potential differences in the

long-term adaptation to ionic and non-ionic osmotic stressors, which up to date have not been

investigated in detail. It is currently also not known if the adaptations besides the accumulation of

potassium and osmolytes are similar or differ depending on the osmotic stressor, and this was

addressed as part of this study.

Using a transposon insertion sequencing (TN-seq) method (Santiago *et al.* 2015), we

determined on a whole genome level genes that are essential or dispensable during long-term

osmotic stress caused by exposure to NaCl, KCl and Sucrose. This yielded very different sets of

essential and dispensable candidate genes*,* providing experimental evidence of the distinct nature

of these stresses and linking a number of previously unknown factors to the osmotic stress response in *S. aureus*. NaCl stress proved to be the most severe of these three stressors and using further TN-seq screens and experimental verification, several unknown and understudied genes were identified to be essential under salt stress. This included *SAUSA300_0957,* coding for a cytoplasmic protein, which we show here is important for peptidoglycan homeostasis in *S. aureus*. Another protein, which we identified as important during salt stress and essential for peptidoglycan synthesis, is the penicillin binding protein Pbp2, again highlighting a key function of peptidoglycan during NaCl-induced salt stress. Taken together, these experiments highlight the bacterial cell wall as a key player in the salt stress response of *S. aureus*.

Results

Different types of osmotic stresses target different sets of genes

In the literature, osmotic stress is a loosely used term for any accumulation of osmolytes, however the type of ion or osmolyte can potentially have a great impact on how bacteria respond. To address this issue, we determined how the stress responses to different commonly used osmolytes compare. To this end, a highly saturated *S. aureus* transposon library with a promoter-less transposon was prepared using a phage-based delivery system as described in (Santiago *et al.* 2015) and subsequently used for transposon sequencing (TN-seq) experiments (see Supplementary Table 1 for details on all TN-seq experiments). The library was propagated for 16 generations in either LB medium with a normal salt concentration of 85 mM NaCl (LBN) or with 0.5 M NaCl, 0.5 M KCl or 1.0 M sucrose added (Figure 1A). The molarity of sucrose was doubled to accommodate for the dissociation of the salts into two ions. The culture challenged with 0.5 M NaCl grew the slowest,

followed by 1.0 M sucrose, whereas the cells grown in 0.5 M KCl grew similarly to the cells grown in

LBN (Figure 1B).

As expected, by using a large and good quality transposon library, transposon insertions were found

under all conditions in genes throughout the whole genome (Figure 1C). Next, the number of TN-

insertions per gene following growth under the NaCl, KCl or sucrose stress condition was compared

to the number of TN-insertions per gene after growth in LBN medium (Supplementary Table 2) and

in this manner conditionally essential (Supplementary Table S3) and dispensable (Supplementary

Table S4) genes identified. Fold-changes in transposon insertions per gene and q-values were

plotted in volcano plots (Figure 1D). From this, it was evident that under KCl stress the number of

essential and dispensable genes was much lower than for NaCl or sucrose stress, indicating a much

less severe effect of KCl on *S. aureus* cells. Sucrose stress also showed a reduced set of essential and

dispensable genes compared to NaCl stress, but considerably higher than under the KCl condition.

This was also reflected when inspecting the gene lists of the top 30 essential or dispensable genes

(Supplementary Table S3 and S4) as in the KCl condition only 15 and 2 genes respectively met the

requirements of q-value ≤ 0.05 and a fold-change of 5. To identify common genes between

conditions, the fold-change stringency was relaxed to 2-fold and the overlap of genes was visualized

in Venn diagrams (Figure 1E, individual genes in Supplementary Table S5). Three genes, namely

SAUSA300_0425 (*USA300HOU_0457*, *mpsA (nuoF)*, a cation transporter of the respiratory chain

(Mayer *et al.* 2015)) *SAUSA300_*0750 (*USA300HOU_0796*, *whiA*, a hypothetical protein, possibly

involved in cell wall synthesis) and SAUSA300_0846 (USA300HOU_0903, a possible sodium:proton

antiporter) were essential in all conditions. In the case of dispensable genes, only one gene,

SAUSA300_1255   (*USA300HOU_1294*,   *mprF/fmtC*),   coding   for   a   phosphatidylglycerol

lysyltransferase involved in the defense against cationic microbial peptides, was identified in all

conditions (Figure 1E, Supplementary Table S5). In order to assess similarities and differences on a

gene function and cellular pathway level, functional Voronoi maps were generated from the

conditionally essential (Figure 2) and dispensable (Supplementary Figure 1) genes regardless of their

p-values. These images highlighted the essentiality of the wall teichoic acid through *tagO* and the

dlt operon (Figure 2A) and the transporters AapA and MgtE in both NaCl (Figure 2A) and KCl (Figure

2B) stress conditions. In the high sucrose condition, these transporters were much less prominent

(Figure 2C). The patterns of essential genes suggested a similar stress response for the ionic stressors

NaCl and KCl that was distinct from sucrose stress. When the dispensable genes were investigated

(Supplementary Figure 1), the number and type of genes in the NaCl condition differed considerably

from KCl and sucrose stress. Under NaCl stress (Supplementary Figure 1A), in addition to the

penicillin binding protein *pbp2*, a large number of genes involved either directly in respiration or

indirectly by their role in ubiquinone synthesis were found to be dispensable (Supplementary

Figures 1A). The number of dispensable genes in the KCl (Supplementary Figure 1B) and sucrose

(Supplementary Figure 1C) conditions were considerably less, indicating differences in the osmotic

stress response between NaCl and KCl. Taken together, the overlaps between conditions were small,

indicating distinct modes of actions for each osmotic stress.

TN-seq is a robust method to identify *S. aureus* genes involved in NaCl stress

Since the addition of NaCl provided the most stringent selection, we focused our further analysis on

the *S. aureus* salt stress response and conducted two additional independent TN-seq screens to

confirm the robustness of this approach (Figure 3A). For replicate A, a multiplexed transposon

library, generated using 6 different transposons possessing different or no promoters (Santiago *et*

al. 2015; Coe *et al.* 2019) was used, and for replicate B the same transposon library as described

above was used. The libraries were propagated in either LBN or LB medium with 0.5 M NaCl and the

expected growth reduction was observed in the presence of 0.5 M NaCl (Figure 3B). The Tn

insertions were distributed evenly throughout the genome in the library grown for 17 generations

in LBN but became more unbalanced after growth in the high salt medium due to the enrichment

of better salt-adapted TN-strains (Supplementary Figure 2). The fold-change of transposon

insertions per gene under the salt stress conditions compared to the LBN condition (input library)

as well as q-values (false discovery rate according to Benjamini-Hochberg) were determined for each

gene (Supplementary Table S6 & S7). The data were again plotted as volcano plots (Figure 3C),

revealing similar numbers of conditional salt essential and dispensable genes in both replicates.

Furthermore, a good overlap of the top essential and dispensable genes was observed between the

replicates confirming the robustness of the TN-Seq screens (Supplementary Tables S8-11).

The TN-seq screens highlight the essentiality of transporters and cell envelope related genes and

the dispensability of respiration genes under high salt conditions

To provide a visual and genome-wide guide to the cellular functions of conditional salt essential or

dispensable genes, Voronoi maps were again generated as described above for the replicate A and

B 0.5 M NaCl datasets (Otto *et al.* 2010; Liebermeister *et al.* 2014). In both replicates, genes coding

for transporters and cell envelope related genes were the most prominent conditionally salt

essential genes (Supplementary Figure S3). More specifically, the TN-Seq data indicated that

inactivation of the putative magnesium transporter MgtE (Schuster *et al.* 2019) and the putative D-

serine/D-alanine/glycine transporter AapA is detrimental under salt stress. Inactivation of AapA has

previously been shown to suppress the temperature sensitivity phenotype seen in an *S. aureus*

fmtC/mprF mutant (Ishibashi *et al.* 2007) and to lead to increased amoxacillin and daptomycin

sensitivity (Mlynek *et al.* 2016; Coe *et al.* 2019). In addition, the *dlt* operon, coding for enzymes

required for the D-alanylation of teichoic acids, an important factor for cationic peptide resistance

(Brown *et al.* 2013), as well as the stationary and stress sigma factor gene *sigB* were found to be

essential.

An even larger number of genes was identified as dispensable during salt stress (Supplementary

Figure 4). Inactivating or reducing the expression of such genes is expected to help *S. aureus* survive

under salt stress. Among these dispensable genes were genes coding for the sodium antiporter

Mnh2, which has previously been shown to transport Na^+/H^+ and K^+/H^+ in high pH medium and to

play an important role in cytoplasmic pH maintenance (Vaish *et al.* 2018). Strikingly, a large number

of genes involved in respiration, such as the *cta/qox/men/hem* operons as well as genes involved

indirectly in respiration through quinone synthesis (shikimate pathway, *aro* operon) were also

identified as dispensable during salt stress. Curiously, the data also reproducibly indicated that

genes coding for the phosphodiesterases GdpP and Pde2 involved in degradation of the signaling

molecule c-di-AMP are dispensable under high salt conditions. Somewhat contradictory, the

construction of a *dacA* mutant coding for the cognate c-di-AMP cyclase DacA was only possible at

high salt concentrations (or in defined chemical medium) (Zeden *et al.* 2018); but altogether these

findings confirm the involvement of c-di-AMP in salt mediated osmotic stress adaption in *S. aureus*.

Confirmation of genes essential during salt stress using defined mutants

To determine to what extent different genes identified in the TN-seq screen contribute to the salt

resistance of *S. aureus*, a set of candidate genes from the top salt essential genes of dataset B were

chosen for further analysis. For this analysis we used defined transposon mutants available from the

S. aureus NTML transposon mutant library (Fey *et al.* 2013) (Supplementary Table S12). Most of the

chosen genes showed similar trends in dataset A, but intriguingly two of the tested genes, *rsbV* and

rsbW, were flagged as highly dispensable in dataset A but as essential in dataset B. Since no strains

with transposon insertions in the *dlt* operon were available in the NTLM transposon mutant library,

the contribution of these genes to the salt stress resistance was not further investigated.

Next, the growth of the different NTML transposon mutant strains was assessed in LBN and LB 0.5

M NaCl medium. In LBN medium, most strains grew similarly to the WT strain JE2, indicating that

inactivation of the respective gene does not greatly affect bacterial growth in normal LB medium

(Figures 4A-D, left panels). Exceptions were strains NE1109 (*sigB*) (Figure 4A), NE1778 (*lcpB*) (Figure

4B) and NE188 (*mfd*) (Figure 4C), cultures of which reached slightly lower final optical densities. At

high osmolarity conditions (LB 0.5 M NaCl), all strains showed a growth defect compared to the WT

strain (Figures 4A-D), validating TN-seq as a method to identify *S. aureus* genes important during

salt stress. Several strains exhibited strong growth defects in the presence of 0.5 M NaCl (Figures 4C

and D) with strain NE1384 (*SAUSA300_0957*) showing extremely reduced growth (Figure 4D). To

confirm that the salt sensitivity was mediated by the inactivation of the genes in questions, we

constructed complementation plasmids for seven of the most promising candidates by either

expressing the gene of interest from its native promoter or an anhydrotetracycline (Atet) inducible

promoter in the transposon mutant strains. While no complementation was observed for four

strains (Supplementary Figures 5A and B), the salt-dependent growth defect could be

complemented for strains carrying mutations in *SAUSA300_0694* (Figure 5A), *SAUSA300_0910*

(*mgtE*) and *SAUSA300_0957* (Figure 5B). *SAUSA300_0694* encodes a hypothetical protein with 6

predicted transmembrane helices but no other identifiable domain motif. MgtE (SAUSA300_0910)

is a predicted magnesium transporter. *SAUSA300_0957* (from here on also referred to as gene *957*)

codes for a cytoplasmic protein of unknown function and was further investigated in this study

because a *957* mutant exhibited the strongest salt-sensitivity phenotype.

Genomic organization of *957* suggests a potential function in cell envelope homeostasis

SAUSA300_0957 encodes a protein with a DUF2538 domain of unknown function. Proteins with this

domain are found mainly in Gram-positive bacteria (Actinobacteria and Firmicutes) (based on an

Interpro search, https://www.ebi.ac.uk/interpro/entry/IPR024469). Although a crystal structure of

the 957 protein has been deposited in the PDB (3KBY), to our knowledge nothing is known about

the actual cellular function of this protein. Transcriptome data (Mäder *et al.* 2016) suggest that gene

957 is in an operon with *lcpB* (Figure 6A). LcpB is one of three wall teichoic acid (WTA) ligases present

in *S. aureus* that attaches WTA to the peptidoglycan polymer (Over *et al.* 2011; Chan *et al.* 2013;

Schaefer *et al.* 2017). Inactivation of LcpB leads to slightly lower WTA levels in the cell wall, whereas

deletion of all three WT ligases completely abolishes WTA anchoring to the cell wall (Chan *et al.*

2013). In addition, LcpB cell wall ligase activity was demonstrated *in vitro* (Schaefer *et al.* 2017).

fmtA is located upstream of the *957* operon and has been proposed to code for an esterase that can

remove D-alanine modifications from teichoic acids (Rahman *et al.* 2016) and is involved in

methicillin resistance (Berger-Bächi *et al.* 1989). Genes coding for a predicted acetyltransferase and

atl coding for the major *S. aureus* autolysin are found immediately downstream of the *957* operon

(Oshida *et al.* 1995). In this early work, gene *957* was first described as an ORF of "unknown

function". Since functionally related genes often cluster together in bacteria, this prompted us to

next look at cell-envelope related phenotypes for a *957* deletion strain.

Deletion of gene *957* causes physiological changes leading to a smaller cell size

As all previous experiments were performed with transposon mutants, we first constructed strain

LAC*Δ*957* with a marker-less in-frame deletion of gene *957*. Strain LAC*Δ*957* had the expected

growth defect in the presence of 0.5 M NaCl, which could be complemented by expressing a

functional copy of *957* (Figure 6B). We then proceeded to test if the *957* mutant displays other

phenotypes and to investigate if gene *957* is involved in cell envelope homeostasis. First, WT LAC*,

the *957* mutant and the complementation strain were grown in LBN and the ratio of colony forming

units to optical density (CDU/OD) was determined. The CFU/OD ratio was significantly larger for the

957 mutant (60.8±6.8 x10^7 CFU/OD) mutant as compared to strain LAC* (37.5±9.3 x10^7 CFU/OD)

when grown in LBN, suggesting a change in cell shape or size (Figure 6C). This phenotype was

restored in the complementation strain (37.3±3.6 x10^7 CFU/OD) (Figure 6C). When the average cell

diameter was determined by microscopy, a small but significant reduction in cell diameter was seen

for the *957* mutant (1.25±0.02 μm) as compared to the WT (1.36±0.03 μm) and the cell size was

restored to wild type levels in the complementation strain (1.35±0.05 nm) (Figure 6D).

Salt sensitivity of *957* is not due to interaction with LcpB but likely due to changes in peptidoglycan crosslinking

Since the gene *957* is on the same transcript as *lcpB* coding for a WTA ligase (Figure 6A), the contribution of WTA to salt stress resistance as well as the contribution of *957* to the attachment of WTA to the cell wall was assessed. Initially the growth of a *tagO* (*tarO/llm*) mutant, a strain unable to produce WTA, was tested in LB and LB 0.5 M NaCl medium. When cells were grown in LBN, they grew normally (Supplementary Figure 6A), but under high salt conditions, the *tagO* mutant showed reduced growth, similar to that of the *957* mutant (Figure 7A). This is consistent with what has been reported previously for a *Staphylococcus epidermidis tagO* mutant (Holland *et al.* 2011) and indicates an important role for WTA during salt stress. When the *lcpB* mutant was propagated in LBN medium, its growth was slightly inhibited (Supplementary Figure 6A). However, in high salt medium (Figure 7A) it grew unimpeded and like WT. Since the *lcpB* mutant did not show a growth defect in high salt medium, this made it unlikely that *957* is a regulator of LcpB activity. To further exclude that gene *957* could be somehow involved in the attachment of WTA to the peptidoglycan, we isolated WTA from strains grown in either LB or LB containing 0.4 M NaCl and separated the polymer on native polyacrylamide gels. The slightly lower concentration of salt was chosen because growth of the *957* and the *tagO* mutants was too strongly inhibited at 0.5 M NaCl. As expected the *tagO* mutant did not produce any WTA (Supplementary Figure 6B, Figure 7B) and consistent with previous studies (Over *et al.* 2011; Chan *et al.* 2013), the *lcpB* mutant exhibited slightly reduced WTA levels compared to all other strains (Supplementary Figure 6B, Figure 7B). There was no reduction of WTA observable in the *957* mutant compared to the WT strain in either LBN (Supplementary Figure 6B & 6C) or LB 0.4 M NaCl (Figure 7B & 7C) but rather a small, complementable increase, suggesting no influence of *957* over LcpB on WTA attachment.

Gene *957* is found upstream of the bi-functional autolysin gene *atl* and therefore we next investigated potential changes in autolytic activity using zymograms. Cell extracts from overnight

cultures of the WT, the *957* mutant and the complementation strain were separated on polyacrylamide gels with embedded *Micrococcus luteus* cells and incubated overnight in buffer to re-fold the autolysins and enable the digestion of the peptidoglycan. A small increase in autolytic activity was detected in the *957* mutant (Figure 7D) for bands around 55 kDa and 70 kDa compared to the WT and complementation strain, indicating a possible change in Atl availability. We also tested the susceptibility of the *957* mutant towards the cell wall active beta-lactam antibiotic oxacillin. In contrast to our expectations, the *957* mutant exhibited a small but significant increase in MIC towards this antibiotic than the WT or the complementation strain (Figure 7E), indicating potential changes to the peptidoglycan structure. To test this, we next analyzed the muropeptide profile of mutanolysin-digested peptidoglycan isolated from the WT, the *957* mutant and complementation strain after growth in LBN or LB 0.4 M NaCl medium. Upon first inspection, no peaks were absent in any condition or strain (Supplementary Figure 6D, Supplementary Figure 6E). Probing further, mono- and multimers of muropeptides up to 7-mers (see Supplementary Figure 6D for how binning was performed) were quantified but no significant changes were found when strains were cultured in LBN medium (Supplementary Figure 6F). However, when the muropeptide profiles from cells cultured in 0.4 M NaCl were compared, a significant decrease in the total amount of di- and trimers and a significant increase in higher multimers was observed for the peptidoglycan isolated from the *957* mutant (Figure 7F), suggesting that sensitivity to NaCl could potentially be caused by higher crosslinking and rigidity of the peptidoglycan in the mutant.

Salt-resistant suppressors possess variations in the transglycosylase domain of Pbp2 and show reduced peptidoglycan crosslinking

Next, we attempted to generate *957* suppressor strains that showed improved growth in the presence of 0.5 M NaCl, with the idea that by mapping and investigating the compensatory mutations, further insight into the cellular functions of protein 957 could be gained. We initially

attempted to raise suppressor strains on solid medium but failed to do so, because the growth

defect of the *957* mutant was not as prevalent on agar plates. We therefore used liquid medium

instead, where we readily obtained suppressor strains. These strains grew as well as or even better

than the WT LAC* strain in the high salt conditions (Figure 8A) and also grew similar to the WT in

LBN (Supplementary Figure 7A). The genome sequences of 10 independent suppressor strains were

determined and single nucleotide polymorphisms (SNPs) in one or two genes could be identified for

each strain (Supplementary table S13), except for one strain where the coverage was insufficient,

and this strain was omitted from further analysis. A common denominator for all suppressor strains

were SNPs in the *pbp2* gene, coding for the Penicillin binding protein 2 (Pbp2) (Figure 8B). *S. aureus*

Pbp2 is a bifunctional enzyme (Wyke 1984; Murakami *et al.* 1994), which possesses transglycosylase

and transpeptidase activity and is involved in peptidoglycan synthesis. All nine discovered SNPs were

unique and led to mutations in the amino acid sequence in the transglycosylase domain but neither

to stop codons, nor frameshifts nor mutations in the transpeptidase domain. The mutations were

mapped (Figure 8B) onto an available Pbp2 structure (3DWK) (Lovering *et al.* 2008) to see if a specific

area of the transglycosylase domain had been targeted, but the amino acid substitutions were found

throughout the molecule and not only in the active site. Interestingly, *pbp2* also appeared to be

considerably less important for growth at 0.5 M NaCl (Supplementary Figure 7B and 7C), although

transposons were mainly localized at the beginning and end of the gene or within the promoter

region.

To confirm that the *pbp2* SNPs were indeed responsible for the suppression of the salt sensitivity of

the *957* mutant, we transferred the *pbp2* SNPs from two strains into a fresh LAC*Δ*957* background

by co-transduction of a transposon in a nearby gene (Supplementary Figure 7D, schematic). As

expected, these "recreated" suppressor strains showed also increased salt resistance (Figure 8C,

Supplementary Figure 8A). In addition, we repaired the *pbp2* gene in two suppressors to the WT

pbp2 allele and this reduced their ability to cope with NaCl stress, all consistent with the hypothesis

that the SNPs in *pbp2* are responsible for the suppression phenotype (Figure 8D, Supplementary

Figure 8B). When the same *pbp2* SNPs were transferred into a WT LAC* strain, a reproducible

growth improvement of the SNP-bearing strains was observed in high salt medium, an indication

that the *pbp2* mutations lead to a general growth improvement of *S. aureus* under NaCl stress

conditions (Figure 8E, Supplementary Figure 8C).

Moenomycin is a phosphoglycolipid antibiotic that inhibits the transglycosylase activity of the *S.*

aureus Pbp2 enzyme (reviewed in (Ostash and Walker 2010)). To test if a decrease in Pbp2

transglycosylase activity results in the improved growth of the *957* mutant in high salt conditions,

cells were grown on LB agar containing 0.5 M NaCl with or without moenomycin. The growth

behaviors of the WT strain and the *957* mutant were similar when grown on solid LB agar containing

0.5 M NaCl but lacking moenomycin (Figure 8F, top panel). As stated above, this discrepancy in the

growth behavior of the *957* mutant in high salt liquid versus solid medium was already noted when

we attempted to generate suppressors on agar plates. On the other hand, in medium supplemented

with 0.02 µg/ml moenomycin, we observed better growth of the *957* mutant compared to the WT

strain (Figure 8F, lower panel). In addition, the *957* mutant also showed improved growth compared

to the two suppressor strains S2 and S4 when moenomycin was added (Figure 8F, lower panel).

These findings are consistent with the idea that a partial inhibition of the glycosyltransferase activity

of Pbp2 can improve the growth of the *957* mutant in high salt medium.

Because the LAC*Δ*957* strain showed increased peptidoglycan crosslinking when compared

to a WT strain, we hypothesized that the peptidoglycan of the suppressor strains would be again

less crosslinked. To test this, the peptidoglycan from two of the suppressor strains grown in LB 0.4

M NaCl was isolated, the muropeptide profiles determined and compared to that of the original *957*

mutant (Figure 8G, Supplementary Figure 8D). As before, the muropeptide profiles looked similar

(Supplementary Figure 8D) but quantification revealed a significant reduction in crosslinked

peptidoglycan fragments and a significantly overrepresentation of monomeric and dimeric

fragments in the suppressor strains as compared to the original *957* mutant (Figure 8G). These

results indicate that the amount of crosslinking of the peptidoglycan polymer is an important factor

in the salt resistance of *S. aureus* and that dysregulation of peptidoglycan crosslinking can lead to

destabilizing effects.

Discussion

In this work, we performed TN-seq screens with *S. aureus* exposed to different osmotic stresses and could show that each osmotic stressor affects a defined but different set of genes. It is therefore important to use the term osmotic stress carefully, as our results highlight that there is not one general osmotic stress response but rather responses tailored to the individual osmolyte. Using TN-seq, we successfully identified several salt tolerance genes in the USA300 *S. aureus* LAC* strain and further characterized gene *957,* a gene without previously assigned function. Our results indicate that the gene product is involved in cell envelope homeostasis but likely not WTA attachment and we show that its absence leads to increased crosslinking of the peptidoglycan. Suppressors of a *957* mutant acquired mutations in *pbp2* and the increase in peptidoglycan crosslinking was reversed in such stains. This demonstrates that the synthesis of the peptidoglycan layer is a tightly regulated process that plays an essential role during osmotic stress.

From our TN-seq data and the use of different osmotic stressors (NaCl, KCl and sucrose), it is evident that although all these compounds exert osmotic stress, the response differs greatly between them (Figure 1, Figure 2 and Supplementary Figure S1). NaCl and KCl conditions share a set of genes that become essential such as the transporters *aapA* and *mgtE* (Figure 2A and B) but differ in the set of dispensable genes (Supplementary Figure S1A and B). It is intriguing that a whole set of respiration related genes become dispensable under NaCl stress but not under KCl stress. This could indicate that respiration should be inhibited or slowed down in high NaCl conditions, possibly by interference with sodium pumps a process less likely to be inhibited by potassium ions. The set of essential genes under sucrose conditions (Figure 2C) was vastly different from that of the NaCl and KCl conditions whereas the set of sucrose dispensable genes (Supplementary Figure S1C) looked similar to the KCl set. This highlights the differences in osmotic stress adaptations and serves as a reminder that the

type of stressor remains important. Therefore, although the term osmotic pressure is useful in some

contexts, it often needs to be taken with a grain of salt.

TN-seq experiments provide a wealth of information about the growth ability of single mutants in a

mixed population and can therefore quickly and accurately determine genes that are required under

certain growth condition. It is however always a concern that a single TN-seq experiment might not

be reproducible due to stochastic extinction of individual mutants or certain growth dynamics. In

this study, we performed three independent NaCl TN-seq experiments using two different libraries

and slightly different experimental setups. Although there were differences between experiments,

overall the data for 0.5 M NaCl LB proved to be rather consistent between different runs (Figure 2A,

Supplementary Figures S1-4, Supplementary Tables S3-S4, S8-S11) suggesting that a single

experiment would be sufficient in most cases.

Because TN-seq relies on the selection of strains adapted to certain growth conditions, competition

between mutants and subsequent depletion of the input library it is to be expected that this method

is ideal for exploring the long-term effects on growth instead of quick, short-term responses to

stresses. To evaluate how our TN-seq results compared to previous results from high-throughput

studies of *S. aureus* cells grown in high salt conditions, we determined the overlaps between our

two 0.5 M NaCl, 17 generation datasets and a microarray data set acquired from cells grown for ~6

generations in 2 M NaCl (Price-Whelan *et al.* 2013). To compare the datasets, the *S. aureus* strain

COL locus tags used in the microarray study were first converted to TCH1516 locus tags where

possible (Supplementary Table S14) and afterwards compared with the final two 0.5 M NaCl, 17

generation TN-seq datasets. The overlap between the downregulated (microarray) and dispensable

(TN-seq) genes was minimal in both replicates (1 and 7) and also very small for the upregulated and

essential genes (both 14). Three operons were consistently essential or upregulated between all

datasets: the *cap5*, *SAUSA300_0771-2*, and the *sda* operon. The *cap5* genes are involved in capsule

production (Chan *et al.* 2014) and this type of capsule has been previously reported to be induced

by the addition of salt (Pöhlmann-Dietze *et al.* 2000). Genes *SAUSA300_0771-2* are hypothetical

membrane proteins and presumably part of a threonine/serine exporter (Interpro database IDs:

IPR010619 and IPR024528) and the *sda* genes are annotated as L-serine dehydratase components

and a regulator protein. Taken together, the low overlap between the microarray and TN-seq data

(Supplementary Table S14) further highlights that in our screen we have identified a number of

previously unknown salt tolerance genes.

Most mutants of potential NaCl essential genes identified in the TN-seq screen proved to be

sensitive to 0.5 M NaCl (Figure 4A-D), confirming the effectiveness of the TN-seq screen. In

particular, previously uncharacterized genes such as *SAUSA300_0694*, *mgtE,* and the *957* gene could

be identified as essential during NaCl stress and their phenotypes could be complemented (Figure

5). In our study, we opted not to experimentally validate any of the dispensable genes, as the

verification of a growth advantage is more difficult to prove than a growth defect, but these genes

will provide interesting starting points for future research. In addition, it will be interesting to

investigate higher concentrations of NaCl to see if the same or different genes are flagged as

essential and dispensable.

Gene *957* was selected for further characterization as part of this study, because it had previously

not been linked to NaCl stress and had not been investigated before. As gene *957* and *lcpB* are co-

transcribed we initially assumed that protein 957 is involved in WTA attachment via LcpB but we

were unable to demonstrate such a link. Instead the *957* mutant exhibited increased muropeptide

crosslinking, which could be suppressed by certain *pbp2* SNPs suggesting involvement in

peptidoglycan synthesis. We can only speculate if the SNPs identified in *pbp2* increase or inhibit the

activity of Pbp2. However, judging by the number of independent SNPs, it seems more likely that

the mutations lead to a decrease rather than an increase in transglycosylase activity. In addition,

the results from the moenomycin sensitivity experiments using a sublethal antibiotic concentration

(Figure 8F) highlights that partial inactivation of the transglycosylase activity of Pbp2 improves the

growth of the *957* strain in high salt conditions. This is consistent with the idea that the obtained

pbp2 SNPs lead to a decrease in transglycosylase activity. The absence of frameshift or non-sense

mutations can be explained by the importance of the C-terminal transpeptidase domain of *pbp2*

(Pinho *et al.* 2001) and this is reflected in the transposon insertion distributions (Supplementary

Figure 7B & C). Intriguingly, the suppressor mutants show reduced growth compared to the *957*

mutant on moenomycin plates, presumably due to transglycosylation activity being reduced to

levels that are detrimental to the cell. At this point it is also unclear how mutations in the

transglycosylase domain alter the muropeptide pattern, since the peptide bonds are made by the

transpeptidase, not the transglycosylase domain. We hypothesize that a decrease in the efficiency

of the glycosylation activity and slowing down the glycan chain synthesis process will reduce the

efficiency of the subsequent transpeptidation process, resulting in decreased peptidoglycan

crosslinking in the *pbp2* SNP strains. It is also noteworthy that the *pbp2* gene becomes dispensable

under NaCl conditions (Supplementary Figure S1A and S4), an additional indicator that Pbp2 activity

needs to be avoided under high salt conditions.

Interestingly, the *957* mutant exhibited a slightly higher oxacillin MIC than the WT. This is in contrast

to what we expected as MICs are determined in 2% (0.342 M) salt medium and this should lead to

a growth disadvantage of the mutant. However, when seen in light of the increase in peptidoglycan

crosslinking, this could explain the growth advantage of the *957* mutant in the presence of this beta-

lactam antibiotic, as this could counter the inhibition of transpeptidases by oxacillin.

Although we were unable to determine the exact molecular mechanism, it is clear from our data

that gene *957* is involved in cell wall synthesis either directly or indirectly. More generally, our work

suggests that the rigidity of the peptidoglycan cell wall is an important factor to counter osmotic

stresses and it is conceivable that the more crosslinked cell wall of the *957* mutant is less elastic and

therefore easier to break under strong osmotic stress caused by NaCl exposure (Summarized in

Figure 9).

In conclusion, we have identified a number of previously uncharacterized factors involved in the osmotic stress response of *S. aureus,* which in particular highlighted the importance of the cell envelope and the data generated will provide a great resource for further studies of the staphylococcal osmotic stress response.

Material and Methods

Growth of bacteria

E. coli and *S. aureus* strains were streaked from frozen stock onto Lysogenic Broth (LB) or Tryptic Soya agar (TSA) plates, respectively. For all *E. coli* and most *S. aureus* experiments, the bacteria were grown in LB medium (LB normal, LBN, 10 g Tryptone, 5 g Yeast extract, 5 g NaCl) pH 7.5 or LB medium with extra 0.5 M NaCl, 0.5 M KCl or 1 M sucrose added. For molecular cloning and some other experiments (as indicated in the text), *S. aureus* strains were grown in TSB medium. Growth curves were done in 125 ml flasks with 20 ml of medium without antibiotics. Cultures were inoculated from overnight cultures to a starting OD_{600} of 0.01 and growth was followed by determining OD_{600} readings every one or two hours.

TN-seq experiments

A previously published highly-saturated *S. aureus* USA300 library with a mix of six different transposons was used (Santiago *et al.* 2015; Coe *et al.* 2019). In addition, a new transposon library with only the blunt transposon, was produced using the same techniques as described in Santiago *et al.* (Santiago *et al.* 2015) containing approximately 1.2 million individual clones.

In the first experiment (NaCl, KCl and sucrose stress), a tube of the blunt library was thawed on ice and used to set up a 20 ml pre-culture in LBN/Erm 10 µg/ml with a starting OD_{600} of 0.1. This starter culture was grown for 1 hour and subsequently used to inoculate 25 ml of either LBN or LB 0.5 M

NaCl without Erm to a starting OD_{600} of 0.00125. When the cultures reached an OD_{600} of

approximately 0.3, the cultures were back-diluted 1:100-fold into 40 ml fresh medium and 10-12 ml

harvested by centrifugation when the back-diluted culture reached an OD_{600} of approx. 1.0 (16

generations). The bacterial pellets were washed once with their respective growth medium and

stored at -20°C for further processing.

In the second experiment (NaCl, Replicate A), the multiplexed promoter transposon library was used

and Erm 5 µg/ml was added to the pre-culture and subsequent cultures in either LBN or LB

containing 0.5 M NaCl. Samples were collected at an OD_{600} of about 1.4 (17 generations).

The third experiment (NaCl, replicate B) was set up similar to replicate A, however the blunt library

was used in this case and Erm 5 µg/ml was only added to the pre-culture.

The library preparation for sequencing was done as described previously (Santiago *et al.* 2015).

Briefly, genomic DNA was isolated, cut with *Not*I, biotinylated adapters ligated, the DNA cut again

with *Mme*I, and fragments ligated to Illumina adapters and the products PCR amplified,

incorporating bar codes and the Illumina adaptor sequences P5 and P7. The samples were

sequenced on an Illumina HiSeq machine after spiking with 40% PhiX DNA. The data analysis was

done using the Tufts Galaxy Server and custom python scripts as described earlier (Santiago *et al.*

2015). To this end, the reads were trimmed up to the barcode and de-multiplexed by strain barcode.

Due to the variability in the DNA cleavage by the *Mme*I restriction enzyme, the reads that could not

be mapped to a barcode were trimmed by an additional base and the process repeated. Reads were

then mapped to the USA300_TCH1516 (CP000730.1) genome and a hop table was generated.

Statistical analysis (Mann-Whitney tests and Benjamini-Hochberg) was done using custom python

scripts (Santiago *et al.* 2015). Further exploration of the data was done using Excel and custom R

scripts.

S. aureus cell diameter measurements

For cell diameter measurements, *S. aureus* strains ANG4054, ANG4340, ANG4341 were grown overnight (14-20 hours) at 37°C with 180 rpm shaking in 5 ml LBN in the presence of 100 ng/ml Atet. 100 µl of these overnight cultures was transferred to a 1.5 ml reaction tube and 1.5 µl of BODIPY-FL-vancomycin (100 µg/ml in phosphate buffered saline) was added to each tube to stain the cell wall. After a 30 min static incubation step at 37°C, 1.5 µl of the suspension was pipetted onto a slide covered with a 1% agarose pad and analyzed by microscopy using an Axio Imager.A2 microscope with an EC Plan-Neofluar 100x/1.30 Oil M27 objective and images recorded using an AxioCam MR R3. Native CZI files were opened in FIJI (Schindelin *et al.* 2012) and cell diameters of 200 cells were measured using the line and measure tool. Only cells without any visible septa were measured. Experiment was conducted five times with independent biological cultures (n=5, 200 cells each).

CFU/OD correlations

The different *S. aureus* strains were grown as described for the cell diameter measurements. Optical densities of cultures were measured as well as 100 µl of 10^{-6} dilutions (made in their respective growth medium) plated onto TSA plates. The next day, the number of colonies were counted and the ratios of OD_{600} to CFU calculated. The experiment was conducted six time with independent biological samples (n=6).

Zymogram assays

The different *S. aureus* strains were grown over night in TSB. The next day, the bacteria from an OD_{600} equivalent of 20 were pelleted by centrifugation for 3 min at 17000 x *g*. The cells were washed twice with 600 µl PBS and subsequently suspended in 50 µl SDS sample buffer. The suspension was boiled for 20 min with interspersed shaking, centrifuged for 5 min at 17000 x *g* and 20 µl of each

sample was loaded onto Zymogram gels, which where were prepared as described previously

(Atilano *et al.* 2014). Gels were stained and de-stained using methylene blue and water. The

experiment was performed twice, and a representative result is shown.

WTA isolation and analysis

Overnight cultures of *S. aureus* strains ANG4054, ANG4340, ANG4341, ANG1575, ANG4290,

ANG4748, ANG4749, ANG4759 were prepared in LBN and used to inoculate 50 ml LBN or LB 0.4 M

NaCl to an OD_{600} of 0.01. The cultures were incubated at 37°C with shaking until they reached an

OD_{600} of 5 to 6. Where appropriate, 10 μg/ml chloramphenicol and 100 ng/ml Atet was also added

to the medium. The bacteria from 20-24 ml culture were harvested by centrifugation for 10 min at

7000 x *g*. The bacterial pellets were washed with 50 mM 2-(N-morpholino) ethanesulfonic acid

(MES) pH 6.5 buffer and stored at -20°C until further use. Pellets were then processed and WTA

separated on a 20% native polyacrylamide gel by electrophoresis on a Biorad Protein XL ii cell as

described previously (Covas *et al.* 2016) and WTA visualized by silver staining according to the

manufacturer protocol. Experiments were done with three biological replicates (n=3) and a

representative result is shown.

Peptidoglycan isolation and analysis

Peptidoglycan was prepared as described previously (Corrigan *et al.* 2011) with the following

modifications: 0.5 L (ANG4290, ANG4382, ANG4384) or 1.0 L (ANG5054, ANG4340, ANG4341) of LB

0.4 M NaCl medium (all strains) or LBN (ANG5054, ANG4340, ANG4341) medium were inoculated

to an OD_{600} of 0.01. For strains carrying derivatives of plasmid piTET, the medium was supplemented

with 100 ng/ml Atet starting from the pre-cultures. The cultures were grown at 37°C with shaking

at 180 rpm until they reached an OD_{600} of 2-3 and bacteria were subsequently harvested by

centrifugation. Chromatography of mutanolysin digested peptidoglycan was performed as

described previously using an Agilent 1260 infinity system (Corrigan *et al.* 2011) and muropeptide

peaks assigned according to de Jonge *et al.* (de Jonge *et al.* 1992). For the muropeptide

quantification, a baseline was drawn, the total peak area determined as well as the peak areas for

mono- and the different multimer peaks and calculated as percentage of the total peak area. The

peak area quantification was done three times for each HPLC chromatogram and average values

were calculated. Experiments were done with three biological replicates (n=3).

Data analysis, software and statistics

Data were processed with a combination of Python 3.6 (https://www.python.org), R 3.3 & 3.5

(https://www.r-project.org/), RStudio 1.1 & 1.2 (https://www.rstudio.com/), Prism 7 and 8

(https://www.graphpad.com), ChemStation OpenLab C.01.05 (https://www.agilent.com/) and

Microsoft Excel 15 and 16 (https://www.office.com/). Voronoi images were drawn using

Proteomaps V2 (Otto *et al.* 2010; Liebermeister *et al.* 2014) using custom TMD files. Microscopy

images were analyzed using FIJI 1.0 (https://fiji.sc/). Statistical analysis was performed with Prism

using appropriate tests as described in the figure legends.

Raising of Δ*957* suppressor strains

Multiple independent overnight cultures of strain LAC*Δ*957* (ANG4290) were prepared in 5 ml LBN.

These cultures were back diluted the next day into 20 ml LB 0.5 M NaCl to a starting OD_{600} of 0.05

and grown for 8-10 hours at 37°C with shaking until the cultures were slightly turbid. 50 µl of each

culture was passed into 5 ml of LBN and grown overnight. The next day, appropriate culture dilutions

were plated onto LBN agar plates and incubated overnight at 37°C. For each independent culture,

multiple single colonies were picked and used to inoculate individual wells of a 96-well microtiter

plates containing 100 µl LBN and the plates were subsequently incubated at 37°C in a 96-well plate

incubator with shaking at 650 rpm. The next day, a culture aliquot was stored at -80°C, and the

cultures were also diluted 1:50 in LB 0.5 M NaCl medium and 20 µl of these diluted cultures used to

inoculate 180 µl of LB 0.5 M NaCl medium. The plates were incubated overnight with shaking at

37°C and the next morning the growth of the potential *957* suppressor strains compared to that of

the original LAC*Δ*957* deletion strain (ANG4290), which showed low growth and the WT LAC*

(ANG1575) strain, which showed good growth. For each lineage, four suppressors that showed good

growth were streaked out from the previously frozen stocks and subsequently single colonies

selected to set up overnight cultures. The deletion of gene *SAUSA300_0957* was confirmed by PCR

and after performing growth curves in culture flasks, one strain that showed significant growth

improvement compared to the original LAC*Δ*957* (ANG4290) strain was selected from each lineage.

These strains were streaked out again for single colonies and used to inoculate an overnight culture,

giving rise to the independently raised suppressor strains ANG4381 through ANG4394. The

compensatory mutations for several of these suppressor strains were subsequently determined by

whole genome sequencing.

Transfer of *pbp2* SNPs by phage transduction

In order to demonstrate that the SNPs in *pbp2* are responsible for the growth rescue of strain

LAC*Δ*957* (ANG4290), we transferred two different *pbp2* SNPs by phage transduction into the

original LAC*Δ*957* strain as well as repaired these SNPs in the suppressor strains. This was done by

placing an Erm-marked transposon in proximity of the *pbp*2 gene allowing at a certain frequency for

the co-transduction of the Erm marker and the *pbp2* SNP. To this end, the NTML transposon mutant

library strain NE789 (Fey *et al.* 2013) was used, which harbors a transposon insertion in

SAUSA300_1332 (putative exonuclease) about 10 kbp upstream of the *pbp2* gene. *SAUSA300_1332*

is expected to be unrelated to salt stress and the cell wall synthesis machinery but close enough to

lead to an intermediate rate of co-transference with the *pbp2* SNPs. The transposon from NE789

was transduced into WT LAC* (Boles *et al.* 2010) and strain LAC*Δ*957* (ANG4290) using phage Φ85

yielding strains ANG4561 and ANG452, respectively. The transposon was also transduced into the

suppressor strains ANG4382 and ANG4384 transferring either *SAUSA300_1332::Tn* only, yielding

strains ANG4527 and 4528 or transferring *SAUSA300_1332::Tn* as well as replacing the *pbp2* SNP

with a WT *pbp2* allele yielding *pbp2* repaired strains ANG4557 and ANG4568. Lysates from strains

ANG4527 and 4528 (suppressors containing *pbp2* SNPs and *SAUSA300_1332::Tn*) were used to

transfer the *pbp2* SNPs back into a clean LAC*Δ*957* strain background yielding strains ANG4624 and

ANG4625 and into WT LAC*, yielding strains ANG4563 and ANG4564. Successful repair or transfer

of the SNPs was checked by PCR and subsequent restriction digest of the product choosing enzymes

that recognize sites present in either the WT or SNP allele (ANG4382 *pbp2* SNP: BsrI. Recognition

site CCWGG, additional site introduced by SNP; ANG4384 *pbp2* SNP: SspI. Recognition site: AATATT,

site missing in SNP).

Determination of oxacillin MICs using Etest strips

S. aureus strains LAC* piTET (ANG4054), LAC*Δ*957* piTET (ANG4340) and the complementation

strain LAC*Δ*957* piTET-*957* (ANG4341) were grown overnight (22 hours) in 5 ml TSB containing 10

µg/ml chloramphenicol and 100 ng/ml Atet. The next day, the cultures were diluted to an OD_{600} of

0.1 in sterile PBS buffer and streaked with cotton swabs onto cation adjusted Mueller-Hinton agar

plates supplemented with 2% NaCl and 100 ng/ml Atet. One M.I.C.Evaluator strip was placed on

each plate and the plates were incubated for 24 hours at 35°C. MICs were then read directly from

the strips. The experiment was done with 4 biological replicates (n=4).

Moenomycin growth improvement test

S. aureus strains LAC* (ANG1575), LAC* Δ*957* (ANG4290), LAC* Δ*957* S2 (ANG4382) and LAC* Δ*957*

S4 (ANG4384) were grown overnight (18 hours) in 5 ml LBN. The next day, the cells were diluted to

an OD_{600} of 0.01 in LBN, grown until mid-exponential phase (OD_{600} 0.4-0.6) and normalized to an

OD_{600} of 0.1. The cells were 10-fold serially diluted and 5 μl of each dilution spotted onto LB agar

containing 0.5 M NaCl with either 0.02 μg/ml moenomycin (mix of moenomycin A, A12, C1, C3 and

C4, Santa Cruz Biotechnology) or no moenomycin. Plates were incubated at 37°C overnight (18-22

hours) and photographed. The experiment was done with 3 biological replicates (n=3) and one

representative result is shown.

Whole genome sequencing

Genomic DNA was either isolated using the Promega Gene Wizard kit according to the manufacturer

instructions or using chloroform-isoamylalcohol as described previously (Schuster and Bertram

2014). DNA was sent off for whole genome sequencing to MicrobesNG, Birmingham, U.K. or libraries

prepared using the Illumina Nextera DNA kit and sequenced at the London Institute of Medical

Sciences. Short reads were trimmed in CLC workbench genomics (Qiagen), then mapped to a custom

S. aureus USA300 LAC* reference genome generated in a previous study (Bowman *et al.* 2016) and

single nucleotide polymorphisms called based on at least 80% frequency of occurrence. This list was

compared to a manually curated list of well non false-positives and these entries were removed.

Nebraska Transposon Mutant Library (NTML) strains and complementation strains

All Nebraska transposon mutant library (NTML) strains and primers used in this study are listed in

Supplementary Tables S12, S15 and S16. Transposon insertions in the respective genes were

confirmed by PCRs. For complementation analysis, NTML strains NE535 (JE2 *SAUSA300_0867*::*Tn*),

NE867 (JE2 *SAUSA300_0483*::*Tn*) were transformed as controls with the empty plasmid pCL55 (Lee

et al. 1991), yielding strains ANG4326 and ANG4328 and strains NE188 (JE2 *SAUSA300_0481*::*Tn*),

NE251 (JE2 SAUSA300_0482::*Tn*), NE526 (JE2 *SAUSA300_0694*::*Tn*), NE736 (JE2

SAUSA300_0910::*Tn*), NE1384 (JE2 *SAUSA300_0957*::*Tn*) with plasmid piTET (Gründling and

Schneewind 2007), yielding ANG4308, ANG4310, ANG4377, ANG4336, ANG4338. The WT strain JE2

(Fey *et al.* 2013) was also transformed with both plasmids yielding JE2 pCL55 (ANG4325) and JE2

piTET (ANG4307).

The complementation plasmid piTET-*481* for strain NE188 (JE2 *SAUSA300_0481::Tn*) was

constructed by amplifying the *SAUSA300_0481* gene using primers P2378/P2379, digesting the

product with AvrII/BglII and ligating the fragment with plasmid piTET cut with the same enzymes.

The plasmid was then introduced into *E. coli* strain XL1-Blue, yielding ANG4139. After shuttling the

plasmid through *E. coli* strain IM08B (Monk *et al.* 2012), creating strain ANG4150, the plasmid was

integrated into the *geh* locus of NE188 (JE2 *SAUSA300_0481*::*Tn*) yielding strain ANG4309. The

complementation plasmid piTET-*482* for strain NE251 (JE2 SAUSA300_0482::*Tn*), piTET-*694* for

strain NE526 (JE2 *SAUSA300_0694*::*Tn*), piTET-*910* for strain NE736 (JE2 *SAUSA300_0910*::*Tn*), and

piTET-*957* for strain NE1384 (JE2 *SAUSA300_0957*::*Tn*) were constructed in a similar manner using

primer pairs P2380/P2381 (*SAUSA300_0482*), P2388/P2389 (*SAUSA300_0694*), P2384/P2385

(*SAUSA300_0910*) and P2386/P2387 (*SAUSA300_0957*), respectively. The complementation

plasmids were recovered in *E. coli* XL1-Blue yielding strains ANG4140 (SAUSA300_0482), ANG4142

(*SAUSA300_0694*), ANG4144 (*SAUSA300_0910*) and ANG4145 (*SAUSA300_0957*), shuttled through

E. coli IM08B giving strains ANG4151 (*SAUSA300_0482*), ANG4153 (*SAUSA300_0694*), ANG4155

(*SAUSA300_0910*) and ANG4156 (*SAUSA300_0957*) and finally introduced in the respective NTML

strain yielding the complementation strains ANG4311 (*SAUSA300_0482*), ANG4378

(*SAUSA300_0694*), ANG4337 (*SAUSA300_0910*) and ANG4339 (*SAUSA300_0957*).

Complementation plasmid pCL55-*483* for strain NE867 (JE2 *SAUSA300_0483*::*Tn*) was made by

fusing PCR products of the operon promoter (in front of *SAUSA300_0481*) amplified with primers

P2588/P2553 and the *SAUSA300_0483* gene, amplified with primers P2554/P2589 together using

primers P2588/P2589. The resulting fragment was cloned into pCL55 using EcoRI/BamHI restriction

sites. The plasmid was recovered in XL1-Blue, creating strain ANG4291, shuttled through *E. coli*

IM08B (ANG4292) and subsequently introduced into NE867 (JE2 *SAUSA300_0483*::*Tn*) to create

strain ANG4327. The complementation plasmid pCL55-*867* for strain NE535 (JE2

SAUSA300_0867::*Tn*) was constructed in a similar manner, using primers P2590/P2556 and

P2557/P2591 to amplify the promoter and *SAUSA300_0867* gene, which were subsequently fused

in a second PCR using primers P2590/P2591. The resulting fragment was cloning into pCL55 using

EcoRI/BamHI restriction sites and the resulting plasmid recovered in *E. coli* XL1-Blue (ANG4293),

shuttling through IM08B (ANG4294) and finally introduced into NE535 (JE2 *SAUSA300_0867*::*Tn*),

creating the complementation strain ANG4329.

Construction of *S. aureus* gene deletion and complementation strains

S. aureus strains with in-frame gene deletions were constructed by allelic exchange using plasmids

pIMAY (Monk *et al.* 2012) and pIMAY* (Schuster *et al.* 2019). The gene deletion plasmids were

designed to contain approximately 1000 bp up- and downstream regions around the deletion site,

amplified from LAC* genomic DNA (Boles *et al.* 2010), and the first and last 30 bp of the open reading

frame to be deleted.

For construction of plasmid pIMAY-Δ*957*, the up- and downstream regions of *SAUSA300_0957* were

amplified using primers P2370/P2371 and P2372/P2373, spliced together in a second PCR using

primers P2370/P2373 and cloned into pIMAY using XmaI/EcoRI. Plasmid pIMAY-Δ957 was recovered

in *E. coli* XL1-Blue, creating strain ANG4147, shuttled through *E. coli* IM08B (ANG4159) and

subsequently introduced into S*. aureus* LAC*Δ*957*. The allelic exchange to delete *SAUSA300_0957*

and to create strain LAC*Δ*957* (ANG4290) was performed as previously described (Monk *et al.*

2012). For complementation analysis, the above described plasmid piTET-*957* was integrated into

the chromosome of strain LAC*Δ*957*, giving rise to the complementation strain LAC*Δ*957* piTET-

957 (ANG4341). As control, the empty plasmid piTET from ANG4163 was also integrated into the

chromosome of LAC*Δ*957* yielding strain LAC*Δ*957* piTET (ANG4340). The *lcpB* (SAUSA300_0958)

gene was deleted in a similar manner using primers P2844/P2845 and P2846/P2847 for the first and

primers P2844/P2847 for the second PCR and cloning the fragment XmaI/EcoRI into pIMAY. Plasmid

pIMAY-Δ*lcpB* was recovered in *E. coli* XL1blue (ANG4740), shuttled through *E. coli* IM08B (ANG4742)

and introduced into the *S. aureus* LAC* (ANG4744) and finally yielding the *lcpB* deletion strain

LAC*Δ*lcpB* (ANG4748).

Plasmid pIMAY*-Δ*957-958* was constructed for the production of an *SAUSA300_0957-0958* (*lcpB*)

double mutant strain in which the first 30 bases of *SAUSA300_0957* are fused to the last 30 bp of

SAUSA300_0958 (*lcpB*). This was done using primers P2844/P2848 and P2849/P2373 in the first and

primers P2844/P2373 in a second PCR. The fragment was cloned using XmaI/EcoRI into pIMAY*

(Schuster *et al.* 2019) and the resulting plasmid pIMAY*-Δ*957-958* recovered in *E. coli* XL1-Blue

(ANG4741). The plasmid was shuttled through *E. coli* IM08B (ANG4743), introduced into *S. aureus*

LAC* (ANG4745) and following the allelic exchange procedure yielding the *S. aureus* strain

LAC*Δ*957-958* (*lcpB*) (ANG4749).

The plasmid pIMAY-Δ*mgtE* was designed to contain up- and downstream regions of

SAUSA300_0910 (*mgtE*) fused together by the first and last 30 bp of the gene in order to create a

LAC*Δ*mgtE* mutant. For this, primer pairs P2374/P2375 and P2376/P2377 were used to amplify up-

and downstream regions and subsequently spliced together by PCR using primers P2374/P2377.

The PCR fragment was digested with XmaI/EcoRI and ligated into pIMAY cut with the same enzymes.

The plasmid pIMAY-Δ*mgtE* was recovered in *E. coli* XL1-Blue yielding strain ANG4146 and shuttled

through *E. coli* IM08B yielding strain ANG4158. The plasmid pIMAY-Δ*mgtE* was then introduced into

LAC* yielding strain LAC*Δ*SAUSA300_0910* (*mgtE*) (ANG4422) after the knockout procedure. To

create an empty vector-containing control strain and a plasmid-based complementation strain, the

plasmid piTET and piTET-*910* from strains ANG4163 and ANG4155 were isolated and used to

transform strain LAC*Δ*SAUSA300_0910* (*mgtE*) (ANG4422), yielding strains ANG4445 (LAC*Δ*mgtE*

piTET) and ANG4446 (LAC*Δ*mgtE* piTET-*mgtE).*

The *tagO* gene (*SAUSA300_0731*) was inactivated using the targetron intron homing system (Yao *et*

al. 2006) or by allelic exchange using pIMAY* (Schuster *et al.* 2019). For the targetron mutagenesis,

plasmid pNL9164-*tagO* was produced by amplifying the upstream targetron fragment with primers

P2887 and the EBS universal primer (P2890) and the downstream fragment using primers

P2888/P2889, the fragments were fused using the IBS primer (P2887) and the EBS1d primer (P2888).

The final fragment was cloned using HindIII/BsrGI into pNL9164 (Yao *et al.* 2006). Plasmid pNL9164-

tagO was recovered in *E. coli* XL1-Blue, yielding strain ANG4703, introduced into *E. coli* IM08B to

give rise to strain ANG4704 and finally introduced into *S. aureus* LAC* (Boles *et al.* 2010) to create

strain LAC* pNL9164-*tagO* (ANG4751). The final *tagO* targetron mutant strain LAC**tagO*::targetron

(ANG4753) was generated using the method described by Yao *et al.* (Yao *et al.* 2006). For the

construction of *S. aureus* strain LAC*Δ*tagO* with an in-frame deletion in *tagO,* the allelic exchange

plasmid pIMAY*-Δ*tagO* was used. To this end, up- and downstream *tagO* fragments were produced

using primer pairs P2883/P2884 and P2885/P2886 and introduced by Gibson cloning into

XmaI/EcoRI pre-cut pIMAY*. Plasmid pIMAY*-Δ*tagO* was recovered in *E. coli* XL1-Blue, generating

strain ANG4755. After shuttling through *E. coli* strain IM08B (ANG4756), the plasmid was introduced

into *S. aureus* LAC* to create strain ANG4757. Finally, the *tagO* locus was deleted by allelic exchange,

generating strain LAC*Δ*tagO* (ANG4759). Because initial generation of the mutant failed, cells were

plated unselectively onto TSA plates and incubated at room temperature (18-25°C) for about two

weeks to enable colony differentiation between WT and Δ*tagO* mutant strains. Upon prolonged

incubation, *tagO* mutant strain colonies have an opaque appearance.

Data access

The whole genome sequencing data were deposited at the European Nucleotide Archive (ENA)

under accession id ERP115099 and TN-seq data in the short read archive (SRA) at the National

Center for Biotechnology Information (NCBI) under BioProject ID PRJNA544248. Analyzed tables can

be found in the supplementary files. For other data, please contact the corresponding author.

Acknowledgements

This research was funded by the Wellcome Trust through grants 100289 and 210671/Z/18/Z to A.G.

and the NIH grant P01AI083214 to S.W.. C.F.S. is funded by the German Research Foundation

[Deutsche Forschungsgemeinschaft (DFG)] through grant SCHU 3159/1–1 and F.C.M.K. was

supported by a Wellcome trust Ph.D. studentship. We thank Microbes NG and the London Institute

of Medical Sciences for providing whole genome sequencing services and the Tufts sequencing

facility for the sequencing services for the TN-seq experiments. We also would like to thank Ute

Bertsche for kind advice on HPLC analyses.

Author contributions

C.F.S., D.M.W., F.C.K. and A.G. performed the research; C.F.S., D.M.W., F.C.M.K., M.S., S.W. and A.G.

analyzed the data; C.F.S. and A.G. designed the research; C.F.S. and A.G. wrote the paper. All authors

approved the final version of the manuscript.

Disclosure declaration

The authors declare not conflict of interest.

References

Adams MR, Moss MO. 2008. 7.14.2 The Organism and its Characteristics. In *Food Microbiology (3rd Edition)*. Royal Society of Chemistry.

Anderson CB, Witter LD. 1982. Glutamine and proline accumulation by *Staphylococcus aureus* with reduction in water activity. *Appl Environ Microbiol* **43**: 1501-1503.

Atilano ML, Pereira PM, Vaz F, Catalão MJ, Reed P, Grilo IR, Sobral RG, Ligoxygakis P, Pinho MG,
Filipe SR. 2014. Bacterial autolysins trim cell surface peptidoglycan to prevent detection by
the *Drosophila* innate immune system. *Elife* **3**: e02277.
Bae JH, Miller KJ. 1992. Identification of two proline transport systems in *Staphylococcus aureus* and
their possible roles in osmoregulation. *Appl Environ Microbiol* **58**: 471-475.
Berger-Bächi B, Barberis-Maino L, Strässle A, Kayser FH. 1989. FemA, a host-mediated factor
essential for methicillin resistance in *Staphylococcus aureus*: molecular cloning and
characterization. *Mol Gen Genet* **219**: 263-269.
Boles BR, Thoendel M, Roth AJ, Horswill AR. 2010. Identification of genes involved in polysaccharide-
independent *Staphylococcus aureus* biofilm formation. *PLoS One* **5**: e10146.
Bowman L, Zeden MS, Schuster CF, Kaever V, Grundling A. 2016. New Insights into the Cyclic Di-
adenosine Monophosphate (c-di-AMP) Degradation Pathway and the Requirement of the
Cyclic Dinucleotide for Acid Stress Resistance in *Staphylococcus aureus*. *J Biol Chem* **291**:
26970-26986.
Brown S, Santa Maria JP, Jr., Walker S. 2013. Wall teichoic acids of gram-positive bacteria. *Annu Rev
Microbiol* **67**: 313-336.
Chan YG, Frankel MB, Dengler V, Schneewind O, Missiakas D. 2013. *Staphylococcus aureus* mutants
lacking the LytR-CpsA-Psr family of enzymes release cell wall teichoic acids into the
extracellular medium. *J Bacteriol* **195**: 4650-4659.
Chan YG, Kim HK, Schneewind O, Missiakas D. 2014. The capsular polysaccharide of *Staphylococcus
aureus* is attached to peptidoglycan by the LytR-CpsA-Psr (LCP) family of enzymes. *J Biol
Chem* **289**: 15680-15690.
Christian JH, Waltho JA. 1964. The Composition of *Staphylococcus aureus* in Relation to the Water
Activity of the Growth Medium. *J Gen Microbiol* **35**: 205-213.
Coe KA, Lee W, Komazin-Meredith G, Meredith TC, Grad YH, Walker S. 2019. Comparative Tn-Seq
reveals common daptomycin resistance determinants in *Staphylococcus aureus* despite
strain-dependent differences in essentiality of shared cell envelope genes. *bioRxiv*
doi:10.1101/648246: 648246.
Corrigan RM, Abbott JC, Burhenne H, Kaever V, Gründling A. 2011. c-di-AMP is a new second
messenger in *Staphylococcus aureus* with a role in controlling cell size and envelope stress.
PLoS Pathog **7**: e1002217.
Covas G, Vaz F, Henriques G, Pinho MG, Filipe SR. 2016. Analysis of Cell Wall Teichoic Acids in
Staphylococcus aureus. *Methods Mol Biol* **1440**: 201-213.
Csonka LN. 1989. Physiological and genetic responses of bacteria to osmotic stress. *Microbiol Rev
53: 121-147.
Csonka LN, Hanson AD. 1991. Prokaryotic osmoregulation: genetics and physiology. *Annu Rev
Microbiol* **45**: 569-606.
de Jonge BL, Chang YS, Gage D, Tomasz A. 1992. Peptidoglycan composition of a highly methicillin-
resistant *Staphylococcus aureus* strain. The role of penicillin binding protein 2A. *J Biol Chem
267: 11248-11254.
Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, Bayles KW. 2013. A genetic resource
for rapid and comprehensive phenotype screening of nonessential *Staphylococcus aureus
genes. *MBio* **4**: e00537-00512.
Graham JE, Wilkinson BJ. 1992. *Staphylococcus aureus* osmoregulation: roles for choline, glycine
betaine, proline, and taurine. *J Bacteriol* **174**: 2711-2716.
Gründling A, Schneewind O. 2007. Genes required for glycolipid synthesis and lipoteichoic acid
anchoring in *Staphylococcus aureus*. *J Bacteriol* **189**: 2521-2530.
Holland LM, Conlon B, O'Gara JP. 2011. Mutation of *tagO* reveals an essential role for wall teichoic
acids in *Staphylococcus epidermidis* biofilm development. *Microbiology* **157**: 408-418.

Ishibashi M, Kurokawa K, Nishida S, Ueno K, Matsuo M, Sekimizu K. 2007. Isolation of temperature-
sensitive mutations in *murC* of *Staphylococcus aureus*. *FEMS Microbiology Letters* **274**: 204-
209.
Kho K, Meredith TC. 2018. Salt-Induced Stress Stimulates a Lipoteichoic Acid-Specific Three-
Component Glycosylation System in *Staphylococcus aureus*. *J Bacteriol* **200**.
Koprivnjak T, Mlakar V, Swanson L, Fournier B, Peschel A, Weiss JP. 2006. Cation-induced
transcriptional regulation of the dlt operon of *Staphylococcus aureus*. *J Bacteriol* **188**: 3622-
3630.
Lee CY, Buranen SL, Ye ZH. 1991. Construction of single-copy integration vectors for *Staphylococcus*
aureus. *Gene* **103**: 101-105.
Liebermeister W, Noor E, Flamholz A, Davidi D, Bernhardt J, Milo R. 2014. Visual account of protein
investment in cellular functions. *Proc Natl Acad Sci U S A* **111**: 8488-8493.
Lovering AL, De Castro L, Strynadka NC. 2008. Identification of dynamic structural motifs involved in
peptidoglycan glycosyltransfer. *J Mol Biol* **383**: 167-177.
Mäder U, Nicolas P, Depke M, Pané-Farré J, Debarbouille M, van der Kooi-Pol MM, Guérin C,
Dérozier S, Hiron A, Jarmer H et al. 2016. *Staphylococcus aureus* Transcriptome Architecture:
From Laboratory to Infection-Mimicking Conditions. *PLoS Genet* **12**: e1005962.
Madiraju MV, Brunner DP, Wilkinson BJ. 1987. Effects of temperature, NaCl, and methicillin on
penicillin-binding proteins, growth, peptidoglycan synthesis, and autolysis in methicillin-
resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* **31**: 1727-1733.
Mayer S, Steffen W, Steuber J, Götz F. 2015. The *Staphylococcus aureus* NuoL-like protein MpsA
contributes to the generation of membrane potential. *J Bacteriol* **197**: 794-806.
Measures JC. 1975. Role of amino acids in osmoregulation of non-halophilic bacteria. *Nature* **257**:
398-400.
Miller KJ, Zelt SC, Bae J-H. 1991. Glycine betaine and proline are the principal compatible solutes of
Staphylococcus aureus. *Current Microbiology* **23**: 131-137.
Mistretta N, Brossaud M, Telles F, Sanchez V, Talaga P, Rokbi B. 2019. Glycosylation of
Staphylococcus aureus cell wall teichoic acid is influenced by environmental conditions. *Sci*
Rep **9**: 3212.
Mitchell P, Moyle J. 1959. Permeability of the envelopes of *Staphylococcus aureus* to some salts,
amino acids, and non-electrolytes. *J Gen Microbiol* **20**: 434-441.
Mlynek KD, Callahan MT, Shimkevitch AV, Farmer JT, Endres JL, Marchand M, Bayles KW, Horswill
AR, Kaplan JB. 2016. Effects of Low-Dose Amoxicillin on *Staphylococcus aureus* USA300
Biofilms. *Antimicrob Agents Chemother* **60**: 2639-2651.
Monk IR, Shah IM, Xu M, Tan MW, Foster TJ. 2012. Transforming the untransformable: application
of direct transformation to manipulate genetically *Staphylococcus aureus* and
Staphylococcus epidermidis. *MBio* **3**.
Murakami K, Fujimura T, Doi M. 1994. Nucleotide sequence of the structural gene for the penicillin-
binding protein 2 of *Staphylococcus aureus* and the presence of a homologous gene in other
staphylococci. *FEMS Microbiol Lett* **117**: 131-136.
Ochiai T. 2000. *Staphylococcus aureus* produces autolysin-susceptible cell walls during growth in a
high-NaCl and low-Ca2+ concentration medium. *Microbiol Immunol* **44**: 97-104.
Onyango LA, Hugh Dunstan R, Roberts TK, Macdonald MM, Gottfries J. 2013. Phenotypic variants of
staphylococci and their underlying population distributions following exposure to stress.
PLoS One **8**: e77614.
Oshida T, Sugai M, Komatsuzawa H, Hong YM, Suginaka H, Tomasz A. 1995. A *Staphylococcus aureus*
autolysin that has an N-acetylmuramoyl-L-alanine amidase domain and an endo-beta-N-
acetylglucosaminidase domain: cloning, sequence analysis, and characterization. *Proc Natl*
Acad Sci U S A **92**: 285-289.

Ostash B, Walker S. 2010. Moenomycin family antibiotics: chemical synthesis, biosynthesis, and
biological activity. *Nat Prod Rep* **27**: 1594-1617.
Otto A, Bernhardt J, Meyer H, Schaffer M, Herbst FA, Siebourg J, Mäder U, Lalk M, Hecker M, Becher
D. 2010. Systems-wide temporal proteomic profiling in glucose-starved Bacillus subtilis. *Nat*
Commun **1**: 137.
Over B, Heusser R, McCallum N, Schulthess B, Kupferschmied P, Gaiani JM, Sifri CD, Berger-Bächi B,
Stutzmann Meier P. 2011. LytR-CpsA-Psr proteins in *Staphylococcus aureus* display partial
functional redundancy and the deletion of all three severely impairs septum placement and
cell separation. *FEMS Microbiol Lett* **320**: 142-151.
Pinho MG, Filipe SR, de Lencastre H, Tomasz A. 2001. Complementation of the essential
peptidoglycan transpeptidase function of penicillin-binding protein 2 (PBP2) by the drug
resistance protein PBP2A in *Staphylococcus aureus*. *J Bacteriol* **183**: 6525-6531.
Pöhlmann-Dietze P, Ulrich M, Kiser KB, Döring G, Lee JC, Fournier JM, Botzenhart K, Wolz C. 2000.
Adherence of *Staphylococcus aureus* to endothelial cells: influence of capsular
polysaccharide, global regulator agr, and bacterial growth phase. *Infect Immun* **68**: 4865-
4871.
Pourkomailian B, Booth IR. 1992. Glycine betaine transport by *Staphylococcus aureus*: evidence for
feedback regulation of the activity of the two transport systems. *Microbiology* **140 (Pt 11)**:
3131-3138.
Price-Whelan A, Poon CK, Benson MA, Eidem TT, Roux CM, Boyd JM, Dunman PM, Torres VJ,
Krulwich TA. 2013. Transcriptional profiling of *Staphylococcus aureus* during growth in 2 M
NaCl leads to clarification of physiological roles for Kdp and Ktr K+ uptake systems. *MBio* **4**.
Rahman MM, Hunter HN, Prova S, Verma V, Qamar A, Golemi-Kotra D. 2016. The *Staphylococcus*
aureus Methicillin Resistance Factor FmtA Is a d-Amino Esterase That Acts on Teichoic Acids.
MBio **7**: e02070-02015.
Santiago M, Matano LM, Moussa SH, Gilmore MS, Walker S, Meredith TC. 2015. A new platform for
ultra-high density *Staphylococcus aureus* transposon libraries. *BMC Genomics* **16**: 252.
Schaefer K, Matano LM, Qiao Y, Kahne D, Walker S. 2017. In vitro reconstitution demonstrates the
cell wall ligase activity of LCP proteins. *Nat Chem Biol* **13**: 396-401.
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,
Saalfeld S, Schmid B et al. 2012. Fiji: an open-source platform for biological-image analysis.
Nat Methods **9**: 676-682.
Schuster CF, Bertram R. 2014. Fluorescence based primer extension technique to determine
transcriptional starting points and cleavage sites of RNases *in vivo*. *J Vis Exp*
doi:10.3791/52134: e52134.
Schuster CF, Howard SA, Gründling A. 2019. Use of the counter selectable marker PheS* for genome
engineering in *Staphylococcus aureus*. *Microbiology* doi:10.1099/mic.0.000791.
Schwan WR, Lehmann L, McCormick J. 2006. Transcriptional activation of the *Staphylococcus aureus*
putP gene by low-proline-high osmotic conditions and during infection of murine and human
tissues. *Infect Immun* **74**: 399-409.
Scybert S, Pechous R, Sitthisak S, Nadakavukaren MJ, Wilkinson BJ, Jayaswal RK. 2003. NaCl-sensitive
mutant of *Staphylococcus aureus* has a Tn917-lacZ insertion in its ars operon. *FEMS Microbiol*
Lett **222**: 171-176.
Sleator RD, Hill C. 2002. Bacterial osmoadaptation: the role of osmolytes in bacterial stress and
virulence. *FEMS Microbiol Rev* **26**: 49-71.
Stapleton MR, Horsburgh MJ, Hayhurst EJ, Wright L, Jonsson IM, Tarkowski A, Kokai-Kun JF, Mond
JJ, Foster SJ. 2007. Characterization of IsaA and SceD, two putative lytic transglycosylases of
Staphylococcus aureus. *J Bacteriol* **189**: 7316-7325.

Stewart CM, Cole MB, Legan JD, Slade L, Schaffner DW. 2005. Solute-specific effects of osmotic
stress on *Staphylococcus aureus*. *J Appl Microbiol* **98**: 193-202.
Stewart CM, Cole MB, Legan JD, Slade L, Vandeven MH, Schaffner DW. 2002. *Staphylococcus aureus*
growth boundaries: moving towards mechanistic predictive models based on solute-specific
effects. *Appl Environ Microbiol* **68**: 1864-1871.
Tong SY, Davis JS, Eichenberger E, Holland TL, Fowler VG, Jr. 2015. *Staphylococcus aureus* infections:
epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev*
28: 603-661.
Townsend DE, Wilkinson BJ. 1992. Proline transport in *Staphylococcus aureus*: a high-affinity system
and a low-affinity system involved in osmoregulation. *J Bacteriol* **174**: 2702-2710.
Vaish M, Price-Whelan A, Reyes-Robles T, Liu J, Jereen A, Christie S, Alonzo F, 3rd, Benson MA, Torres
VJ, Krulwich TA. 2018. Roles of *Staphylococcus aureus* Mnh1 and Mnh2 Antiporters in Salt
Tolerance, Alkali Tolerance, and Pathogenesis. *J Bacteriol* **200**.
Vijaranakul U, Nadakavukaren MJ, de Jonge BL, Wilkinson BJ, Jayaswal RK. 1995. Increased cell size
and shortened peptidoglycan interpeptide bridge of NaCl-stressed *Staphylococcus aureus*
and their reversal by glycine betaine. *J Bacteriol* **177**: 5116-5121.
Vijaranakul U, Xiong A, Lockwood K, Jayaswal RK. 1998. Cloning and nucleotide sequencing of a
Staphylococcus aureus gene encoding a branched-chain-amino-acid transporter. *Appl*
Environ Microbiol **64**: 763-767.
Vollmer W, Blanot D, de Pedro MA. 2008. Peptidoglycan structure and architecture. *FEMS Microbiol*
Rev **32**: 149-167.
Wengender PA, Miller KJ. 1995. Identification of a PutP proline permease gene homolog from
Staphylococcus aureus by expression cloning of the high-affinity proline transport system in
Escherichia coli. *Appl Environ Microbiol* **61**: 252-259.
Wertheim HF, Melles DC, Vos MC, van Leeuwen W, van Belkum A, Verbrugh HA, Nouwen JL. 2005.
The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infect Dis* **5**: 751-762.
Wetzel KJ, Bjorge D, Schwan WR. 2011. Mutational and transcriptional analyses of the
Staphylococcus aureus low-affinity proline transporter OpuD during *in vitro* growth and
infection of murine tissues. *FEMS Immunol Med Microbiol* **61**: 346-355.
Wyke AW. 1984. Isolation of five penicillin-binding proteins from *Staphylococcus aureus*. *FEMS*
Microbiology Letters **22**: 133-138.
Yao J, Zhong J, Fang Y, Geisinger E, Novick RP, Lambowitz AM. 2006. Use of targetrons to disrupt
essential and nonessential genes in *Staphylococcus aureus* reveals temperature sensitivity
of Ll.LtrB group II intron splicing. *RNA* **12**: 1271-1281.
Zeden MS, Schuster CF, Bowman L, Zhong Q, Williams HD, Gründling A. 2018. Cyclic di-adenosine
monophosphate (c-di-AMP) is required for osmotic regulation in *Staphylococcus aureus* but
dispensable for viability in anaerobic conditions. *J Biol Chem* **293**: 3180-3200.

Figure legends

Fig. 1. Different sets of genes are conditionally essential for the growth of *S. aureus* under NaCl, KCl or sucrose stress conditions.

(A) Workflow of the TN-seq experiment. A TN-seq library was pre-cultured and then used to inoculate LBN, LB containing 0.5 M NaCl, 0.5 M KCl or 1 M sucrose. Cells were kept in exponential phase for 16 generations, harvested and transposon insertion sites determined by high throughput sequencing.

(B) Growth curves. For the TN-seq experiment (n=1), the growth of an *S. aureus* library cultured in LBN, LB 0.5M NaCl, LB 0.5 M KCl or LB 1.0 M sucrose medium was followed by taking OD_{600} measurements at timed intervals. The cultures were back-diluted once when they reached an OD_{600}

of approximately 0.3. The plotted OD_{600} values were calculated by multiplying the measured OD_{600}

with the dilution factor. The dotted line indicates the 16 generation cut-off at which cultures were

harvested.

(C) Circular plot showing the transposon insertion density along the *S. aureus* genome under

different osmotic stress conditions. The two outer rings depict genes located on the (+) or (-) strand

in *S. aureus* strain USA300 FPR3757. The inner four rings show the histograms of transposon

insertions on a per gene basis after growth of the library in LBN (orange), 0.5 M NaCl (brown), 0.5

M KCl (blue) and 1 M sucrose (pink) medium for 16 generations.

(D) Volcano plots of conditionally essential and dispensable genes following growth of *S. aureus* in

0.5 M NaCl, 0.5 M KCl or 1 M sucrose medium and compared to LBN conditions. q-values represent

Benjamini-Hochberg corrected p-values, and $\log_2$ (fold changes) indicate essential genes (negative

values) or dispensable genes (positive values). The black horizontal line indicates a q-value of 0.05,

which was deemed the significance level. Vertical lines indicate 2- (green), 5- (orange) or 10- (red)

fold changes in either direction. Each dot represents one gene and coloring follows the fold-change

scheme whenever the q-value threshold was met. Very small q-values were truncated to fit onto

the graph. The TN-seq experiments were conducted once (n=1).

(E) Venn diagrams showing overlaps of conditionally essential and dispensable genes during

prolonged NaCl, KCl or sucrose stress. Genes with a 2-fold decrease (essential genes) or a 2-fold

increase (dispensable genes) in transposon insertions under each stress condition compared to the

LBN condition and a q-value of ≤0.05 were determined and the overlap between these gene lists

displayed in Venn diagrams.

Conditionally essential genes

Fig. 2. Voronoi diagram-based visualization of conditionally essential *S. aureus* genes when

exposed to different osmotic stressors underlines differences.

Conditionally essential genes of *S. aureus* strains grown either in 0.5 M NaCl (A), 0.5 M KCl (B) or 1.0

M sucrose (C) were mapped to cellular functions. Area sizes were adjusted to their essentiality

regardless of p- or q-value to visualize differences on a genome wide level. The larger the area, the

988   higher the number of transposon insertions in the LBN control condition as compared to the stress

989   conditions. Colors of polygons indicate cellular functions as detailed in (D).

Fig. 3. TN-Seq screen reveals essential and dispensable genes during prolonged NaCl stress in *S. aureus*.

(A) Workflow of TN-seq experiments. A saturated transposon library was pre-cultured for one hour,

then used to inoculate LB medium containing either normal levels (0.086 M) or 0.5 M of NaCl. After

17 generations, cells were harvested, and transposon insertion sites determined by high throughput

sequencing.

(B) Growth curve. Growth of the *S. aureus* library culture in LBN and LB 0.5 M medium was followed

by determining OD_{600} readings. Dotted line indicates 17 generation threshold. Cultures were back-

diluted once when they reached an OD of approximately 0.3 and the optical densities shown are

calculated from measured ODs time dilution factor. Culture from replicate B (n=1) is shown and a

similar growth profile was seen for replicate A.

(C) Volcano plots of essential and dispensable genes in NaCl conditions compared to LBN. Negative

log_2 fold-changes indicate essential and positive log_2 fold-changes dispensable genes. q-value stands

for Benjamini-Hochberg false discovery rate. Black horizontal line indicates q-value of 0.05 (cut-off)

and red, orange and green vertical lines 10-, 5- and 2-fold differences in either direction. Colored

circles indicate genes for which a significant change in the number of transposon insertions was

observed above q-value cutoff and follow the same color code as the vertical lines. Very small q-

values were truncated to fit onto the graph. Replicate A is shown in left panel and replicate B in right

panel (n=1 for each plot).

Fig. 4. Growth curves of *S. aureus* strains with transposon insertions in potential salt essential genes.

S. aureus strain JE2 (WT) and strains from the Nebraska Transposon Mutant Library (NTML) containing transposon insertions in potential salt essential genes were grown in LBN (left column) or 0.5 M NaCl LB medium (right column) and their growth monitored over 8 hours. Transposon mutant strains shown in panels (A) and (B) had a mild growth defect while strains shown in panels (C) and (D) had a stronger growth defect when grown in the high salt medium. NE numbers

1020   correspond to the NTML mutant number. Locus tag number and where available gene names with

1021   transposon insertion site are given in parenthesis. Growth curves were performed in triplicates

1022   (n=3) and means and SDs were plotted.

1023

Fig. 5. Growth curves and complementation analysis of *S. aureus* strains with transposon insertions in salt essential genes.

S. aureus strains (A) JE2 (WT), *SAUSA300_0694::Tn* mutant with empty plasmid and complementation strain or (B) JE2 (WT), *SAUSA300_0910* (*mgtE*), *SAUSA300_0957::Tn* with empty plasmid and respective complementation strains were grown in LBN (left panels) or LB 0.5 M NaCl medium (right panels) supplemented with 100 ng/ml Atet and OD_{600} readings determined at timed intervals. Growth curves were performed in triplicates (n=3) and means and SDs of OD_{600} readings were plotted.

1032

1033
1034
Fig. 6. Gene *SAUSA300_0957* is located in a genomic region with cell envelope genes and its

deletion leads to salt sensitivity and other phenotypic changes.

(A) Schematic of the *S. aureus* USA300 FPR3757 chromosomal region with gene *SAUSA300_0957*

(gene *957*). Gene *957* is encoded in an operon with *lcpB*, which codes for a wall teichoic acid ligase.

fmtA, a proposed D-amino esterase acting on teichoic acid and genes coding for a predicted

acyltransferase and the major autolysin Atl are located up and downstream of this operon. Putative

promoters are shown as angled arrows (adapted from (Mäder *et al.* 2016)) and black bars indicate

the corresponding transcripts.

(B) Growth curves. *S. aureus* strains LAC* piTET (WT), LAC*Δ*957* piTET (Δ*957*) and the

complementation strain LAC*Δ*957* piTET-*957* (compl.) were grown in LBN or LB containing 0.5 M

NaCl medium and 100 ng/ml Atet. The growth was monitored by determining OD_{600} readings the

means and SDs from three independent experiments were plotted.

(C) Determination of CFU/OD ratios. OD_{600} values as well as CFU/ml were determined for overnight

cultures of the *S. aureus* strains described in panel (B) and the means and SDs of the CFU per ml

OD_{600} of 1 from five independent experiments were plotted.

(D) Cell size measurements. The cell walls of the *S. aureus* strains described in panel (B) were stained

with fluorescently labelled vancomycin and the cells subsequently observed under a microscope.

The diameters of 200 cells were measured and the means calculated. This average of the means and

SDs from five independent experiments were plotted.

For statistical analysis, a Kruskal-Wallis one-way ANOVA test was performed followed by a Dunn's

post-hoc test to determine p-values. Asterisks (*) indicate $p \leq 0.05$ and two asterisks (**) $p \leq 0.01$ and

ns=not significant.

1057

Fig. 7. Inactivation of gene 957 leads to an increase in peptidoglycan crosslinking.

(A) Growth curves. *S. aureus* strains LAC* (WT), LAC*Δ*957* (Δ*957*), LAC*Δ*lcpB*, (Δ*957*), and

LAC*Δ*tagO* (Δ*tagO*) were grown in LB 0.5 M NaCl medium and OD$_{600}$ readings determined at timed

intervals. The means and SDs from three biological replicates were plotted.

(B) Detection of WTA on silver stained gels. WTA was isolated from *S. aureus* strains described in

panel (A) following growth in LB 0.4 M NaCl medium. The WTA was separated by electrophoresis

and visualized by silver stain. The experiment was performed three times and one representative

gel image is shown.

(C) Detection of WTA on silver stained gels. Same as (B) but using the strains LAC* piTET (WT),

LAC*Δ*957* piTET (Δ*957*), and the complementation strain LAC*Δ*957* piTET-*957* (compl.) grown in LB

0.4 M NaCl medium also containing 100 ng/ml Atet.

(D) Zymogram gel. Cell extracts prepared from *S. aureus* strains described in panel (C) were

separated on a gel containing heat killed *Micrococcus luteus* cells. Autolysins were renatured and

the zones of lysis visualized by methylene blue staining. The experiment was performed twice and

one experiment is shown.

(E) Determination of oxacillin MICs. Oxacillin MICs were determined for *S. aureus* strains described

in panel (C) using Etest strips. The median and SDs from four biological replicates were plotted.

(F) Muropeptide analysis. The same *S. aureus* strains as described in (C) were grown in LB 0.4 M

NaCl medium also containing 100 ng/ml Atet, peptidoglycan isolated and digested with mutanolysin

and the resulting muropeptide fragments separated by HPLC. Representative chromatograms are

shown in Supplementary Figures S6E. The peak area of individual peaks (corresponding to monomer

up to 8-mers and above) were quantified and the means and SDs from three biological replicates

plotted.

For statistical analysis of the oxacillin MICs in (E) a Kruskal-Wallis one-way ANOVA and a Dunn's

post-hoc test was performed and for the muropeptide analysis in (F) a two-way ANOVA and a

Dunnett's post-hoc test. One asterisk (*) indicates $p \leq 0.05$, two asterisks (**) $p \leq 0.01$ and four

asterisks (****) $p \leq 0.0001$. ns=not significant.

1086

1087

Fig. 8. The growth and peptidoglycan defect observed for the *957* mutant can be rescued by

compensatory mutations in *pbp2*.

(A) Growth curves. *S. aureus* strains LAC* (WT), LAC*Δ*957* (Δ*957*) and the LAC*Δ*957* suppressors

S1-10 (S1 to 10) were grown in LB 0.5 M NaCl medium, OD_{600} readings determined and the means

and SDs of four biological replicates plotted.

(B) Schematic of Pbp2 with amino acid substitutions identified. Top: Structure of the *S. aureus*

penicillin binding protein Pbp2 (PDB: 3DWK, (Lovering *et al.* 2008)) with amino acids that are altered

in the obtained suppressor strains shown in red. Bottom: Schematic of the Pbp2 enzyme with the

transglycosylase and transpeptidase domains as well as the observed amino acid changes in the

transglycosylase domain indicated.

(C) Growth curves using *957* mutant strains containing *pbp2* alleles from suppressors S2 and S4.

Growth curves were performed in LB 0.5 M NaCl medium and plotted as described in panel (A) but

using *S. aureus* strains LAC* *1332*::Tn (WT), LAC*Δ*957* *1332*::Tn (Δ*957*), LAC*Δ*957* *1332*::Tn *pbp2*

SNP S2 (Δ*957 pbp2* SNP S2), LAC*Δ*957* suppressor S4 *1332*::Tn (Δ*957 pbp2* SNP S4).

(D) Growth curves using *957* suppressors with their *pbp2* gene repaired to WT. Growth curves were

performed in LB 0.5 M NaCl medium and plotted as described in panel (A) but using *S. aureus* strains

LAC* *1332*::Tn (WT), LAC*Δ*957* *1332*::Tn (Δ*957*), LAC*Δ*957* suppressor S2 *1332*::Tn (S2), LAC*Δ*957*

suppressor S4 *1332*::Tn (S4), LAC*Δ*957* suppressor S2 *1332*::Tn repaired WT *pbp2* (S2 WT *pbp2*),

and LAC*Δ*957* suppressor S4 *1332*::Tn repaired WT *pbp2* (S4 WT *pbp2*).

(E) Growth curves using WT strains carrying *pbp2* SNPs. *S. aureus* strains LAC* *1332*::Tn (WT), LAC*

1332::Tn *pbp2* SNP S2 (WT *pbp2* SNP S2) and LAC* *1332*::Tn *pbp2* SNP S4 (WT *pbp2* SNP S4) were

grown in LB 0.5 M NaCl medium and OD_{600} readings determined and the means and SDs of three

biological replicates plotted.

(F) Bacterial growth on moenomycin supplemented agar plates. *S. aureus* strains LAC* (WT),

LAC*Δ*957* (Δ*957*), LAC*Δ*957* Suppressor S2 (S2) and S4 LAC*Δ*957* Suppressor S4 (S4) were grown

to exponential phase, normalized to an OD_{600} of 0.1 and plated either neat (n) or in 10-fold dilutions

onto LB agar containing 0.5 M NaCl without (top) or with (bottom) moenomycin. Shown is one

representative image of three biological replicates.

(G) Muropeptide analysis. *S. aureus* strains LAC*Δ*957* (Δ*957*), LAC*Δ*957* Suppressor S2 (S2) and S4

LAC*Δ*957* Suppressor S4 (S4) were grown in 0.4 M NaCl LB medium and muropeptides quantified

as described in Fig 5F. Representative chromatograms are shown in Supplementary Figure S8D. The

means and SDs from three biological replicates were plotted. For statistical analysis a two-way

ANOVA and Dunnett's post-hoc test was performed. One asterisk (*) indicates p≤0.05, two asterisks

(**) p≤0.01 and four asterisks (****) p≤0.0001.

1122

Fig. 9. Proposed model of 957 activity during NaCl stress.

S. aureus is normally able to grow in medium containing up to 2.5 M NaCl. The deletion of gene *957*

leads to a strong growth defect in high salt conditions and changes in the peptidoglycan structure.

We speculate that the observed increase in peptidoglycan crosslinking will lead to an increased

rigidity of the cell wall and therefore increased cell wall damage when these cells are exposed to

osmotic stress conditions.

Supplementary Figures

Supplementary Fig. 1. Voronoi diagram-based visualization of conditionally dispensable *S. aureus* genes when exposed to different osmotic stressors.

Conditionally dispensable genes of *S. aureus* strains grown either in 0.5 M NaCl (A), 0.5 M KCl (B) or 1.0 M sucrose (C) were mapped to cellular functions with area sizes adjusted to their essentiality

1136   regardless of p- or q-value. The larger the area, the higher the number of transposon insertions in

1137   relation to the LBN control. *recU* was scaled to 1/10 of the original ratio to preserve space. Colors

1138   of polygons indicate cellular functions as detailed in (D).

1139

1140

Supplementary Fig. 2. Circos plots showing transposon insertions in the *S. aureus* genome

following growth under different conditions.

The outer two bands represent genes on the forward (blue) and reverse (green) strand, respectively.

The inner three rings represent transposon insertions per gene following the growth of *S. aureus* in

in either LBN (orange) or LB 0.5 M NaCl (brown) for 17 generations. Each circular plot represents

one individual experiment (n=1).

1147

Conditionally essential genes at 0.5 M NaCl and 17 generations

1148

Supplementary Fig. 3. Voronoi diagram-based visualization of *S. aureus* conditionally essential

genes when exposed to 0.5 M NaCl.

Conditionally essential *S. aureus* genes from replicate A (A) and replicate B (B) grown in 0.5 M NaCl

for 17 generations were mapped to cellular functions with area sizes adjusted to their essentiality

regardless of p- or q-value. The larger the area, the lower the number of transposon insertions in

stress condition as compared to the LBN control condition. Colors of polygons indicate cellular

functions as detailed in (C).

Conditionally dispensable genes at 0.5 M NaCl and 17 generations

1157

Supplementary Fig. 4. Voronoi diagram-based visualization of *S. aureus* conditionally dispensable

genes when exposed to 0.5 M NaCl.

Conditionally dispensable *S. aureus* genes from replicate A (A) and replicate B (B) grown in 0.5 M

NaCl for 17 generations were mapped to cellular functions with area sizes adjusted to their

dispensability regardless of p- or q-value. The larger the area, the higher the number of transposon

insertions in the stress condition as compared to the LBN control. *recU* was scaled to 1/5 of the

original ratio to preserve space. Colors of polygons indicate cellular functions as detailed in (C).

1165

Supplementary Fig. 5. Complementation analysis using *S. aureus* mutants with transposon insertions in putative salt resistance genes.

1166

1167

(A) Growth curves. *S. aureus* strains JE2 (WT), NE867 and NE535 with integrated pCL55 and *S. aureus* strains NE867 and NE535 with respective complementation plasmids were grown in either LBN or LB 0.5 M NaCl and their growth monitored by determining OD_{600} readings.

1168

1169

1170

(B) As in (A) but using strains JE2 piTET (WT) or NE188 and NE251 containing plasmid piTET or the respective complementation plasmid. Strains were grown in LBN or 0.5 M NaCl LB medium supplemented with 100 ng/ml Atet. All experiments were conducted three times and the means and standard deviations were plotted.

1171

1172

1173

1174

1175

1176

Supplementary Fig. 6. Investigating the involvement of gene *957* in cell wall homeostasis.

(A) Growth curves using strains with mutations in WTA synthesis genes. *S. aureus* strains LAC* (WT),

LAC*Δ*957* (Δ*957*), LAC*Δ*lcpB* (Δ*lcpB*) and LAC*Δ*tagO* (Δ*tagO*) were grown in LBN and growth

monitored by measuring OD_{600} readings. The experiment was performed three times and means

and standard deviations were plotted.

(B) Detection of WTA. WTA was isolated from the strains described in (A) following growth in LBN

medium, then subjected to electrophoresis on polyacrylamide gels and visualized by silver staining.

One representative result is shown out of three independent experiments.

(C) Detection of WTA. Experimental setup is the same as in (B) but using the strains LAC* piTET (WT),

LAC*Δ*957* piTET (Δ*957*) and LAC* Δ*957* piTET-*957* (compl.).

(D) Muropeptide profile. The same strains as described in panel (C) were grown in LBN medium and

the peptidoglycan isolated, digestion with mutanolysin and the muropeptides subsequently

separated by HPLC. Shown are representative chromatograms from three independent

experiments. Retention time ranges of mono-, di- and higher multi-mers are indicated by vertical

lines.

(E) Muropeptide profiles. As in (D) but bacteria were grown in LB 0.4 M NaCl.

(F) Muropeptide quantification. The areas under the peaks from the chromatograms shown in panel

(D) and two additional replicates were quantified for each mono-, di- or multimer area. Bars

represent mean and error bars standard deviations from three independent experiments. For

statistical analyses, a two-way ANOVA was performed, followed by a Dunnett's post-hoc test but no

significant changes were present between either the wild type and the mutant or the wild type and

the deletion strain. Where induction was necessary, the medium was supplemented with 100 ng/µl

Atet.

Supplementary Fig. 7. Growth of Δ957 suppressor strains and schematic of the co-transduction strategy for complementation analysis.

(A) Growth curves of Δ957 suppressor strains. Strains LAC* (WT), LAC*Δ957 (Δ957) and LAC*Δ957 suppressors S1 trough S10 (S1-S10) were grown in LBN medium and their growth monitored by

taking OD$_{600}$ readings. The experiment was performed three times and means and standard

deviations were plotted.

(B) TN insertion map. Transposon insertions from replicate A (ANG dataset) were mapped to the

recU/pbp2 operon from a library grown in LBN (red) or LB medium containing 0.5 M NaCl salt (blue).

(C) TN insertion map. As in (B) but using replicate B (FCK dataset).

(D) Schematic of phage co-transduction experiment to either repair the *pbp2* SNP in the suppressor

strains (left branch) or to transduce the *pbp2* suppressor mutations into a clean *957* mutant (right

branch). A phage lysate was prepared using an NTML library strain with a transposon insertion in an

unrelated gene (*SAUSA300_1332*) located approximately 10 kb from the *pbp2* SNPs present in the

suppressor strains. This lysate was used for transductions using different LAC*Δ*957* suppressor

strains as recipient strains. With about 50% frequency, the *pbp2* SNPs were exchanged to the wild

type allele by transducing *SAUSA300_1332*::*Tn* and the WT *pbp2* allele producing strain LAC*Δ*957*

suppressor *1332*::*Tn pbp2* repaired (left branch). In the other cases, the *SAUSA300_1332*::*Tn* region

was transferred to a suppressor strain without replacing the *pbp2* SNP (right branch). A phage lysate

prepared on this intermediate strain (LAC*Δ*957* suppressor *1332*::*Tn*) was then used to transduce

the *pbp2* suppressor SNP into a clean LAC*Δ*957* strain, resulting in the clean suppressor strain

LAC*Δ*957 1332*::Tn *pbp2* SNP transduced.

1222

Supplementary Fig. 8. Growth and peptidoglycan analysis of *957* mutant and suppressor strains.

(A) Growth curves using *957* mutants with transduced *pbp2* SNPs. Strains LAC* *1332*::Tn (WT), LAC*Δ*957* *1332*::Tn (Δ*957*), LAC*Δ*957* *1332*::Tn *pbp2* SNP S2 (Δ*957 pbp2* SNP S2), LAC*Δ*957* suppressor S4 *1332*::Tn (Δ*957 pbp2* SNP S4) were grown in LBN medium and growth monitored by

determining OD$_{600}$ readings. The means and standard deviations from three independent

experiments were plotted.

(B) Growth curves using *957* suppressor strains carrying a repaired WT *pbp2* gene. Same as in (A)

but using strains LAC* *1332*::Tn (WT), LAC*Δ*957 1332*::Tn (Δ*957*), LAC*Δ*957* suppressor S2 *1332*::Tn

(S2), LAC*Δ*957* suppressor S4 *1332*::Tn (S4), LAC*Δ*957* suppressor S2 *1332*::Tn repaired WT *pbp2*

(S2 WT *pbp2*), and LAC*Δ*957* suppressor S4 *1332*::Tn repaired WT *pbp2* (S4 WT *pbp2*).

(C) Growth curves using WT strains carrying *pbp2* SNP mutations. Same as in (A) but using strains

LAC* *1332*::Tn (WT), LAC* *1332*::Tn *pbp2* SNP S2 (WT *pbp2* SNP S2) and LAC* *1332*::Tn *pbp2* SNP S4

(WT *pbp2* SNP S4).

(D) Muropeptide profiles. Strains LAC*Δ*957* (Δ*957*), LAC*Δ*957* Suppressor S2 (S2) and S4 LAC*Δ*957*

Suppressor S4 (S4) were grown in LB medium with 0.4 M NaCl, the peptidoglycan was extracted,

digested with mutanolysin, and the muropeptide separated by HPLC. Retention ranges for

monomers, di- and higher oligomers are indicated by vertical lines. The experiment was conducted

with three times and one representative result is shown.

1241