Common neurodegeneration-associated proteins are physiologically expressed by antigen-presenting cells and are interconnected via the inflammation/autophagy-related proteins TRAF6 and SQSTM1

Serge Nataf*[1, 2, 3], Marine Guillen[2] and Laurent Pays[1, 2, 3]

[1]CarMeN Laboratory, INSERM U1060, INRA U1397, INSA de Lyon, Lyon-Sud Faculty of Medicine, University of Lyon, Pierre-Bénite, France.
[2]University of Lyon 1, Lyon, France.
[3]Banque de Tissus et de Cellules des Hospices Civils de Lyon, Hôpital Edouard Herriot, Lyon, France.

* Corresponding author: serge.nataf@inserm.fr

ABSTRACT

There is circumstantial evidence that, under neurodegenerative conditions, peptides deriving from aggregated or misfolded specific proteins elicit adaptive immune responses. On another hand, several genes involved in familial forms of neurodegenerative diseases were found to exert key innate immune functions. However, whether or not such observations are causally linked remains unknown. To start addressing this issue, we followed a systems biology strategy based on the mining of large proteomics and immunopeptidomics databases. First, we retrieved the expression patterns of common neurodegeneration-associated proteins in two professional antigen-presenting cells, namely B-cells and dendritic cells. Surprisingly, we found that under physiological conditions, numerous neurodegeneration-associated proteins are abundantly expressed by human B-cells. A survey of the human proteome allowed us to map a unique protein-protein interaction network linking common neurodegeneration-associated proteins and their first shell interactors in human B-cells. Surprisingly, network connectivity analysis identified two major hubs that both relate with inflammation and autophagy, namely TRAF6 (TNF Receptor Associated Factor 6) and SQSTM1 (Sequestosome-1). Moreover, the mapped network in B-cells also comprised two other hub proteins involved in both inflammation and autoimmunity: HSPA8 (Heat Shock Protein Family A Member 8 also known as HSC70) and HSP90AA1 (Heat Shock Protein 90 Alpha Family Class A Member 1). Based on these results, we then explored the Immune Epitope Database "IEDB-AR" and actually found that a large share of neurodegeneration-associated proteins were previously reported to provide endogenous MHC class II-binding peptides in human B-cells. Of note, peptides deriving from amyloid beta A4 protein, sequestosome-1 or profilin-1 were reported to bind multiple allele-specific MHC class II molecules. In contrast, peptides deriving from microtubule-associated

protein tau, presenilin 2 and serine/threonine-protein kinase TBK1 were exclusively reported

to bind MHC molecules encoded by the HLA-DRB1 1501 allele, a recently-identified

susceptibility gene for late onset Alzheimer's disease. Finally, we observed that the whole list

of proteins reported to provide endogenous MHC class II-binding peptides in human B-cells

is specifically enriched in neurodegeneration-associated proteins. Overall, our work indicates

that immunization against neurodegeneration-associated proteins might be a physiological

process which is shaped, at least in part, by B-cells.

INTRODUCTION

Multiple studies have now established that neurodegenerative disorders are not cell-autonomous. The pathophysiological processes leading to neurodegeneration involve and target not only neurons but also glial cells, including astrocytes, microglia and oligodendrocytes (1–3). Moreover, beyond central nervous system (CNS) cells, the adaptive immune system has emerged as a potentially important player. When considering only T-cell responses, T-cell reactivity against amyloid beta peptides was already reported more than a decade ago in both patients suffering from Alzheimer's disease (AD) and aged healthy subjects (4). Recent works further demonstrated that alpha-synuclein (SNCA)-derived peptides elicit helper and cytotoxic T-cell responses in a subgroup of Parkinson's disease (PD) patients harboring specific major histocompatibility complex (MHC) alleles (5,6). Similarly, in animal models of neurodegenerative disorders, T-cell responses against peptides deriving from neurodegeneration-associated proteins were also demonstrated. In particular, in the MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) model of PD, pathogenic brain-infiltrating T-cells directed against SNCA were found to target the substantia nigra of diseased mice (7,8). Along this line, in a mouse model of tauopathy, CD8 cytotoxic T-cells directed against tau protein were shown to infiltrate the hippocampus and to drive cognitive alterations (9). Finally, besides AD and PD, several studies provided evidence that T-cells are activated during the course of Huntington's disease (HD), amyotrophic lateral sclerosis (ALS) and/or their animal models (10–13). Nevertheless, the putative autoantigens targeted under those conditions remain unknown.

Extending the functional results mentioned above, genome-wide association studies (GWAS) have unraveled the central position of autoimmunity-related genes in the genetic susceptibility landscape of neurodegenerative diseases (14–19). This is notably the case for

HLA-DRB1 alleles which were recently demonstrated to confer increased risks of developing

PD (15) or late-onset AD (14). Also, two genes involved in familial forms of PD, namely *PINK1*

(PTEN-induced putative kinase 1) and *PRKN* (Parkin RBR E3 Ubiquitin Protein Ligase also

named PARK2 or Parkin) were shown to regulate the presentation of mitochondria-derived

antigens in the context of MHC class I molecules (20). Finally, innate immune functions were

demonstrated for several genes causatively-linked to familial forms of PD or ALS. These

comprise *C9ORF72* (21–23), *LRRK2* (Leucine-rich repeat kinase 2) (24–27), *GRN* (Granulin

Precursor) (28,29) as well as *PRKN* and *PINK1* (30–34).

However, advocating for the role of autoimmunity in neurodegenerative disorders

requires yet addressing several important issues. In particular, it appears crucial to

determine how T-cells directed against neurodegeneration-associated antigens are primed

in the periphery. The extent to which autoimmunity against neurodegeneration-associated

antigens might be a physiological event needs also to be assessed. Last but not least, a global

view on the expression of neurodegeneration-associated proteins by professional antigen-

presenting cells (APCs) is still lacking. In an attempt to address these issues, we used here a

systems biology approach embracing a large range of previously-published experimental

data. We notably explored in normal human APCs the expression patterns and reported

protein-protein interactions of the most common neurodegeneration-associated proteins.

Overall, our results indicate that in human B-cells, a large majority of neurodegeneration-

associated proteins are expressed at the protein level and form a complex network centered

on the inflammation/autophagy-related molecules SQSTM1 (Sequestosome-1) and TRAF6

(TNF Receptor Associated Factor 6). Finally, the analysis of MHC class II immunopeptidome

databases provides evidence that neurodegeneration-associated proteins expressed by

human B-cells are a source of endogenous peptides which are presented in the context of

HLA class II molecules.

MATERIALS AND METHODS

Workflow

A scheme summarizing the workflow followed in the present work is shown in Figure 1.

Figure 1: Workflow of the sudy. The workflow starts from the upper central grey rectangle. Other rectangles (yellow or red) frame the main results obtained following each of the analytical steps which are briefly described in ellipse shapes (green or blue). Terms in italics correspond to the name of the bioinformatics tools used for each analytical step. MHC: major histocompatibility complex; APCs: antigen-presenting cells.

Data mining methods and bioinformatics tools

All the bioinformatics analyses were performed at least 3 times between March 2018 and

September 2019. Databases, bioinformatic tools and corresponding tasks performed in this

study are described below.

- The Genetics Home Reference website (35), is a regularly updated consumer health

resource from the National Library of Medicine. It provides information to the general public

about the effects of genetic variation on human health. In the present paper, we used the

"Genetics Home Reference" website to select without *a priori* the most common genetic

variations/alterations that are linked to the following neurodegenerative disorders:

Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease,

fronto-temporal dementia (FTD).

- The enrichment web platform Enrichr (36) perform enrichment analyses from queried lists

of genes. The Enrichr website allows surveying simultaneously 132 libraries gathering

245,575 terms and their associated lists of genes or proteins. Enrichment analysis tools

provided by the Enrichr bioinformatics platform provides adjusted P-values computed from

the Fisher's exact test. We focused our analysis on the "Jensen DISEASES" ontology library

(37) which is based exclusively on text-mining and allow determining whether a list of genes

is significantly associated with specific disease-related terms.

- The BioGrid database (38) compiles 29 169 experimentally-proven protein-protein

interactions connecting 23 098 human proteins. Querying the BioGrid database allows

retrieving for any given human protein the current list (update in 2019) of published

experimentally-identified human protein partners.

- The Human Proteome Map database (39) compiles protein expression data obtained by

mass spectrometry from human normal tissues and cells including a total of 85 human blood

samples from which the protein profiles of 6 blood-circulating cell types were established. In

parallel, we also explored 3 recently-published mass spectrometry datasets reporting on the

protemics profiles of blood-derived human B-cells (40), blood-derived human dendritic cells

(DCs) (41) and cultured monocyte-derived human dendritic cells (MoDCs)(42).

- The Immune Epitope Database and Analysis Resource (IEDB-AR) (43) compiles experimental

data on antibody and T-cell epitopes in humans, non-human primates and other animal

species. We followed a 3 step strategy as described below.

Step 1: retrieval of a list of human MHC class II binding peptides and their parent

proteins

- in the "Epitope" tab, the "Any Epitopes" item was marked

- in the "Assay" tab, the "MHC Ligand Assays" item was marked

- in the "Antigen" tab, "Homo sapiens" was entered in the "Organism" line

- in the "MHC restriction" tab, the "MHC class II" item was marked

- in the "Host" tab, the "Humans" item was marked

- in the "Disease" tab, the "Any Disease" item was marked.

Search was then launched and, from the results page, the "Assays" tab was selected

and a list of currently known ligands (peptides and parent proteins) of human MHC

molecules was retrieved.

Step 2: filtering the results

We retained only results which were both: i) obtained by mass spectrometry analysis

of peptides eluted from MHC class II molecules and ii) generated from cells of the B-

cell lineage, in the absence of immunization or stimulation protocols.

Step 3: checking the reference IDs associated to each identified parent protein

For each selected study and/or set of results, the reference ID associated to each

parent protein was checked on the UniProt website (44) in order to avoid

redundant or obsolete IDs.

- The Cytoscape software (45) is an open source software allowing to visualize complex

networks. We used the function "Network analysis" to identify hubs i.e objects exhibiting the

highest number of degrees (connections to other objects) in the generated networks.

Statistics

When not embedded in bioinformatics webtools, statistics for enrichment analyses were

performed using the Fisher exact test. In particular, we assessed for several identified hub

proteins whether their retrieved list of partners (obtained via the BioGrid database) was

statistically enriched in neurodegeneration-associated proteins. To calculate enrichment

factors, we set the expected reference ratio as 0.002 which corresponds to the number of

common neurodegeneration-associated proteins (i.e 48 according to the "Home Genetics

Reference" website) over the number of coding genes for which interactors can be retrieved

from the BioGrid database (i.e 23098). The obtained p-values were then adjusted using

Bonferroni correction. The same approach was used to determine whether lists of parent

proteins from which derive human MHC class II-binding peptides are significantly enriched in

neurodegeneration-associated proteins.

RESULTS

1- Neurodegeneration-associated proteins form a unique interaction network in which

PRKN is the main hub

From the most recently updated version of the human proteome compiled in the "BioGrid"

database, we extracted the currently published and experimentally-proven protein-protein

interactions connecting the most common neurodegeneration-associated proteins. Self-

interactions were excluded from our analysis. From these retrieved interactions we were

able to build and visualize a protein network which encompasses 35 (72%) of the 48 most

common neurodegeneration-associated proteins (Figure 2). Interestingly, network analysis

showed that PRKN harbored the highest number of interacting partners (n=17). Moreover,

when considering the whole list of known PRKN protein partners, the calculated enrichment

factor in neurodegeneration associated protein reached 18.97 and was highly significant

(Fisher exact test p-value: 3.35E^{-15}). This result points to a yet unnoticed property of PRKN as

a major hub protein connecting a large array of proteins involved in not only PD but also ALS,

AD, Huntington's disease and/or FTD.

Figure 2: Mapping of the protein-protein interaction network linking common
neurodegeneration-associated proteins irrespective of the cell type considered. A survey of
the human proteome was performed by querying the protein-protein interaction database BioGRID

(38). Each node represents a protein indicated by the corresponding gene symbol and each edge
represents an experimentally-demonstrated protein-protein interaction. In this network, the "PRKN"
(Parkin) node, highlighted in yellow, exhibits the highest degree (i.e the highest number of direct
interactors). White nodes correspond to the first-shell partners of PRKN. Grey nodes are not direct
interactors of PRKN.
199

2- A majority of common neurodegeneration-associated proteins are abundantly

expressed in human B-cells

We then explored proteomics databases to determine whether neurodegeneration-

associated proteins are expressed by APCs under physiological conditions. Our search was

restricted to 2 cell lineages which harbor demonstrated antigen-presenting functions: B-cells

(46–48) and dendritic cells (DCs) (49–51). Results were combined with those retrieved from

proteomics data obtained in 3 independent studies by the mass spectrometry analysis of

human B-cells (40) or human dendritic cells (41,42) . Since mass spectrometry is a not a

sensitive technique, one may consider that detected proteins are abundantly expressed. As

shown in table 1, it is striking that 34 (70%) of 48 common neurodegeneration-associated

proteins are detectable by mass spectrometry in human B-cells (Table 1). In human DCs, 20

(41%) of 48 common neurodegeneration-associated proteins were detected and all of these

were also detected in human B-cells. Conversely, 14 common degeneration-associated

proteins were reported to be detected in B-cells but not DCs. These include notably

apolipoprotein E, amyloid beta A4 protein, presenilin-1 and alpha-synuclein.

Table 1

Gene Symbol	Protein ID	Blood-derived B-cells (39)	Blood-derived B-cells (40)	Blood-derived DCs (41)	Cultured MoDCs (42)
APOE	P02649	detected	detected	0	0
APP	P05067	detected	0	0	0
ATXN2	Q99700	detected	detected	0	0
CHMP2B	Q9UQN3	detected	detected	0	detected
DCTN1	Q14203	detected	detected	0	detected

FIG4	Q92562	detected	detected	0	0
FUS	P35637	detected	detected	detected	detected
GBA	P04062	0	detected	0	detected
GRN	P28799	detected	detected	0	0
HNRNPA1	P09651	detected	0	0	detected
HTT	P42858	detected	detected	detected	detected
LRRK2	Q5S007	detected	detected	detected	0
MATR3	P43243	detected	detected	detected	detected
OPTN	Q96CV9	detected	detected	0	0
PARK7	Q99497	detected	detected	detected	detected
PFN1	P07737	detected	detected	detected	detected
PRPH	P41219	detected	0	0	detected
PSEN1	P49768	detected	0	0	0
SETX	Q7Z333	detected	detected	0	detected
SIGMAR1	Q99720	detected	detected	0	0
SNCA	P37840	detected	0	0	0
SOD1	P00441	detected	detected	detected	detected
SPG11	Q96JI7	detected	detected	0	0
SQSTM1	Q13501	detected	detected	0	detected
TARDBP	Q13148	detected	0	detected	detected
TBK1	Q9UHD2	detected	detected	0	detected
TBP	P20226	detected	detected	0	0
TRPM7	Q96QT4	0	detected	0	0
TUBA4A	P68366	detected	0	0	detected
UBQLN2	Q9UHD9	detected	detected	0	0
UCHL1	P09936	detected	detected	0	0
VAPB	O95292	detected	detected	detected	detected
VCP	P55072	detected	detected	detected	detected
VPS35	Q96QK1	detected	detected	detected	detected

Table 1: Expression pattern of common neurodegeneration-associated proteins in human B-cells and human dendritic cells. The proteomics profiles of human circulating B-cells, circulating dendritic cells (DCs) or cultured monocyte-derived dendritic cells (MoDCs) were retrieved from 4 independent mass spectrometry studies (39–42). We found that 34 from 48 common neurodegeneration-associated proteins were detected by mass spectrometry in at least 1 of the explored studies. Interestingly, 14 common degeneration-associated proteins (highlighted in yellow) were reported to be detected in B-cells but not DCs. These include notably apolipoprotein E, amyloid beta A4 protein, presenilin-1 and alpha-synuclein. In contrast, when detected in DCs, common neurodegeneration-associated proteins were constantly detected in B-cells.

Of note, in both human B-cells and DCs, *PRKN* was not detected at the protein level by mass

spectrometry. This finding urged us to identify the main hub protein(s) which may

interconnect abundantly-expressed neurodegeneration-associated proteins in human B-

cells.

3- Neurodegeneration-associated proteins expressed by human B-cells form a unique

interaction network in which the inflammation/autophagy-related protein SQSTM1 is the

main hub

From the interaction network depicted in Figure 2 we only retained nodes corresponding to

neurodegeneration-associated genes that are detectable by mass spectrometry in human B-

cells. We observed that in this cell type, 24 neurodegeneration-associated proteins are

predicted   to   form   a   tight   interaction   network   which   is   centered   on   the

inflammation/autophagy-related protein SQSTM1.

Figure 5: Mapping of the protein-protein interaction network linking common
neurodegeneration-associated proteins in human B-cells. A survey of the human proteome

was performed by querying the protein-protein interaction database BioGRID (38). Each node
represents a protein indicated by the corresponding gene symbol and each edge represents an
experimentally-demonstrated protein-protein interaction. In this network, the "SQSTM1"
(Sequestosome-1) node, highlighted in yellow, exhibits the highest degree (i.e the highest number of
direct interactors). White nodes correspond to the first-shell partners of SQSTM1 in human B-cells.
Grey nodes are expressed by B-cells but are not direct interactors of SQSTM1.
249

4- The protein network formed by neurodegeneration-associated proteins connects with a

set of first shell interactors among which the B-cell inflammation/autophagy-related

protein TRAF6 is the main hub

To extend these data, we then retrieved from the "BioGrid" database the currently known

and experimentally-demonstrated interactors of common neurodegeneration-associated

proteins, irrespective of their levels of expression in human B-cells. From this list, we

extracted candidate hub proteins interacting with more than 10 neurodegeneration-

associated proteins. For each of these candidate hubs, total lists of protein interactors were

then retrieved from BioGrid and a Fisher exact test was applied so to determine if

neurodegeneration-associated proteins were actually significantly enriched in each

established list. By this mean, we were able to assign an enrichment factor and an associated

p-value to each candidate hub. This approach allowed us to identify 20 hub proteins (Table

2) with which common neurodegeneration-associated proteins are specifically connected.

Among these 20 hub proteins, 10 are expressed at the protein level as assessed via the

"Human Proteome Map" database (39). Of note, from the 10 hub proteins exhibiting the

most significant and highest enrichment factors, 8 are abundantly expressed by human B-

cells. These include: i) the heat shock proteins HSPA8 (also named HSC70), HSPA4 (also

named HSP70) and HSP90AA1 (also named HSP90) which are all involved in both antigen

presentation by MHC class II molecules (52–54) and the modulation of T-cell responses (55–

57) and ii) YWHAZ and YWHAQ proteins also known as 14-3-3 protein zeta and theta

respectively, which bind MHC class II molecules (58,59) and are implicated in various

neurodegenerative diseases (60–62). Finally, unexpectedly, the most significant hub we

identified was TRAF6 (TNF receptor associated factor 6), an inflammation/autoimmunity-

related molecule (63,64) playing a major role in the control of B-cell activation (65,66).

Indeed, the whole list of TRAF6 interactors comprises 17 neurodegeneration-associated

proteins which corresponds to an enrichment factor of 27.68 and an adjusted p-value of

$1.57E^{-16}$.

Table 2

Gene symbol	Protein ID	Protein partners	ND-associated protein partners	Enrichment factor	Adjusted p-value
TRAF6	Q9Y4K3	307	17	27.68	1.37E-16
PRKN	O60260	448	17	18.97	6.71E-14
HSPA4	P34932	363	13	17.90	1.58E-10
SQSTM1	Q13501	294	12	20.40	2.03E-10
HSPA8	P11142	708	16	11.29	8.20E-10
YWHAZ	P63104	412	12	14.56	9.38E-9
HSP90AA1	P07900	839	15	8.93	6.65E-08
YWHAQ	P27348	476	11	11.55	4.74E-07
CYLD	Q9NQC7	620	12	9.67	8.15E-07
CTNNB1	P35222	616	11	8.92	6.00E-06
UBC	P0CG48	1050	14	6.66	6.54E-06
TP53	P04637	1062	14	6.59	7.46E-06
CUL7	Q14999	658	11	8.35	1.13E-05
EGFR	P00533	1224	14	5.71	3.73E-05
RNF4	P78317	1251	14	5.59	4.75E-05
MCM2	P49736	940	11	5.85	3.09E-04
BRCA1	P38398	962	11	5.71	3.80E-04
NTRK1	P04629	1948	16	4.10	4.54E-04
CUL3	Q13618	1227	13	5.29	1.75E-04
ESR2	Q92731	2249	13	2.89	4.67E-02

Legend: A survey of the human proteome was performed by querying the protein-protein interaction database BioGRID (38). Candidate hub proteins interacting with more than 10 common neurodegeneration-associated proteins were retrieved. The lists of partners currently demonstrated for each of these candidate hubs were then retrieved. For each of these lists, a factor of enrichment in neurodegeneration-associated proteins was established along with an associated Fisher's exact test p-value, as described in the Materials and Methods section. From left to right, gene symbols and corresponding protein IDs of the identified hub proteins are shown in the first and second column, the associated total numbers of partners and total numbers of neurodegeneration (ND)-associated

286 protein interactors are shown in the third and fourth column. The corresponding enrichment factors
287 and adjusted p-values are shown in the last two columns. Proteins that are detectable by mass
288 spectrometry in human B-cells, according to the "Human Proteome Map" database (39), are
289 highlighted in yellow.
290

However, one may argue that the list of currently-known TRAF6 partners might be enriched

in not only neurodegeneration-associated proteins but also many other sets of proteins

which do not relate with neurodegeneration. To address this issue, we performed on the

whole list of TRAF6 partners a non *a priori* enrichment analysis using the "JENSEN Disease"

text-mining webtool (37) embedded in the "Enrichr" analysis platform (36). Results shown in

Table 3 indicate that the 10 disease-related terms with which TRAF6 partners are the most

significantly associated comprise the terms "Frontotemporal dementia",

"Neurodegenerative disease" and "Pick's disease". This finding points to a specific link

between TRAF6 and neurodegeneration.

Table 3

Disease Enrichment	
Term	Adjusted p-value
Cancer	5.09E-7
Frontotemporal dementia	2.69E-6
Hypohidrotic ectodermal dysplasia	2.75E-6
Toxic encephalopathy	3.20E-6
Mulibrey nanism	3.21E-6
Neurodegenerative disease	3.45E-6
Arthritis	4.14E-6
Influenza	8.77E-6
Pick's disease	1.93E-4
Leishmaniasis	3.72E-4

Legend: A survey of the human proteome was performed by querying the protein-protein interaction
database BioGRID (38). The list of currently known and experimentally-demonstrated TRAF6 partners
was retrieved. From this list of interactors, an enrichment analysis with the "JENSEN Disease" text-
mining webtool (37) embedded in the "Enrichr" analysis platform (36) was then performed. Shown
are the disease-related terms exhibiting the 10 most statistically significant enrichments. Terms
relating with neurodegeneration are highlighted in yellow.
307

As a control and to further establish the specificity of our findings, we assessed whether

similar results would be obtained from a list of common demyelination-associated proteins

i.e. proteins commonly considered as candidate autoantigens in multiple sclerosis (67). We

found that neither TRAF6 nor the other hubs linking common neurodegeneration-associated

proteins exhibited lists of protein partners which were significantly enriched in common

demyelination-associated proteins (data supplement 1).

From these results we then built and visualized a B-cell protein-protein interaction

network encompassing the most significant hub proteins and their neurodegeneration-

associated interactors expressed in B-cells (Figure 5). We observed that from the 32

neurodegeneration-associated proteins expressed by B-cells, 22 (68%) are first shell

interactors of HSPA4, HSPA8, TRAF6 or SQSTM1. These results unravel a yet unknown

function of these molecules as major hub proteins connecting in B-cells a large array of

proteins involved in PD, ALS, AD, Huntington's disease and/or FTD.

Figure 6: Mapping of the protein-protein interaction network linking common neurodegeneration-associated proteins and their hub protein partners in human B-cells. A survey of the human proteome was performed by querying the protein-protein interaction database BioGRID (38). Only proteins expressed by human B-cells according to the "Human Proteome Map" database (39) were taken into account. Each node represents a protein indicated by its corresponding gene symbol. The 4 hub proteins expressed by B-cells and whose partners exhibited the most significant and highest enrichment factors in neurodegeneration-associated proteins are highlighted in yellow. The neurodegeneration-associated proteins expressed by B-cells and interacting with TRAF6 are shown in white ellipses. Other neurodegeneration-associated proteins involved in this network are shown in grey ellipses.

5- Common neurodegeneration-associated proteins provide a source of endogenous MHC class II-binding peptides in human B-cells

While peptides presented by MHC class II molecules are classically generated by the

proteolysis of phagocytized exogenous antigens, the presentation of endogenous peptides

by MHC class II molecules is an alternate pathway which has been robustly documented (68–

71). We thus surveyed the Immune Epitope DataBase (IEDB-AR) (43) to assess whether

peptides derived from neurodegeneration-associated proteins had been previously

identified as binding MHC class II molecules in immunopeptidome studies which used human

cells of the B-cell lineage as a source of endogenous peptides. Of note, we excluded studies

assessing the MHC binding of exogenously-provided specific peptides and retained only

works relying on the systematic mass spectrometry-based identification of peptides eluted

from MHC class II molecules. In addition, we excluded experiments in which immunization or

stimulation protocols were applied to B-cells. On this basis, we retained 19 studies (data

supplement 2) which were performed on cells of the B-cell lineage including predominantly

Epstein-Barr virus (EBV)-transformed B-cells. When screening these studies, we found that

23 out of 48 common neurodegeneration-associated proteins were reported to provide

endogenous MHC class II-binding peptides in B-cells (Table 4 and data supplement 2). The

351 most frequently identified neurodegeneration-associated parent proteins are abundantly

352 expressed in B-cells and comprise notably the proteins encoded by: i) the AD-related genes

353 *APP* and *PSEN1*, ii) the ALS/FTD-related genes *PFN1*, *SQSTM1*, *GRN*, *SOD1* and *VCP* and iii)

354 the PD-related proteins *PARK7* and *GBA* (Figure 1).

355 **Table 4**

Neurodegeneration-associated proteins	Reported MHC class II restrictions
Amyloid beta A4 protein (*APP*)	HLA class II, HLA-DQ, HLA-DR, HLA-DR3, HLA-DQ7.5, HLA-DQ2.5, HLA-DQ2.2, HLA-DRB1*15:01 (HLA-DR15), HLA-DRB5*01:01 (HLA-DR51), HLA-DRB1*04:05 (HLA-DR4), HLA-DQA1*05:01/DQB1*02:01 (HLA-DQ2.5)
Sequestosome-1 (*SQSTM1*)	HLA class II, HLA-DR, HLA-DQ, HLA-DR1, HLA-DQ7.5, HLA-DQ2.5, HLA-DQ2.2, HLA-DR15/DR51, HLA-DQA1*05:01/DQB1*02:01 (HLA-DQ2.5)
Profilin-1 (*PFN1*)	HLA class II, HLA-DR, HLA-DQ, HLA-DR1, HLA-DR3, HLA-DRB1*04:05 (HLA-DR4), HLA-DR15/DR51
Superoxide dismutase [Cu-Zn] (*SOD1*)	HLA class II, HLA-DR, HLA-DR1, HLA-DQ7.5, HLA-DQ2.5, HLA-DQ2.2
Transitional endoplasmic reticulum ATPase (*VCP*)	HLA class II, HLA-DR, HLA-DR1, HLA-DQ7.5, HLA-DQ2.5, HLA-DQ2.2
Lysosomal acid glucosylceramidase (*GBA)*	HLA class II, HLA-DR, HLA-DQ, HLA-DR1, HLA-DR15, HLA-DRB1*04:05 (HLA-DR4)
Presenilin-1 (*PSEN1*)	HLA class II, HLA-DQ, HLA-DQ2.2, HLA-DR15/DR51, HLA-DRB1*04:05 (HLA-DR4)
Ubiquilin-2 (*UBQLN2*)	HLA class II, HLA-DR1, HLA-DQ7.5,

	HLA-DQ2.2, HLA-DRB1*04:05 (HLA-DR4)
Major prion protein (*PRNP*)	HLA-DQ, HLA-DR, HLA-DR1, HLA-DQ7.5, HLA-DQ2.2
Vacuolar protein sorting-associated protein 35 (*VPS35*)	HLA class II, HLA-DR , HLA-DR15/DR51, HLA-DQA1*05:01/DQB1*02:01 (HLA-DQ2.5)
Optineurin (*OPTN*)	HLA-DR1, HLA-DQ7.5, HLA-DQ2.5, HLA-DQ2.2
Protein deglycase DJ-1 (*PARK7*)	HLA class II, HLA-DR1, HLA-DRB1*04:01 (HLA-DR4), HLA-DRB1*04:05 (HLA-DR4)
Progranulin (*GRN*)	HLA class II, HLA-DQ, HLA-DQ7.5
Spatacsin (*SPG11*)	HLA-DQ7.5, HLA-DQ2.2
Ubiquitin carboxyl-terminal hydrolase isozyme L1 (*UCHL1*)	HLA class II, HLA-DR
RNA-binding protein FUS (*FUS*)	HLA class II
Probable cation-transporting ATPase 13A2 (*ATP13A2*)	HLA-DR
Heterogeneous nuclear ribonucleoprotein A1 (*HNRNPA1*)	HLA class II
Charged multivesicular body protein 2b (*CHMP2B*)	HLA-DQ7.5
Huntingtin (*HTT*)	HLA-DR
Serine/threonine-protein kinase TBK1 (*TBK1*)	HLA-DR15/DR51
Presenilin-2 (*PSEN2*)	HLA-DR15/DR51
Microtubule-associated protein tau (*MAPT*)	HLA-DR15/DR51

Legend: The "IEDB-AR" immune epitope database (43) was screened in order to retain only publications reporting on the systematic mass spectrometry identification of endogenous peptides binding MHC-class II molecules in the human B-cell lineage. Only results obtained on human cells of the B-cell lineage in the absence of immunization or stimulation protocols were taken into account. The retrieved list of parent proteins and corresponding reported MHC class II restriction of derived peptides was then crossed with the list of 48 common neurodegeneration-associated proteins we established. The protein names and gene symbols are shown in the left column; the corresponding reported MHC class II restrictions of derived peptides are shown in the right column. When needed, aliases frequently used in the HLA class II genotype/serotype nomenclature are indicated in brackets.

6- Hub molecules linking common neurodegeneration-associated proteins provide a source of endogenous MHC class II-binding peptides in human B-cells

From the 19 relevant B-cell studies we retained on IEDB-AR, we also attempted to determine whether the hub molecules we identified as linking neurodegeneration-associated proteins in B-cells (Table 2) were, in parallel, reported to provide endogenous ligands for MHC class II molecules in B-cells. We found that from 10 candidate hubs abundantly expressed by B-cells,

372   8 were reported to provide endogenous ligands for MHC class II molecules in human B-cells

373   (Table 5 and data supplement 2).

375   **Table 5**

Hub molecules linking common neurodegeneration-associated proteins	Reported MHC class II restriction
Heat shock cognate 71 kDa protein (*HSPA8*)	HLA class II, HLA-DR, HLA-DQ, HLA-DR1, HLA-DQ7.5, HLA-DQ2.5, HLA-DQ2.2, HLA-DR3, HLA-DRB1*15:01 (HLA-DR15), HLA-DR15/DR51, HLA-DQA1*05:01/DQB1*02:01 (HLA-DQ2.5), HLA-DRB1*04:01 (HLA-DR4), HLA-DRB1*04:05 (HLA-DR4)
Sequestosome-1 (*SQSTM1*)	HLA class II, HLA-DR, HLA-DQ, HLA-DR1, HLA-DQ7.5, HLA-DQ2.5, HLA-DQ2.2, HLA-DR15/DR51, HLA-DQA1*05:01/DQB1*02:01 (HLA-DQ2.5)
Heat shock protein HSP 90-alpha (*HSP90AA1*)	HLA class II, HLA-DR, HLA-DQ, HLA-DQ7.5, HLA-DQ2.5, HLA-DQ2.2, HLA-DRB1*04:05 (HLA-DR4), HLA-DRB1*15:01 (HLA-DR15) HLA-DQA1*05:01/DQB1*02:01 (HLA-DQ2.5)
14-3-3 protein zeta/delta (*YWHAZ*)	HLA class II, HLA-DR, HLA-DQ, HLA-DR1, HLA-DRB1*04:05 (HLA-DR4)
14-3-3 protein theta (*YWHAQ*)	HLA class II, HLA-DQ, HLA-DRB1*04:05 (HLA-DR4)
Heat shock 70 kDa protein 4 (*HSPA4*)	HLA-DR, HLA-DR3
Ubiquitin carboxyl-terminal hydrolase CYLD (*CYLD*)	HLA-DRB1*04:05 (HLA-DR4)
TNF receptor-associated factor 6 (*TRAF6*)	HLA-DR1

Legend: The "IEDB-AR" immune epitope database (43) was screened in order to retain only
publications reporting on the systematic mass spectrometry identification of endogenous peptides
binding MHC-class II molecules in the human B-cell lineage. Only results obtained in the absence of
immunization or stimulation protocols were taken into account. The retrieved list of parent proteins
and corresponding reported MHC class II restriction of derived peptides was then crossed with the
list of 11 hub molecules being abundantly expressed by human B-cells and linking common
neurodegeneration-associated. The protein names and corresponding gene symbols are shown in the
left column; the corresponding reported MHC class II restrictions of derived peptides are shown in

the right column. When needed, aliases frequently used in the HLA class II genotype/serotype
nomenclature are indicated in brackets.
386

Among these 8 hubs, HSPA8 was the parent protein being the most frequently identified as

providing endogenous peptides which bind MHC class II molecules in human B-cells (Table

5).

7- In human B-cells, the list of parent proteins from which derive endogenous MHC class II-

binding peptides is specifically enriched in neurodegeneration-associated proteins

The whole list of parent proteins identified as providing MHC class II-binding endogenous

peptides in B-cells was retrieved from IEDB-AR and an enrichment analysis was performed to

determine whether such a list was significantly enriched in neurodegeneration-associated

proteins. We found that genes coding for neurodegeneration-associated proteins

encompassed 0.62% (22 out of 3523) of the whole genes coding for such parent proteins

(data supplement 3), which corresponds to an enrichment of 2.58 (p-value = 0.0006, Fisher

exact test) when considering the whole number of human protein-coding genes as roughly

20 000 (72). To confirm these results and assess their level of specificity, we used the

"JENSEN Disease" text-mining webtool (37) and performed an unsupervised enrichment

analysis from the whole list of genes coding for parent proteins previously identified as

providing MHC class II-binding peptides in human B-cells (data supplement 3). From this list

of 3522 genes, 96 (2.72%) were annotated with the term "Neurodegenerative disease" (data

supplement 3) which corresponds to the second most significant enrichment, after the term

"Arthritis" (Table 6).

Table 6

Disease Enrichment	
Term	Adjusted p-value
Arthritis	3.94E-8
Neurodegenerative disease	1.80E-7
Lupus erythematosus	4.92E-7
Tetanus	2.07E-5
Human immunodeficiency virus infectious disease	4.80E-5
Diamond-Blackfan anemia	7.10E-5
Coronary artery disease	1.57E-4
Fabry disease	2.08E-4
Inclusion-cell disease	2.16E-4
Neuropathy	1.97E-4

Legend: The "IEDB-AR" immune epitope database (43) was screened in order to retain only
publications reporting on the systematic mass spectrometry identification of endogenous peptides
binding MHC class II molecules in the human B-cell lineage. Only results obtained in the absence of
immunization or stimulation protocols were taken into account. The parent proteins from which
derive the identified bound peptides were retrieved and the list of corresponding coding genes was
submitted to an enrichment analysis with the "JENSEN Disease" text-mining webtool (37) embedded
in the "Enrichr" analysis platform (36). The 10 most significant terms associated with this list of genes
are shown. The second most significant enrichment is observed for the term "Neurodegenerative
disease" (term and corresponding adjusted p-value highlighted in yellow).

418
419

Since the HLA-DRB1 1501 allele (corresponding to the HLA-DR15 serotype) was recently

identified as a risk factor for sporadic forms of late onset AD (14), we retrieved the whole list

of parent proteins (and corresponding coding genes) which, in human B-cells, were

previously reported to provide peptides that bind HLA-DRB1 1501-encoded MHC class II

molecules. Importantly, such a list was significantly enriched in neurodegeneration-

associated proteins (enrichment factor: 5.34; p-value = 0.0003, Fisher exact test). Moreover,

when this list of parent proteins was submitted to an unsupervised enrichment analysis, the

term "Neurodegenerative disease" was found to reach the highest level of statistical

significance (Table 7).

Table 7

Disease Enrichment	
Term	Adjusted p-value
Neurodegenerative disease	3.31E-6
Allergic hypersensitivity disease	1.70E-4
Influenza	1.34E-4
Inclusion-cell disease	1.64E-3
Diamond-Blackfan anemia	2.12E-3
Gangliosidosis	1.77E-3
Lysosomal storage disease	2.55E-3
Lupus erythematosus	2.41E-3
Dementia	2.57E-3
Collagen disease	2.62E-3

Legend: The "IEDB-AR" immune epitope database (43) was screened in order to retain only publications reporting on the systematic mass spectrometry identification of endogenous peptides which bind HLA-DRB1 1501-encoded MHC class II molecules in the human B-cell lineage. Only results obtained in the absence of immunization or stimulation protocols were taken into account. The parent proteins from which derive the identified bound peptides were retrieved and the list of corresponding coding genes was submitted to an enrichment analysis with the "JENSEN Disease" text-mining webtool (37) embedded in the "Enrichr" analysis platform (36). Shown are the disease-related terms exhibiting the 10 most statistically significant enrichments. Terms relating with neurodegeneration are highlighted in yellow.

Finally, based on the IEDB-AR survey we performed, HLA-DRB1 1501-encoded MHC class II molecules are the only HLA class II molecules which, in human B-cells, were reported to bind endogenous peptides deriving from microtubule-associated protein tau, presenilin-2 or serine/threonine-protein kinase TBK1 (Table 4 and data supplement 2).

DISCUSSION

In the present work, we mined large publically-available databases to provide experiment-based evidence of a link between neurodegeneration and autoimmunity. Using a systems biology approach we report that a large range of common neurodegeneration-associated proteins: i) are expressed by human B-cells under physiological conditions, ii) form a comprehensive and functionally-relevant protein-protein interaction network in human B-cells and iii) provide endogenous peptides which bind MHC class-II molecules in human B-cells. Patients suffering from neurodegenerative conditions exhibit T cell- and/or antibody-mediated responses directed against major neurodegeneration-associated proteins such as amyloid beta A4 protein, alpha-synuclein and tau protein (4–6,73). However, naturally-occurring antibodies against amyloid-beta (74), alpha-synuclein (75–77) and tau protein (73,77,78) were also demonstrated in cohorts of healthy subjects. Similarly, apart from any pathological context, robust T-cell responses against peptides deriving from tau protein were recently demonstrated to widely occur in the general population (79).These findings suggest that the autoimmune processes described in patients with neurodegenerative conditions might be shaped by pre-existing physiological autoimmune responses directed against common neurodegeneration-associated proteins. It is worth noting that, while physiological autoimmunity was firmly demonstrated nearly 50 years ago (80–82), the intimate nature of the links bridging physiological autoimmunity to its pathological counterpart is still matter of debate. Numerous functions have been assigned to physiological autoimmunity (83,84), including, more recently, a support to cognition (85–90). In line with these findings, our data indicate that a specific set of brain antigens expressed by B-cells and involved in neurodegenerative diseases might prime a neuroprotective and, possibly, cognition-promoting T-cell response under physiological conditions. Of note, B-cells

are now recognized as professional APCs and, as such, are able to prime naïve CD4 T-cells

with a similar efficiency than DCs (91,92). The autophagy of cytosolic and nuclear proteins in

B-cells was shown to provide a continuous source of endogenous MHC class-II ligands (93).

Moreover, such autophagosome-derived peptides induce the proliferation of autologous T-

cells under *in vitro* conditions (94). However, that neurodegeneration-associated proteins

provide MHC class II-binding endogenous peptides in B-cells neither prove that T-cells are

actually primed against such peptides *in vivo* nor that peptide deriving from misfolded

neurodegeneration-associated proteins are presented by B-cells under neurodegenerative

conditions. Moreover, even if it was actually the case, the phenotype of autoreactive T-cells

generated via such a mechanism would need to be explored. More generally, one has to

keep in mind that no consensus has been currently reached regarding the phenotype of

physiological autoreactive T-cells. Thus, both autoreactive Tregs and autoreactive pro-

inflammatory T-cells belong to the physiological T-cell repertoire and were both found to

exert neuroprotective effects (95–98). Finally, the picture might be even more tangling since

Tregs were demonstrated to prevent autoimmunity by restraining autophagy in APCs (99).

Several genes involved in familial forms of neurodegenerative disorders exert key

functions in the autophagy pathway. These notably comprise PRKN, PINK1 and SQSTM1

(100–103). In the recent years, major works provided evidence that, in neurons and immune

cells, functional defects in such genes hamper mitophagy (a specialized form of autophagy),

stimulate the inflammasome pathway and foster the presentation of mitochondrial antigens

by MHC class I molecules (33,104,105). However, these findings do not render account for

the existence of HLA class II-restricted T-cell responses against neurodegeneration-

associated proteins. Moreover, antigens targeted by autoimmunity during

neurodegenerative conditions are far from deriving only from the mitochondrial

compartment. In this regard, our work suggests that in B-cells, the inflammation/autophagy-

related molecules SQSTM1 and TRAF6 are crucially involved in the presentation of

neurodegeneration-associated antigens by MHC class II molecules.

The mining of previously published mass spectrometry analyses performed on MHC

class II-eluted peptides showed that, in human B-cells, several neurodegeneration-

associated proteins provide endogenous peptides which bind a large range of MHC class II

alleles. This is notably the case for amyloid beta A4 protein, sequestosome-1 and profilin-1.

Similarly, peptides deriving from HSPA8, a hub molecule which links a high number of

neurodegeneration-associated proteins, bind multiple MHC class II alleles. Our observations

suggest that these 4 molecules are likely to elicit immune responses in a large range of the

human population. Determining whether or not such antigens are targeted by cognition-

promoting autoimmunity is a potentially important issue. On another hand, in human B-

cells, several neurodegeneration-associated antigens appear to provide endogenous

peptides harboring an allele-specific MHC class II restriction in B-cells. For e.g. endogenous

peptides deriving from microtubule-associated protein tau, PSEN2 and Serine/threonine-

protein kinase TBK1 were exclusively reported to bind HLA-DRB1 1501-encoded MHC class II

molecules under the experimental conditions described above. Since the HLA-DRB1 1501

allele is a susceptibility gene for late onset AD (14), this may prove to be of interest in the

context of AD pathophysiology. However, additional studies are clearly needed in order to

establish, under strictly comparable conditions, the endogenous immunopeptidomes of

distinct MHC class II alleles in B-cells. Moreover, it should be reminded that the endogenous

immunopeptidome of MHC class II molecules does not match the corresponding repertoire

of exogenous peptides (106). Since a widespread T-cell reactivity against Tau peptides was

recently demonstrated in the general human population (79), one may anticipate that

individuals bearing the HLA-DRB1 1501 allele exhibit functional specificities regarding their

anti-Tau T-cell responses.

We previously proposed that the genetic polymorphism of the HLA-DRB1 locus,

which, among primates, is extremely high in the human species (107), might allow the allele-

specific presentation of distinct sets of "brain superautoantigens" (108,109) leading, in turn,

to the development and maintenance of distinct sets of cognition-promoting T-cells. The

present work indicates that common neurodegeneration-associated proteins might

represent an important share of brain superautoantigens. Interestingly, recent magnetic

resonance imaging studies reported that, in healthy subjects, specific HLA-DR alleles

correlate with the volume ranges of specific brain structures (110,111). Future studies

should be designed to determine whether HLA-DR polymorphism might match both brain

structural features and the diversity of T-cell responses against common neurodegeneration-

associated proteins. Finally, in the human B-cell lineage, MHC class II-binding endogenous

peptides are highly significantly and specifically enriched in peptides deriving from common

neurodegeneration-associated proteins. This result raises the intriguing possibility that a

main function of physiological autoimmunity could be to control the blood-circulating levels

of aggregated forms of amyloid beta A4 protein, alpha-synuclein, tau protein and possibly

other neurodgeneration-associated proteins.

Acknowledgments: no specific funds were used for this study. SN performed the

bioinformatics analyses and wrote the paper. MG and LP performed quality controls of

bioinformatics analyses and wrote the paper. We thank Céline Auxenfans and Pascale Pascal

from the "Bank of Tissues and Cells" of the "Hospices Civils de Lyon" for leaving us the

opportunity to develop a bioinformatics research group. This paper can be freely accessed

via the BioRxiv platform at the following URL: http://biorxiv.org/cgi/content/short/690677v1

Conflicts of interest statement: The authors declare that there is no conflict of interest

regarding the publication of this article.

REFERENCES

1. Ilieva H, Polymenidou M, Cleveland DW. Non–cell autonomous toxicity in neurodegenerative disorders: ALS and beyond. *J Cell Biol* (2009) **187**:761–772. doi:10.1083/jcb.200908164

2. Scekic-Zahirovic J, Oussini H El, Mersmann S, Drenner K, Wagner M, Sun Y, Allmeroth K, Dieterlé S, Sinniger J, Dirrig-Grosch S, et al. Motor neuron intrinsic and extrinsic mechanisms contribute to the pathogenesis of FUS-associated amyotrophic lateral sclerosis. *Acta Neuropathol* (2017) **133**:887–906. doi:10.1007/s00401-017-1687-9

3. Boillee S, Yamanaka K, Lobsiger CS, Copeland NG, Jenkins NA, Kassiotis G, Kollias G, Cleveland DW. Onset and Progression in Inherited ALS Determined by Motor Neurons and Microglia. *Science (80-)* (2006) **312**:1389–1392. doi:10.1126/science.1123511

4. Monsonego A, Zota V, Karni A, Krieger JI, Bar-Or A, Bitan G, Budson AE, Sperling R, Selkoe DJ, Weiner HL. Increased T cell reactivity to amyloid β protein in older humans and patients with Alzheimer disease. *J Clin Invest* (2003) **112**:415–422. doi:10.1172/JCI18104

5. Sulzer D, Alcalay RN, Garretti F, Cote L, Kanter E, Agin-Liebes J, Liong C, McMurtrey C, Hildebrand WH, Mao X, et al. T cells from patients with Parkinson's disease recognize α-synuclein peptides. *Nature* (2017) **546**:656–661. doi:10.1038/nature22815

6. Garretti F, Agalliu D, Lindestam Arlehamn CS, Sette A, Sulzer D. Autoimmunity in Parkinson's Disease: The Role of α-Synuclein-Specific T Cells. *Front Immunol* (2019) **10**:303. doi:10.3389/fimmu.2019.00303

7. Benner EJ, Banerjee R, Reynolds AD, Sherman S, Pisarev VM, Tsiperson V, Nemachek C, Ciborowski P, Przedborski S, Mosley RL, et al. Nitrated α–Synuclein Immunity Accelerates Degeneration of Nigral Dopaminergic Neurons. *PLoS One* (2008) **3**:e1376.

doi:10.1371/journal.pone.0001376

8.   Brochard V, Combadière B, Prigent A, Laouar Y, Perrin A, Beray-Berthat V, Bonduelle

O, Alvarez-Fischer D, Callebert J, Launay J-M, et al. Infiltration of CD4+ lymphocytes

into the brain contributes to neurodegeneration in a mouse model of Parkinson

disease. *J Clin Invest* (2008) **119**:182–92. doi:10.1172/JCI36470

9.   Laurent C, Dorothée G, Hunot S, Martin E, Monnet Y, Duchamp M, Dong Y, Légeron F-

P, Leboucher A, Burnouf S, et al. Hippocampal T cell infiltration promotes

neuroinflammation and cognitive decline in a mouse model of tauopathy. *Brain* (2017)

140:184–200. doi:10.1093/brain/aww270

10.   Sheean RK, McKay FC, Cretney E, Bye CR, Perera ND, Tomas D, Weston RA, Scheller KJ,

Djouma E, Menon P, et al. Association of Regulatory T-Cell Expansion With

Progression of Amyotrophic Lateral Sclerosis. *JAMA Neurol* (2018) **75**:681.

doi:10.1001/jamaneurol.2018.0035

11.   Beers DR, Zhao W, Appel SH. The Role of Regulatory T Lymphocytes in Amyotrophic

Lateral Sclerosis. *JAMA Neurol* (2018) **75**:656. doi:10.1001/jamaneurol.2018.0043

12.   Coque E, Salsac C, Espinosa-Carrasco G, Varga B, Degauque N, Cadoux M, Crabé R,

Virenque A, Soulard C, Fierle JK, et al. Cytotoxic CD8 [+] T lymphocytes expressing ALS-

causing SOD1 mutant selectively trigger death of spinal motoneurons. *Proc Natl Acad*

Sci (2019) **116**:2312–2317. doi:10.1073/pnas.1815961116

13.   Niemelä V, Burman J, Blennow K, Zetterberg H, Larsson A, Sundblom J. Cerebrospinal

fluid sCD27 levels indicate active T cell-mediated inflammation in premanifest

Huntington's disease. *PLoS One* (2018) **13**:e0193492.

doi:10.1371/journal.pone.0193492

14.   Kunkle BW, Grenier-Boley B, Sims R, Bis JC, Damotte V, Naj AC, Boland A, Vronskaya

593   M, van der Lee SJ, Amlie-Wolf A, et al. Genetic meta-analysis of diagnosed Alzheimer's

594   disease identifies new risk loci and implicates Aβ, tau, immunity and lipid processing.

595   *Nat Genet* (2019) **51**:414–430. doi:10.1038/s41588-019-0358-2

596  15.  Hollenbach JA, Norman PJ, Creary LE, Damotte V, Montero-Martin G, Caillier S,

597   Anderson KM, Misra MK, Nemat-Gorgani N, Osoegawa K, et al. A specific amino acid

598   motif of *HLA-DRB1* mediates risk and interacts with smoking history in Parkinson's

599   disease. *Proc Natl Acad Sci* (2019) **116**:7419–7424. doi:10.1073/pnas.1821778116

600  16.  Witoelar A, Jansen IE, Wang Y, Desikan RS, Gibbs JR, Blauwendraat C, Thompson WK,

601   Hernandez DG, Djurovic S, Schork AJ, et al. Genome-wide Pleiotropy Between

602   Parkinson Disease and Autoimmune Diseases. *JAMA Neurol* (2017) **74**:780.

603   doi:10.1001/jamaneurol.2017.0469

604  17.  Broce I, Karch CM, Wen N, Fan CC, Wang Y, Hong Tan C, Kouri N, Ross OA, Höglinger

605   GU, Muller U, et al. Immune-related genetic enrichment in frontotemporal dementia:

606   An analysis of genome-wide association studies. *PLOS Med* (2018) **15**:e1002487.

607   doi:10.1371/journal.pmed.1002487

608  18.  Pottier C, Ren Y, Perkerson RB, Baker M, Jenkins GD, van Blitterswijk M, DeJesus-

609   Hernandez M, van Rooij JGJ, Murray ME, Christopher E, et al. Genome-wide analyses

610   as part of the international FTLD-TDP whole-genome sequencing consortium reveals

611   novel disease risk factors and increases support for immune dysfunction in FTLD. *Acta*

612   *Neuropathol* (2019) **137**:879–899. doi:10.1007/s00401-019-01962-9

613  19.  Raj T, Rothamel K, Mostafavi S, Ye C, Lee MN, Replogle JM, Feng T, Lee M, Asinovski N,

614   Frohlich I, et al. Polarization of the Effects of Autoimmune and Neurodegenerative

615   Risk Alleles in Leukocytes. *Science (80-)* (2014) **344**:519–523.

616   doi:10.1126/science.1249547

20.   Matheoud D, Sugiura A, Bellemare-Pelletier A, Laplante A, Rondeau C, Chemali M,

Fazel A, Bergeron JJ, Trudeau L-E, Burelle Y, et al. Parkinson's Disease-Related Proteins

PINK1 and Parkin Repress Mitochondrial Antigen Presentation. *Cell* (2016) **166**:314–

327. doi:10.1016/j.cell.2016.05.039

21.   Lall D, Baloh RH. Microglia and C9orf72 in neuroinflammation and ALS and

frontotemporal dementia. *J Clin Invest* (2017) **127**:3250–3258. doi:10.1172/JCI90607

22.   ORourke JG, Bogdanik L, Yanez A, Lall D, Wolf AJ, Muhammad AKMG, Ho R, Carmona

S, Vit JP, Zarrow J, et al. C9orf72 is required for proper macrophage and microglial

function in mice. *Science (80-)* (2016) **351**:1324–1329. doi:10.1126/science.aaf1064

23.   Nataf S, Pays L. Gene co-expression analysis unravels a link between C9orf72 and RNA

metabolism in myeloid cells. *Acta Neuropathol Commun* (2015) **3**:64.

doi:10.1186/s40478-015-0242-y

24.   Fava VM, Manry J, Cobat A, Orlova M, Van Thuc N, Ba NN, Thai VH, Abel L, Alcaïs A,

Schurr E, et al. A Missense LRRK2 Variant Is a Risk Factor for Excessive Inflammatory

Responses in Leprosy. *PLoS Negl Trop Dis* (2016) **10**:e0004412.

doi:10.1371/journal.pntd.0004412

25.   Wong AYW, Oikonomou V, Paolicelli G, De Luca A, Pariano M, Fric J, Tay HS, Ricciardi-

Castagnoli P, Zelante T. Leucine-Rich Repeat Kinase 2 Controls the Ca2+/Nuclear

Factor of Activated T Cells/IL-2 Pathway during Aspergillus Non-Canonical Autophagy

in Dendritic Cells. *Front Immunol* (2018) **9**:210. doi:10.3389/fimmu.2018.00210

26.   Härtlova A, Herbst S, Peltier J, Rodgers A, Bilkei-Gorzo O, Fearns A, Dill BD, Lee H,

Flynn R, Cowley SA, et al. LRRK2 is a negative regulator of *Mycobacterium tuberculosis*

phagosome maturation in macrophages. *EMBO J* (2018) **37**:e98694.

doi:10.15252/embj.201798694

27.   Kozina E, Sadasivan S, Jiao Y, Dou Y, Ma Z, Tan H, Kodali K, Shaw T, Peng J, Smeyne RJ.

Mutant LRRK2 mediates peripheral and central immune responses leading to

neurodegeneration in vivo. *Brain* (2018) **141**:1753–1769. doi:10.1093/brain/awy077

28.   Nielsen SR, Quaranta V, Linford A, Emeagi P, Rainer C, Santos A, Ireland L, Sakai T,

Sakai K, Kim Y-S, et al. Macrophage-secreted granulin supports pancreatic cancer

metastasis by inducing liver fibrosis. *Nat Cell Biol* (2016) **18**:549–560.

doi:10.1038/ncb3340

29.   Nguyen AD, Nguyen TA, Singh RK, Eberlé D, Zhang J, Abate JP, Robles A, Koliwad S,

Huang EJ, Maxfield FR, et al. Progranulin in the hematopoietic compartment protects

mice from atherosclerosis. *Atherosclerosis* (2018) **277**:145–154.

doi:10.1016/j.atherosclerosis.2018.08.042

30.   Carneiro FRG, Lepelley A, Seeley JJ, Hayden MS, Ghosh S. An Essential Role for ECSIT in

Mitochondrial Complex I Assembly and Mitophagy in Macrophages. *Cell Rep* (2018)

22:2654–2666. doi:10.1016/j.celrep.2018.02.051

31.   Zhou J, Yang R, Zhang Z, Liu Q, Zhang Y, Wang Q, Yuan H. Mitochondrial Protein PINK1

Positively Regulates RLR Signaling. *Front Immunol* (2019) **10**:1069.

doi:10.3389/fimmu.2019.01069

32.   Xu Y, Shen J, Ran Z. Emerging views of mitophagy in immunity and autoimmune

diseases. *Autophagy* (2019)1–15. doi:10.1080/15548627.2019.1603547

33.   Zhong Z, Umemura A, Sanchez-Lopez E, Liang S, Shalapour S, Wong J, He F, Boassa D,

Perkins G, Ali SR, et al. NF-κB Restricts Inflammasome Activation via Elimination of

Damaged Mitochondria. *Cell* (2016) **164**:896–910. doi:10.1016/j.cell.2015.12.057

34.   Sliter DA, Martinez J, Hao L, Chen X, Sun N, Fischer TD, Burman JL, Li Y, Zhang Z,

Narendra DP, et al. Parkin and PINK1 mitigate STING-induced inflammation. *Nature*

(2018) **561**:258–262. doi:10.1038/s41586-018-0448-9

35.   National Library of Medicine (US). Genetics Home Reference. Bethesda (MD): The

Library; Available from: https://ghr.nlm.nih.gov. Available at: https://ghr.nlm.nih.gov

36.   Kuleshov M V., Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, Koplev S,

Jenkins SL, Jagodnik KM, Lachmann A, et al. Enrichr: a comprehensive gene set

enrichment analysis web server 2016 update. *Nucleic Acids Res* (2016) **44**:W90–W97.

doi:10.1093/nar/gkw377

37.   Pletscher-Frankild S, Pallejà A, Tsafou K, Binder JX, Jensen LJ. DISEASES: Text mining

and data integration of disease–gene associations. *Methods* (2015) **74**:83–89.

doi:10.1016/j.ymeth.2014.11.020

38.   Oughtred R, Stark C, Breitkreutz B-J, Rust J, Boucher L, Chang C, Kolas N, O'Donnell L,

Leung G, McAdam R, et al. The BioGRID interaction database: 2019 update. *Nucleic*

Acids Res (2019) **47**:D529–D541. doi:10.1093/nar/gky1079

39.   Kim M-S, Pinto SM, Getnet D, Nirujogi RS, Manda SS, Chaerkady R, Madugundu AK,

Kelkar DS, Isserlin R, Jain S, et al. A draft map of the human proteome. *Nature* (2014)

509:575–581. doi:10.1038/nature13302

40.   Johnston HE, Carter MJ, Larrayoz M, Clarke J, Garbis SD, Oscier D, Strefford JC, Steele

AJ, Walewska R, Cragg MS. Proteomics Profiling of CLL Versus Healthy B-cells Identifies

Putative Therapeutic Targets and a Subtype-independent Signature of Spliceosome

Dysregulation. *Mol Cell Proteomics* (2018) **17**:776–791.

doi:10.1074/mcp.RA117.000539

41.   Worah K, Mathan TSM, Vu Manh TP, Keerthikumar S, Schreibelt G, Tel J, Duiveman-

de Boer T, Sköld AE, van Spriel AB, de Vries IJM, et al. Proteomics of Human Dendritic

Cell Subsets Reveals Subset-Specific Surface Markers and Differential Inflammasome

Function. *Cell Rep* (2016) **16**:2953–2966. doi:10.1016/j.celrep.2016.08.023

42.   Strasser L, Dang H-H, Schwarz H, Asam C, Ferreira F, Horejs-Hoeck J, Huber CG.

Unbiased Quantitative Proteomics Reveals a Crucial Role of the Allergen Context for

the Activation of Human Dendritic Cells. *Sci Rep* (2017) **7**:16638. doi:10.1038/s41598-

017-16726-2

43.   Dhanda SK, Mahajan S, Paul S, Yan Z, Kim H, Jespersen MC, Jurtz V, Andreatta M,

Greenbaum JA, Marcatili P, et al. IEDB-AR: immune epitope database—analysis

resource in 2019. *Nucleic Acids Res* (2019) **47**:W502–W506. doi:10.1093/nar/gkz452

44.   UniProt Consortium. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids*

Res (2019) **47**:D506–D515. doi:10.1093/nar/gky1049

45.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B,

Ideker T. Cytoscape: A Software Environment for Integrated Models of Biomolecular

Interaction Networks. *Genome Res* (2003) **13**:2498–2504. doi:10.1101/gr.1239303

46.   Bretou M, Kumari A, Malbec O, Moreau HD, Obino D, Pierobon P, Randrian V, Sáez PJ,

Lennon-Duménil A-M. Dynamics of the membrane-cytoskeleton interface in MHC

class II-restricted antigen presentation. *Immunol Rev* (2016) **272**:39–51.

doi:10.1111/imr.12429

47.   Yuseff M-I, Pierobon P, Reversat A, Lennon-Duménil A-M. How B cells capture, process

and present antigens: a crucial role for cell polarity. *Nat Rev Immunol* (2013) **13**:475–

86. doi:10.1038/nri3469

48.   Abbas AK. Antigen presentation by B lymphocytes: mechanisms and functional

significance. *Semin Immunol* (1989) **1**:5–12. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/15630954

49.   Villadangos JA, Schnorrer P. Intrinsic and cooperative antigen-presenting functions of

dendritic-cell subsets in vivo. *Nat Rev Immunol* (2007) **7**:543–55. doi:10.1038/nri2103

50.   Guermonprez P, Valladeau J, Zitvogel L, Théry C, Amigorena S. Antigen presentation

and T cell stimulation by dendritic cells. *Annu Rev Immunol* (2002) **20**:621–67.

doi:10.1146/annurev.immunol.20.100301.064828

51.   Steinman RM, Hawiger D, Nussenzweig MC. Tolerogenic dendritic cells. *Annu Rev*

Immunol (2003) **21**:685–711. doi:10.1146/annurev.immunol.21.120601.141040

52.   Zhou D, Li P, Lin Y, Lott JM, Hislop AD, Canaday DH, Brutkiewicz RR, Blum JS. Lamp-2a

Facilitates MHC Class II Presentation of Cytoplasmic Antigens. *Immunity* (2005)

22:571–581. doi:10.1016/j.immuni.2005.03.009

53.   Mycko MP, Cwiklinska H, Szymanski J, Szymanska B, Kudla G, Kilianek L, Odyniec A,

Brosnan CF, Selmaj KW. Inducible Heat Shock Protein 70 Promotes Myelin

Autoantigen Presentation by the HLA Class II. *J Immunol* (2004) **172**:202–213.

doi:10.4049/jimmunol.172.1.202

54.   Houlihan JL, Metzler JJ, Blum JS. HSP90alpha and HSP90beta isoforms selectively

modulate MHC class II antigen presentation in B cells. *J Immunol* (2009) **182**:7451–8.

doi:10.4049/jimmunol.0804296

55.   van Eden W, Jansen MAA, Ludwig I, van Kooten P, van der Zee R, Broere F. The Enigma

of Heat Shock Proteins in Immune Tolerance. *Front Immunol* (2017) **8**:1599.

doi:10.3389/fimmu.2017.01599

56.   Cohen IR. Autoantibody repertoires, natural biomarkers, and system controllers.

Trends Immunol (2013) **34**:620–5. doi:10.1016/j.it.2013.05.003

57.   Quintana FJ, Cohen IR. Heat shock proteins as endogenous adjuvants in sterile and

septic inflammation. *J Immunol* (2005) **175**:2777–82.

doi:10.4049/jimmunol.175.5.2777

58. Kuwana T, Peterson PA, Karlsson L. Exit of major histocompatibility complex class II-

invariant chain p35 complexes from the endoplasmic reticulum is modulated by

phosphorylation. *Proc Natl Acad Sci U S A* (1998) **95**:1056–61.

doi:10.1073/pnas.95.3.1056

59. Buschow SI, van Balkom BWM, Aalberts M, Heck AJR, Wauben M, Stoorvogel W. MHC

class II-associated proteins in B-cell exosomes and potential functional implications for

exosome biogenesis. *Immunol Cell Biol* (2010) **88**:851–856. doi:10.1038/icb.2010.64

60. Umahara T, Uchihara T, Tsuchiya K, Nakamura A, Iwamoto T, Ikeda K, Takasaki M. 14-

3-3 proteins and zeta isoform containing neurofibrillary tangles in patients with

Alzheimer's disease. *Acta Neuropathol* (2004) **108**:279–86. doi:10.1007/s00401-004-

0885-4

61. Kawamoto Y, Akiguchi I, Nakamura S, Honjyo Y, Shibasaki H, Budka H. 14-3-3 proteins

in Lewy bodies in Parkinson disease and diffuse Lewy body disease brains. *J*

Neuropathol Exp Neurol (2002) **61**:245–53. doi:10.1093/jnen/61.3.245

62. Jia B, Wu Y, Zhou Y. 14-3-3 and aggresome formation: implications in

neurodegenerative diseases. *Prion* (2014) **8**:173–177. doi:10.4161/pri.28123

63. Magilnick N, Reyes EY, Wang W-L, Vonderfecht SL, Gohda J, Inoue J, Boldin MP. *miR-*

146a – Traf6 regulatory axis controls autoimmunity and myelopoiesis, but is

dispensable for hematopoietic stem cell homeostasis and tumor suppression. *Proc*

Natl Acad Sci (2017) **114**:E7140–E7149. doi:10.1073/pnas.1706833114

64. Brenke JK, Popowicz GM, Schorpp K, Rothenaigner I, Roesner M, Meininger I, Kalinski

C, Ringelstetter L, R'kyek O, Jürjens G, et al. Targeting TRAF6 E3 ligase activity with a

small-molecule inhibitor combats autoimmunity. *J Biol Chem* (2018) **293**:13191–

13203. doi:10.1074/jbc.RA118.002649

65.   Ahonen CL, Manning EM, Erickson LD, O'Connor BP, Lind EF, Pullen SS, Kehry MR,

Noelle RJ. The CD40-TRAF6 axis controls affinity maturation and the generation of

long-lived plasma cells. *Nat Immunol* (2002) **3**:451–456. doi:10.1038/ni792

66.   Jalukar S V., Hostager BS, Bishop GA. Characterization of the Roles of TNF Receptor-

Associated Factor 6 in CD40-Mediated B Lymphocyte Effector Functions. *J Immunol*

(2000) **164**:623–630. doi:10.4049/jimmunol.164.2.623

67.   Ben-Nun A, Kaushansky N, Kawakami N, Krishnamoorthy G, Berer K, Liblau R, Hohlfeld

R, Wekerle H. From classic to spontaneous and humanized models of multiple

sclerosis: Impact on understanding pathogenesis and drug development. *J Autoimmun*

(2014) **54**:33–50. doi:10.1016/j.jaut.2014.06.004

68.   Adorini L, Moreno J, Momburg F, Hämmerling GJ, Guéry JC, Valli A, Fuchs S.

Exogenous peptides compete for the presentation of endogenous antigens to major

histocompatibility complex class II-restricted T cells. *J Exp Med* (1991) **174**:945–8.

doi:10.1084/jem.174.4.945

69.   Leung CSK. Endogenous Antigen Presentation of MHC Class II Epitopes through Non-

Autophagic Pathways. *Front Immunol* (2015) **6**:464. doi:10.3389/fimmu.2015.00464

70.   Coulon P-G, Richetta C, Rouers A, Blanchet FP, Urrutia A, Guerbois M, Piguet V,

Theodorou I, Bet A, Schwartz O, et al. HIV-Infected Dendritic Cells Present Endogenous

MHC Class II-Restricted Antigens to HIV-Specific CD4+ T Cells. *J Immunol* (2016)

197:517–32. doi:10.4049/jimmunol.1600286

71.   Rudensky AY, Maric M, Eastman S, Shoemaker L, DeRoos PC, Blum JS. Intracellular

assembly and transport of endogenous peptide-MHC class II complexes. *Immunity*

(1994) **1**:585–94. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7600287

72.   Pertea M, Shumate A, Pertea G, Varabyou A, Breitwieser FP, Chang Y-C, Madugundu

AK, Pandey A, Salzberg SL. CHESS: a new human gene catalog curated from thousands

of large-scale RNA sequencing experiments reveals extensive transcriptional noise.

Genome Biol (2018) **19**:208. doi:10.1186/s13059-018-1590-2

73. Bartos A, Fialová L, Švarcová J. Lower Serum Antibodies Against Tau Protein and Heavy

Neurofilament in Alzheimer's Disease. *J Alzheimers Dis* (2018) **64**:751–760.

doi:10.3233/JAD-180039

74. Taguchi H, Planque S, Nishiyama Y, Szabo P, Weksler ME, Friedland RP, Paul S.

Catalytic antibodies to amyloid beta peptide in defense against Alzheimer disease.

Autoimmun Rev (2008) **7**:391–7. doi:10.1016/j.autrev.2008.03.004

75. Huang Y, Xie X, Ji M, Yu X, Zhu J, Zhang L, Liu X, Wei C, Li G, Liu R. Naturally occurring

autoantibodies against α-synuclein rescues memory and motor deficits and

attenuates α-synuclein pathology in mouse model of Parkinson's disease. *Neurobiol*

Dis (2019) **124**:202–217. doi:10.1016/j.nbd.2018.11.024

76. Albus A, Gold M, Bach J-P, Burg-Roderfeld M, Jördens M, Kirchhein Y, Kronimus Y,

Mengel D, Zerr I, Dodel R. Extending the functional characteristics of naturally

occurring autoantibodies against β-Amyloid, Prion Protein and α-Synuclein. *PLoS One*

(2018) **13**:e0202954. doi:10.1371/journal.pone.0202954

77. Kuhn I, Rogosch T, Schindler TI, Tackenberg B, Zemlin M, Maier RF, Dodel R, Kronimus

Y. Serum titers of autoantibodies against α-synuclein and tau in child- and adulthood.

J Neuroimmunol (2018) **315**:33–39. doi:10.1016/j.jneuroim.2017.12.003

78. Krestova M, Hromadkova L, Ricny J. Purification of Natural Antibodies Against Tau

Protein by Affinity Chromatography. *Methods Mol Biol* (2017) **1643**:33–44.

doi:10.1007/978-1-4939-7180-0_4

79. Lindestam Arlehamn CS, Pham J, Alcalay RN, Frazier A, Shorr E, Carpenter C, Sidney J,

Dhanwani R, Agin-Liebes J, Garretti F, et al. Widespread Tau-Specific CD4 T Cell

Reactivity in the General Population. *J Immunol* (2019) **203**:84–92.

doi:10.4049/jimmunol.1801506

80.   Cohen IR, Wekerle H. REGULATION OF AUTOSENSITIZATION: THE IMMUNE

ACTIVATION AND SPECIFIC INHIBITION OF SELF-RECOGNIZING THYMUS-DERIVED

LYMPHOCYTES. *J Exp Med* (1973) **137**:224–238. doi:10.1084/jem.137.2.224

81.   Cohen IR, Wekerle H. Autosensitization of lymphocytes against thymus reticulum cells.

Science (1972) **176**:1324–5. doi:10.1126/science.176.4041.1324

82.   Cohen IR, Globerson A, Feldman M. AUTOSENSITIZATION IN VITRO. *J Exp Med* (1971)

133:834–845. doi:10.1084/jem.133.4.834

83.   Coutinho A, Kazatchkine MD, Avrameas S. Natural autoantibodies. *Curr Opin Immunol*

(1995) **7**:812–8. Available at: http://www.ncbi.nlm.nih.gov/pubmed/8679125

84.   Schwartz M, Cohen IR. Autoimmunity can benefit self-maintenance. *Immunol Today*

(2000) **21**:265–8. Available at: http://www.ncbi.nlm.nih.gov/pubmed/10825737

85.   Schwartz M, Baruch K. Breaking peripheral immune tolerance to CNS antigens in

neurodegenerative diseases: Boosting autoimmunity to fight-off chronic

neuroinflammation. *J Autoimmun* (2014) **54**:8–14. doi:10.1016/j.jaut.2014.08.002

86.   Kipnis J, Cardon M, Strous RD, Schwartz M. Loss of autoimmune T cells correlates with

brain diseases: possible implications for schizophrenia? *Trends Mol Med* (2006)

12:107–112. doi:10.1016/j.molmed.2006.01.003

87.   Radjavi A, Smirnov I, Derecki N, Kipnis J. Dynamics of the meningeal CD4+ T-cell

repertoire are defined by the cervical lymph nodes and facilitate cognitive task

performance in mice. *Mol Psychiatry* (2014) **19**:531–532. doi:10.1038/mp.2013.79

88.   Derecki NC, Cardani AN, Yang CH, Quinnies KM, Crihfield A, Lynch KR, Kipnis J.

Regulation of learning and memory by meningeal immunity: a key role for IL-4. *J Exp*

Med (2010) **207**:1067–1080. doi:10.1084/jem.20091419

89.   Brynskikh A, Warren T, Zhu J, Kipnis J. Adaptive immunity affects learning behavior in

mice. *Brain Behav Immun* (2008) **22**:861–869. doi:10.1016/j.bbi.2007.12.008

90.   Kipnis J, Cohen H, Cardon M, Ziv Y, Schwartz M. T cell deficiency leads to cognitive

dysfunction: Implications for therapeutic vaccination for schizophrenia and other

psychiatric conditions. *Proc Natl Acad Sci* (2004) **101**:8180–8185.

doi:10.1073/pnas.0402268101

91.   Roche PA, Furuta K. The ins and outs of MHC class II-mediated antigen processing and

presentation. *Nat Rev Immunol* (2015) **15**:203–16. doi:10.1038/nri3818

92.   Schuijs MJ, Hammad H, Lambrecht BN. Professional and "Amateur" Antigen-

Presenting Cells In Type 2 Immunity. *Trends Immunol* (2019) **40**:22–34.

doi:10.1016/j.it.2018.11.001

93.   Schmid D, Münz C. Innate and Adaptive Immunity through Autophagy. *Immunity*

(2007) **27**:11–21. doi:10.1016/j.immuni.2007.07.004

94.   Mohme M, Hotz C, Stevanović S, Binder T, Lee J-H, Okoniewski M, Eiermann T,

Sospedra M, Rammensee H-G, Martin R. HLA-DR15-derived self-peptides are involved

in increased autologous T cell proliferation in multiple sclerosis. *Brain* (2013)

136:1783–1798. doi:10.1093/brain/awt108

95.   Kipnis J, Avidan H, Caspi RR, Schwartz M. Dual effect of CD4+CD25+ regulatory T cells

in neurodegeneration: A dialogue with microglia. *Proc Natl Acad Sci* (2004)

101:14663–14669. doi:10.1073/pnas.0404842101

96.   Baruch K, Rosenzweig N, Kertser A, Deczkowska A, Sharif AM, Spinrad A, Tsitsou-

Kampeli A, Sarel A, Cahalon L, Schwartz M. Breaking immune tolerance by targeting

Foxp3+ regulatory T cells mitigates Alzheimer's disease pathology. *Nat Commun*

(2015) **6**:7967. doi:10.1038/ncomms8967

97.   Baek H, Ye M, Kang G-H, Lee C, Lee G, Bin D, Jung J, Kim H, Lee S, Kim JS, et al.

Neuroprotective effects of

CD4⁺CD25⁺Foxp3<sup>+</su

p> regulatory T cells in a 3xTg-AD Alzheimer’s disease model.

Oncotarget (2016) **7**: doi:10.18632/oncotarget.12469

98.   Dansokho C, Ait Ahmed D, Aid S, Toly-Ndour C, Chaigneau T, Calle V, Cagnard N,

Holzenberger M, Piaggio E, Aucouturier P, et al. Regulatory T cells delay disease

progression in Alzheimer-like pathology. *Brain* (2016) **139**:1237–1251.

doi:10.1093/brain/awv408

99.   Alissafi T, Banos A, Boon L, Sparwasser T, Ghigo A, Wing K, Vassilopoulos D, Boumpas

D, Chavakis T, Cadwell K, et al. Tregs restrain dendritic cell autophagy to ameliorate

autoimmunity. *J Clin Invest* (2017) **127**:2789–2804. doi:10.1172/JCI92079

100. Majcher V, Goode A, James V, Layfield R. Autophagy receptor defects and ALS-FTLD.

Mol Cell Neurosci (2015) **66**:43–52. doi:10.1016/j.mcn.2015.01.002

101. Ciechanover A, Kwon YT. Protein Quality Control by Molecular Chaperones in

Neurodegeneration. *Front Neurosci* (2017) **11**:185. doi:10.3389/fnins.2017.00185

102. Springer W, Kahle PJ. Regulation of PINK1-Parkin-mediated mitophagy. *Autophagy*

(2011) **7**:266–278. doi:10.4161/auto.7.3.14348

103. Vives-Bauza C, Przedborski S. Mitophagy: the latest problem for Parkinson's disease.

Trends Mol Med (2011) **17**:158–165. doi:10.1016/j.molmed.2010.11.002

104. Baden P, Deleidi M. Mitochondrial Antigen Presentation: A Vacuolar Path to

Autoimmunity in Parkinson's Disease. *Trends Immunol* (2016) **37**:719–721.

881   doi:10.1016/j.it.2016.08.016

882 105. Roberts RF, Fon EA. Presenting mitochondrial antigens: PINK1, Parkin and MDVs steal

883   the show. *Cell Res* (2016) **26**:1180–1181. doi:10.1038/cr.2016.104

884 106. Ooi JD, Petersen J, Tan YH, Huynh M, Willett ZJ, Ramarathinam SH, Eggenhuizen PJ,

885   Loh KL, Watson KA, Gan PY, et al. Dominant protection from HLA-linked autoimmunity

886   by antigen-specific regulatory T cells. *Nature* (2017) **545**:243–247.

887   doi:10.1038/nature22329

888 107. Bergström TF, Josefsson A, Erlich HA, Gyllensten U. Recent origin of HLA-DRB1 alleles

889   and implications for human evolution. *Nat Genet* (1998) **18**:237–42.

890   doi:10.1038/ng0398-237

891 108. Nataf S. Evolution, immunity and the emergence of brain superautoantigens.

892   *F1000Research* (2017) **6**:171. doi:10.12688/f1000research.10950.1

893 109. Nataf S. Autoimmunity as a Driving Force of Cognitive Evolution. *Front Neurosci* (2017)

894   **11**:582. doi:10.3389/fnins.2017.00582

895 110. James LM, Christova P, Lewis SM, Engdahl BE, Georgopoulos A, Georgopoulos AP.

896   Protective Effect of Human Leukocyte Antigen (HLA) Allele DRB1*13:02 on Age-

897   Related Brain Gray Matter Volume Reduction in Healthy Women. *EBioMedicine* (2018)

898   **29**:31–37. doi:10.1016/j.ebiom.2018.02.005

899 111. Wang Z-X, Wan Y, Tan L, Liu J, Wang H-F, Sun F-R, Tan M-S, Tan C-C, Jiang T, Tan L, et

900   al. Genetic Association of HLA Gene Variants with MRI Brain Structure in Alzheimer's

901   Disease. *Mol Neurobiol* (2017) **54**:3195–3204. doi:10.1007/s12035-016-9889-z