Abundance, fatty acid composition and saturation index of neutral lipids are significantly

different between isogenic primary and metastatic colon cancer cell lines

Rimsha Munir[1,2], Jan Lisec[3], Carsten Jaeger[4, 5] and Nousheen Zaidi[1,2*]

[1]Cancer Biology Lab, MMG, University of the Punjab, Lahore, Pakistan, [1]Cancer Research Centre, University of
the Punjab, Lahore, Pakistan, [3]Bundesanstalt für Materialforschung und -prüfung (BAM), Berlin, [4]Charité -
Universitätsmedizin Berlin, Molekulares Krebsforschungszentrum (MKFZ), Augustenburger Platz 1, 13353
Berlin, [5] Berlin Institute of Health (BIH), Anna-Louisa-Karsch-Straße 2, 10178 Berlin, Germany.

*Corresponding author: Nousheen Zaidi

Email: nosheen.mmg@pu.edu.pk, nzzaidi@yahoo.com

Abstract

Lipid droplets, the dynamic organelles that store triglycerides (TG) and cholesterol esters (CE), are

highly accumulated in colon cancer cells. This work studies the TG and CE subspecies profile in

colon carcinoma cells lines, SW480 derived from primary tumor, and SW620 derived from a

metastasis of the same tumor. It was previously reported that the total TG and CE content is

dramatically higher in SW620 cells; however, TG and CE subspecies profile has not been

investigated in detail. The presented work confirms that total TG and CE content is significantly

higher in SW620 cells. Moreover, the FA-composition of TGs is significantly altered in SW620 cells,

with significant decrease in the abundance of saturated triglycerides. This resulted in significantly

decreased TG saturation index in SW620 cells. The saturation index of CEs was also significantly

decreased in SW620 cells. The saturation indices of some other major lipid classes were either

similar or only moderately different between SW480 and SW620 cells. We also compared the

expression of metabolic genes that may regulate these changes in the lipidomic profiles.

Keywords: Cancer metabolism, colon cancer, fatty acid saturation, lipid droplets, lipidomics,

triglycerides.

Introduction

Lipidomic landscape of colorectal cancer (CRC) cells is significantly modified in comparison to that in their non-malignant counterparts (reviewed in [1]). Primary and metastatic colon cancer cells also display differences in their lipidomic profiles [2]. Previous works have taken advantage of SW480/SW620 cell line pair, which represents an accepted model to study metastatic progression [3]. SW480 is derived from a primary tumor, and SW620 is derived from a metastasis of the same tumor [4]. Multiple differences in the lipidomic profiles of these two cell lines have been noted [2]. For instance; increased levels of plasmanylcholine, while decreased levels of plasmenylethanolamine, C-16 containing sphingomyelin and ceramide were observed [2]. It has been shown that , SW620 cells display dramatic increase in total triglyceride (TG) and cholesterol ester (CE) levels [2]. However, the TG and CE subspecies profiles has not been explored in detail. Studying the TG and CE profile CRC cells is particularly important because these neutral lipids from hydrophobic core of lipid droplets (LDs) –that are highly accumulated in CRC cells [5]. Previous works have shown that exposure to excessive poly unsaturated fatty acids (PUFAs) induces LD formation in both SW480 and SW620 cells [6]. Interestingly, SW620 cells accumulate PUFA-enriched CEs, while SW480 cells accumulates PUFA-enriched TGs [6]. This indicates differential regulation of TG and CE content in SW480 and SW620 cells. Herein, we study the differences in TG- and CE-subspecies profiles and saturation index (SI) between SW480 and SW620 cells. Moreover, the expression of various metabolic genes that may bring about these lipidomic changes were also analyzed.

Materials and Methods

Cell culture

The SW480 and SW620, cell lines were purchased from American Type Culture Collection and were

maintained in DMEM (Gibco, 31966-021) supplemented with 10% fetal bovine serum (FBS) (Sigma,

F75240) and penicillin-streptomycin solution (Corning, 30-002-CI). Cell cultures were maintained

in the atmosphere of 5% CO_2 and 37°C. Cell lines were commercially authenticated (Eurofins,

Germany) and mycoplasma tested prior to submission of this manuscript.

Gene Expression Analysis

For quantitative RT-PCR, total RNA was extracted from cell pellets using Quick-RNA™ MiniPrep Plus

(Zymo Research). All RNA samples were reverse-transcribed into cDNA using SuperScript™ III

Reverse Transcriptase (Thermo Scientific, 18080093) and Oligo(dT)18 Primers (Thermo Scientific,

SO131). Quantitative PCR was performed using a TaqMan™ Gene Expression Master Mix (4369016,

Applied Biosystems) *via* StepOne Real-Time PCR Systems (Applied Biosystems). The TaqMan Gene

Expression assays used were Hs01005622_m1 (fatty acid synthase, FASN), Hs00168352_m1 (3-

hydroxy-3-methylglutaryl-CoA reductase, HMGCR), Hs00996004_m1 (monoglyceride lipase,

MGLL), Hs00173425_m1 (lipoprotein lipase, LPL) and Hs00354519_m1 (CD36). The expression of

each gene was normalized to the expression of GADPH (Hs02786624_g1). Microarray data were

obtained from publicly accessible gene expression profile database GEO

(http://www.ncbi.nlm.nih.gov/geo/). Provenzani *et al* [7] performed a microarray study, analyzing

differential gene expression in SW480 and SW620 cells (data accessible at NCBI GEO database,

accession GSE2509)

Lipid Extractions

First, the cell pellets were washed with 0.5 mL 0.9% NaCl. For extraction of lipids the pellets were

resuspended in 1 ml ice-cold MMC (1:1:1 v/v/v methanol/MTBE/chloroform). Samples were

incubated on an ultrasonic bath for two minutes. Phase separation was induced by adding 300 μL

MS-grade water. After 10 min incubation, the samples were centrifuged for 10 min at 1000 rpm and

the upper (organic) phase was collected. Then 200 μL of collected organic phase were dried in a

vacuum rotator and stored at –20 °C until analysis.

Determination of Total Triglyceride and Cholesterol Ester Contents

Total triglyceride content in the lipid extracts was spectrophotometrically determined using

commercially available kit (Analyticon Biotechnologies AG, Catalogue # 5052) against a calibration-

curve generated using known concentrations of triglyceride standard (SUPELCO, 17811-1AMP).

Cholesteryl esters were quantified using Cholesterol/ Cholesteryl Ester Quantitation Kit (Abcam,

ab65359) was used according to manufacturer's guidelines.

Lipidomic Profiling and Data Analysis

Dried sample extracts were reconstituted in 100 μL 2:1:1 v/v/v isopropanol/acetonitrile/water. 5

μL aliquots were injected into an ACQUITY I-class ultra-performance liquid chromatography

(UPLC) system (Waters, Germany) coupled to an Impact II high-resolution quadrupole time-of-

flight mass spectrometer (Bruker Daltonik GmbH, Germany). Chromatographic separation was

achieved by gradient elution (%A: 0 min, 60; 1.2 min, 57; 1.26 min, 50; 7.2 min, 46; 7.26 min, 30;

10.8 min, 0; 12.96 min, 0; 13.02 min, 60; 14.4 min, 60) using a buffered solvent system (A: 60:40

v/v acetonitrile/water, B: 90:10 v/v isopropanol:water, both with 10 mM ammonium formate and

0.1% formic acid) and a 2.1 mm × 75 mm × 1.7 μm CSH-C18 column (Waters, Germany) equipped

with an 0.2 μm pre-column in-line filter. The flow rate was 0.5 mL min-1 and column temperature

was 55°C. Electrospray ionization (ESI) conditions were as follows: polarity (+), capillary voltage,

4500 V, end plate offset, 500 V, nebulizer pressure, 2.5 bar, dry gas (N2) flow, 8 L/min. Ion transfer

parameters were set to: Funnel 1 RF, 200 Vpp, Funnel 2 RF, 200 Vpp, Hexapole RF, 50 Vpp,

Quadrupole Ion Energy, 5 eV, Low Mass, 100 m/z, Collision Energy, 8.0 eV, Pre Pulse Storage, 6.0 μs,

Stepping Mode, Basic, Collision RF, 500-1000 Vpp, Transfer Time, 60-100 µs, Timing, 50/50,

Collision Energy, 100-250%. Alternating MS and MS/MS scans were acquired using a Sequential

Windowed Acquisition of All Theoretical Fragment Ion Mass Spectra (SWATH) scheme (m/z 350-

975, width 25 Da). For internal calibration, Na formate clusters were spiked into the LC effluent at

the end of each run. After data acquisition, files were converted to Analysis Base File (ABF) format

using a publicly available converter (Reifycs, Japan) and imported into MS-DIAL (Tsugawa et al.

2015). MS-DIAL parameter settings were as follows: Soft Ionization, Data independent MS/MS,

Centroid data, Positive ion mode, Lipidomics. Detailed analysis settings were left at default, except

for Retention time end (10 min), Alignment Retention time tolerance (0.2 min), Identification

Retention time tolerance (3 min) and Identification score cut off (60%).

For each of 6 replicate samples per cell line we calculated within each major lipid class the relative

distribution of lipids containing none (SFA0), exactly one (SFA1) and two or more saturated fatty

acids (SFA2). To this end, the measured ion intensities for the groups (SFA0, 1 and 2) were summed

up and normalized to the total sum of this lipid class. Mean values and standard deviations for the

SFA distribution were calculated over the 6 replicates per cell line. Saturation indices were

calculated as a ratio of total saturated fatty acid content to total unsaturated fatty acid content [8]

within each major lipid class. Total content of saturated fatty acids was calculated by summing up

the intensities of all lipids multiplied by the number of saturated fatty acids present within each

lipid. Total content of un-saturated fatty acids in each individual lipid class was calculated by

summing up the intensities of all lipids multiplied by the number of un-saturated fatty acids present

within each lipid. To compare SI over lipid classes we normalized with respect to the SI value

obtained for SW480.

Statistical analysis

The differences between groups were analyzed by ANOVA or t-test (paired or unpaired), where

applicable. Statistical analyses and graphical representations for lipidomic data and quantitative

RT-PCR data were performed using the R software environment 3.4.2 (http://cran.r-project.org/)

or MetaboAnalyst 3.5 (http://www.metaboanalyst.ca/faces/home.xhtml). P-values <0.05 were

considered statistically significant and indicated when different.

Results and Discussion

A previous study has shown a dramatic increase in abundance of total cholesterol esters (CEs) and

triglycerides (TGs) in SW620 cells in comparison to SW480 cells [2]. We also assessed the total TG

and CE content and observed that SW620 cells display 2.5-3 fold increase in the levels these storage

lipids (**supplementary figure 1**). Next, the fatty acid (FA) composition of TGs and CEs was

compared. We observed that in SW480 cells the triglycerides (TGs) with ≥2 saturated fatty acids

(SFA) were in the highest proportion (i.e. 61%). The proportion of TGs containing 1 or 0 SFA was

35% and 4%, respectively. These proportions were significantly altered in SW620 cells in which the

proportion of TGs with ≥2 saturated fatty acids was decreased to 40%, while the proportions of TGs

containing 1 or 0 SFA were respectively increased to 53% and 7%. In agreement to these results,

the TG saturation index (SI) was markedly higher in SW480 cells than in SW620 cells. On the other

hand, FA-composition and SI of diacylglycerols (DGs), the direct precursors to TGs, was only slightly

altered. The FA-composition of CEs also showed minor changes, with slight shift in the proportions

of CEs containing 0 or 1 SFA. However, the SI of CEs was also significantly decreased in SW620 cells.

We also compared the FA composition of the most abundant membrane lipids,

phosphatidylcholines (PCs) and phosphatidylethanolamine (PEs), in SW480 and SW620 cells. No

substantial differences were observed in fatty acid composition and SI of PCs and PEs (**Figure 1**).

Hence, TGs constitute the major lipid class that displays changes in FA composition in SW620 cells

when compared to SW480 cells.

To understand the molecular mechanisms regulating these changes in the lipidomic profiles, we

performed expression analyses and also used publicly accessible gene expression profile database

GEO (http://www.ncbi.nlm.nih.gov/geo/). **Figure 2** displays overview of the major cancer-

associated lipid metabolism pathways. The fold-changes in the expression levels of various

genes/enzymes are indicated with a colored-box on *left-side* of the respective gene symbol. For

color-codes see color-key on *top-left* corner of **Figure 2**. As discussed above, the differences in the

FA composition of TGs was the most significant difference between lipidomic profiles of SW480 and

SW620 cells, with SW620 cells displaying significantly lower TG saturation-index. A recent study

has shown that gene silencing of diglyceride acyltransferases (DGATs) –enzymes that catalyze

formation of TGs from DGs and Acyl CoA– induces increase in TG saturation-index [9]. Hence,

increased expression of DGAT might induce decreased SI in TGs. Our analyses revealed that

expression of DGAT was slightly decreased (1-1.5 Fold) in SW620 cells (**Figure 2, TG Synthesis**).

Hence, DGAT expression might not be responsible for the changes in TG saturation-index of SW620

cells.

Next, we looked at other lipid metabolism pathways that have been implicated in regulation of lipid

saturation-indices. Highly active *de novo* FA synthesis induces increased SFAs or MUFAs levels in

cancer cell cells [10]. The expression of FA synthesis genes is up-regulated in SW620 cells, hence

this pathway is also not contributing in decreased SI in these cells (**Figure 2, FA Synthesis**).

Major genes of intracellular lipolytic pathway –including PNPLA2, HSL and MGLL– are all down

regulated in SW620 cells (**Figure 2, Intracellular Lipolysis**). This may explain the dramatic

increase in total triglyceride (TG) levels in SW620 cells. However, specific decrease in ≥2 SFA

containing TGs could not be explained by this mechanism.

The expression of ACADM and ACADL –the enzymes that catalyze the first step of FA oxidation in

mitochondria– is differentially regulated in SW620 cells. The expression of ACADM was up

regulated, while the expression of ACADL was down-regulated in SW620 in comparison to SW480

cells (**Figure 2, FA Oxidation**). This observation is of special interest in this context, because

ACADL is known to mediate unsaturated FA oxidation, whereas ACDM prefers saturated FAs as

substrates. Hence, it is possible that in SW620 cells there is an increased oxidation of SFAs, while

decreased oxidation of unsaturated FAs. This may cause selective depletion of SFAs in the stored

lipids and induced decreased SI of TGs and CEs. Recent studies have shown that disruptions in lipid

metabolism or nutrient/oxygen supply also drastically affects FA-composition of TGs in cancer

cells, while the FA-composition of other lipid classes is not significantly affected [9,11]. Hence, FA-

composition of TGs is most markedly affected by the changes in cancer stage, tumor

microenvironment and de novo FA synthesis. Further studies are required to understand the

significance of fatty acid composition of TGs in cancer progression and survival.

Abbreviations

FASN, fatty acid synthase; HMGCR, 3-hydroxy-3-methylglutaryl-CoA reductase; MGLL, monoacylglycerol lipase; LPL, lipoprotein lipase; PC, Phosphatidylcholines; PE, Phosphatidylethanolamines; PPE, Plasmenylphosphatidylethanolamines; PPC, Plasmenylphosphatidylcholines; TG, Triacylglycerols; CE, Cholesterol esters and DG, Diacylglycerols.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of Data and Material

The datasets supporting the conclusions of this article are included within the article and supplementary files.

Competing Interests

The authors declare no potential competing interests.

Funding

This work was financially supported by Alexander von Humboldt Foundation (Georg Forster Research Fellowship for Experienced Researchers awarded to N. Zaidi) and University of the Punjab under Projects for Cancer research Centre (PI: N. Zaidi). Funding bodies did not play any role in study designing; data collection, analysis, & interpretation or manuscript writing.

Acknowledgments

We are grateful to Mr. Banaras Masih for his technical assistance for this project.

245 **Authors' Contributions**

246 This study was designed by NZ. Experimental work was done by NZ, CJ and RM; and data was

247 analyzed and interpreted by JL, CJ, NZ and RM. The manuscript was written by RM and NZ; and

248 reviewed and approved by all authors.

References

[1] A. Pakiet, J. Kobiela, P. Stepnowski, T. Sledzinski, A. Mika, Changes in lipids composition and metabolism in colorectal cancer: a review, Lipids Health Dis 18 (2019) 29.

[2] C.J. Fhaner, S. Liu, H. Ji, R.J. Simpson, G.E. Reid, Comprehensive lipidome profiling of isogenic primary and metastatic colon adenocarcinoma cell lines, Anal Chem 84 (2012) 8917-8926.

[3] R.E. Hewitt, A. McMarlin, D. Kleiner, R. Wersto, P. Martin, M. Tsokos, G.W. Stamp, W.G. Stetler-Stevenson, Validation of a model of colon cancer progression, J Pathol 192 (2000) 446-454.

[4] A. Leibovitz, J.C. Stinson, W.B. McCombs, 3rd, C.E. McCoy, K.C. Mazur, N.D. Mabry, Classification of human colorectal adenocarcinoma cell lines, Cancer Res 36 (1976) 4562-4569.

[5] L. Tirinato, C. Liberale, S. Di Franco, P. Candeloro, A. Benfante, R. La Rocca, L. Potze, R. Marotta, R. Ruffilli, V.P. Rajamanickam, M. Malerba, F. De Angelis, A. Falqui, E. Carbone, M. Todaro, J.P. Medema, G. Stassi, E. Di Fabrizio, Lipid droplets: a new player in colorectal cancer stem cells unveiled by spectroscopic imaging, Stem Cells 33 (2015) 35-44.

[6] S.A. Schonberg, A.G. Lundemo, T. Fladvad, K. Holmgren, H. Bremseth, A. Nilsen, O. Gederaas, K.E. Tvedt, K.W. Egeberg, H.E. Krokan, Closely related colon cancer cell lines display different sensitivity to polyunsaturated fatty acids, accumulate different lipid classes and downregulate sterol regulatory element-binding protein 1, Febs j 273 (2006) 2749-2765.

[7] A. Provenzani, R. Fronza, F. Loreni, A. Pascale, M. Amadio, A. Quattrone, Global alterations in mRNA polysomal recruitment in a cell model of colorectal cancer progression to metastasis, Carcinogenesis 27 (2006) 1323-1333.

[8] D. Ackerman, S. Tumanov, B. Qiu, E. Michalopoulou, M. Spata, A. Azzam, H. Xie, M.C. Simon, J.J. Kamphorst, Triglycerides Promote Lipid Homeostasis during Hypoxic Stress by Balancing Fatty Acid Saturation, Cell Rep 24 (2018) 2596-2605.e2595.

[9] D. Ackerman, S. Tumanov, B. Qiu, E. Michalopoulou, M. Spata, A. Azzam, H. Xie, M.C. Simon, J.J. Kamphorst, Triglycerides Promote Lipid Homeostasis during Hypoxic Stress by Balancing Fatty Acid Saturation, Cell reports 24 (2018) 2596-2605. e2595.

[10] E. Rysman, K. Brusselmans, K. Scheys, L. Timmermans, R. Derua, S. Munck, P.P. Van Veldhoven, D. Waltregny, V.W. Daniëls, J. Machiels, De novo lipogenesis protects cancer cells from free radicals and chemotherapeutics by promoting membrane lipid saturation, Cancer research (2010) 0008-5472. CAN-0009-3871.

[11] J. Lisec, C. Jaeger, R. Rashid, R. Munir, N. Zaidi, Cancer cell lipid class homeostasis is altered under nutrient-deprivation but stable under hypoxia, BMC Cancer 19 (2019) 501.

Figure Legends

Figure 1: Comparison of fatty acid composition and saturation indices in major lipid classes

in SW480 and SW620 cells. The figure is structured in five rows and three columns. Rows show

data for the major lipid classes Triglycerides (TG), Diglycerides (DG), Cholesterol Esters (CE),

Phosphatidylethanolamines (PE) and Phosphatidylcholines (PC), Left column displays lipid class ID

and a legend indicating the number of lipid-subspecies within each lipid class containing different

numbers of saturated fatty acids (0, 1 or ≥2SFAs). Center column shows the relative amount (in per

cent of summed ion intensity) of lipid subspecies within both SW cell lines. Numeric values for

mean and standard deviation calculated over 6 replicate samples per cell line are given within the

bars. Right column displays the saturation indices (cf. Method section) of the respective lipid class

normalized to SW480.

Figure 2: Fold-changes in expression levels of various genes in major lipid metabolism

pathways in SW620 *vs.* SW480 cells. The figure highlights all the key lipid metabolism pathways

activated in cancer cells. Major pathways are shown as boxes without outlines. The systematic

names of these pathways are given in each box. The fold-changes in the expression levels of various

genes/enzymes in SW620 cells compared with SW480 cells are indicated with a colored-box on *left-*

side of the respective gene symbol. For color-codes see color-key on *top-left* corner. The expression

data for these analyses were extracted from at NCBI GEO database, accession GSE2509. For some

markers gene expression analysis was performed (**Supplementary Figure S2**). The symbols within

fold-change boxes represent the data source (G; GEO database, S, Supplementary Figure S2).

Abbreviations: ACACA, acetyl-CoA carboxylase 1; ACACB, acetyl-CoA carboxylase 2; ACSS2, acyl

Co-A synthetase-2; ADFP, adipose differentiation protein; ATGL, adipose triglyceride lipase; FA,

fatty acids; FABP3, fatty acid binding protein 3; FABP7, fatty acid binding protein 7; FFA, free fatty

acids; FASN, fatty acid synthase; H, hypoxia; HIF-1α, hypoxia-inducible factor 1-alpha; HIF-2α,

hypoxia-inducible factor 2α; HIG2, hypoxia-inducible gene 2 protein; HMGCR, 3-hydroxy-3-

methylglutaryl-CoA reductase; LD, lipid droplet; LCAD, long-Chain Specific acyl-CoA

dehydrogenase; MAG, monoacylglycerol ; MCAD, Medium-chain acyl-CoA dehydrogenase; MUFA,

monounsaturated fatty acids; PBMCs, peripheral blood mononuclear cells; PC,

Phosphatidylcholines; PE, phosphatidylethanolamines; Pcho, propargyl-choline; PI,

phosphatidylinositol; PS, phosphatidylserine; PUFA, polyunsaturated fatty acids; SCD, stearoyl-CoA

Desaturase; SFA, saturated fatty acids; SREBP, sterol regulatory element-binding proteins; TG,

triglycerides.

Supplementary Figure S1: Fold-changes in total triglyceride(TG) and cholesterol ester (CE) content in SW480 vs. SW620 cells.

Supplementary Figure S2: Expression of selected genes from de novo lipid synthesis or lipid uptake/degradation pathways in SW480 and SW620 cell lines. Box plots showing log2 transformed and median normalized values of FASN, HMGCR and MGLL. LPL and CD36 expressions were not detected. The levels of the different transcripts were measured in 3 to 6 samples by qPCR. The results show the distribution of corresponding transcripts relative to GAPDH, with the box indicating the 25th–75th percentiles, with the median indicated. line. The whiskers show the range.