1

3 # Y-chromosome haplotypes drive variation in size and age at maturity in male
4 # Chinook salmon

8 Garrett J. McKinney[1a*], James E Seeb[1b], Carita E. Pascal[1c], Daniel E. Schindler[1d], Sara E. Gilk-Baumer[2e],
9 Lisa W. Seeb[1f]

[1] School of Aquatic and Fishery Sciences, University of Washington, 1122 NE Boat Street, Box 355020,

Seattle WA 98195-5020, USA.

[2] Alaska Department of Fish and Game, 333 Raspberry Road, Anchorage, AK, 99518, USA

[*]**Corresponding author:** Garrett J. McKinney (email: gjmckinn@uw.edu, phone: 1-765-430-3272).
Current address: NOAA National Marine Fisheries Service, Northwest Fisheries Science Center, 2725
Montlake Blvd E, Seattle, WA 98112

Emails: [a] gjmckinn@uw.edu; [b]jseeb@uw.edu; [c]cpascal@uw.edu; [d] deschind@uw.edu;
[e]sara.gilk@alaska.gov ; [f]lseeb@uw.edu

Keywords: Chinook salmon, Y chromosome, haplotypes

ABSTRACT

Variation in size and age at maturity is an important component of life history in salmon that is influenced

both by environmental and genetic factors. Large size confers a direct reproductive advantage through

increased fecundity and egg quality in females, while larger males gain a reproductive advantage by

monopolizing access to females. In addition, variation in size and age at maturity in males can be

associated with different reproductive strategies; younger smaller males may gain reproductive success by

sneaking in among mating pairs. In both sexes there is a trade-off between older age and increased

reproductive success and increased risk of mortality by delaying reproduction. We used RADseq data for

21 populations of Alaska Chinook salmon (*Oncorhynchus tshawytscha*) and identified four Y-

chromosome haplotypes that showed regional and population-specific variation in frequency. These

haplotypes exhibited associations with size at maturity in multiple populations suggesting that the lack of

recombination between X and Y-chromosomes have allowed Y-chromosome haplotypes to capture

different alleles that influence size at maturity. Ultimately, conservation of life history diversity in

Chinook salmon may require conservation of Y-chromosome haplotype diversity.

Introduction

Variation in life history within populations is common across taxa and is often associated with alternative strategies for increasing fitness. This includes partial migration, where some individuals of a population migrate while others remain resident (Chapman et al. 2011), reproductive morphs that exhibit different mating strategies or sexually selected traits (Shuster 1989, Johnston et al. 2013, Küpper et al. 2015), age and size at maturity (Gibbons et al. 1981), and even length of life-span such as annual vs perennial plants (Hall et al. 2006). While life history variation is often assumed to be under the influence of many genes of small effect, examples of a simpler genetic basis of life history variation are increasingly being found, such as single genes that influence age at maturity and sexually selected traits (Johnston et al. 2013, Barson et al. 2015) and chromosome inversions contributing to annual versus perennial live history and variation in migration (Pearse et al. 2014, Twyford and Friedman 2015). The mechanism underlying variation in life history has important implications for how life history diversity is maintained under different selective regimes.

Variation in life history strategies is exhibited by many salmon species with size and age at maturity being an important component of this variation. In females, large body size confers a direct reproductive advantage through increased fecundity and egg quality (Healey and Heard 1984). In contrast, variation in size and age at maturity in males can be associated with different reproductive strategies that exhibit frequency dependent fitness. For example, older larger males gain reproductive success by monopolizing access to females while younger smaller males gain reproductive success by sneaking in among mating pairs (Healey and Heard 1984, Berejikian et al. 2010). For both sexes, there is a trade-off to delayed maturation where increased reproductive success is tempered by an increased risk of mortality before reproduction. The optimal age at maturity for a population should represent the balance between reproductive benefits and mortality costs of delayed maturation (Healey 1986), and forces that change these costs or benefits could result in shifts in age composition. Nonetheless, considerable diversity in age at maturation is maintained within many populations, presumably as a bet-hedging strategy to spread risks over the life-cycle of these fish.

Age at maturity in salmon is generally thought to be a threshold trait that is dependent either upon reaching a minimum size at age or upon growth rate at key periods (Healey 1991, Thorpe 2007). Environmental factors that influence growth rate have been shown to influence age at maturity in multiple salmon species. In the wild, studies show correlations between ocean conditions such as temperature and productivity and patterns of age at maturity (Otero et al. 2012, Siegel et al. 2017). In experimental

settings, age at maturity was manipulated through changing temperature (Heath et al. 1994, Harstad et al.

2018) or food ration (Rowe and Thorpe 1990, Larsen et al. 2006).

In addition to environmental effects, multiple lines of evidence show a genetic component to age

at maturity. Age at maturity is highly heritable in multiple salmon species (Gjerde 1984, Gall et al. 1988,

Hankin et al. 1993, Heath et al. 1994), and Quantitative Trait Locus (QTL) and Genome-Wide

Association (GWAS) studies identified genomic regions associated with variation in age at maturity

(Barson et al. 2015, Kodama et al. 2018, Micheletti and Narum 2018, Waters et al. 2018). Studies also

demonstrated that offspring of alternative male phenotypes exhibit different growth rates (Garant et al.

2002, Berejikian et al. 2011). In Atlantic salmon (*Salmo salar*), individuals with different life histories

exhibit differing maturation thresholds that are genetically based (Aubin Horth and Dodson 2004).

Despite these findings, the genetic basis of maturation age in most salmonids remains poorly understood.

One complicating factor is that the genetic architecture underlying variation in age at maturity

appears to vary among salmonid species and even among populations within a species. In Atlantic

salmon, age at maturity is strongly influenced by a single gene (VGLL3) (Ayllon et al. 2015, Barson et al.

2015). This gene exhibits sex-dependent dominance which facilitates sexually antagonistic selection

(Barson et al. 2015). While this gene explained 39% of the phenotypic variability in European Atlantic

salmon, studies in North American Atlantic salmon and in Pacific salmon have not found a similar

association (Micheletti and Narum 2018, Boulding et al. 2019).

Chinook salmon (*Oncorhynchus tshawytscha*) are the largest of the Pacific salmon and follow

various life history strategies spending 0 to 2 years in fresh water and 1 to 4 or more years in the ocean

(Riddell et al. 2018).   Male Chinook salmon exhibit significant variation in size and age at maturity

(Healey 1991) that is linked to differential reproductive tactics and is likely controlled by both

environmental and genetic components (i.e. sneaker vs. dominant males, Berejikian et al. 2010, Young et

al. 2013). Historically, males predominantly matured at age four or older; however, many populations

throughout North America recently experienced marked declines in size and age at maturity (Lewis et al.

2015, Ohlberger et al. 2018). Explanations for decreased age at maturity have focused on the impacts of

fisheries induced evolution or changing environmental conditions (Hard et al. 2008, Kendall et al. 2014,

Siegel et al. 2017) though this explanation is not particularly consistent with the data (Ohlberger et al.

2018). While the genetic control of age at maturity in Chinook salmon is still poorly understood, past

studies offer clues to genomic regions that may be associated with maturation age. In particular, Heath et

al. (2002) identified a strong sex-linked component to age at maturity in Chinook salmon, suggesting the

influence of genes on the Y-chromosome.

The X and Y chromosomes in most salmonid species are morphologically undifferentiated

(Davidson et al. 2009), and available sequence data suggests that the primary difference between sex

chromosomes is an insertion containing the sex-determining gene (SDY, Yano et al. 2013). Despite a

lack of large-scale differentiation, the X- and Y-chromosomes could show sequence divergence due to

sex-specific patterns of recombination. Recombination in females takes place along the full length of the

chromosome while recombination in males is strongly localized to telomeric regions (Lien et al. 2011),

restricting recombination between the X- and Y-chromosomes. Reduced recombination between sex

chromosomes is supported by a 33Mb signal of sex association observed in Atlantic salmon (Kijas et al.

2018). In addition to facilitating divergence between sex chromosomes, sex-specific recombination could

lead to the formation of different Y-chromosome haplotypes and the capture of adaptive genetic variants.

We identified four Y-chromosome haplotypes in Chinook salmon from Alaska that showed

regional and population-specific variation in frequency. These haplotypes showed associations with size

at maturity in multiple populations suggesting that the lack of recombination between X and Y-

chromosomes has allowed Y-chromosome haplotypes to capture different genetic variants influencing

size and age at maturity.

Materials and Methods

RAD Y-chromosome Haplotypes

We used existing RADseq data to examine patterns of genetic variation in the sex chromosome.

RADseq data for 21 populations of Alaska Chinook salmon were obtained from previous studies (Larson

et al. 2014, Dann et al. 2018, McKinney et al. 2018, McKinney et al. In Revision). RADseq data from

Larson et al. (2014) and McKinney et al. (2018) are available in NCBI SRA accessions SRP034950 and

SRP129894; raw data from Dann et al. (2018) and McKinney et al. (In Revision) are available from those

authors upon request. Locations of populations ranged from Cook Inlet to the Upper Yukon River and

include a total of 1,082 samples (Table 1). RADseq data were processed with Stacks V1.7 (Catchen et al.

2011, Catchen et al. 2013) using default settings with the following exceptions: process_radtags (-c -r -q

-filter_illumina -t 94), ustacks (-m 2 -M 2, --model_type bounded --bound_high 0.05), cstacks (-n 2). The

catalog of variation from McKinney et al. (In Revision) was used for consistent locus names among this

and previous studies. A total of six individuals per population from Cook Inlet were added to this catalog

to allow for additional allelic variation.

Paralogs are common in salmonid genomes due to an ancestral whole-genome duplication

(Allendorf and Thorgaard 1984) but cannot be reliably genotyped in typical RADseq studies because of

insufficient read depth (McKinney et al. 2018). Paralogs were identified using *HDplot* (McKinney et al.

2017) and removed from further analysis. Loci with more than 10% missing data and a minor allele

frequency (MAF) < 0.05 were also removed from analysis. Finally, loci were aligned to the Chinook

salmon genome (Christensen et al. 2018) using Bowtie2 (Langmead and Salzberg 2012) to determine

genomic position; only loci that aligned to the sex chromosome (Ots17, Phillips et al. 2013) were retained

for analysis.

Patterns of variation on the sex chromosome were visualized using principal component analysis

(PCA). Clusters of samples on the PCA were identified visually; markers that differentiated clusters of

samples were identified by estimating F_{ST} per marker between clusters using Genepop (Rousset 2008).

Markers that differentiated clusters showed high linkage disequilibrium (LD) within clusters, suggesting

inhibited recombination and exhibited genotype patterns consistent with Y-chromosome haplotypes. Full

details are available in supplemental materials and methods (File S1).

Markers that differentiated clusters and appeared to be diagnostic for Y-chromosome haplotypes

were made into an amplicon panel (GT-seq , Campbell et al. 2015) for expanded sampling. Markers that

were unlikely to amplify single genomic regions were filtered prior to primer design following the

methods of McKinney et al. (In Revision). Primers were designed using batch primer3 (You et al. 2008),

and genome sequences were used to extend the original RAD sequence for primer design where

necessary.

Expanded sampling

Additional samples ranging from Alaska to California were genotyped using the GT-seq panel to

establish the geographic distribution of the haplotypes. In addition to the Y-haplotype markers, this panel

included the sex identification marker Ots_sexy3-1 (Hess et al. 2016a) to confirm that Y-chromosome

haplotypes were present only in male fish. A total of 1,341 samples from 27 populations were genotyped

(Table 1). Samples were sequenced on a HiSeq 4000, and data were processed and genotyped using GT-

score (McKinney et al. In Revision) available at https://github.com/gjmckinney/GTscore.

Y-chromosome haplotype analyses

Y-chromosome haplotype assignments were combined for GT-seq and RADseq samples to

characterize the distribution and frequency of Y-chromosome haplotypes. Haplotypes were assigned to

each sample based on multi-locus genotype patterns (File S1). Data for length and age at maturity were

obtained from Alaska Department of Fish and Game (ADFG) for populations in Alaska and compared

166 with Y-chromosome haplotype data to determine if there were relationships between Y-chromosome

167 haplotype and length and age at maturity. Significance of associations between Y-chromosome

168 haplotypes and length were assessed using a Tukey test while significance of associations between Y-

169 chromosome haplotypes and age at maturity were assessed using a chi-square test. There are multiple

170 methods of reporting age in salmon (Koo 1962); we report freshwater and ocean age for each individual

171 using European notation, so an individual with an age of 1.3 would have spent 1 year growing in

172 freshwater after emergence followed by 3 years in the ocean for a total age of 5 years.

173 Results

174 RADseq Y-haplotype discovery

175  Analysis of the Y-chromosome showed multiple patterns consistent with Y-chromosome

176 haplotype blocks. Individuals resolved into multiple distinct clusters when PCA was done on the sex

177 chromosome (Figure S1-S3). Samples showed clustering by region (Cook Inlet vs Western Alaska) and

178 also clustering by sex with multiple clusters of male samples (Figure S4). Some putative females were

179 present in male clusters and vice versa; however, individuals had been visually sexed based on external

180 features which is known to have variable accuracy (Lozorie and McIntosh 2014). Ranking loci by F_{ST}

181 and examining genotype patterns revealed a set of 50 SNPs with $F_{ST} > 0.15$ that differentiated clusters

182 (Table S2.1). A general pattern was observed for loci with high F_{ST} among clusters; within each region

183 one cluster contained samples that were homozygous for a single allele for all high F_{ST} loci while the

184 other clusters showed a pattern of high to fixed heterozygosity for different sets of high F_{ST} loci. In Cook

185 Inlet, the homozygous cluster contained phenotypic females while heterozygous clusters contained

186 phenotypic males (Figure S4). A total of four different patterns of fixed heterozygosity were observed

187 throughout Yukon River, Western Alaska, and Cook Inlet populations suggesting the presence of

188 conserved Y-chromosome haplotype blocks throughout western and southcentral Alaska (Figures 1, S5,

189 S6). Within Alaska, Y-chromosome haplotypes showed regional variation in frequency with the AK Y1

190 haplotype being most frequent in Western Alaska and the AK Y3 haplotype the most frequent in Cook

191 Inlet (Figure 2).

192 GT-seq Y-chromosome haplotype expanded sampling

193  A GT-seq panel was developed to genotype Y-chromosome haplotype markers for a set of

194 samples representing the North American range of Chinook salmon. A total of 23 of the 50 RAD loci

195 passed filtering criteria prior to primer design; of these 14 RAD loci were successfully converted to GT-

196 seq assays (Table S1, Table S2.2). Genotyping of the expanded sample set confirmed that Y-

197 chromosome haplotypes were sex-specific; 200 (81%) of the genetic males from Alaska were assigned a

Y-chromosome haplotype and no genetic females had genotype patterns consistent with the Y-
chromosome haplotype blocks (Table S2.2). Males that were not assigned Y-chromosome haplotypes
were concentrated in Southeast Alaska; 40 of the 46 unassigned males were from the Little Port Walter
and Pullen Creek populations. Only 16% of males could be assigned a haplotype in Little Port Walter
and a single male from Pullen Creek could be assigned a haplotype. Excluding these two populations
results in 97% haplotype assignment of male Chinook salmon in Alaska.   For the remaining six
unassigned males, four were ambiguous for AK Y2 vs AK Y3 haplotypes and two were homozygous for
all Y-chromosome haplotype markers. Distribution of Y-chromosome haplotypes varied regionally: Y-
chromosome haplotype blocks identified in Alaska were not found in Chinook salmon outside of Western
and Southcentral Alaska and had only a rare occurrence in Southeast Alaska.

Size and Age at Maturity

Size at maturity data were available for eight of the populations in this study; four populations
were genotyped using RADseq data, three were genotyped using GT-seq data, and one was genotyped
using both RADseq and GT-seq. Populations were grouped by region (Yukon River, Western Alaska,
and Cook Inlet) for visualization. Boxplots of size at maturity for each Y-chromosome haplotype showed
a consistent relationship throughout western and southcentral Alaska. The AK Y1 and AK Y2 haplotypes
had the smallest individuals, the AK Y3 haplotype was associated with intermediate sized fish, and the
AK Y4 haplotype was associated with the largest individuals in each region (Figure 3). In the Yukon
River, individuals with the AK Y4 haplotype were significantly larger ($p<0.05$) than fish with the AK Y1
or AK Y2 haplotype, in Western Alaska, the fish with the AK Y3 haplotype were significantly larger fish
than the AK Y1 haplotype, and in Cook Inlet, size differences between the AK Y1, AK Y3, and AK Y4
haplotypes were statistically significant ($p<0.05$) (Figure 3). Despite a tendency for differences in size
distributions, the length of fish with the AK Y1 haplotype was not significantly different than the AK Y3
haplotype in the Yukon River, and the AK Y4 haplotype was not significantly different than other
haplotypes in Western Alaska. This is likely due to low samples size for these haplotypes.

Age at maturity data were available for the Lower Yukon Test Fishery. Histograms of age at
maturity for each Y-chromosome haplotype revealed that the AK Y3 haplotype had approximately three
times as many 1.3 fish as 1.4 fish while the Y4 haplotype had nearly even proportions of 1.3 and 1.4 fish
(Figure 4). While suggestive, this result was not significant, possibly due to the low samples size both of
age 1.4 fish and of fish with the AK Y4 haplotype. Boxplots of size at age showed that fish with the AK
Y4 haplotype were larger for age 1.3 and 1.4 than fish with the AK Y3 haplotype. Results were
statistically significant for age 1.3 fish but not age 1.4, possibly due to low sample size. The smallest fish

had the AK Y1 and AK Y2 haplotypes; however, only a single fish had both size and age data for these
haplotypes.

Discussion

Size and age at maturity are ecologically and evolutionarily important traits in Chinook salmon.
Numerous studies have examined ongoing declines in age at maturity; however, it has been difficult to
disentangle the interactions of environmental and genetic causes of this decline.   Size and age associated
markers and genes have previously been identified in genetic studies of Chinook salmon (Micheletti and
Narum 2018, Waters et al. 2018); however, results were not consistent across populations, and no markers
were located on the sex chromosome.   We show that a conserved set of Y-chromosome haplotypes is
associated with variation in size and age at maturity in Chinook salmon across Western and Southcentral
Alaska. This observation opens a new line of research into the genetics of age at maturity in salmonids.

Range-wide distribution of haplotypes

Chinook salmon are represented by multiple genetically distinct lineages throughout the species
range (Waples et al. 2004, Beacham et al. 2006, Moran et al. 2012). These lineages often show little gene
flow among them due to differences in geographic range or spawn timing, and this isolation may result in
different sets of Y-chromosome haplotypes and patterns of recombination across lineages. The Y-
chromosome haplotypes we identified through extended linkage disequilibrium were consistently
observed throughout Chinook salmon populations from the Upper Yukon River south to Little Port
Walter in Southeast Alaska. The occurrences become rarer in Southeast Alaska, and no individuals south
of Southeast Alaska could be assigned haplotypes suggesting that the haplotypes identified within Alaska
are regionally restricted. This corresponds with observed breakpoints between Chinook salmon lineages
near Cape Fairweather (Templin et al. 2011) which is approximately 200 miles northwest of Little Port
Walter and 100 miles west of Pullen Creek. It is likely that other haplotype blocks and genomic regions
of low recombination exist outside of Alaska; this could be determined by examining reduced-
representation or whole-genome sequence data from additional populations.

Populations throughout Alaska showed variation in haplotype frequency which may be
ecologically significant given the association between haplotypes and size and age at maturity.
Populations from Western Alaska and the Lower Yukon River had a greater proportion of the AK Y1 and
AK Y2 haplotypes which were associated with smaller fish. Populations in the Upper Yukon River and
Cook Inlet were primarily composed of AK Y3 and AK Y4 haplotypes which were associated with larger
fish. While we did not have enough samples with size data to characterize size distributions within
region, this finding is consistent with a long-term analysis of Chinook salmon returns by Lewis et al.

(2015) who reported smaller fish on average in Kuskokwim and Nushagak river populations from
Western Alaska relative to Yukon River and Cook Inlet populations. In addition, the Cook Inlet
populations sampled in this study are adjacent to the Kenai River which has historically produced large
Chinook salmon (Lewis et al. 2015, Schoen et al. 2017). The relationship between size and haplotype also
varied by region, with the Yukon River having the smallest difference in sizes between haplotypes and
Cook Inlet having the largest difference in sizes (Figure 3). Taken together, these results suggest that
differing frequencies of Y-chromosome haplotypes may contribute to regional variation in size of
Chinook salmon, and that the effect of haplotype on size can vary between regions, potentially due to
other genetic influence or different environmental conditions.

Importance of Y-haplotypes for life history diversity

271
Male Chinook salmon exhibit life history diversity related to maturation age with dominant males
successfully spawning by monopolizing access to females and jacks obtaining reproductive success by
sneaking in among mating pairs. This diversity has been eroding as the proportion of dominant (age 4+)
males declines and jacks (3 years or younger) increase in frequency (Ohlberger et al. 2018). The alternate
life histories of male Chinook salmon exhibit frequency dependent fitness which in theory should exhibit
stable proportions; however, this assumes populations are at equilibrium. Male sockeye salmon
(*Oncorhynchus nerka*) exhibit similar frequency dependent life history variation but persistent
demographic variation occurs as a result of strong selection events coupled with variation in recruitment
(DeFilippo et al. 2019). If jacks are produced by specific Y-chromosome haplotypes, then fishing
practices that increase the proportion of jacks may result in demographic shifts and loss of age diversity
that are difficult to recover. The markers we developed can be used to characterize historic (i.e., from
archived samples) and current Y-chromosome haplotype diversity in Alaska Chinook salmon to
determine if demographic shifts correspond with shifts in frequencies of Y-chromosome haplotypes.
Ultimately, conservation of life history diversity in Chinook salmon may require conservation of Y-
chromosome haplotype diversity.

Hypotheses of population structure and delineation of management units using genetic data are
typically based on genome-wide analyses consistent with the assumption that major life history traits are
controlled by many genes with small effects.   Waples and Lindley (2018) recently commented on the
new challenges facing existing conservation frameworks when associations are identified between one or
a very few genes and key life history traits. Their comment was prompted by the recent identification of
SNPs from a GREB1L gene that explain a large proportion of the variation associated with adult
migration time in steelhead (*O. mykiss*) and Chinook salmon (Hess et al. 2016b, Prince et al. 2017).
Conservation of the Y-chromosome haplotype shares similar challenges to the GREB1L situation.

Waples and Lindley (2018) pose a series of key questions to help provide an informed basis for decisions

or management actions.   Among other key questions, they argue that a full understanding of the

distribution of the variation in space and time is needed and that investigations into the genes and

mechanisms responsible for the life history variation should be initiated. In the case of the Y-haplotypes,

additional questions exist such as: whether the specific allelic variants at these SNP loci are important or

is the strong linkage disequilibrium among loci a signal from other adaptive genes in these regions; do

haplotype blocks exhibit consistent phenotypes in different environments and with different genetic

backgrounds; and if haplotype blocks are found in other regions of the Chinook salmon range do they

function in a similar manner to those suggested by the results of this study.

Conclusion

We identified Y-chromosome haplotypes that are associated with size, and likely age, at maturity

in Chinook salmon throughout Alaska. These haplotypes were restricted to northern Alaska Chinook

salmon where the most diversity in age-at-maturity exists, and likely represent a subset of the total

diversity across the species range. Interestingly, this region holds the most stable populations of Chinook

salmon in the Northeast Pacific (Griffiths et al. 2014), possibly due to the bet-hedging benefits of

diversity in age at maturity. It is possible that each Chinook salmon lineage has a specific set of

haplotypes and relationships between haplotypes and size/age at maturity may differ by lineage. The

discovery of Y-chromosome haplotypes and their potential effect on life history variation in Chinook

salmon may help understand the causes and consequences of the recent declines in size and age of adult

Chinook salmon, trends that are most pronounced in the region with the highest haplotype diversity.

Ongoing efforts to understand the causes of these declines point to a size-specific mortality filter on

maturing fish, but also requires an unknown evolutionary basis (Ohlberger et al. In Review). The results

presented here point to such a mechanism for the genetic control of changes in size-at-age and age-at-

maturity in Chinook salmon, as future changes in environmental conditions and selective fishing will lead

to further demographic responses in this economically and ecologically important species

Acknowledgements

We would like to thank Chris Habicht, Bill Templin, Wes Larson, and Todd Seamons for helpful

discussions and support for this research. This research was partially funded by North Pacific Research

Board grant 1724-000, AKSSF grants 44812, 44913 and 44914, and Pacific States Marine Fisheries

Commission grant # 16-108G to Alaska Department of Fish and Game. GJM was partially supported by

a National Research Council postdoctoral fellowship. The statements, findings, and conclusions are those

of the authors and do not necessarily reflect the views of the NOAA, the U.S. Department of Commerce,

or the ADFG.

Data Accessibility

329   GT-seq data is available in NCBI SRA XXXXXXXXXX.

Supporting Information

File S1: Supplementary materials, methods, and results for identification and characterization of Y-
chromosome haplotypes.

Table S1: Primer and probe sequence for GT-seq loci that were used for expanded genotyping of Y-
chromosome haplotypes.

Table S2.1: F_{ST}, F_{IS}, and allele frequencies for RAD loci associated with differentiation between clusters
in RADseq data.

Table S2.2: Genotypes for loci successfully developed into GT-seq. Genotypes are included for all
samples, RADseq and GT-seq.

Tables and Figures

Table 1. Populations used for Y-chromosome haplotype identification. RADseq samples were used for haplotype discovery while GT-seq samples were used to confirm that haplotypes were male-specific and to examine the geographic distribution of haplotypes. For populations with both RADseq and GT-seq samples, samples sizes for RADseq are given first. Frequency of Y-chromosome haplotypes are given for populations in Alaska; Y-chromosome haplotypes could not be assigned outside of Alaska.

Map Number	Population	Data Type	Region	Lat	Lon	N	AK Y1	AK Y2	AK Y3	AK Y4
1	Big Salmon River	RADseq	Yukon River	61.87	-134.92	49	0.00	0.00	0.79	0.21
2	Kantishna River	GT-seq	Yukon River	64.76	-149.97	48	0.00	0.00	1.00	0.00
3	Lower Yukon Test Fishery	GT-seq	Yukon River	62.79	-164.81	95	0.04	0.02	0.75	0.19
4	Anvik River	RADseq	Yukon River	62.68	-160.21	56	0.40	0.08	0.48	0.04
5	Tubutulik River	RADseq	Western Alaska	64.74	-161.89	56	0.39	0.48	0.10	0.03
6	Necons River	RADseq	Western Alaska	61.10	-153.85	47	0.11	0.05	0.21	0.63
7	George River	RADseq/GT-seq	Western Alaska	61.90	-157.71	48/47	0.16	0.05	0.38	0.41
8	Kogrukluk River	RADseq	Western Alaska	60.84	-157.85	59	0.58	0.21	0.13	0.08
9	Aniak River	RADseq	Western Alaska	61.06	-159.18	47	0.15	0.54	0.23	0.08
10	Kisaralik River	RADseq	Western Alaska	60.86	-161.24	48	0.64	0.27	0.09	0.00
11	Kwethluk River	RADseq	Western Alaska	60.81	-161.45	47	0.26	0.41	0.30	0.04
12	Kanektok River	RADseq/GT-seq	Western Alaska	59.75	-161.93	48/47	0.29	0.25	0.45	0.00
13	Arolik River	RADseq	Western Alaska	59.69	-161.88	48	0.33	0.11	0.48	0.07
14	Goodnews River	RADseq	Western Alaska	59.25	-161.36	47	0.14	0.10	0.43	0.33
15	Togiak River	RADseq	Western Alaska	59.09	-160.37	48	0.21	0.05	0.63	0.11
16	Iowithla River	RADseq	Western Alaska	59.18	-158.06	48	0.19	0.06	0.65	0.10
17	Stuyahok River	RADseq	Western Alaska	59.68	-156.17	48	0.23	0.19	0.35	0.23
18	Koktuli River	RADseq	Western Alaska	59.94	-156.43	56	0.29	0.21	0.38	0.12
19	Karluk River	GT-seq	Western Alaska	57.57	-154.38	48	0.00	0.00	1.00	0.00
20	Montana Creek	GT-seq	Southcentral Alaska	62.18	-149.95	48	0.23	0.00	0.31	0.46
21	Chuitna River	RADseq	Southcentral Alaska	61.20	-151.66	57	0.10	0.03	0.76	0.10
22	Sucker Creek	RADseq	Southcentral Alaska	61.51	-150.83	57	0.06	0.00	0.88	0.06
23	Talachulitna River	RADseq	Southcentral Alaska	61.62	-151.15	57	0.09	0.04	0.65	0.22
24	Theodore Creek	RADseq	Southcentral Alaska	61.49	-151.09	56	0.10	0.03	0.84	0.03
25	Coal Creek	RADseq	Southcentral Alaska	61.62	-151.76	55	0.00	0.00	0.67	0.33
26	Pullen Creek*	GT-seq	Southeast Alaska	59.45	-135.32	48	0.00	0.00	1.00	0.00
27	Little Port Walter-Unuk River stock**	GT-seq	Southeast Alaska	56.09	-131.06	48	0.00	0.00	0.00	1.00
-	Harrison River	GT-seq	British Columbia	49.28	-121.92	48	NA	NA	NA	NA
-	Big Qualicum Hatchery	GT-seq	British Columbia	49.40	-124.62	48	NA	NA	NA	NA
-	Kitwanga River	GT-seq	British Columbia	55.10	-128.09	48	NA	NA	NA	NA
-	Kitsumkalum River	GT-seq	British Columbia	54.52	-128.66	48	NA	NA	NA	NA
-	Morice River	GT-seq	British Columbia	54.41	-126.75	48	NA	NA	NA	NA
-	Rapid River Hatchery	GT-seq	Idaho	45.35	-116.40	48	NA	NA	NA	NA
-	McCall Fish Hatchery South Fork Salmon River	GT-seq	Idaho	44.67	-115.71	48	NA	NA	NA	NA
-	Marblemount Fish Hatchery	GT-seq	Washington	48.52	-121.42	48	NA	NA	NA	NA
-	Soos Creek Hatchery	GT-seq	Washington	47.31	-122.16	48	NA	NA	NA	NA
-	Columbia River at Wells Hatchery	GT-seq	Washington	47.95	-119.87	48	NA	NA	NA	NA
-	Quinault Lake Pens	GT-seq	Washington	47.47	-123.89	48	NA	NA	NA	NA
-	Lyons Ferry Hatchery	GT-seq	Washington	46.59	-118.22	48	NA	NA	NA	NA
-	Wenatchee River at Tumwater Dam	GT-seq	Washington	47.61	-120.72	48	NA	NA	NA	NA
-	Spring Creek Hatchery	GT-seq	Washington	45.73	-121.55	48	NA	NA	NA	NA
-	Rock Creek, Umpqua River	GT-seq	Oregon	43.34	-123.00	48	NA	NA	NA	NA
-	Cole River Hatchery, Rogue River	GT-seq	Oregon	42.66	-122.69	48	NA	NA	NA	NA
-	Cedar Creek Hatchery	GT-seq	Oregon	45.22	-123.84	48	NA	NA	NA	NA
-	McKenzie Hatchery	GT-seq	Oregon	44.11	-122.68	48	NA	NA	NA	NA
-	Coleman National Fish Hatchery	GT-seq	California	40.40	-122.14	48	NA	NA	NA	NA

*Only one male from Pullen Creek was assigned a Y-chromosome haplotype. **Only four out of 20 males from Little Port Walter Unuk River Stock were assigned a Y-chromosome haplotype.

Figure 1. Plot of genotype patterns for loci successfully developed into GT-seq assays for Y-
chromosome haplotype identification. The position of each marker along the Y-chromosome is shown by
vertical lines. The heatmap shows individual genotypes color-coded for homozygous (0), heterozygous
(1) and alternate homozygous (2). Samples are grouped by assigned haplotype on the Y-axis, with each
row representing an individual fish. This plot includes samples genotyped with RADseq and with GT-
seq. The AK Y1, AK Y2, and AK Y3 haplotypes share a common set of core markers up to ~6Mb but
vary in markers with extended linkage. The AK Y4 haplotype is fixed for alternate alleles for a unique
set of markers. Females exhibit two distinct genotype patterns for these SNPs, either nearly fixed
homozygosity for most markers or the presence of all three possible genotypes.

Figure 2. Frequency of Y-chromosome haplotypes throughout Alaska. Locations of populations are
approximate to prevent overlap of pie charts. Population names are given in Table 1. Note that the
location for the Lower Yukon Test Fishery (population 3) indicates where fish were caught on their return
to the Yukon River and that these samples may represent fish from throughout the Yukon River.

Figure 3. Distribution of size at maturity for each Y-chromosome haplotype for Alaska Chinook salmon.
Female (F) size at maturity is included for comparison. Samples sizes for each haplotype are given above
the boxplots. Within each region, Y-chromosome haplotypes with statistically different lengths are
represented by different letters. Haplotypes with two letters (i.e., ab) do not have statistically different
size distributions from haplotypes with a or b.

Figure 4. Distribution of A) age at maturity and B) size at age for each Y-chromosome haplotype in the
Yukon River. Within each age class, Y-chromosome haplotypes with statistically different lengths are
represented by different letters. Sample sizes for each haplotype are given above the boxplots.

References

Allendorf, F. W., and G. H. Thorgaard. 1984. Tetraploidy and the evolution of salmonid fishes. Pages 1-53 *in* B. Turner, editor. Evolutionary Genetics of Fishes. Plenum Publishing Corporation.

Aubin Horth, N., and J. J. Dodson. 2004. Influence of Individual Body Size and Variable Thresholds on the Incidence of a Sneaker Male Reproductive Tactic in Atlantic Salmon. Evolution **58**:136-144.

Ayllon, F., E. Kjaerner-Semb, T. Furmanek, V. Wennevik, M. F. Solberg, G. Dahle, G. L. Taranger, K. A. Glover, M. S. Almen, C. J. Rubin, R. B. Edvardsen, and A. Wargelius. 2015. The vgll3 Locus Controls Age at Maturity in Wild and Domesticated Atlantic Salmon (*Salmo salar L.*) Males. PLoS Genet **11**:e1005628.

Barson, N. J., T. Aykanat, K. Hindar, M. Baranski, G. H. Bolstad, P. Fiske, C. Jacq, A. J. Jensen, S. E. Johnston, S. Karlsson, M. Kent, T. Moen, E. Niemela, T. Nome, T. F. Naesje, P. Orell, A. Romakkaniemi, H. Saegrov, K. Urdal, J. Erkinaro, S. Lien, and C. R. Primmer. 2015. Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. Nature **528**:405-408.

Beacham, T. D., K. L. Jonsen, J. Supernault, M. Wetklo, L. Deng, and N. Varnavskaya. 2006. Pacific rim population structure of Chinook salmon as determined from microsatellite analysis. Transactions of the American Fisheries Society **135**:1604-1621.

Berejikian, B. A., D. M. Van Doornik, and J. J. Atkins. 2011. Alternative Male Reproductive Phenotypes Affect Offspring Growth Rates in Chinook Salmon. Transactions of the American Fisheries Society **140**:1206-1212.

Berejikian, B. A., D. M. Van Doornik, R. C. Endicott, T. L. Hoffnagle, E. P. Tezak, M. E. Moore, and J. Atkins. 2010. Mating success of alternative male phenotypes and evidence for frequency-dependent selection in Chinook salmon, Oncorhynchus tshawytscha. Canadian Journal of Fisheries and Aquatic Sciences **67**:1933-1941.

Boulding, E. G., K. P. Ang, J. A. K. Elliott, F. Powell, and L. R. Schaeffer. 2019. Differences in genetic architecture between continents at a major locus previously associated with sea age at sexual maturity in European Atlantic salmon. Aquaculture **500**:670-678.

Campbell, N. R., S. A. Harmon, and S. R. Narum. 2015. Genotyping-in-Thousands by sequencing (GT-seq): a cost effective SNP genotyping method based on custom amplicon sequencing. Mol Ecol Resour **15**:855-867.

Catchen, J., P. A. Hohenlohe, S. Bassham, A. Amores, and W. A. Cresko. 2013. Stacks: an analysis tool set for population genomics. Mol Ecol **22**:3124-3140.

Catchen, J. M., A. Amores, P. Hohenlohe, W. Cresko, and J. H. Postlethwait. 2011. Stacks: building and genotyping Loci de novo from short-read sequences. G3 (Bethesda) **1**:171-182.

Chapman, B. B., C. Brönmark, J.-Å. Nilsson, and L.-A. Hansson. 2011. The ecology and evolution of partial migration. Oikos **120**:1764-1775.

Christensen, K. A., J. S. Leong, D. Sakhrani, C. A. Biagi, D. R. Minkley, R. E. Withler, E. B. Rondeau, B. F. Koop, and R. H. Devlin. 2018. Chinook salmon (*Oncorhynchus tshawytscha*) genome and transcriptome. PLoS One **13**:e0195461.

Dann, T. H., C. Habicht, W. D. Templin, L. W. Seeb, G. J. McKinney, and J. S. Seeb. 2018. Identification of genetic markers useful for mixed stock analysis of Chinook salmon in Cook Inlet, Alaska. Alaska Department of Fish and Game, Division of Commercial Fisheries, Anchorage.

Davidson, W. S., T. K. Huang, K. Fujiki, K. R. von Schalburg, and B. F. Koop. 2009. The Sex Determining Loci and Sex Chromosomes in the Family Salmonidae. Sexual Development **3**:78-87.

DeFilippo, L. B., D. E. Schindler, J. Ohlberger, K. L. Schaberg, M. B. Foster, D. Ruhl, and A. E. Punt. 2019. Recruitment variation disrupts the stability of alternative life histories in an exploited salmon population. Evol Appl **12**:214-229.

Gall, G. A. E., J. Baltodano, and N. Huang. 1988. Heritability of age at spawning for rainbow trout. Aquaculture **68**:93-120.

Garant, D., P.-M. Fontaine, S. P. Good, J. J. Dodson, and L. Bernatchez. 2002. The influence of male
parental identity on growth and survival of offspring in Atlantic salmon (*Salmo salar*).
Evolutionary Ecology Research **4**:537-549.
Gibbons, J. W., R. D. Semlitsch, J. L. Greene, and J. P. Schubauer. 1981. Variation in Age and Size at
Maturity of the Slider Turtle (*Pseudemys scripta*). The American Naturalist **117**:841-845.
Gjerde, B. 1984. Response to individual selection for age at sexual maturity in Atlantic salmon.
Aquaculture **38**:229-240.
Griffiths, J. R., D. E. Schindler, J. B. Armstrong, M. D. Scheuerell, D. C. Whited, R. A. Clark, R.
Hilborn, C. A. Holt, S. T. Lindley, J. A. Stanford, and E. C. Volk. 2014. Performance of salmon
fishery portfolios across western North America. Journal of Applied Ecology **51**:1554-1563.
Hall, M. C., C. J. Basten, and J. H. Willis. 2006. Pleiotropic quantitative trait loci contribute to population
divergence in traits associated with life-history variation in Mimulus guttatus. Genetics **172**:1829-
1844.
Hankin, D. G., J. W. Nicholas, and T. W. Downey. 1993. Evidence for inheritance of age of maturity in
Chinook salmon (*Oncorhynchus tshawytscha*). Canadian Journal of Fisheries and Aquatic
Sciences **50**:347-358.
Hard, J. J., M. R. Gross, M. Heino, R. Hilborn, R. G. Kope, R. Law, and J. D. Reynolds. 2008.
Evolutionary consequences of fishing and their implications for salmon. Evolutionary
Applications **1**:388-408.
Harstad, D. L., D. A. Larsen, J. Miller, I. Adams, D. K. Spangenberg, S. Nance, L. Rohrbach, J. G.
Murauskas, and B. R. Beckman. 2018. Winter-Rearing Temperature Affects Growth Profiles,
Age of Maturation, and Smolt-to-Adult Returns for Yearling Summer Chinook Salmon in the
Upper Columbia River Basin. North American Journal of Fisheries Management **38**:867-885.
Healey, M. C. 1986. Optimum size and age at maturity in pacific salmon and effects of size-selective
fisheries. Can. Pec. Publ. Fish. Aquat. Sci. **89**:39-52.
Healey, M. C. 1991. Life history of chinook salmon (*Oncorhynchus tshawytscha*). Pages 311-393 *in* L.
Margolis and C. Groot, editors. Pacific salmon life histories. UBC Press, Vancouver, B.C.
Healey, M. C., and W. R. Heard. 1984. Inter- and Intra-Population Variation in the Fecundity of Chinook
Salmon (Oncorhynchus tshawytscha) and its Relevance to Life History Theory. Canadian Journal
of Fisheries and Aquatic Sciences **41**:476-483.
Heath, D. D., R. H. Devlin, J. W. Heath, and G. K. Iwama. 1994. Genetic, environmental and interaction
effects on the incidence of jacking in *Oncorhynchus tshawytscha* (Chinook salmon). Heredity
(Edinb) **72**:146-154.
Heath, D. D., L. Rankin, C. A. Bryden, J. W. Heath, and J. M. Shrimpton. 2002. Heritability and Y-
chromosome influence in the jack male life history of chinook salmon (*Oncorhynchus
tshawytscha*). Heredity (Edinb) **89**:311.
Hess, J. E., N. R. Campbell, A. P. Matala, D. J. Hasselman, and S. P. Narum. 2016a. Genetic assessment
of Columbia River stocks, 4/1/2014-3/31/2105 annual report, 2008-907-00. CRITFC. Available
from http://www.critfc.org/wp-content/uploads/2016/04/16-03.pdf.
Hess, J. E., J. S. Zendt, A. R. Matala, and S. R. Narum. 2016b. Genetic basis of adult migration timing in
anadromous steelhead discovered through multivariate association testing. Proc Biol Sci **283**.
Johnston, S. E., J. Gratten, C. Berenos, J. G. Pilkington, T. H. Clutton-Brock, J. M. Pemberton, and J.
Slate. 2013. Life history trade-offs at a single locus maintain sexually selected genetic variation.
Nature **502**:93-95.
Kendall, N. W., U. Dieckmann, M. Heino, A. E. Punt, and T. P. Quinn. 2014. Evolution of age and length
at maturation of Alaskan salmon under size-selective harvest. Evolutionary Applications **7**:313-
322.
Kijas, J., S. McWilliam, M. Naval Sanchez, P. Kube, H. King, B. Evans, T. Nome, S. Lien, and K.
Verbyla. 2018. Evolution of Sex Determination Loci in Atlantic Salmon. Scientific Reports
8:5664.

Kodama, M., J. J. Hard, and K. A. Naish. 2018. Mapping of quantitative trait loci for temporal growth
and age at maturity in coho salmon: Evidence for genotype-by-sex interactions. Marine Genomics
38:33-44.
Koo, T. S. Y. 1962. Age determination in salmon. Pages 37-48 *in* T. S. Y. Koo, editor. Studies of Alaska
red salmon. University of Washington Press,, Seattle.
Küpper, C., M. Stocks, J. E. Risse, N. dos Remedios, L. L. Farrell, S. B. McRae, T. C. Morgan, N.
Karlionova, P. Pinchuk, Y. I. Verkuil, A. S. Kitaysky, J. C. Wingfield, T. Piersma, K. Zeng, J.
Slate, M. Blaxter, D. B. Lank, and T. Burke. 2015. A supergene determines highly divergent male
reproductive morphs in the ruff. Nature Genetics **48**:79.
Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods **9**:357-
359.
Larsen, D. A., B. R. Beckman, C. R. Strom, P. J. Parkins, K. A. Cooper, D. E. Fast, and W. W. Dickhoff.
2006. Growth Modulation Alters the Incidence of Early Male Maturation and Physiological
Development of Hatchery-Reared Spring Chinook Salmon: A Comparison with Wild Fish.
Transactions of the American Fisheries Society **135**:1017-1032.
Larson, W. A., L. W. Seeb, M. V. Everett, R. K. Waples, W. D. Templin, and J. E. Seeb. 2014.
Genotyping by sequencing resolves shallow population structure to inform conservation of
Chinook salmon (*Oncorhynchus tshawytscha*). Evol Appl **7**:355-369.
Lewis, B., W. S. Grant, R. E. Brenner, and T. Hamazaki. 2015. Changes in Size and Age of Chinook
Salmon *Oncorhynchus tshawytscha* Returning to Alaska. PLoS One **10**:e0130184-e0130184.
Lien, S., L. Gidskehaug, T. Moen, B. J. Hayes, P. R. Berg, W. S. Davidson, S. W. Omholt, and M. P.
Kent. 2011. A dense SNP-based linkage map for Atlantic salmon (Salmo salar) reveals extended
chromosome homeologies and striking differences in sex-specific recombination patterns. BMC
Genomics **12**:615.
Lozorie, J. D., and B. C. McIntosh. 2014. Sonar esimation of salmon passage in the Yukon River near
Pilot Station, 2012. Alaska Department of Fish and Game.
McKinney, G. J., C. E. Pascal, W. D. Templine, S. E. Gilk-Baumer, T. H. Dann, L. W. Seeb, and J. E.
Seeb. In Revision. Dense SNP panels resolve closely related Chinook salmon populations. In
Revision.
McKinney, G. J., R. K. Waples, C. E. Pascal, L. W. Seeb, and J. E. Seeb. 2018. Resolving allele dosage in
duplicated loci using genotyping-by-sequencing data: A path forward for population genetic
analysis. Mol Ecol Resour **18**:570-579.
McKinney, G. J., R. K. Waples, L. W. Seeb, and J. E. Seeb. 2017. Paralogs are revealed by proportion of
heterozygotes and deviations in read ratios in genotyping-by-sequencing data from natural
populations. Mol Ecol Resour **17**:656-669.
Micheletti, S. J., and S. R. Narum. 2018. Utility of pooled sequencing for association mapping in
nonmodel organisms. Mol Ecol Resour **18**:825-837.
Moran, P., D. J. Teel, M. A. Banks, T. D. Beacham, M. R. Bellinger, S. M. Blankenship, J. R. Candy, J.
C. Garza, J. E. Hess, S. R. Narum, L. W. Seeb, W. D. Templin, C. G. Wallace, and C. T. Smith.
2012. Divergent life-history races do not represent Chinook salmon coast-wide: the importance of
scale in Quaternary biogeography. Canadian Journal of Fisheries and Aquatic Sciences **70**:415-
435.
Ohlberger, J., D. E. Schindler, E. J. Ward, T. E. Walsworth, and T. E. Essington. In Review. Resurgence
of an apex marine predator and the decline in prey body size.
Ohlberger, J., E. J. Ward, D. E. Schindler, and B. Lewis. 2018. Demographic changes in Chinook salmon
across the Northeast Pacific Ocean. Fish and Fisheries **19**:533-546.
Otero, J., A. J. Jensen, J. H. L'Abee-Lund, N. C. Stenseth, G. O. Storvik, and L. A. Vollestad. 2012.
Contemporary ocean warming and freshwater conditions are related to later sea age at maturity in
Atlantic salmon spawning in Norwegian rivers. Ecol Evol **2**:2192-2203.

Pearse, D. E., M. R. Miller, A. Abadia-Cardoso, and J. C. Garza. 2014. Rapid parallel evolution of
standing variation in a single, complex, genomic region is associated with life history in
steelhead/rainbow trout. Proc Biol Sci **281**:20140012.
Phillips, R. B., L. K. Park, and K. A. Naish. 2013. Assignment of Chinook salmon (Oncorhynchus
tshawytscha) linkage groups to specific chromosomes reveals a karyotype with multiple
rearrangements of the chromosome arms of rainbow trout (Oncorhynchus mykiss). G3 (Bethesda,
Md.) **3**:2289-2295.
Prince, D. J., S. M. O'Rourke, T. Q. Thompson, O. A. Ali, H. S. Lyman, I. K. Saglam, T. J. Hotaling, A.
P. Spidle, and M. R. Miller. 2017. The evolutionary basis of premature migration in Pacific
salmon highlights the utility of genomics for informing conservation. Sci Adv **3**:e1603198.
Riddell, B. R., R. D. Brodeur, A. V. Bugaev, P. Moran, J. M. Murphy, J. A. Orsi, M. trudel, L. A.
Weitkamp, B. K. Wells, and A. C. Wertheimer. 2018. Ocean ecology of Chinook Salmon. Pages
55-696 *in* R. J. Beamish, editor. The ocean ecology of Pacific salmon and trout. American
Fisheries Society, Bethesda, Maryland.
Rousset, F. 2008. Genepop'007: A complete re-implementation of the Genepop software for Windows
and Linux. Mol Ecol Resour **8**:103-106.
Rowe, D., and J. E. Thorpe. 1990. Suppression of maturation in male Atlantic salmon parr (*Salmon salar*
L.) by reduction in feeding and growth during spring months. Aquaculture **86**:291-313.
Schoen, E. R., M. S. Wipfli, E. J. Trammell, D. J. Rinella, A. L. Floyd, J. Grunblatt, M. D. McCarthy, B.
E. Meyer, J. M. Morton, J. E. Powell, A. Prakash, M. N. Reimer, S. L. Stuefer, H. Toniolo, B. M.
Wells, and F. D. W. Witmer. 2017. Future of Pacific Salmon in the Face of Environmental
Change: Lessons from One of the World's Remaining Productive Salmon Regions. Fisheries
42:538-553.
Shuster, S. M. 1989. Male alternative reproductive strategies in a marine isopod crustacean (*paracerceis*
sculpta): The use of genetic markers to measure differences in fertilization success among alpha-,
beta-, and gamma-males. Evolution **43**:1683-1698.
Siegel, J. E., M. V. McPhee, and M. D. Adkison. 2017. Evidence that marine temperatures influence
growth and maturation of Western Alaskan Chinook salmon. Marine and Coastal Fisheries **9**:441-
456.
Templin, W. D., J. E. Seeb, J. R. Jasper, A. W. Barclay, and L. W. Seeb. 2011. Genetic differentiation of
Alaska Chinook salmon: the missing link for migratory studies. Mol Ecol Resour **11 Suppl**
1:226-246.
Thorpe, J. E. 2007. Maturation responses of salmonids to changing developmental opportunities. Marine
Ecology Progress Series **335**:285-288.
Twyford, A. D., and J. Friedman. 2015. Adaptive divergence in the monkey flower Mimulus guttatus is
maintained by a chromosomal inversion. Evolution **69**:1476-1486.
Waples, R. S., and S. T. Lindley. 2018. Genomics and conservation units: The genetic basis of adult
migration timing in Pacific salmonids. Evolutionary Applications **11**:1518-1526.
Waples, R. S., D. J. Teel, J. M. Myers, and A. R. Marshall. 2004. Life-history divergence in Chinook
salmon: Historic contingency and parallele evolution. Evolution **58**:386-403.
Waters, C. D., J. J. Hard, M. S. O. Brieuc, D. E. Fast, K. I. Warheit, C. M. Knudsen, W. J. Bosch, and K.
A. Naish. 2018. Genomewide association analyses of fitness traits in captive-reared Chinook
salmon: Applications in evaluating conservation strategies. Evol Appl **11**:853-868.
Yano, A., B. Nicol, E. Jouanno, E. Quillet, A. Fostier, R. Guyomard, and Y. Guiguen. 2013. The sexually
dimorphic on the Y-chromosome gene (sdY) is a conserved male-specific Y-chromosome
sequence in many salmonids. Evolutionary applications **6**:486-496.
You, F. M., N. Huo, Y. Q. Gu, M. C. Luo, Y. Ma, D. Hane, G. R. Lazo, J. Dvorak, and O. D. Anderson.
2008. BatchPrimer3: a high throughput web application for PCR and sequencing primer design.
BMC Bioinformatics **9**:253.

Young, B., D. V. Conti, and M. D. Dean. 2013. Sneaker "jack" males outcompete dominant "hooknose"
males under sperm competition in Chinook salmon (*Oncorhynchus tshawytscha*). Ecology and
Evolution **3**:4987-4997.