Aquatic biodiversity enhances multiple nutritional benefits to humans

Joey R. Bernhardt[1,2]*, Mary I. O'Connor [1]

[1]Department of Zoology and Biodiversity Research Centre, University of British Columbia,
Vancouver, Canada.

[2]Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT, USA.

*Correspondence to: joey.bernhardt@biodiversity.ubc.ca.

ORCIDs:

Joey R. Bernhardt: 0000-0003-1824-2801

Mary I. O'Connor: 0000-0001-9583-1592

Classification: Biological Sciences/Ecology; Sustainability Science

Keywords: seafood, biodiversity-ecosystem functioning, ecosystem services

Author contributions: JRB conceived the study, collected and curated the data. JRB developed
software, analyzed and visualized the data with input from MIO. JRB wrote the original draft of
the manuscript, JRB and MIO contributed to writing and editing. JRB and MIO supervised the
study.

22
23 **This PDF file includes:**
24 Main Text
25 Figures 1 to 6

Abstract

Humanity depends on biodiversity for health, well-being and a stable environment. As biodiversity change accelerates, we are still discovering the full range of consequences for human health and well-being. Here, we test the hypothesis -- derived from biodiversity - ecosystem functioning theory -- that species richness and ecological functional diversity allow seafood diets to fulfill multiple nutritional requirements, a condition necessary for human health. We analyzed a newly synthesized dataset of 7245 observations of nutrient and contaminant concentrations in 801 aquatic animal taxa, and found that species with different ecological traits have distinct and complementary micronutrient profiles, but little difference in protein content. The same complementarity mechanisms that generate positive biodiversity effects on ecosystem functioning in terrestrial ecosystems also operate in seafood assemblages, allowing more diverse diets to yield increased nutritional benefits independent of total biomass consumed. Notably, nutritional metrics that capture multiple micronutrients essential for human well-being depend more strongly on biodiversity than common ecological measures of function such as productivity, typically reported for grasslands and forests. Further, we found that increasing species richness did not increase the amount of protein in seafood diets, and also increased concentrations of toxic metal contaminants in the diet. Seafood-derived micronutrients are important for human health and are a pillar of global food and nutrition security. By drawing upon biodiversity-ecosystem functioning theory, we demonstrate that ecological concepts of biodiversity can deepen our understanding of nature's benefits to people and unite sustainability goals for biodiversity and human well-being.

Significance statement:

Food security is not simply about maintaining yields, it is also about the need for a stable supply

of nutritionally diverse foods. Obtaining nutritious food is a major challenge facing humanity

and diverse aquatic ecosystems can help meet this goal. To test how aquatic biodiversity affects

human health, we assembled a new dataset of nutrients, contaminants and ecological traits of 801

aquatic species. We used ecological models to quantify the role of species richness and

ecological functional diversity and found that these biodiversity dimensions enhanced seafood

micronutrient provisioning by the same mechanisms that link biodiversity to productivity in

grasslands, forests and other systems. Our results underscore the need to minimize aquatic

biodiversity loss to sustain and improve human well-being.

Introduction

Species losses and range shifts due to climate change, harvesting, and other human activities are altering aquatic biodiversity locally and globally (1–5). In aquatic ecosystems, not only are some species severely depleted because of overfishing or habitat loss (3, 6–8), the ecosystem-level dimensions of biodiversity such as the total number of species and their functional diversity have also changed (9). Beyond the loss of particular species, changes in ecosystem-level dimensions of biodiversity threaten numerous ecosystem services to humans, which include the cultural, economic or health benefits people derive from nature (10–13). In many regions, such as tropical coastal systems, the cumulative impacts of human activities are severe and associated with strong declines in taxonomic and ecological functional diversity (6) and coincide with regions with high dependence of people upon wild-caught seafood for food and nutrition (14). In temperate regions, where some coastal communities depend on local wild seafood harvests to meet their nutritional needs (15, 16), species richness may be increasing as species recover from exploitation and warmer oceans allow species to expand their ranges into new territory (1, 2, 17).

There is growing concern that biodiversity change leads to changes in human health and well-being (10, 13, 18). Specific and quantitative links between aquatic biodiversity and human health that distinguish contributions of species diversity from those of biomass, as predicted by biodiversity-ecosystem functioning theory, have not been established. At a time of unprecedented global change and increasing reliance on seafood to meet nutritional demands (19), there is an urgent need to understand how changing aquatic ecosystem structure may alter the provisioning of seafood-derived human nutrition.

Seafood, consisting of wild-caught marine and freshwater finfish and invertebrates, provides an

important source of protein and calories to humans. Additionally, unlike staple foods such as rice

or other grains, seafood can address multiple dimensions of food and nutritional security

simultaneously by providing essential micronutrients, such as vitamins, minerals and

polyunsaturated essential fatty acids critical to human health (19–22). Given the multiple

attributes of seafood valuable to human health, it is possible that the diversity of an aquatic

assemblage, distinct from the inclusion of any particularly nutritious species, could support

human well-being consistent with a large body of evidence for biodiversity's major contributions

to ecological functions (11, 23–26). Dietary diversity is a basic tenet of a nutritious diet (27) and

it is widely appreciated that diets composed of more food groups and more species are more

nutritious (28–31). Ecological measures of dietary diversity (diet diversity, species richness,

functional diversity and Simpson's index of evenness) have been associated with nutritional

value of diets in a range of contexts (27, 29, 32–38). These studies rely on relationships between

species included in the diet (or other food intake measures) and nutritional adequacy of reported

diets. However, a simple correlation between dietary diversity and a measure of dietary benefits

provides only partial support for a claim that biodiversity benefits human well-being consistent

with the same ecological processes by which biodiversity supports numerous ecosystem

functions and services (23, 26). We build upon this foundation of empirical relationships

between diet diversity and diet quality by placing this question in the quantitative ecological

theoretical framework that relates biodiversity to function (24, 25), thereby laying the

groundwork for additional development of links between biodiversity science and our

understanding of human well-being.

Ecological theory predicts that biodiversity can be ecologically and economically important,

apart from the importance of total biomass or the presence of particular species (23, 39).

According to theory and over 500 explicit experimental tests (23, 40, 41) diversity in ecological

communities and agricultural systems enhances ecosystem functioning by two mechanisms: (i)

more diverse assemblages may outperform less diverse assemblages *of the same density or*

biomass of individuals because more diverse assemblages will include more of the possible

species and are therefore more likely to include high-performing species, assuming random

processes of including species from the species pool (a *selection effect*), or (ii) more diverse

assemblages of a given density (or biomass) contain species with complementary functional

traits, allowing them to function more efficiently (a *complementarity effect*) (25, 39). For aquatic

animals, increased diversity enhances productivity of fish biomass (42) and also enhances

temporal stability of biomass production and total yields (43, 44), providing economic and

nutritional benefits to humans related to increased stability of harvests and production of biomass

for consumption (43). However, when considering aquatic species from the perspective of

human nutrition, functions other than biomass production become relevant because total seafood

biomass consumption is not predictive of micronutrient benefits from seafood (45, 46).

Here we test a hypothesis central to ecological theory in the 21st century: whether biodiversity

per se (species richness and ecological functional diversity), distinct from the identities and

abundance of species, enhances human well-being (**Figure 1**). We chose a measure of human

well-being distinct from provision of protein, calories or total yields - the micronutrient benefits

of seafood. For increasing biodiversity *per se* (as opposed to increasing total seafood

consumption) to enhance nutritional benefits as predicted by biodiversity-ecosystem functioning

theory (25, 47), the amounts of various micronutrients within edible tissues must differ among

species and, furthermore, micronutrient concentrations must trade off among species, such that

species that have relatively high concentrations of some nutrients also have relatively low

concentrations of others (25). Specifically, a 'biodiversity effect' (*sensu (25)*) on nutritional

benefits requires that concentrations of multiple nutrients are negatively correlated with each

other, or uncorrelated, when compared among species, creating a complementary distribution of

micronutrients across species. In contrast, if micronutrient concentrations in edible tissue are

positively correlated for multiple micronutrients across species such that, for example, a species

containing high amounts of iron also has a high essential fatty acid concentration, thereby

containing multiple nutrients in high concentrations simultaneously, seafood species or

ecological functional *diversity* in the diet would not be important. In the case of positive

correlations among micronutrient concentrations, the ecosystem service of nutritional benefits

would be enhanced by consuming more fish biomass or by selecting a few highly nutritious

species, without considering species richness or ecological functional diversity.

Figure 1. Aquatic biodiversity increases human well-being because edible species have distinct

and complementary multi-nutrient profiles (**A**) and differ in mean micro- and macronutrient

content (shown here relative to 10% and 25% thresholds of recommended dietary allowance

(RDA) guidelines) for representative finfish (*Abramis brama*, *Mullus surmuletus*), mollusc

(*Mytilus galloprovincialis*) and crustacean species (*Nephrops norvegicus*). Biodiversity –

ecosystem functioning theory predicts that nutritional benefits, including the number of nutrient

RDA targets met per 100g portion (*NT*; i, iii) and minimum portion size (P_{min}; ii, iv) (**B, E**), are

enhanced with increasing seafood species richness. Orange dots in panels B and E correspond to

potential diets high and low biodiversity levels. Seafood consumers with limited access to

seafood each day may not reach RDA targets if diets are low in diversity (**F** vs **C**; grey shading

indicates proportion of population that meets nutrient requirements). DHA: docosahexaenoic

acid, EPA: eicosapentaenoic acid.

We aimed to bridge two distinct theoretical frameworks - the biodiversity-ecosystem functioning

theory and human nutrition science - by quantitatively testing for effects of aquatic species

richness and ecological functional diversity (48, 49) in seafood diets on nutritional benefits via

complementarity or selection effects. We used the public health measure of recommended

dietary allowance (RDA) index to quantify nutritional benefits. RDAs are nutrient-based

reference values that indicate the average daily dietary intake level that is sufficient to meet the

nutrient requirement of nearly all (97 to 98 percent) healthy individuals in a particular life stage

and gender group (50). Here we used the RDA for females aged 19-50 (SI Appendix, Tables S1,

S2). We measured nutritional value in terms of concentrations relative to RDAs, and we refer to

these recommended amounts (or portions thereof) as 'RDA targets' (SI Appendix, Table S1,

Table S2, Methods 1.1). We quantified nutritional value in two ways: 1) the minimum amount of

seafood tissue (in grams, g) required to meet given RDA targets (either for a single nutrient, or

five micronutrients simultaneously; referred to as 'minimum portion size required', P_{min} (SI

Appendix, Table S1, Equation A1, Methods 1.1) and 2) the number of nutrients that meet an

RDA target in a single 100g seafood portion (NT, Table S1, Equation A2). By considering

nutritional value per unit biomass in both metrics, we avoided confounding diversity of seafood

consumed with the total amount consumed (Methods 1.3). We first tested two hypotheses: (i)

seafood species richness increases NT due to complementarity in nutrient concentrations among

species and (ii) seafood species richness increases the nutritional value of a 100g edible portion

of seafood, thereby lowering the minimum portion size, P_{min}, and improving the efficiency with

which seafood consumers reach nutritional targets (**Figure 1**). Following biodiversity-ecosystem

functioning theory, we predicted that increased species richness is correlated with ecological

functional diversity (51) in potential seafood diets, and that ecological functional diversity is

related to diversity in the concentration of essential elements and fatty acids that have nutritional

value to human consumers, such that species and ecological functional diversity yields increased

nutritional benefits. We also tested the hypothesis (iii) that seafood diversity increases total

intake of heavy metal contaminants, because some aquatic animals are known to bioaccumulate

toxic metals in their tissues. For this reason, variation in bioaccumulation among species could

lead to a biodiversity effect on contaminant intake that is detrimental to human health.

In a global analysis of over 5040 observations of nutrient concentrations in 547 aquatic species

(SI Appendix, **Figure S1**), we considered provision of nutritional benefits to human consumers.

To assess whether the relationships between biodiversity and human nutrition benefits depend on

the geographic extent (global, local) over which seafood are harvested or accessed (11), we

tested whether seafood species richness is associated with higher nutritional value at local scales

(*vs* global scale) in traditional indigenous seafood diets in North America (SI Appendix, Methods

1.4). Seafood is critical for indigenous groups, who on average consume seafood at a rate that is

15 times higher than the global average *per capita* consumption rate (16). To test our hypotheses

at the geographic scale of local consumer communities, we complemented our global analysis

with additional analyses of 25 – 57 species in fourteen geographically constrained groups of

species consumed together as part of traditional indigenous diets (SI Appendix, Methods 1.4).

Results

Diversity of seafood nutrient concentrations

Biodiversity effects via complementarity or selection require that species differ in their

functional traits. We found that the global species pool was highly diverse with regard to

concentrations of the micronutrients iron, zinc, calcium, and two fatty acids, docosahexaenoic

acid (DHA) and eicosapentaenoic acid (EPA), in edible fish tissue relative to RDAs for those

micronutrients (**Figure 2**; micronutrient geometric coefficients of variation (CV): ln(iron) = 3.28,

ln(calcium) = 3.56, ln(EPA) = 2.62, ln(zinc) = 3.02, ln(DHA) = 2.17; note the log scale). We

observed limited variation in protein concentrations (ln(protein) CV = 0.04). The frequency

distribution of trait values, such as nutrient concentrations, across species may indicate the

potential strength of biodiversity effects, with lognormal distributions (such as we observed for

micronutrients) more likely to confer strong effects of biodiversity than normal distributions with

low dispersion (as we observed for protein). Most species did not meet a single micronutrient

RDA in a 100g portion: fewer than half of the 547 species we examined reached an RDA target

of 10% RDA for calcium, iron and the essential fatty acid EPA in a standard 100g edible portion

of a single species (SI Appendix, Table S3).

Figure 2. Variation in nutrient concentrations differs among taxonomic groups. (**A**) Frequency of reported protein, fat and micronutrient (including eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA)) content in 100 g of the edible portion of 547 seafood species.

Note the x-axis is plotted on a log scale. (**B**) Proportion of species, and number shown on each

bar, with available data that reach 10% of RDA targets for any one, two or up to five of the

micronutrients examined here.

Biodiversity increased the nutritional content of an edible portion of seafood

We found that seafood species richness not only enhanced nutritional value for consumers

selecting seafood from our global species dataset, but that seafood species richness *per se* was

essential to meeting nutritional targets for seafood diets with limited biomass consumption.

Increasing seafood species richness allowed simulated diets to reach more RDA targets per 100g

of tissue, so that nutritional value increased with species richness even as total biomass

consumption (e.g., total seafood portion size) remained constant. We quantified the minimum

amount of seafood, in grams, that would be required to reach an RDA target (SI Appendix, Table

S2) at each of ten levels of species richness (referred to as minimum portion size, P_{min}, for which

lower values signify higher nutrition benefits to consumers per gram seafood consumed,

Methods 1.3). We then estimated the biodiversity effect using Equation 3 (Methods 2.3), in

which b_{Pmin} is the scaling coefficient that describes how function (here, P_{min}) varies with species

richness; higher absolute values of b_f (where f is an ecosystem function) indicate a steeper

relationship between biodiversity and function, and can be used to compare 'biodiversity effects'

among studies and systems (12, 52). As species richness increased in potential diets, P_{min}

declined, and RDA targets for each micronutrient were achieved with less total seafood intake

(**Figure 3A**, $b_{Pmin} < 0$ for every micronutrient: calcium -0.32 (95% CI -0.35, -0.28), iron -0.24

(95% CI -0.27, -0.22), zinc -0.26 (95% CI -0.28, -0.23), EPA -0.25 (95% CI -0.27, -0.23) and

DHA -0.22 (95% CI -0.23, -0.21)). Increasing species richness reduced the minimum portion

size required, P_{min}, in our sample diets, independent of systematic changes in the identity of

species included (see Methods 2.2), because the diets were assembled using random samples of

the species pool. The restricted variation in protein concentrations (**Figure 2A**), combined with

high levels of protein in all edible tissues, lead to no benefit, and a minimal detrimental effect of

seafood species richness on protein provisioning (**Figure 3A***, b_{Pmin} = 0.0071 95% CI 0.0062,

0.0080). In other words, the ecosystem service of protein provisioning was adequately provided

by total seafood edible biomass, and not improved by species richness or even species identity.

These findings are consistent with demonstrations that variety and diversity in diets is important

for nutrition (28, 29), but extend these findings to show seafood species richness allows

consumers to gain more nutritional benefit without consuming more total seafood biomass, and

explicitly relates this pattern with general effects of biodiversity in ecological systems.

Figure 3. Aquatic biodiversity enhances nutritional benefits. (**A**) Seafood species richness

improves the efficiency with which human diets can meet RDA targets by reducing the minimum

portion size required, P_{min}, to meet RDA targets (measured in grams of seafood). P_{min} is shown

for five micronutrients and protein separately (points are median values for calcium, iron, zinc,

EPA, DHA and protein, lines show the fit of Equation 3 to the data and shading refers to 95%

CI) as well as for five micronutrients simultaneously (top purple line). Colours corresponding to

each nutrient in panel A are shown in panel B. (**B**) Estimates (± 95% CI) for the scaling

parameter that relates species richness to P_{min} (b_{Pmin}) (Equation 3). (**C**) Species richness increases

the number of distinct nutrient RDA targets met in a 100g seafood portion (*NT*); black line and

95% confidence interval correspond to the fit of Equation 2 to mean *NT* derived from resampled

diets from the global seafood species pool. Flower plots in **C** summarize micronutrient

concentrations relative to 10% RDA (grey circle) in two representative diets at low and high

species richness levels. Data shown in panels A and C are derived from $n = 1000$ resampled

diets.

We then considered the effects of seafood species richness on the provisioning of multiple

nutrients simultaneously (Methods 1.3). This is referred to as a multifunctional benefit of

biodiversity (53, 54), and takes into account possible trade-offs or correlations among functions,

in this case, concentrations of micronutrients. For some ecosystem services (e.g. water quality or

ecotourism), benefits of biodiversity accumulate when multiple ecosystem functions are

considered simultaneously (54–56). Consistent with biodiversity-ecosystem functioning theory,

we found that in the case of a multifunctional metric of an ecosystem service defined from the

human beneficiary's perspective (i.e. multiple micronutrient targets reached simultaneously),

biodiversity benefits for the multifunctional service are greater than for individual functions

(b_{Pmin} for all five micronutrients simultaneously = -0.42 (95% CI -0.47, -0.38) vs. single nutrients

b_{Pmin} range from -0.32 (95% CI -0.35, -0.28) for calcium to -0.22 for DHA (95% CI -0.23, -

0.21)) that comprise the ecosystem service (**Figure 3A, B**). Increasing seafood species richness

from one to ten species in 1000 simulated, resampled diets drawn from our global species pool

allowed diets to meet RDA targets for five essential microelements and fatty acids

simultaneously more than twice as efficiently (i.e. a median of 494.19 g of tissue required with

one species vs. median of 213.34 g of tissue required with ten species) (**Figure 3A, B**). Then, we

assessed the effects of biodiversity when the total biomass of seafood consumed was held

constant, at 100g, by counting the number of nutrients for which RDA targets were reached in a

100g portion (*NT*). We found positive effects of biodiversity on the number of nutrients that met

RDA targets, *NT*, in a single 100g portion (**Figure 3C**): diets with higher species richness

reached more nutritional targets (higher *NT*) per 100g serving than diets of the same fish biomass

comprising fewer species (b_{NT} = 0.20 (95% CI: 0.20-0.21) **Figure 3C**). These findings were

robust to different RDA target levels (for P_{min}, they were independent of RDA target level, and

for *NT* they were positive over the range from 1% RDA to 40% RDA per 100g portion, SI

Appendix, **Figure S2**). These results reveal biodiversity effects of seafood quantitatively

comparable with the widely recognized relationship between biodiversity and productivity (26,

47, 52), and demonstrate a benefit of biodiversity for human nutritional well-being over and

above the benefits of consuming a particular amount (biomass) or identity of aquatic species.

Increasing seafood diversity increased contaminant exposure

We considered a range of trace elements, which, at high concentrations are known to be harmful

to human health (57). We focused on four heavy metals considered as contaminants

(methylmercury, cadmium, arsenic and lead) for which there exist public health guidelines for

upper tolerable limits and for which seafood is potentially a major source of dietary intake (SI

Appendix**, Figure S1, Table S2**). We examined the concentrations of these elements in the

muscle tissues of 353 seafood species (thereby excluding parts such as viscera, liver and bones).

We found that the same mechanisms that lead to a positive relationship between seafood species

richness and nutritional benefits also contributed to exposure to a wider range of contaminants

with higher species richness. We observed high levels of variation in contaminant concentrations

across species (**Figure 4A-D**), and as we observed for micronutrients, these distributions were

often right-skewed such that most species contained low contaminant concentrations and few

contained high concentrations. On average, muscle tissue concentrations of contaminants were

weakly positively correlated across species, such that species that contained high concentrations

of one contaminant also contained high concentrations of another (average pairwise correlation =

0.09) (SI Appendix, **Figure S3**). When considering multiple contaminants simultaneously,

increasing seafood species richness increased the number of contaminants which exceeded their

upper tolerable limits (PTDI) in a 100g portion, referred to as *NC* (SI Appendix, Table S1

Equation A4), (b_{NC} = 0.10, 95% CI 0.084, 0.12) (**Figure 4E**). When we considered the effects of

biodiversity on exposure to each contaminant separately, we found that increasing species

richness generally increased contaminant content per 100g, but the strength of this effect varied

among contaminants. For example, increasing species richness from one to ten species was

associated with doubling methylmercury concentrations on average, thereby reducing the

maximum portion size before exceeding PTDI, referred to as P_{max}, (SI Appendix, Table S1

Equation A3) (b_{Pmax} = -0.25 95% CI -0.27, -0.22). For lead, however, the biodiversity effect was

weaker (an order of magnitude smaller than for methylmercury). Increasing species richness

from one to ten species was associated with only a 10% increase in average lead concentrations,

and a small reduction in maximum portion size, P_{max} (b_{Pmax}= -0.039, 95% CI -0.049, -0.034). For

cadmium, b_{Pmax} = -0.12 (95% CI -0.13, -0.12), and for arsenic, b_{Pmax} = -0.17 (95% CI -0.19, -

0.14). These differences in the strength of biodiversity effects among contaminants are linked to

the shape of the distribution of their concentrations among fish species (i.e. normal vs.

lognormal) (**Figure 4A-D**), and the more skewed the distribution, the stronger the biodiversity

effect.

Figure 4. Frequency histograms of concentration of arsenic (**A**), cadmium (**B**), lead (**C**) and

methylmercury (**D**) in edible muscle tissues of North American aquatic species relative to

Provisional Tolerable Daily Intakes (PTDI). (**E**) Increasing seafood species richness increases

the number of contaminants which exceed the upper tolerable limit (PTDI) in a 100g portion

(*NC*). Black line indicates the mean *NC* from 1000 bootstrapped samples of seafood diets

sampled from the North American seafood contaminant dataset and grey shading refers to 95%

confidence intervals. Slope, $b_{NC} = 0.10$, 95% CI 0.084, 0.12.

Biodiversity benefits were consistent at local and global scales

Consistent with the positive biodiversity effects we observed when sampling diets from a global

seafood species pool, we also found benefits of seafood diversity in a local context. We analyzed

the effects of biodiversity on nutrient content in fourteen traditional indigenous North American

diets of seafood harvested and consumed locally. We found a consistent, beneficial effect of

biodiversity on *NT* and *P_{min}*, although the magnitude of the biodiversity effect was generally

lower at the local scale than the global scale (SI Appendix, **Figure S4**) (global b_{Pmin} = -0.42

(95% CI -0.47, -0.38) vs mean local b_{Pmin} = -0.25 ± 0.0091 S.E. and global b_{NT} = 0.20 (95% CI

0.20, 0.21) vs mean local b_{NT} = 0.097 ± 0.0082 S.E.). This finding is consistent with lower

nutritional functional diversity (NFD, Table S1) *sensu* (34) (mean local *NFD* = 2.77 ± 0.17 S.E.

vs. global *NFD* = 3.87) and higher nutritional functional evenness in local diets (mean local

NFEve = 0.82 ± 0.0037 S.E. vs. global *NFEve* = 0.57) (SI Appendix, Methods section 3, SI

Appendix, **Figure S5**), suggesting that functional consequences of changes to diversity in local

seafood diets may be buffered by higher nutritional redundancy among species.

Nutritional traits covary with ecological traits

We found that nutrient concentrations varied substantially among species in ways that differed

for different nutrients (**Figure 2**) -- a condition necessary for biodiversity *per se* to increase

nutritional benefits. The diversity that we observed in the nutrient content of edible portions

(**Figure 2**) was partly explained by ecological attributes and functional traits: habitat, trophic

position, body size, diet source and feeding mode (SI Appendix, Tables S4-S8). When

considering all five micronutrients (calcium, iron, zinc, EPA and DHA) together, finfish,

crustaceans and molluscs differed significantly in their multi-nutrient profiles (SI Appendix,

Table S1, PERMANOVA, $F_{2,103} = 3.429$, $p = 0.006$). Among finfish, nutrient concentrations

depended on which tissues were included in the edible portions (significant 'body part' effect

shown in SI Appendix, Tables S4, S6-S11). Finfish species whose edible portions included

organs such as liver or bones, had higher nutrient concentrations in the edible portion than those

whose edible portions were restricted to muscle tissue (ANOVA $p < 0.01$ for calcium, iron and

zinc concentrations; SI Appendix, **Figure S6**). Principal components analysis of multiple

nutrient concentrations in species showed that essential element (calcium, iron and zinc)

concentrations were typically negatively correlated with essential fatty acid concentrations (EPA

and DHA) (SI Appendix, **Figure S7**), allowing complementarity among species to increase

nutritional benefits. Specifically, high EPA and DHA concentrations traded off against low

calcium and zinc concentrations (and vice-versa) (negative pairwise Pearson correlation

coefficients; SI Appendix, **Figure S7**). When considering muscle tissues or muscle and skin

tissues of finfish only (thereby eliminating the influence of body parts such as bones on nutrient

concentration), concentrations of calcium, iron, zinc, EPA and DHA were associated with

ecological traits across species, including habitat and diet source (e.g. demersal vs. pelagic),

body size and trophic position (**Figure 5**). Relationships between species' nutrient tissue

concentrations and their habitats and trophic positions have been predicted by ecological

stoichiometry theory (58). Relationships between ecological traits and nutrient concentrations

differed for different nutrients. For example, tissue concentrations decreased with body size for

calcium, but not for the other microelements, EPA or DHA (**Figure 5,** SI Appendix, Figures S8-

S10, Tables S4 - S8). Species at lower trophic positions had higher zinc concentrations in their

tissues than species at higher trophic positions, but we did not observe this relationship for other

nutrients (SI Appendix, Tables S4 - S8). These examples illustrate that trade-offs and variation in

nutrient concentrations across species were associated with variation in different ecological traits

and roles that species play in ecosystems.

Figure 5. Nutrient concentrations in finfish muscle tissue vary with ecological traits in ways that

differ among the essential trace elements (calcium, iron, zinc) and the essential fatty acids (EPA

and DHA). Model-averaged standardized regression coefficients and 95% confidence intervals

from phylogenetic least squares regression (see SI Appendix, Methods 4 for full model

description) are shown for samples including muscle or muscle and skin tissues only. Traits for

which there is no symbol did not appear in any of the models in the top 95% set (with cumulative

sum of Akaike weights <= 0.95). Open symbols indicate coefficient estimates for which 95%

confidence intervals do not encompass zero. Note that x-axes differ across panels for clarity of

presentation. Number of species: $n = 155$ for EPA, $n = 159$ for DHA, $n = 104$ for calcium, $n = 99$

for iron, and $n = 90$ for zinc. For model results including tissues other than muscle tissue, see SI

Appendix, Tables S4-S11.

Nutritional value was linked to diversity of ecological functions

Ecological functional diversity of a species assemblage captures variation in traits and ecological

roles of species (48, 49), and is understood to play an important role in the relationship between

biodiversity and ecosystem function (**Figure 6A**). We assessed the relationship between

ecological functional diversity of seafood species diets (independent from species richness) and

nutritional value. Consistent with observations for ecosystem functions such as productivity and

biomass, nutritional benefits and ecological functional diversity were positively related, such that

seafood diets with higher ecological functional diversity also provided higher nutritional value

(i.e. were more likely to reach five micronutrient RDA targets simultaneously, $NT = 5$) (**Figure**

6B). Ecological functional diversity increased with species richness (SI Appendix, **Figure S11**),

and higher levels of ecological functional diversity were also associated with lower minimum

portion size required (SI Appendix, **Figure S12**). Because aquatic assemblages with higher

ecological functional diversity have been shown to exploit more diverse resources, transform and

transport energy and materials more efficiently, produce higher yields, and be more productive

and resilient over time (42, 59–61), it is possible that the provisioning of multiple micronutrients

occurs in tandem with a range of other ecological functions.

Figure 6. A) Ecological functional diversity captures variation in the ecological traits and

ecological roles of species, and can be quantified as the average of the distances between

species' trait values (circles) and the centre of functional trait space (square) (indicated by z)

(adapted from (93)). B) Seafood diets with higher levels of ecological functional diversity are

associated with higher levels of nutritional benefits (i.e. higher number of RDA targets per 100g

portion). Probability of reaching five micronutrient RDA targets simultaneously in a single 100g

portion ($NT = 5$, light orange line) increases as the ecological functional diversity of the diet

increases. Probabilities predicted from ordinal logistic regression (log odds of functional

diversity = 2.08, 95% CI (0.52, 3.64), $n = 1000$ resampled diets).

Discussion

Biodiversity of seafood provides high levels of nutritional benefits to humans because nutrient

concentrations vary substantially among species in ways that differ for different nutrients

(**Figures 2, 5,** SI Appendix, **Figure S7**). The effects of species richness observed for nutritional

benefits equal or exceed mean observed diversity effects documented for plant and forest species

richness and productivity (12, 40, 52, 62). Our findings build on evidence showing that

biodiversity in marine systems enhances ecological functions by extending this paradigm to

human nutrition. In agro-ecosystems, specific combinations of species such as corn, pumpkin

and beans have been planted to exploit complementary traits (shade provisioning, nitrogen

fixation and biomass production), to attain higher yields, more resilient crops and enhanced

nutritional benefits (35). By demonstrating that nutritional benefits of seafood can be understood

as a consequence of seafood species richness and ecological functional diversity, we have shown

that diverse seafood assemblages also provide nutritional and other ecological benefits

simultaneously.

Our analysis provides new and robust evidence that biodiversity is critical to multifunctionality

of ecosystem services when function thresholds are grounded *a priori* in multivariate metrics

meaningful for human well-being such as RDA. Our approach overcomes the critique that

multifunctionality is not enhanced by biodiversity (63), but rather a statistical artifact of how

multifunctionality was commonly estimated. Our findings are robust to a range of RDA target

levels (SI Appendix, **Figure S2**). In the case of P_{min}, we found that the benefit of biodiversity

was consistent across all RDA target levels considered, and in the case of *NT*, we found

beneficial effects of biodiversity across a range levels of nutritional value that are significant for

human nutrition (i.e. 1-40% RDA per 100g portion), highlighting the importance of species

richness in seafood diets. More generally, ecosystem service benefits, as defined in metrics of

human wellbeing rather than the traits of the species pool under consideration (e.g., biomass or

stability of the food web), typically are produced by several underlying ecosystem functions

(54). The strong effects of diversity on multifunctional benefits observed here may also apply to

relationships between diversity and other services e.g., desired filtration rates of pollutants in

wetlands (64), or desired pest consumption rates in agricultural systems (65).

Physiological and ecological mechanisms underlying nutritional benefits and risks

When comparing the magnitude of the biodiversity-ecosystem function scaling parameter, b,

among the nutrients and contaminants, we found that the magnitude of b was higher for nutrients

with more skewed distributions of tissue concentrations (**Figure 2**, **Figure 4**). For all of these

highly skewed distributions, we observed a strong biodiversity effect (i.e. high values of b).

However, for protein and lead in muscle tissues, we observed symmetrical tissue concentration

distributions. In these two cases, the biodiversity effect was either non-existent (in the case of

protein) or very weak (in the case of lead). This finding suggests that biodiversity effects may

increase when the trait distribution is skewed and includes some species with extreme trait

values, known as 'functionally specialized' and 'functionally unique' species (9, 66, 67).

Understanding the drivers of the tissue concentration distribution among species may lend

insights into how biodiversity effects may change as the nutrient under consideration changes or

as the environment changes.

Increasing seafood species richness increases nutritional benefits as well as contaminant

exposure. Increasing biodiversity can have negative consequences for some ecosystem functions,

despite positive consequences for others. On balance, the benefits of diversity may outweigh the

negative effects on function when considering multiple functions together (68). Here we found

that the same mechanisms (lognormal and complementary tissue concentration distributions

among species) that contributed to the positive relationship between biodiversity and human

nutritional benefits also applied to contaminant exposure. As a result, increasing seafood species

richness comes with both benefits and risks (69–71). However, interpreting our results in the

context of public health outcomes is complicated by the fact that epidemiological evidence on

health outcomes of contaminant exposure is mixed and likely dependent on complex social and

health risks. The health risks of contaminant exposure depend on other health factors such as

smoking or nutrients in the diet (72, 73) and disease status. Complex interactions among multiple

diet and health risk factors (74) were beyond the scope of this study, but would be necessary to

understand exposure risks from seafood consumption in any particular community. Nonetheless,

our results suggest that while increasing biodiversity increases contaminant exposure, it also

increases nutritional benefit, and reduces the portion sizes required to meet nutritional demands.

Finding a balance between seafood biodiversity, seafood biomass consumption, and the resulting

risks and benefits will be critical for both human and ecosystem health.

Aquatic biodiversity and food security in a changing world

Seafood-derived nutrition plays an important role in food security. The link that we have

demonstrated between seafood biodiversity (species richness and ecological functional diversity)

and nutrition in an ecological framework unites three of the United Nations Sustainable

Development Goals focused on biodiversity, hunger and well-being (75). More than two billion

people suffer from micronutrient deficiencies (76, 77), and many of the most nutritionally

vulnerable populations – those that are deficient in essential micronutrients during particularly

sensitive stages of life (i.e. pregnancy, breastfeeding and childhood) - may rely heavily on local

aquatic ecosystems to meet their nutritional demands (15, 19, 22, 78). These populations may

have access to a limited amount of locally available seafood tissue each day, suggesting that

nutritional efficiency (i.e. lower P_{min}) provided by biodiversity in wild-caught seafood may be

particularly important for these populations. Regions of high nutritional vulnerability continue to

experience major changes in biodiversity and ecosystem structure (6, 17, 79), and climate change

is compounding threats to the sustainability of capture fisheries (2, 80). Our results suggest that

declines in these aspects of diversity in wild aquatic ecosystems could make achieving

sustainability goals for food security via seafood even more difficult. Seafood diets composed of

more species, and groups of species with higher levels of ecological functional diversity, are

more likely to provide more nutrients per unit biomass than less diverse seafood diets, while also

maintaining high levels of ecosystem function (26). This finding bridges the growing

understanding of hidden hunger and food security with the large theoretical and empirical

understanding of relationships between biodiversity, ecosystem function and benefits to people.

Biodiversity in natural aquatic systems can be maintained by reducing pollution and overharvest

and by allowing ecosystems to adapt to climate change, and these measures could also benefit

humanity directly through seafood provisioning.

Conclusions

Nutritional value appears to be derived from ecological diversity, suggesting links between the

complexity of aquatic ecosystems and their capacity to produce nutritional benefits. While the

role of seafood is well-recognized as an important source of protein in the human diet, the role of

seafood biodiversity as an important aspect of the provision of essential micronutrients has been

overlooked. Our results reveal that aspects of ecological structure including species and

ecological functional diversity enhance nutritional benefits, while also increasing contaminant

exposure, thereby linking the processes that structure ecosystems with their potential benefits

and risks to human nutrition and health.

Methods

To test our hypotheses about aquatic species diversity and potential nutritional benefits for human well-being, we assembled a new database of nutritional values by synthesizing observations from existing data (SI Appendix Methods, **Figure S1**). To build the database, we identified quantitative and comparable measures of nutritional content for the edible portions of aquatic species (SI Appendix Methods 1.1). We then identified metrics for relating nutrient content to human health (Methods 1.1, 1.3). We repeated this exercise for contaminants (Methods 1.2). Next, we adapted an approach from biodiversity - ecosystem functioning theory for quantitatively assessing the potential nutritional value of seafood diets varying in their species composition and diversity for multiple nutrients separately and simultaneously (Methods 2.1). To test predictions of biodiversity-ecosystem functioning theory, we simulated potential diets at global and local scales and fit models to test for effects of species richness and functional diversity (Methods 2.2 - 2.3). SI Appendix, **Figure S1** provides a graphical overview of our analyses.

1. Metrics

1.1 Quantifying nutritional value for single nutrients

We characterized an aquatic species' nutritional value by drawing on two well-established nutritional metrics: nutrient concentration (mg/100 g edible portion) and Recommended Dietary Allowances (RDA). RDAs are developed following health guidelines to quantify the recommended amount of a particular nutrient required to maintain health (50). We used RDAs established by the Food and Nutrition Board of the United States Institute of Medicine to

quantify the daily intake level of a nutrient needed to meet the requirements of 97–98% of

healthy adults (females aged 19-50) (50) (SI Appendix, Table S2). We refer to an RDA-defined

threshold as an *RDA target* (SI Appendix, Tables S1 and S2). We calculated a ratio of the

nutrient content in a 100g edible portion relative to the RDA (or fraction thereof) for that

nutrient. For many species, nutrient concentrations in edible tissues provided only small fractions

of the RDA (SI Appendix, Table S2). Following the IOM, we chose an RDA target as 10% of

RDA for a given nutrient in a single portion (SI Appendix, Table S2) because this is a minimum

threshold for a food to be considered of nutritional benefit (50). We defined the minimum

portion size required, P_{min}, as the minimum amount of edible seafood issue (g) required to reach

the RDA target for a given nutrient (or a set of nutrients, Methods 1.3) (SI Appendix, Table S1,

Equation A1). We quantified the sensitivity of our biodiversity findings to different threshold

levels of RDA for both single nutrient analyses and multi-nutrient analyses (Methods 2.2) and

found that for P_{min}, they did not vary with threshold level, and for *NT*, they varied with threshold,

but were robust to RDA threshold levels between 1% and 40% RDA (SI Appendix, **Figure S2**).

We examined data on concentrations of macronutrients including protein and fat, as well as five

micronutrients (n = 5041 observations of nutrient concentrations, SI Appendix, **Figure S1A**):

metals beneficial at low concentrations but toxic at high concentrations (zinc and iron), one

beneficial mineral (calcium) and the polyunsaturated fatty acids eicosapentaenoic acid (EPA)

and docosahexaenoic acid (DHA). We chose these five micronutrients because we required that

RDA standards exist for each one (50) and that each nutrient has known functions in organismal

physiology, and are considered 'biologically essential' because they are required by organisms to

grow or reproduce. The concentration of biologically essential nutrients in organisms' tissues are

controlled homeostatically by organisms (81, 82), and therefore might be biologically related to

ecological trait values we considered in our trait analysis.

1.2 Quantifying contaminant exposure in terms of human health risks

We characterized an aquatic species' contaminant content relative to established public health

guidelines for exposure. We used the Provisional Tolerable Weekly Intake (PTWI) developed by

the FAO/WHO Expert Committee on Food Additives (JEFCA) (57), which estimates the amount

of a substance in air, food, soil or drinking water that can be assimilated weekly per unit body

weight over a lifetime without appreciable health risk (57). To make this metric comparable to

RDA targets defined above, we quantified contaminant exposure per day by dividing the PTWI

by seven and using the body weight of a 70-kg person, to calculate a daily tolerable limit for use

in our analyses, which we refer to as Provisional Tolerable Daily Intake, PTDI (SI Appendix,

Table S1). In our analyses, we consider a tolerable upper limit to be exceeded if an edible

portion contains 100% or more of the PTDI. We examined the sensitivity of this choice of 100%

threshold by considering lower values (e.g. 50%) and found that for all values between 50% and

100% of PTDI our findings remain consistent.

1.3 Defining nutritional benefits and risks for multiple nutrients or contaminants across diverse

species groups

To test our hypothesis that nutritional value may depend on seafood diversity in a way that

considers multiple nutrients at the same time, we used two metrics that considered multiple

micronutrients simultaneously (**Figure 1**): (i) the minimum portion size introduced above, P_{min},

(SI Appendix, Table S1, Equation A1), which quantifies the amount of tissue, in grams, required

to reach the RDA target for 5 nutrients simultaneously and (ii) number of nutrients for which

RDA targets are met in a standard 100g edible portion, NT (SI Appendix, Table S1, Equation

A2). Note that our NT differs from another measure 'Nutritional functional diversity, NFD' (34)

which is a measure of nutritional profile in multivariate trait space and does not explicitly

quantify the number of nutrient RDA targets met in a portion. Our NT is a measure of potential

nutritional value and is derived from the biodiversity-ecosystem function perspective to allow us

to compare nutritional value with other functions that depend on biodiversity (e.g., productivity,

resource cycling). This measure does not consider the potential physiological interactions or co-

benefits of consuming multiple nutrients that would be considered in metabolic models for

human health. To test our hypothesis that contaminant exposure also varies with seafood

diversity, we defined P_{max}, as the amount of tissue beyond which a tolerable upper limit (PTDI)

for a given contaminant would be reached (SI Appendix, Table S1, Equation A3, SI Appendix

Methods 2). Analogous to the NT metric defined above for the nutrients, we defined the number

of distinct PTDIs in a 100g portion as NC (Table S1, Equation A4, SI Appendix Methods 2). We

chose to standardize NT and NC relative to 100g of seafood per day because 100g edible portion

is a standard metric used in the food and nutrition literature (37, 83) and consumption rates of

100g per day is within the range of daily consumption rates for many communities that rely

heavily on seafood to meet nutritional needs and for whom seafood may also pose substantial

contaminant exposure risks (16, 45, 84, 85).

2. **Statistical analyses and hypothesis testing**

2.1 Modeling the biodiversity effect for sample diets

To quantitatively compare effects of biodiversity among possible seafood diets varying in

species richness, we modeled species richness effects as a power function:

$$Y = aS^b \tag{1}$$

where the parameter b is referred to as the 'biodiversity effect' and describes the relationship

between a change in species richness, S, and a measure of function, Y, such as P_{min} or NT, where

a is a constant (12, 52).

2.2 Simulating seafood diets and estimating nutritional benefits at different seafood species

richness levels

We tested the effect of species richness, S, on nutritional value by randomly assembling diets

from the global seafood species pool at varying levels of species richness (SI Appendix, **Figure**

S1B). In our analyses of NT, we kept the total biomass constant at each level of species richness

(100g). In analyses using the global dataset (547 species) (SI Appendix Methods 1.1), we

assembled diets from the entire global species pool, choosing species at random without

replacement. This way of simulating diets certainly ignores economic, social and cultural factors

that affect which species people consume, but allows us to consider the potential effect of

biodiversity on diets before diets are filtered by these other processes. To assess potential effects

of biodiversity on nutritional value for populations that consume seafood locally from a

restricted species pool, we sampled diets from species contained within traditional diets in

fourteen indigenous cultures in North America (SI Appendix, Methods 1.4).

We created sample diets by sampling 10 species at random from the global species pool and then

assembling seafood diets from all possible combinations of these ten randomly chosen species at

10 levels of species richness (1-10) to generate 1023 simulated diets (SI Appendix, **Figure S1B,**

i).   At each level of species richness, we assembled diets following the typical experimental

design employed to test the hypothesis that biodiversity affects ecosystem functioning, analogous

to a biodiversity-ecosystem function experiment with a replacement design (86), where species'

abundances in the diet decline proportionally as species richness increases such that each species

contributed an equal proportion of biomass. For each diet at each level of species richness (i.e. n

$= 1023$), we calculated P_{min}, (Methods 1.1, 1.3, SI Appendix, Table S1 Equation A1**,** SI

Appendix, **Figure S1C, ii**) (for either: one of six possible nutrient targets individually (protein,

calcium, iron, zinc, EPA and DHA), or five micronutrients (calcium, iron, zinc, EPA and DHA

targets simultaneously) (Methods 1.3). To estimate NT (Methods 1.3, SI Appendix, **Figure S1C,**

ii), we quantified the number of distinct nutrient RDA targets by assigning each diet ($n = 1023$) a

set of 0's or 1's according to whether that combination met the RDA target for each nutrient (see

SI Appendix, Table S1, Equation A2). This approach allowed us to explore how likely it would

be for potential human diets containing different numbers of seafood species to reach RDA

targets for a given number of micronutrients (NT ranges between 0 and 5), assuming that seafood

species were included in the human diet at different levels of richness at random. At each level of

species richness, we averaged P_{min} and NT. We then repeated this process of random sampling

ten species from the species pool, assembling diets at each level of richness, estimating metrics

of nutritional value and averaging at each richness level 1000 times, yielding 1000 estimates of

each metric at each richness level (SI Appendix, **Figure S1C, iii**).

2.3 Testing hypotheses that biodiversity enhances nutritional benefits

We tested the hypothesis that complementarity in nutrient concentrations among species

increases nutritional benefits by increasing NT (SI Appendix, **Figure S1D**). We quantified the

effect of seafood species richness, S, on NT at each richness level estimated in Methods 2.2 ($n =$

1000 estimates of NT per richness level), by fitting a power function of the form shown in

Equation 1,

$$NT = aS^{b_{NT}} \tag{2}$$

where the parameter b_{NT} describes the relationship between a change in species richness, S, and a

change in NT, and a is a constant.

We tested the hypothesis that complementarity in nutrient concentrations among species reduces

the minimum portion size required, P_{min} (Methods 1.1, 1.3, 2.2), by estimating the effect of

species richness, S, on P_{min} ($n = 1000$ estimates of P_{min} per richness level) at each richness level

using

$$P_{min} = aS^{b_{Pmin}} \tag{3}$$

where the parameter b_{Pmin} describes the relationship between a change in species richness, S, and

a change in P_{min}, and a is a constant (SI Appendix, **Figure S1D**).

We estimated *a* and *b* in Equations 2 and 3 using non-linear regression using the *nls.LM* function

in the *minpack.lm* package in R (87). We conducted all analyses in R version 3.3.2 (88). To

quantify uncertainty in parameter estimates associated with the fit of Equations 2 and 3 to our

estimates of function (Methods 2.2), we calculated bootstrapped confidence ($n = 1000$

bootstraps) intervals using non-parametric bootstrapping of mean centered-residuals using the

nlsBoot function in the R package *nlstools* (89). For both P_{min} and *NT*, we tested the hypothesis

that biodiversity enhances nutritional benefits by assessing whether the estimate of the scaling

exponent, *b*, had confidence intervals not overlapping zero.

We tested the hypothesis that nutrient concentrations are related to species' ecological traits in

two ways: 1) testing whether multi-nutrient profiles (i.e. concentrations of all five

micronutrients) differ among major phylogenetic groups using PERMANOVA (SI Appendix,

Methods 4) and 2) whether differences in single nutrient concentrations differ with species'

ecological traits using phylogenetic least squares regression (SI Appendix, Methods 4). We

quantified the relationship between nutritional benefits and ecological functional diversity (EFD,

SI Appendix, Table S1) by estimating *NT*, P_{min} and *EFD*, estimated as functional dispersion, of

diets simulated from the global species pool (SI Appendix, Methods 5).

2.4. Uncertainties:

There are several sources of uncertainty in our analyses. First, there are substantial sources of

uncertainty in food composition estimates. The data in our dataset meet international standards

for data quality and standardization, meaning that we followed guidelines for checking food

composition data and converting units, denominators and expressions (90). Still, tissue

concentrations may vary depending on analytical techniques, labs, season, diet of the animal, life

stage etc. Some of these sources of uncertainty (e.g. differences in analytical techniques) are

unavoidable consequences of synthesizing previously published data collected across many labs.

We assumed that these uncertainties in the data were randomly distributed over our

geographically and taxonomically diverse dataset. Further uncertainty is associated with how

well our set of 547 species represents the global pool of seafood consumed. We do not know

whether our sample is random or biased, though we can say that our dataset includes 41 of the 67

most consumed species worldwide (as determined by FAO production volumes (91), species

with capture production of 150 000 tonnes or more, after removing species for which the

majority of production volume is diverted to fish meal and oil (92), SI Appendix, Table S13). A

remaining source of variation among samples is likely due to natural sources of variation

associated with seasonal and other sources of temporal variability, which we consider to be an

important component of biodiversity.

Acknowledgments: We thank P. Thompson, A. Gehman, W. Cheung, M. Ruckelshaus and C.

Harley, M. Deith, L. Dee, and F. Isbell for helpful comments on previous versions of this

manuscript. Portions of this manuscript were adapted from and based on work contained

in the doctoral thesis of J.R. Bernhardt from the University of British Columbia.

Funding: Funding was provided by a Vanier Canada Graduate Scholarship to JRB.

Competing interests: We declare no competing interests.

Data and materials availability: Data and code are available at

https://github.com/JoeyBernhardt/Nutrient_analysis, archived at 10.5281/zenodo.4474988.

Data are available on Dryad (https://doi.org/10.5061/dryad.rn8pk0p8t).

References

1.   R. Elahi, *et al.*, Recent trends in local-scale marine biodiversity reflect community structure and
human impacts. *Curr. Biol.* **25**, 1938–1943 (2015).

2.   M. L. Pinsky, B. Worm, M. J. Fogarty, J. L. Sarmiento, S. A. Levin, Marine taxa track local climate
velocities. *Science* **341**, 1239–1242 (2013).

3.   D. J. McCauley, *et al.*, Marine defaunation: animal loss in the global ocean. *Science* **347**, 1255641
(2015).

4.   FAO, *The State of the World's Biodiversity for Food and Agriculture*, J. Bélanger, D. Pilling, Eds.
(FAO Commission on Genetic Resources for Food and Agriculture Assessments, 2019).

5.   IPBES, *Global assessment report on biodiversity and ecosystem services of the Intergovernmental*
Science-Policy Platform on Biodiversity and Ecosystem Services, E. S. Brondizio, J. Settele, S. Díaz,
H. T. Ngo, Eds. (2019).

6.   G. P. Jones, M. I. McCormick, M. Srinivasan, J. V. Eagle, Coral decline threatens fish biodiversity
in marine reserves. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 8251–8253 (2004).

7.   R. A. Myers, B. Worm, Rapid worldwide depletion of predatory fish communities. *Nature* **423**, 280–
283 (2003).

8.   J. A. Hutchings, R. A. Myers, What Can Be Learned from the Collapse of a Renewable Resource?
Atlantic Cod, Gadus morhua, of Newfoundland and Labrador. *Can. J. Fish. Aquat. Sci.* **51**, 2126–
2146 (1994).

9.   C. Pimiento, *et al.*, Functional diversity of marine megafauna in the Anthropocene. *Sci Adv* **6**,
eaay7650 (2020).

10.   S. S. Myers, *et al.*, Human health impacts of ecosystem alteration. *Proc. Natl. Acad. Sci. U. S. A.*
110, 18753–18760 (2013).

11.   F. Isbell, *et al.*, Linking the influence and dependence of people on biodiversity across scales. *Nature*
546, 65–72 (2017).

12.   M. I. O'Connor, *et al.*, A general biodiversity--function relationship is mediated by trophic level.
Oikos **126**, 18–31 (2017).

13.   E. Chivian, "Biodiversity: Its Importance to Human Health" (Center for Health and the Global
Environment Harvard Medical School, 2002).

14.   FAO, *The State of Food Security and Nutrition in the World: Building Resilience for Peace and*
Food Security, Rome: Food and Agriculture Organization of the United Nations, Ed. (2017).

15.   H. V. Kuhnlein, O. Receveur, Local Cultural Animal Food Contributes High Levels of Nutrients for
Arctic Canadian Indigenous Adults and Children. *J. Nutr.* **137**, 1110–1114 (2007).

16.   A. M. Cisneros-Montemayor, D. Pauly, L. V. Weatherdon, Y. Ota, A Global Estimate of Seafood
Consumption by Coastal Indigenous Peoples. *PLoS One* **11**, e0166681 (2016).

17. J. G. Molinos, *et al.*, Climate velocity and the future global redistribution of marine biodiversity.
Nat. Clim. Chang. **6**, 83–88 (2016).

18. WHO, *Connecting global priorities: biodiversity and human health: a state of knowledge review*,
World Health Organization and Secretariat of the Convention on Biological Diversity, Ed. (2015).

19. C. D. Golden, *et al.*, Fall in fish catch threatens human health. *Nature* **534**, 317–320 (2016).

20. B. Belton, S. H. Thilsted, Fisheries in transition: Food and nutrition security implications for the
global South. *Global Food Security* **3**, 59–66 (2014).

21. FAO, *The State of World Fisheries and Aquaculture 2018 - Meeting the sustainable development
goals* (2018).

22. N. Kawarazuka, C. Béné, Linking small-scale fisheries and aquaculture to household nutritional
security: an overview. *Food Security* **2**, 343–357 (2010).

23. B. J. Cardinale, *et al.*, Biodiversity loss and its impact on humanity. *Nature* **486**, 59–67 (2012).

24. D. Tilman, *et al.*, The Effects of Plant Composition and Diversity on Ecosystem Processes. *Science*
277, 1302–1305 (1997).

25. M. Loreau, Biodiversity and ecosystem functioning: recent theoretical advances. *Oikos* **91**, 3–17
(2000).

26. J. E. Duffy, C. M. Godwin, B. J. Cardinale, Biodiversity effects in the wild are common and as
strong as key drivers of productivity. *Nature* **549**, 261–264 (2017).

27. M. T. Ruel, Operationalizing dietary diversity: a review of measurement issues and research
priorities. *J. Nutr.* **133**, 3911S–3926S (2003).

28. E. A. Frison, J. Cherfas, T. Hodgkin, Agricultural Biodiversity Is Essential for a Sustainable
Improvement in Food and Nutrition Security. *Sustain. Sci. Pract. Policy* **3**, 238–253 (2011).

29. J. Fanzo, D. Hunter, T. Borelli, F. Mattei, *Diversifying Food and Diets: Using Agricultural
Biodiversity to Improve Nutrition and Health* (Routledge, 2013).

30. E. A. Frison, I. F. Smith, T. Johns, J. Cherfas, P. B. Eyzaguirre, Agricultural biodiversity, nutrition,
and health: making a difference to hunger and nutrition in the developing world. *Food Nutr. Bull.* **27**,
167–179 (2006).

31. Á. Toledo, B. Burlingame, Biodiversity and nutrition: A common path toward global food security
and sustainable development. *J. Food Compost. Anal.* **19**, 477–483 (2006).

32. A. D. Jones, Critical review of the emerging research evidence on agricultural biodiversity, diet
diversity, and nutritional status in low- and middle-income countries. *Nutr. Rev.* **75**, 769–782 (2017).

33. C. Lachat, *et al.*, Dietary species richness as a measure of food biodiversity and nutritional quality of
diets. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 127–132 (2018).

34. R. Remans, *et al.*, Assessing nutritional diversity of cropping systems in African villages. *PLoS One*
6 (2011).

35. F. A. J. DeClerck, J. Fanzo, C. Palm, R. Remans, Ecological approaches to human nutrition. *Food
Nutr. Bull.* **32**, S41–50 (2011).

36. D. Penafiel, C. Lachat, R. Espinel, P. Van Damme, P. Kolsteren, A systematic review on the
contributions of edible plant and animal biodiversity to human diets. *EcoHealth* **8**, 381–399 (2011).

37. R. Remans, S. A. Wood, N. Saha, T. L. Anderman, R. S. DeFries, Measuring nutritional diversity of
national food supplies. *Global Food Security* **3**, 174–182 (2014).

38. S. A. Wood, Nutritional functional trait diversity of crops in south-eastern Senegal. *J. Appl. Ecol.* **55**,
81–91 (2018).

39. M. Loreau, A. Hector, Partitioning selection and complementarity in biodiversity experiments.
Nature **412**, 72–76 (2001).

40. P. B. Reich, *et al.*, Impacts of biodiversity loss escalate through time as redundancy fades. *Science*
336, 589–592 (2012).

41. B. J. Cardinale, Biodiversity improves water quality through niche partitioning. *Nature* **472**, 86–89
(2011).

42. L. E. Dee, *et al.*, Functional diversity of catch mitigates negative effects of temperature variability on
fisheries yields. *Proc. Biol. Sci.* **283** (2016).

43. R. Hilborn, T. P. Quinn, D. E. Schindler, D. E. Rogers, Biocomplexity and fisheries sustainability.
Proc. Natl. Acad. Sci. U. S. A. **100**, 6564–6568 (2003).

44. S. Wang, M. Loreau, Biodiversity and ecosystem stability across scales in metacommunities. *Ecol.
Lett.* **19**, 510–518 (2016).

45. J. R. Bogard, *et al.*, Higher fish but lower micronutrient intakes: Temporal changes in fish
consumption from capture fisheries and aquaculture in Bangladesh. *PLoS One* **12** (2017).

46. C. C. Hicks, *et al.*, Harnessing global fisheries to tackle micronutrient deficiencies. *Nature* **574**, 95–
98 (2019).

47. D. Tilman, The Ecological Consequences of Changes in Biodiversity: A Search for General
Principles. *Ecology* **80**, 1455 (1999).

48. D. Tilman, Functional diversity. *Encyclopedia of biodiversity* **3**, 109–120 (2001).

49. C. Violle, *et al.*, Let the concept of trait be functional! *Oikos* **116**, 882–892 (2007).

50. IOM, "Dietary reference intakes for calcium and vitamin d" (2011).

51. F. Micheli, B. S. Halpern, Low functional redundancy in coastal marine assemblages. *Ecol. Lett.* **8**,
391–400 (2005).

52. B. J. Cardinale, *et al.*, The functional role of producer diversity in ecosystems. *Am. J. Bot.* **98**, 572–
592 (2011).

53. J. E. K. Byrnes, *et al.*, Investigating the relationship between biodiversity and ecosystem
multifunctionality: Challenges and solutions. *Methods in Ecology and Evolution* **5**, 111–124 (2014).

54. P. Manning, *et al.*, Redefining ecosystem multifunctionality. *Nature Ecology & Evolution* **2**, 427–
436 (2018).

55. A. Hector, R. Bagchi, Biodiversity and ecosystem multifunctionality. *Nature* **448**, 188–190 (2007).

56. L. Gamfeldt, H. Hillebrand, P. R. Jonsson, Multiple functions increase the importance of
biodiversity for overall ecosystem functioning. *Ecology* **89**, 1223–1231 (2008).

57. WHO, *Evaluation of Certain Contaminants in Food: Seventy-third Report of the Joint FAO/WHO*
Expert Committee on Food Additives (2011).

58. R. W. Sterner, J. J. Elser, *Ecological stoichiometry: the biology of elements from molecules to the*
biosphere (Princeton University Press, 2002).

59. J. S. Lefcheck, *et al.*, Tropical fish diversity enhances coral reef functioning across multiple scales.
Sci Adv **5**, eaav6420 (2019).

60. C. Mora, *et al.*, Global human footprint on the linkage between biodiversity and ecosystem
functioning in reef fishes. *PLoS Biol.* **9**, e1000606 (2011).

61. J. S. Lefcheck, J. E. Duffy, Multitrophic functional diversity predicts ecosystem functioning in
experimental assemblages of estuarine consumers (2015) https:/doi.org/10.1890/14-1977.1.

62. J. Liang, *et al.*, Positive biodiversity-productivity relationship predominant in global forests. *Science*
354, aaf8957 (2016).

63. L. Gamfeldt, F. Roger, Revisiting the biodiversity--ecosystem multifunctionality relationship.
Nature Ecology & Evolution **1**, s41559–017 (2017).

64. T. Boyer, S. Polasky, Valuing urban wetlands: A review of non-market valuation studies. *Wetlands*
24, 744–755 (2004).

65. D. S. Karp, *et al.*, Forest bolsters bird abundance, pest control and coffee yield. *Ecol. Lett.* **16**, 1339–
1347 (2013).

66. D. Mouillot, N. A. J. Graham, S. Villéger, N. W. H. Mason, D. R. Bellwood, A functional approach
reveals community responses to disturbances. *Trends Ecol. Evol.* **28**, 167–177 (2013).

67. R. P. Leitao, *et al.*, Rare species contribute disproportionately to the functional structure of species
assemblages. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20160084 (2016).

68. J. S. Lefcheck, *et al.*, Biodiversity enhances ecosystem multifunctionality across trophic levels and
habitats. *Nat. Commun.* **6**, 6936 (2015).

69. M. C. Sheehan, *et al.*, Global methylmercury exposure from seafood consumption and risk of
developmental neurotoxicity: a systematic review. *Bull. World Health Organ.* **92**, 254–269F (2014).

70. H. V. Kuhnlein, H. M. Chan, Environment and contaminants in traditional food systems of northern
indigenous peoples. *Annu. Rev. Nutr.* **20**, 595–626 (2000).

71. P. W. Davidson, *et al.*, Methylmercury and neurodevelopment: longitudinal analysis of the
Seychelles child development cohort. *Neurotoxicol. Teratol.* **28**, 529–535 (2006).

72. D. Mozaffarian, E. B. Rimm, Fish intake, contaminants, and human health: evaluating the risks and
the benefits. *JAMA* **296**, 1885–1899 (2006).

73. D. Kinloch, H. Kuhnlein, D. C. Muir, Inuit foods and diet: a preliminary assessment of benefits and
risks. *Sci. Total Environ.* **122**, 247–278 (1992).

74. M. O. Gribble, *et al.*, Mercury, selenium and fish oils in marine food webs and implications for
human health. *J. Mar. Biol. Assoc. U. K.* **96**, 43–59 (2016).

75. UN, *Transforming our world : the 2030 Agenda for Sustainable Development*, UN General
Assembly, Ed. (2015).

76. J. C. Ruel-Bergeron, *et al.*, Global Update and Trends of Hidden Hunger, 1995-2011: The Hidden
Hunger Index. *PLoS One* **10**, e0143497 (2015).

77. J. Fanzo, *et al.*, 2018 Global Nutrition Report: Shining a light to spur action on nutrition (2018)
(June 12, 2020).

78. J. R. Bogard, *et al.*, Nutrient composition of important fish species in Bangladesh and potential
contribution to recommended nutrient intakes. *J. Food Compost. Anal.* **42**, 120–133 (2015).

79. W. W. L. Cheung, *et al.*, Shrinking of fishes exacerbates impacts of global ocean changes on marine
ecosystems. *Nat. Clim. Chang.* **3**, 254–258 (2013).

80. L. A. Rogers, *et al.*, Shifting habitats expose fishing communities to risk under climate change. *Nat.*
Clim. Chang. **9**, 512–516 (2019).

81. R. Karimi, C. L. Folt, Beyond macronutrients: element variability and multielement stoichiometry in
freshwater invertebrates. *Ecol. Lett.* **9**, 1273–1283 (2006).

82. M. T. Arts, M. T. Brett, M. Kainz, *Lipids in Aquatic Ecosystems* (Springer Science & Business
Media, 2009).

83. R. DeFries, *et al.*, Global nutrition. Metrics for land-scarce agriculture. *Science* **349**, 238–240
(2015).

84. N. Roos, M. A. Wahab, C. Chamnan, S. H. Thilsted, The Role of Fish in Food-Based Strategies to
Combat Vitamin A and Mineral Deficiencies in Developing Countries. *J. Nutr.* **137**, 1106–1109
(2007).

85. EPA, Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories (2000).

86. J. E. Byrnes, J. J. Stachowicz, The consequences of consumer diversity loss: different answers from
different experimental designs. *Ecology* **90**, 2879–2888 (2009).

87. T. V. Elzhov, K. M. Mullen, A.-N. Spiess, B. Bolker, *minpack.lm: R interface to the Levenberg-*
Marquardt nonlinear least-squares algrothim found in MINPACK, pluss support for bounds (2013).

88. R Core Team, R: A Language and Environment for Statistical Computing (2019).

89. F. Baty, *et al.*, A Toolbox for Nonlinear Regression in R : The Package nlstools. *J. Stat. Softw.* **66**,
1–21 (2015).

90. Food and Agriculture Organization/ International Network of Food Data Systems (INFOODS),
"FAO / INFOODS Guidelines for Checking Food Composition Data prior to the publication of a
User Table / Database - Version 1 . 0" (2012).

91. Food and Agriculture Organization of the United Nations, "FAOSTAT. Calculated from food
balance sheets" (2016).

92. T. Cashion, F. Le Manach, D. Zeller, D. Pauly, Most fish destined for fishmeal production are food-
grade fish. *Fish Fish* **18**, 837–844 (2017).

93. E. Laliberté, P. Legendre, A distance-based framework for measuring functional diversity from
multiple traits. *Ecology* **91**, 299–305 (2010).