Aquatic biodiversity enhances multiple nutritional benefits to humans

Joey R. Bernhardt[1]*, Mary I. O'Connor [1]

[1]Department of Zoology and Biodiversity Research Centre, University of British Columbia,
Vancouver, Canada, V6T 1Z4.

*Correspondence to: joey.bernhardt@biodiversity.ubc.ca.

ORCIDs:

Joey R. Bernhardt: 0000-0003-1824-2801

Mary I. O'Connor: 0000-0001-9583-1592

Classification: Biological Sciences/Ecology

Keywords: seafood, biodiversity-ecosystem functioning, ecosystem services

15
16 **Author contributions:** JRB conceived the study, collected and curated the data. JRB analyzed
17 and visualized the data with input from MIO. JRB wrote the original draft of the manuscript,
18 JRB and MIO contributed to writing and editing. MIO supervised the study.
19
20 **This PDF file includes:**
21 Main Text
22 Figures 1 to 4
23
24

Abstract

Global food security relies on protein and essential micronutrients provided by seafood (Latham

1997, Golden et al. 2016). The importance of seafood-derived micronutrients to human health

suggests that consuming diverse aquatic species could enhance human well-being (Penafiel et al.

2011). Biodiversity provides important ecosystem benefits to humanity (Cardinale et al. 2012),

causing concern that declining biodiversity may affect human health (Myers et al. 2013). A key

scientific question remains unanswered - does increasing species diversity in seafood diets

improve their ability to meet nutritional needs? Here we used biodiversity-ecosystem functioning

theory to test whether increasing species diversity allows seafood diets to fulfill multiple

nutritional requirements simultaneously, a condition necessary for human health. We found that

aquatic species with different ecological traits have distinct and complementary micronutrient

profiles. The same complementarity mechanisms that generate positive biodiversity effects on

ecosystem functioning in terrestrial ecosystems also operate in seafood assemblages, allowing

for more diverse diets to yield increased nutritional benefits. Notably, nutritional metrics that

capture multiple micronutrients essential for human well-being depend more strongly on

biodiversity than ecological measures of function such as productivity. In contrast to the

micronutrients, we found that increasing species diversity did not increase the amount of protein

in seafood diets. We unify biodiversity-ecosystem functioning theory and human nutrition,

demonstrating a direct link between multiple nutritional benefits of biodiversity and an

ecosystem service underpinning human wellbeing. Our findings demonstrate that minimizing

biodiversity loss at local and global scales will benefit global food and nutrition security.

Significance statement:

Food security is not simply about maintaining yields, it is also the need for a stable supply of

nutritionally diverse foods. Obtaining nutritious food is a major challenge facing humanity, and

aquatic ecosystems can help meet this goal. From the perspective of human nutrition, how much

biodiversity is enough biodiversity? How ecological processes influence the capacity of aquatic

ecosystems to provide nutritionally diverse diets is largely unknown. We found that aquatic

biodiversity enhanced nutritional benefits, because edible species showed distinct and

complementary multi-nutrient profiles. Protein supply was independent of biodiversity, but the

supply of micronutrients increased with biodiversity. Extending the multifunctional benefits of

biodiversity to human nutrition underscores the need to minimize biodiversity loss for the

benefits of humanity.

Introduction

Species losses and range shifts due to climate change, harvesting and other human activities alter aquatic biodiversity locally and globally (6, 7), affecting ecosystem functions and services to humans (8, 9). Seafood, consisting of wild-caught marine and freshwater finfish and invertebrates, not only provides an important source of protein and calories to human consumers, but also provides essential micronutrients, such as vitamins and minerals (1, 2, 10, 11). Though it is appreciated that biodiversity (variety and composition of species) in diets can lead to a more nutritious or beneficial diet overall (3, 12–18), aquatic biodiversity *per se* is only nutritionally valuable if consuming a more diverse seafood diet is more beneficial than simply consuming more seafood biomass. Understanding the value of biodiversity - distinct from total biomass - is a critical challenge as patterns of biodiversity continue to change in the Anthropocene.

Ecological theory predicts that biodiversity can be ecologically and economically important, apart from the importance of total biomass or the presence of particular species (4, 19). Experiments and theory have shown that for a given species pool and ecological context (e.g., prairie grasses, crops or intertidal seaweeds) diversity in ecological communities and agricultural systems enhances ecosystem functioning by two mechanisms: (i) more diverse assemblages may outperform less diverse assemblages of the same biomass because more diverse assemblages are more likely to include high performing species (a *selection effect*), or (ii) more diverse assemblages contain species with complementary functional traits that are not correlated or are correlated negatively among species, allowing them to function more efficiently (a *complementarity effect*) (20, 21). For aquatic animals, increased diversity enhances productivity of fish biomass (22) and also enhances temporal stability of biomass production and total yields

(22–24), providing economic and nutritional benefits to humans related to increased stability of

harvests and production of biomass for consumption (23). When considering aquatic species

from the perspective of human consumers however, functions other than biomass production

become relevant. In addition to providing an important source of calories and protein in the

human diet, seafood is increasingly recognized for its value in providing micronutrients, which

are often scarce in the human diet (25, 26) but critical for human health. Total seafood biomass

yields are not predictive of micronutrient benefits, suggesting that micronutrient content does not

increase proportionally with community biomass or total seafood intake (26).

Relationships between biodiversity (broadly defined to include abundance and biomass of

species) and ecosystem function have been extended to ecosystem services, which include the

cultural, economic or health benefits people derive from nature (5, 8, 27). Though there is clear

evidence that certain species, such as salmon, provide important ecosystem services to humans

(28), here we are interested in whether aquatic biodiversity *per se* (distinct from the identities

and abundance of species) provides ecosystem services in the form of human nutrition, beyond

total yields. For biodiversity *per se* (as opposed to increasing total seafood consumption) to

enhance nutritional benefits as predicted by biodiversity-ecosystem functioning theory,

micronutrient profiles must differ across species and their concentrations must be either

uncorrelated or negatively correlated (21). Specifically, a 'biodiversity effect' (sensu (21)) of

nutritional benefits requires that some seafood species contain high concentrations of some

micronutrients while other species contain high concentrations of different micronutrients (i.e.

nutrient concentrations are negatively correlated or uncorrelated across species), creating a

complementary distribution of micronutrients across species. In contrast, if key micronutrients

are positively correlated across species such that, for example, a species containing high amounts

of iron also has a calcium concentration, diversity in the diet would not be important. In the case

of positive correlations among micronutrients, the ecosystem service of nutritional benefits

would be enhanced by consuming more fish biomass or by selecting a few highly nutritious

species, rather than by biodiversity *per se*.

Differences in species' micronutrient content (29), if complementary and not positively

correlated across species, could lead to positive effects of seafood diversity for human nutritional

well-being. Furthermore, because micronutrient content does not appear to be correlated with

protein or calorie content (26), there is no reason to expect that a positive effect of diversity on

biomass or protein would imply a similar relationship between diversity and nutritional benefits.

We hypothesized that aquatic biodiversity leads to nutritional complementarity in human diets of

seafood, conferring potential benefits for humans via complementarity or selection effects, as

predicted by biodiversity-ecosystem functioning theory. We quantified these benefits using

recommended dietary allowance (RDA) indices, which are nutrient-based reference values that

indicate the average daily dietary intake level that is sufficient to meet the nutrient requirement

of nearly all (97 to 98 percent) healthy individuals in a particular life stage and gender group

(30). Here we used the RDA for females aged 19-50 (SI Appendix Table S1). We consider

nutritional benefits provided by seafood as how much nutrition is provided in a single standard

amount of seafood (such as a 100g portion of seafood). We ask whether consuming a more

diverse set of species actually increases the nutritional content of a potential seafood diet.

Though it is possible to consider alternative ways to increase seafood derived nutrition in a diet,

such as increasing total seafood biomass consumed or increasing consumption of certain species,

these approaches do not necessarily implicate biodiversity as an important element of the

ecosystem service. Furthermore, from an economic or cultural perspective, increasing total

seafood consumption is also not always feasible or desirable, especially if seafood is costly or

only forms a small part of meals.

We tested the hypothesis that aquatic biodiversity confers nutritional benefits through

complementarity in nutrient concentrations among species. We quantified nutritional benefits in

two ways: 1) the number of distinct nutritional needs met in a single 100g portion (referred to as

'nutritional diversity', N_D) and 2) the amount of seafood tissue required to meet a given

nutritional threshold (referred to as 'nutritional efficiency', N_E). We tested whether 1) aquatic

biodiversity enhances nutritional diversity (N_D) in potential seafood diets and 2) biodiversity

increases the nutritional content of an edible portion of seafood, thereby improving the efficiency

(N_E) with which seafood consumers reach nutritional targets (Fig 1). We predicted that increased

species richness (one measure of biodiversity) in potential seafood diets yields increased

nutritional benefits, and that variation in nutrient concentrations among species is related to

species' ecological traits. In a global analysis of 430 aquatic species, we considered provision of

nutritional benefits to human consumers accessing worldwide seafood markets. To assess the

extent to which the relationships between biodiversity and human nutrition benefits depend on

spatial scale (8), we tested whether seafood biodiversity is associated with higher nutritional

benefits at local scales in fourteen traditional indigenous seafood diets in North America. Fish is

critical for indigenous groups, who on average consume fish at a rate of 15 times higher than the

global average (31). We quantified variation in nutrient concentrations in edible tissues of 430

commonly consumed aquatic species in the global species pool, and 25 – 57 species in fourteen

local dietary species pools. We extend biodiversity-ecosystem functioning theory to a

multivariate ecosystem service (32) that is defined from the perspective of the human beneficiary

and is directly relevant to human health.

Fig 1. Aquatic biodiversity increases human well-being because edible species have distinct and
complementary multi-nutrient profiles (A) and differ in mean micro- and macronutrient content
(shown here relative to 10% and 25% thresholds of recommended dietary allowance (RDA)
guidelines for representative finfish (*Abramis brama*, *Mullus surmuletus*), mollusc (*Mytilus
galloprovincialis*) and crustacean species (*Nephrops norvegicus*). Biodiversity – ecosystem
functioning theory predicts that nutritional benefits, including nutritional diversity (N_D; i, iii) and
nutritional efficiency (N_E; ii, iv) (B, E), are enhanced with increasing dietary species richness.
Seafood consumers with limited access to seafood each day may not reach RDA targets if diets
are low in diversity (F vs C; orange shading indicates proportion of population that meets
nutrient requirements). DHA: docosahexaenoic acid, EPA: eicosapentaenoic acid.

Results and Discussion

Biodiversity and nutritional benefits of edible aquatic species

We found that for a given amount of seafood, biodiversity not only enhances nutritional benefits

for consumers selecting seafood from our global species dataset, but that biodiversity *per se* is

essential to meeting nutritional targets. To assess the contribution of species diversity and traits

to the ecosystem service of promoting human well-being, we assessed species' protein, fat, and

micronutrient concentrations in edible tissues of a standard amount - 100g - relative to a measure

of requirements for these nutritional elements set by public health organizations - the

recommended dietary allowance (RDA) targets ((30); Methods). The global species pool was

highly diverse with regard to concentrations of the micronutrients iron, zinc, calcium, and two

fatty acids DHA and EPA in edible fish tissue relative to RDA targets for those micronutrients

(Fig 2; micronutrient geometric CVs: $\ln(\text{iron}) = 3.97$, $\ln(\text{calcium}) = 3.25$, $\ln(\text{EPA}) = 2.52$,

$\ln(\text{zinc}) = 2.10$, $\ln(\text{DHA}) = 1.70$). There was very little diversity in species' protein

concentrations relative to RDA targets (Fig 2; $\ln(\text{protein})$ geometric coefficient of variation

(geometric CV) = 0.03). Most species did not meet a single micronutrient RDA threshold: when

we considered each micronutrient separately we found that fewer than half of the 430 species

reached a single-nutrient threshold of 10% of the daily RDA target (30) for calcium, iron and the

essential fatty acid EPA in a standard 100g portion of a single species (Table S2).

Fig 2. Variation in nutrient concentrations differs among taxonomic groups. (A) Frequency of
reported protein, fat and micronutrient (including eicosapentaenoic acid (EPA), docosahexaenoic
acid (DHA)) content in 100 g of the edible portion of 430 seafood species (references in Dataset
S1). (B) Proportion of species, and number shown on each bar, with available data that reach
10% of RDI targets for any one, two or up to five of the micronutrients examined here.

Nutritional traits covary with ecological traits

Substantial variation in nutrient concentrations in edible portions among species can be explained partly by major ecological attributes and traits: taxonomic group, latitude, body size, diet breadth and feeding habits (Tables S3-S7). Finfish, crustaceans and molluscs differed significantly in their multi-nutrient profiles (PERMANOVA, $F_{2,103} = 3.429$, $p = 0.006$). Among finfish, concentrations of calcium, iron and zinc in edible tissue decreased with increasing body size (Fig 3, negative slopes, $p < 0.05$). Variation in protein and fat was poorly explained by species' ecological traits ($R^2 = 0.023$, 0.09, for protein and fat). In addition to ecological traits, finfish species that are eaten whole, or whose edible portions include organs such as skin, liver or bones, have higher nutrient concentrations in the edible portion than those whose edible portions are restricted to muscle tissue ($R^2 = 0.60$, $F_{5, 251} = 76.24$, $p < 0.01$; Fig S3). Nutrient concentrations were typically weakly negatively correlated or uncorrelated with each other among species (Fig S4), allowing complementarity among species to increase nutritional benefits.

211
Fig 3. Micronutrient concentrations are significantly negatively related to body size (length, cm)
in finfish. Negative slopes from phylogenetic least squares: calcium: -0.63 (95% CI -0.94, -0.33,
n = 89), iron: -0.31 (95% CI -0.61, -0.0075, n = 92), zinc: -0.33 (95% CI -0.52, -0.13, n = 86).

Aquatic biodiversity effects on nutritional benefits to humans

To quantify the effect of biodiversity on nutritional benefits, we first quantified the amount of

seafood, in grams, that would be required to reach a given nutrient intake target (10% of RDA,

Table S1) at each of ten levels of species richness (nutritional efficiency, N_E, for which lower

values signify higher nutrition benefits to consumers). We then estimated the biodiversity effect

using Equation 1, in which b_E is the scaling coefficient that describes how function (i.e. N_E)

varies with species richness. As species richness increased in potential diets, 10% RDA for any

micronutrient was achieved with less total seafood intake (Fig 4A, b_E < 0 for every

micronutrient: calcium -0.43 (95% CI -0.47, -0.40), iron -0.40 (95% CI -0.43, -0.36), zinc -0.21

(95% CI -0.21, -0.23), EPA -0.25 (95% CI -0.26, -0.24) and DHA -0.21 (95% CI -0.21, -0.20)).

This indicates that increasing species richness enhanced nutritional efficiency (N_E) in our sample

diets, independent of any changes in the identity of species included, since the diets were

assembled using a random sample of the species pool. All species reached the 10% protein RDA

target, so that there was no benefit of seafood diversity for protein (Fig 4A, b_E = 0.0092 95% CI

0.0086, 0.010). In other words, the ecosystem service of protein provision to diets is adequately

provided by seafood edible biomass, and not by diversity or even species identity.

Fig 4. Aquatic biodiversity enhances nutritional benefits at global and local scales. A) Seafood
species richness improves the efficiency with which human diets can meet 10% of RDA by
reducing the grams of seafood required to meet nutritional thresholds. Nutritional efficiency (N_E)
is shown for five micronutrients and protein separately (points are median values for calcium,
iron, zinc, EPA, DHA and protein, lines show the fit of Equation 1 to the data and shading refers
to 95% CI) as well as for five micronutrients simultaneously (purple line labeled '5
Micronutrients'); estimates for the b_E parameter (± standard error, inset) describe the strength of
the biodiversity effect. B) Increasing dietary species richness sourced from global seafood
markets increases the efficiency with which five micronutrient nutritional targets are reached;
solid circles are median values, as plotted in the purple line in A. C) Increasing species richness

increases the efficiency with which five micronutrient nutritional targets are reached at local and
global scales; shaded areas are 95% confidence intervals about the mean; black lines are the fit of
Equation 1 for each of fourteen traditional indigenous diets in North America, green line is for a
diet sourced from the global seafood market. D) Species richness increases the number of
distinct RDA targets met in a 100g seafood portion (nutritional diversity, N_D) in local and global
seafood diets; lines are the fit of Equation 2, color coding as in C.

Multi-functional biodiversity benefits of aquatic species

We then considered the effects of seafood species richness on the provisioning of multiple

nutrients simultaneously. This is referred to as a multifunctional benefit of biodiversity (32–34),

and takes into account possible trade-offs or correlations among functions, in this case,

concentrations of micronutrients. For some ecosystem services (e.g. water quality or

ecotourism), benefits of biodiversity accumulate when multiple ecosystem functions are

considered simultaneously (32, 35, 36). We treated each micronutrient concentration (relative to

10% RDA for that micronutrient, Table S1) as one ecosystem function. Consistent with

biodiversity-ecosystem functioning theory (37), we found that in the case of a multifunctional

metric of an ecosystem service defined from the human perspective (i.e. multiple micronutrient

targets reached simultaneously), biodiversity benefits for the multifunctional service are greater

than for individual functions (b_E for all five micronutrients simultaneously = -0.49 (95% CI -

0.50, -0.48) vs. single nutrients b_E range from -0.43 (95% CI -0.47, -0.40) for calcium to -0.21

for EPA (95% CI -0.21, -0.20)) that comprise the ecosystem service (Fig 4A). Increasing species

diversity from one to five species in 1000 simulated, resampled hypothetical diets drawing from

our global species pool allowed consumers to meet 10% of RDA for five essential microelements

and fatty acids simultaneously more than twice as efficiently (i.e. a median of 485.83g of tissue

required with one species vs. median of 216.96g of tissue required with five species) (Fig 4B, C).

Then, we assessed the effects of biodiversity when the total seafood biomass is held constant, at

100g, by counting the number of distinct RDA thresholds in a 100g portion (N_D). We also found

positive effects of biodiversity on the number of distinct nutritional functions (10% of RDA

reached) in a single 100g portion (Fig 4D): more diverse diets reached more nutritional targets

(higher N_D) per 100g serving than diets of the same fish biomass comprising fewer species ($b_D =$

0.21 (95% CI: 0.18-0.24) Fig 4D). These results demonstrate a benefit of biodiversity for human

nutritional well-being, over and above the benefits of consuming a particular amount (biomass)

or identity of aquatic species.

Our analysis provides new and robust evidence that biodiversity is critical to multifunctionality

of ecosystem services when function thresholds are grounded *a priori* in multivariate metrics

meaningful for human well-being such as RDA. Our approach overcomes the critique that

multifunctionality is not enhanced by biodiversity (37), but rather a statistical artifact of how

multifunctionality was commonly estimated. Our findings are robust to multiple RDA threshold

levels (Fig S1), and the biodiversity effect is strongest at a threshold of approximately 28% of

RDA (Fig S1). In this case, the positive effects of biodiversity are greatest at levels of nutritional

benefit that are significant for human nutrition, highlighting the importance of species diversity

in seafood diets. More generally, ecosystem service benefits, as defined in metrics of human

wellbeing rather than the traits of the species pool under consideration (e.g., biomass or stability

of the food web), typically are produced by several underlying ecosystem functions (32). The

strong effects of diversity on multifunctional benefits observed here may also apply to

relationships between diversity and other services e.g., desired filtration rates of pollutants in

wetlands (38), or desired pest consumption rates in agricultural systems (39).

Biodiversity benefits are consistent from local to global scales

Consistent with the positive biodiversity effects we observed when assuming consumers have access to global seafood markets, we also found benefits of seafood diversity in a local context. We analyzed the effects of biodiversity in fourteen traditional indigenous North American diets of seafood harvested locally, and found a consistent, positive effect of biodiversity on N_D and N_E, although the magnitude of the biodiversity effect was generally lower at the local scale than the global scale (Fig 4C-D, Fig S2, S5) (global $b_D = 0.21$ (95% CI 0.18, 0.24) vs mean local $b_D = 0.14 \pm 0.0083$ S.E. and global $b_E = -0.50$ (95% CI -0.52, -0.47) vs mean local $b_E = -0.16 \pm 0.0091$ S.E.). This finding is consistent with lower nutritional functional diversity (mean local $FD = 2.77 \pm 0.17$ S.E. vs. global $FD = 3.87 \pm 0.0096$ S.E.) and higher nutritional functional evenness in local diets (mean local $FEve = 0.76 \pm 0.01$ S.E. vs. global $FEve = 0.71 \pm 0.0018$ S.E.) (Fig S2, S5), suggesting that functional consequences of changes to diversity in local seafood diets may be buffered by higher redundancy among species. Given increasing trends towards homogenization of the global food supply (40) including aquaculture (41, 42), this local-scale finding highlights the importance of local species diversity in the diets of vulnerable populations (26).

Potential benefits of aquatic biodiversity for human well being

The problem of 'hidden hunger', in which people have access to sufficient calories but insufficient essential micronutrients, is becoming increasingly recognized as a major contributor to the burden of disease in developing countries (43, 44). Indeed, more than two billion people suffer from micronutrient deficiencies, and it is the aim of the United Nations Sustainable Development Goals (SDGs) to end all forms of malnutrition (44). Micronutrients in fish lead to a

variety of health benefits, including lowered risk of cardiovascular disease, improved maternal

health and pregnancy outcomes, increased early childhood physical and cognitive development.

Polyunsaturated fatty acids such as EPA and DHA are important for cardiovascular and brain

health. The amount of seafood in the human diet is often limited, either by costs or because

traditionally harvested species are being replaced by other 'junk foods' (45). In light of this, we

have focused here on the nutritional benefits potentially provided by seafood biodiversity over

and above the services provided by consuming seafood in general, and by consuming particular

highly nutritious species. Seafood consumption is influenced by many factors including cultural

preferences, social status, and economic pressures, none of which are considered here. Still, our

study adds to our knowledge of seafood and nutrition by quantifying the contribution of

biodiversity *per se* to meeting nutritional needs, and this biodiversity effect occurs in addition to

how diversity enhances and stabilizes total yields (22). This finding bridges the growing

understanding of hidden hunger and food security with the large theoretical and empirical

understanding of relationships between biodiversity, ecosystem function and benefits to people.

Aquatic biodiversity and food security in a changing world

Maintaining the diversity of global fisheries is important for ensuring food and nutrition security.

A diverse seafood species pool feeds not only local communities but also seafood markets

worldwide, and aquatic species contain micronutrients not found in other foods. Many of the

most nutritionally vulnerable populations – those that are deficient in essential micronutrients

during particularly sensitive stages of life (i.e. pregnancy, breastfeeding and childhood) may rely

on local ecosystems to meet their nutritional demands (2, 25, 29). These populations may have

access to a limited amount of locally available fish tissue each day or to fish from a subset of

habitat types, suggesting that for these populations nutritional efficiency may be particularly

important. In tropical regions, fish diversity in coastal regions has plummeted in recent decades

(46) characterized by two-fold declines in body sizes of fish (47, 48). These regions are also

regions of high nutritional vulnerability and reliance on locally harvested seafood (49). However,

as the seafood trade becomes increasingly global (50, 51), seafood-derived nutrition available to

consumers participating in the global market may be related to globally harvested seafood

biodiversity. As a result, changes to local biodiversity and resultant impacts on human nutrition

may be buffered by access to global seafood markets. Together, our results suggest that in the

context of global change, understanding and protecting the potential for nature to support diverse

and productive aquatic ecosystems has direct and immediate benefits to humanity.

Conclusions

Nutritional diversity appears to be derived from ecological diversity, suggesting links between

the complexity of aquatic ecosystems and their capacity to produce nutritional benefits. While

the role of seafood is well recognized as an important source of protein in the human diet, the

role of seafood as a source of essential micronutrients is often overlooked. This study provides

the first global assessment of patterns in the macro- and micronutrient content of aquatic species.

Our results reveal that not all fish species are equally nutritionally valuable to humans and that

aspects of ecological structure, including species and trait diversity enhance nutritional diversity,

thereby linking the processes that structure ecosystems with the benefits they provide.

Methods

Quantifying nutritional value in terms of human health benefits

We characterized an aquatic species' nutritional profile by drawing on two well-established

nutritional metrics: nutrient concentration (nutrient content/100g edible portion) and

Recommended Dietary Allowances (RDA). RDAs are developed following health guidelines,

and quantify the recommended amount of a particular nutrient required to maintain health (30).

The RDAs used here were established by the Food and Nutrition Board of the United States

Institute of Medicine, and quantify the daily intake level of a nutrient needed to meet the

requirements of 97–98% of healthy adults (females aged 19-50) (30). We defined the nutritional

value of a fish species for each nutrient in terms of the nutrient content in an edible portion

relative to the RDA for that nutrient. For some aquatic species, micronutrients are in such high

concentrations in edible tissues that a single 100g portion contains the entire RDA for that

nutrient, while edible tissues from other species provide only small fractions of the RDA. We

considered thresholds of nutritional benefit (function) between 1% and 100% of the RDA,

although we focus on 10% of RDA since it is a minimum threshold for a food to be considered

of nutritional benefit (30). We defined an RDA target as 10% of RDA for a given nutrient. We

characterized nutritional profiles in two ways: first, in terms of concentrations of each nutrient

(single nutrient profile), and secondly, in terms of multiple nutrients simultaneously (i.e. multi-

nutrient profile).

Defining nutritional benefits

We quantified the effect of biodiversity on nutritional benefits in two ways (Fig 1): 1) nutritional

efficiency, N_E, which quantifies the amount of tissue, in grams, required to reach a given number

of RDA targets for different nutrients simultaneously and 2) nutritional diversity, N_D, which

quantifies the number of distinct RDA thresholds for different nutrients met in a standard 100g

edible portion. Nutritional benefit increases with decreasing values of N_E, since N_E quantifies the

grams required to reach RDA targets, and fewer grams required is better from the perspective of

human nutrition. We quantified N_D in an arbitrary daily diet, assuming that the seafood diet

contains 100g of seafood per day.

Literature search and data collection

We assembled a dataset of published nutrient concentrations in edible portions of 430 aquatic

species. We analyzed tissue concentrations of nutrients for which RDA standards exist and that

are implicated in a range of biologically important processes that affect organismal growth and

reproduction, and therefore may potentially relate ecological function with human nutritional

wellbeing. We examined macronutrients including protein and fat, as well as five micronutrients:

metals beneficial at low concentrations but toxic at high concentrations (zinc and iron), one

beneficial mineral (calcium) and the polyunsaturated fatty acids eicosapentaenoic acid (EPA)

and docosahexaenoic acid (DHA). We aimed to include as many marine and freshwater species

as possible covering a wide geographic extent. We searched peer-reviewed literature for

analytical food composition values as well as the Food and Agriculture Organization's Global

Food Composition Database for Fish and Shellfish (52). For finfish, we restricted our analysis to

include only edible portions of wild, raw fish (excluding prepared or farmed seafood items). We

included both farmed and wild mollusc species because mollusc farming does not typically

involve additional food inputs, which could influence tissue nutrient composition. For each

species, we noted which body parts are included in the edible portion and season of collection.

For each sample, we noted the location of origin (e.g. latitude and longitude). To address

inconsistencies in fatty acid data reporting, we standardized fatty acid measurements using the

fatty acid conversion factors proposed by Nowak et al. 2014 (53). When there were multiple

observations available for a single species, we averaged nutrient concentrations across the

observations. We did not include data from national food composition tables because these data

usually report seafood data with a generic food description, which does not allow for a clear

description of which fish tissues are included in the edible portion. For each species with

nutritional data, we collected ecological trait information from FishBase (54) and SeaLifeBase

(55). We included body size (maximum length), fractional trophic position, temperature

preference (using latitude as a proxy), diet breadth and feeding mode.

Statistical analyses and hypothesis testing

Hypothesis 1: Biodiversity enhances nutritional benefits

We tested the effect of species diversity (quantified as the number of species, or species richness,

S), on nutritional diversity and efficiency by randomly assembling diets from the global seafood

species pool, keeping the portion size constant (100 grams) across all levels of diversity. In

analyses using the global dataset (430 species), we assumed that human populations have access

to the entire global species pool and choose species at random. Though this assumption certainly

ignores economic, social and cultural factors that affect which species people consume, in the

absence of detailed diet information for the majority of the world's populations this is a

necessary assumption. To assess potential effects of biodiversity on nutritional benefits for

populations that consume seafood locally and not as part of the global seafood market, we

sampled diets from species contained within traditional diets in fourteen indigenous cultures in

North America. We used species lists for local diets that were obtained from an ethnographic

database of traditional animal foods of indigenous peoples of northern North America, and only

included species that were harvested wild and whose nutrient compositions were analyzed when

raw (Table S7, (56)). To avoid confounding differences in biodiversity effects at global and local

scales with the sizes of the species pools at each of these scales, we matched the size of the

global species pool to the average size of the local species pools (40 species).

We calculated two metrics of nutritional benefits for hypothetical diets comprising species drawn

randomly from either the global dataset (global diet) or local datasets (local diets):

1) Nutritional efficiency (N_E). We tested the hypothesis that complementarity in nutrient

concentrations among species enhances N_E by estimating the effect of species richness on N_E (the

'biodiversity effect'). To estimate the 'biodiversity effect' on N_E, we took an approach to

modeling dietary species composition that is analogous to a biodiversity-ecosystem function

experiment with a replacement design, where species abundances decline proportionally as

species richness increases. From the global species pool, we sampled ten species at random and

then assembled seafood diets from all possible combinations of these ten randomly chosen

species at 10 levels of species richness (1-10). We repeated this process of sampling ten species

from the global species pool and then assembling all possible diets 1000 times. For each

combination of species in each dietary diversity level (1-10 species), we calculated the number

of grams required to reach a given RDA target (either: one of six possible nutrient targets

individually, or five micronutrient (calcium, iron, zinc, EPA and DHA) targets simultaneously).

We quantified the effect of species richness in a diet on nutritional efficiency, N_E, by fitting a

power function to these bootstrapped nutritional efficiency estimates:

$$N_E = aS^{b_E} \tag{1}$$

where the parameter b_E describes the relationship between a change in species richness, S, and a

change in N_E, and a is a constant (in units of grams). Since N_E is measured in grams required to

reach a given RDA target, and fewer grams required is better from the perspective of human

nutrition, then a benefit of biodiversity would be reflected in a negative b_E (i.e. N_E, measured in

grams of tissue required, decreases with species richness). For each nutrient individually, and for

all five micronutrients together, we estimated the exponent parameter, b_E, using non-linear

regression using the *nls.LM* function in the *minpack.lm* package in R (57). To quantify

uncertainty in parameter estimates associated with sampling from the pool of observed nutrient

content values, we calculated bootstrapped confidence intervals using non-parametric

bootstrapping of mean centered-residuals using the *nlsBoot* function in the R package *nlstools*

(58).

2) Nutritional diversity (N_D). To test the hypothesis that complementarity in nutrient

concentrations among species increases nutritional benefits by increasing the number of distinct

RDA targets in a 100g portion, we constructed recommended dietary allowance (RDA) target

accumulation curves. These are analogous to species accumulation curves used in ecological

studies to assess patterns of beta-diversity, or species turnover, in ecological community

composition data. We assessed turnover of nutrients content in edible tissues among fish species.

Each fish species is associated with a set of 0s and 1s corresponding to whether or not it achieves

10% of RDA for each of five micronutrients (equivalent to a species presence-absence matrix in

community composition data), sampled with replacement 1000 times. This approach allowed us

to explore how likely it would be for human diets containing different numbers of fish species to

reach a given number of micronutrient RDA targets (N_D ranges between 0 and 5), assuming that

fish species were included in the human diet at random. We quantified the effect of biodiversity

on nutritional diversity, N_D, by fitting a power function,

481 $$N_D = aS^{b_D}$$ (2)

where the parameter b_D describes the relationship between a change in species richness, S, and a

change in N_D (i.e. the number of RDA targets reached per average 100g portion), and a is a

constant.

For both N_E and N_D, we tested the hypothesis that biodiversity enhances nutritional benefits by

assessing whether the estimate of the scaling exponent, b, had confidence intervals not

overlapping zero. We concluded that biodiversity enhanced nutritional benefits if b_E was

negative and b_D was positive.

We tested the hypothesis that nutritional functional diversity drives positive biodiversity effects

by testing for a positive association between nutritional functional diversity and the biodiversity

scaling exponent, b. We hypothesized that nutritional functional diversity would be higher at the

global scale than the local scale, because the global species pool contains more ecological and

biogeographic diversity. To assess levels of nutritional functional diversity among species, we

calculated functional diversity (*FD*) (59). *FD* is based on an assessment of the entire functional

diversity of a group represented as a functional dendrogram, and *FD* allows estimation of

complementarity among species' nutrient concentrations (i.e. nutritional functional traits) using

the dendrogram. We treated the concentration of each micronutrient (calcium, iron, zinc, EPA

and DHA) as a functional trait. We also quantified functional evenness metric (*FEve*) using the

FD package in R, which normally quantifies the evenness of abundance in a functional trait

space. Here, we used *FEve* to quantify the evenness in concentration of nutrients across species

(60). To compare *FD* and *FEve* at the global and local scales, we first subsampled 40 species

(the average species pool at the local scale) from the global pool, then calculated the functional

diversity metrics on the subsample, and repeated this process 1000 times. Using this same

approach, we calculated levels of 'expected' *FD* and *FEve* for each local diet by choosing

random subsets of the global pool with sample size equal to the species pool in each local diet,

and repeated this process 1000 times (Fig S5).

Hypothesis 2: Diversity in nutrient concentrations of the edible portion is related to

ecological diversity estimated as species' ecological traits.

We tested the hypothesis that nutrient concentrations are related to species' ecological traits in

two ways: 1) testing whether multi-nutrient profiles (i.e. concentrations of all five

micronutrients) differ among major phylogenetic groups and 2) whether differences in single

nutrient concentrations differ with species ecological traits. We examined variation in multi-

nutrient profiles among seafood species using the *vegan* package in R (61). We ln transformed

nutrient concentration data to achieve normality. Differences in multi-nutrient profiles were

visualized through non-metric dimensional scaling (NMDS) using the *metaMDS* function. The

ordination ran for 1000 iterations, and the stress score of 0.032 for the final solution was

sufficiently low to enable reliable interpretation in the two dimensions (62). First, we tested the

hypothesis that major phylogenetic groups correlated with functional differences in life history,

resource use and ecology (i.e. finfish, mollusc, and crustacean) differ in their multi-nutrient

profiles via permutational multivariate ANOVA (PERMANOVA) using the *adonis* function (999

permutations) based on Bray-Curtis dissimilarity matrices. We used an overall (three-way)

PERMANOVA first to investigate phylogenetic group effects on nutrient profile of a species'

edible tissue.

To test for associations between species' ecological functional traits and their nutrient

concentrations, we modeled the relationship between traits and ln(nutrient concentration) with

phylogenetic least squares regression (PGLS). The full model included the entire set of trait

predictors as fixed effects:

$ln(nutrient) = \beta_0 + \beta_1 \times ln(body\ size) + \beta_2 \times latitude + \beta_3 \times trophic\ position + \beta_4 \times feeding\ mode +$

$\beta_5 \times diet\ breadth + \varepsilon$

Unlike in ordinary least squares (OLS), which assumes there is no covariance structure in the

error term, ε, (all species are independent from one another, and residuals from closely related

species are not more similar on average than residuals from distantly related species), PGLS

assumes that the residuals are non-independent, and that the expected covariance is related to the

shared evolutionary history between the species. To assess whether the relationships between

species' traits and their nutritional profiles were associated, we fit multiple regression models

using PGLS using the *gls* function in the *nlme* package in R using the Brownian motion model of

evolution. This approach allowed us to account for phylogenetic non-independence by using

shared ancestry as weights on the elements of the residual variance-covariance matrix used in the

model. We created a supertree by combining phylogenies that included the species of finfish,

molluscs and crustaceans in our nutrient dataset using the *rotl* package in R (63), which is an

interface to the Open Tree of Life (64). We computed branch lengths according to taxonomic

depth (65) using the *compute.brlen* function in the *ape* package in R (66).

We created models from subsets of the full model that represented hypotheses based on the

known physiological roles of micronutrients and their relationships to our set of predictors

(Tables S3-S7, S10). To avoid issues associated with multicollinearity of predictor variables, we

excluded other possible variables if they were highly correlated (i.e. correlation coefficient >

0.6). We identified the best subset of models using the Akaike Information Criterion, adjusted

for small sample sizes (AICc). We used AICc, ∂aic and Akaike weights (*w*) to compare models.

We ranked models based on *w*, and selected the set of models that produced a cumulative w >

0.95, meaning that we are 95% confident that the chosen set includes the best model (67). In

cases where we could not obtain measurements of all traits for all species, we performed model

selection on reduced datasets without missing values.

To account for model uncertainty in the ecological trait correlation analyses, we performed

model averaging of coefficients in all models that produced a cumulative $w > 0.95$, and included

zeros as coefficients when variables did not enter a given model (67). We conducted our model

selection and averaging analyses with the *MuMIn* package (68) and all other analyses in R

version 3.3.2 (69).

Uncertainties:

There are several sources of uncertainty in our analyses. First, there are substantial sources of

uncertainty in food composition estimates. The data in our dataset meet international standards

for data quality and standardization, meaning that we followed guidelines for checking food

composition data and converting units, denominators and expressions (70). Still, tissue

concentrations may vary depending on analytical techniques, labs, season, diet of the animal, life

stage etc. Some of these sources of uncertainty (e.g. differences in analytical techniques) are

unavoidable consequences of synthesizing previously published data collected across many labs.

We assumed that these uncertainties in the data were randomly distributed over our

geographically and taxonomically diverse dataset. Further uncertainty is associated with how

well our set of 430 species represents the global pool of seafood consumed. We do not know

whether our sample is random or biased, though we can say that our dataset includes 41 of the 67

most consumed species worldwide (as determined by FAO production volumes (71), species

with capture production of 150 000 tonnes or more, after removing species for which the

majority of production volume is diverted to fish meal and oil (72), Table S9). A remaining

source of variation among samples is likely due to natural sources of variation associated with

seasonal and other sources of temporal variability, which we consider to be an important

component of biodiversity.

Acknowledgments: We thank P. Thompson, A. Gehman, W. Cheung, M. Ruckelshaus and C.
Harley and M. Deith for helpful comments on previous versions of this manuscript.
589
590 **Funding:** Funding was provided by a Vanier Canada Graduate Scholarship to JRB.
591
592 **Competing interests:** We declare no competing interests.
593
594 **Data and materials availability:** All data and code will be archived at the Dryad digital
595   repository.

References

1.   M. C. Latham, *Human nutrition in the developing world* (Food & Agriculture Org., 1997).

2.   C. D. Golden, *et al.*, Fall in fish catch threatens human health. *Nature* **534**, 317–320 (2016).

3.   D. Penafiel, C. Lachat, R. Espinel, P. Van Damme, P. Kolsteren, A systematic review on
the contributions of edible plant and animal biodiversity to human diets. *EcoHealth* **8**, 381–
399 (2011).

4.   B. J. Cardinale, *et al.*, Biodiversity loss and its impact on humanity. *Nature* **486**, 59–67
(2012).

5.   S. S. Myers, *et al.*, Human health impacts of ecosystem alteration. *Proc. Natl. Acad. Sci. U.*
S. A. **110**, 18753–18760 (2013).

6.   R. Elahi, *et al.*, Recent trends in local-scale marine biodiversity reflect community structure
and human impacts. *Curr. Biol.* **25**, 1938–1943 (2015).

7.   M. L. Pinsky, B. Worm, M. J. Fogarty, J. L. Sarmiento, S. A. Levin, Marine taxa track local
climate velocities. *Science* **341**, 1239–1242 (2013).

8.   F. Isbell, *et al.*, Linking the influence and dependence of people on biodiversity across
scales. *Nature* **546**, 65–72 (2017).

9.   M. I. O'Connor, *et al.*, A general biodiversity--function relationship is mediated by trophic
level. *Oikos* **126**, 18–31 (2017).

10.   B. Belton, S. H. Thilsted, Fisheries in transition: Food and nutrition security implications
for the global South. *Global Food Security* **3**, 59–66 (2014).

11.   FAO, *The State of World Fisheries and Aquaculture 2018 - Meeting the sustainable*
development goals (2018).

12.   A. D. Jones, Critical review of the emerging research evidence on agricultural biodiversity,
diet diversity, and nutritional status in low- and middle-income countries. *Nutr. Rev.* **75**,
769–782 (2017).

13.   C. Lachat, *et al.*, Dietary species richness as a measure of food biodiversity and nutritional
quality of diets. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 127–132 (2018).

14.   R. Remans, *et al.*, Assessing nutritional diversity of cropping systems in African villages.
PLoS One **6** (2011).

15.   F. A. J. DeClerck, J. Fanzo, C. Palm, R. Remans, Ecological approaches to human nutrition.
Food Nutr. Bull. **32**, S41–50 (2011).

16.   M. T. Ruel, Operationalizing dietary diversity: a review of measurement issues and research

priorities. *J. Nutr.* **133**, 3911S–3926S (2003).

17. FAO, Compendium of Indicators for Nutrition-Sensitive Agriculture (2013).

18. J. Fanzo, D. Hunter, T. Borelli, F. Mattei, *Diversifying Food and Diets: Using Agricultural*
Biodiversity to Improve Nutrition and Health (Routledge, 2013).

19. M. Loreau, N. Mouquet, A. Gonzalez, Biodiversity as spatial insurance in heterogeneous
landscapes. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 12765–12770 (2003).

20. M. Loreau, A. Hector, Partitioning selection and complementarity in biodiversity
experiments. *Nature* **412**, 72–76 (2001).

21. M. Loreau, Biodiversity and ecosystem functioning: recent theoretical advances. *Oikos* **91**,
3–17 (2000).

22. J. E. Duffy, J. S. Lefcheck, R. D. Stuart-Smith, S. A. Navarrete, G. J. Edgar, Biodiversity
enhances reef fish biomass and resistance to climate change. *Proc. Natl. Acad. Sci. U. S. A.*
113, 6230–6235 (2016).

23. R. Hilborn, T. P. Quinn, D. E. Schindler, D. E. Rogers, Biocomplexity and fisheries
sustainability. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 6564–6568 (2003).

24. S. Wang, M. Loreau, Biodiversity and ecosystem stability across scales in
metacommunities. *Ecol. Lett.* **19**, 510–518 (2016).

25. N. Kawarazuka, C. Béné, Linking small-scale fisheries and aquaculture to household
nutritional security: an overview. *Food Security* **2**, 343–357 (2010).

26. J. R. Bogard, *et al.*, Higher fish but lower micronutrient intakes: Temporal changes in fish
consumption from capture fisheries and aquaculture in Bangladesh. *PLoS One* **12** (2017).

27. P. Balvanera, *et al.*, Linking Biodiversity and Ecosystem Services: Current Uncertainties
and the Necessary Next Steps. *Bioscience* **64**, 49–57 (2014).

28. D. E. Schindler, *et al.*, Pacific salmon and the ecology of coastal ecosystems. *Front. Ecol.*
Environ. **1**, 31–37 (2003).

29. J. R. Bogard, *et al.*, Nutrient composition of important fish species in Bangladesh and
potential contribution to recommended nutrient intakes. *J. Food Compost. Anal.* **42**, 120–
133 (2015).

30. Institute of Medecine, *Dietary Reference Intakes for Calcium and Vitamin D* (National
Academies Press, 2012; http://nationalacademies.org/hmd/~/media/Files/Activity
Files/Nutrition/DRI-Tables/2_ RDA and AI Values_Vitamin and Elements.pdf?la=en), vol.
130.

31. A. M. Cisneros-Montemayor, D. Pauly, L. V. Weatherdon, Y. Ota, A Global Estimate of

Seafood Consumption by Coastal Indigenous Peoples. *PLoS One* **11**, e0166681 (2016).

32. P. Manning, *et al.*, Redefining ecosystem multifunctionality. *Nature Ecology & Evolution* **2**,
427–436 (2018).

33. J. E. K. Byrnes, *et al.*, Investigating the relationship between biodiversity and ecosystem
multifunctionality: Challenges and solutions. *Methods in Ecology and Evolution* **5**, 111–124
(2014).

34. M. E. Mastrangelo, *et al.*, Concepts and methods for landscape multifunctionality and a
unifying framework based on ecosystem services. *Landsc. Ecol.* **29**, 345–358 (2014).

35. A. Hector, R. Bagchi, Biodiversity and ecosystem multifunctionality. *Nature* **448**, 188–190
(2007).

36. L. Gamfeldt, H. Hillebrand, P. R. Jonsson, Multiple functions increase the importance of
biodiversity for overall ecosystem functioning. *Ecology* **89**, 1223–1231 (2008).

37. L. Gamfeldt, F. Roger, Revisiting the biodiversity--ecosystem multifunctionality
relationship. *Nature Ecology & Evolution* **1**, s41559–017 (2017).

38. T. Boyer, S. Polasky, Valuing urban wetlands: A review of non-market valuation studies.
Wetlands **24**, 744–755 (2004).

39. D. S. Karp, *et al.*, Forest bolsters bird abundance, pest control and coffee yield. *Ecol. Lett.*
16, 1339–1347 (2013).

40. C. K. Khoury, *et al.*, Increasing homogeneity in global food supplies and the implications
for food security. *Proceedings of the National Academy of Sciences* **111**, 4001–4006 (2014).

41. J. Bostock, *et al.*, Aquaculture: global status and trends. *Philos. Trans. R. Soc. Lond. B Biol.*
Sci. **365**, 2897–2912 (2010).

42. C. M. Duarte, *et al.*, Will the oceans help feed humanity? *Bioscience* **59**, 967–976 (2009).

43. G. Kennedy, G. Nantel, P. Shetty, The scourge of "hidden hunger": global dimensions of
micronutrient deficiencies. *Food Nutrition and Agriculture* **32**, 8–16 (2003).

44. K. von Grebmer A. Saltzman E. Birol D. Wiesmann N. Prasai S. Yin Y. Yohannes P.
Menon J. Thompson A. Sonntag, 2014 Global hunger index: The challenge of hidden
hunger (2014) (September 2, 2019).

45. H. V. Kuhnlein, O. Receveur, Local Cultural Animal Food Contributes High Levels of
Nutrients for Arctic Canadian Indigenous Adults and Children. *J. Nutr.* **137**, 1110–1114
(2007).

46. G. P. Jones, M. I. McCormick, M. Srinivasan, J. V. Eagle, Coral decline threatens fish
biodiversity in marine reserves. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 8251–8253 (2004).

47.   J. G. Molinos, *et al.*, Climate velocity and the future global redistribution of marine
biodiversity. *Nat. Clim. Chang.* **6**, 83–88 (2016).

48.   W. W. L. Cheung, *et al.*, Shrinking of fishes exacerbates impacts of global ocean changes
on marine ecosystems. *Nat. Clim. Chang.* **3**, 254–258 (2013).

49.   E. H. Allison, *et al.*, Vulnerability of national economies to the impacts of climate change
on fisheries. *Fish Fish* **10**, 173–196 (2009).

50.   J. A. Gephart, M. L. Pace, Structure and evolution of the global seafood trade network.
Environ. Res. Lett. **10**, 125014 (2015).

51.   R. A. Watson, B. S. Green, S. R. Tracey, A. Farmery, T. J. Pitcher, Provenance of global
seafood. *Fish Fish* **17**, 585–595 (2016).

52.   Food and Agriculture Organization of the United Nations, "FAO/INFOODS Global Food
Composition Database for Fish and Shellfish Version 1.0- uFiSh1.0" (2016).

53.   V. Nowak, D. Rittenschober, J. Exler, U. R. Charrondiere, Proposal on the usage of
conversion factors for fatty acids in fish and shellfish. *Food Chem.* **153**, 457–463 (2014).

54.   R. A. D. P. Froese, "Fishbase version (06/2017)."

55.   M. L. D. A. D. P. Palomares, "SeaLifeBase version (06/2017)."

56.   H. V. Kuhnlein, M. M. Humphries, Traditional Animal Foods of Indigenous Peoples of
Northern North America (2017).

57.   T. V. Elzhov, K. M. Mullen, A.-N. Spiess, B. Bolker, *minpack.lm: R interface to the*
Levenberg-Marquardt nonlinear least-squares algrothim found in MINPACK, pluss support
for bounds (2013).

58.   F. Baty, *et al.*, A Toolbox for Nonlinear Regression in R : The Package nlstools. *J. Stat.*
Softw. **66**, 1–21 (2015).

59.   O. L. Petchey, K. J. Gaston, Functional diversity (FD), species richness and community
composition. *Ecol. Lett.* **5**, 402–411 (2002).

60.   S. Villéger, N. W. H. Mason, D. Mouillot, New multidimensional functional diversity
indices for a multifaceted framework in functional ecology. *Ecology* **89**, 2290–2301 (2008).

61.   J. Oksanen, *et al.*, Package "vegan." *Community ecology package, version* **2** (2013).

62.   P. Legendre, L. Legendre, *Numerical Ecology* (1998).

63.   F. Michonneau, J. W. Brown, D. J. Winter, rotl : an R package to interact with the Open
Tree of Life data. *Methods Ecol. Evol.* **7**, 1476–1481 (2016).

64.   C. E. Hinchliff, *et al.*, Synthesis of phylogeny and taxonomy into a comprehensive tree of

life. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 12764–12769 (2015).

65. A. Grafen, The phylogenetic regression. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **326**, 119–
157 (1989).

66. E. Paradis, J. Claude, K. Strimmer, APE: Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics* **20**, 289–290 (2004).

67. K. P. Burnham, D. R. Anderson, *Information and likelihood theory: a basis for model
selection and inference* (2002).

68. K. Barton, MuMIn: Multi-model inference. R package version 1.9. 13. *The Comprehensive
R Archive Network (CRAN), Vienna, Austria* (2013).

69. R Core Team, R: A Language and Environment for Statistical Computing (2017).

70. Food and Agriculture Organization/ International Network of Food Data Systems
(INFOODS), "FAO / INFOODS Guidelines for Checking Food Composition Data prior to
the publication of a User Table / Database - Version 1 . 0" (2012).

71. Food and Agriculture Organization of the United Nations, "FAOSTAT. Calculated from
food balance sheets" (2016).

72. T. Cashion, F. Le Manach, D. Zeller, D. Pauly, Most fish destined for fishmeal production
are food-grade fish. *Fish Fish* **18**, 837–844 (2017).

Figure legends

Fig 1. Aquatic biodiversity increases human well-being because edible species have distinct and complementary multi-nutrient profiles (A) and differ in mean micro- and macronutrient content (shown here relative to 10% and 25% thresholds of recommended dietary allowance (RDA) guidelines for representative finfish (*Abramis brama, Mullus surmuletus*), mollusc (*Mytilus galloprovincialis*) and crustacean species (*Nephrops norvegicus*). Biodiversity – ecosystem functioning theory predicts that nutritional benefits, including nutritional diversity (N_D; i, iii) and nutritional efficiency (N_E; ii, iv) (B, E), are enhanced with increasing dietary species richness. Seafood consumers with limited access to seafood each day may not reach RDA targets if diets are low in diversity (F vs C; orange shading indicates proportion of population that meets nutrient requirements). DHA: docosahexaenoic acid, EPA: eicosapentaenoic acid.

Fig 2. Variation in nutrient concentrations differs among taxonomic groups. (A) Frequency of reported protein, fat and micronutrient (including eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA)) content in 100 g of the edible portion of 430 seafood species (references in Dataset S1). (B) Proportion of species, and number shown on each bar, with available data that reach 10% of RDI targets for any one, two or up to five of the micronutrients examined here.

Fig 3. Micronutrient concentrations are significantly negatively related to body size (length, cm) in finfish. Negative slopes from phylogenetic least squares: calcium: -0.63 (95% CI -0.94, -0.33, *n* = 89), iron: -0.31 (95% CI -0.61, -0.0075, *n* = 92), zinc: -0.33 (95% CI -0.52, -0.13, *n* = 86).

Fig 4. Aquatic biodiversity enhances nutritional benefits at global and local scales. A) Seafood

species richness improves the efficiency with which human diets can meet 10% of RDA by

reducing the grams of seafood required to meet nutritional thresholds. Nutritional efficiency (N_E)

is shown for five micronutrients and protein separately (points are median values for calcium,

iron, zinc, EPA, DHA and protein, lines show the fit of Equation 1 to the data and shading refers

to 95% CI) as well as for five micronutrients simultaneously (purple line labeled '5

Micronutrients'); estimates for the b_E parameter (± standard error, inset) describe the strength of

the biodiversity effect. B) Increasing dietary species richness sourced from global seafood

markets increases the efficiency with which five micronutrient nutritional targets are reached;

solid circles are median values, as plotted in the purple line in A. C) Increasing species richness

increases the efficiency with which five micronutrient nutritional targets are reached at local and

global scales; shaded areas are 95% confidence intervals about the mean; black lines are the fit of

Equation 1 for each of fourteen traditional indigenous diets in North America, green line is for a

diet sourced from the global seafood market. D) Species richness increases the number of

distinct RDA targets met in a 100g seafood portion (nutritional diversity, N_D) in local and global

seafood diets; lines are the fit of Equation 2, color coding as in C.