Blood meal analysis of tsetse flies (*Glossina pallidipes*: Glossinidae) reveals higher host fidelity on wild compared with domestic hosts

Manun Channumsin[1*], Marc Ciosi[2], Dan Masiga[3], Harriet Auty[4], C. Michael Turner[5], Elizabeth Kilbride[4] and Barbara K. Mable[4*]

*Correspondence: manun_wo@rmutto.ac.th, barbara.mable@glasgow.ac.uk.

[1] Faculty of Veterinary Medicine, Rajamangala University of Technology Tawan-Ok, Chonburi, 20110, Thailand

[2] Institute of Molecular, Cell and Systems Biology (MCSB), University of Glasgow, University Place, Glasgow G12 8QQ, UK

[3] International Centre of Insect Physiology and Ecology (ICIPE), P.O. Box 30772, 00100 Nairobi, Kenya

[4] Institute of Biodiversity, Animal Health and Comparative Medicine (BAHCM), Graham Kerr Building, University of Glasgow, University Place, Glasgow G12 8QQ, UK

[5] Institute of Infection Immunity and Inflammation (III), University of Glasgow, University Place, Glasgow G12 8QQ, UK

Keywords: blood meal, tsetse flies, Trypanosomes, parasites, host, host population structure, African trypanosomiasis.

Abstract

Changes in climate and land use can alter risk of transmission of parasites between domestic hosts and wildlife, particularly when mediated by vectors that can travel between populations. Here we focused on tsetse flies (genus *Glossina*), the cyclical vectors for both Human African Trypanosomiasis (HAT) and Animal African Trypanosomiasis (AAT). The aims of this study were to investigate: 1) the diversity of vertebrate hosts that flies fed on; 2) whether host feeding patterns varied in relation to type of hosts, tsetse feeding behaviour, site or tsetse age and sex; and 3) if there was a relationship between trypanosome detection and host feeding behaviours or host types. Sources of blood meals of *Glossina pallidipes* were identified by sequencing of the mitochondrial cytochrome b gene and analyzed in relationship with previously determined trypanosome detection in the same flies. In an area dominated by wildlife but with seasonal presence of livestock (Nguruman), 98% of tsetse fed on single wild host species, whereas in an area including a mixture of resident domesticated animals, humans and wildlife (Shimba Hills), 52% of flies fed on more than one host species. Multiple Correspondence Analysis revealed strong correlations between feeding pattern, host type and site but these were resolved along a different dimension than trypanosome status, sex and age of the flies. Our results suggest that individual *G. pallidipes* in interface areas may show higher feeding success on wild hosts when available but often feed on both wild and domesticated hosts. This illustrates the importance of *G. pallidipes* as a vector connecting the sylvatic and domestic cycles of African trypanosomes.

Introduction

In sub-Saharan Africa, changes in land use increase encroachment of domestic livestock into areas that are primarily managed to conserve wildlife. This increases risks that livestock will be exposed to a wider range of parasites, with potentially important consequences for disease burden and control. Wildlife can represent 'reservoir communities[31,71] for multi-host pathogens that could spill-over into domesticated animals. Domesticated animals infected by wildlife pathogen could in turn show more severe disease, given limited opportunity for host-pathogen coevolution in novel hosts. This could be particularly true for vector-mediated transmission, where movement of the vectors could facilitate parasite sharing across interface areas, even if fences are used to reduce contact between domestic and wild hosts.

One particularly complex system where this could be important to understand is trypanosome-mediated diseases transmitted by tsetse flies in Africa. Although there are multiple species of tsetse flies that can transmit multiple species of trypanosomes, *Glossina pallidipes* is the most economically important species in East Africa [21], because it is the main vector of Animal African Tryanosomiasis (AAT) and it is also a vector of Human African Trypanosomiasis(HAT). Wild animals have been reported as reservoir hosts both for AAT[3,57] and HAT[25,26,34,73,74] but the extent of transmission across the wildlife-livestock interface remains unclear.

Tsetse flies (genus *Glossina*) are generalist blood-feeders on a wide variety of vertebrate host species, including mammals, reptiles and birds[72]. Importantly, both male and female tsetse feed throughout their lifetimes. There is thus high potential for vector-mediated connection between parasite sylvatic and domestic cycles in wildlife-livestock interface areas if tsetse flies take meals from different host species

at each feeding opportunity. However, the likelihood that an individual tsetse feeds

on different types of hosts where they occur sympatrically, compared to feeding

predominantly on a single species, has not been clearly established and so the

relative risks of increased trypanosome infections in livestock living near wildlife

remains a critical gap in knowledge[6]. Although three trypanosome species are

traditionally associated with the disease in livestock (*T. brucei*, *T. congolense*, and *T.*

vivax), a higher diversity has been identified in wildlife[4], which could potentially

increase risks of disease if transmission from wildlife to domesticated animals is

common.

Few studies have attempted to combine investigation of host-feeding patterns of

individual flies, trypanosome infection, and intrinsic factors of tsetse flies distributed

in different regions. Identification of hosts through blood meal analyses is a highly

useful tool that has been used to predict host preferences and feeding behaviours

across a wide range of vectors[24,28,39] A commonly used approach has been to use

polymerase chain reaction (PCR)-based techniques to amplify and sequence host

DNA from blood meal contents in the guts of fed flies. This has largely been based

on mitochondrial genes due to their high copy number and the extensive databases

available due to their use as universal markers for DNA barcoding[33,42,60]. For

example, in the Serengeti ecosystem in Tanzania, which holds a high number and

wide range of wild animals, an investigation of blood meal composition in tsetse flies

based on sequencing of the cytB gene compared to surveys of host density revealed

strong preferences for particular wild hosts, which varied by species of fly (*G.*

swynnertoni vs *G. pallidipes*[5]. This clearly demonstrated the value of relating feeding

patterns to the diversity of hosts present. However, trypanosome prevalence was not

quantified in these studies and domestic hosts were not present in the study area;

so, relative host preferences for wildlife compared to livestock was not determined.

Feeding activity, where individual flies feed consecutively on different types of hosts,

could alter relative risk of transmission of trypanosomes. More frequent feeding

might occur, for example, if flies are disrupted while feeding or if they abandon a

host that they perceive to be unsuitable or that shows defensive behaviour[63,68]. The

dominance of nonpreferred hosts in a particular geographic area could thus result in

more frequent host switching and so increased rates of multiple feeding and

potentially higher exposure to a diverse range of parasites. In East Africa, *G.*

pallidipes is widespread and has been demonstrated to feed on a wide range of

hosts, including bovines[19,20,54,58,69], suids[11,58], elephants[54], antelopes[1] and cattle[54].

Warthogs, bushbuck and African buffalo have been suggested as the preferred

hosts[13,19,43,45,54,58] but this varies by geographic region[5,22,54,65,66] and relative

preference for domestic and wild hosts has not been specifically assessed.

In a previous study, we established that the prevalence of trypanosomes among

tsetse flies in two regions of Kenya (Nguruman and the Shimba Hills) showed

complex relationships with geographic location, tsetse specific factors (age, sex and

fly species), species of trypanosome and the presence of an endosymbiont[17]. The

main aim of the current study was to assess whether some of the variation in the

detection of trypanosomes across sites could be explained by differences in host

feeding patterns. Specifically, we aimed to determine: 1) the diversity of vertebrates

tsetse fed on at sites where different types of host were present; 2) whether host

feeding patterns varied in relation to type of hosts or intrinsic tsetse factors (i.e. age

and sex); and 3) if there was a relationship between trypanosome detection and host

feeding patterns, host types or tsetse-specific factors.

Methods

Sampling and tsetse fly characterisation

The *G. pallidipes* samples are described in Channumsin *et al.*[17], where details of the sampling strategy are provided (see Extended data 1[18] for locations of the traps). NG2G traps baited with acetone and cow urine[15] were used for collecting tsetse flies from three sites that differ in anticipated levels of relative abundance of livestock and wildlife, with the sampling effort (number of traps) determined by the relative abundance of flies in the area. Two sites were sampled in the Shimba Hills National Reserve (Kwale County, in the coastal region of Kenya), which is a relatively small (250 km^2) protected area separated from surrounding agricultural areas by a wildlife fence. There is extensive habitat for tsetse flies, including on the park boundaries. Buffalo Ridge is within the fenced wildlife protected area in the middle of a thicket forest, where many tourists visit all year, while Zungu Luka has a woodland type of vegetation, and is located on the border of the park close to a permanently human-inhabited rural area with resident livestock. In contrast, the Nguruman region contains lowland woodland patches surrounded by open savannah; habitats, which have been found to host a large number of *G. pallidipes* and *G. longipennis*[16]. The sampling site (Mukinyo) is at the border of the Olkiramatian group ranch, which is a wildlife conservancy without fences, where the distribution of domestic and wild tsetse hosts overlap when livestock are grazed in the area but there is no permanent human settlement close by.

Characteristics of the flies and presence of trypanosomes were previously determined by Channumsin *et al.*[17]. Sex and species of flies were determined based on morphological characters. Age was estimated based on a wing fray score where increased damage indicates increasing age[37]., Whole flies were preserved in 95%

ethanol and stored at -20°C. Presence of trypanosomes in mouth parts and

proboscis of the flies collected was determined using general primers targeting the

ITS-1 region of the rDNA array (CF: 5' CCGGAAGTTACCGATATTG 3' and BR: 5'

TTGCTGCGTTCTTCAACGAA 3'[56], that allow identification of trypanosome species

based on size of amplicons, as described in Channumsin et al.[17]. Although multiple

species of tsetse were used in the previous study, here we focused on individuals

identified morphologically as G. pallidipes (N = 577) and screened trypanosomes in

DNA that had been extracted from abdomens. All flies sampled were used, rather

than selecting individuals that had appeared to have fed recently.

Identification of diversity of hosts and feeding patterns
153

from G. pallidipes blood meals
154

We used primers developed by Kocher et al.[40] targeting a 359 bp fragment of the

mitochondrial gene cytochrome B (cytb) gene in mammals (Cb1: 5'

CCATCCAACATCTCAGCATG   ATGAAA   3'   and   Cb2:   5'

GCCCCTCAGAATGATATTTGTCCTCA 3') which enabled direct comparison with

two previous studies[4,54] and because they showed more reliable amplification in a

pilot study[75] than primers targeting the mitochondrial cytochrome C oxygenase 1

(CO1) gene (VF1d-t1 and VR1d-t1)[36]. During processing for DNA extractions, in

order to reduce risk of contamination, the dissected tissues were cleaned 2-3 times

with 95% ethanol, then left to dry, before moving to new individual microtubes with

liquid nitrogen for sample crushing and DNA extraction using DNeasy® blood and

tissue kits (Qiagen Inc., Paisley, UK). PCR cycling was carried out in 25 µl reaction

mixtures containing: 1X PCR buffer; 0.2 mM dNTP mixture; 1.5 mM $MgCl_2$ (Thermo

Scientific); 0.5 µM of each primer; 1 unit of Taq DNA polymerase (Invitrogen Inc,

Carlsbad, CA., USA); and 2 µl tsetse abdomen DNA template. Samples were pre-

heated at 94°C for 5 min, denatured at 94°C for 30 sec, annealed at 55°C for 45 sec,

then extended at 72°C for 30 sec, with 35 cycles of the amplification and a final

extension at 72°C for 10 min[54]. PCR products were visualised using 1.5%

UltraPure^{TM} Agarose gels (Invitrogen, Paisley) with 2% Ethidium Bromide

(Invitrogen, Paisley) in 1X TBE buffer (108 g of Tris Base, 55 g of Boric acid and 40

ml of 0.5 M EDTA). Results were visualised and analysed on a gel documentation

system (UVIpro Platinum, UVITEC, Cambridge, UK or GeneDoc, BioRad Inc, UK).

PCR products of the expected size (359 bp) yielding ≥ 20 ng were cleaned using

ExoSAP-IT PCR Clean-up Kits (GE Healthcare). In cases where the yield of PCR

products was lower than this threshold, multiple PCR products were concentrated

and QIAquick Gel Extraction Kits (Qiagen Inc, Paisley, UK) were applied to extract

the PCR products from agarose gels. All purified samples were sent for Sanger

sequencing in both forward and reverse directions, using the Sequencing Service at

the University of Dundee (MRC I PPU, School of Life Sciences, University of

Dundee, Scotland, www.dnaseq.co.uk) using Applied Biosystems Big-Dye Ver 3.1

chemistry on an Applied Biosystems model 3730 automated capillary DNA

sequencer.

Base-calling was manually corrected, sequences were aligned and consensus

sequences for forward and reverse primers for each individual generated using

Sequencher version 5.3 (Gene Codes Corporation, Ann Arbor, MI USA). The Basic

Local Alignment Tool (BLASTn)[2], was used to identify the closest matching

sequences in the GenBank database to determine the host identity of each

consensus sequence. Chromatographs with only single peaks based on direct

sequences were classified as "single host feeding". Sequences that were still clearly

readable but showed more than one peak at multiple positions were classified as

"multiple host feeding". While the difficulty of resolving the phase of genetic variants

precluded identification of all hosts from direct sequencing of multiple-peak products,

the dominant host was determined based on BLASTn analysis of the most prominent

peaks.

Host-feeding patterns were confirmed in a subset of samples by cloning using

TOPO®-TA Cloning Kits (Invitrogen, UK), with at least six plasmids of each sample

sent for sequencing, after purifying using QIAprep Spin Miniprep Kits (Qiagen Inc,

Paisley, UK). Ten samples whose chromatographs showed double or triple peaks at

single positions in the direct sequences were cloned to confirm that multiple peaks

were due to feeding on multiple host species rather than poor quality sequences

(five flies from Buffalo Ridge; three from Zungu Luka; two from Mukinyo). An

additional seven samples that appeared to have fed on single hosts but with some

ambiguous peaks were also cloned and sequenced (five from Buffalo Ridge and two

from Zungu Luka).

To enable of assessment of variation in the type of hosts fed on across sites,

dominant hosts resolved were classified as "domestic" (including livestock or

companion animals), "human", or "wild".

In order to assess infraspecific diversity in hosts fed on across the sites, sequences

were first exported to Se-Al version 2.0[61] to manually align and prune sequences to

the same length. DNAsp version 5.0[47] was then used to resolve variants into unique

haplotypes within host species. Minimum spanning networks were plotted using

PopArt[46], to indicate relative frequencies of host haplotypes across the sampling

sites.

Variables influencing tsetse feeding behaviour

Generalised linear models (using the glm function, as implemented in the lme4 package[8] using R version 4.0.2[67] were used to test whether variation in feeding behaviour of the flies (single vs multiple hosts, modelled as a binary response variable) was influenced by the type of the dominant host (domestic, human, or wildlife), tsetse sex and age (sex as a continuous variable based on awing fray score averaged across the two wings of an individual), or site (Buffalo Ridge, Zungu Luka, Mukinyo). Interactions between the type of host with sex, age and site were also considered in the full model. Model selection was performed using likelihood ratio tests to find the minimum model that best explained the data. Odds ratios were calculated from the coefficients of the final model using the "oddsratio" package in R[64]. To check the appropriateness of the binomial model, overdispersion was assessed by checking that the ratio of the residual deviance to the degrees of freedom in the final model was below 1. The fit of the final model was assessed by McFadden's pseudo-R^2, defined as $1 - LL(\text{final model})/LL(\text{null model})$, where LL = log likelihood[52].

Prevalence of *Trypanosoma spp.* in relation to *G. pallidipes* feeding patterns

A similar statistical approach was used to test whether the presence of trypanosomes was explained by host type or feeding behaviour while considering possible influences of fly sex and age, or site based on conclusions from our previous study[17]. Since we were specifically interested in whether tsetse feeding behaviour affected trypanosome detection, pairwise interactions were considered between the type of host and the feeding pattern with age, sex and sampling site of

the flies. Model selection and fit were performed as described for the feeding pattern

models. Given the wide range of hosts that the flies feed on, the influence of

particular host species on trypanosome prevalence was considered only

qualitatively.

To specifically visualise whether feeding patterns or dominant host types were

related to trypanosome prevalence when accounting for geographic location and

tsetse sex and age, Multiple Correspondence Analysis (MCA), as implemented in

the FactoMineR package (version 1.30[44]) was used. For this analysis, age was

considered as a categorical variable by classifying individuals into the following age

categories: "young" (wing fray score 1–2.5); "juvenile" (3.0-4.0) and "old" (4.5–6.0)

based on the average score between the two wings for each individual. Other

variables were: site, presence or absence of *Trypanosoma spp*, sex, feeding pattern

(single vs multiple) and dominant host type (domestic, human or wild). Variation

along pairs of principal component axes was visualised using "ggplot2()"[27] in R[67].

Results

Diversity of hosts identified from *G. pallidipes* blood meals

From 573 *G. pallidipes*, 128 flies showed no evidence of a recent blood meal based

on lack of amplification products following screening with the Cb1 and Cb2 primers.

These samples were excluded from analyses (Table 1). The remaining 445 flies

showed amplified products of the expected size, which were sequenced and used to

classify feeding status (Table 1; Extended data 2[18]). For 197 of the 247 samples for

which dominant hosts could be resolved to the species level, a single amplification

product was apparent in the chromatographs; these were classified as having

recently fed on a single host. Cloning of seven of these samples confirmed

amplification of DNA from only a single host species (Extended data 2 and 3[18]). The

chromatographs for 53 samples clearly showed multiple peaks that could be

confidently attributed to feeding on multiple hosts rather than poor sequence quality

and the dominant host could be resolved through BLASTn analysis of the strongest

peaks. This represented 37% (Buffalo Ridge), 31% (Zungu Luka) and 51%

(Mukinyo) of the samples screened at the three sites (Table 1). Cloning of 10 of

these PCR products confirmed amplification of DNA from more than one host, with

up to four different host species identified in single flies (Extended data 3[18]).

We took a conservative approach to classifying feeding patterns: chromatograms of

the remaining 198 samples from which amplification products were obtained were

not considered of sufficient quality to reliably determine the source of the blood

meals; these were classified as "unidentified". While many of these would likely

represent multiple feeding, we wanted to avoid confounding with poor sequence

quality so they were classified as fed but not identified (Table 1; Extended data 2[18]);

only samples with confident dominant host calls were included in the statistical

analyses.

Host composition of blood meals varied across sites (Table 2), with buffalo

dominating in the two wildlife protected areas (Buffalo Ridge and Mukinyo) and

humans predominating in the site bordering the SHNR (Zungu Luka), where no

buffalo feeds were detected (Figure 2). Mukinyo had a wider range of wild hosts

identified in blood meals than Buffalo Ridge but elephants, antelope and warthog

were found at both sites. Flies from Buffalo Ridge also shared most of the same

domestic host species as Zungu Luka, suggesting that flies moved across the

fenced interface to feed. Across all sites, only a single fly (from Mukinyo) was

confirmed to have fed on domestic cattle.

Table 1 Summary of blood meal analysis results based on direct sequencing. Cytb negative samples were classified as "unfed flies"
but were not considered in the analyses since they could represent lack of amplification rather than lack of feeding. Single host feeding
refers to cases where the cytb sequence had only single chromatograph peaks. Multiple host feeding were samples for which cytb was
amplified but the sequences showed multiple peaks and the dominant sequence could be identified to species, classified as domestic
animals, humans or wildlife. Flies showing strongly amplified cytb PCR products but for which the number or type of host species could
not be confirmed due to poor sequencing quality are labelled as "not identified". The number of flies that tested positive for the presence
of trypanosomes is indicated in parentheses. The human samples that were potential contaminants (haplotype 1; Figure 3) were
excluded.

Site	Single Feeding			Multiple Feeding			Not identified	"Unfed"	Total Screened
	Domestic	Human	Wild	Domestic	Human	Wild			
Buffalo Ridge	0	1 (1)	31 (8)	9 (5)	9 (3)	5 (3)	62 (23)	33 (10)	150 (53)
Zungu Luka	1 (1)	6 (2)	8 (5)	12 (7)	11 (8)	1 (1)	60 (31)	29 (22)	128 (77)
Mukinyo	1 (1)	0	146 (60)	0	0	3 (2)	76 (22)	66 (20)	292 (105)

301
302

Table 2 Dominant host species resolved from blood meal analysis *G. pallidipes* sampled from the Shimba Hills (Buffalo Ridge and Zungu Luka) and the Nguruman region of Kenya, based on direct sequencing of cytb. Homozygous amplicons were classified as "single" feeding whereas sequences with multiple peaks were classified as having fed on "multiple" hosts. The dominant host was identified based on BLASTn. The relative abundance of the various host species is expressed as the total % of sequences for which the dominant host could be identified within that site. Species are ordered by relative abundance of wild and domestic hosts.

Site	Buffalo Ridge			Zungu Luka			Mukinyo		
Host	Single	Multiple	Total %	Single	Multiple	Total %	Single	Multiple	Total %
Buffalo	25	3	50.9	0	0	0.0	105	2	71.3
Elephant	2	1	5.5	1	1	5.1	29	1	20.0
Antelope	2	1	5.5	6	0	15.4	1	0	0.7
Warthog	2	0	3.6	1	0	2.6	5	0	3.3
Giraffe	0	0	0.0	0	0	0.0	4	0	2.7
Hyaena	0	0	0.0	0	0	0.0	2	0	1.3
Human[a]	1	9	18.2	6	11	43.6	0	0	0.0
Goat	0	7	12.7	1	7	20.5	0	0	0.0
Mouse	0	2	3.6	0	4	10.3	0	0	0.0
Chicken	0	0	0.0	0	1	2.6	0	0	0.0
Cattle	0	0	0.0	0	0	0.0	1	0	0.7
Total	32	23		14	25		147	3	

[a] Excluding potential contaminants

In addition to identifying just the species of host from the blood meals, we found

intraspecific variation in mtDNA haplotypes within host species (Figure 1; Extended

data 4[18]). Single haplotypes were found for all of the domestic hosts identified:

mouse (*Mus musculus*), chickens (*Gallus gallus*), goat (*Capra hircus*) and cattle (*Bos*

taurus). Three human haplotypes were identified, with the majority showing similarity

to cytb sequences identified from tsetse blood meals in the Serengeti, Tanzania

(type 2; n = 21) or Zambia (type 3; n = 3) (Extended data 4[18]). However, three

samples with evidence of only a single host matched an Asian haplotype from

Taiwan (type 1: one from each of the three sites), which is the ethnic origin of the

primary researcher; these three were excluded from analyses because they were

suspected laboratory contaminants. Three additional samples were identified as

human but the sequences were not clean enough to resolve the haplotype because

they were all identified in flies that appear to have fed multiple times. There was

extensive variation in haplotype diversity among the wild hosts but this was not

always related to their relative abundance in the samples (Figure 1; Table 2).

Figure 1. Minimum spanning networks indicating intraspecific diversity and relative frequency of haplotypes between populations from this study for: a) buffalo; b) elephants; c) antelope; d) warthogs; and e) humans. Note that human type 1 matched the ethnic origin of the main investigator (Asian); samples with this haplotype were considered as contaminants and excluded from analyses. The two antelope sequences labelled "unknown" were found only in single clones, with a more dominant host predominating, and had no close match using BLASTn. Circle sizes are proportional to the frequency of each haplotype (see Extended data 4[18] for values); notches on branches indicate the number of nucleotide substitutions separating haplotypes; colours represent the population of origin (white = Buffalo Ridge; grey = Mukinyo; black = Zungu Luka). Three haplotypes were found in giraffes but they differed by only single nucleotide and were each found in only one or two individuals so they are not shown here.

Variables influencing tsetse feeding behaviour

A qualitative summary of variation in feeding behaviours (single vs multiple) of flies

in relation to their sex, age, sample site and type of host fed on is provided in Figure

2. Although there was variation in the sex and age distribution of flies across sites

(Extended data 5[18]), the most striking pattern distinguishing single and multiple

feeding was in relation to differences in the type of hosts fed on.

The two sites from the Shimba Hills (Buffalo Ridge and Zungu Luka) showed a

higher proportion of flies that appear to have fed on multiple hosts than the site from

Nguruman (Mukinyo): 41.8% from Buffalo Ridge; 61.5% from Zungu Luka, compared

with 2.0% from Mukinyo. However, this appeared to be influenced by the type of host

(Table 2; Figure 2). Buffalo Ridge showed a predominance of flies that had fed on

wild hosts (65.5%) and most individuals with a dominant domestic or human host

had fed on multiple species (18/19, compared with 5/36 for wild hosts). Cloning

revealed that all five of the individuals classified as multiple feeding had fed on

humans and at least one other domestic animal; four of the individuals had also fed

on a wild host (Extended data 3[18]). In contrast, for Zungu Luka, humans comprised

43% of dominant hosts identified compared with 33% domestic and only 23% wild

animals. Similar to Buffalo Ridge, the majority of flies feeding on non-wild hosts fed

on more than one host species (23/30), compared with only 1/9 of the flies for which

dominant sequences were identified as wild hosts. All three multiple feeding flies

cloned from this site had fed on humans, with one also having fed on both domestic

(goat, mouse) and wild (antelope) hosts, one on a single wild host (bushbuck) and

another on a domestic host (chicken) (Extended data 3[18]). At Mukinyo, 99% of flies

had fed on wild hosts, with only a single fly identified as having recently fed on a

domestic host (identified as single feeding on cattle) and no human hosts were

detected. Moreover, only 3/149 flies with dominant wild hosts had fed on more than

one host species (confirmed by cloning for two of the individuals; Extended data 3[18]).

Figure 2 Feeding patterns of flies across sites in relation to their age, sex, and

dominant host type (domestic, human, wild). Age of flies was estimated based on

average wing fray scores across the two wings or an individual fly, with increasing

damage indicating relatively older flies. Feeding patterns were based on whether

sequence chromatograms indicated amplification of a blood meal from a single host

(black) or more than one host (red); the sex of the flies was determined visually

(female = circles; male = triangles). Only values greater than 0 have been plotted.

Flies from all age classes at Mukinyo fed predominantly on single wild host species,

with no evidence of feeding on humans and only a single mid-age female feeding on

a domestic cow. In contrast, feeding on a mixture of domestic and wild hosts was

found for all age classes at the Shimba Hills sites, Buffalo Ridge and Zungu Luka

and multiple feeding was more frequent than single feeding, except for wild hosts.

Using the type of feeding behaviour (single vs multiple hosts) as a binary response

variable, the final model selected by maximum likelihood included a highly significant

effect of type of host (LRT: χ^2 = 52.0, df = 2, p = 5.09e-12) and a significant effect of

site (LRT: χ^2 = 13.2, df = 5, p = 0.001). Examining the odds ratios (OR) indicated a

substantially lower incidence of multiple feeding on wild compared to domestic hosts

(OR = 0.009; CI = 0.001-0.045) but similar incidence in humans and domestic hosts

(OR = 0.232: CI = 0.031-1.151). As might be expected based on the difference in

distribution of hosts, Mukinyo showed lower levels of multiple feeding (OR = 0.081;

CI = 0.014-0.323) than Buffalo Ridge, whereas there was little difference between

the two Shimba Hill sites (OR = 0.407; CI = 0.081-1.528). Comparing the residual

deviance (104.12 on 239 df) and null deviance (247.49 on 243 df) indicated that

there was no evidence for over-dispersion and McFadden's pseudo-R^2 was 0.58,

indicating a relatively good fit to the data that the final model explained.

393 Prevalence of *Trypanosoma spp.* in relation to *G.*

394 *pallidipes* feeding patterns

Across sites, 44% (n = 107) of the flies for which hosts could be identified to species

tested positive for trypanosomes with 54% (29/53) associated with dominant

domestic hosts and 41% (79/194) with wild (Table 1). Of flies feeding on multiple

hosts, 58% tested positive for trypanosomes, compared to 40% that had fed on

single hosts, but this was influenced by the higher rate of infection in Zungu Luka

(61.5%), where single feeding was rare, compared to in Buffalo Ridge (36%) and

Mukinyo (42%). It was more difficult to interpret patterns by host species because of

the large differences in their relative abundance (Extended data 6[18]).

As found in our previous study[17], generalised linear models using trypanosome

presence as a response variable were difficult to interpret. All of the interactions

considered except for that between feeding pattern and site significantly explained

variation in trypanosome detection ($p<0.01$). However, testing the fit of the final

model based on pseudo-R^2 (0.08) indicated that only a small amount of the variation

in trypanosome presence was explained. The residual deviance (303.22 on 225 df)

also suggested over dispersion.

Figure 3 Detection of trypanosomes across sites in relation to fly feeding

behaviour, age and site. Trypanosome detection (black = negative, red = positive)

is indicated in relation to feeding pattern (circle = single, triangle = multiple), with

separate plots by type of host and site. Generalised linear mixed models indicated

multiple significant pairwise interactions between feeding behaviours and other

tsetse-specific variables. Sex was involved in a significant interaction with type of

host but not feeding pattern, but it has been excluded here to more clearly

demonstrate the complicated interactions between the other variables.

For this reason, multivariate ordination analyses were used to visualise associations

between variables. Based on MCA analyses, strong correlations among site of *G.*

pallidipes collection, host feeding pattern and type of host were apparent in

dimension 1 (Figure 4; Extended data 7[18]). In contrast, trypanosome status was

resolved primarily along dimensions 2 and 3, as were sex and age of the flies; a

positive association was found between trypanosome positive samples and juvenile

male flies, while trypanosome negative flies tended to be found in young female flies.

Figure 4 Multiple Correspondence Analysis (MCA), showing associations of

dimension 1 (Dim 1; 29.79 % of the variance) and 2 (Dim 2; 12.32% of the variance)

in relation to age category (young, juvenile, old), feeding pattern (single or multiple),

host type (domestic or wildlife), sex (male or female), site (Buffalo Ridge, Zungu

Luka or Mukinyo), and *Trypanosoma spp*. status (positive or negative). This Figure

clearly shows the strong association between feeding pattern and host type, driven

by the differences in fly behaviour at Mukinyo compared to Zungu Luka resolved

along dimension 1. Old flies were also highly correlated with multiple feeding of

domestic and human hosts at Zungu Luka Trypanosome status was not explained

by variation along dimension 1 but was more related to sex and age of younger flies
resolved along dimension 2. Buffalo Ridge was differentiated from the other two
populations along both dimensions 1 and 2.

440 **Discussion**

Based on detailed sequence analysis of mitochondrial gene amplicons, our results

suggest that individual tsetse flies (*G. pallidipes*) vary markedly in their feeding

patterns. In particular, we found that flies feeding on wild hosts tended to show

higher feeding success (based on evidence for amplification of only a single host

species in blood meals) than those feeding on domestic animals and humans.

Although site also influenced patterns of feeding, this was somewhat confounded by

the relative abundance of wild hosts that were fed on between the two regions

compared. Although previous studies have found a similar diversity of hosts as we

found based on analyses using the same cytochrome b primers[40] or other mtDNA

regions[54], we are not aware of other studies that differentiated single from multiple

feeding based on analysis of sequence chromatograms. Moreover, blood meal

analyses do not typically assess within-host diversity; our haplotype analysis

suggests that there is potential to use feeding arthropods as "flying syringes"[12] not

only for identification of hosts but also could be used to make inferences about host

population structure. Our results were not able to clearly test whether host feeding

patterns or type of host influenced prevalence of trypanosomes in individual flies. As

in our previous study[17], trypanosome presence was explained by interactions

between multiple variables. We had hypothesised that some of this complexity might

be reduced by including feeding behaviours, but they also were found to influence

variation dependent on other variables. Multivariate analysis using MCA suggested

that prevalence of trypanosomes was correlated with sex and age of the flies

whereas feeding pattern was correlated with type of host and geographic location.

Together, these results suggest that differences in host communities in different

regions could influence the risk of transmission between vectors and hosts in

complex ways and highlight the potential for increased transmission risk in interface

areas where both livestock and domestic hosts coexist.

Host diversity in *G. pallidipes* blood meals

We identified the dominant hosts for 46% of the *G. pallidipes* samples screened

(56% of the samples that showed positive amplification products), which is

comparable or higher than previous studies using the same primers[5,12]. We found

extensive variation among the species fed on in two different geographic areas. In

the Shimba Hills, where a fenced wildlife protected area is located within a few km of

human settlements, flies fed on both domestic and wild hosts, with blood meals from

both host types detectable within individual flies. In contrast, in the Nguruman region,

only a single fly was identified that had fed on a domestic host. This is consistent

with Muturi *et al.*[54], who also did not identify domestic hosts in their survey of the

Nguruman region, despite finding predominantly cattle blood meals at a site

surveyed in Uganda. The results from Nguruman may be due to sampling time and

the large-scale shifts in cattle grazing sites according to season (Masiga,

unpublished). Snow *et al.*[65] suggested that, even though flies in areas dominated by

cattle fed readily on these domestic hosts, a positive correlation between the number

of wild herbivores and the abundance of *G. pallidipes* suggested that feeding

success was poor on local livestock (based on a low density of flies where cattle

were numerous). The use of insecticides on cattle also could help to explain

decreased density of tsetse traversing from wildlife-protected areas through to

livestock dominated areas in interface areas[48,49]

Surprisingly, no domestic cattle were detected in our study from the Shimba Hills,

despite the proximity to settlements with mixed herds of cattle, sheep and goats [55].

However, domestic hosts were also not identified in the Shimba Hills region in a

previous study based on host detection using haemagglutinin assays[65]. This could

indicate that flies avoid cattle when more favourable hosts are present. However,

there also could be seasonal differences, as trypanosome prevalence in cattle was

found to be high (33.9%) in Kwale County, in a previous study that also found *G.*

pallidipes at high abundance[51]. At Mukinyo a single individual fed on domestic cattle

but there was a very low proportion of flies that fed on multiple hosts (2%) and a

predominance of buffalo (71%) among the samples where the dominant host could

be identified. It is possible that buffalo are abundant hosts that are easy to feed on

and so flies could learn to return to the same host species[14]. Our results in general

are consistent with higher feeding success on wild compared to domestic hosts.

Our finding of African buffalo as the main hosts of *G. pallidipes* in Nguruman and

Buffalo Ridge supports previous reports that ruminants are attractive to adult *G.*

pallidipes, *G. fuscipes* and *G. brevipalpis*[1,30]. However, host selection has been

found to vary extensively by population (Extended data 7[18]). Differences across

studies could be due to differences in methodology but also could be due to

microhabitat differences[66], such as seasonal variation in host availability, the

vegetation type or cover at particular sites, or particular environmental conditions in

different years, which affects overlap of habitat and activities between tsetse flies

and hosts[1,19]. It is interesting that no buffalo blood meals were detected at Zungu

Luka, despite its close proximity (~ 20 km) to Buffalo Ridge, where buffalo are

abundant. This could suggest that flies feeding in human settlements move into the

park to feed on wildlife but once feeding on their preferred wildlife, they do not move

out into the human-settled regions or that flies tend to dwell proximally to where

bloodmeals are readily available. It would be interesting to quantify relative

abundance of hosts of different types and directionality of movements to test this

hypothesis. Specific choice tests between domestic and wild hosts also could reveal

important information about preferences that could inform control interventions[70] as

has been done for malaria-carrying mosquitos[50]. Nevertheless, the finding of flies

collected in the same traps feeding on both wild and domestic hosts emphasizes the

high potential for cross- feeding between these host types when they occur

sympatrically.

Humans have been suggested as inappropriate hosts because they camouflage

their odours, apply chemical repellents, and react strongly to tsetse bites, which

could result in unsuccessful feeding[9,10,29] that could lead to host switching.

Hargrove[29] found that the presence of humans not only repelled tsetse flies but also

inhibited the landing response to approach other potential hosts nearby. Most of the

mtDNA haplotypes we identified from human samples were consistent with those

expected regionally and one haplotype was shared with previous published

sequences from the Serengeti (Auty et al. 2016a; Extended data 4[18]); we also did

not find evidence of feeding on humans in Mukinyo. Although measures were taken

to rule out contamination, these results are surprising and patterns of tsetse feeding

in areas of higher human density should be investigated further.

532   Variables influencing tsetse feeding behaviour

We found that the propensity for feeding on single compared to more than one host

species was highly influenced by the type of host fed on, with more single feeding on

wild hosts than on humans or domestic host. Cloning and sequencing revealed that

some flies feeding on domestic or human hosts had fed on up to four different host

species and confirmed that single feeding was more common in flies feeding on

wildlife. Theoretically, the number of clones could be used to predict which host was

last fed on, but this would also depend on the rate of feeding of the fly (e.g. if they

were interrupted and switched hosts very rapidly, more than one blood meal might

have a similar DNA concentration) and lack of bias in PCR amplifications. There also

could be behavioural differences that could result in detection biases: 1) flies might

feed more thoroughly on their preferred hosts (such as buffalo), increasing the blood

meal volume from that host; 2) flies might feed multiple times on the same host

species occurring at high local densities (suggested here by the presence of multiple

haplotypes of the same host species in some cases); or 3) hosts might differ in

effective defence mechanisms, resulting in low blood meal volumes due to

interrupted feeding[68]. If feeding on an initial host is interrupted or too low quality

("unsuccessful"), flies might switch hosts. Unsuccessful feeding of tsetse flies on

cattle have been attributed to host defence, such as twitching the skin, flicking the

tail, flicking the ears, and kicking or stamping[63]. Wild animals might react less to

tsetse flies feeding and/or be surrounded by less other biting insects than

domesticated animals. Nevertheless, our results suggest higher host fidelity (or

feeding success) when feeding on wild, compared to domestic, hosts.

Prevalence of *Trypanosoma spp.*

There was not a clear association between prevalence of trypanosomes and type of

host or host-feeding patterns in the tsetse flies. In our previous study Channumsin *et*

al.[17], we found that trypanosome prevalence was explained by complicated

interactions between age, sex and sampling site of the tsetse flies. Here we found

that detection of trypanosomes was also significantly influenced by interactions with

these tsetse-specific variables with both host type and feeding patterns. This made it

difficult to test our hypothesis that the tsetse feeding behaviour might explain some

of the variation in trypanosome detection. Specifically, we hypothesised that feeding

on multiple hosts could increase risk of trypanosome infection in flies. However, this

was not apparent in the multivariate analysis using MCA (Figure 4) suggested a

stronger correlation among feeding pattern, host type and site than with

trypanosome status, sex and age of tsetse flies. The blood meals we analysed also

only reflect the most recent feeds and so likely do not reflect their overall feeding

history. Bouyer *et al.*[14], suggested that repeated feeding on the same host species

was likely to increase risk of trypanosome transmission within species, but to

decrease risk between species. There is some evidence that trypanosome infection

might influence feeding success and feeding behaviour of the flies, but it is not

conclusive[45]. For example, high numbers of *T. congolense*, which attach to the

cuticle of the proboscis, could interrupt feeding and result in more frequent probing[38].

Alternatively, nutritional status of the flies could affect their relative susceptibility to

trypanosome establishment[41]. Thus, an association between the frequency of

feeding and trypanosome infection status should be further studied in laboratory

experiments to test whether trypanosome infection causes a feeding pattern change

or differences in feeding patterns promote trypanosome infection. Nevertheless, our

results did not suggest an increased prevalence of trypanosomes in communities

where both domestic and wild hosts were fed on that would suggest increased risks

in livestock interface regions.

Amplicon-based blood meal analyses

Although blood meal analyses provide a powerful tool for investigating feeding

behaviours of haemotophagous insects, the potential for biases in any PCR-based

approach deserves consideration. For example, we found that a higher proportion of

hosts could be resolved from Mukinyo than the other areas, which could be due to

the dominance of wild hosts but could also be due to higher fidelity of the primers

used on the species of hosts detected. Previous studies comparing the relative

reliability of cytb and COI mtDNA e.g. Muturi *et al.*[54] have found that neither alone

amplifies products from all potential host types present. In the Shimba Hills, although

goats were identified from flies sampled from both sites, there were also additional

samples that matched goats in BLAST analyses that were not included in the

analysis because it was difficult to determine whether the sequences represented

multiple feeding or just poor sequence quality. Moreover, blood meal analyses rely

on completeness of reference databases. We found several cytochrome b

haplotypes that were closest to antelope in BLAST but the similarity was too low to

resolve to species (93%); this lack of reference sequences could have led to

underestimates of host usage in previous blood meal analyses. Analyses of blood

meals also do not typically consider the possibility of amplification of nuclear copies

of mitochondrial genes (numts), the presence of which can vary dramatically across

vertebrate species[32]. It was for these reasons that we took a conservative approach

to interpreting feeding patterns based on blood meals by only considering

sequences where the dominant host could be clearly identified by direct sequencing

(or cloning), While this meant that we likely underestimated the rate of feeding on

multiple host species, a clear pattern remained that fewer ambiguous sequences

were found at Mukinyo, where wild hosts dominated, than at the other sites (26% vs

44%, respectively).

There has been a recent shift towards using deep sequencing approaches for

amplicon-based host identification[35,62], which would allow more rigorous testing of

potential biases and could also allow simultaneous targeting of hosts and

trypanosomes by using multiplexed approaches[23]. Non-PCR based assays such as

high-resolution melting point analysis have already shown high promise as

alternatives for blood meal analysis[22,59]. However, deep sequencing following

enrichment approaches rather than PCR amplification, such as hybrid sequence

capture[7,53], have the potential to not only provide a more comprehensive analysis of

host diversity, but could allow clearer interpretation of relative read numbers in

relation to feeding patterns.

Conclusions

Identification of the hosts that *G. pallidipes* fed on based on direct PCR sequencing

revealed evidence for both use of a wide range of hosts and multiple feeding bouts

by individual flies. However, in wildlife dominated areas, there was a much stronger

tendency for flies to feed on single host species compared to sites where domestic

hosts were more commonly fed on, with individual flies feeding on up to four different

detectable host species. If this indicates that domestic animals are not preferred

hosts, this could have important implications for understanding risk of transmission

of trypanosomes between wildlife and livestock in interface areas. Our results also

demonstrate the value of detailed sequence analysis of blood meals of

haematophagous insects to include not only identification of the host species but

patterns of feeding by individual flies in relation to their sex, age and habitat. The

increased accessibility of deep sequencing approaches opens up new possibilities

for more detailed assessments, which might also include the ability to predict the

timing or success of feeding on different hosts based on relative read depths.

Data Availability

637 Sequences have been deposited to Genbank, with accession numbers MN148732-

638 MN148768 (Extended data 4[18]).

639 Acknowledgments

This work was funded by the Wellcome Trust (grant number: 093692) to the

University of Glasgow. For the purpose of Open Access, the author has applied a

CC BY public copyright licence to any Author Accepted Manuscript version arising

from this submission. The authors are grateful to the Kenya Wildlife Service for

permission to perform this study in protected areas. Esther Waweru and James Kabii

provided technical and administrative assistance at *icipe*. James Kabii, Moses Tanju,

Loya Suleyman, John Andoke (deceased) and his team (*icipe*) provided invaluable

expertise and help for the trapping of tsetse. We thank DNA Sequencing & Services

(MRC I PPU, School of Life Sciences, University of Dundee, Scotland,

www.dnaseq.co.uk) for DNA sequencing. The comments from three anonymous

reviewers on a previous draft substantially improved the manuscript.

References

1. Allsopp R, Baldry DAT, Rodrigues C. **The influence of game animals on the distribution and feeding habits of Glossina pallidipes in the Lambwe Valley.** *B World Health Organ.* 1972; **47**(6):795-809.
2. Altschul SF, Gish W, Miller W, *et al.*: **Basic Local Alignment Search Tool.** *J Mol Biol.* 1990; **215**(3):403-10.
3. Anderson NE, Mubanga J, Fevre EM, *et al.*: **Characterisation of the Wildlife Reservoir Community for Human and Animal Trypanosomiasis in the Luangwa Valley, Zambia.** *Plos Neglect Trop D.* 2011; **5**(6)
4. Auty H, Anderson NE, Picozzi K, *et al.*: **Trypanosome diversity in wildlife species from the serengeti and Luangwa Valley ecosystems.** *PLoS Negl Trop Dis.* 2012; **6**(10):e1828.
5. Auty H, Cleaveland S, Malele I, *et al.*: **Quantifying heterogeneity in host-vector contact: tsetse (*Glossina swynnertoni* and *G. pallidipes*) host choice in Serengeti National Park, Tanzania.** *PLoS One.* 2016; **11**(10):e0161291.
6. Auty H, Morrison LJ, Torr SJ, *et al.*: **Transmission dynamics of Rhodesian sleeping sickness at the interface of wildlife and livestock areas.** *Trends in Parasitology.* 2016; **32**(8):608-21.
7. Barrow LN, Allen JM, Huang X, *et al.*: **Genomic sequence capture of haemosporidian parasites: Methods and prospects for enhanced study of host-parasite evolution.** *Molecular Ecology Resources.* 2019; **19**(2):400-10.
8. Bates D, Mächler M, Bolker B, *et al.*: **Fitting Linear Mixed-Effects Models Using lme4.** *Journal of Statistical Software.* 2015; **67**(1):1-48.
9. Baylis M. **Effect of defensive behaviour by cattle on the feeding success and nutritional state of the tsetse fly, *Glossina pallidipes* (Diptera: Glossinidae).** *Bulletin of Entomological Research.* 1996; **86**(4):329-36.
10. Baylis M, Nambiro CO. **The effect of cattle infection by *Trypanosoma congolense* on the attraction, and feeding success, of the tsetse fly *Glossina pallidipes*.** *Parasitology.* 1993; **106 (Pt 4)**(4):357-61.
11. Bett B, Irungu P, Nyamwaro SO, *et al.*: **Estimation of tsetse challenge and its relationship with trypanosomosis incidence in cattle kept under pastoral production systems in Kenya.** *Veterinary Parasitology.* 2008; **155**(3–4):287-98.
12. Bitome-Essono PY, Ollomo B, Arnathau C, *et al.*: **Tracking zoonotic pathogens using bloodsucking flies as 'flying syringes'.** *Elife.* 2017; **6**
13. Boakye DA, Tang J, P. T, *et al.*: **Identification of bloodmeals in haematophagous Diptera by cytochrome B heteroduplex analysis.** *Medical and veterinary entomology.* 1999; **13**:282-87.
14. Bouyer J, Pruvot M, Bengaly Z, *et al.*: **Learning influences host choice in tsetse.** *Biology Letters.* 2007; **3**:113-16.
15. Brightwell R, Dransfield RD, Kyorku C. **Development of a low-cost tsetse trap and odour baits for *Glossina pallidipes* and *G. longipennis* in Kenya.** *Medical and veterinary entomology.* 1991; **5**(153-164)
16. Brightwell R, Dransfield RD, Stevenson P, *et al.*: **Changes over twelve years in populations of *Glossina pallidipes* and *Glossina longipennis* (Diptera: Glossinidae) subject to varying trapping pressure at Nguruman, south-west Kenya.** *Bulletin of Entomological Research.* 1997; **87**:349-70.
17. Channumsin M, Ciosi M, Masiga D, *et al.*: **Sodalis glossinidius presence in wild tsetse is only associated with presence of trypanosomes in complex**

interactions with other tsetse-specific factors. *BMC Microbiology*. 2018; **18**(1):163.

18. Channumsin M, Ciosi M, Masiga DK, *et al.*: **Supplementary Information for Blood meal analysis of tsetse flies (Glossina pallidipes: Glossinidae) reveals higher host fidelity on wild compared with domestic hosts.** 2021;

19. Clausen PH, Adeyemi I, Bauer B, *et al.*: **Host preferences of tsetse (Diptera: *Glossinidae*) based on bloodmeal identifications.** *Medical and veterinary entomology*. 1998; **12**(2):169-80.

20. England EC, Baldry DAT. **The hosts and trypanosome infection rates of *Glossina pallidipes* in the Lambwe and Roo valleys.** *B World Health Organ*. 1972; **47**(6):785-88.

21. England EC, Baldry DAT. **Observations on relative attractiveness to *Glossina pallidipes* of different animal baits, a tsetse Trap, and a fly-round patrol.** *B World Health Organ*. 1972; **47**(6):789-93.

22. Farikou O, Njiokou F, Simo G, *et al.*: **Tsetse fly blood meal modification and trypanosome identification in two sleeping sickness foci in the forest of southern Cameroon.** *Acta Tropica*. 2010; **116**(1):81-88.

23. Gaithuma A, Yamagishi J, Hayashida K, *et al.*: **Blood meal sources and bacterial microbiome diversity in wild-caught tsetse flies.** *Scientific Reports*. 2020; **10**(1):5005.

24. Gariepy TD, Lindsay R, Ogden N, *et al.*: **Identifying the last supper: utility of the DNA barcode library for bloodmeal identification in ticks.** *Molecular Ecology Resources*. 2012; **12**(4):646-52.

25. Geigy R, Kauffman.M, Jenni L. **Wild Mammals as Reservoirs for Rhodesian Sleeping Sickness in Serengeti, 1970-71**. *T Roy Soc Trop Med H*. 1973; **67**(2):284-86.

26. Geigy R, Mwambu PM, Kauffman.M. **Sleeping sickness survey in Musoma District, Tanzania. IV. Examination of wild mammals as a potential reservoir for *T. rhodesiense*.** *Acta Tropica*. 1971; **28**:211-20.

27. Ginestet C. **ggplot2: Elegant Graphics for Data Analysis.** *J R Stat Soc a Stat*. 2011; **174**:245-45.

28. Haouas N, Pesson B, Boudabous R, *et al.*: **Development of a molecular tool for the identification of leishmania reservoir hosts by blood meal analysis in the insect vectors.** *Am J Trop Med Hyg*. 2007; **77**(6):1054-59.

29. Hargrove JW. **The effect of human presence on the behaviour of tsetse (*Glossina spp.*) (Diptera, *Glossinidae*) near a stationary ox.** *Bulletin of Entomological Research*. 1976; **66**(1):173-78.

30. Harraca V, Syed Z, Guerin PM. **Olfactory and behavioural responses of tsetse fies, *Glossina spp.*, to rumen metabolites.** *Journal of Comparative Physiology A*. 2009; **195**(9):815-24.

31. Haydon DT, Cleaveland S, Taylor LH, *et al.*: **Identifying Reservoirs of Infection: A Conceptual and Practical Challenge.** *Emerging Infectious Diseases*. 2002; **8**(12):1468-73.

32. Hazkani-Covo E, Zeller RM, Martin W. **Molecular poltergeists: mitochondrial DNA copies (numts) in sequenced nuclear genomes.** *PLoS genetics*. 2010; **6**(2):e1000834.

33. Hebert PDN, Cywinska A, Ball SL, *et al.*: **Biological identifications through DNA barcodes.** *Proceedings of the Royal Society of London Series B-Biological Sciences*. 2003; **270**(1512):313-21.

34. Heisch RB, McMahon JP, Mansonbahr PE. **The isolation of Trypanosoma rhodesiense from a bushbuck.** *Br Med J*. 1958; **2**(5106):1203-04.

35. Hoffmann C, Stockhausen M, Merkel K, *et al.*: **Assessing the feasibility of fly based surveillance of wildlife infectious diseases.** *Scientific Reports.* 2016; **6**:37952.

36. Ivanova NV, zemlak TS, Hanner rH, *et al.*: **Universal primer cocktails for fish DNA barcoding.** *Molecular Ecology Notes.* 2007; **7**:544–48.

37. Jackson CHN. **An artificially isolated generation of tsetse flies (*Diptera*).** *Bulletin of Entomological Research.* 1946; **37**(02):291-99.

38. Jenni L, Molyneux DH, Livesey JL, *et al.*: **Feeding-behavior of tsetse flies infected with Salivarian trypanosomes.** *Nature.* 1980; **283**(5745):383-85.

39. Kent RJ. **Molecular methods for arthropod bloodmeal identification and applications to ecological and vector-borne disease studies.** *Molecular Ecology Resources.* 2009; **9**(1):4-18.

40. Kocher TD, Thomas WK, Meyer A, *et al.*: **Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers.** *Proceedings of the National Academy of Sciences of the United States of America.* 1989; **86**:6196-200.

41. Kubi C, Van den Abbeele J, De Deken R, *et al.*: **The effect of starvation on the susceptibility of teneral and non-teneral tsetse flies to trypanosome infection.** *Medical and veterinary entomology.* 2006; **20**(4):388-92.

42. Lah EFC, Ahamad M, Haron MS, *et al.*: **Establishment of a molecular tool for blood meal identification in Malaysia.** *Asian Pacific Journal of Tropical Biomedicine.* 2012; **2**(3):223-27.

43. Langridge WP, Kernaghan RJ, Glover PE. **A review of recent knowledge of the ecology of the main vectors of trypanosomiasis.** *B World Health Organ.* 1963; **28**:671-701.

44. Le S, Josse J, Husson F. **FactoMineR: An R package for multivariate analysis.** *Journal of Statistical Software.* 2008; **25**(1):1-18.

45. Leak SGA. Tsetse biology and ecology. The role in the epidemiology and control of trypanosmiasis. London: CABI Publishing, 1998.

46. Leigh JW, Bryant D. **POPART: full-feature software for haplotype network construction.** *Methods Ecol Evol.* 2015; **6**(9):1110-16.

47. Librado P, Rozas J. **DnaSP v5: a software for comprehensive analysis of DNA polymorphism data.** *Bioinformatics.* 2009; **25**(11):1451-2.

48. Lord JS, Lea RS, Allan FK, *et al.*: **Assessing the effect of insecticide-treated cattle on tsetse abundance and trypanosome transmission at the wildlife-livestock interface in Serengeti, Tanzania.** *Plos Neglect Trop D.* 2020; **14**(8):e0008288.

49. Lord JS, Torr SJ, Auty HK, *et al.*: **Geostatistical models using remotely-sensed data predict savanna tsetse decline across the interface between protected and unprotected areas in Serengeti, Tanzania.** *J Appl Ecol.* 2018; **55**(4):1997-2007.

50. Lyimo IN, Ferguson HM. **Ecological and evolutionary determinants of host species choice in mosquito vectors.** *Trends in Parasitology.* 2009; **25**(4):189-96.

51. Mbahin N, Affognon H, Andoke J, *et al.*: **Parasitological prevalence of bovine trypanosomiasis in Kubo Division of Kwale Country of Coastal: baseline survey** *American Journal of Animal and Veterinary Sciences.* 2013; **8**(1):28-36.

52. McFadden D. **The measurement of urban travel demand.** *Journal of Public Economics.* 1974; **3**(4):303-28.

53. Metsky HC, Siddle KJ, Gladden-Young A, *et al.*: **Capturing sequence diversity**
in metagenomes with comprehensive and scalable probe design. *Nat*
Biotechnol. 2019; **37**(2):160-+.
54. Muturi CN, Ouma JO, Malele, II, *et al.*: **Tracking the feeding patterns of**
tsetse flies (Glossina genus) by analysis of bloodmeals using
mitochondrial cytochromes genes. *PLoS One.* 2011; **6**(2):e17284.
55. Njenga SM, Mwandawiro CS, Muniu E, *et al.*: **Adult population as potential**
reservoir of NTD infections in rural villages of Kwale district, Coastal
Kenya: implications for preventive chemotherapy interventions policy.
Parasites & Vectors. 2011; **4**(1):1-6.
56. Njiru ZK, Constantine CC, Guya S, *et al.*: **The use of ITS1 rDNA PCR in**
detecting pathogenic African trypanosomes. *Parasitology Research.* 2005;
95(3):186-92.
57. OIE. Trypanosomosis (tsetse-transmitted). April 2013 ed, 2013.
58. Okoth SO, Kokwaro ED, Kirugu JM, *et al.*: *Glossina pallidipes* **and host**
interactions: implications of host preference on tranmission Rick of
Rhodesian Sleeping Sickness in Kenya. *Trends in Applied Sciences*
Research. 2007; **2**(5):386-94.
59. Omondi D, Masiga DK, Ajamma YU, *et al.*: **Unraveling host-vector-arbovirus**
Interactions by two-gene high resolution melting mosquito bloodmeal
analysis in a Kenyan wildlife-livestock Interface. *PloS one.* 2015;
10(7):e0134375-e75.
60. Pradhan V, Kamble Y, Ladniya V, *et al.*: **A overview of species identification**
by DNA barcoding. *International Journal of Current Microbiology and*
Applied Sciences 2015; **4**(4):127-40.
61. **Se-Al: Sequence Alignment Editor** [program]. Oxford, UK: Department of
Zoology, University of Oxford. 1996.
62. Reeves L, Gillett-Kaufman J, Kawahara A, *et al.*: **Barcoding blood meals: New**
vertebrate-specific primer sets for assigning taxonomic identities to host
DNA from mosquito blood meals. *PLoS Neglected Tropical Disease.* 2018;
12(8):e0006767.
63. Schofield S, Torr SJ. **A comparison of the feeding behaviour of tsetse and**
stable flies. *Medical and veterinary entomology.* 2002; **16**(2):177-85.
64. Schratz P. **R package 'oddsratio': Odds ratio calculation for GAM(M)s &**
GLM(M)s, version: 1.0.2. 2017;
65. Snow WF, Tarimo SA, Staak C, *et al.*: **The feeding-habits of the Tsetse,**
Glossina pallidipes **Austen on the South Kenya Coast, in the context of**
its host range and trypanosome infection-rates in other parts of East-
Africa. *Acta Tropica.* 1988; **45**(4):339-49.
66. Spath J. **Feeding patterns of three sympatric tsetse species (Glossina spp,)**
(Diptera : Glossinidae) in the preforest zone of Cote d'Ivoire. *Acta*
Tropica. 2000; **75**(1):109-18.
67. Team RC. **R: A Language and Environment for Statistical Computing. R**
Foundation for Statistical Computing. Vienna2016.
68. Torr SJ, Wilson PJ, Schofield S, *et al.*: **Application of DNA markers to identify**
the individual-specific hosts of tsetse feeding on cattle. *Medical Vet*
Entomology. 2001; **15**(1):78 - 86.
69. Turner DA. **The population ecology of** *Glossina pallidipes* **Austen (Diptera:**
Glossinidae) **in the Lambwe Valley, Kenya. I. Feeding behaviour and**
activity patterns. *Bulletin of Entomological Research.* 1987; **77**(02):317-33.
70. Vale GA. **Feeding responses of tsetse flies (Diptera-Glossinidae) to stationary**
hosts. *Bulletin of Entomological Research.* 1977; **67**(4):635-49.

71. Viana M, Mancy R, Biek R, *et al.*: **Assembling evidence for identifying**
reservoirs of infection. *Trends in Ecology & Evolution.* 2014; **29**(5):270-79.
72. Weitz B. **The feeding habits of *Glossina*.** *Bulletin of World Health*
Organization. 1963; **28**:711-29.
73. Welburn SC, Picozzi K, Fevre EM, *et al.*: **Identification of human-infective**
trypanosomes in animal reservoir of sleeping sickness in Uganda by
means of serum-resistance-associated (SRA) gene. *Lancet.* 2001;
358(9298):2017-19.
74. Welburn SC, Picozzi K, Kaare M, *et al.*: **Control options for human sleeping**
sickness in relation to the animal reservoir of disease. *Conservation and*
Development Interventions at the Wildlife/Livestock Interface. 2005:55-61.
75. Wongserepipatana M. **Prevalence and associations of *Trypanosoma* spp. and**
***Sodalis glossinidius* with intrinsic factors of tsetse flies.** University of
Glasgow. 2016.
869
870

871