Individual Variation in Brain Network Topography is Linked to Schizophrenia

Symptomatology

Running title: Network Topography in Schizophrenia

Nawaz, Uzma*[12], Lee, Ivy*[12]; Beermann, Adam[12]; Eack, Shaun[4]; Keshavan, Matcheri[12];

Brady Jr., Roscoe^[123].

[1] Department of Psychiatry, Beth Israel Deaconess Medical Center, Boston, MA, United

States.

[2] Department of Psychiatry, Harvard Medical School, Boston, MA, USA.

[3] Schizophrenia and Bipolar Disorder Program, McLean Hospital, Belmont, MA, USA.

[4] University of Pittsburgh, School of Social Work and Department of Psychiatry,
Pittsburgh, PA, USA.

* These authors contributed equally to this manuscript

^ Corresponding author

Address correspondence to:

Roscoe Brady Jr.

75 Fenwood Road

Boston, MA, 02115

Phone: (617)-754-1261

Fax: (617)-754-1250

robrady@bidmc.harvard.edu

23
24 This work is not peer reviewed
25
26
27

Abstract

Background: Resting state fMRI (rsfMRI) demonstrates that the brain is organized into distributed networks. Numerous studies have examined links between psychiatric symptomatology and network functional connectivity. Traditional rsfMRI analyses assume that the spatial organization of networks is invariant between individuals. This dogma has recently been overturned by the demonstration that networks show significant variation between individuals. We tested the hypothesis that previously observed relationships between schizophrenia negative symptom severity and network connectivity are actually due to individual differences in network spatial organization.

Methods: 44 participants diagnosed with schizophrenia underwent rsfMRI scans and clinical assessments. A multivariate pattern analysis determined how whole brain functional connectivity correlates with negative symptom severity at the individual voxel level.

Results: Brain connectivity to a region of the right dorso-lateral pre-frontal cortex correlates with negative symptom severity. This finding results from individual differences in the topographic distribution of two networks: the default mode network (DMN) and the task positive network (TPN). Both networks demonstrate strong (r~0.49) and significant (p<0.001) relationships between topography and symptom severity. For individuals with low symptom severity, this critical region is part of the DMN. In highly symptomatic individuals, this region is part of the TPN.

Conclusion: Previously overlooked individual variation in brain organization is tightly linked to differences in schizophrenia symptom severity. Recognizing critical links between network topography and pathological symptomology may identify key circuits

52 that underlie cognitive and behavioral phenotypes. Individual variation in network

53 topography likely guides different responses to clinical interventions that rely on

54 anatomical targeting (e.g. TMS).

Introduction

The brain is organized into distributed resting-state networks whose spatial distribution are typically visualized using group averaging[1, 2]. Individual variation in network topography among healthy control participants has been demonstrated previously[3-8]. Until very recently, the relevance of individual network topography to cognitive and behavioral variation has been undetermined but recent reports have linked individual topography to several cognitive and behavioral phenotypes[9-11]. To date these reports have examined these phenotypes in healthy populations[9, 10] or have examined normative processes (e.g. social cognition) in clinical populations[11]. However, the following question has been left unaddressed: Is individual network topography linked to clearly pathological (i.e. disease-specific) processes such as psychiatric symptomatology?

A very provocative idea espoused by Bijsterbosch et al. is that previously reported cross-subject variation in functional connectivity are mis-interpretations of variation in the spatial organization of brain networks[9]. We therefore sought to simultaneously address two aims: First, to determine if psychiatric symptomatology is linked to brain network topography, and second, to re-examine previously reported functional connectivity findings to determine if these findings are instead explained by individual topographic variation.

In a previous manuscript we used an unbiased, data-driven approach to find functional MRI correlates of negative symptom severity[12]. Although psychotic symptoms (e.g. hallucinations, delusions) are the most readily recognizable symptoms of schizophrenia, they are not the best predictors of functional status (e.g. independent

housing and employment). Rather, the severity of negative symptoms such as

amotivation, expressive deficits, and anhedonia best predict functional outcomes and

overall quality of life of individuals with schizophrenia[13-15]. In our prior manuscript, we

observed that the most significant resting state functional connectivity correlate of

negative symptom severity was connectivity between the bilateral dorsolateral prefrontal

cortex and the Default Mode Network (DMN). Here we re-examine this result in light of

newly discovered relationships between functional connectivity and network topography.

We examined whether negative symptom severity is linked to individual network

topography. Specifically, we looked at the spatial organization of two distributed brain

networks in relation to a critical region in the DLPFC.

89
90   **Methods**
91
92   **Participants**
93
Participants included in the following analysis were recruited for an Institutional

Review Board approved clinical trial (NCT01561859) at Beth Israel Deaconess Medical

Center (Boston, MA) and University of Pittsburg (Pittsburg, PA). Prior to enrollment in

this longitudinal, randomized study, all participants gave their written informed consent.

The data included in the analysis were collected at baseline (i.e. prior to trial

intervention). Clinical and imaging data from 80 participants with schizophrenia were

analyzed. After selection for very low-motion scans, a total of 44 participants were

retained for further analysis.

Diagnosis was determined using the Structured Clinical Interview for the DSM-IV

(SCID)[16]. Participants were assessed by raters trained in the administration and scoring

of the SCID. Inclusion criteria for all participants were: (1) age 18-45 years; (2) current

IQ $\geq$ 80 as assessed using the WASI-II[17]; and (3) the ability to read (sixth grade level or

higher) and speak fluent English. Additional inclusion criteria for participants with a

psychotic disorder were (1) a diagnosis of schizophrenia or schizoaffective disorder

verified using the SCID interview[18]; (2) time since first psychotic symptom of $\leq$ 8 years;

and (3) clinically stabilized on antipsychotic medication (assessed via SCID in

consensus conferences). Healthy comparison (HC) participants did not meet criteria for

any Axis I psychiatric disorder currently or historically.

Exclusion criteria were (1) significant neurological or medical disorders that may

produce cognitive impairment (e.g., seizure disorder, traumatic brain injury); (2)

persistent suicidal or homicidal behavior; (3) a recent history of substance abuse or

dependence (within the past 3 months); (4) any MRI contraindications and (5) decisional

incapacity requiring a guardian.

Demographic and clinical data for participants included in the analysis can be

found in Table 1.

137 **Table 1. Demographic and Clinical Information for Participants**
138

	Participants (n = 44)
Mean Age (SD)	24.40 (±4.52)
Sex	
Male	29
Female	15
Race	
Caucasian	27
African American	6
Other	11
Diagnosis	
Schizophrenia (SZ)	35
Schizoaffective Disorder (SA)	9
Mean CPZE mg (SD)	306.03 (±232.14)
Mean FD mm (SD)	0.14 (±0.08)
Mean SANS Scores	33.73

139
140 Abbreviations
141 CPZE- Chlorpromazine Equivalents
142 FD- Framewise Displacement
143 SD- Standard Deviation
144

Behavioral Assessment

Negative symptom severity was assessed by the Scale for the Assessment of

Negative Symptoms (SANS)[19].

MRI Data Collection

Boston site: Data were acquired on 3T Siemens Trio (TIM upgrade) scanners using a

standard head coil. The echoplanar imaging parameters were as follows: repetition

time, 3000 milliseconds; echo time, 30 milliseconds; flip angle, 85°; 3 x 3 x 3-mm

voxels; and 47 axial sections collected with interleaved acquisition and no gap.

Structural data included a high-resolution T1 image. In addition, all participants

underwent a resting fMRI run with the instructions 'remain still, stay awake, and keep

your eyes open'. Each functional run lasted 6.2 minutes (124 time points).

Pittsburgh site: Data were acquired on a 3T Siemens Verio scanner using a standard

head coil. The echoplanar imaging parameters were as follows: repetition time, 3000

milliseconds; echo time, 30 milliseconds; flip angle, 85°; 3 x 3 x 3-mm voxels; and 45

axial sections collected with interleaved acquisition and no gap. Structural data included

a high-resolution T1 image (voxel size of 1.0 × 1.0 × 1.2 mm, TR 2300 ms, TI = 900 ms,

TE = 2.89 ms, flip angle = 90°, FOV = 256 mm, 256 × 256 matrix, 160 slices, slice

thickness = 1.2 mm). In addition, all participants underwent a resting fMRI run that

lasted 6.2 minutes (124 time points).

Functional MRI Data Pre-Processing

The rsfMRI data was processed using DPABI[20]. The first 4 images were removed

to reduce the effects of scanner signal stabilization. Scans with head motion exceeding

3 mm or 3° of maximum rotation through the resting-state run were discarded.

Functional and structural images were co-registered. Structural images were then

normalized and segmented into gray, white and CSF partitions using the DARTEL

technique[21]. A Friston 24-parameter model[22] was used to regress out head motion

effects from the realigned data. CSF and white matter signals, global signal and, the

linear trend were also regressed out. The decision to regress the global signal was

based on prior demonstration that the combination of this step plus volume-wise

'scrubbing' for head 'micromovements' is an effective strategy for removal of motion

artifacts[23].

After realigning, slice timing correction, and co-registration, framewise

displacement (FD) was calculated for all resting state volumes[24]. All volumes within a

scan with a FD greater than 0.2-mm were censored. Any scan with > 50% of volumes

requiring censoring was discarded. After nuisance covariate regression, the resultant

data were band-pass filtered to select low frequency (0.01 – 0.08Hz) signals. Filtered

data were normalized by DARTEL into MNI space and then smoothed by a Gaussian

kernel of 8mm[3] full-width at half maximum (FWHM). Voxels within a gray matter mask

were used for further analyses.

After preprocessing, 36 scans were removed from analysis on the basis of head

motion leaving a total of 44 participants with schizophrenia or schizoaffective disorder

across all sites in the study (Table 1).

Imaging Analysis

Multivariate Distance Matrix Regression:

We performed a connectome-wide association study using multivariate distance

matrix regression (MDMR) as originally described in Shehzad et al.[25] Briefly, MDMR

tests every voxel to determine if whole-brain connectivity to that voxel is more similar in

individuals with a similar SANS score than in individuals with a dissimilar SANS score. A

step-wise description of MDMR can be found in Supplementary Material.

This analysis identifies anatomical regions where SANS score is significantly

related to functional connectivity. At each brain voxel, MDMR calculates a distance

metric r between subjects by first calculating the temporal correlation in BOLD signal

between that voxel and every other brain voxel. The generated set of temporal

correlations is then correlated with the analogous set of temporal correlations from a

different subject. The resulting between-subject correlation determines a distance metric

that is a measure of between-subject similarity (or dissimilarity) in terms of functional

connectivity at that voxel. Notably, this process disregards spatial information about the

voxels that gave rise to between-individual distances. For example, two individuals may

be very distant (dissimilar) in the functional connectivity of a voxel in the precuneus, but

is their dissimilarity driven by differences in precuneus connectivity to the mPFC or

temporal lobe or parietal lobe or all of the above? MDMR as implemented by Shehzad

et al. does not display this information[25].

To visualize this spatial information requires follow-up seed-based connectivity

analysis. Shehzad et al. and others have termed this follow-up analysis 'post-hoc'

testing to make clear that this should not be considered independent hypothesis testing

nor should it be considered independent validation of the original MDMR finding[25-28]. As

in these prior studies, we conducted the MDMR analysis to find anatomical regions

where connectivity significantly varied with SANS score and then conducted follow-up

seed-based connectivity analysis to examine the spatial distribution of these

connectivity differences.

ROI based connectivity Analysis:

Following the creation and analysis of distance maps in MDMR, DPABI was used

to extract the time course of the BOLD signals from rsfMRI scans in reference to a

specific ROI identified in the MDMR process. Whole brain connectivity maps were

generated and the resulting z-transformed Pearson's correlation coefficients were used

in the program, SPM12 ("SPM - Statistical Parametric Mapping,"

http://www.fil.ion.ucl.ac.uk/spm). For the ROI generated from MDMR above, we

regressed the maps against SANS score to generate spatial maps of how whole brain

functional connectivity to the ROI varies with SANS scores. SPM generated T contrasts

maps were created using Caret[31]. Correlations between ROI to ROI connectivity values

and SANS score were calculated in R.

Results

Resting State Connectivity to the Bilateral DLPFC Is Correlated to Negative

Symptom Severity

From the MDMR analysis of 44 participants, two regions were identified where

resting state functional connectivity correlated to SANS scores (Figure 1). Functional

connectivity to the bilateral middle frontal gyrus (Brodmann Area 9) in the dorsolateral

prefrontal cortex covaried significantly with negative symptom severity. The right

dorsolateral prefrontal cortex (peak voxel z-stat 3.82 at MNI coordinates: x36, y24, z30)

demonstrated a more significant relationship between functional connectivity and

negative symptom severity than the left (peak voxel z-stat 2.95 at MNI coordinates: x-

33, y30, z42) (Figure 1F).

Figure 1. Multivariate Distance Matrix Regression identifies brain voxels whose
functional connectivity varies with negative symptom severity.
MDMR procedure: **A.** rsfMRI and SANS are collected from each participant. **B.** For each
participant a functional connectivity map is generated to an individual voxel. **C.**
Voxelwise temporal correlations between participants are used to generate a Pearson's
correlation *r* and a distance metric *d*. This is repeated for all participants to generate a
matrix of between subject distances. **D.** The distance matrix is centered and an
ANOVA-like test is used to generate an F-statistic to assess the relationship between a
predictor variable (SANS score) and dissimilarities in functional connectivity at that
voxel. **E.** This process is repeated for every voxel. This results in a whole brain map of
how significantly functional connectivity is related to emotional intelligence. Permutation
testing then identifies whole-brain significant clusters in connectivity-SANS score
relationships. **F.** In our sample of 44 participants with schizophrenia or schizoaffective
disorder, we identified the bilateral DLPFC (Brodmann Area 9) as regions where
resting-state connectivity correlated significantly with SANS score. In this image,
connectivity is thresholded at a voxelwise level of p<.005 and extent threshold of p<.05.
256

The Spatial Distribution of Symptom-Correlated Connectivity to the DLPFC

Identifies the Default Mode Network and Task-Positive Network

We performed follow-up analysis using the right DLPFC in a seed-based

connectivity analysis to determine the spatial distribution and directionality of

connectivity that gave rise to this result. This analysis revealed two apparent patterns of

functional connectivity: higher SANS (i.e. more severe symptoms) correlated with

increasing intrinsic connectivity between the right DLPFC and the "task-positive

network" (TPN) (Figure 2A)[32]. Inversely, lower negative symptom severity correlated

with increasing functional connectivity between the right DLPFC and the "default mode

network" (DMN) (Figure 2B)[1].

Figure 2. In a follow-up analysis, a seed-region was placed in the right dorsolateral
prefrontal cortex (ROI) in all subjects and then this seed-based connectivity map was
correlated with SANS scores to identify locations where increasing connectivity to
dorsolateral prefrontal cortex corresponds to more severe symptoms (red) and
decreased connectivity corresponds to less severe symptoms (blue). Here, we observe

275 that right DLPFC connectivity to the Task-Positive Network (TPN) is correlated with
276 higher SANS score (A) and right DLPFC connectivity to the Default Mode Network
277 (DMN) is correlated with lower SANS score (B). Color bar = T-statistic.
278

Individual Variation in Network Topography Explains Connectivity-Symptom

Relationships in the DLPFC

We plotted the relationship between negative symptom severity and functional

connectivity for both networks as a scatter plot in Figure 3C & 3D. As expected from the

maps in Figure 2A & 2B, we observed that a higher SANS score was correlated with

increased functional connectivity between the right DLPFC and the TPN and decreased

connectivity to the DMN. Both scatter plots of connectivity cross the axis of 0

connectivity i.e. activity in this DLPFC region is intrinsically correlated to the DMN in

some individuals and is intrinsically correlated to the TPN in others. Furthermore, this

network membership is organized along an axis of negative symptom severity with this

critical zone included in the spatial distribution of the DMN in the least symptomatic

individuals and, in the most symptomatic individuals this region is included in the TPN.

291
Figure 3. Negative Symptom Severity is Linked to Network Topography
A. Plot of SANS score (y-axis) and functional connectivity between the rDLPFC ROI
and the DMN network. **B.** Plot of SANS score (y-axis) and functional connectivity
between the rDLPFC and TPN network. Notably, both graphs demonstrate correlations
that cross the reference line of null functional connectivity, meaning that the network
association of this region appears to shift from being a member of the DMN to a
member of the TPN as one moves along the distribution from less symptomatic
individuals to more symptomatic individuals. Network topography correlates were highly
significant (p<.001) for both networks.

We sought to re-visualize the relationship between symptom severity, the critical

right DLPFC region, and the whole-brain spatial organization of the DMN. To facilitate

this, participants were grouped by symptom severity. Forty-four participants were split

into three groups based on their SANS score with group 1 consisting of the lowest

scorers (i.e. least symptomatic) and 3 with the highest scores (most symptomatic). We

examined the spatial distribution of the DMN in each group and visualized this in

relation to the previously identified right DLPFC (Figure 4). As would be expected from

Figure 3, we observe the spatial extent of the DMN does not include the identified

DLPFC node in the most symptomatic participants. The DLPFC node of the DMN is

organized along an anterior-posterior spatial gradient organized along a least

symptomatic to most symptomatic axis. The spatial extent of the TPN (data not shown)

demonstrated the inverse relationship i.e. the TPN extending into the DLPFC critical

region in the most symptomatic participants.

Figure 4. Whole Brain DMN Spatial Topography in Relation to Negative Symptom
Severity. Shown is the distribution of DMN correlated voxels in the participants grouped
by SANS score i.e. lowest (least symptomatic) 20%, middle 60% and highest (most
symptomatic) 20% of the sample. In least symptomatic participants the rDLPFC was
part of the DMN. In the most symptomatic (Group 3), this region was a part of the TPN.

Groups with intermediate severity were intermediate in their connectivity strength to the
DMN. Scale: Connectivity strengths (Pearson's *r*) less than 0.25 are shown in dark red.
Higher strengths in bright red. The MDMR identified right DLPFC region is shown in
yellow.
325

Discussion

We conducted an entirely data-driven connectome wide study examining the

relationship between negative symptoms severity and functional connectivity. Bilateral

dorsolateral prefrontal cortex regions were identified as having the most significant

relationship between functional connectivity and psychopathology. Follow-up testing

demonstrated that this result is driven by the correlation between DLPFC activity and

two distributed brain networks: the Default Mode Network (DMN) and the Task Positive

Network (TPN).

Analyses that assume inter-individual homogeneity in the spatial distribution of

these networks would misinterpret this result as the product of individual variation in the

temporal correlation in activity between these networks and the DLPFC. Instead, we

observe that this result is entirely driven by individual variation in spatial organization

(i.e. network topography) within this region.

More specifically, the degree to which this region was a member of one of two

large scale resting-state networks, the DMN and TPN, was directly linked to symptom

severity.

The implication of this result is that recently observed links between variation in

network topography and normative cognitive / behavioral phenotypes also apply to

disease-specific symptoms[9-11]. More broadly, this result lends support to the hypothesis

proposed by by Bijsterbosch et al. that previously reported cross-subject variations in

functional connectivity are in fact misinterpretations of variation in the spatial

organization of brain networks[9].

It is notable that in both this manuscript and our previous report on social

cognition[11], we demonstrated that the *most significant* connectivity-phenotype

relationships in the entire brain were driven by variation in network topography. The

increasing prevalence of fully data-driven connectivity studies will determine if this (i.e.

spatial variation driving the most significant connectivity-phenotype relationships) is a

general rule.

Returning to this specific result, what are the implications of schizophrenia

symptoms being linked to network organization within a narrowly circumscribed portion

of the DLPFC? The DLPFC has been implicated in an enormous variety of cognitive

processes and thus links between DLPFC organization and negative symptomatology

have face validity. From a pragmatic/ experimental perspective, our finding has clearer

significance: An enormous body of literature has explored the activation of the DLPFC

in a variety of neuroimaging tasks. Although our result was generated using task-free

connectivity data, the at-rest spatial organization of the TPN would be expected to

represent the spatial organization of the networks activated under task conditions. The

implication of our finding is that there is considerable variability in the organization of

this network in a population diagnosed with schizophrenia. This would almost certainly

confound task-based studies that rely on group-based comparisons. Put simply: In

participants with schizophrenia, the DLPFC region activated by a task may lie in very

different MNI coordinates even after normalization to a standard template. This result

would likely be interpreted as hypoactivation simply because of a lack of overlap among

a group when compared to a groups of healthy comparison participants.

Our results also have clear implications for therapeutic interventions that rely on

the precise spatial targeting. In a previous publication, we demonstrated that repeated

application of intermittent Theta Burst Stimulation (iTBS) targeted to the cerebellum can

restore connectivity between cerebellar and DLPFC nodes of the DMN and, in the

process, can reduce negative symptom severity[12]. In contrast, one of the largest studies

of rTMS applied to the left DLPFC for negative symptom amelioration did not observe a

significant difference between sham and real rTMS[33]. A re-analysis of that trial

demonstrated that response in the real TMS group was linked to within-trial change in

brain structure with responders showing an increase in the volume of DMN-related

regions including hippocampus, parahippocampus, and precuneus[34].

We hypothesize that this result is directly linked to our network topography result

in the following manner: In Wobrock et al., rTMS was targeted to EEG position F3. This

scalp target lies directly adjacent to the left DLPFC region of substantial individual

network topography variation[33]. In the Wobrock et al. trial's selection of a participant

population with high negative symptom burden, the implication of our results presented

here would be that this target would overlie the DMN only in a subset of participants.

We hypothesize that rTMS response was partially determined by individual DMN

network topography: When the TMS coil position coincided with the topography of an

individual participant's DMN, symptom improvement and DMN network related

structural changes followed.

This logic, applied broadly, may explain individual variation in response to

anatomically targeted interventions in other disorders e.g. rTMS for depression. Prior

claims linking group-average connectivity maps of rTMS targets to antidepressant

response may in fact, represent movement of coil position from regions of high

individual variation in network topography to regions where a majority of patients show

shared topography of disease relevant networks[35]. This remains a conjecture as we

have not analyzed differences in network topography among depressed candidates for

TMS.

Finally, our work contains a cautionary note for imaging studies examining

connections between connectivity and individual phenotypes. The most frequently used

approaches to identifying individual-level network topography do so by generating

individual parcellations of the cortex using a "winner-take-all" approach i.e. spontaneous

activity in a given cortical region is frequently correlated with *multiple* networks. In a

"winner-take-all" or "hard" parcellation approach, the network "identity" assigned to that

region is that of the single network most strongly connected to that region. Subsequent

analyses then typically analyze results in terms of connectivity to that "winner" network.

While this approach allows for easy visualization of individually parcellated

connectomes, this approach *continues* to obfuscate links between network topography

and phenotypes. Specifically: The region we identify in this manuscript lies at the

intersection of several different networks. At the resolution used for rsfMRI imaging

these networks all show varying levels correlation within the voxels identified in our

DLPFC region. While the topography of two of these networks (DMN and TPN) are very

significantly linked to symptomatology, a winner-take-all analysis would actually have

assigned this parcel of cortex to the FPCN (Supplementary Material Figure 1). The

consequence of this is likely that subsequent analysis of connectivity-phenotype

relationships would be misinterpreted as inter-network connectivity i.e. FPCN-DMN and

FPCN-TPN connectivity correlations with symptomatology. This is, in fact, *exactly* what

was described in a recent publication (Figure 2, Baker et al.[36]) which uses an individual

"winner-take-all" parcellation approach in an independent dataset to elucidate

connectome- negative symptom relationships.

In summary, we demonstrate that inter-individual variation in disease phenotypes

is linked to individual network spatial organization. We present a model for how this

variation in cortical organization is likely to explain previously observed variation in

response to targeted neuromodulation. We also highlight how commonly used individual

parcellation approaches continue to obfuscate significant network topography-

phenotype relationships.

Acknowledgements
Funding:
This work was supported by the National Institutes of Health (K23 MH100623 and R01
MH116170 to RB, R01 MH92440 to SE, MK)

431
432

References

1. Yeo BT, Krienen FM, Sepulcre J, et al. The organization of the human cerebral cortex
estimated by intrinsic functional connectivity. *J Neurophysiol* Sep 2011;106(3):1125-
1165.
2. Power JD, Cohen AL, Nelson SM, et al. Functional network organization of the human
brain. *Neuron* Nov 17 2011;72(4):665-678.
3. Mueller S, Wang D, Fox MD, et al. Individual variability in functional connectivity
architecture of the human brain. *Neuron* Feb 6 2013;77(3):586-595.
4. Laumann TO, Gordon EM, Adeyemo B, et al. Functional System and Areal Organization
of a Highly Sampled Individual Human Brain. *Neuron* Aug 5 2015;87(3):657-670.

5. Poldrack RA, Laumann TO, Koyejo O, et al. Long-term neural and physiological
phenotyping of a single human. *Nat Commun* Dec 9 2015;6:8885.

6. Wang D, Buckner RL, Fox MD, et al. Parcellating cortical functional networks in
individuals. *Nat Neurosci* Dec 2015;18(12):1853-1860.

7. Gordon EM, Laumann TO, Gilmore AW, et al. Precision Functional Mapping of Individual
Human Brains. *Neuron* Aug 16 2017;95(4):791-807 e797.

8. Gordon EM, Laumann TO, Adeyemo B, Gilmore AW, Nelson SM, Dosenbach NUF,
Petersen SE. Individual-specific features of brain systems identified with resting state
functional correlations. *Neuroimage* Feb 1 2017;146:918-939.

9. Bijsterbosch JD, Woolrich MW, Glasser MF, Robinson EC, Beckmann CF, Van Essen DC,
Harrison SJ, Smith SM. The relationship between spatial configuration and functional
connectivity of brain regions. *Elife* Feb 16 2018;7.

10. Kong R, Li J, Orban C, et al. Spatial Topography of Individual-Specific Cortical Networks
Predicts Human Cognition, Personality, and Emotion. *Cereb Cortex* Jun 1
2019;29(6):2533-2551.

11. Ling G, Lee I, Guimond S, et al. Individual variation in brain network topology is linked to
emotional intelligence. *Neuroimage* Jan 8 2019;189:214-223.

12. Brady RO, Jr., Gonsalvez I, Lee I, et al. Cerebellar-Prefrontal Network Connectivity and
Negative Symptoms in Schizophrenia. *Am J Psychiatry* Jan 30
2019:appiajp201818040429.

13. Ho BC, Nopoulos P, Flaum M, Arndt S, Andreasen NC. Two-year outcome in first-episode
schizophrenia: predictive value of symptoms for quality of life. *Am J Psychiatry* Sep
1998;155(9):1196-1201.

14. Rabinowitz J, Levine SZ, Garibaldi G, Bugarski-Kirola D, Berardo CG, Kapur S. Negative
symptoms have greater impact on functioning than positive symptoms in schizophrenia:
analysis of CATIE data. *Schizophr Res* May 2012;137(1-3):147-150.

15. Robertson BR, Prestia D, Twamley EW, Patterson TL, Bowie CR, Harvey PD. Social
competence versus negative symptoms as predictors of real world social functioning in
schizophrenia. *Schizophr Res* Dec 2014;160(1-3):136-141.

16. First MB, New York State Psychiatric Institute. Biometrics Research D. *Structured clinical
interview for DSM-IV-TR axis I disorders : SCID-I*. New York, N.Y.: Biometrics Research
Dept., New York State Psychiatric Institutute; 2007.

17. Hays JR, Reas DL, Shaw JB. Concurrent validity of the Wechsler abbreviated scale of
intelligence and the Kaufman brief intelligence test among psychiatric inpatients.
Psychological reports 2002;90(2):355-359.

18. First MB. *Structured Clinical Interview for DSM-IV Axis I Disorders: Patient Edition
(February 1996 Final), SCID-I/P*: Biometrics Research Department, New York State
Psychiatric Institute; 1998.

19. Andreasen NC. The Scale for the Assessment of Negative Symptoms (SANS): conceptual
and theoretical foundations. *Br J Psychiatry Suppl* Nov 1989(7):49-58.

20. Yan CG, Wang XD, Zuo XN, Zang YF. DPABI: Data Processing & Analysis for (Resting-
State) Brain Imaging. *Neuroinformatics* Jul 2016;14(3):339-351.

21. Ashburner J. A fast diffeomorphic image registration algorithm. *Neuroimage* Oct 15
2007;38(1):95-113.

22.　Friston KJ, Williams S, Howard R, Frackowiak RS, Turner R. Movement-related effects in
fMRI time-series. *Magn Reson Med* Mar 1996;35(3):346-355.

23.　Yan CG, Cheung B, Kelly C, et al. A comprehensive assessment of regional variation in
the impact of head micromovements on functional connectomics. *Neuroimage* Aug 1
2013;76:183-201.

24.　Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. Spurious but systematic
correlations in functional connectivity MRI networks arise from subject motion.
Neuroimage Feb 1 2012;59(3):2142-2154.

25.　Shehzad Z, Kelly C, Reiss PT, et al. A multivariate distance-based analytic framework for
connectome-wide association studies. *Neuroimage* Jun 2014;93 Pt 1:74-94.

26.　Sharma A, Wolf DH, Ciric R, et al. Common Dimensional Reward Deficits Across Mood
and Psychotic Disorders: A Connectome-Wide Association Study. *Am J Psychiatry* Jan 31
2017:appiajp201616070774.

27.　Shanmugan S, Wolf DH, Calkins ME, et al. Common and Dissociable Mechanisms of
Executive System Dysfunction Across Psychiatric Disorders in Youth. *Am J Psychiatry*
May 1 2016;173(5):517-526.

28.　Satterthwaite TD, Vandekar SN, Wolf DH, et al. Connectome-wide network analysis of
youth with Psychosis-Spectrum symptoms. *Mol Psychiatry* Dec 2015;20(12):1508-1515.

29.　Zapala MA, Schork NJ. Multivariate regression analysis of distance matrices for testing
associations between gene expression patterns and related variables. *Proc Natl Acad Sci*
U S A Dec 19 2006;103(51):19430-19435.

30.　Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance
neuroimages. *Comput Biomed Res* Jun 1996;29(3):162-173.

31.　Van Essen DC, Drury HA, Dickson J, Harwell J, Hanlon D, Anderson CH. An integrated
software suite for surface-based analyses of cerebral cortex. *J Am Med Inform Assoc*
Sep-Oct 2001;8(5):443-459.

32.　Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME. The human brain
is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl*
Acad Sci U S A Jul 5 2005;102(27):9673-9678.

33.　Wobrock T, Guse B, Cordes J, et al. Left prefrontal high-frequency repetitive transcranial
magnetic stimulation for the treatment of schizophrenia with predominant negative
symptoms: a sham-controlled, randomized multicenter trial. *Biol Psychiatry* Jun 1
2015;77(11):979-988.

34.　Hasan A, Wobrock T, Guse B, et al. Structural brain changes are associated with
response of negative symptoms to prefrontal repetitive transcranial magnetic
stimulation in patients with schizophrenia. *Mol Psychiatry* Jun 2017;22(6):857-864.

35.　Fox MD, Buckner RL, White MP, Greicius MD, Pascual-Leone A. Efficacy of transcranial
magnetic stimulation targets for depression is related to intrinsic functional connectivity
with the subgenual cingulate. *Biol Psychiatry* Oct 01 2012;72(7):595-603.

36.　Wang D, Li M, Wang M, et al. Individual-specific functional connectivity markers track
dimensional and categorical features of psychotic illness. *Mol Psychiatry* Nov 15 2018.

529
530
531

532
533
534

535   Conflicts of interest:

537   All authors report no conflicts of interest

Supplemental Information

Multivariate Distance Matrix Regression:

As previously described [26-28], this analysis occurs in several stages: First, a seed-to-voxel connectivity map is generated by using an individual voxel's BOLD signal time-course to calculate the temporal Pearson's correlation coefficients between that voxel and all other gray matter voxels (Figure 1B). These maps are generated for every participant. In the second stage, the temporal correlation coefficients for each voxel in the connectivity map is correlated with the values of corresponding voxels in maps generated from every other participant. This Pearson's correlation coefficient, r is a measure of the similarity of whole-brain connectivity to that voxel between patients. This value is used to calculate between-subject distance (or dissimilarity) using the metric d_{ij} $= \sqrt{2(1 - r_{ij})}$ where i and j are two subjects and r is the correlation coefficient above. (Figure 1C)[29]. The next stage tests the relationship between a given variable (here SANS score) and the inter-subject distances in connectivity generated in the previous stage.

Broadly speaking, this process consists of ANOVA-like hypothesis testing where the tested relationship is between a variable of interest and a matrix of distances. This process was originally termed multivariate distance matrix regression by Zapala and Schork and used to test associations between gene expression and related variables [29]. Shezhad et al. then used their framework for testing the relationship between variables of interest and a matrix of distances where the matrix is between-subject similarity in whole-brain functional connectivity.

This test consists of first forming a distance matrix $A = (-\frac{1}{2} d_{ij}^2)_{1 \leq i,j \leq n}$ among n participants where d = the between subject distance metric calculated above. Next, this matrix is used to create a Gower's centered matrix $\mathbf{G} = \left(\mathbf{I} - \frac{1}{n}\mathbf{1}\mathbf{1}^T\right) A \left(\mathbf{I} - \frac{1}{n}\mathbf{1}\mathbf{1}^T\right)$, in which n is the number of participants, $\mathbf{I}$ is the $n \times n$ identity matrix, and $\mathbf{1}$ is a vector of n 1s. The F statistic for assessing the relationship between a predictor variable (e.g. SANS score) and dissimilarities in connectivity is calculated as follows: For m predictor variables, let $\mathbf{X}$ be a $n \times m$ design matrix of predictor values, and let $\mathbf{H} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T$ be the associated $n \times m$ "hat" matrix. $F = \frac{\text{tr}(\mathbf{H}\mathbf{G})/(m-1)}{\text{tr}[(\mathbf{I}-\mathbf{H})\mathbf{G}]/(n-m)}$ (Figure 1D) [25]. This process is repeated for every voxel.

The result is a whole brain map of how significantly SANS scores are related to functional connectivity at every voxel (Figure 1E). Covariates included in this analysis included scanner site, age, sex, mean FD (as a subject-level correction for motion effects[24]), and prescribed antipsychotic dosage (chlorpromazine equivalents, CPZE). To correct for multiple comparisons, a nonparametric permutation was calculated for voxels that exceeded the significance threshold of p < 0.005 and clusters of such of sizes at a

threshold of p < 0.05 [30], with a null distribution calculated from 5000 such permutations, as in prior MDMR studies.

Supplemental Figure 1

Supplemental Figure 1. Winner-Take-All Parcellations Obfuscate Topology-Phenotype Relationships. Several approaches to individual parcellation use a "winner-take-all" approach in assigning cortical territory to one canonical resting state network or another. We explored the consequence of this approach when applied to our dataset. In A) the connectivity strength between the right DLPFC region of interest and three

resting state networks (DMN, TPN and FPCN, Fronto-Parietal Control Network) were plotted for every individual in the dataset. Using a winner-take-all parcellation, this region would have been assigned to the FPCN in 75% of participants. In B) we demonstrate that within-network connectivity between the right DLPFC region and the FPCN is not correlated with symptom severity. We conclude that an analysis of connectivity-symptom relationships using a winner-take-all approach would then likely be misinterpreted as correlations between symptoms and *inter*-network connectivity i.e. between the FPCN and both the DMN and TPN when, in fact, symptom-connectivity relationships at this region can be entirely described by DMN and TPN topology irrespective of the FPCN.