Colicin E1 binds to TolC as an open hinge to penetrate the outer-membrane

Authors: S. Jimmy Budiardjo[1], Jacqueline J. Stevens[2], Anna L. Calkins[3], Ayotunde P. Ikujuni[2], Virangika K. Wimalasena[2], Emre Firlar[4], David A. Case[5], Julie S. Biteen[3], Jason T. Kaelber[4] and Joanna S.G. Slusky[1,2,*]

Affiliations:

[1] Center for Computational Biology, The University of Kansas, 2030 Becker Dr., Lawrence, KS 66047

[2] Department of Molecular Biosciences, The University of Kansas, 1200 Sunnyside Ave. Lawrence KS 66045

[3] Department of Chemistry, University of Michigan, Ann Arbor MI 48109-1055.

[4] Rutgers CryoEM & Nanoimaging Facility and Institute for Quantitative Biomedicine, Rutgers, The State University of New Jersey, Piscataway, NJ 08854, USA

[5] Department of Chemistry and Chemical Biology, Rutgers University, Piscataway, NJ 08854 USA

*Correspondence: slusky@ku.edu.

ABSTRACT

The double membrane architecture of Gram-negative bacteria forms a barrier that is effectively impermeable to extracellular threats. Bacteriocin proteins evolved to exploit the accessible, surface-exposed proteins embedded in the outer membrane to deliver cytotoxic cargo. Colicin E1 is a bacteriocin produced by, and lethal to, *Escherichia coli* that hijacks the outer membrane proteins TolC and BtuB to enter the cell. Here we capture the colicin E1 translocation domain inside its membrane receptor, TolC, by high-resolution cryoEM. Colicin E1 binds stably to TolC as an open hinge through the TolC pore—an architectural rearrangement from colicin E1's unbound conformation. This binding is stable in live *E. coli* cells as indicated by single-molecule fluorescence microscopy. Finally, colicin E1 fragments binding to TolC plugs the channel,

29   inhibiting its native efflux function as an antibiotic efflux pump and heightening susceptibility to

30   four antibiotic classes.

32   **INTRODUCTION**

33   In Gram-negative bacteria, the concentric structures of the outer membrane, cell wall, and

cytoplasmic membrane protect the cell from extracellular threats. Of these protective structures,

the outer membrane forms a particularly formidable barrier[4], owing to the impermeability of the

lipopolysaccharide (LPS) layer that constitutes the outer membrane[5]. The primary means by

which external molecules can gain access to the cell is through the ~100 varieties of barrel-

shaped proteins that are embedded in each bacterium outer membrane[6] and whose diverse

functions include the transport of molecules across the membrane—specifically, the import of

nutrients and metabolites and the export of toxins and waste.

41   Because outer membrane proteins (OMPs) are accessible from outside the cell,

bacteriophage and bacterial toxins have evolved to exploit OMPs to initiate delivering cargo

across the outer membrane. Bacteriocins hijack the OMPs of a target bacterium to cross its

impermeable outer membrane and kill the bacterium. Colicins are *E. coli*-specific bacteriocins,

protein toxin systems through which bacteria engage in bacterial warfare with other, similar

bacteria. Although colicins differ in their receptor targets and killing mechanisms, most colicins

share a common tri-domain architecture, comprising the following components: (i) an N-terminal

translocation (T) domain, (ii) a receptor-binding (R) domain, and (iii) a C-terminal cytotoxic (C)

domain (Figure 1A). Much of what is known of E colicin import has been determined through

studies of the colicin E3 and E9 as reviewed by Cramer et al.[7] Import is initialized by R domain

binding to the vitamin B12 transporter, BtuB, with high affinity[8,9]; this binding localizes the

colicin onto the outer membrane. Once the colicin is tethered to the outer membrane surface, the

T domain initiates translocation using the secondary OMP receptor OmpF to access TolA/Pal

system for group A colicins or the Ton system group B colicins. In most cases, the T domain

requires an OMP distinct from the R domain target.[10] A handful of colicins have been

structurally characterized with their OMP counterparts, although not in their entirety. Previous

structures of short N-terminal fragments of the T domain of ColE9 with OmpF[11] and the The R

domains of ColE2/E3 and Ia with BtuB[9,12] and Cir[8], respectively showed that T domains fully

penetrate deeply into lumen of the outer membrane receptors while R domains interact with the

extracellular loop regions of their receptors. Studies of the bacteriocin pyocin S5 from

Pseudomonas aeruginosa suggest that bacteriocin architectures and mechanisms may be

conserved across all Gram-negative species.

Colicin E1 is a bacteriocin secreted by *E. coli* that enters the periplasm of neighboring

cells and forms a pore on the cytoplasmic membrane leading to membrane depolarization and

cell death. Colicin E1 uses TolC, the outer membrane component of the acridine efflux pump, as

the T domain receptor[13] and BtuB as the R domain receptor[14]. A high-resolution x-ray structure

exists of only the cytotoxic domain of colicin E1[15] but not the T or R domains, thus no

mechanistic insights can be gained about initial steps of import. Domain swapping experiments

between colicin A and E1 identified regions of the protein that interact with specific outer

membrane proteins. Early studies of pore forming colicins, including E1, showed that cytotoxic-

domain-induced K^+ efflux on the inner membrane can be reversed after it has begun by

subsequent addition of trypsin to cells. This, in addition to trypsin's inability to cross the outer

membrane, led to the belief that portions of colicin remain tethered to their OMP partners as the

cytotoxic domain depolarizes the cytoplasmic membrane.[16-19]

In order to understand the early stages of colicin E1 import, functional studies of truncations of colicin E1 which lack the cytotoxic domain have been characterized *in vitro*. Residues 100-120 of colicin E1 (termed the 'TolC box', Figure 1A) have been shown to be necessary but not sufficient for binding TolC. Peptides that include the TolC box co-elute with TolC[20] and disrupt channel conductance[20,21]. Moreover, *E. coli* exposure to TolC-box-containing peptides can prevent subsequent binding and cytotoxicity of full-length colicin E1[22]. Circular dichroism of the T domain indicates that it exists as a helical hairpin in solution (similar to other colicin T domains)[23] and that the proline at the center of the TolC box forms its apex[20]. This measurement led to the proposal, known as the "pillar model,"[20] that the T domain inserts into the TolC barrel as a helical hairpin where the N and C-termini are pointing to the cell exterior. According to this model, the hairpin stuck in TolC acts as a buttress to facilitate the cytotoxic domain entry directly through the membrane.

A competing model for colicin E1 import, known as the "total thread" model[7,20], posits that the protein unfolds and passes through TolC N-terminus-first as an unstructured peptide, followed by refolding in the periplasm. In this model the binding between the intrinsically unstructured colicin N-termini and periplasmic proteins[22,24] creates a pulling force that results in the translocation of the whole colicin.

Here we use cryogenic electron microscopy (cryoEM) to solve the high-resolution structure of the colicin E1 T domain bound to TolC. We find that colicin E1 binds stably to TolC, not as a helical hairpin but as a single-pass folded helix with the N-terminus inside the periplasm. Additionally, we find that the colE1 TR domain binds TolC *in vivo* as well. Using single-molecule fluorescence microscopy, we find that colE1-TR, lacking the cytotoxic domain, remains stalled on the outer membrane and does not fully translocate into cells. Lastly, we

98   leverage this stalling of ColE1 to block the native TolC function as an antibiotic efflux pump.

99   Through real-time efflux assays, minimum inhibitory concentration (MIC) experiments, we find

100   that colE1-TR and colE1-T are able to inhibit TolC-mediated efflux. We find that this plugging

101   of TolC reduces the amount of antibiotics required to inhibit the bacterial growth—indicating

102   that this colicin E1 fragment reduces the antibiotic resistance conferred by TolC.

Figure 1. Cell-surface localization of colicin E1 fragments. (A) Architecture of full-length colicin E1 showing domains and their known binding partners. Three truncation constructs used in this study (B) CryoEM structure of TolC embedded in nanodiscs. Side, top, and bottom views are colored by local resolution, as computed in cryoSPARC from the final half-maps. The side view is cropped to display the particle interior. (C) SEC chromatogram of colE1-T (red line) and TolC (purple line). The arrow indicates the co-elution (black line) fractions that were analyzed by SDS-PAGE. On the SDS-PAGE gel (bottom), red arrows indicate the presence of the colicin E1 construct that has co-eluted with TolC. (D) The CryoEM structure of colE1-T bound to TolC and colored by local resolution as in (B).

RESULTS

CryoEM structure of TolC embedded in nanodiscs

To determine the structural details of colicin E1 binding to TolC, we solved the cryoEM

(cryogenic electron microscopy) structure of TolC embedded in nanodiscs with and without

added colE1-T (PDB 6WXH and 6WXI, respectively). We recently reported a high-yield TolC

preparation by refolding from inclusion bodies[25] and inserted refolded TolC into nanodiscs. The

cryoEM structure of refolded TolC alone (Figure 1B) is similar to the previously published

crystal structures of natively derived TolC alone[26] or in complex with hexamminecobalt[27] but

with more splayed loops at the periplasmic opening (residues 165-175). There was local

variation in

Table 1 Cryo-EM data collection, refinement, and validation statistics

	TolC (EMDB-21960) (PDB 6WXI)	TolC + colE1 (EMDB-21959) (PDB 6WXH)
Data collection and processing		
Electron microscope	Talos Arctica	Talos Arctica
Magnification	130,000×	205,000×
Voltage (kV)	200	200
Electron exposure (e–/Å^2)	33.04	35.96
Defocus range (μm)	0.5-2.6	0.4-2.1
Pixel size (Å)	1.038	0.6578
Symmetry imposed	C$_3$	C$_1$ (C$_3$)
Initial particle images (no.)	2,092,678	339,779
Final particle images (no.)	778,220	179,834
Map resolution (Å) FSC threshold	2.84 0.143	3.09 (2.81) 0.143
Refinement		
Initial model used (PDB code)	6WXH	1TQQ
Model resolution (Å) FSC threshold	3.11 0.5	3.37 0.5

R.m.s. deviations		
Bond lengths (Å)	0.0072	0.012
Bond angles (°)	1.11	1.76
Validation		
MolProbity score	1.33	0.50
Clashscore	4.42	0.00
Poor rotamers (%)	1.40	0.18
Ramachandran plot		
Favored (%)	98.12	98.31
Allowed (%)	1.88	1.32
Disallowed (%)	0	0.37

resolution within the model with lower resolution associated with the extracellular/periplasmic

ends and the nanodisc scaffold protein (Figure 1B). The lower residue resolution at these

apertures may indicate dynamics not captured in the x-ray crystal structures.

We next determined binding of colicin E1-T domain to capture the complex for structure

determination. Residues 1-190 (colE1-T) which span the translocation domain and are known to

bind to TolC (Figure 1A). We determined colE1-T:TolC binding *in vitro* via co-elution by size

exclusion chromatography (SEC) as previously described[28]. When colE1-T and TolC were

mixed, we observed a leftward shift in the TolC peak and a decrease in intensity associated with

the colE1-T peak indicating that a subset of the population has migrated with TolC (Figure 1C).

We analyzed the peak (Figure 1C arrow) using SDS-PAGE and found the presence of both

colE1-T and TolC. The unimodal shifted peak observed with colE1-T:TolC indicates that there is

a single species of fully bound TolC. When resolved by cryoEM, addition of colE1-T to TolC

breaks the three-fold channel symmetry and the additional protein is observed running all the

way through the TolC barrel as a single-pass, all-α-helical chain spanning more than 130 Å

(Figure 1D). The maps refined to nominal global resolutions of 2.84 Å and 3.09 Å for the TolC

and colE1-T:TolC, respectively (Table 1).

131    The colE1 T domain inserts into TolC with its amino terminus pointing inwards through

132  the periplasmic iris, and 2D class-averages show that the carboxy terminus of the helix continues

133  and projects out into the extracellular space (Figure 2A). 85 of the 190 colE1-T residues could be

134  modeled to this density (residues 46-131) (Figure 2B). No such regular protrusion was seen for

135  the glycine-rich colE1 amino-terminus, which we expect to be disordered in the periplasmic

136  space[20,24,29]. The colE1 chain binds TolC asymmetrically, primarily contacting only one of the

137  three TolC chains.

Figure 2. Colicin E1 binds to TolC as a single-pass kinked helix. (A) 2D class averages of colE1-T bound to TolC imbedded in nanodiscs (B) Cutaway view of the TolC interior (gray). ColE1-T (red) binds asymmetrically in an open-hinge conformation. The hinge region of colE1T including P110 shown in stick representation (inset). (C and D) The dilated terminal apertures. The apo cryoEM structure of TolC (subunits colored light blue, light green, and yellow) compared to holo cryoEM structure (subunits colored dark blue, dark green and orange). (C) Periplasmic aperture. (D) Extracellular aperture.

Compared to the unbound state, asymmetric colE1 binding dilates the periplasmic TolC

aperture (Figure 2C) and, to a lesser extent, the extracellular aperture (Figure 2D) so that they

can accommodate colE1 in the absence of any other proteins or motive force. Although the

terminal apertures dilate and are filled by the ligand and the TolC box forms specific interactions

with the TolC inner wall, the large internal volume of TolC is not fully occupied by colE1.

In solution, unbound colicin E1 is a two-helix hairpin as indicated by far UV circular

dichroism (CD)[28]. However, our cryoEM reconstruction of colicin E1 in complex with TolC

shows that colE1-T hinges open at the TolC box to an extended conformation upon binding to

TolC (Figure 2B). The colE1-T helices have a short kink in the middle around proline 110,

precisely at the proposed turn location in the hairpin model (Figure 2B inset). This area is also

the center of the TolC box that has been known to be critical for the colE1:TolC interaction[20,22].

Colicin E1 makes residue-specific interactions with TolC

TolC forms a large rigid conduit in the outer membrane and periplasm (Figure

3A) with a water filled lumen. The hydrophilic nature of the channel[30] coupled with the

inflexibility of the outer-membrane-embedded barrel largely precludes the formation of any

hydrophobic interfaces that are typically the basis of protein-protein interfaces involving helical

peptides. Yet, we did observe specific polar contacts between colicin E1's TolC box and the

TolC barrel.

To obtain the most accurate atomic model for interpretation of atomic interactions between

peptides, we utilized map-restrained molecular dynamics in model refinement[31]. The refined

model had improved chemical plausibility (for instance, MolProbity[32] score improved from 1.25

to 0.50) and polar contact were more easily identified. Because this refinement improved the

Figure 3. Inter-chain contacts between colE1T and TolC. (A) Molecular dynamics simulation refined structure of the colE1-T:TolC complex colored by chain (colE1-T: light red, TolC: light blue, light green, light gold) (B-C) Polar interaction network of colE1-T and TolC on the C-terminal side of the proline kink in the β-barrel region. (B) near the extracellular opening of TolC and(C) before the transition to the TolC periplasmic helical region. Residues involved in forming polar interactions and proline 110 are shown in sticks. (D) ColE1-T spanning the TolC helical barrel does not make full contact with the side of the TolC barrel. The cavity (black spheres) between colicin (light red) and one chain of TolC (light blue) was detected using GHECOM[1-3]

concordance between the map and model, we conclude that the use of cryoEM density as a

restraint was successful in preventing overfitting. Specifically, the EMRinger[33] score improved

from 2.77 to 3.81 and the map-model FSC=0.5 improved from 3.44Å to 3.37Å.

The atomic model showed significant polar interactions form between colicin E1 and

TolC at the apertures and around the TolC box. The acidic patch of colicin E1 near the

extracellular aperture contains two arginine residues (R129 and R131) that form hydrogen bond

networks with one TolC chain each (Figure 3B). In addition, colicin E1 residues T113, A116,

N119, N120 (all part of the TolC box) establish polar interactions with TolC residues G67, D69,

S90 and K305 on one TolC chain (Figure 3C, light red and light blue) while colicin H117

interacts with TolC F310 on a neighboring TolC chain (Figure 3C, light red and light gold). By

contrast with the β-barrel of TolC, in the interior of periplasmic helical barrel of TolC, colicin E1

does not make full contact with the α-helical barrel interior and there is a gap between the two

proteins (Figure 3D). This is in agreement with previous studies that showed a colicin truncation

(residues 1-100) that ends just before the beginning of the TolC box does not bind to TolC[20] or

prevent cytotoxicity of full-length colicin in cells[22].

Moreover, this structure is consistent with reports that mutations at TolC sites G43 or

S257 abrogate colE1 activity[34] as these residues are contact sites between TolC and colE1

(Figure S1). Conversely, the absence of any effect of mutating colicin R103 and R108[22], is

consistent with interaction those residues have with the barrel solvent rather than TolC. The

newly solved structure, in combination with previous work[20,22], indicates the specificity of

colicin E1 binding to TolC is encoded in the portion that binds to TolC β-barrel.

Membrane localization of colE1 truncations

Since we find the colE1 T domain binds to TolC, we investigated if colicin E1 fragments

bind to the native TolC in *E. coli* cells. Though ColE1 C domain activity—depolarizing the inner

membrane—requires the C domain to pass through the outer membrane[35], it remains unclear if

the colE1-TR domains translocate as well. We first determined *in vitro* binding of colE1-TR,

which includes residues 1-364 (colE1-TR) (Figure 1A) which contains the N-terminal portion

that binds to TolC and the C-terminal portion that binds to BtuB. ColE1-TR aggregates at the salt

concentration (40 mM) used for the co-elution experiment used for colE1-T so we increased the

NaCl concentration to 200 mM. ColE1-T still binds to TolC at 250 mM NaCl although to a much

lesser degree (Figure S2). Unlike colE1-T (Figure 1C), colE1-TR shows a bimodal distribution

indicative of a mixed population of bound and unbound TolC (Figure 4A). Using an extracellular

protease digestion assay, we assessed whether colE1-T and colE1-TR translocate into the cell or

localize on the cell surface. ColE1 fragments were incubated with cells, washed, and treated with

or without trypsin to digest any extracellularly bound protein to cells. When probing for the C-

terminal epitope by immunoblotting, there was no detectible colE1-T binding (Figure 4B, left),

though there was detectible binding of ColE1-TR (Figure 4B, right). Moreover, we found that if

colE1-TR is incubated with cells and subsequently treated with increasing trypsin concentrations,

the colE1-TR band disappeared, indicating that the colicin E1 fragment was localized to the

outer membrane surface (Figure 4B, right). The colE1-TR band of the intact cells may be slightly

less intense than the colE1 band for lysed cells due to some aggregation and precipitation of the

intact cell and proteins bound to it when boiling the whole cells for loading the SDS-PAGE. The

periplasmic control SurA was not degraded at any trypsin concentration unless the cells were

lysed before trypsin digestion[36].

Since the proteolysis experiments are not sensitive enough to detect if only a few

molecules have entered the cell, we further probed the interaction between the cell membrane

and surface-localized colE1-TR with single-molecule fluorescence microscopy using the

fluorescent dye Cy3. This method is able to detect single molecules on the cell surface as well as

those within the cell indicative of those molecules having crossed the outer membrane[37,38]. When

colE1-TR-Cy3 was added to the extracellular environment of WT BW25113 *E. coli* (containing

TolC), distinct puncta (Figure 4C, left) formed on 94% of the cells ($n = 111$) (Figure 4D). In a

$\Delta tolC$ strain, puncta were observed on only 18% of cells ($n = 99$) (Figure 4C, right; Figure 4D).

As a $\Delta btuB$ control strain, we used BL21 (DE3) which is known to have a premature stop codon

at *btuB* residue 58[39]. Puncta were observed on only 3% of cells lacking BtuB ($n = 105$) (Figure

4D and S3A. In WT and $\Delta tolC$ cells that featured puncta, colE1-TR formed puncta consistent

Figure 4. Colicin E1-TR localizes on the outside of the cell. (A) SEC chromatogram of colE1-TR (red line) and TolC (purple line). The arrow indicates the co-elution (black line) fractions that were analyzed by SDS-PAGE. On the SDS-PAGE gels (right), red arrow indicates the presence of colicin E1 construct that has co-eluted with TolC (purple arrow). (B) Extracellular protease digestion assay with two colicin E1 truncation constructs, each labeled with a C-terminal His-Tag. Periplasmic SurA was used as a membrane integrity control. (C) Fluorescence image of colE1-TR-Cy3 overlaid on outlines of living *E. coli* cells for WT and $\Delta tolC$. Red arrow points to a punctum. (D) Cell counts where colE1-TR-Cy3 punctum formation was observed for WT, $\Delta tolC$, and $\Delta btuB$.

with ~20 molecules. The observed size and number of molecules are in agreement with previous studies of BtuB puncta[40,41]. To rule out punctum formation as an artifact of Cy3 conjugation, we found colE1-TR-GFP displayed the same cluster formation as colE1-TR-Cy3 (Figure S3B). No other single protein binding events were detected aside from the puncta on either WT or Δ*tolC* cells.

Fluorescently labeled pyocins, the colicin analog in *Pseudomonas*, have previously been used to detect translocation across the outer membrane of *Pseudomonas aeruginosa*[42]. Here we use an analogous experiment with fluorescently labeled colE1 to determine cellular localization in *E. coli*. In time courses, bound colE1-TR puncta remained immobile for > 5 minutes (Movie S1). This is consistent with continued association of colE1-TR with membrane-embedded BtuB, which has limited mobility[43]. This result indicates that colE1-TR does not fully translocate[42], because if colE1-TR entered the periplasmic space it would freely diffuse on these timescales.

Colicin constructs lacking the R domain (colE1-T-Cy3) showed no detectable binding either to WT or Δ*tolC* cells (Figure S3C), indicating that the TolC-colE1-T interaction is much weaker than the BtuB-colE1-TR interaction.

Because we only see one punctum per cell, we anticipate that some BtuB and TolC remain unbound because of the geometric constraints of punctum formation. Though BtuB and TolC need to be in close proximity for colE1 binding to occur, when BtuB clusters together in groups of 12 or more, it may exclude TolC. These cluster geometries would therefore lower the number of full binding sites available for the T and R domains of colicin E1.

ColE1-TR inhibits efflux and makes *E. coli* more susceptible to antibiotics

239   Since the colicin truncations are stalled on their respective OMP receptors, we

240   investigated if this stalling could disrupt native TolC efflux. Real-time efflux inhibition by

Figure 5. Colicin E1 fragments inhibit efflux and potentiate antibiotics.
(A-C) Effect of colicin E1 fragments on efflux: WT with no protein (green), ΔtolC (red),
WT + 10 μM (orange), WT + 100 μM protein (blue). (A) colE1-T$_{100-143}$, (B) colE1-T, (C)
colE1-TR. (D) Antibiotic susceptibilities in the absence (green) and presence of colE1-TR
added to WT at 10 μM (orange) and 100 μM (blue). MICs for ΔtolC (red) are included as a
reference. Data are biological replicates reported as the mean and individual data points.

colicin E1 fragments was assessed using a live-cell assay with N-(2-naphthyl)-1-naphthylamine

(NNN)-dye, which is effluxed by the acridine efflux pump and fluoresces when it is localized

inside the cell[44]. NNN efflux can be turned off by the protonophore CCCP, which neutralizes the

proton motive force and allows NNN to accumulate within the cell. Active efflux can then be

monitored by the fluorescence decay once proton motive force is reenergized by glucose

addition[44-48].

We assessed the ability of colicin E1 truncations to plug TolC by monitoring real-time

NNN efflux. Cells exposed to colE1-T$_{100-143}$ did not show lower real-time efflux than untreated

cells (Figure 5A). This observation is notable in light of the fact that, in previous conductance

studies, similar peptides were shown to bind TolC[22,28] and to occlude the channel[21,28].

Cells exposed to colE1-T showed less decay in final fluorescence than untreated cells,

indicating that colE1-T partially inhibits efflux (Figure 5B). Finally, exposure to colE1-TR

produced no decrease in fluorescence, showing full inhibition of efflux (Figure 5C).

Because colE1-TR completely inhibits NNN efflux, we evaluated the capacity of colE1-

TR to potentiate antibiotics representing four different classes that are known TolC substrates:

kanamycin, ciprofloxacin, erythromycin, and benzalkonium chloride (from aminoglycosides,

fluoroquinolones, macrolides, and quaternary ammonium compounds, respectively). An effective

TolC plug will reduce the concentration required to inhibit growth as antibiotics remain trapped

within the cell. WT *E. coli* cells that were exposed to 100 μM colE1-TR with each of these

antibiotics showed significantly lower minimum inhibitory concentrations (MICs), the lowest

concentration of antibiotic that inhibits visible growth, than cells exposed to the antibiotics alone

(Fig. 5D): exposure to 100 μM colE1-TR made WT *E. coli* ~2 to 7-fold more susceptible to these

antibiotics (Table S1). Engagement of TolA by the N-terminal TolA box of colicin E9 has

previously been shown to cause outer membrane defects[49] which would lead to enhanced

antibiotic susceptibility. To rule out outer membrane-mediated defects caused by the TolA box in

colE1, we created a truncation of colE1-TR (colE1-TR$_{\Delta1\text{-}40}$) lacking the N-terminal 40 residues

including the TolA box. The MIC of colE1-TR and colE1-TR$_{\Delta1\text{-}40}$ are identical (Figure S4)

indicating that colE1-TR engagement of TolA does not contribute to the observed antibiotic

susceptibility.

Discussion

There are two conflicting models of colE1 translocation: 1) the total thread model in which the entire colicin unfolds and threads through TolC[22,50] and 2) the pillar model, in which colicin E1 inserts into TolC as a helical hairpin, facilitating LPS-mediated self-translocation of the colicin cytotoxic domain[28]. Our data support aspects of both models. The belief in a closed hinge conformation of bound colE1 prompted two arguments against the total thread mechanism: 1) the colE1-T closed hinge conformation is too wide to fit through TolC and 2) both ends would face the extracellular milieu[28,50]. The unanticipated colE1-T hinge opening in the bound state resolves these objections: colicin threads into the TolC barrel with the amino terminus pointing into the periplasm, as the total thread model hypothesized. However, our trypsin digests and single-molecule fluorescence images show colE1-TR stalls at the outer membrane. Though we cannot exclude the possibility that full length protein translocation depends on the C domain, we observe—as hypothesized by the pillar model—that colicin sits inside TolC but does not translocate. However, the architecture of the complex differs from that hypothesized by the pillar model. These results are in agreement with early studies of pore-forming colicins in which trypsin added to the extracellular environment can reverse colicin activity. Digestibility by trypsin, as well as the ability of colE1-T/TolC to reconstitute a monodisperse stable complex *in vitro* are consistent with the colicin remain in in contact with its outer membrane receptors while the cytotoxic domain depolarizes the inner membrane.

In addition to demonstrating a colicin insertion mechanism, our observations form the basis of a means to manipulate bacterial import and efflux. Though colicin E1 confers some level of antibiotic potentiation, the relatively high concentration of colE1-TR needed to inhibit efflux may be explained by a combination of the geometric constraints of creating large clusters of

BtuB and TolC, low colE1-T-TolC affinity[51], and residually unblocked pore, even in the bound state. Because of their limited potency, our colE1-T and colE1-TR fragments are not likely to be practical in direct applications of antibiotic potentiation. However, our proof-of-concept findings offer a potential roadmap for further development. A more potent TolC binder would not need the R domain anchor for affinity. Such a binder could be designed using the atomic details of the colE1-T-TolC structure as a basis. Moreover, this scaffold can be used for efflux pump inhibitors for at least five other bacterial organisms. Each of these organisms has a structurally characterized outer membrane efflux pump that is homologous and structurally similar to TolC[52].

More broadly, there are at least nine known outer membrane protein receptors for bacteriocins. These outer membrane proteins have a variety of functions including adhesion, iron transport, and general import[10,53]. Using this same strategy with fragments derived from other bacteriocins may additionally allow for controlled inhibition of other bacterial functions.

Acknowledgments We gratefully acknowledge Daniel Montezano, Pinakin Sukthankar, Rik Dhar, Dwight Deay, Scott Lovell, Matthias Wolf, Alexander Little, and Heather Shinogle for discussions and feedback, Mark Richter for the use of his fluorometer, Rajeev Misra for the pTrc vector containing the TolC gene, Vasileios Petrou for guidance on nanodiscs, Chamani Perera for peptide synthesis, Karen Marom for editorial guidance. We thank the Office of Advanced Research computing (OARC) at Rutgers for high-performance computing resources. Funding: NIH award R21-GM128022 to JSB, NIGMS awards DP2GM128201, P20GM113117, P20GM103638 and the Gordon and Betty Moore Inventor Fellowship to JSGS, and NIGMS awards P20 GM103418 and 2K12GM063651 to SJB.

Author contributions Conceptualization: SJB, JSGS; Methodology: SJB, JTK, JSGS, JSB; Writing: SJB, JSGS, JSB, JTK; Editing: SJB, JSGS, JSB, JTK; Investigation: SJB, JJS, ALC, API, VKW, EF, DAC, and JTK.

Competing interests The authors declare no competing interests.

References

1. Kawabata, T. Detection of multiscale pockets on protein surfaces using mathematical morphology. *Proteins* **78**, 1195-211 (2010).
2. Kawabata, T. Detection of cave pockets in large molecules: Spaces into which internal probes can enter, but external probes from outside cannot. *Biophys Physicobiol* **16**, 391-406 (2019).
3. Kawabata, T. & Go, N. Detection of pockets on protein surfaces using small and large probe spheres to find putative ligand binding sites. *Proteins* **68**, 516-29 (2007).
4. Alexander, C. & Rietschel, E.T. Invited review: Bacterial lipopolysaccharides and innate immunity. *Journal of Endotoxin Research* **7**, 167-202 (2001).
5. Kamio, Y. & Nikaido, H. Outer membrane of Salmonella typhimurium: accessibility of phospholipid head groups to phospholipase c and cyanogen bromide activated dextran in the external medium. *Biochemistry* **15**, 2561-70 (1976).
6. Freeman, J.T.C. & Wimley, W.C. TMBB-DB: a transmembrane β-barrel proteome database. *Bioinformatics* **28**, 2425-2430 (2012).
7. Cramer, W.A., Sharma, O. & Zakharov, S.D. On mechanisms of colicin import: the outer membrane quandary. *Biochem J* **475**, 3903-3915 (2018).
8. Buchanan, S.K. et al. Structure of colicin I receptor bound to the R-domain of colicin Ia: implications for protein import. *EMBO J* **26**, 2594-604 (2007).
9. Kurisu, G. et al. The structure of BtuB with bound colicin E3 R-domain implies a translocon. *Nat Struct Biol* **10**, 948-54 (2003).
10. Cascales, E. et al. Colicin Biology. *Microbiology and Molecular Biology Reviews* **71**, 158-229 (2007).
11. Housden, N.G. et al. Directed epitope delivery across the Escherichia coli outer membrane through the porin OmpF. *Proc Natl Acad Sci U S A* **107**, 21412-7 (2010).
12. Sharma, O. et al. Structure of the complex of the colicin E2 R-domain and its BtuB receptor. The outer membrane colicin translocon. *J Biol Chem* **282**, 23163-70 (2007).
13. Benedetti, H. et al. Individual domains of colicins confer specificity in colicin uptake, in pore-properties and in immunity requirement. *J Mol Biol* **217**, 429-39 (1991).
14. Benedetti, H. et al. Comparison of the Uptake Systems for the Entry of Various BtuB Group Colicins into Escherichia coli. *Microbiology* **135**, 3413-3420 (1989).
15. Elkins, P., Bunker, A., Cramer, W.A. & Stauffacher, C.V. A mechanism for toxin insertion into membranes is suggested by the crystal structure of the channel-forming domain of colicin E1. *Structure* **5**, 443-58 (1997).
16. Plate, C.A. & Luria, S.E. Stages in colicin K action, as revealed by the action of trypsin. *Proc Natl Acad Sci U S A* **69**, 2030-4 (1972).
17. Dankert, J., Hammond, S.M. & Cramer, W.A. Reversal by trypsin of the inhibition of active transport by colicin E1. *J Bacteriol* **143**, 594-602 (1980).
18. Dankert, J.R., Uratani, Y., Grabau, C., Cramer, W.A. & Hermodson, M. On a domain structure of colicin E1. A COOH-terminal peptide fragment active in membrane depolarization. *J Biol Chem* **257**, 3857-63 (1982).
19. Benedetti, H., Lloubes, R., Lazdunski, C. & Letellier, L. Colicin A unfolds during its translocation in Escherichia coli cells and spans the whole cell envelope when its pore has formed. *EMBO J* **11**, 441-7 (1992).
20. Zakharov, S.D., Wang, X.S. & Cramer, W.A. The Colicin E1 TolC-Binding Conformer: Pillar or Pore Function of TolC in Colicin Import? *Biochemistry* **55**, 5084-94 (2016).

21. Zakharov, S.D. et al. Colicin occlusion of OmpF and TolC channels: outer membrane
translocons for colicin import. *Biophys J* **87**, 3901-11 (2004).
22. Jakes, K.S. The Colicin E1 TolC Box: Identification of a Domain Required for Colicin
E1 Cytotoxicity and TolC Binding. *J Bacteriol* **199**(2017).
23. Wiener, M., Freymann, D., Ghosh, P. & Stroud, R.M. Crystal structure of colicin Ia.
Nature **385**, 461-4 (1997).
24. Housden, N.G. et al. Intrinsically disordered protein threads through the bacterial outer-
membrane porin OmpF. *Science* **340**, 1570-4 (2013).
25. Budiardjo, S.J. et al. High-Yield Preparation of Outer Membrane Protein Efflux Pumps
by in Vitro Refolding is Concentration Dependent. *J Membr Biol* **254**, 41-50 (2021).
26. Koronakis, V., Sharff, A., Koronakis, E., Luisi, B. & Hughes, C. Crystal structure of the
bacterial membrane protein TolC central to multidrug efflux and protein export. *Nature*
405, 914-9 (2000).
27. Higgins, M.K. et al. Structure of the ligand-blocked periplasmic entrance of the bacterial
multidrug efflux protein TolC. *J Mol Biol* **342**, 697-702 (2004).
28. Zakharov, S.D., Wang, X.S. & Cramer, W.A. The Colicin E1 TolC-Binding Conformer:
Pillar or Pore Function of TolC in Colicin Import? *Biochemistry* **55**, 5084-5094 (2016).
29. Housden, N.G. et al. Directional Porin Binding of Intrinsically Disordered Protein
Sequences Promotes Colicin Epitope Display in the Bacterial Periplasm. *Biochemistry*
57, 4374-4381 (2018).
30. Dhar, R., Feehan, R. & Slusky, J.S.G. Membrane Barrels Are Taller, Fatter, Inside-Out
Soluble Barrels. *J Phys Chem B* **125**, 3622-3628 (2021).
31. Wu, X., Subramaniam, S., Case, D.A., Wu, K.W. & Brooks, B.R. Targeted
conformational search with map-restrained self-guided Langevin dynamics: application
to flexible fitting into electron microscopic density maps. *J Struct Biol* **183**, 429-440
(2013).
32. Williams, C.J. et al. MolProbity: More and better reference data for improved all-atom
structure validation. *Protein Sci* **27**, 293-315 (2018).
33. Barad, B.A. et al. EMRinger: side chain-directed model and map validation for 3D cryo-
electron microscopy. *Nat Methods* **12**, 943-6 (2015).
34. Masi, M., Vuong, P., Humbard, M., Malone, K. & Misra, R. Initial steps of colicin E1
import across the outer membrane of Escherichia coli. *J Bacteriol* **189**, 2667-76 (2007).
35. Brunden, K.R., Uratani, Y. & Cramer, W.A. Dependence of the conformation of a colicin
E1 channel-forming peptide on acidic pH and solvent polarity. *J Biol Chem* **259**, 7682-7
(1984).
36. Besingi, R.N. & Clark, P.L. Extracellular protease digestion to evaluate membrane
protein cell surface localization. *Nat Protoc* **10**, 2074-80 (2015).
37. Tuson, H.H. & Biteen, J.S. Unveiling the inner workings of live bacteria using super-
resolution microscopy. *Anal Chem* **87**, 42-63 (2015).
38. Al-Husini, N. et al. BR-Bodies Provide Selectively Permeable Condensates that
Stimulate mRNA Decay and Prevent Release of Decay Intermediates. *Mol Cell* **78**, 670-
682 e8 (2020).
39. Studier, F.W., Daegelen, P., Lenski, R.E., Maslov, S. & Kim, J.F. Understanding the
differences between genome sequences of Escherichia coli B strains REL606 and
BL21(DE3) and comparison of the E. coli B and K-12 genomes. *J Mol Biol* **394**, 653-80
(2009).

40. Rassam, P. et al. Supramolecular assemblies underpin turnover of outer membrane proteins in bacteria. *Nature* **523**, 333-6 (2015).

41. Chavent, M. et al. How nanoscale protein interactions determine the mesoscale dynamic organisation of bacterial outer membrane proteins. *Nat Commun* **9**, 2846 (2018).

42. White, P. et al. Exploitation of an iron transporter for bacterial protein antibiotic import. *Proc Natl Acad Sci U S A* **114**, 12051-12056 (2017).

43. Kleanthous, C., Rassam, P. & Baumann, C.G. Protein-protein interactions and the spatiotemporal dynamics of bacterial outer membrane proteins. *Curr Opin Struct Biol* **35**, 109-15 (2015).

44. Bohnert, J.A., Schuster, S., Szymaniak-Vits, M. & Kern, W.V. Determination of real-time efflux phenotypes in Escherichia coli AcrB binding pocket phenylalanine mutants using a 1,2'-dinaphthylamine efflux assay. *PLoS One* **6**, e21196 (2011).

45. Bohnert, J.A., Karamian, B. & Nikaido, H. Optimized Nile Red efflux assay of AcrAB-TolC multidrug efflux system shows competition between substrates. *Antimicrob Agents Chemother* **54**, 3770-5 (2010).

46. Seeger, M.A. et al. Engineered disulfide bonds support the functional rotation mechanism of multidrug efflux pump AcrB. *Nat Struct Mol Biol* **15**, 199-205 (2008).

47. Iyer, R., Ferrari, A., Rijnbrand, R. & Erwin, A.L. A fluorescent microplate assay quantifies bacterial efflux and demonstrates two distinct compound binding sites in AcrB. *Antimicrob Agents Chemother* **59**, 2388-97 (2015).

48. Misra, R., Morrison, K.D., Cho, H.J. & Khuu, T. Importance of Real-Time Assays To Distinguish Multidrug Efflux Pump-Inhibiting and Outer Membrane-Destabilizing Activities in Escherichia coli. *J Bacteriol* **197**, 2479-88 (2015).

49. Rassam, P. et al. Intermembrane crosstalk drives inner-membrane protein organization in Escherichia coli. *Nat Commun* **9**, 1082 (2018).

50. Cramer, W.A., Sharma, O. & Zakharov, S.D. On mechanisms of colicin import: the outer membrane quandary. *Biochemical Journal* **475**, 3903-3915 (2018).

51. Taylor, R., Burgner, J.W., Clifton, J. & Cramer, W.A. Purification and characterization of monomeric Escherichia coli vitamin B12 receptor with high affinity for colicin E3. *J Biol Chem* **273**, 31113-8 (1998).

52. Franklin, M.W. et al. Efflux Pumps Represent Possible Evolutionary Convergence onto the beta-Barrel Fold. *Structure* **26**, 1266-1274.e2 (2018).

53. Kleanthous, C. Swimming against the tide: progress and challenges in our understanding of colicin translocation. *Nat Rev Microbiol* **8**, 843-8 (2010).

54. Hagn, F., Nasr, M.L. & Wagner, G. Assembly of phospholipid nanodiscs of controlled size for structural studies of membrane proteins by NMR. *Nat Protoc* **13**, 79-98 (2018).

55. Wiegand, I., Hilpert, K. & Hancock, R.E. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat Protoc* **3**, 163-75 (2008).

56. Daury, L. et al. Tripartite assembly of RND multidrug efflux pumps. *Nat Commun* **7**, 10731 (2016).

57. Weis, F. & Hagen, W.J.H. Combining high throughput and high quality for cryo-electron microscopy data collection. *Acta Crystallogr D Struct Biol* **76**, 724-728 (2020).

58. Mastronarde, D.N. Advanced Data Acquisition From Electron Microscopes With SerialEM. . *Microscopy and Microanalysis* **24**, 864-865 (2018).

59. Rice, W.J. et al. Routine determination of ice thickness for cryo-EM grids. *J Struct Biol* **204**, 38-44 (2018).

60. Budiardjo, S.J. et al. High yield preparation of outer-membrane protein efflux pumps by in vitro refolding is concentration dependent. *bioRxiv*, 2020.09.14.296756 (2020).

61. Punjani, A., Rubinstein, J.L., Fleet, D.J. & Brubaker, M.A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).

62. Zivanov, J. et al. New tools for automated high-resolution cryo-EM structure determination in RELION-3. *Elife* **7**(2018).

63. Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *J Struct Biol* **192**, 216-21 (2015).

64. Wagner, T. et al. SPHIRE-crYOLO is a fast and accurate fully automated particle picker for cryo-EM. *Commun Biol* **2**, 218 (2019).

65. Cardone, G., Heymann, J.B. & Steven, A.C. One number does not fit all: mapping local variations in resolution in cryo-EM reconstructions. *J Struct Biol* **184**, 226-36 (2013).

66. Afonine, P.V. et al. Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr D Struct Biol* **74**, 531-544 (2018).

67. Croll, T.I. ISOLDE: a physically realistic environment for model building into low-resolution electron-density maps. *Acta Crystallogr D Struct Biol* **74**, 519-530 (2018).

68. Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).

69. Maier, J.A. et al. ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J Chem Theory Comput* **11**, 3696-713 (2015).

70. Nguyen, H., Roe, D.R. & Simmerling, C. Improved Generalized Born Solvent Model Parameters for Protein Simulations. *J Chem Theory Comput* **9**, 2020-2034 (2013).

71. Goddard, T.D. et al. UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci* **27**, 14-25 (2018).

RESOURCE AVAILABILITY

STAR METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
THE HisTag mAb Mouse	GenScript	Cat#A00186-100; RRID:19J001957
anti-SurA Rabbit polyclonal	Cusabio	Cat#CSB-PA359693HA01ENV; RRID:E0109A
IRDye 800CW goat anti-mouse IgG secondary	LI-COR	Cat#925-32210; RRID:C81106-01
IRDye 800CW Donkey anti-Rabbit	LI-COR	Cat#925-32213; RRID:C80829-10
Bacterial and Virus Strains		
BW25113	Coli Genetic Stock Center	Cat#7636

JW5503-1	Coli Genetic Stock Center	Cat#11430
BL21(DE3)	Coli Genetic Stock Center	Cat#12504
Chemicals, Peptides, and Recombinant Proteins		
TolC	This paper	N/A
ColE1-T	This paper	N/A
ColE1-TR	This paper	N/A
ColE1-T100-143	This paper	N/A
Deposited Data		
TolC alone map	This paper	EMDB-21960 PDB 6WXI
TolC-colE1-T complex map	This paper	EMDB-21959 PDB 6WXH
Recombinant DNA		
pTrc99a-NoSS-TolC	This paper	Addgene
pET303-colE1-T	This paper	Addgene
pET303-colE1-TR	This paper	Addgene
MSP1D1	Addgene	Cat#20061
Software and Algorithms		
cryoSPARC 2 v2.12	Punjani, Rubinstein et al. 2017	https://cryosparc.com/
phenix v1.14-3260	Afonine, Poon et al. 2018	https://www.phenix-online.org/
serialEM v3.7	Mastronarde 2018	https://bio3d.colorado.edu/SerialEM/
coot v0.8.9.2-re 7766	Emsley, Lohkamp et al. 2010	https://www2.mrc-lmb.cam.ac.uk/personal/pemsley/coot/
ISOLDE v1.0b4.dev1	Croll 2018	https://isolde.cimr.cam.ac.uk/static/isolde/doc/tools/ISOLDE.html
UCSF ChimeraX	Goddard, Huang et al. 2018	https://www.cgl.ucsf.edu/chimerax/
PyMOL	Schrödinger, LLC	https://pymol.org/2/

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources should be directed to and will be fulfilled by the Lead Contact, Joanna Slusky (slusky@ku.edu).

Data and materials availability

All strains used are commercially available. All plasmids available through Addgene. CryoEM maps and models have been deposited with accession codes EMD-21960, EMD-21959, PDB ID 6WXI, and PDB ID 6WXH. All other data is available in the main text or the supplementary materials.

METHOD DETAILS

***E. coli* strains.** *E. coli* strains BW25113 and JW5503-1 were purchased from the Coli Genetic Stock Center (CGSC). JW5503-1 is a tolC732(del)::kan from the parent strain BW25113. BL21(DE3) were used for expression of the colicin constructs and TolC. BL21(DE3) has a premature stop codon at residue 58 of the *btuB* gene and therefore we used it as a Δ*btuB* strain for microscopy.

Expression and Purification. The gene for colE1-TR was synthesized as a gBlock (Integrated DNA Technologies) and cloned into pET303. Inverse PCR was used to delete the R domain and produce colicin E1-T. The gene for colicin E1-TR-GFP was produced by inserting GFP at the C terminus of colE1-TR.

Plasmids were transformed into *E. coli* BL21(DE3) cells and plated on LB + agar + 100 μg/mL carbenicillin. Single colonies were inoculated into 50 mL LB broth with 100 μg/mL carbenicillin and grown overnight at 37 °C with shaking at 250 r.p.m. Proteins were expressed by inoculating 1L of TB supplemented with 0.4% glycerol, 10 mM $MgCl_2$ and 100 μg/mL carbenicillin with 20 mL of the overnight culture. The culture was grown at 37 °C to an OD600 of 2.0 and induced with 1 mM IPTG. Expression cultures were then grown at 15 °C for 24 hours and harvested at 4,000 g for 30 minutes at 4 °C. Cell pellets were resuspended at 3 mL/g of cell pellet in lysis buffer (TBS, 5 mM $MgCl_2$, 10 mM imidazole, 1mM PMSF, 10 μg/mL DNase, 0.25 mg/mL lysozyme) and lysed via sonication (2 minutes, 2s on, 8s off, 40% amplitude, QSonica Q500 with 12.7 mm probe) in an ice bath. Lysates were centrifuged at 4,000 g for 10 minutes to remove un-lysed cells and debris. The supernatant was centrifuged again at 50,400 g in a Beckman Coulter J2-21 for 1 hour at 4 °C. Clarified lysates were applied to a 5 mL HisTrap FF column and purified using an ÄKTA FPLC system with a 20 column volume wash step with binding buffer (TBS, 25 mM imidazole) and eluted using a linear gradient from 0-50% elution buffer (TBS, 500 mM imidazole) in 10 column volumes. Concentrated proteins were loaded onto a HiLoad Superdex 16/60 200 pg gel filtration column and eluted into phosphate buffered saline (PBS) pH 7.4.

TolC expression and purification was conducted in the same manner for preparation for cryoEM and for SEC. The gene for full-length TolC (a generous gift from R. Misra) was cloned into pTrc99a with the signal sequence deleted for expression into inclusion bodies. Plasmids were transformed into BL21(DE3) and plated on LB + agar + 100 μg/mL carbenicillin. A single colony was picked and grown in LB at 37 °C with shaking at 250 r.p.m. overnight. In the morning, 1L of LB was inoculated with 20 mL of the overnight culture and grown at 37 °C with shaking at 250 r.p.m. until the culture reached an OD600 of 0.6, at which point protein expression was induced with 1mM IPTG for an additional 4 hours. Cells were then harvested at 4,000g for 30 minutes at 4 °C. Cell pellets were resuspended in mL of lysis buffer (TBS, 5 mM $MgCl_2$, 0.2 mg/mL lysozyme, 5 μg/mL DNase, 1mM PMSF) at 3 mL/1g of cell pellet and lysed via sonication (4 minutes, 2s on, 8s off, 40% amplitude, QSonica Q500 with 12.7 mm probe) in an ice bath. Cell lysates were centrifuged at 4,000 g for 30 minutes at 4 °C to harvest inclusion bodies. Inclusion body pellets were resuspended in inclusion body wash buffer (20 mM Tris pH 8.0, 0.5 mM EDTA, 1% Triton X-100) and centrifuged at 4,000g for 30 minutes at 4 °C. The inclusion body wash was repeated two more times. A final wash was done in 20 mM Tris pH 8.0 and inclusion bodies were stored at -20 °C. Inclusion bodies were solubilized in 20 mM Tris pH

8.0, 8M urea at 500 μM. N-octylpolyoxyethylene was added to 5 mL of solubilized TolC to a final concentration of 10% detergent and pipetted into a Slide-A-Lyzer G2 dialysis cassette with a 10,000 molecular weight cut off. Refolding was initiated by dialysis in 5L of 20 mM Tris pH 8.0, 100 mM NaCl at 4 °C with stirring overnight. TolC was centrifuged at 15,200 g for 10 minutes at 4 °C to remove aggregates. The supernatant was filtered through a 0.22 μm filter, concentrated to 2 mL, applied onto a HiLoad 16/60 Superdex 200 pg column on an ÄKTA Pure FPLC system, and eluted with 1.5 column volumes in 20 mM Tris pH 8.0, 100 mM NaCl, 0.05% n-dodecyl-β-D-maltoside. TolC containing fractions were pooled and concentrated to 300 μM in Amicon centrifugal filters with molecular weight cutoff of 30kDa.

Membrane scaffold protein MSP1D1. MSP1D1 in pET28a was purchased from Addgene and was expressed and purified as previously described[54].

Peptide Synthesis. ColE1-T100-143 was synthesized using standard Fmoc chemistry with a CEM liberty blue microwave peptide synthesizer. The peptides were cleaved using a solution of 92.5:2.5:2.5:2.5 TFA:TIPS:H2O:DoDt and the crude peptides where purified using HPLC. Analytical HPLC traces were acquired using an Agilent 1100 quaternary pump and a Hamilton PRP-1 (polystyrene-divinylbenzene) reverse phase analytical column (7 μm particle size, 4 mm x 25 cm) with UV detection at 215 nm. The eluents were heated to 45 °C to reduce separation of rotational isomers, and elution was achieved with gradients of water/ acetonitrile (90:10 to 0:100 containing 0.1% TFA) over 20 min. Low-resolution mass spectra (LRMS) were obtained using a Waters Micromass ZQ 4000 instrument with ESI+ ionization

Extracellular Protease Digestion. Protein localization after exogenous protein addition to whole cells was determined as previously described[36].

Single-Molecule Microscopy. Cysteine mutations were introduced at the C-terminus before the histidine tag for fluorophore conjugation. These constructs were purified as described in the expression and purification section with the addition of 1 mM TCEP in all buffers. All subsequent steps were performed with limited exposure to light and in amber tubes. Cyanine3 (Cy3) maleimide (Lumiprobe) was reconstituted in DMSO. Fluorophore labeling was achieved by mixing a 20-fold molar excess of Cy3 maleimide to protein and incubating overnight at 4 °C. Free dye was removed by gel filtration on a Sephadex NAP-10 G-25 column. Simultaneously to the dye removal, the sample was buffer exchanged into storage buffer (PBS pH 7.4, 1 mM DTT, 1 mM EDTA). The degree of labeling was determined spectrophotometrically from the concentrations of the dye and protein solutions using their respective extinction coefficients, ε, as described by their manufacturers or for the proteins as estimated by Expasy ProtParam (Cy3 ε_{548nm} = 162,000 L mol-1 cm-1; colE1-T-E192C ε_{280nm} = 9,970 L mol-1 cm-1; colE1-TR-E366C ε_{280nm} = 14,440 L mol-1 cm-1). Labeling efficiencies were ~75% and ~85% for colE1-T-E192C and colE1-TR-E366C, respectively. Protein concentrations were adjusted according to the percentage of labeled protein.

Cultures of *E. coli* (WT, ΔtolC, or BL21(DE3)) were grown in LB medium at 37 °C with shaking (180 r.p.m.) overnight, then transferred to MOPS minimal medium (Teknova) with 0.2% glycerol and 1.32 mM K2HPO4, and grown at 37 °C for 13 h. The sample was transferred to MOPS medium and grown to turbidity at 37 °C overnight. A 1-mL aliquot of culture was

centrifuged for 2 min at 4,850 g to pellet the cells. The pellet was washed in 1 mL MOPS and centrifuged a second time. The supernatant was then removed, and the cell pellet was resuspended in 500 µL MOPS. A 1.0 µL droplet of concentrated cells was placed onto a glass slide. Then, a 1.0 µL droplet of 1 µg/mL colicin E1 protein construct stock was added to the cells. The droplet was covered by an agarose pad (1% agarose in MOPS media) and a second coverslip.

Samples were imaged at room temperature using wide-field epifluorescence microscopy with sensitivity to detect single dye molecules as described previously [37]. Briefly, fluorescence was excited by a 561-nm laser (Coherent Sapphire 560-50) for Cy3 or a 488-nm laser (Coherent Sapphire 488-50) for GFP. The lasers were operated at low power densities (1 – 2 W/cm2), and fluorescence was imaged with an Olympus IX71 inverted microscope with a 100x, 1.40-NA oil-immersion objective and appropriate excitation, emission, and dichroic filters. A Photometrics Evolve electron multiplying charge-coupled device (EMCCD) camera with > 90% quantum efficiency captured the images at a rate of 20 frames per second. Each detector pixel corresponds to a 49 nm × 49 nm area of the sample.

Co-elution. The interaction of TolC and colicin E1-T or -TR were determined by co-elution on an SEC column. Purified TolC and colicin E1-T or -TR were mixed at a 1:2 molar ratio and incubated at room temperature for 1 hour before loading onto a Superdex 200 Increase 10/300 GL column (GE Healthcare). The protein was eluted with 1.5 column volumes into 20 mM Tris pH 8.0, 40 mM NaCl, 0.05% n-dodecyl-β-D-maltoside for colE1-T. For colE1-TR the NaCl concentration was increased to 200 mM to prevent precipitation. Elution fractions were collected every 0.5 mL. Peak fractions were concentrated to 20 µL and analyzed by 4-20% SDS-PAGE.

Real-Time Efflux. Real-time efflux activity in the presence of colE1-TR was determined as previously described with some modifications[44,45]. Cells were resuspended to OD600 1.5 in cold PBS with and without 10-100 µM colicin proteins and incubated for 15 minutes on ice. To turn off efflux, 100µM carbonyl cyanide m-chlorophenyl hydrazone (CCCP) was added. After an additional 15 minutes the efflux dye NNN was added to the cells to 10 µM. The cells were incubated at 25 °C with shaking at 140 r.p.m. for 2 hours. Cells were harvested at 3,500 g for 5 minutes and washed once in 20 mM potassium phosphate buffer pH 7 with 1mM MgCl2. Cell concentrations were adjusted to OD600 1.0 and placed on ice. Then, 2 mL of the cell suspension was loaded into a quartz cuvette (Agilent Technologies). Fluorescence was measured with an Agilent Cary Eclipse fluorescence spectrophotometer with slit widths at 5 and 10 nm for excitation wavelength of 370 nm and emission wavelength of 415 nm. Fluorescence measurements were taken every 1 second. After 100 seconds, 50 mM glucose was added to re-energize the cells and initiate efflux, and fluorescence data were collected for an additional 600 seconds.

Minimum Inhibitory Concentrations (MICs). MICs were determined using the broth dilution method[55] with some modifications in 96 well plate format using LB media in 100 µL well volumes. Cultures were grown at 37 °C with shaking at 250 r.p.m. and OD600 was read on a Biotek plate reader after 20 hours. MICs are defined by the lowest concentration that prevents visible growth. We chose an OD600 of >0.1 as the cutoff for growth. We report MICs as the mean of 3 or 6 biological replicates with each replicate plotted (Figure 3D, Table S1). Due to the

2-fold discretized nature of concentration ranges used to determine MICs we do not report statistical significance values as is typical of MIC reporting.

Reconstitution of TolC into Amphipol. Amphipol A8-35 (Anatrace) was solubilized in water at 33 mgs/mL. 1 mL of TolC at 0.5 mg/mL was mixed with 0.75 µL of A8-35 at 33 mg/mL for a 5-fold weight excess and incubated at room temperature for 30 minutes. Bio-Beads SM-2 resin that was washed in methanol and equilibrated with 20 mM Tris, 40 mM NaCl was added to the reaction mixture at 0.5 g/mL to initiate detergent exchange for A8-35 and incubated with rotation at 4 °C overnight. The mixture was transferred to a tube with fresh Bio-Beads and incubated at 4 °C for an additional 4 hours. The reaction mixture was loaded onto a HiLoad 16/60 Superdex 200 pg column on an ÄKTA Pure FPLC system and eluted with 1.5 column volumes in 20 mM Tris pH 8.0, 40 mM NaCl to remove free A8-35 and detergent. For colicin-bound TolC in A8-35, colicin E1-T was added to the reaction mixture at a >2 molar excess prior to gel filtration. TolC or colicin-bound TolC in A8-35 was concentrated to 2-4 mg/mL for cryoEM.

Reconstitution of TolC into lipid nanodiscs. POPC (Avanti Polar Lipids) in chloroform was dried under a stream of nitrogen and freeze dried to remove residual chloroform. Lipids were reconstituted to 50 mM in cholate buffer (20 mM Tris pH 8.0, 100 mM NaCl, 0.5 mM EDTA, 100 mM cholate) in an amber glass vial. The vial was submerged under a stream of warm water until the solution became clear. Lipids, membrane scaffold protein, and TolC were mixed in a 36:1:0.4 ratio as previously described[56]. Final concentrations were 4.5 mM POPC, 125 µM MSP1D1, 50 µM TolC in a 2 mL reaction with cholate brought up to 20 mM and dodecyl-maltoside up to 0.1%. The reaction mixture was incubated on ice for 1 hour. Bio-Beads SM-2 resin that was washed in methanol and equilibrated with 20 mM Tris, 40 mM NaCl were added to the reaction mixture at 0.5 g/mL to initiate nanodisc formation and incubated with rotation at 4°C overnight. The mixture was transferred to a tube with fresh Bio-Beads and incubated at 4 °C for an additional 4 hours. The reaction mixture was loaded onto a HiLoad 16/60 Superdex 200 pg column on an ÄKTA Pure FPLC system and eluted with 1.5 column volumes in 20 mM Tris pH 8.0, 40 mM NaCl to separate TolC inserted into nanodiscs from empty nanodiscs. For colicin-bound TolC in nanodiscs, colicin E1-T was added to the reaction mixture at a >2 molar excess prior to gel filtration. TolC or colicin-bound TolC in nanodiscs was concentrated to 2-4 mg/mL for cryoEM.

Cryoelectron microscopy. 3 µL of protein solution (TolC/colE1-T in amphipol or TolC/colE1-T in nanodiscs) was diluted to approximately 1.05 mg/mL concentration, applied to a glow-discharged TEM grid, and plunge-frozen in ethane using a Vitrobot Mark IV (FEI Company) with grade 595 filter paper (Ted Pella). Glow discharge was performed in ambient atmosphere at 0.39 mBar pressure. Imaging was performed using a Talos Arctica (FEI Company) operated at 200 kV with energy-filter and K2 Summit (Gatan, Inc.) for detection. To collect multiple images per hole while maintaining parallel illumination conditions, a nonstandard 20 µm condenser aperture was used to image TolC-colE1-T in nanodiscs. At nominal magnification of 205,000×, images were acquired in counting mode with a calibrated pixel size of 0.6578 Å. Fresnel fringes attributable to the beam interaction with the aperture were often seen in images. Some investigators have moved the microscope stage and altered the nominal objective lens true focus point to generate a fringe-free condition[57]. In this study, imaging at 205,000× with a 20 µm aperture yielded better results than imaging at 130,000× with a 50 µm aperture; at 130,000× with

a 20 μm aperture the fringes were extremely severe due to the larger field of view, so a full
dataset was not collected with those conditions. TolC in nanodiscs (without colE1-T) was
imaged at 130,000× with a 50μm condenser aperture (Table S2).

Micrographs were collected with SerialEM[58] using in-house modifications of the scripts of Chen
Xu (sphinx-emdocs.readthedocs.io). Briefly, multishot imaging was configured with 4 images
per hole for each of 16 holes; intermediate-magnification montages of grid squares were
acquired; points were selected manually for collection of 64 images per point; images were
acquired using coma-compensated image shift as gain- and dark-corrected LZW-compressed
TIFs. Side, top, and oblique views were seen in areas of thin ice. During screening, ice thickness
was estimated at 17-30nm by the method of I_0/I_{ZLP}[59].

The collection of micrographs of TolC without colicin at 130,000× magnification has been
previously described[60].

3D reconstruction and modeling. Final reconstructions were obtained using cryoSPARC 2[61].
1,018 micrographs were collected of amphipol-embedded TolC/colE1-T. Micrographs were
motion-corrected using RELION 3[62]. CTF parameters were determined by means of *ctffind*[63].
115,362 particles were selected with crYOLO[64]. 2D classification revealed that many particles
had aberrant morphology and only 24,624 (21%) were selected for 3D reconstruction. *Ab initio*
reconstruction in cryoSPARC 2[61] was effective at recovering a map whose shape was similar to
that of previously described TolC trimers. However, the data could only be refined to a nominal
global resolution of 6.0 Å, and lumenal density was insufficiently resolved. 4,492 micrographs
were then collected of nanodisc-embedded TolC/colE1-T and processed similarly. Of the
339,779 particles detected by cryoSPARC Live, 179,834 (53%) were in good classes. Although
there were slightly fewer particles per micrograph in the nanodisc dataset, more particles per
micrograph were usable. Beginning with the *ab initio* model and mask derived from the
amphipol data, this particle set was refined by cryoSPARC 2 non-uniform refinement with or
without imposed C_3 symmetry. The maps refined to nominal global resolutions of 2.81 Å and
3.09 Å for the symmetrized and asymmetric maps, respectively. There was local variation in
resolution within the map, with consistent, high resolution in the middle of TolC and lower
resolution at the lids and for colE1-T. Local resolution was computed in cryoSPARC by the
locally windowed FSC method[65] and rendered with UCSF Chimera. To reduce the voxel-based
values to averages for four regions of the complex, the local resolution map was masked to
include only voxels within 3 Å of a modeled atom and then the median value was calculated for
those voxels closest to colE1, closest to TolC residues 168-172 and 386-390, closest to TolC
residues 285-301 and 76-82, or closest to other TolC residues. Furthermore, it is notable that
while the nanodisc appears as a double-belt in the symmetrized map, in the asymmetric map the
nanodisc protein mostly appears on the side of TolC that is bound to colE1-T. One possible
explanation is that, despite masking, nanodisc asymmetry is a confounder of the asymmetric
refinement and is one source of heterogeneity in the data. Another possibility is that the C-
terminus of colE1-T forms an interaction with the nanodisc, causing preferential alignment of the
nanodisc with respect to the TolC/colE1-T complex.

196,158 particles of TolC without colicin were obtained as previously described[60]. Homogeneous refinement yielded a structure at 2.89 Å; local motion correction and global CTF refinement yielded a final map at 2.84 Å.

Modeling was initiated by rigidly docking a crystal structure of TolC in complex with hexamminecobalt (1TQQ)[27] into the symmetrized map density. Automated, semi-automated, and manual real-space refinement was performed using phenix[66], ISOLDE[67], and coot[68]. For TolC with colE1-T, additional refinement was performed in AMBER using the cryoEM density map as a restraint.

Although additional residues are present at the TolC C-terminus, these were not modeled because the density becomes unsharp after residue 428. Blurred density in the map suggests that the C-terminus follows helix 3 towards the periplasmic end of the molecule. ColE1-T was modeled *ab initio*. The 3-fold symmetrized map contains density at ~1/3rd occupancy for colE1-T and this density contains some high-resolution information not present in the asymmetric map, except near the TolC lid regions where symmetrization overlays colE1-T density with TolC density at a threefold-related position. First, polyalanine helices were placed in the helical density in coot. Cross-correlation coefficients for both helices are higher with the N-terminus oriented towards the periplasm, and the Christmas tree appearance was observed indicating that this is the correct chain orientation. An estimate of the registration was made by visual inspection of potential anchor residues. Finally, the hinge region was filled in using phenix and coot. This completed chain was refined against the asymmetric map in ISOLDE. Iteration between phenix, coot, and ISOLDE was continued until acceptable fit to density was achieved. In the case of TolC with colE1-T, the map was further improved by combining map information with molecular dynamics force fields[31]. Briefly, starting with the phenix/coot/ISOLDE-refined model, we performed restrained simulated annealing in AMBER, heating from 0K to 300K for 0.2 nsec, holding constant temperature for 1 nsec, and then cooling to 0K over 0.2 nsec. The cryoEM density map is utilized as a restraint potential in the annealing so that both map information and AMBER force field information are simultaneously utilized to obtain an optimum model consistent with the data[31]. The protein force field used the ff14SB force field[69] and a generalized Born implicit solvent model with *igb=8*[70], and a nonbonded cutoff of 20 Å. The relative weight of real-space map-based restraints and the force field was fixed using *fcons=0.02*. For colE1-T, information from the symmetrized map was integrated into the modeling procedure during manual remodeling in coot, but map-based refinement in phenix, ISOLDE, and AMBER were against the asymmetric map. TolC without colE1-T was modeled similarly but using the TolC-colE1-T structure as a starting point instead of 1TQQ, and without final AMBER refinement.

Molecular representations were generated with Chimera, ChimeraX[71] or PyMOL (Schrödinger, LLC).

Supplemental Information

767
768
769

770 **Colicin E1 binds to TolC as an open hinge to penetrate the outer-membrane**

772 S. Jimmy Budiardjo, Jacqueline J. Stevens, Anna L. Calkins, Ayotunde P. Ikujuni, Virangika K.
773 Wimalasena, Emre Firlar, David A. Case, Julie S. Biteen, Jason T. Kaelber and Joanna S.G.
774 Slusky

S257

G43

10Å

Figure S1. TolC box density cropped from the fully asymmetric cryoEM map of the TolC/colE1-T complex. Mutations at positions G43 and S257 in the TolC (grey sticks) barrel make direct contact with colE1-T (green). These mutations were previously shown to abolish WT colE1 function[34]. Mutations to bulkier side chains introduce a steric clash that prevents binding.

Figure S2. Co-elution of colE1-T with TolC at 200 mM NaCl (black). Under this higher salt
concentration, when TolC (purple) and colE1-T (red) are mixed, there is a smaller peak shift than
that seen with colE1-TR and the presence of a shoulder (black arrow). SDS-PAGE of the
shoulder shows presence of both TolC (purple arrow) and colE1-T (red arrow). Although some
binding was detected, this higher salt concentration prevents full binding as indicated by a much
fainter band for colE1-T than seen at the lower salt concentration (Figure 4A).

A colE1-TR-Cy3

B

C colE1-T-Cy3

Figure S3 Single-molecule microscopy. Fluorescence images overlaid on outlines of living *E. coli* cells from phase-contrast microscopy for WT and Δ*tolC* for colE1-TR-GFP and counts of cells where colE1-TR-GFP punctum formation was observed. (A) 97% of cells showed no binding of Cy3-labeled colE1-TR to Δ*btuB* (B) ColE1-TR-GFP forms similar puncta as Cy3-labeled ColE1-TR (main text Figure 4C). (C) No binding of Cy3-labeled colE1-T to WT or Δ*tolC* cells was detected with Cy3-labeled colE1-T. Scale bars: 2 μm.

Figure S4. Minimum Inhibitory Concentration of Erythromycin with colE1-TR TolA box deletion construct. The MIC for erythromycin with 100 µM colE1-TR or colE1-TR$_{\Delta 1\text{-}40}$ is identical indicating that TolA engagement by colE1-TR does not contribute to antibiotic susceptibility.

817 **Table S1.**

818 Mean minimum inhibitory concentrations (MICs) of antimicrobials in the presence of colE1-TR.

	Erythromycin (µg/mL)	Ciprofloxacin (ng/mL)	Kanamycin (µg/mL)	Benzalkonium Chloride (µg/mL)
WT	625.00	56.25	21.88	45.83
ΔtolC	7.03	15.63	NA	6.88
WT + 10 µM colE1-TR	364.58	75.00	28.14	25
WT + 100 µM colE1-TR	312.50	28.13	6.25	6.25

820
821

822   **Movie S1.**

823   A representative 8-second clip of colE1-TR bound to a live cell. ColE1-TR localizes on, and

824   remains bound to, the extracellular surface of *E. coli*. Fluorescence movie of Cy3-labeled colE1-

825   TR on living WT *E. coli* overlaid on outline of the *E. coli* cell from phase-contrast microscopy.

826   Continuous imaging at 25 frames per second. Scale bar: 2 μm.