Title

Colicin E1 opens its hinge to plug TolC

Authors

S. Jimmy Budiardjo[1], Jacqueline J. Stevens[2], Anna L. Calkins[3], Ayotunde P. Ikujuni[2], Virangika K. Wimalasena[2], Emre Firlar[4], David A. Case[5], Julie S. Biteen[3], Jason T. Kaelber[4] and Joanna S.G. Slusky[1,2,*]

Affiliations

[1] Center for Computational Biology, The University of Kansas, 2030 Becker Dr., Lawrence, KS 66047.

[2] Department of Molecular Biosciences, The University of Kansas, 1200 Sunnyside Ave. Lawrence KS 66045.

[3] Department of Chemistry, University of Michigan, Ann Arbor MI 48109-1055.

[4] Rutgers CryoEM & Nanoimaging Facility and Institute for Quantitative Biomedicine, Rutgers University, Piscataway, NJ 08854, USA.

[5] Department of Chemistry and Chemical Biology, Rutgers University, Piscataway, NJ 08854 USA.

*Correspondence: slusky@ku.edu.

Abstract

The double membrane architecture of Gram-negative bacteria forms a barrier that is effectively impermeable to extracellular threats. Bacteriocin proteins evolved to exploit the accessible, surface-exposed proteins embedded in the outer membrane to deliver cytotoxic cargo. Colicin E1 is a bacteriocin produced by, and lethal to, *Escherichia coli* that hijacks the outer membrane proteins TolC and BtuB to enter the cell. Here we capture the colicin E1 translocation domain inside its membrane receptor, TolC, by high-resolution cryoEM, the first reported structure of a bacteriocin bound to TolC. Colicin E1 binds stably to TolC as an open hinge through the TolC pore—an architectural rearrangement from colicin E1's unbound conformation.

This binding is stable in live *E. coli* cells as indicated by single-molecule fluorescence

microscopy. Finally, colicin E1 fragments binding to TolC plugs the channel, inhibiting its native

efflux function as an antibiotic efflux pump and heightening susceptibility to three antibiotic

classes. In addition to demonstrating that these protein fragments are useful starting points for

developing novel antibiotic potentiators, this method could be expanded to other colicins to

inhibit other outer membrane protein functions.

41
42   **Introduction**
43
In Gram-negative bacteria, the concentric structures of the outer membrane, cell wall, and

cytoplasmic membrane protect the cell from extracellular threats. Of these protective structures,

the outer membrane forms a particularly formidable barrier(*5*), owing to the impermeability of the

lipopolysaccharide (LPS) layer that constitutes the outer membrane(*7*). The primary means by

which external molecules can gain access to the cell is through the ~100 varieties of barrel-shaped

proteins that are embedded in each bacterium outer membrane(*8*) and whose diverse functions

include the transport of molecules across the membrane—specifically, the import of nutrients and

metabolites and the export of toxins and waste.

Because outer membrane proteins (OMPs) are accessible from outside the cell,

bacteriophage and bacterial toxins have evolved to exploit OMPs to initiate delivering cargo

across the outer membrane. Bacteriocins hijack the OMPs of a target bacterium to cross its

impermeable outer membrane and kill the bacterium. Colicins are *E. coli*-specific bacteriocins,

protein toxin systems through which bacteria engage in bacterial warfare with other, similar

bacteria. Although colicins differ in their receptor targets and killing mechanisms, most colicins

share a common tri-domain architecture, comprising the following components: (i) an N-terminal

translocation (T) domain, (ii) a receptor-binding (R) domain, and (iii) a C-terminal cytotoxic (C)

domain (Figure 1A). Much of what is known of E colicin import has been determined through

studies of the colicin E3 and E9 as reviewed by Cramer et al.(*9*) Import is initialized by R domain

binding to the vitamin B12 transporter, BtuB, with high affinity(*10, 11*); this binding localizes the

colicin onto the outer membrane. Once the colicin is tethered to the outer membrane surface, the

T domain initiates translocation using the secondary OMP receptor OmpF to access TolA/Pal

system for group A colicins or the Ton system group B colicins. In most cases, the T domain

requires an OMP distinct from the R domain target.(*12*) A handful of colicins have been

structurally characterized with their OMP counterparts, although not in their entirety. Previous

structures of short N-terminal fragments of the T domain of ColE9 with OmpF(*13*) and the The R

domains of ColE2/E3 and Ia with BtuB(*11, 14*) and Cir(*10*), respectively showed that T domains

fully penetrate deeply into lumen of the outer membrane receptors while R domains interact with

the extracellular loop regions of their receptors. Studies of the bacteriocin pyocin S5 from

Pseudomonas aeruginosa suggest that bacteriocin architectures and mechanisms may be

conserved across all Gram-negative species.

Colicin E1 is a bacteriocin secreted by *E. coli* that enters the periplasm of neighboring

cells and forms a pore on the cytoplasmic membrane leading to membrane depolarization and cell

death. Colicin E1 uses TolC, the outer membrane component of the acridine efflux pump, as the T

domain receptor(*15*) and BtuB as the R domain receptor(*16*). A high-resolution x-ray structure

exists of only the cytotoxic domain of colicin E1(*17*) but not the T or R domains, thus no

mechanistic insights can be gained about initial steps of import. Domain swapping experiments

(*15*) between colicin A and E1 identified regions of the protein that interact with specific outer

membrane proteins. Early studies of pore forming colicins, including E1, showed that cytotoxic-

domain-induced K^+ efflux on the inner membrane can be reversed after it has begun by

subsequent addition of trypsin to cells. This, in addition to trypsin's inability to cross the outer

membrane, led to the belief that portions of colicin remain tethered to their OMP partners as the

cytotoxic domain depolarizes the cytoplasmic membrane.(*18-21*)

In order to understand the early stages of colicin E1 import, functional studies of

truncations of colicin E1 which lack the cytotoxic domain have been characterized *in vitro*.

Residues 100-120 of colicin E1 (termed the 'TolC box', Figure 1A) have been shown to be

necessary but not sufficient for binding TolC. Peptides that include the TolC box co-elute with

TolC(*22*) and disrupt channel conductance(*22, 23*). Moreover, *E. coli* exposure to TolC-box-

containing peptides can prevent subsequent binding and cytotoxicity of full-length colicin E1(*24*).

Circular dichroism of the T domain indicates that it exists as a helical hairpin (closed hinge) in

solution similar to other colicin T domains(*25*) and that the proline at the center of the TolC box

forms its apex(*22*). This measurement led to the proposal, known as the "pillar model,"(*22*) that

the T domain inserts into the TolC barrel as a helical hairpin where the N and C-termini are

pointing to the cell exterior. According to this model, the hairpin stuck in TolC acts as a buttress

to facilitate the cytotoxic domain entry directly through the membrane.

A competing model for colicin E1 import, known as the "total thread" model(*9, 22*), posits

that the protein unfolds and passes through TolC N-terminus-first as an unstructured peptide,

followed by refolding in the periplasm. In this model the binding between the intrinsically

unstructured colicin N-termini and periplasmic proteins(*24, 26*) creates a pulling force that results

in the translocation of the whole colicin.

Here we use cryogenic electron microscopy (cryoEM) to solve the first high-resolution

structure of a bacteriocin, colicin E1, bound to TolC (*27*). We find that colicin E1 binds stably to

TolC, not as a helical hairpin but as a single-pass folded helix with the N-terminus inside the

periplasm. Additionally, we find that the colE1 TR domain binds TolC *in vivo* as well. Using

single-molecule fluorescence microscopy, we find that colE1-TR, lacking the cytotoxic domain,

remains stalled on the outer membrane and does not fully translocate into cells. Lastly, we

leverage this stalling of ColE1 to block the native TolC function as an antibiotic efflux pump.

Because they are accessible from outside the cell, OMPs are attractive targets for the development

of novel antibiotics, and research has begun to reveal the therapeutic potential of interfering with

OMP structure and function. Recently, a monoclonal antibody was found to inhibit OMP folding

with bactericidal effects(*28*).

Here by determining the structure and mechanism of colE1 insertion we establish an

alternative approach for targeting OMPs—the development of molecular plugs that block OMP

pores. Such plugs would allow for the manipulation of bacterial transport, providing a means of

either starving the bacterium by preventing the import of valuable nutrients or poisoning the

bacterium by preventing the export of toxins. Through real-time efflux assays, minimum

inhibitory concentration (MIC) experiments, we find that colE1-TR and colE1-T are able to

inhibit TolC-mediated efflux. We find that this plugging of TolC reduces the amount of

antibiotics required to inhibit the bacterial growth—indicating that this colicin E1 fragment

reduces the antibiotic resistance conferred by TolC.

Results

124
CryoEM structure of TolC embedded in nanodiscs

To determine the structural details of colicin E1 binding to TolC, we solved the cryoEM

(cryogenic electron microscopy) structure of TolC embedded in nanodiscs with and without

added colE1-T (PDB 6WXH and 6WXI, respectively). We recently reported a high-yield TolC

preparation by refolding from inclusion bodies(*29*) and inserted refolded TolC into nanodiscs.

The cryoEM structure of refolded TolC alone (Figure 1B) is similar to the previously published

crystal structures of natively derived TolC alone(*30*) or in complex with hexamminecobalt(*31*)

but with more splayed loops at the periplasmic opening (residues 165-175). There was local

variation in resolution within the model with lower resolution associated with the

extracellular/periplasmic ends and the nanodisc scaffold protein (Figure 1B). The lower residue

resolution at these apertures may indicate dynamics not captured in the x-ray crystal structures.

Figure 1. Cell-surface localization of colicin E1 fragments. (**A**) Architecture of full-length colicin E1 showing domains and their known binding partners. Three truncation constructs used in this study (**B**) CryoEM structure of TolC embedded in nanodiscs. Side, top, and bottom views are colored by local resolution, as computed in cryoSPARC from the final half-maps. The side view is cropped to display the particle interior. (**C**) SEC chromatogram of colE1-T (red line) and TolC (purple line). The arrow indicates the co-elution (black line) fractions that were analyzed by SDS-PAGE. On the SDS-PAGE gel (bottom), red arrows indicate the presence of the colicin E1 construct that has co-eluted with TolC. (**D**) The CryoEM structure of colE1-T bound to TolC and colored by local resolution as in (B).

We next determined binding of colicin E1-T domain to capture the complex for structure

determination. Residues 1-190 (colE1-T) which span the translocation domain and are known to

bind to TolC (Figure 1A). We determined colE1-T:TolC binding *in vitro* via co-elution by size

exclusion chromatography (SEC) as previously described(*6*). When colE1-T and TolC were

mixed, we observed a leftward shift in the TolC peak and a decrease in intensity associated with

the colE1-T peak indicating that a subset of the population has migrated with TolC (Figure 1C).

We analyzed the peak (Figure 1C arrow) using SDS-PAGE and found the presence of both colE1-

T and TolC. The unimodal shifted peak observed with colE1-T:TolC indicates that there is a

single species of fully bound TolC. When resolved by cryoEM, addition of colE1-T to TolC

breaks the three-fold channel symmetry and the additional protein is observed running all the way

through the TolC barrel as a single-pass, all-α-helical chain spanning more than 130 Å (Figure

1D). The maps refined to nominal global resolutions of 2.84 Å and 3.09 Å for the TolC and

colE1-T:TolC, respectively (Table S1).

The colE1 T domain inserts into TolC with its amino terminus pointing inwards through

the periplasmic iris, and 2D class-averages show that the carboxy terminus of the helix continues

and projects out into the extracellular space (Figure 2A). 85 of the 190 colE1-T residues could be

modeled to this density (residues 46-131) (Figure 2B). No such regular protrusion was seen for

the glycine-rich colE1 amino-terminus, which we expect to be disordered in the periplasmic

space(*22, 26, 32*). The colE1 chain binds TolC asymmetrically, primarily contacting only one of

the three TolC chains.

Compared to the unbound state, asymmetric colE1 binding dilates the periplasmic TolC

aperture (Figure 2C) and, to a lesser extent, the extracellular aperture (Figure 2D) so that they can

accommodate colE1 in the absence of any other proteins or motive force. Although the terminal

apertures dilate and are filled by the ligand and the TolC box forms specific interactions with the

TolC inner wall, the large internal volume of TolC is not fully occupied by colE1.

In solution, unbound colicin E1 is a two-helix hairpin as indicated by far UV circular

dichroism (CD)(*6*). A homology model built using I-TASSER(*4*) also predicts the closed hinge

conformation with proline 110 at the apex (Figure 2E top). However, our cryoEM reconstruction

of colicin E1 in complex with TolC shows that colE1-T hinges open at the TolC box to an

extended conformation upon binding to TolC (Figure 2B, and 2E bottom). The colE1-T helices

have a short kink in the middle around proline 110, precisely at the proposed turn location in the

hairpin model (Figure 2B inset). This area is also the center of the TolC box that has been known

to be critical for the colE1:TolC interaction(*22, 24*).

Figure 2. Colicin E1 binds to TolC as a single-pass kinked helix. (**A**) 2D class averages of colE1-T bound to TolC imbedded in nanodiscs (**B**) Cutaway view of the TolC interior (gray). ColE1-T (red) binds asymmetrically in an open-hinge conformation. The hinge region of colE1T including P110 shown in stick representation (inset). (**C and D**) The dilated terminal apertures. The apo cryoEM structure of TolC (subunits colored light blue, light green, and yellow) compared to holo cryoEM structure (subunits colored dark blue, dark green and orange). (**C**) Periplasmic aperture. (**D**) Extracellular aperture. (**E**) In the unbound state, colE1-T (red cylinders) exists as a closed hinge. When bound, colE1-T is in an open-hinge conformation through TolC with the N-terminus into the pore of TolC (grey). The homology model of colE1-T in its solution state was built using I-TASSER(*4*) and is consistent with previous experiments(*6*).

Colicin E1 makes residue-specific interactions with TolC

170 TolC forms a large rigid conduit in the outer membrane and periplasm (Figure 3A) with a

171 water filled lumen. The hydrophilic nature of the channel(*33*) coupled with the inflexibility of the

172 outer-membrane-embedded barrel largely precludes the formation of any hydrophobic interfaces

173   that are typically the basis of protein-protein interfaces involving helical peptides. Yet, we did

174   observe specific polar contacts between colicin E1's TolC box and the TolC barrel.

Figure 3. Inter-chain contacts between colE1T and TolC. (**A**) Molecular dynamics simulation refined structure of the colE1-T:TolC complex colored by chain (colE1-T: light red, TolC: light blue, light green, light gold) (**B-C**) Polar interaction network of colE1-T and TolC on the C-terminal side of the proline kink in the β-barrel region. (**B**) near the extracellular opening of TolC and (**C**) before the transition to the TolC periplasmic helical region. Residues involved in forming polar interactions and proline 110 are shown in sticks. (**D**) ColE1-T spanning the TolC helical barrel does not make full contact with the side of the TolC barrel. The cavity (black spheres) between colicin (light red) and one chain of TolC (light blue) was detected using GHECOM(*1-3*)

To obtain the most accurate atomic model for interpretation of atomic interactions

between peptides, we utilized map-restrained molecular dynamics in model refinement(*34*). The

refined model had improved chemical plausibility (for instance, MolProbity(*35*) score improved

from 1.25 to 0.50) and polar contact were more easily identified. Because this refinement

improved the concordance between the map and model, we conclude that the use of cryoEM

density as a restraint was successful in preventing overfitting. Specifically, the EMRinger(*36*)

score improved from 2.77 to 3.81 and the map-model FSC=0.5 improved from 3.44Å to 3.37Å.

The atomic model showed significant polar interactions form between colicin E1 and

TolC at the apertures and around the TolC box. The acidic patch of colicin E1 near the

extracellular aperture contains two arginine residues (R129 and R131) that form hydrogen bond

networks with one TolC chain each (Figure 3B). In addition, colicin E1 residues T113, A116,

N119, N120 (all part of the TolC box) establish polar interactions with TolC residues G67, D69,

S90 and K305 on one TolC chain (Figure 3C, light red and light blue) while colicin H117

interacts with TolC F310 on a neighboring TolC chain (Figure 3C, light red and light gold). By

contrast with the β-barrel of TolC, in the interior of periplasmic helical barrel of TolC, colicin E1

does not make full contact with the α-helical barrel interior and there is a gap between the two

proteins (Figure 3D). This is in agreement with previous studies that showed a colicin truncation

(residues 1-100) that ends just before the beginning of the TolC box does not bind to TolC(22) or

prevent cytotoxicity of full-length colicin in cells(24).

Moreover, this structure is consistent with reports that mutations at TolC sites G43 or

S257 abrogate colE1 activity(37) as these residues are contact sites between TolC and colE1

(Figure S1). Conversely, the absence of any effect of mutating colicin R103 and R108(24), is

consistent with interaction those residues have with the barrel solvent rather than TolC. The

newly solved structure, in combination with previous work(22, 24), indicates the specificity of

colicin E1 binding to TolC is encoded in the portion that binds to TolC β-barrel.

Membrane localization of colE1 truncations

Since we find the colE1 T domain binds to TolC, we investigated if colicin E1 fragments

bind to the native TolC in *E. coli* cells. Though ColE1 C domain activity—depolarizing the inner

membrane—requires the C domain to pass through the outer membrane(38), it remains unclear if

the colE1-TR domains translocate as well. We first determined *in vitro* binding of colE1-TR,

which includes residues 1-364 (colE1-TR) (Figure 1A) which contains the N-terminal portion that

binds to TolC and the C-terminal portion that binds to BtuB. ColE1-TR aggregates at the salt

concentration (40 mM) used for the co-elution experiment used for colE1-T so we increased the

NaCl concentration to 200 mM. ColE1-T still binds to TolC at 200 mM NaCl although to a much

lesser degree (Figure S2). Unlike colE1-T (Figure 1C), colE1-TR shows a bimodal distribution

indicative of a mixed population of bound and unbound TolC (Figure 4A). Using an extracellular

protease digestion assay, we assessed whether colE1-T and colE1-TR translocate into the cell or

localize on the cell surface. ColE1 fragments were incubated with cells, washed, and treated with

or without trypsin to digest any extracellularly bound protein to cells. When probing for the C-

terminal epitope by immunoblotting, there was no detectible colE1-T binding (Figure 4B, left),

though there was detectible binding of ColE1-TR (Figure 4B, right). Moreover, we found that if

colE1-TR is incubated with cells and subsequently treated with increasing trypsin concentrations,

the colE1-TR band disappeared, indicating that the colicin E1 fragment was localized to the outer

membrane surface (Figure 4B, right). The periplasmic control SurA was not degraded at any

trypsin concentration unless the cells were lysed before trypsin digestion(*39*).

Since the proteolysis experiments are not sensitive enough to detect if only a few

molecules have entered the cell, we further probed the interaction between the cell membrane and

surface-localized colE1-TR with single-molecule fluorescence microscopy using the fluorescent

dye Cy3. This method is able to detect single molecules on the cell surface as well as those within

the cell indicative of those molecules having crossed the outer membrane(*40, 41*). When colE1-

TR-Cy3 was added to the extracellular environment of WT BW25113 *E. coli* (containing TolC),

distinct puncta (Figure 4C, left) formed on 94% of the cells (number of cells, $n = 111$) (Figure

4D); cells with observed puncta most often featured a single punctum, though a small fraction had 2

puncta. On average, the WT cells had 1.2 puncta. In a Δ*tolC* strain, puncta were observed on only

18% of cells ($n = 99$) (Figure 4C, right; Figure 4D); on average, the Δ*tolC* cells had 0.2 puncta.

As a Δ*btuB* control strain, we used BL21 (DE3) which is known to have a premature stop codon

at *btuB* residue 58(*42*). Puncta were observed on only 3% of cells lacking BtuB ($n = 105$) (Figure

4D and S3A); on average, the Δ*btuB* cells had 0 puncta. In WT and Δ*tolC* cells that featured

puncta, colE1-TR formed puncta consistent with ~20 molecules based on dividing the puncta

intensity by the fluorescence intensity of the last fluorescent molecule before photobleaching (*43*).

The observed size and number of molecules are in agreement with previous studies of BtuB

clusters (*44, 45*) and the punctum locations within the cell were variable. To rule out punctum

formation as an artifact of Cy3 conjugation, we found colE1-TR-GFP displayed the same cluster

formation as colE1-TR-Cy3 (Figure S3B). No other single protein binding events were detected

aside from the puncta on either WT or Δ*tolC* cells.

Fluorescently labeled pyocins, the colicin analog in *Pseudomonas*, have previously been

Figure 4. Colicin E1-TR localizes on the outside of the cell. (**A**) SEC chromatogram of colE1-TR (red line) and TolC (purple line). The arrow indicates the co-elution (black line) fractions that were analyzed by SDS-PAGE. On the SDS-PAGE gels (right), red arrow indicates the presence of colicin E1 construct that has co-eluted with TolC (purple arrow). (**B**) Extracellular protease digestion assay with two colicin E1 truncation constructs, each labeled with a C-terminal His-Tag. Periplasmic SurA was used as a membrane integrity control. (**C**) Fluorescence image of colE1-TR-Cy3 overlaid on outlines of living *E. coli* cells from phase-contrast microscopy for WT and Δ*tolC*. Red arrow points to a punctum. (**D**) Cell counts where colE1-TR-Cy3 punctum formation was observed for WT, Δ*tolC*, and Δ*btuB*. Number of cells observed, *n* = 111, 91, 105, respectively.

used to detect translocation across the outer membrane of *Pseudomonas aeruginosa*(*46*). Here we

use an analogous experiment with fluorescently labeled colE1 to determine cellular localization in

E. coli. In time courses, bound colE1-TR puncta remained immobile for > 5 minutes (Movie S1

corresponds to the first 8 seconds of data used to attain the WT image in Figure 4C). This is

consistent with continued association of colE1-TR with membrane-embedded BtuB, which has

limited mobility(*47*). This result indicates that colE1-TR does not fully translocate(*46*), because if

colE1-TR entered the periplasmic space it would freely diffuse on these timescales.

Colicin constructs lacking the R domain (colE1-T-Cy3) showed no detectable binding

either to WT or Δ*tolC* cells (Figure S3C), indicating that the TolC-colE1-T interaction is much

weaker than the BtuB-colE1-TR interaction.

Because we only see one punctum per cell, we anticipate that some BtuB and TolC remain

unbound because of the geometric constraints of punctum formation. Though BtuB and TolC

need to be in close proximity for colE1 binding to occur, when BtuB clusters together in groups

of 12 or more, it may exclude TolC. These cluster geometries would therefore lower the number

of full binding sites available for the T and R domains of colicin E1.

ColE1-TR inhibits efflux and makes *E. coli* more susceptible to antibiotics

Since the colicin truncations are stalled on their respective OMP receptors, we

investigated if this stalling could disrupt native TolC efflux. Real-time efflux inhibition by colicin

E1 fragments was assessed using a live-cell assay with N-(2-naphthyl)-1-naphthylamine (NNN)-

dye, which is effluxed by the acridine efflux pump and fluoresces when it is localized inside the

cell(*48*). NNN efflux can be turned off by the protonophore CCCP, which neutralizes the proton

motive force and allows NNN to accumulate within the cell. Active efflux can then be monitored

by the fluorescence decay once proton motive force is reenergized by glucose addition(*48-52*).

We assessed the ability of colicin E1 truncations to plug TolC by monitoring real-time

NNN efflux. Cells exposed to colE1-T$_{100-143}$ did not show lower real-time efflux than untreated

cells (Figure 5A). This observation is notable in light of the fact that, in previous conductance

studies, similar peptides were shown to bind TolC(*6, 24*) and to occlude the channel(*6, 23*).

Cells exposed to colE1-T showed less decay in final fluorescence than untreated cells,

indicating that colE1-T partially inhibits efflux (Figure 5B). Finally, exposure to colE1-TR

produced no decrease in fluorescence, showing full inhibition of efflux (Figure 5C).

Figure 5. Colicin E1 fragments inhibit efflux and potentiate antibiotics.
(**A-C**) Effect of colicin E1 fragments on efflux: WT with no protein (green), $\Delta tolC$ (red),
WT + 10 μM (orange), WT + 100 μM protein (blue). (**A**) colE1-T$_{100-143}$, (**B**) colE1-T, (**C**)
colE1-TR. (**D**) Antibiotic susceptibilities in the absence (green) and presence of colE1-TR
added to WT at 10 μM (orange) and 100 μM (blue). MICs for $\Delta tolC$ (red) are included as a
reference. Data are biological replicates reported as the mean and individual data points.

Because colE1-TR completely inhibits NNN efflux, we evaluated the capacity of colE1-

TR to potentiate antibiotics representing three different classes that are known TolC substrates:

ciprofloxacin, erythromycin, and benzalkonium chloride (from fluoroquinolones, macrolides, and

quaternary ammonium compounds, respectively). An effective TolC plug will reduce the

concentration required to inhibit growth as antibiotics remain trapped within the cell. WT *E. coli*

cells that were exposed to 100 μM colE1-TR with each of these antibiotics showed significantly

lower minimum inhibitory concentrations (MICs), the lowest concentration of antibiotic that

inhibits visible growth, than cells exposed to the antibiotics alone (Figure 5D): exposure to 100

μM colE1-TR made WT *E. coli* ~2 to 7-fold more susceptible to these antibiotics (Table S2).

Exposure of WT cells to 10μM colE1-TR was sufficient to potentiate erythromycin. Engagement

of TolA by the N-terminal TolA box of colicin E9 has previously been shown to cause outer

membrane defects(*53*) which would lead to enhanced antibiotic susceptibility. To rule out outer

membrane-mediated defects caused by the TolA box in colE1, we created a truncation of colE1-

TR (colE1-TR$_{\Delta 1-40}$) lacking the N-terminal 40 residues including the TolA box. The MIC of

colE1-TR and colE1-TR$_{\Delta 1-40}$ are identical (Figure S4) indicating that colE1-TR engagement of

TolA does not contribute to the observed antibiotic susceptibility.

Discussion

There are two conflicting models of colE1 translocation: 1) the total thread model in

which the entire colicin unfolds and threads through TolC(*24, 54*) and 2) the pillar model, in

which colicin E1 inserts into TolC as a helical hairpin, facilitating LPS-mediated self-

translocation of the colicin cytotoxic domain(*6*). Our data support aspects of both models. The

belief in a closed hinge conformation of bound colE1 prompted two arguments against the total

thread mechanism: 1) the colE1-T closed hinge conformation is too wide to fit through TolC and

2) both ends would face the extracellular milieu(*6, 54*). The unanticipated colE1-T hinge opening

in the bound state resolves these objections: colicin threads into the TolC barrel with the amino

terminus pointing into the periplasm, as the total thread model hypothesized. However, our

trypsin digests and single-molecule fluorescence images show colE1-TR stalls at the outer

membrane. Though we cannot exclude the possibility that full length protein translocation

depends on the C domain, we observe—as hypothesized by the pillar model—that colicin sits

inside TolC but does not translocate. However, the architecture of the complex differs from that

hypothesized by the pillar model. These results are in agreement with early studies of pore-

forming colicins in which trypsin added to the extracellular environment can reverse colicin

activity. Digestibility by trypsin, as well as the ability of colE1-T/TolC to reconstitute a

monodisperse stable complex *in vitro* are consistent with the colicin remain in in contact with its

outer membrane receptors while the cytotoxic domain depolarizes the inner membrane.

In addition to demonstrating a colicin insertion mechanism, our observations form the basis of a means to manipulate bacterial import and efflux. Though colicin E1 confers some level of antibiotic potentiation, the relatively high concentration of colE1-TR needed to inhibit efflux may be explained by a combination of the geometric constraints of creating large clusters of BtuB and TolC, low colE1-T-TolC affinity(*55*), and residually unblocked pore, even in the bound state. Because of their limited potency, our colE1-T and colE1-TR fragments are not likely to be practical in direct applications of antibiotic potentiation. However, our proof-of-concept findings offer a potential roadmap for further development. A more potent TolC binder would not need the R domain anchor for affinity. Such a binder could be designed using the atomic details of the colE1-T-TolC structure as a basis. Moreover, this scaffold can be used for efflux pump inhibitors for at least five other bacterial organisms. Each of these organisms has a structurally characterized outer membrane efflux pump that is homologous and structurally similar to TolC(*56*).

More broadly, there are at least nine known outer membrane protein receptors for bacteriocins. These outer membrane proteins have a variety of functions including adhesion, iron transport, and general import(*12, 57*). Using this same strategy with fragments derived from other bacteriocins may additionally allow for controlled inhibition of other bacterial functions.

Materials and Methods

E. coli **strains.** *E. coli* strains BW25113 and JW5503-1 were purchased from the Coli Genetic Stock Center (CGSC). JW5503-1 is a tolC732(del)::kan from the parent strain BW25113. BL21(DE3) were used for expression of the colicin constructs and TolC. BL21(DE3) has a premature stop codon at residue 58 of the *btuB* gene and therefore we used it as a Δ*btuB* strain for microscopy.

Expression and Purification. The gene for colE1-TR was synthesized as a gBlock (Integrated DNA Technologies) and cloned into pET303. Inverse PCR was used to delete the R domain and

produce colicin E1-T. The gene for colicin E1-TR-GFP was produced by inserting GFP at the C

terminus of colE1-TR.

Plasmids were transformed into *E. coli* BL21(DE3) cells and plated on LB + agar + 100 µg/mL

carbenicillin. Single colonies were inoculated into 50 mL LB broth with 100 µg/mL carbenicillin

and grown overnight at 37 °C with shaking at 250 r.p.m. Proteins were expressed by inoculating

1L of TB supplemented with 0.4% glycerol, 10 mM $MgCl_2$ and 100 µg/mL carbenicillin with 20

mL of the overnight culture. The culture was grown at 37 °C to an OD600 of 2.0 and induced

with 1 mM IPTG. Expression cultures were then grown at 15 °C for 24 hours and harvested at

4,000 g for 30 minutes at 4 °C. Cell pellets were resuspended at 3 mL/g of cell pellet in lysis

buffer (TBS, 5 mM $MgCl_2$, 10 mM imidazole, 1mM PMSF, 10 µg/mL DNase, 0.25 mg/mL

lysozyme) and lysed via sonication (2 minutes, 2s on, 8s off, 40% amplitude, QSonica Q500 with

12.7 mm probe) in an ice bath. Lysates were centrifuged at 4,000 g for 10 minutes to remove un-

lysed cells and debris. The supernatant was centrifuged again at 50,400 g in a Beckman Coulter

J2-21 for 1 hour at 4 °C. Clarified lysates were applied to a 5 mL HisTrap FF column and purified

using an ÄKTA FPLC system with a 20 column volume wash step with binding buffer (TBS, 25

mM imidazole) and eluted using a linear gradient from 0-50% elution buffer (TBS, 500 mM

imidazole) in 10 column volumes. Concentrated proteins were loaded onto a HiLoad Superdex

16/60 200 pg gel filtration column and eluted into phosphate buffered saline (PBS) pH 7.4.

TolC expression and purification was conducted in the same manner for preparation for cryoEM

and for SEC. The gene for full-length TolC (a generous gift from R. Misra) was cloned into

pTrc99a with the signal sequence deleted for expression into inclusion bodies. Plasmids were

transformed into BL21(DE3) and plated on LB + agar + 100 µg/mL carbenicillin. A single colony

was picked and grown in LB at 37 °C with shaking at 250 r.p.m. overnight. In the morning, 1L of

LB was inoculated with 20 mL of the overnight culture and grown at 37 °C with shaking at 250

r.p.m. until the culture reached an OD600 of 0.6, at which point protein expression was induced

with 1mM IPTG for an additional 4 hours. Cells were then harvested at 4,000g for 30 minutes at 4

°C. Cell pellets were resuspended in mL of lysis buffer (TBS, 5 mM $MgCl_2$, 0.2 mg/mL

lysozyme, 5 µg/mL DNase, 1mM PMSF) at 3 mL/1g of cell pellet and lysed via sonication (4

minutes, 2s on, 8s off, 40% amplitude, QSonica Q500 with 12.7 mm probe) in an ice bath. Cell

lysates were centrifuged at 4,000 g for 30 minutes at 4 °C to harvest inclusion bodies. Inclusion

body pellets were resuspended in inclusion body wash buffer (20 mM Tris pH 8.0, 0.5 mM

EDTA, 1% Triton X-100) and centrifuged at 4,000g for 30 minutes at 4 °C. The inclusion body

wash was repeated two more times. A final wash was done in 20 mM Tris pH 8.0 and inclusion

bodies were stored at -20 °C. Inclusion bodies were solubilized in 20 mM Tris pH 8.0, 8M urea at

500 µM. N-octylpolyoxyethylene was added to 5 mL of solubilized TolC to a final concentration

of 10% detergent and pipetted into a Slide-A-Lyzer G2 dialysis cassette with a 10,000 molecular

weight cut off. Refolding was initiated by dialysis in 5L of 20 mM Tris pH 8.0, 100 mM NaCl at

4 °C with stirring overnight. TolC was centrifuged at 15,200 g for 10 minutes at 4 °C to remove

aggregates. The supernatant was filtered through a 0.22 µm filter, concentrated to 2 mL, applied

onto a HiLoad 16/60 Superdex 200 pg column on an ÄKTA Pure FPLC system, and eluted with

1.5 column volumes in 20 mM Tris pH 8.0, 100 mM NaCl, 0.05% n-dodecyl-β-D-maltoside.

TolC containing fractions were pooled and concentrated to 300 µM in Amicon centrifugal filters

with molecular weight cutoff of 30kDa.

Membrane scaffold protein MSP1D1. MSP1D1 in pET28a was purchased from Addgene and was

expressed and purified as previously described(*58*).

Peptide Synthesis. ColE1-T100-143 was synthesized using standard Fmoc chemistry with a

CEM liberty blue microwave peptide synthesizer. The peptides were cleaved using a solution of

92.5:2.5:2.5:2.5 TFA:TIPS:H2O:DoDt and the crude peptides where purified using HPLC.

Analytical HPLC traces were acquired using an Agilent 1100 quaternary pump and a Hamilton

PRP-1 (polystyrene-divinylbenzene) reverse phase analytical column (7 µm particle size, 4 mm x

25 cm) with UV detection at 215 nm. The eluents were heated to 45 °C to reduce separation of

rotational isomers, and elution was achieved with gradients of water/ acetonitrile (90:10 to 0:100

containing 0.1% TFA) over 20 min. Low-resolution mass spectra (LRMS) were obtained using a

Waters Micromass ZQ 4000 instrument with ESI+ ionization

Extracellular Protease Digestion. Protein localization after exogenous protein addition to whole

cells was determined as previously described (*39*) with one modification. Samples for intact cells

were lysed prior to loading on SDS-PAGE by adding 0.2 mg/mL lysozyme and incubating for 15

minutes followed by five freeze thaw cycles in liquid nitrogen.

Single-Molecule Microscopy. Cysteine mutations were introduced at the C-terminus before the

histidine tag for fluorophore conjugation. These constructs were purified as described in the

expression and purification section with the addition of 1 mM TCEP in all buffers. All subsequent

steps were performed with limited exposure to light and in amber tubes. Cyanine3 (Cy3)

maleimide (Lumiprobe) was reconstituted in DMSO. Fluorophore labeling was achieved by

mixing a 20-fold molar excess of Cy3 maleimide to protein and incubating overnight at 4 °C. Free

dye was removed by gel filtration on a Sephadex NAP-10 G-25 column. Simultaneously to the

dye removal, the sample was buffer exchanged into storage buffer (PBS pH 7.4, 1 mM DTT, 1

mM EDTA). The degree of labeling was determined spectrophotometrically from the

concentrations of the dye and protein solutions using their respective extinction coefficients, ε, as

described by their manufacturers or for the proteins as estimated by Expasy ProtParam (Cy3

ε_{548nm} = 162,000 L mol-1 cm-1; colE1-T-E192C ε_{280nm} = 9,970 L mol-1 cm-1; colE1-TR-

E366C ε_{280nm} = 14,440 L mol-1 cm-1). Labeling efficiencies were ~75% and ~85% for colE1-

T-E192C and colE1-TR-E366C, respectively. Protein concentrations were adjusted according to

the percentage of labeled protein.

Cultures of *E. coli* (WT, ΔtolC, or BL21(DE3)) were grown in LB medium at 37 °C with shaking

(180 r.p.m.) overnight, then transferred to MOPS minimal medium (Teknova) with 0.2% glycerol

and 1.32 mM K_2HPO_4, and grown at 37 °C for 13 h. The sample was transferred to MOPS

medium and grown to turbidity at 37 °C overnight. A 1-mL aliquot of culture was centrifuged for

2 min at 4,850 g to pellet the cells. The pellet was washed in 1 mL MOPS and centrifuged a

second time. The supernatant was then removed, and the cell pellet was resuspended in 500 μL

MOPS. A 1.0 μL droplet of concentrated cells was placed onto a glass slide. Then, a 1.0 μL

droplet of 1 μg/mL colicin E1 protein construct stock was added to the cells. The droplet was

covered by an agarose pad (1% agarose in MOPS media) and a second coverslip.

Samples were imaged at room temperature using wide-field epifluorescence microscopy with

sensitivity to detect single dye molecules as described previously (40). Briefly, fluorescence was

excited by a 561-nm laser (Coherent Sapphire 560-50) for Cy3 or a 488-nm laser (Coherent

Sapphire 488-50) for GFP. The lasers were operated at low power densities (1 – 2 W/cm2), and

fluorescence was imaged with an Olympus IX71 inverted microscope with a 100x, 1.40-NA

phase-contrast oil-immersion objective and appropriate excitation, emission, and dichroic filters.

A Photometrics Evolve electron multiplying charge-coupled device (EMCCD) camera with >

90% quantum efficiency captured the images at a rate of 20 frames per second. Each detector

pixel corresponds to a 49 nm × 49 nm area of the sample. Phase-contrast images of cells were

segmented to attain *E. coli* cell outlines using a custom MATLAB script (The MathWorks,

Natick, MA).

Co-elution. The interaction of TolC and colicin E1-T or -TR were determined by co-elution on an

SEC column. Purified TolC and colicin E1-T or -TR were mixed at a 1:2 molar ratio to ensure an

excess of colicin to saturate TolC binding sites and incubated at room temperature for 1 hour

before loading onto a Superdex 200 Increase 10/300 GL column (GE Healthcare). The protein

was eluted with 1.5 column volumes into 20 mM Tris pH 8.0, 40 mM NaCl, 0.05% n-dodecyl-β-

D-maltoside for colE1-T. For colE1-TR the NaCl concentration was increased to 200 mM to

prevent precipitation. Elution fractions were collected every 0.5 mL. Peak fractions were

concentrated to 20 µL and analyzed by 4-20% SDS-PAGE.

Real-Time Efflux. Real-time efflux activity in the presence of colE1-TR was determined as

previously described with some modifications(*48, 49*). Cells were resuspended to OD600 1.5 in

cold PBS with and without 10-100 µM colicin proteins and incubated for 15 minutes on ice. To

turn off efflux, 100µM carbonyl cyanide m-chlorophenyl hydrazone (CCCP) was added. After an

additional 15 minutes the efflux dye NNN was added to the cells to 10 µM. The cells were

incubated at 25 °C with shaking at 140 r.p.m. for 2 hours. Cells were harvested at 3,500 g for 5

minutes and washed once in 20 mM potassium phosphate buffer pH 7 with 1mM $MgCl_2$. Cell

concentrations were adjusted to OD600 1.0 and placed on ice. Then, 2 mL of the cell suspension

was loaded into a quartz cuvette (Agilent Technologies). Fluorescence was measured with an

Agilent Cary Eclipse fluorescence spectrophotometer with slit widths at 5 and 10 nm for

excitation wavelength of 370 nm and emission wavelength of 415 nm. Fluorescence

measurements were taken every 1 second. After 100 seconds, 50 mM glucose was added to re-

energize the cells and initiate efflux, and fluorescence data were collected for an additional 600

seconds. Figure 5A-C, reflects time after glucose addition.

Minimum Inhibitory Concentrations (MICs). MICs were determined using the broth dilution

method(*59*) with some modifications in 96 well plate format using LB media in 100 µL well

volumes. Cultures were grown at 37 °C with shaking at 250 r.p.m. and OD600 was read on a

Biotek plate reader after 20 hours. MICs are defined by the lowest concentration that prevents

visible growth. We chose an OD600 of >0.1 as the cutoff for growth. We report MICs as the

mean of 3 or 6 biological replicates with each replicate plotted (Figure 3D, Table S2). Due to the

2-fold discretized nature of concentration ranges used to determine MICs we do not report

statistical significance values as is typical of MIC reporting.

Reconstitution of TolC into Amphipol. Amphipol A8-35 (Anatrace) was solubilized in water at 33 mgs/mL. 1 mL of TolC at 0.5 mg/mL was mixed with 0.75 µL of A8-35 at 33 mg/mL for a 5-fold weight excess and incubated at room temperature for 30 minutes. Bio-Beads SM-2 resin that was washed in methanol and equilibrated with 20 mM Tris, 40 mM NaCl was added to the reaction mixture at 0.5 g/mL to initiate detergent exchange for A8-35 and incubated with rotation at 4 °C overnight. The mixture was transferred to a tube with fresh Bio-Beads and incubated at 4 °C for an additional 4 hours. The reaction mixture was loaded onto a HiLoad 16/60 Superdex 200 pg column on an ÄKTA Pure FPLC system and eluted with 1.5 column volumes in 20 mM Tris pH 8.0, 40 mM NaCl to remove free A8-35 and detergent. For colicin-bound TolC in A8-35, colicin E1-T was added to the reaction mixture at a >2 molar excess prior to gel filtration. TolC or colicin-bound TolC in A8-35 was concentrated to 2-4 mg/mL for cryoEM.

Reconstitution of TolC into lipid nanodiscs. POPC (Avanti Polar Lipids) in chloroform was dried under a stream of nitrogen and freeze dried to remove residual chloroform. Lipids were reconstituted to 50 mM in cholate buffer (20 mM Tris pH 8.0, 100 mM NaCl, 0.5 mM EDTA, 100 mM cholate) in an amber glass vial. The vial was submerged under a stream of warm water until the solution became clear. Lipids, membrane scaffold protein, and TolC were mixed in a 36:1:0.4 ratio as previously described(*60*). Final concentrations were 4.5 mM POPC, 125 µM MSP1D1, 50 µM TolC in a 2 mL reaction with cholate brought up to 20 mM and dodecyl-maltoside up to 0.1%. The reaction mixture was incubated on ice for 1 hour. Bio-Beads SM-2 resin that was washed in methanol and equilibrated with 20 mM Tris, 40 mM NaCl were added to the reaction mixture at 0.5 g/mL to initiate nanodisc formation and incubated with rotation at 4°C overnight. The mixture was transferred to a tube with fresh Bio-Beads and incubated at 4 °C for an additional 4 hours. The reaction mixture was loaded onto a HiLoad 16/60 Superdex 200 pg column on an ÄKTA Pure FPLC system and eluted with 1.5 column volumes in 20 mM Tris pH 8.0, 40 mM NaCl to separate TolC inserted into nanodiscs from empty nanodiscs. For colicin-

bound TolC in nanodiscs, colicin E1-T was added to the reaction mixture at a >2 molar excess

prior to gel filtration. TolC or colicin-bound TolC in nanodiscs was concentrated to 2-4 mg/mL

for cryoEM.

Cryoelectron microscopy. 3 μL of protein solution (TolC/colE1-T in amphipol or TolC/colE1-T

in nanodiscs) was diluted to approximately 1.05 mg/mL concentration, applied to a glow-

discharged TEM grid, and plunge-frozen in ethane using a Vitrobot Mark IV (FEI Company) with

grade 595 filter paper (Ted Pella). Glow discharge was performed in ambient atmosphere at 0.39

mBar pressure. Imaging was performed using a Talos Arctica (FEI Company) operated at 200 kV

with energy-filter and K2 Summit (Gatan, Inc.) for detection. To collect multiple images per hole

while maintaining parallel illumination conditions, a nonstandard 20 μm condenser aperture was

used to image TolC-colE1-T in nanodiscs. At nominal magnification of 205,000×, images were

acquired in counting mode with a calibrated pixel size of 0.6578 Å. Fresnel fringes attributable to

the beam interaction with the aperture were often seen in images. Some investigators have moved

the microscope stage and altered the nominal objective lens true focus point to generate a fringe-

free condition(61). In this study, imaging at 205,000× with a 20 μm aperture yielded better results

than imaging at 130,000× with a 50 μm aperture; at 130,000× with a 20 μm aperture the fringes

were extremely severe due to the larger field of view, so a full dataset was not collected with

those conditions. TolC in nanodiscs (without colE1-T) was imaged at 130,000× with a 50μm

condenser aperture (Table S1).

Micrographs were collected with SerialEM(62) using in-house modifications of the scripts of

Chen Xu (sphinx-emdocs.readthedocs.io). Briefly, multishot imaging was configured with 4

images per hole for each of 16 holes; intermediate-magnification montages of grid squares were

acquired; points were selected manually for collection of 64 images per point; images were

acquired using coma-compensated image shift as gain- and dark-corrected LZW-compressed

TIFs. Side, top, and oblique views were seen in areas of thin ice. During screening, ice thickness

was estimated at 17-30nm by the method of I_0/I_{ZLP}(*63*).

The collection of micrographs of TolC without colicin at 130,000× magnification has been

previously described(*64*).

3D reconstruction and modeling. Final reconstructions were obtained using cryoSPARC 2(*65*).

1,018 micrographs were collected of amphipol-embedded TolC/colE1-T. Micrographs were

motion-corrected using RELION 3(*66*). CTF parameters were determined by means of

ctffind(*67*). 115,362 particles were selected with crYOLO(*68*). 2D classification revealed that

many particles had aberrant morphology and only 24,624 (21%) were selected for 3D

reconstruction. *Ab initio* reconstruction in cryoSPARC 2(*65*) was effective at recovering a map

whose shape was similar to that of previously described TolC trimers. However, the data could

only be refined to a nominal global resolution of 6.0 Å, and lumenal density was insufficiently

resolved. 4,492 micrographs were then collected of nanodisc-embedded TolC/colE1-T and

processed similarly. Of the 339,779 particles detected by cryoSPARC Live, 179,834 (53%) were

in good classes. Although there were slightly fewer particles per micrograph in the nanodisc

dataset, more particles per micrograph were usable. Beginning with the *ab initio* model and mask

derived from the amphipol data, this particle set was refined by cryoSPARC 2 non-uniform

refinement with or without imposed C_3 symmetry. The maps refined to nominal global resolutions

of 2.81 Å and 3.09 Å for the symmetrized and asymmetric maps, respectively. There was local

variation in resolution within the map, with consistent, high resolution in the middle of TolC and

lower resolution at the lids and for colE1-T. Local resolution was computed in cryoSPARC by the

locally windowed FSC method(*69*) and rendered with UCSF Chimera. To reduce the voxel-based

values to averages for four regions of the complex, the local resolution map was masked to

include only voxels within 3 Å of a modeled atom and then the median value was calculated for

those voxels closest to colE1, closest to TolC residues 168-172 and 386-390, closest to TolC

residues 285-301 and 76-82, or closest to other TolC residues. Furthermore, it is notable that

while the nanodisc appears as a double-belt in the symmetrized map, in the asymmetric map the

nanodisc protein mostly appears on the side of TolC that is bound to colE1-T. One possible

explanation is that, despite masking, nanodisc asymmetry is a confounder of the asymmetric

refinement and is one source of heterogeneity in the data. Another possibility is that the C-

terminus of colE1-T forms an interaction with the nanodisc, causing preferential alignment of the

nanodisc with respect to the TolC/colE1-T complex.

196,158 particles of TolC without colicin were obtained as previously described(*64*).

Homogeneous refinement yielded a structure at 2.89 Å; local motion correction and global CTF

refinement yielded a final map at 2.84 Å.

Modeling was initiated by rigidly docking a crystal structure of TolC in complex with

hexamminecobalt (1TQQ)(*31*) into the symmetrized map density. Automated, semi-automated,

and manual real-space refinement was performed using phenix(*70*), ISOLDE(*71*), and coot(*72*).

For TolC with colE1-T, additional refinement was performed in AMBER using the cryoEM

density map as a restraint.

Although additional residues are present at the TolC C-terminus, these were not modeled because

the density becomes unsharp after residue 428. Blurred density in the map suggests that the C-

terminus follows helix 3 towards the periplasmic end of the molecule. ColE1-T was modeled *ab*

initio. The 3-fold symmetrized map contains density at ~$1/3^{rd}$ occupancy for colE1-T and this

density contains some high-resolution information not present in the asymmetric map, except near

the TolC lid regions where symmetrization overlays colE1-T density with TolC density at a

threefold-related position. First, polyalanine helices were placed in the helical density in coot.

Cross-correlation coefficients for both helices are higher with the N-terminus oriented towards the

periplasm, and the Christmas tree appearance was observed indicating that this is the correct chain

orientation. An estimate of the registration was made by visual inspection of potential anchor

residues. Finally, the hinge region was filled in using phenix and coot. This completed chain was

refined against the asymmetric map in ISOLDE. Iteration between phenix, coot, and ISOLDE was

continued until acceptable fit to density was achieved. In the case of TolC with colE1-T, the map

was further improved by combining map information with molecular dynamics force fields(*34*).

Briefly, starting with the phenix/coot/ISOLDE-refined model, we performed restrained simulated

annealing in AMBER, heating from 0K to 300K for 0.2 nsec, holding constant temperature for 1

nsec, and then cooling to 0K over 0.2 nsec. The cryoEM density map is utilized as a restraint

potential in the annealing so that both map information and AMBER force field information are

simultaneously utilized to obtain an optimum model consistent with the data(*34*). The protein

force field used the ff14SB force field(*73*) and a generalized Born implicit solvent model with

igb=8(*74*), and a nonbonded cutoff of 20 Å. The relative weight of real-space map-based

restraints and the force field was fixed using *fcons=0.02*. For colE1-T, information from the

symmetrized map was integrated into the modeling procedure during manual remodeling in coot,

but map-based refinement in phenix, ISOLDE, and AMBER were against the asymmetric map.

TolC without colE1-T was modeled similarly but using the TolC-colE1-T structure as a starting

point instead of 1TQQ, and without final AMBER refinement.

Molecular representations were generated with Chimera, ChimeraX(*75*) or PyMOL (Schrödinger,

LLC).

575
References
577

1.   T. Kawabata, Detection of multiscale pockets on protein surfaces using mathematical
morphology. *Proteins* **78**, 1195-1211 (2010).
2.   T. Kawabata, Detection of cave pockets in large molecules: Spaces into which internal
probes can enter, but external probes from outside cannot. *Biophys Physicobiol* **16**, 391-
406 (2019).
3.   T. Kawabata, N. Go, Detection of pockets on protein surfaces using small and large probe
spheres to find putative ligand binding sites. *Proteins* **68**, 516-529 (2007).
4.   J. Yang *et al.*, The I-TASSER Suite: protein structure and function prediction. *Nat
Methods* **12**, 7-8 (2015).
5.   C. Alexander, E. T. Rietschel, Invited review: Bacterial lipopolysaccharides and innate
immunity. *Journal of Endotoxin Research* **7**, 167-202 (2001).

6.   S. D. Zakharov, X. S. Wang, W. A. Cramer, The Colicin E1 TolC-Binding Conformer:
Pillar or Pore Function of TolC in Colicin Import? *Biochemistry* **55**, 5084-5094 (2016).

7.   Y. Kamio, H. Nikaido, Outer membrane of Salmonella typhimurium: accessibility of
phospholipid head groups to phospholipase c and cyanogen bromide activated dextran in
the external medium. *Biochemistry* **15**, 2561-2570 (1976).

8.   J. T. C. Freeman, W. C. Wimley, TMBB-DB: a transmembrane β-barrel proteome
database. *Bioinformatics* **28**, 2425-2430 (2012).

9.   W. A. Cramer, O. Sharma, S. D. Zakharov, On mechanisms of colicin import: the outer
membrane quandary. *Biochem J* **475**, 3903-3915 (2018).

10.   S. K. Buchanan *et al.*, Structure of colicin I receptor bound to the R-domain of colicin Ia:
implications for protein import. *EMBO J* **26**, 2594-2604 (2007).

11.   G. Kurisu *et al.*, The structure of BtuB with bound colicin E3 R-domain implies a
translocon. *Nat Struct Biol* **10**, 948-954 (2003).

12.   E. Cascales *et al.*, Colicin Biology. *Microbiology and Molecular Biology Reviews* **71**, 158-
229 (2007).

13.   N. G. Housden *et al.*, Directed epitope delivery across the Escherichia coli outer
membrane through the porin OmpF. *Proc Natl Acad Sci U S A* **107**, 21412-21417 (2010).

14.   O. Sharma *et al.*, Structure of the complex of the colicin E2 R-domain and its BtuB
receptor. The outer membrane colicin translocon. *J Biol Chem* **282**, 23163-23170 (2007).

15.   H. Benedetti *et al.*, Individual domains of colicins confer specificity in colicin uptake, in
pore-properties and in immunity requirement. *J Mol Biol* **217**, 429-439 (1991).

16.   H. Benedetti *et al.*, Comparison of the Uptake Systems for the Entry of Various BtuB
Group Colicins into Escherichia coli. *Microbiology* **135**, 3413-3420 (1989).

17.   P. Elkins, A. Bunker, W. A. Cramer, C. V. Stauffacher, A mechanism for toxin insertion
into membranes is suggested by the crystal structure of the channel-forming domain of
colicin E1. *Structure* **5**, 443-458 (1997).

18.   C. A. Plate, S. E. Luria, Stages in colicin K action, as revealed by the action of trypsin.
Proc Natl Acad Sci U S A **69**, 2030-2034 (1972).

19.   J. Dankert, S. M. Hammond, W. A. Cramer, Reversal by trypsin of the inhibition of active
transport by colicin E1. *J Bacteriol* **143**, 594-602 (1980).

20.   J. R. Dankert, Y. Uratani, C. Grabau, W. A. Cramer, M. Hermodson, On a domain
structure of colicin E1. A COOH-terminal peptide fragment active in membrane
depolarization. *J Biol Chem* **257**, 3857-3863 (1982).

21.   H. Benedetti, R. Lloubes, C. Lazdunski, L. Letellier, Colicin A unfolds during its
translocation in Escherichia coli cells and spans the whole cell envelope when its pore has
formed. *EMBO J* **11**, 441-447 (1992).

22.   S. D. Zakharov, X. S. Wang, W. A. Cramer, The Colicin E1 TolC-Binding Conformer:
Pillar or Pore Function of TolC in Colicin Import? *Biochemistry* **55**, 5084-5094 (2016).

23.   S. D. Zakharov *et al.*, Colicin occlusion of OmpF and TolC channels: outer membrane
translocons for colicin import. *Biophys J* **87**, 3901-3911 (2004).

24.   K. S. Jakes, The Colicin E1 TolC Box: Identification of a Domain Required for Colicin E1
Cytotoxicity and TolC Binding. *J Bacteriol* **199**, (2017).

25.   M. Wiener, D. Freymann, P. Ghosh, R. M. Stroud, Crystal structure of colicin Ia. *Nature*
385, 461-464 (1997).

26.   N. G. Housden *et al.*, Intrinsically disordered protein threads through the bacterial outer-
membrane porin OmpF. *Science* **340**, 1570-1574 (2013).

27.   A preprint of this work was first published on *bioRxiv* (692251) in July 2019 and updated
with the structure May 2021.

28.   K. M. Storek *et al.*, Monoclonal antibody targeting the beta-barrel assembly machine of
Escherichia coli is bactericidal. *Proc Natl Acad Sci U S A* **115**, 3692-3697 (2018).

29.   S. J. Budiardjo *et al.*, High-Yield Preparation of Outer Membrane Protein Efflux Pumps
640   by in Vitro Refolding is Concentration Dependent. *J Membr Biol* **254**, 41-50 (2021).
641   30.   V. Koronakis, A. Sharff, E. Koronakis, B. Luisi, C. Hughes, Crystal structure of the
642   bacterial membrane protein TolC central to multidrug efflux and protein export. *Nature*
643   **405**, 914-919 (2000).
644   31.   M. K. Higgins *et al.*, Structure of the ligand-blocked periplasmic entrance of the bacterial
645   multidrug efflux protein TolC. *J Mol Biol* **342**, 697-702 (2004).
646   32.   N. G. Housden *et al.*, Directional Porin Binding of Intrinsically Disordered Protein
647   Sequences Promotes Colicin Epitope Display in the Bacterial Periplasm. *Biochemistry* **57**,
648   4374-4381 (2018).
649   33.   R. Dhar, R. Feehan, J. S. G. Slusky, Membrane Barrels Are Taller, Fatter, Inside-Out
650   Soluble Barrels. *J Phys Chem B* **125**, 3622-3628 (2021).
651   34.   X. Wu, S. Subramaniam, D. A. Case, K. W. Wu, B. R. Brooks, Targeted conformational
652   search with map-restrained self-guided Langevin dynamics: application to flexible fitting
653   into electron microscopic density maps. *J Struct Biol* **183**, 429-440 (2013).
654   35.   C. J. Williams *et al.*, MolProbity: More and better reference data for improved all-atom
655   structure validation. *Protein Sci* **27**, 293-315 (2018).
656   36.   B. A. Barad *et al.*, EMRinger: side chain-directed model and map validation for 3D cryo-
657   electron microscopy. *Nat Methods* **12**, 943-946 (2015).
658   37.   M. Masi, P. Vuong, M. Humbard, K. Malone, R. Misra, Initial steps of colicin E1 import
659   across the outer membrane of Escherichia coli. *J Bacteriol* **189**, 2667-2676 (2007).
660   38.   K. R. Brunden, Y. Uratani, W. A. Cramer, Dependence of the conformation of a colicin
661   E1 channel-forming peptide on acidic pH and solvent polarity. *The Journal of biological*
662   *chemistry* **259**, 7682-7687 (1984).
663   39.   R. N. Besingi, P. L. Clark, Extracellular protease digestion to evaluate membrane protein
664   cell surface localization. *Nat Protoc* **10**, 2074-2080 (2015).
665   40.   H. H. Tuson, J. S. Biteen, Unveiling the inner workings of live bacteria using super-
666   resolution microscopy. *Anal Chem* **87**, 42-63 (2015).
667   41.   N. Al-Husini *et al.*, BR-Bodies Provide Selectively Permeable Condensates that Stimulate
668   mRNA Decay and Prevent Release of Decay Intermediates. *Mol Cell* **78**, 670-682 e678
669   (2020).
670   42.   F. W. Studier, P. Daegelen, R. E. Lenski, S. Maslov, J. F. Kim, Understanding the
671   differences between genome sequences of Escherichia coli B strains REL606 and
672   BL21(DE3) and comparison of the E. coli B and K-12 genomes. *J Mol Biol* **394**, 653-680
673   (2009).
674   43.   S. Bakshi, A. Siryaporn, M. Goulian, J. C. Weisshaar, Superresolution imaging of
675   ribosomes and RNA polymerase in live Escherichia coli cells. *Mol Microbiol* **85**, 21-38
676   (2012).
677   44.   P. Rassam *et al.*, Supramolecular assemblies underpin turnover of outer membrane
678   proteins in bacteria. *Nature* **523**, 333-336 (2015).
679   45.   M. Chavent *et al.*, How nanoscale protein interactions determine the mesoscale dynamic
680   organisation of bacterial outer membrane proteins. *Nat Commun* **9**, 2846 (2018).
681   46.   P. White *et al.*, Exploitation of an iron transporter for bacterial protein antibiotic import.
682   *Proc Natl Acad Sci U S A* **114**, 12051-12056 (2017).
683   47.   C. Kleanthous, P. Rassam, C. G. Baumann, Protein-protein interactions and the
684   spatiotemporal dynamics of bacterial outer membrane proteins. *Curr Opin Struct Biol* **35**,
685   109-115 (2015).
686   48.   J. A. Bohnert, S. Schuster, M. Szymaniak-Vits, W. V. Kern, Determination of real-time
687   efflux phenotypes in Escherichia coli AcrB binding pocket phenylalanine mutants using a
688   1,2'-dinaphthylamine efflux assay. *PLoS One* **6**, e21196 (2011).

49. J. A. Bohnert, B. Karamian, H. Nikaido, Optimized Nile Red efflux assay of AcrAB-TolC
multidrug efflux system shows competition between substrates. *Antimicrob Agents*
Chemother **54**, 3770-3775 (2010).
50. M. A. Seeger *et al.*, Engineered disulfide bonds support the functional rotation mechanism
of multidrug efflux pump AcrB. *Nat Struct Mol Biol* **15**, 199-205 (2008).
51. R. Iyer, A. Ferrari, R. Rijnbrand, A. L. Erwin, A fluorescent microplate assay quantifies
bacterial efflux and demonstrates two distinct compound binding sites in AcrB.
Antimicrob Agents Chemother **59**, 2388-2397 (2015).
52. R. Misra, K. D. Morrison, H. J. Cho, T. Khuu, Importance of Real-Time Assays To
Distinguish Multidrug Efflux Pump-Inhibiting and Outer Membrane-Destabilizing
Activities in Escherichia coli. *J Bacteriol* **197**, 2479-2488 (2015).
53. P. Rassam *et al.*, Intermembrane crosstalk drives inner-membrane protein organization in
Escherichia coli. *Nat Commun* **9**, 1082 (2018).
54. W. A. Cramer, O. Sharma, S. D. Zakharov, On mechanisms of colicin import: the outer
membrane quandary. *Biochemical Journal* **475**, 3903-3915 (2018).
55. R. Taylor, J. W. Burgner, J. Clifton, W. A. Cramer, Purification and characterization of
monomeric Escherichia coli vitamin B12 receptor with high affinity for colicin E3. *J Biol*
Chem **273**, 31113-31118 (1998).
56. M. W. Franklin *et al.*, Efflux Pumps Represent Possible Evolutionary Convergence onto
the beta-Barrel Fold. *Structure* **26**, 1266-1274.e1262 (2018).
57. C. Kleanthous, Swimming against the tide: progress and challenges in our understanding
of colicin translocation. *Nat Rev Microbiol* **8**, 843-848 (2010).
58. F. Hagn, M. L. Nasr, G. Wagner, Assembly of phospholipid nanodiscs of controlled size
for structural studies of membrane proteins by NMR. *Nat Protoc* **13**, 79-98 (2018).
59. I. Wiegand, K. Hilpert, R. E. Hancock, Agar and broth dilution methods to determine the
minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat Protoc* **3**, 163-
175 (2008).
60. L. Daury *et al.*, Tripartite assembly of RND multidrug efflux pumps. *Nat Commun* **7**,
10731 (2016).
61. F. Weis, W. J. H. Hagen, Combining high throughput and high quality for cryo-electron
microscopy data collection. *Acta Crystallogr D Struct Biol* **76**, 724-728 (2020).
62. D. N. Mastronarde, Advanced Data Acquisition From Electron Microscopes With
SerialEM. . *Microscopy and Microanalysis* **24**, 864-865 (2018).
63. W. J. Rice *et al.*, Routine determination of ice thickness for cryo-EM grids. *J Struct Biol*
204, 38-44 (2018).
64. S. J. Budiardjo *et al.*, High yield preparation of outer-membrane protein efflux pumps by
in vitro refolding is concentration dependent. *bioRxiv*,
2020.2009.2014.296756 (2020).
65. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).
66. J. Zivanov *et al.*, New tools for automated high-resolution cryo-EM structure
determination in RELION-3. *Elife* **7**, (2018).
67. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *J Struct Biol* **192**, 216-221 (2015).
68. T. Wagner *et al.*, SPHIRE-crYOLO is a fast and accurate fully automated particle picker
for cryo-EM. *Commun Biol* **2**, 218 (2019).
69. G. Cardone, J. B. Heymann, A. C. Steven, One number does not fit all: mapping local
variations in resolution in cryo-EM reconstructions. *J Struct Biol* **184**, 226-236 (2013).
70. P. V. Afonine *et al.*, Real-space refinement in PHENIX for cryo-EM and crystallography.
Acta Crystallogr D Struct Biol **74**, 531-544 (2018).

71. T. I. Croll, ISOLDE: a physically realistic environment for model building into low-resolution electron-density maps. *Acta Crystallogr D Struct Biol* **74**, 519-530 (2018).

72. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).

73. J. A. Maier *et al.*, ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J Chem Theory Comput* **11**, 3696-3713 (2015).

74. H. Nguyen, D. R. Roe, C. Simmerling, Improved Generalized Born Solvent Model Parameters for Protein Simulations. *J Chem Theory Comput* **9**, 2020-2034 (2013).

75. T. D. Goddard *et al.*, UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci* **27**, 14-25 (2018).

Acknowledgments

We gratefully acknowledge Daniel Montezano, Pinakin Sukthankar, Rik Dhar, Dwight Deay, Scott Lovell, Matthias Wolf, Alexander Little, Heather Shinogle, and Sarah Noga for discussions and feedback, Mark Richter for the use of his fluorometer, Rajeev Misra for the pTrc vector containing the TolC gene, Vasileios Petrou for guidance on nanodiscs, Chamani Perera for peptide synthesis, Karen Marom for editorial guidance. We thank the Office of Advanced Research computing (OARC) at Rutgers for high-performance computing resources.

Funding:
NIH award R21-GM128022 to JSB
NIGMS awards DP2GM128201, P20GM113117, P20GM103638 and the Gordon and Betty Moore Inventor Fellowship to JSGS,
NIGMS awards P20 GM103418 and 2K12GM063651 to SJB.

Author contributions:
Conceptualization: SJB, JSGS
Methodology: SJB, JTK, JSGS, JSB
Investigation: SJB, JJS, ALC, API, VKW, EF, DAC, JTK
Writing—original draft: SJB, JSGS
Writing—review & editing: SJB, JSGS, JSB, JTK

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials." All strains used are commercially available. All plasmids available through Addgene. CryoEM maps and models have been deposited with accession codes EMD-21960, EMD-21959, PDB ID 6WXI, and PDB ID 6WXH. All other data is available in the main text or the supplementary materials.

Supplementary Materials for

Colicin E1 opens its hinge to plug TolC

S. Jimmy Budiardjo, Jacqueline J. Stevens, Anna L. Calkins, Ayotunde P. Ikujuni, Virangika K. Wimalasena, Emre Firlar, David A. Case, Julie S. Biteen, Jason T. Kaelber and Joanna S.G. Slusky[*]

*Corresponding author. Email: slusky@ku.edu

This PDF file includes:

      Figs. S1 to S4
      Tables S1 to S2
      Movie caption S1

Other Supplementary Materials for this manuscript include the following:
[use this section only if you have movies, audio or data files]
      Movie S1

812 **Fig. S1.**

S257

G43

10Å

813
814
815 **Figure S1. TolC box density** cropped from the fully asymmetric cryoEM map of the TolC/colE1-T complex.
816 Mutations at positions G43 and S257 in the TolC (grey sticks) barrel make direct contact with colE1-T (green). These
817 mutations were previously shown to abolish WT colE1 function(*37*). Mutations to bulkier side chains introduce a
818 steric clash that prevents binding.
819

Fig. S2.

Figure S2. Co-elution of colE1-T with TolC at 200 mM NaCl (black). Under this higher salt
concentration, when TolC (purple) and colE1-T (red) are mixed, there is a smaller peak shift than
that seen with colE1-TR and the presence of a shoulder (black arrow). SDS-PAGE of the shoulder
shows presence of both TolC (purple arrow) and colE1-T (red arrow). Although some binding
was detected, this higher salt concentration prevents full binding as indicated by a much fainter
band for colE1-T than seen at the lower salt concentration (Figure 4A).

829 **Fig. S3.**

830
Figure S3 Single-molecule microscopy. Fluorescence images overlaid on outlines of living *E.*
coli cells from phase-contrast microscopy for WT and Δ*tolC* for colE1-TR-GFP and counts of
cells where colE1-TR-GFP punctum formation was observed. (A) 97% of cells showed no
binding of Cy3-labeled colE1-TR to Δ*btuB* (B) ColE1-TR-GFP forms similar puncta as Cy3-
labeled ColE1-TR (main text Figure 4C). (C) No binding of Cy3-labeled colE1-T to WT or Δ*tolC*
cells was detected with Cy3-labeled colE1-T. Scale bars: 2 μm.
837
838

839   **Fig. S4.**

Erythromycin

Figure S4. Minimum Inhibitory Concentration of Erythromycin with colE1-TR TolA box
deletion construct. The MIC for erythromycin with 100 μM colE1-TR or colE1-TR$_{\Delta1-40}$ is
identical indicating that TolA engagement by colE1-TR does not contribute to antibiotic
susceptibility.

848 **Table S1.**

849 Cryo-EM data collection, refinement, and validation statistics

	TolC (EMDB-21960) (PDB 6WXI)	TolC + colE1 (EMDB-21959) (PDB 6WXH)
Data collection and processing		
Electron microscope	Talos Arctica	Talos Arctica
Magnification	130,000×	205,000×
Voltage (kV)	200	200
Electron exposure (e–/Å^2)	33.04	35.96
Defocus range (μm)	0.5-2.6	0.4-2.1
Pixel size (Å)	1.038	0.6578
Symmetry imposed	C_3	C_1 (C_3)
Initial particle images (no.)	2,092,678	339,779
Final particle images (no.)	778,220	179,834
Map resolution (Å)	2.84	3.09 (2.81)
FSC threshold	0.143	0.143
Refinement		
Initial model used (PDB code)	6WXH	1TQQ
Model resolution (Å)	3.11	3.37
FSC threshold	0.5	0.5
R.m.s. deviations		
Bond lengths (Å)	0.0072	0.012
Bond angles (°)	1.11	1.76
Validation		
MolProbity score	1.33	0.50
Clashscore	4.42	0.00
Poor rotamers (%)	1.40	0.18
Ramachandran plot		
Favored (%)	98.12	98.31
Allowed (%)	1.88	1.32
Disallowed (%)	0	0.37

850
851

852   **Table S2.**

853   Mean minimum inhibitory concentrations (MICs) of antimicrobials in the presence of colE1-TR.

	Erythromycin (µg/mL)	Ciprofloxacin (ng/mL)	Benzalkonium Chloride (µg/mL)
WT	625.00	56.25	50
ΔtolC	7.03	10.16	4.125
WT + 10 µM colE1-TR	312.50	75.00	50
WT + 100 µM colE1-TR	312.50	28.13	25

855
856
857

Movie S1.

A representative 8-second movie of colE1-TR bound to a live cell. ColE1-TR localizes on, and remains bound to, the extracellular surface of *E. coli*. Fluorescence movie of Cy3-labeled colE1-TR on living WT *E. coli* overlaid on outline of the *E. coli* cell from phase-contrast microscopy. Continuous imaging at 25 frames per second; the movie plays in real time at the same speed. This movie corresponds to the first 8 seconds of data used to attain the WT image in Figure 4C. Scale bar: 2 μm.

866
867
868
869