Evaluation of longitudinal time-lapsed *in vivo* micro-CT for monitoring fracture healing in

mouse femur defect models

Esther Wehrle[1], Duncan C Tourolle né Betts[1], Gisela A Kuhn[1], Ariane C Scheuren[1], Sandra

Hofmann[1,2], Ralph Müller[1*]

[1] Institute for Biomechanics, ETH Zurich, Zurich, Switzerland, [2] Department of Biomedical

Engineering and Institute for Complex Molecular Systems, Eindhoven University of

Technology, The Netherlands.

Corresponding author:

Ralph Müller, PhD

Institute for Biomechanics

ETH Zurich

Leopold-Ruzicka-Weg 4

8093 Zurich, Switzerland

Email: ram@ethz.ch

Abstract

Longitudinal *in vivo* micro-computed tomography (micro-CT) is of interest to non-invasively capture the healing process of individual animals in preclinical fracture healing studies. However, it is not known whether longitudinal imaging itself has an impact on callus formation and remodeling. In this study, a scan group received weekly micro-CT measurements (week 0-6), whereas controls were only scanned post-operatively and at week 5 and 6. Registration of consecutive scans using a branching scheme (bridged vs. unbridged defect) combined with a two-threshold approach enabled assessment of localized bone turnover and mineralization kinetics relevant for monitoring callus remodeling. Weekly micro-CT application did not significantly change any of the assessed callus parameters in the defect and periosteal volumes. This was supported by histomorphometry showing only small amounts of cartilage residuals in both groups, indicating progression towards the end of the healing period. Also, immunohistochemical staining of Sclerostin, previously associated with mediating adverse radiation effects on bone, did not reveal differences between groups.

The established longitudinal *in vivo* micro-CT-based approach allows monitoring of healing phases in mouse femur defect models without significant effects of anesthesia, handling and radiation on callus properties. Therefore, this study supports application of longitudinal *in vivo* micro-CT for healing-phase-specific monitoring of fracture repair in mice.

Introduction

Adequate monitoring and characterization of the healing process is important in preclinical fracture healing studies. One recent approach to non-invasively capture the formation and remodeling of the osseous fracture callus is the repeated application of *in vivo* micro-computed tomography (micro-CT)[1-3], allowing for three-dimensional assessment of callus structures over time. A further development is the consecutive registration of time-lapsed *in vivo* callus measurements[4]. In a recent study, we developed a method for registration of time-lapsed *in vivo* scans in a femur defect model in mice using a branching scheme (registration of whole scan for bridged defects; separate registration of the two fragments for unbridged defects)[5], so that we are now able to also assess dynamic parameters such as bone formation and resorption. In combination with two- and multi-threshold approaches applied in recent studies[5,6], this allows for better understanding of localized bone turnover and mineralization kinetics during callus formation and remodeling. Whereas many studies focused on understanding the early healing phases (inflammation, repair) with the aim of increasing bone formation and achieving earlier bone union, recent studies also indicate a potential to improve fracture healing outcome via modulation of callus remodeling[7-9]. To address this, time-lapsed *in vivo* micro-CT-based monitoring approaches seem particularly suitable, as not only bone formation but also bone resorption can be reliably monitored over time. Furthermore, as each animal can be followed individually throughout the healing process with low variance in the assessed parameters, animal numbers can be reduced compared to well-established cross-sectional studies with endpoint three-dimensional micro-CT and two-dimensional histological callus evaluation.

In non-fractured bone, longitudinal time-lapsed *in vivo* micro-CT has been increasingly used

to monitor changes in bone properties associated with different diseases and external

factors, e.g. estrogen-deficiency[10-12], mechanical-(un)loading[13-15], and drug application[16].

However, several studies indicate that anesthesia, cumulative radiation dosage and stress

due to the required handling for the CT measurements may have effects on animal well-

being and on trabecular and cortical bone properties[11,17-21]. Isoflurane is the most commonly

used inhalation anesthetic in rodents[22], characterized by a quick on- and offset of anesthesia

and low metabolism rate[22]. According to the EU Directive 2010/63, the severity of repeated

isoflurane anesthesia can be categorized as mild, although repeated anesthesia was

considered worse than a single session with sex-dependent differences in perceiving the

severity of a procedure[17]. Specifically, female mice were shown to be more susceptible to

anesthesia-induced effects on well-being compared to male mice[17]. Nevertheless, in both

sexes repeated isoflurane anesthesia caused only short-term mild distress and impairment

of well-being, mainly in the immediate post-anesthetic period[17]. Radiation has also been

shown to have dosage-dependent effects on bone cells *in vivo* and *in vitro*: Whereas high

dose x-ray radiation (2.5-8Gy) was associated with reduced osteoblast and osteoclast

proliferation[23-25], lower doses (<2Gy) had a stimulatory effect on osteoclasts. Some studies

reported radiation-associated effects on structural bone parameters with stronger and

earlier effects seen in trabecular bone compared to cortical bone, whereas other studies did

not see any changes[11,19,21,26,27]. Similarly, pre- and post-operative radiation treatments have

been associated with changes in fracture healing. Whereas high dose x-ray radiation (5-8Gy)

was associated with an impairment of fracture healing, lower doses (0.5-1Gy) accelerated

endochondral and intramembranous ossification during the repair process[28-31]. These

findings indicate the importance of study-specific adaptation of micro-CT protocols, and to

protect animals scanning times and radiation settings should be minimized.

Recently, several fracture healing studies have applied longitudinal *in vivo* micro-CT to

monitor callus formation under different conditions (e.g. osteoporosis, polytrauma) and

treatments (e.g. mechanical loading, pharmaceuticals)[2-4,32,33]. However, none of these

studies assessed, whether longitudinal imaging itself has an impact on callus formation and

remodeling. As several studies have reported adverse effects of longitudinal imaging during

normal bone remodeling (e.g. reduction in trabecular and cortical thickness)[11,19,21,27], there is

a need to also assess imaging-associated impact on the callus formation during the highly

metabolically active process of fracture healing.

Therefore, the objectives of this study were to establish an *in vivo* micro-CT based approach

for longitudinal monitoring of fracture healing in a mouse femur defect model and to assess

the combined effect of radiation, anesthesia and handling associated with weekly time-

lapsed micro-CT measurements on callus properties during the remodeling phase of fracture

healing.

Results

In order to enable longitudinal monitoring of fracture healing in single animals, we

established a time-lapsed *in vivo* micro-CT based approach for mouse femur defect models.

By registration of consecutive scans, structural and dynamic callus parameters can be

followed in three callus sub-volumes (defect center, defect periphery, cortical fragment

periphery) and the adjacent cortical fragments over time (Figure 1). To capture potential

effects of consecutive micro-CT measurements on callus properties, bone parameters of a

scan group subjected to weekly measurements between weeks 0 and 6 were compared to

control animals, that were scanned only post-operatively (d0) and after 5 and 6 weeks,

respectively (for detailed study design see Supplementary Fig. S1 online).

General physical observation

All mice recovered rapidly from surgery. During the healing period, in both groups, the

animals' body weight did not significantly change compared to pre-operative values without

significant differences between the two groups (see Supplementary Fig. S1 online). We also

assessed the social interaction between mice (animals sitting in groups vs. separated from

others) and the nesting behavior (start of nest building after surgery). Specifically, all animals

started to build nests on the day of surgery and were found in groups in the nest in the

morning of post-operative day 1. Throughout the further healing period social interaction

between mice and nesting behavior did not differ from pre-surgical observations and was

similar for animals of the scan and control groups.

Volumes of interest (VOI) for evaluation by time-lapsed *in vivo* micro-CT

In order to exclude bias in the further micro-CT analyses, we compared the size of the

different VOIs (depicted in Fig. 1) in the control and the scan group. One animal from the

scan group could not be included in the analysis due to failure in VOI generation caused by

too little cortical bone being present in the field of view (for details on VOI generation see [5]).

The VOIs encompassed the following volume for the control (n=8) and scan group (n=10):

$2.51\pm0.34mm^3$ vs. $2.60\pm0.34mm^3$ for the defect center (DC), $17.26\pm2.21mm^3$ vs.

18.73±4.64mm^3 for the defect periphery (DP), 1.86±0.48mm^3 vs. 1.73±0.45mm^3 for the

cortical fragments (FC), 14.47±3.30mm^3 vs. 13.10±2.66mm^3 for the fragment periphery (FP).

The total volume (TOT=DC+DP+FC+FP) between the inner pins of the fixator was

36.10±2.81mm^3 for the control and 36.15±3.74mm^3 for the scan group. No significant

differences in volume were detected in any of the VOIs between groups.

Longitudinal monitoring of fracture healing by time-lapsed _in vivo_ micro-CT

In the scan group (n=10), the repeated micro-CT scans (1x/week, Fig. 2) covered the period

from the day of the defect surgery (d0) until post-operative week 6 with distinct callus

characteristics indicative of the different healing phases (inflammation, repair, remodeling;

significant weekly differences in bone parameters indicated in Fig. 3). From week 0-1 to

week 1-2 a significant 5.7x increase in bone formation was detected (Fig. 3a) in the total VOI

(TOT=DC+DF+FC+FP, p<0.0001; Fig. 1), indicating progression from the inflammation to the

reparative phase. This led to a significant gain in bone volume by week 2 (BV/TV$_{week2}$: 39±7%

vs. BV/TV$_{week0}$: 25±3%, p=0.0134; Fig. 3b). From week 1-2 to week 2-3 a significant 2.8x

increase in resorptive activities (p=0.0020; Fig. 3a) was seen, indicating the progression from

the repair to the remodeling phase. Two weeks after surgery the highly mineralized bone

fraction in the TOT VOI was significantly lower compared to the post-operative

measurement (BV$_{645}$/BV$_{395}$ in week 2: 59±5% vs. BV$_{645}$/BV$_{395}$ in week 0: 84±1%, _p_<0.0001),

indicating formation of mineralized callus of low density. From post-operative week 2

onwards, the highly mineralized bone fraction in the TOT VOI gradually increased in all

subsequent weeks of the healing period reaching statistical significance by week 5

(BV$_{645}$/BV$_{395}$ in week 2: 59±5% vs. BV$_{645}$/BV$_{395}$ in week 5: 79±3%, p=0.0134; Fig. 3c).

In order to better capture the regions where bone is mainly formed and resorbed, we evaluated the different VOIs separately (Table 1): In the early post-operative phase from week 0-1 to week 1-2 a strong onset of bone formation was seen in the DC and FP sub-volumes, leading to significant 11.8x and 3.4x gain in mineralized tissue from week 0 to week 2 for DC (p=0.0090) and FP (p=0.0091), respectively. This indicates that both intra-cortical as well as periosteal callus formation takes place in this femur defect model. In both VOIs bone formation triggered the initiation of bone resorption from week 2-3. In detail, there was a significant 6.2x and 2.9x increase in bone resorption from week 1-2 to week 2-3 in the DC (p=0.0040) and the FP VOI (p=0.0079), respectively. Compared to the DC and FP VOIs, the initiation of bone formation was much less pronounced in the DP sub-volume with less deposition of mineralized tissue leading to only little peripheral callus formation in this VOI (week 2-6) and subsequently low bone resorption activities from week 2-3 to week 5-6. In all three regions (DC, DP, FP) the fraction of highly mineralized bone considerably increased from week 2 to week 5 (+192%, +774%, +227%), indicating callus maturation. In this femur defect model, callus formation and remodeling mainly took place in the defect region (DC) with only little peripheral callus formation and remodeling (DP). Looking at the cortical fragments (FC), a significant 3.2x increase in resorptive activities was detected in week 1-2 compared to week 0-1 (p=0.0421), whereas no significant weekly change in bone formation activities was seen in this region throughout the assessed healing period. This resulted in a significant 24% reduction in bone volume from week 0 to week 4 (p=0.0091). The fraction of highly mineralized bone also gradually decreased in the remaining osseous tissue reaching

statistical significance by week 3 (BV_{645}/BV_{395} in week 3 reduced by 9% compared to week 0,

p=0.0027).

Influence of the longitudinal _in vivo_ micro-CT protocol on callus properties

To assess the combined effects of radiation, anesthesia and handling associated with weekly

micro-CT measurements (week 0 - week 6) on callus properties, callus parameters of the

scan group were compared to control animals that were scanned only directly post-

operatively (d0) and after 5 and 6 weeks (Fig. 4). To exclude bias in the further micro-CT

analyses, we assessed the post-operative bone volume (BV/TV on d0) and did not see any

significant differences between groups: 1.82±0.39% (control) vs. 1.47±0.15% (scan) in DC,

0.16±0.12% (control) vs. 0.13±0.06% (scan) in DP, 56.82±2.31% (control) vs. 57.14±6.99%

(scan) in FC, 4.66±0.30% (control) vs. 4.74±0.76% (scan) in FP, 26.92±4.69% (control) and

24.90±2.88% (scan) in TOT.

During the remodeling phase (post-operative weeks 5-6), no significant difference in bone

turnover was seen in the total volume (TOT) between the inner pins of the fixator (BFR:

0.76±0.23% in controls, 0.64±0.15% in scan group; BRR: 0.82±0.25% in controls, 0.78±0.26%

in scan group). Also, bone volume did not significantly change due to the applied micro-CT

protocol with a similar fraction of highly mineralized bone after 5 weeks (BV_{645}/BV_{395}: 79±3%

in controls, 79±3% in scan group) and 6 weeks (BV_{645}/BV_{395}: 81±3% in controls, 82±4% in

scan group).

As radiation effects are known to vary for different bone cell types and locations[30,34,35], we

assessed the four sub-volumes separately (Table 2). The applied micro-CT protocol did not

significantly affect bone formation and resorption activities in the callus VOIs (DC, DP and FP)

from week 5-6 with similar bone volume observed in week 5 and 6 for controls and scanned

animals. In both groups, bone volume remained stable from week 5 to week 6, whereas the

density of the mineralized tissue increased during the same period (DC, DP, FP). Looking at

the adjacent cortical fragments (FC) also no significant differences were detected in any of

the assessed parameters between the control group and the scan group. Similarly to the

callus VOIs, bone volume also remained stable in this region from week 5 to week 6.

However, in contrast to the callus VOIs, the density of the mineralized tissue in the cortical

VOI FC did not change from week 5 to week 6. In week 6, the FC VOI comprised 33% and

35% of the osseous tissue in the total VOI (TOT) for the control and scan group, respectively.

In the DC VOI, 37% (control group) and 36% (scan group) of the total bone volume were

seen. Less osseous tissue was detected in the two peripheral VOIs, 11% (control group) and

12% (scan group) in the DP and 19% (control group) and 17% (scan group) in the FP VOI.

As group comparisons were only performed from week 5-6, we particularly focused on the

defect VOIs (DC+DP) which are most important for evaluating later healing time points

during the remodeling phase of fracture healing. No significant differences in bone turnover,

bone volume and mineralization were seen between the scan and control group (Figure 4 d-

f, Table 2). In addition, according to the standard clinical evaluation of X-rays, the number of

bridged cortices per callus was evaluated in two perpendicular planes and animals with ≥3

bridged cortices were categorized as healed. By week 5, cortical bridging occurred in 75% of

the control animals and 64% of the scanned animals (Table 3). However, we noticed that

non-unions only occurred in defects of ≥1.5mm length. Although the mean defect length

(1.45mm±0.16 in controls, 1.47mm±0.16 in scan group) did not significantly differ between

groups (p=0.816), the percentage of animals with defects ≥1.5mm was different in the

control and the scan group (55% vs. 38%; Table 3). Taking into account only these defects

(≥1.5mm length), both groups had a non-union rate of 67% indicating no significant effect of

the chosen micro-CT protocol on clinical fracture healing outcome.

Histology

Histomorphometric analysis of Safranin-O stained sections did not show any significant

differences in tissue composition (bone, cartilage, fibrous tissue, bone marrow) in the former

defect region between the control and the scan group six weeks after osteotomy (Fig. 5,

Supplementary Fig. S2 online). The bone tissue fraction was 38+7% and 32+6% for the

control and the scan group, respectively. Hardly any cartilage residuals (< 1% of total area)

were present in both groups indicating progression of the healing process in the final

remodeling stage (Supplementary Fig. S3 online).

As radiation-mediated decreased bone formation by osteoblasts has previously been

associated with the inhibition of osteoanabolic Wnt-signaling[36], we assessed Sclerostin

expression in the defect region by immunohistochemistry. Visual inspection did not reveal

any differences in staining intensity and amount of Sclerostin between the scan and control

group (Fig. 5).

Discussion

In this study, longitudinal *in vivo* micro-CT was applied for monitoring the process of fracture

healing in mice. In addition, the combined effect of radiation, anesthesia and handling

associated with the established imaging approach on callus parameters was assessed. Until

now, preclinical fracture healing studies predominantly have a cross-sectional study design

with end-point callus evaluation in sub-groups of animals at different time points during the

healing period. Only recently, several studies in murine diaphyseal femur defect models

stabilized by external fixation[2-4,32] or intramedullary nails[33], have applied longitudinal *in vivo*

micro-CT to monitor the healing progression in each animal over time. These studies were

able to consecutively assess callus parameters (e.g. callus volume and density) at specific

time points during the healing process for up to 12 weeks[2-4,32,33]. By registering consecutive

scans, we were now able to also include dynamic parameters such as bone formation and

resorption in our micro-CT based monitoring approach for fracture healing (for details on

method see [5]). This allows for characterization of the different healing phases seen by

changes in formation and resorption in the osseous callus volume. Compared to histology-

based determination of the bone formation rate via *in vivo* application of fluorescent dyes

(e.g. calcein, alizarin), our micro-CT-based analysis with registration of consecutive scans also

allowed for assessment of the bone resorption rate in individual animals. Furthermore,

implementation of a two-threshold approach allowed to monitor mineralization kinetics of

the fracture callus. Our micro-CT-based approach is therefore particularly suited to better

understand callus remodeling under different conditions (e.g. premedication with

bisphosponates and associated inhibition of callus remodeling) and to assess

pharmacological and mechanical therapies targeting callus remodeling.

Specifically, we saw that the initiation of bone formation (maximum in week 2-3), indicating

the onset of the reparative phase, triggered bone resorption (maximum in week 3-4) with

maximum osseous callus volumes in week 3. These observations are in line with cross-

sectional histomorphometric analyses in similar defect models, where maximum callus area

was seen in weeks 2 and 3[37-39]. From week 4 to week 6, bone formation and resorption

continuously decreased to levels (BFR ≤ 1%/day; BRR ≤ 0.8%/day) seen in normal bone

remodeling of trabecular and cortical bone in mice[14,40-43], indicating proceeding towards the

end of the healing process. Whereas these remodeling activities also led to a decrease in

callus volume from week 4 to week 6, the fraction of highly mineralized bone increased

during the same period indicating advanced callus maturation. These observations were

further supported by histology with no remaining cartilage in the defect region of unions and

only a small amount of cartilage residuals in some non-unions. The findings are also in

accordance with previous longitudinal studies of up to 4 weeks in mice[2,3,32] and 8 weeks in

rats[4] mainly focusing on the inflammation and reparative phase. Similar to our study, the

bone volume continuously increased during the 4/8 week healing period, although no

remodeling to the original bone geometry was seen due to the shorter observation period.

One study in ovariectomized rats with a 12-week post-operative monitoring period detected

maximum osseous callus volumes by week 6, which diminished thereafter, whereas bone

mineral density continuously increased until week 12, similar to our findings. This indicates

that during the reparative phase first low-density bone is formed, which is then further

mineralized during the remodeling phase of fracture healing. Looking at the functional

fracture healing outcome, complete cortical bridging was seen in 64% of the animals, which

is similar to other studies (60-62%) using similar femur defect models with relatively stiff

external fixation (fixator stiffness measured in our study 24N/mm) in mice[2,44]. These

commercially available rigid external fixators are widely used in fracture healing studies as

they also allow for the assessment of biomaterials in non-union defects (induced by

increasing the defect size). Borgiani et al.[37] have recently extensively characterized the

healing process (week 1-3) using external fixators of similar stiffness and observed

endochondral ossification processes, albeit less pronounced compared to semi-rigid fixation.

When comparing the outcome of these studies using rigid fixation, it is important to consider

differences in defect sizes: $0.7mm^2$, 1.19 ± 0.25 mm[44], $1.47\pm0.16mm$ (this study). In contrast

to the other studies, we saw the manifestation of non-unions only in defects ≥1.5mm. This

might be partially due to the fact that in some studies the exact defect length might have

differed from the reported value (saw diameter used for osteotomy) and that non-union

formation was assessed irrespective of defect length. Overall, all longitudinal fracture

healing studies using *in vivo* micro-CT were able to follow the healing progression in single

animals, thereby reducing variance of outcome parameters compared to cross-sectional

studies, also allowing for reduction of animals per group considering the 3R principles of

animal welfare[45,46]. However, when applying longitudinal *in vivo* micro-CT, the effect of

repeated anesthesia, handling and radiation associated with the scans on the study's main

outcome parameters and the general well-being of the animals have to be considered.

In this study, isoflurane was used as anesthetic agent for the *in vivo* micro-CT scans, which is

the most commonly used inhalation anesthetic in longitudinal imaging studies with a fast on-

and offset of anesthesia and low metabolism rate[17]. Despite its general use, isoflurane has

been associated with several adverse effects, such as hepatic degenerative changes[47],

immunomodulation[48], oxidative DNA damage[49] as well as alterations in expression profiles of

oncogene and tumor suppressor genes in the bone marrow[50], which might potentially also

affect fracture healing and the general well-being of the animals. A recent study by

Hohlbaum et al. (2017)[17], which assessed the impact of repeated isoflurane anesthesia (6

times for 45 min at an interval of 3–4 days) and the associated handling on the well-being of

adult C57BL/6JRj mice, categorized the degree of distress as mild according to the EU

Directive 2010/63 with only short-term impairment of well-being, mainly in the immediate

post-anesthetic period. Therefore, isoflurane-based inhalation anesthesia and the associated

handling of the animal is suggested for longitudinal studies with multiple anesthetic sessions

per animal.

In this study, the applied micro-CT settings (55 kVp, 145 µA, 350 ms integration time, 500

projections per 180°, 21 mm field of view (FOV), scan duration ca. 15 min) were adapted

from well-established protocols used for longitudinal monitoring of bone adaptation and

implant integration in tail vertebrae in mice[14,51]. These settings are similar to protocols used

in 2 of the 5 longitudinal fracture healing studies published so far[3,4], while the 3 other

studies did not specify the CT settings[2,32,33]. The radiation dosage (CT dose index, CTDI)

associated with each scan in the current study was previously estimated to be 0.67 Gy per

scan[51,52]. A radiation control experiment using similar settings did not show detrimental

effects of 5 weekly *in vivo* micro-CT scans on bone development in tail vertebrae of mice[52].

However, some studies in non-fractured bone indicate cumulative radiation-associated

effects on bone properties particularly when using high radiation doses (>2G)[19]. The

radiation effects also depend on animal age[19,27] and ovariectomy (OVX)[11], with younger and

ovariectomized animals being more susceptible to radiation. So far, only one study assessed

radiation-associated effects of longitudinal *in vivo* micro-CT on bone healing in a uni-cortical

tibia burr-hole defect model, and did not find significant radiation effects on bone

properties[1]. In this study, similar CT settings (45kVp, 133 µA, 200 ms integration time, 1000

projections per 180°) were used; however, due to the smaller dimensions of the defect and

resulting shorter scanning time, the radiation dosage (0.36 Gy) was lower compared to our

protocol (0.64 Gy).

In the current study, we assessed the combined imaging-associated impact (anesthesia,

handling, radiation; 1x/week for 6 weeks) on callus formation and remodeling in a

diaphyseal femur defect model in adult female C57BL/6J mice. We did not see any significant

imaging-associated changes in bone volume and turnover in the fracture callus during the

remodeling phase (post-operative week 5 and 6). Furthermore, no significant differences in

callus mineralization between scanned (d0, week1-6) and control animals (d0, week 5+6) in

any of the assessed callus volumes (TOT, DC, DP, FP) were observed. The distribution of the

osseous callus volume into the three sub-volumes was also not significantly different

between groups, indicating similar healing patterns. In addition, when separately assessing

the cortical fragments between the inner pins (FC), also no significant differences in any of

the assessed parameters were detected between groups. These 3D micro-CT based results

were supported by 2D histomorphometric analysis, which also showed no significant

differences in tissue composition (bone, cartilage, fibrous tissue, bone marrow) in the former

defect region. In respect to clinical healing outcome, the percentage of unions in animals

above the sub-critical defect length was 67% in both groups with restoration of the

medullary cavity. Defects without cortical bridging only showed small cartilage residuals

(<1%) as visualized by Safranin-O staining, indicating manifestation of non-union formation.

In the defect regions of unions, no cartilage was seen, indicating progression towards the

end of the healing period.

So far only few studies have focused on the underlying mechanisms of radiation-induced

effects on bone mainly focusing on changes in expression profiles of osteoblast and

osteoclast differentiation markers[30,35]. Recently, three consecutive *in vivo* studies by

Chandra et al.[36,53,54] showed, that radiation-induced bone loss[53,54] is mediated via Sclerostin

inhibition of osteoanabolic Wnt-signaling[36]. First, they showed that focal radiation increases

Sclerostin expression in trabecular bone in mice. Using a Sclerostin antibody, they were then

able to block the radiation-induced deterioration in structural bone properties and

consistently, trabecular bone in Sclerostin null mice was resistant to radiation. Based on

these comprehensive analyses, we also assessed Sclerostin by immunohistochemistry, but

did not see any differences in number of stained osteocytes, staining intensity and pattern

between the scan and the control group in any of the assessed callus regions.

In summary, we did not see significant differences in a total of 10 assessed CT and

histological parameters in the total volume (TOT) between the inner pins of the fixator as

well as in the different callus sub-volumes (DC, DP, FP) and the adjacent cortical fragments

(FC) between the scan and the control group. The sub-volume specific analysis also allowed

capturing of distinct features only present in one region (e.g. significant post-operative

increase in bone resorption in FC), that showed contrary progression compared to the

overall picture (significant post-operative increase in bone formation in DC, DP and FP). This

also shows the importance of VOI selection. The other longitudinal fracture healing studies

either assessed a VOI consisting of the defect as well as adjacent bone fragments, or then

only the callus in the defect region including the last slice of intact cortex[2,3], or the callus in

the defect and periosteal region combined[4] was evaluated. An overall VOI including the

defect region and cortical fragments can give an overview of callus characteristics, but sub-

volume-specific differences might be missed. Therefore, analysis of different endosteal and

periosteal callus regions as well as the adjacent cortical volume is favorable.

The current study has several limitations. We based our evaluation of longitudinal time-

lapsed *in vivo* imaging for monitoring fracture healing on micro-CT and histological

assessment of callus properties. Histomorphometry confirmed the micro-CT results.

However, contouring of non-union defects is challenging and might be user dependent as

indicated previously[55]. This could be addressed in future studies by implementing 2D-3D

registration approaches[56], allowing to transfer the automatically computed VOI borders

generated for the micro-CT evaluation to the 2D histology analyses. We did not assess

systemic effects associated with the repeated *in vivo* micro-CT measurements and isoflurane

anesthesia. Due to the required post-operative scan defined in our ethical license, we did

not include a second control group only receiving one micro-CT measurement at the

endpoint. In respect to the underlying mechanisms associated with radiation effects on

fracture healing, we chose to assess Sclerostin by immunohistochemistry, due to the

comprehensive available literature. As proposed by Chandra et al.[36], future studies could

extend the analysis to other Wnt signaling inhibitors (e.g. DKK1) with the potential to find

further targets for prevention of radiation-mediated bone loss. As we only included female

animals in the study, our findings could be affected by the estrous cycle of the animals,

which in future could be addressed by staging of the cycle using vaginal smears[57]. We

minimized the risk of cycle-associated variance by using individual ventilated cages, which

have been associated with the Lee–Boot effect (suppression or prolongation of estrous

cycles) in group-housed adult female mice isolated from male animals[58]. In addition, recent

literature indicates that unstaged female rodents are no more variable than males (revised

in [59]).

Despite these limitations, the longitudinal *in vivo* micro-CT based approach established in

this study allows monitoring of the different healing phases in mouse femur defect models

without significant anesthesia-, handling- and radiation-associated effects on callus

properties. By registering consecutive scans of the defect region for each animal and

implementing a two-threshold approach, data on bone turnover and mineralization kinetics

can be obtained, which is important for targeting impaired callus remodeling under different

conditions (e.g. premedication with bisphosphonates) with pharmacological and mechanical

therapies. Importantly, repeated anesthesia, handling and radiation associated with the

scans did not impair callus formation and remodeling. Therefore, this study supports the

application of longitudinal *in vivo* micro-CT for healing-phase specific monitoring of fracture

repair in mice. Further studies should evaluate the potential of this micro-CT-based

monitoring approach for healing phase-specific discrimination of normal and impaired

healing conditions.

Methods

Study design

A micro-CT based approach for longitudinal *in vivo* monitoring of fracture healing was

established for a mouse femur defect model (for details on study design see Supplementary

Table S1 online). All mice received a femur defect and post-operative micro-CT scans (vivaCT

40, ScancoMedical, Brüttisellen, Switzerland) were performed. The scan group then received

weekly scans of the defect area (weeks 1-6). To assess the combined effect of radiation,

anesthesia and handling associated with weekly micro-CT measurements, controls were only

scanned after 5 weeks. Control animals then received another scan in post-operative week 6

to enable the assessment of both, static and dynamic bone parameters during the final

remodeling phase of fracture healing.

<u>Animals</u>

All animal procedures were approved by the authorities (licence number: 36/2014;

Kantonales Veterinäramt Zürich, Zurich, Switzerland). We confirm that all methods were

carried out in accordance with relevant guidelines and regulations (ARRIVE guidelines and

Swiss Animal Welfare Act and Ordinance (TSchG, TSchV)). To study adult fracture healing

female 12 week-old C57BL/6J mice were purchased from Janvier (Saint Berthevin Cedex,

France) and housed in the animal facility of the ETH Phenomics Center (EPIC; 12h:12h light-

dark cycle, maintenance feed (3437, KLIBA NAFAG, Kaiseraugst, Switzerland), 5

animals/cage) for 8 weeks. At an age of 20 weeks, all animals received a femur defect by

performing an osteotomy (group 1: control group, defect length - 1.45mm ± 0.16, n=8; group

2: scan group, defect length -1.47mm ± 0.16, n=11; housing after surgery: 2-3 animals/cage;

for details on study design see Supplementary Table 1 online). All defect surgeries were

performed by the same veterinarian. Perioperative analgesia (25 mg/L, Tramal®,

Gruenenthal GmbH, Aachen, Germany) was provided via the drinking water two days before

surgery until the third post-operative day. For surgery and micro-CT scans, animals were

anesthetized with isoflurane (induction/maintenance: 5%/1-2% isoflurane/oxygen).

Perioperative handling and monitoring was performed by the surgeon and one other

veterinarian.

Femur osteotomy

In all animals an external fixator (Mouse ExFix, RISystem, Davos, Switzerland; stiffness:

24N/mm) was positioned at the craniolateral aspect of the right femur and attached using

four mounting pins. First, the most distal pin was inserted approximately 2mm proximal to

the growth plate, followed by placement of the most proximal and the inner pins.

Subsequently, a femur defect was created using 2 Gigli wire saws.

Time-lapsed *in vivo* micro-CT

Immediate post-surgery correct positioning of the fixator and the defect was visualized using

a vivaCT 40 (Scanco Medical AG, Brüttisellen, Switzerland) (isotropic nominal resolution: 10.5

μm; 2 stacks of 211 slices; 55 kVp, 145 μA, 350 ms integration time, 500 projections per

180°, 21 mm field of view (FOV), scan duration ca. 15 min). Subsequently, the fracture callus

and the adjacent bone between the inner pins of the fixator were scanned weekly using the

same settings. Scans were registered consecutively using a branching scheme (registration of

whole scan for bridged defects; separate registration of the two fragments for unbridged

defects)[5]. Subsequently, morphometric indices (bone volume - BV, bone volume/total

volume – BV/TV, bone formation rate – BFR, bone resorption rate - BRR) were computed

(threshold: 395mg HA/cm^3; for details on methods see [5]). To assess mineralization

progression, a second threshold (645mg HA/cm^3) was applied and the ratio between highly

and lowly mineralized tissue (BV_{645}/BV_{395}) was calculated. The two selected thresholds are

included in our recently developed multidensity threshold approach[5]. Similar thresholds

have previously been used in fracture healing studies[60,61]. The lower threshold (395mg

HA/cm^3) was selected to capture early callus formation, based on its previous application for

assessment of lowly mineralized tissue in the metabolically active trabecular

compartment[62,63]. The choice of the second threshold (645 mg HA/cm^3) for monitoring callus

mineralization was based on thresholds previously applied to well-mineralized cortical

bone[62,63]. It has to be considered, that micro-CT-based assessment of the mineralization

process is known to be affected by the polychromatic cone beam ($\leq$10% error in the

quantification of the local bone mineral density[64]). We used an aluminium filter to further

reduce the effect of beam hardening[65] and applied scanning settings previously evaluated

for monitoring *in vivo* mineralization kinetics[66]. According to the standard clinical evaluation

of X-rays, the number of bridged cortices per callus was evaluated in two perpendicular

planes (UCT Evaluation V6.5-1, Scanco Medical AG, Brüttisellen, Switzerland). A ''healed

fracture'' was considered as having a minimum of at least three bridged cortices per callus.

For evaluation, four volumes of interest (VOIs) were defined, which were created

automatically from the post-operative measurement (Fig. 1): defect center (DC), defect

periphery (DP), cortical fragment center (FC), and fragment periphery (FP). Data were

normalised to the central VOIs: DC/DC, DP/DC, FC/FC, FP/FC.

Histology

Histological analyses were performed in a sub-set of animals (n=6/group). On day 42, femora

were excised, the femoral head was removed and the samples were placed in 4% neutrally

buffered formalin for 24 hours and subsequently decalcified in 12.5% EDTA for 10-14 days.

The samples were embedded in paraffin and 4.5 µm longitudinal sections were stained with

Safranin-O/Fast Green: Weigert's iron haematoxylin solution (HT1079, Sigma-Aldrich, St.

Louis, MO) - 4min, 1:10 HCl-acidified 70% ethanol - 10s, tap water - 5min, 0.02% Fast Green

(F7258, Sigma-Aldrich, St. Louis, MO) - 3min, 1% acetic acid - 10s, 0.1% Safranin-O (84120,

Fluka, St. Louis, MO) - 5min. Images were taken with Slide Scanner Pannoramic 250 (3D

Histech, Budapest, Hungary) at 20x magnification.

For immunohistochemical staining of Sclerostin, nonspecific sites were blocked (1% BSA/PBS

+ 1% rabbit serum) for 60 min at room temperature. Subsequently, the sections were

incubated with the primary antibody against Sclerostin (AF1589, R&D Systems, Minneapolis,

MN; 1:150 in 1%BSA/PBS + 0.2% rabbit serum) overnight at 4°C. To detect the primary

antibody, a secondary biotinylated rabbit anti-goat-IgG antibody (BAF017, R&D Systems,

Minneapolis, MN) was added for 1 h at room temperature. For signal amplification, the

slides were incubated with avidin-biotin complex (PK-6100 Vector Laboratories, Burlingame,

CA) for 30 min. Diaminobenzidine (Metal Enhanced DAB Substrate Kit, 34065 ThermoFisher

Scientific, Waltham, MA) was used as detection substrate. Counterstaining was performed

with FastGreen (F7258, Sigma-Aldrich, St. Louis, MO). Species-specific IgG was used as

isotype control. Images were taken with Slide Scanner Pannoramic 250 (3D Histech,

Budapest, Hungary) at 40x magnification.

For histomorphometric analysis, the callus and adjacent cortical fragments between the

inner pins of the fixator were manually contoured on Safranin-O stained sections and the

tissue area of this region of interest (ROI) was quantified using commercial software

(n=6/group; Fiji[67]). Similarly, the different tissue types (bone, cartilage, fibrous tissue and

bone marrow) in the ROI were manually contoured and quantified.

Statistics

CT analysis and histomorphometry: Data were tested for normal distribution (Shapiro-Wilk-

Test) and homogeneity of variance (Levene-Test). Depending on the test outcome, group

comparisons (scan versus control group) of data derived at single time points were

performed by two-tailed Student's t-test or Mann-Whitney U-test (IBM SPSS Statistics

Version 23). For statistical evaluation of repeated measurements (scan group) dependent on

results from normality and variance tests, either one-way analysis of variance (ANOVA) with

Bonferroni correction or Friedman test with Dunn correction for multiple comparisons

(GraphPad Prism 7) were performed. Two-way ANOVA was used for longitudinal comparison

of the body weight between the control and the scan group. The level of significance was set

at $p < 0.05$.

Acknowledgements

The authors gratefully acknowledge support from the EU (BIODESIGN FP7-NMP-2012-

262948 and ERC Advanced MechAGE ERC-2016-ADG-741883). E. Wehrle received funding

from the ETH Postdoctoral Fellowship Program (MSCA-COFUND, FEL-25_15-1).

Author Contributions Statement

The study was designed by E.W., G.A.K., S.H. and R.M.. The experiments were performed

by E.W., G.A.K. and A.C.S.. Data analyses were performed by E.W. and D.C.B.. The

manuscript was written by E.W. and reviewed and approved by all authors.

Data availability

All necessary data generated or analyzed during the present study are included in this

published article and its Supplementary Information files (preprint available on BioRxiv

(BIORXIV/2019/692343). Additional information related to this paper may be requested

from the authors.

Competing Interests

The authors declare no competing interests.

References

1   Taiani, J. T. *et al.* Embryonic stem cell therapy improves bone quality in a model of impaired
fracture healing in the mouse; tracked temporally using in vivo micro-CT. *Bone* **64**, 263-272
(2014).
2   Locher, R. J. *et al.* Traumatic brain injury and bone healing: radiographic and biomechanical
analyses of bone formation and stability in a combined murine trauma model. *J. Musculoskel.*
Neuron. **15**, 309-315 (2015).
3   Graef, F. *et al.* Impaired fracture healing with high non-union rates remains irreversible after
traumatic brain injury in leptin-deficient mice. *J. Musculoskel. Neuron.* **17**, 78-85 (2017).
4   Mehta, M., Checa, S., Lienau, J., Hutmacher, D. & Duda, G. N. In vivo tracking of segmental
bone defect healing reveals that callus patterning is related to early mechanical stimuli. *Eur.*
Cell Mater. **24**, 358-371 (2012).
5   Tourolle né Betts, D. C. *et al.* The association between mineralised tissue formation and the
mechanical local in vivo environment: Time-lapsed quantification of a mouse defect healing
model. Preprint at https://doi.org/10.1101/721365 (2019).
6   Hauser, M., Siegrist, M., Keller, I. & Hofstetter, W. Healing of fractures in osteoporotic bones
in mice treated with bisphosphonates - A transcriptome analysis. *Bone* **112**, 107-119 (2018).
7   Bao, Q. W. *et al.* An appropriate Wnt/beta-catenin expression level during the remodeling
phase is required for improved bone fracture healing in mice. *Sci. Rep.-Uk* **7**, doi:ARTN
269510.1038/s41598-017-02705-0 (2017).

8   Hegde, V., Jo, J. E., Andreopoulou, P. & Lane, J. M. Effect of osteoporosis medications on
fracture healing. *Osteoporosis Int* **27**, 861-871 (2016).
9   Schindeler, A., McDonald, M. M., Bokko, P. & Little, D. G. Bone remodeling during fracture
repair: The cellular picture. *Semin. Cell. Dev. Biol.* **19**, 459-466 (2008).
10   Boyd, S. K., Davison, P., Muller, R. & Gasser, J. A. Monitoring individual morphological
changes over time in ovariectomized rats by in vivo micro-computed tomography. *Bone* **39**,
854-862 (2006).
11   Klinck, R. J., Campbell, G. M. & Boyd, S. K. Radiation effects on bone architecture in mice and
rats resulting from in vivo micro-computed tomography scanning. *Med. Eng. Phys.* **30**, 888-
895 (2008).
12   Lambers, F. M., Kuhn, G., Schulte, F. A., Koch, K. & Muller, R. Longitudinal Assessment of In
Vivo Bone Dynamics in a Mouse Tail Model of Postmenopausal Osteoporosis. *Calcified Tissue*
Int. **90**, 108-119 (2012).
13   David, V. *et al.* Noninvasive in vivo monitoring of bone architecture alterations in hindlimb-
unloaded female rats using novel three-dimensional microcomputed tomography. *J. Bone*
Miner. Res. **18**, 1622-1631 (2003).
14   Lambers, F. M., Schulte, F. A., Kuhn, G., Webster, D. J. & Muller, R. Mouse tail vertebrae
adapt to cyclic mechanical loading by increasing bone formation rate and decreasing bone
resorption rate as shown by time-lapsed in vivo imaging of dynamic bone morphometry.
Bone **49**, 1340-1350 (2011).
15   Schulte, F. A., Lambers, F. M., Kuhn, G. & Muller, R. In vivo micro-computed tomography
allows direct three-dimensional quantification of both bone formation and bone resorption
parameters using time-lapsed imaging. *Bone* **48**, 433-442 (2011).
16   Kuhn, G., Weigt, C., Lambers, F. M., Schulte, F. A. & Muller, R. Bisphosphonate treatment
does not impair load adaptation in osteopenic mouse bones. *Bone* **50**, S156-S156 (2012).
17   Hohlbaum, K. *et al.* Severity classification of repeated isoflurane anesthesia in C57BL/6JRj
mice-Assessing the degree of distress. *Plos One* **12**, doi:ARTN
e017958810.1371/journal.pone.0179588 (2017).
18   Tremoleda, J. L. *et al.* Imaging technologies for preclinical models of bone and joint disorders.
Ejnmmi Res **1**, doi:Artn 1110.1186/2191-219x-1-11 (2011).
19   Laperre, K. *et al.* Development of micro-CT protocols for in vivo follow-up of mouse bone
architecture without major radiation side effects. *Bone* **49**, 613-622 (2011).
20   Hildebrandt, I. J., Su, H. & Weber, W. A. Anesthesia and other considerations for in vivo
imaging of small animals. *ILAR J* **49**, 17-26 (2008).
21   Mustafy, T., Benoit, A., Londono, I., Moldovan, F. & Villemure, I. Can repeated in vivo micro-
CT irradiation during adolescence alter bone microstructure, histomorphometry and
longitudinal growth in a rodent model ? *Plos One* **13**, doi:ARTN
e020732310.1371/journal.pone.027323 (2018).
22   Richardson, C. A. & Flecknell, P. A. Anaesthesia and post-operative analgesia following
experimental surgery in laboratory rodents: Are we making progress? *Atla-Altern. Lab Anim.*
33, 119-127 (2005).
23   Dudziak, M. E. *et al.* The effects of ionizing radiation on osteoblast-like cells in vitro. *Plast.*
Reconstr. Surg. **106**, 1049-1061 (2000).
24   Matsumura, S., Jikko, A., Hiranuma, H., Deguchi, A. & Fuchihata, H. Effect of X-ray irradiation
on proliferation and differentiation of osteoblast. *Calcif. Tissue Int.* **59**, 307-308 (1996).
25   Scheven, B. A. A., Burger, E. H., Kawilarangdehaas, E. W. M., Wassenaar, A. M. & Nijweide, P.
J. Effects of Ionizing Irradiation on Formation and Resorbing Activity of Osteoclasts In vitro.
Lab. Invest. **53**, 72-79 (1985).

26   Brouwers, J. E. M., Van Rietbergen, B. & Huiskes, R. No effects of in vivo micro-CT radiation
on structural parameters and bone marrow cells in proximal tibia of Wistar rats detected
after eight weekly scans. *J. Orthop. Res.* **25**, 1325-1332 (2007).

27   Willie, B. M. *et al.* Diminished response to in vivo mechanical loading in trabecular and not
cortical bone in adulthood of female C57Bl/6 mice coincides with a reduction in deformation
to load. *Bone* **55**, 335-346 (2013).

28   Jegoux, F., Malard, O., Goyenvalle, E., Aguado, E. & Daculsi, G. Radiation effects on bone
healing and reconstruction: interpretation of the literature. *Oral Surg. Oral Med.* **109**, 173-
184 (2010).

29   Donneys, A. *et al.* Deferoxamine Restores Callus Size, Mineralization, and Mechanical
Strength in Fracture Healing after Radiotherapy. *Plast. Reconstr. Surg.* **131**, 711e-719e
(2013).

30   Chen, M. *et al.* Low-Dose X-Ray Irradiation Promotes Osteoblast Proliferation, Differentiation
and Fracture Healing. *Plos One* **9**, doi:ARTN e10401610.1371/journal.pone.0104016 (2014).

31   Wu, Y. R. *et al.* External Beam Irradiation Preferentially Inhibits the Endochondral Pathway of
Fracture Healing: A Rat Model. *Clin. Orthop. Relat. R.* **476**, 2076-2090 (2018).

32   Tsitsilonis, S. *et al.* The effect of traumatic brain injury on bone healing: an experimental
study in a novel in vivo animal model. *Injury* **46**, 661-665 (2015).

33   Wang, A. Y. *et al.* Effect of cervus and cucumis peptides on osteoblast activity and fracture
healing in osteoporotic bone. *Evid. Based Compl. Alt.* **2014**, 958908 (2014).

34   O'Donovan, D. A. *et al.* Radiation-Induced Craniofacial Bone Growth Inhibition: Acute and
Long-Term Effects on Bone Histopathology with and without Cytoprotection. *Plast. Reconstr.*
Surg **129**, 636e-645e (2012).

35   Zhang, J. *et al.* Differences in responses to X-ray exposure between osteoclast and osteoblast
cells. *J. Radiat. Res.* **58**, 791-802 (2017).

36   Chandra, A. *et al.* Suppression of Sclerostin Alleviates Radiation-Induced Bone Loss by
Protecting Bone-Forming Cells and Their Progenitors Through Distinct Mechanisms. *Journal*
of Bone and Mineral Research **32**, 360-372 (2017).

37   Borgiani, E. *et al.* Age-Related Changes in the Mechanical Regulation of Bone Healing Are
Explained by Altered Cellular Mechanoresponse. *J. Bone Miner. Res.*, doi:10.1002/jbmr.3801
(2019).

38   Kruck, B. *et al.* Sclerostin Neutralizing Antibody Treatment Enhances Bone Formation but
Does Not Rescue Mechanically Induced Delayed Healing. *J. Bone Miner. Res.* **33**, 1686-1697
(2018).

39   Bundkirchen, K. *et al.* Hemorrhagic shock alters fracture callus composition and activates the
IL6 and RANKL/OPG pathway in mice. *J. Trauma Acute Care* **85**, 359-366 (2018).

40   Jilka, R. L. The Relevance of Mouse Models for Investigating Age-Related Bone Loss in
Humans. *J. Gerontol. a-Biol* **68**, 1209-1217 (2013).

41   Miyagawa, K. *et al.* A novel underuse model shows that inactivity but not ovariectomy
determines the deteriorated material properties and geometry of cortical bone in the tibia of
adult rats. *J. Bone Miner. Metab.* **29**, 422-436 (2011).

42   Lotinun, S. *et al.* Osteoclast-specific cathepsin K deletion stimulates S1P-dependent bone
formation. *J. Clin. Invest.* **123**, 666-681 (2013).

43   Birkhold, A. I. *et al.* The Periosteal Bone Surface is Less Mechano-Responsive than the
Endocortical. *Sci. Rep.-Uk* **6**, doi:ARTN 2348010.1038/srep23480 (2016).

44   Zwingenberger, S. *et al.* Establishment of a femoral critical-size bone defect model in
immunodeficient mice. *J. Surg. Res.* **181**, e7-e14 (2013).

45   Russel, W. M. S. & Burch, R. L. *The Principles of Humane Experimental Technique* (1959).

46   Tannenbaum, J. & Bennett, B. T. Russell and Burch's 3Rs Then and Now: The Need for Clarity
in Definition and Purpose. *J. Am. Assoc. Lab. Anim.* **54**, 120-132 (2015).

47 Stevens, W. C. *et al.* Comparative Toxicities of Halothane, Isoflurane, and Diethyl-Ether at
Subanesthetic Concentrations in Laboratory-Animals. *Anesthesiology* **42**, 408-419 (1975).
48 Stollings, L. M. *et al.* Immune Modulation by Volatile Anesthetics. *Anesthesiology* **125**, 399-
411 (2016).
49 Kim, H. *et al.* Oxidative damages in the DNA, lipids, and proteins of rats exposed to
isofluranes and alcohols. *Toxicology* **220**, 169-178 (2006).
50 Kadar, B. *et al.* Effects of isoflurane exposure on oncogene and tumour suppressor gene
expressions in vital organs of CBA/CA mice. *In Vivo* **21**, 861-865 (2007).
51 Li, Z. H., Kuhn, G., Schirmer, M., Muller, R. & Ruffoni, D. Impaired bone formation in
ovariectomized mice reduces implant integration as indicated by longitudinal in vivo micro-
computed tomography. *Plos One* **12**, doi:ARTN e018483510.1371/journal.pone.0184835
(2017).
52 Lambers, F. M. *et al.* Bone adaptation to cyclic loading in murine caudal vertebrae is
maintained with age and directly correlated to the local micromechanical environment. *J.*
Biomech. **48**, 1179-1187 (2015).
53 Chandra, A. *et al.* PTH prevents the adverse effects of focal radiation on bone architecture in
young rats. *Bone* **55**, 449-457 (2013).
54 Chandra, A. *et al.* PTH1-34 alleviates radiotherapy-induced local bone loss by improving
osteoblast and osteocyte survival. *Bone* **67**, 33-40 (2014).
55 Freeman, T. A., Patel, P., Parvizi, J., Antoci, V. & Shapiro, I. M. Micro-CT Analysis with Multiple
Thresholds Allows Detection of Bone Formation and Resorption during Ultrasound-Treated
Fracture Healing. *J. Orthop. Res.* **27**, 673-679 (2009).
56 Taylor, C., Scheuren, A., Trüssel, A. & R., M. 3D local in vivo environment (LivE) imaging for
single cell protein analysis of bone tissue. *Curr. Dir. Biomed. Eng.* **2**, 449-453 (2016).
57 Hofman, M. *et al.* Effect of neurokinin-1-receptor blockage on fracture healing in rats. *Sci.*
Rep.-Uk **9**, doi:ARTN 974410.1038/s41598-019-46278-6 (2019).
58 Nevalainen, T. Animal Husbandry and Experimental Design. *Ilar Journal* **55**, 392-398 (2014).
59 Beery, A. K. Inclusion of females does not increase variability in rodent research studies. *Curr.*
Opin. Behav. Sci. **23**, 143-149 (2018).
60 O'Neill, K. R. *et al.* Micro-computed tomography assessment of the progression of fracture
healing in mice. *Bone* **50**, 1357-1367 (2012).
61 Morgan, E. F. *et al.* Micro-computed tomography assessment of fracture healing:
Relationships among callus structure, composition, and mechanical function. *Bone* **44**, 335-
344 (2009).
62 Wehrle, E. *et al.* Distinct Frequency Dependent Effects of Whole-Body Vibration on Non-
Fractured Bone and Fracture Healing in Mice. *J. Orthop. Res.* **32**, 1006-1013 (2014).
63 Wehrle, E. *et al.* The impact of low-magnitude high-frequency vibration on fracture healing is
profoundly influenced by the oestrogen status in mice. *Dis. Model. Mech.* **8**, 93-104 (2015).
64 Mulder, L., Koolstra, J. H. & Van Euden, T. M. G. J. Accuracy of MicroCT in the quantitative
determination of the degree and distribution of mineralization in developing bone. *Acta*
Radiol. **45**, 769-777 (2004).
65 Meganck, J. A., Kozloff, K. M., Thornton, M. M., Broski, S. M. & Goldstein, S. A. Beam
hardening artifacts in micro-computed tomography scanning can be reduced by X-ray beam
filtration and the resulting images can be used to accurately measure BMD. *Bone* **45**, 1104-
1116 (2009).
66 Lukas, C. *et al.* Mineralization kinetics in murine trabecular bone quantified by time-lapsed in
vivo micro-computed tomography. *Bone* **56**, 55-60 (2013).
67 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods*
9, 676-682, doi:10.1038/Nmeth.2019 (2012).

 Figure Legends

 Figure 1. Volumes of interest (VOIs) for micro-CT evaluation of callus and adjacent bone:

 Defect center (DC - red), defect periphery (DP - yellow), fragment center (FC - blue),

 fragment periphery (FP - orange).

 Figure 2. Representative images (full image and cut; threshold: 645 mg HA/cm^3) of the

 defect region from animals of the scan group (week 0-6): union defect (top panel), non-

 union defect (bottom panel). Visualization of bone formation (orange) and resorption (blue)

 via registration of micro-CT scans from week 5 and 6.

 Figure 3. Micro-CT based evaluation of bone parameters in the scan group using different

 VOIS: total VOI (TOT) - defect and adjacent bone fragments (top), defect VOI (DC+DP) –

 defect center and periphery (bottom). a+d. Bone formation rate (solid) and bone resorption

 rate (dashed line) in the femur defect (TV) given in percent per day. b+e: Bone volume (BV)

 normalized to TV (DC+FC for TOT and DC for DC+DP). c+f: Degree of bone mineralization

 given as ratio of bone volume with a density ≥645 mg HA/cm^3 to the total osseous volume in

 the defect (threshold ≥395 mg HA/cm^3). n=10; * indicates $p < 0.05$ between consecutive

 weeks determined by Friedman test with Dunn correction for multiple comparisons (a-e)/

 repeated measurements ANOVA with Bonferroni correction (f).

 Figure 4. Micro-CT based evaluation of bone parameters in the scan (red) and control group

 (blue) using different VOIS: total VOI (TOT: DC+DP+FC+FP) - defect and adjacent bone

 fragments (top), defect VOI (DC+DP) – defect center and periphery (bottom). a+d. Formed

 (solid) and resorbed (empty) bone volume (BV) in the femur defect (TV) given in percent per

day. b+e: Bone volume (BV) normalized to TV (DC+FC for TOT and DC for DC+DP). c+f:

Degree of bone mineralization given as ratio of bone volume with a density ≥645 mg HA/cm^3

to the total osseous volume in the defect (threshold ≥395 mg HA/cm^3). Control group: n=8,

scan group: n=10. * indicates $p < 0.05$ determined by two-tailed Student's t-test (a-e; f: week

6)/Mann-Whitney U-test (f: week 5).

Figure 5. Representative longitudinal sections of fractured femora 6 weeks after defect

surgery of unions and non-unions in the scan and control group. Top panel: Safranin-O

staining - overview images (row 1), scale bar = 2 mm; area between inner pins of fixator (row

2), scale bar = 500 µm; bottom panel: Sclerostin staining – area centered between inner pins

of fixator (row 3), scale bar = 100 µm; Isotype control for Sclerostin staining (row 4), scale

bar = 100 µm.

Tables

Table 1. Micro-CT based evaluation of bone parameters in the scan group using different VOIS

		Post-operative week								Post-operative week							
		0-1	1-2	2-3	3-4	4-5	5-6	p-value #		0	1	2	3	4	5	6	p-value #
DC	BFR [%/day]	0.23 ± 0.10*	2.24 ± 1.20	2.40 ± 1.15	1.30 ± 0.49	0.63 ±0.14	0.41 ± 0.09	< 0.0001	BV/TV [%]	1 ± 0	3 ± 1	17 ± 9*	28 ± 15	30 ± 14	29 ± 12	28 ± 10	< 0.0001
	BRR [%/day]	0.06 ± 0.02	0.14 ± 0.04*	0.88 ± 0.36	1.06 ± 0.85	0.75 ± 0.43	0.52 ± 0.25	< 0.0001	BV_{645}/BV_{395} [%]	1 ± 1	34 ± 10	26 ± 6	49 ± 7	67 ± 6	76 ± 6	80 ± 6	< 0.0001
DP	BFR [%/day]	0.03 ± 0.02*	0.38 ± 0.16	1.07 ± 0.88	0.68 ± 0.52	0.28 ± 0.20	0.18 ± 0.11	< 0.0001	BV/TV [%]	0 ± 0	0 ± 0	3 ± 1	9 ± 7	11 ± 9	11 ± 9	11 ± 8	< 0.0001
	BRR [%/day]	0.01 ± 0.0	0.03 ± 0.02	0.19 ± 0.08	0.34 ± 0.31	0.30 ± 0.23	0.18 ± 0.16	< 0.0001	BV_{645}/BV_{395} [%]	2 ± 3	24 ± 11	8 ± 2	37 ± 9	61 ± 10	71 ± 7	74 ± 9	< 0.0001
FC	BFR [%/day]	0.22 ± 0.08	0.26 ± 0.15	0.31 ± 0.12	0.44 ± 0.19	0.45 ± 0.11	0.40 ± 0.12	0.0019	BV/TV [%]	57 ± 7	57 ± 6	53 + 6	48 ± 5	44 ± 5	41 ± 6	40 ± 7	< 0.0001
	BRR [%/day]	0.24 ± 0.17*	0.78 ± 0.31	1.13 ± 0.30	1.01 ± 0.30	0.80 ± 0.25	0.60 ± 0.21	< 0.0001	BV_{645}/BV_{395} [%]	95 ± 1	92 ± 1	89 ±2	87 ± 1	86 ± 2	86 ± 2	86 ± 3	< 0.0001
FP	BFR [%/day]	0.43 ± 0.09*	1.81 ± 0.58	1.77 ± 0.76	0.85 ± 0.41	0.51 ± 0.28	0.33 ± 0.20	< 0.0001	BV/TV [%]	5 ± 1	6 ± 1	16 ± 4	21 ± 7	20 ± 8	20 ± 9	20 ± 9	< 0.0001
	BRR [%/day]	0.20 ± 0.09	0.40 ± 0.13*	1.16 ± 0.30	0.90 ± 0.38	0.57 ± 0.22	0.35 ± 0.14	< 0.0001	BV_{645}/BV_{395} [%]	0 ± 0	34 ± 9	23 ± 7	52 ± 7	69 ± 5	75 ± 4	78 ± 3	< 0.0001

DC = defect center, DP = defect periphery, FC = cortical fragment center, FP = cortical fragment periphery; BFR – bone formation rate; BRR – bone resorption rate; BV/TV - bone
volume normalized to DC for DC, DP and FC for FC, FP; BV645/BV395 - degree of bone mineralization given as ratio of bone volume with a density ≥645 mg HA/cm^3 to the total
osseous volume (threshold: 395mg HA/cm^3); *, significant difference compared to subsequent week (p < 0.05); #, p-value over time as tested with repeated measurements
ANOVA with Bonferroni correction/Friedman test with Dunn correction for multiple comparisons, n=10

Table 2. CT evaluation of bone parameters in the different volumes of interest (VOI) for the
scan and control group

VOI	Parameter	Evaluation time point/period	Control group	Scan group
DC	BFR [%/day]	week 5-6	0.59 ± 0.24	0.41 ± 0.09
	BRR [%/day]	week 5-6	0.62 ± 0.28	0.52 ± 0.25
	BV/TV [%]	week 5	35 ± 11	29 ± 12
		week 6	35 ± 11	28 ± 10
	BV_{645}/BV_{395} [%]	week 5	76 ± 2	76 ± 6
		week 6	80 ± 3	80 ± 6
DP	BFR [%/day]	week 5-6	0.18 ± 0.12	0.18 ± 0.11
	BRR [%/day]	week 5-6	0.15 ± 0.11	0.18 ± 0.16
	BV/TV [%]	week 5	11 ± 8	11 ± 9
		week 6	11 ± 8	11 ± 8
	BV_{645}/BV_{395} [%]	week 5	71 ± 7	71 ± 7
		week 6	75 ± 6	74 ± 9
FC	BFR [%/day]	week 5-6	0.44 ± 0.19	0.40 ± 0.12
	BRR [%/day]	week 5-6	0.63 ± 0.19	0.60 ± 0.21
	BV/TV [%]	week 5	42 ± 6	41 ± 6
		week 6	41 ± 6	40 ± 7
	BV_{645}/BV_{395} [%]	week 5	84 ± 3	86 ± 2
		week 6	84 ± 3	86 ± 3
FP	BFR [%/day]	week 5-6	0.32 ± 0.09	0.33 ± 0.20
	BRR [%/day]	week 5-6	0.30 ± 0.09	0.35 ± 0.14
	BV/TV [%]	week 5	21 ± 8	20 ± 9
		week 6	21 ± 8	20 ± 9
	BV_{645}/BV_{395} [%]	week 5	76 ± 5	75 + 4
		week 6	79 ± 4	78 + 3

DC - defect center; DP - defect periphery; FC – fragment center; FP – fragment periphery; BFR – bone
formation rate; BRR – bone resorption rate; BV/TV - bone volume normalized to DC for DC, DP and FC
for FC, FP. BV_{645}/BV_{395} - degree of bone mineralization given as ratio of bone volume with a density
≥645 mg HA/cm^3 to the total osseous volume (threshold: 395mg HA/cm^3). Control/scan group -
n=8/10; no significant differences were detected between the scan group and the control group (p ≥
0.05), assessed by two-tailed Student's t-test or Mann-Whitney U-test.

Table 3. Number of bridged cortices per callus evaluated in two perpendicular planes and
number of mice with successful fracture healing (≥ 3 bridged cortices) evaluated 5 and 6
weeks after the defect surgery.

| | Number of bridged cortices | | | | | Fracture healing outcome[#] | |
Group	0	1	2	3	4	Not healed	Healed
Scan							
all animals	3	0	1	0	7	4 (36%)	7 (64%)
defects ≥ 1.5mm	3	0	1	0	2	4 (67%)	2 (33%)
Control							
all animals	2	0	0	0	6	2 (25%)	6 (75%)
defects ≥ 1.5mm	2	0	0	0	1	2 (67%)	1 (33%)

775 [#], percentages of healed/not healed fractures per group given in brackets

Figure 1

Figure 2

Scan group	week 0	week 0-1	week 1-2	week 2-3	week 3-4	week 4-5	week 5-6
Union							
Non-Union							

Figure 3

Figure 4

Figure 5

	Scan		Control	
	Union	**Non-Union**	**Union**	**Non-Union**
Safranin-O Femur overview				
Safranin-O Defect region				
Sclerostin Defect center				
Sclerostin Isotype control Defect center				