Evaluation of longitudinal time-lapsed *in vivo* micro-CT for monitoring fracture healing in

mouse femur defect models

Esther Wehrle[1], Duncan C Betts[1], Gisela A Kuhn[1], Ariane C Scheuren[1], Sandra Hofmann[1,2],

Ralph Müller[1*]

[1] Institute for Biomechanics, ETH Zurich, Zurich, Switzerland, [2] Department of Biomedical

Engineering and Institute for Complex Molecular Systems, Eindhoven University of

Technology, The Netherlands.

Corresponding author:

Ralph Müller, PhD

Institute for Biomechanics

ETH Zurich

Leopold-Ruzicka-Weg 4

8093 Zurich, Switzerland

Email: ram@ethz.ch

Abstract

Longitudinal *in vivo* micro-computed tomography (micro-CT) is of interest to non-invasively capture the healing process of individual animals in preclinical fracture healing studies. However, as adverse imaging effects associated with anesthesia, handling, and radiation have been reported in some cases, this study assessed imaging-associated effects on fracture healing in a mouse femur defect model. A scan group received weekly micro-CT measurements (week 0-6), whereas controls were only scanned post-operatively and at week 5 and 6. Registration of consecutive scans enabled assessment of bone turnover with distinct characteristics of the different healing phases. Weekly micro-CT application did not significantly change any of the assessed callus parameters in defect and periosteal volumes. This was supported by histology showing only small amounts of cartilage residuals in both groups, indicating progression towards the end of the healing period. Also, immunohistochemical staining of Sclerostin, previously associated with mediating adverse radiation effects on bone, did not reveal differences between groups.

The established longitudinal *in vivo* micro-CT-based approach allows monitoring of healing phases in mouse femur defect models without significant effects of anesthesia, handling and radiation on callus properties. Therefore, this study supports application of longitudinal *in vivo* micro-CT for healing-phase-specific monitoring of fracture repair in mice.

Introduction

Adequate monitoring and characterization of the healing process is important in preclinical fracture healing studies. One recent approach to non-invasively capture the formation and remodeling of the osseous fracture callus is the repeated application of *in vivo* micro-computed tomography (micro-CT) [1-3]. This approach allows three-dimensional assessment of callus structures and by registering consecutive scans, dynamic parameters such as bone formation and resorption can be evaluated [4,5]. As each animal can be followed individually throughout the healing process with low variance in the assessed parameters, animal numbers can be reduced compared to well-established cross-sectional studies with endpoint micro-CT and two-dimensional histological callus evaluation.

In non-fractured bone, longitudinal time-lapsed *in vivo* micro-CT has been increasingly used to monitor changes in bone properties associated with different diseases and external factors, e.g. estrogen-deficiency [6-8], mechanical-(un)loading [9-11], and drug application [12]. However, several studies indicate that anesthesia, cumulative radiation dosage and stress due to the required handling for the CT measurements may have effects on animal well-being and bone properties [7,13-16]. According to the EU Directive 2010/63, the severity of repeated isoflurane anesthesia can be categorized as mild, although repeated anesthesia was considered worse than a single session with sex-dependent differences in perceiving the severity of a procedure [13]. Specifically, female mice were shown to be more susceptible to anesthesia-induced effects on well-being compared to male mice. Nevertheless, in both sexes repeated isoflurane anesthesia caused only short-term mild distress and impairment of well-being, mainly in the immediate post-anesthetic period. Radiation has also been shown to have dosage-dependent effects on bone cells *in vivo* and *in vitro* (12): Whereas

high dose x-ray radiation (2.5-8Gy) was associated with reduced osteoblast and osteoclast

proliferation [9,17-19], lower doses (<2Gy) had a stimulatory effect on osteoclasts. Some studies

reported radiation-associated effects on structural bone parameters, whereas other studies

did not see any changes [7,20]. These findings indicate the importance of study-specific

adaptation of micro-CT protocols, and to protect animals scanning times and radiation

settings should be minimized.

Recently, several fracture healing studies have applied longitudinal *in vivo* micro-CT to

monitor callus formation [2,3,21-23]. However, as adverse effects of longitudinal imaging on the

development of bone properties have been reported in the literature in some cases during

normal bone remodeling, there is a need to also assess imaging-associated impact on the

callus formation during the highly metabolically active process of fracture healing.

Therefore, the objectives of this study were to establish an *in vivo* micro-CT based approach

for longitudinal monitoring of fracture healing in a mouse femur defect model and to assess

the combined effect of radiation, anesthesia and handling associated with weekly time-

lapsed micro-CT measurements on callus properties during the remodeling phase of fracture

healing.

<u>Results</u>

<u>General physical observation</u>

All mice recovered rapidly from surgery. During the healing period, in both groups, the

animals' body weight did not significantly change compared to pre-operative values without

significant differences between the two groups (see Supplementary Fig. S1 online). Social

interaction between mice and nesting behaviour did not differ from pre-surgical

observations and was similar for animals of the scan and control groups.

Volumes of interest (VOI) for evaluation by time-lapsed *in vivo* micro-CT

One animal from the scan group could not be included in the analysis due to failure in VOI

generation caused by too little cortical bone being present in the field of view. The four

different VOIs (depicted in Fig. 1) encompassed the following volume for the control (n=8)

and scan group (n=10): $2.51 \pm 0.34 mm^3$ vs. $2.60 \pm 0.34 mm^3$ for the defect center (DC),

$17.26 \pm 2.21 mm^3$ vs. $18.73 \pm 4.64 mm^3$ for the defect periphery (DP), $1.86 \pm 0.48 mm^3$ vs.

$1.73 \pm 0.45 mm^3$ for the cortical fragments (FC), $14.47 \pm 3.30 mm^3$ vs. $13.10 \pm 2.66 mm^3$ for the

fragment periphery (FP). The total volume (TOT) between the inner pins of the fixator was

$36.10 \pm 2.81 mm^3$ for the control and $36.15 \pm 3.74 mm^3$ for the scan group. No significant

differences in volume were detected in any of the VOIs between groups.

Longitudinal monitoring of fracture healing by time-lapsed *in vivo* micro-CT

In the scan group (n=10), the repeated micro-CT scans (1x/week, Fig. 2) covered the period

from the day of the defect surgery (d0) until post-operative week 6 with distinct callus

characteristics indicative of the different healing phases (inflammation, repair, remodeling).

From week 0-1 to week 1-2 a significant 5.7x increase in bone formation was detected (Fig.

3A) in the total VOI (TOT=DC+DF+FC+FP, p<0.0001; Fig. 1), indicating progression from the

inflammation to the reparative phase. This led to a significant gain in bone volume by week 2

(BV/TV$_{week2}$: 39±7% vs. BV/TV$_{week0}$: 25±3%, p=0.0134; Fig. 3b). As bone formation triggers the

onset of bone resorption, a significant 2.8x increase in resorptive activities was seen from

week 1-2 to week 2-3 (p=0.0020), indicating the progression to the remodeling phase. Two

weeks after surgery the highly mineralized bone fraction in the TOT VOI was significantly

lower compared to the post-operative measurement (BV_{645}/BV_{395} in week 2: 59±5% vs.

BV_{645}/BV_{395} in week 0: 84±1%, *p*<0.0001), indicating formation of mineralized callus of low

density. From postoperative week 2 onwards, the highly mineralized bone fraction in the

TOT VOI gradually increased in all subsequent weeks of the healing period reaching

statistical significance by week 5 (BV_{645}/BV_{395} in week 2: 59±5% vs. BV_{645}/BV_{395} in week 5:

79±3%, p=0.0134; Fig. 3c).

In order to better capture the regions where bone is mainly formed and resorbed, we

evaluated the different VOIs separately (Table 1): In the early postoperative phase from

week 0-1 to week 1-2 a strong onset of bone formation was seen in the DC and FP sub-

volumes, leading to significant 11.8x and 3.4x gain in mineralized tissue from week 0 to week

2 for DC (p=0.0090) and FP (p=0.0091), respectively. This indicates that both intra-cortical as

well as periosteal callus formation takes place in this femur defect model. In both VOIs bone

formation triggered the initiation of bone resorption from week 2-3. In detail, there was a

significant 6.2x and 2.9x increase in bone resorption from week 1-2 to week 2-3 in the DC

(p=0.0040) and the FP VOI (p=0.0079), respectively. Compared to the DC and FP VOIs, the

initiation of bone formation was much less pronounced in the DP sub-volume with less

deposition of mineralized tissue leading to only little peripheral callus formation in this VOI

(week 2-6) and subsequently low bone resorption activities from week 2-3 to week 5-6. In all

three regions (DC, DP, FP) the fraction of highly mineralized tissue considerably increased

from week 2 to week 5 (+192%, +774%, +227%), indicating callus maturation. In this femur

defect model, callus formation and remodeling mainly took place in the defect region (DC)

with only little peripheral callus formation and remodeling (DP). Looking at the cortical

fragments (FC), a significant 3.2x increase in resorptive activities was detected in week 1-2

compared to week 0-1 (p=0.0421), whereas no significant weekly change in bone formation

activities was seen in this region throughout the assessed healing period. This resulted in a

significant 24% reduction in bone volume from week 0 to week 4 (p=0.0091). The fraction of

highly mineralized bone also gradually decreased in the remaining osseous tissue reaching

statistical significance by week 3 (BV_{645}/BV_{395} in week 3 reduced by 9% compared to week 0,

p=0.0027).

Influence of the longitudinal _in vivo_ micro-CT protocol on callus properties

To assess the combined effects of radiation, anesthesia and handling associated with weekly

micro-CT measurements (week 0 - week 6) on callus properties, callus parameters of the

scan group were compared to control animals that were scanned only directly post-

operatively (d0) and after 5 and 6 weeks (Fig. 4).

Post-operatively (d0), no significant group differences in bone volume (BV/TV) were found in

any of the VOIs: 1.82±0.39% (control) vs. 1.47±0.15% (scan) in DC, 0.16±0.12% (control) vs.

0.13±0.06% (scan) in DP, 56.82±2.31% (control) vs. 57.14±6.99% (scan) in FC, 4.66±0.30%

(control) vs. 4.74±0.76% (scan) in FP. Likewise, no group differences in bone volume were

seen in the total volume (TOT; 26.92±4.69% in controls and 24.90±2.88% in the scan group).

During the remodeling phase (post-operative weeks 5-6), no significant difference in bone turnover was seen in the total volume (TOT) between the inner pins of the fixator (BFR: 0.76±0.23% in controls, 0.64±0.15% in scan group; BRR: 0.82±0.25% in controls, 0.78±0.26% in scan group). Also, bone volume did not significantly change due to the applied micro-CT protocol with a similar fraction of highly mineralized tissue after 5 weeks (BV_{645}/BV_{395}: 79±3% in controls, 79±3% in scan group) and 6 weeks (BV_{645}/BV_{395}: 81±3% in controls, 82±4% in scan group).

Regarding the four sub-volumes (Table 2), the applied micro-CT protocol did not significantly affect bone formation and resorption activities in the callus VOIs (DC, DP and FP) from week 5-6 with similar bone volume observed in week 5 and 6 for controls and scanned animals. In both groups, bone volume remained stable from week 5 to week 6, whereas the density of the mineralized tissue increased during the same period (DC, DP, FP). Looking at the adjacent cortical fragments (FC) also no significant differences were detected in any of the assessed parameters between the control group and the scan group. Similarly to the callus VOIs, bone volume also remained stable in this region from week 5 to week 6. However, in contrast to the callus VOIs, the density of the mineralized tissue in the cortical VOI FC did not change from week 5 to week 6. In week 6, the FC VOI comprised 33% and 35% of the osseous tissue in the total VOI (TOT) for the control and scan group, respectively. In the DC VOI, 37% (control group) and 36% (scan group) of the total bone volume were seen. Less osseous tissue was detected in the two peripheral VOIs, 11% (control group) and 12% (scan group) in the DP and 19% (control group) and 17% (scan group) in the FP VOI.

As group comparisons were only performed from week 5-6, we particularly focused on the

defect VOIs (DC+DP) which are most important for evaluating later healing time points

during the remodeling phase of fracture healing. No significant differences in bone turnover,

bone volume and mineralization were seen between the scan and control group. In addition,

according to the standard clinical evaluation of X-rays, the number of bridged cortices per

callus was evaluated in two perpendicular planes and animals with ≥3 bridged cortices were

categorized as healed. By week 5, cortical bridging occurred in 75% of the control animals

and 64% of the scanned animals (Table 3). However, non-unions only occurred in defects of

≥1.5mm length. Taking into account only these defects (≥1.5mm length), both groups had a

non-union rate of 67% indicating no significant effect of the chosen micro-CT protocol on

clinical fracture healing outcome.

Histology

As shown by Safranin-O staining six weeks after osteotomy no cartilage residuals were

present in the defect region in the bridged defects indicating normal progression of the

healing process in the final remodeling stage. In the non-union defects, only small areas of

cartilage were seen in the defect region. No differences were seen by visual inspection

between animals from the scan and control group (Fig. 5).

As radiation-mediated decreased bone formation by osteoblasts has previously been

associated with the inhibition of osteoanabolic Wnt-signaling [24], we assessed Sclerostin

expression in the defect region by immunohistochemistry. Visual inspection did not reveal

any differences in staining intensity and amount of Sclerostin between the scan and control

group (Fig. 5).

Discussion

In this study, longitudinal *in vivo* micro-CT was applied for monitoring the process of fracture

healing in mice. In addition, the combined effect of radiation, anesthesia and handling

associated with the established imaging approach on callus parameters was assessed. Until

now, preclinical fracture healing studies predominantly have a cross-sectional study design

with end-point callus evaluation in sub-groups of animals at different time points during the

healing period. Only recently, several studies in murine diaphyseal femur defect models

stabilized by external fixation [2,3,21,22] or intramedullary nails [23], have applied longitudinal *in*

vivo micro-CT to monitor the healing progression in each animal over time. These studies

were able to consecutively assess callus parameters (e.g. callus volume and density) at

specific time points during the healing process for up to 12 weeks [2,3,21-23]. By registering

consecutive scans, we were now able to include dynamic parameters such as bone

formation and resorption in our micro-CT based monitoring approach for fracture healing.

This allows for characterization of the different healing phases seen by changes in formation

and resorption in the osseous callus volume.

Specifically, we saw that the initiation of bone formation (maximum in week 2-3), indicative

of the onset of the reparative phase, triggered bone resorption (maximum in week 3-4) with

maximum osseous callus volumes in week 3. From week 4 to week 6, bone formation and

resorption continuously decreased. Whereas these remodeling activities also led to a

decrease in callus volume from week 4 to week 6, the density of the mineralized tissue in the

callus increased during the same period indicating advanced callus maturation. These

observations were further supported by histology with no remaining cartilage in the defect

region of unions and only a small amount of cartilage residuals in some non-unions. The

findings are also in accordance with previous longitudinal studies of up to 4 weeks in mice

[2,3,22] and 8 weeks in rats [21] mainly focusing on the inflammation and reparative phase.

Similar to our study, the bone volume continuously increased during the 4/8 week healing

period, although no remodeling to the original bone geometry was seen due to the shorter

observation period. One study in ovariectomized rats with a 12-week post-operative

monitoring period detected maximum osseous callus volumes by week 6, which diminished

thereafter, whereas bone mineral density continuously increased until week 12, similar to

our findings. This indicates that during the reparative phase first low-density bone is formed,

which is then further mineralized during the remodeling phase of fracture healing. Looking

at the functional fracture healing outcome, complete cortical bridging was seen in 64% of

the animals, which is similar to other studies (60-62%) using similar femur defect models

with relatively stiff external fixation (fixator stiffness measured in our study 24N/mm) in

mice [2,25]. However, differences in defect sizes have to be considered: 0.7mm [2], 1.19±0.25

mm [25], 1.47±0.16mm (this study). In contrast to the other studies, we saw the manifestation

of non-unions only in defects ≥1.5mm. This might be partially due to the fact that in some

studies the exact defect length might have differed from the reported value (saw diameter

used for osteotomy) and that non-union formation was assessed irrespective of defect

length. Overall, all longitudinal fracture healing studies using *in vivo* micro-CT were able to

follow the healing progression in single animals, thereby reducing variance of outcome

parameters compared to cross-sectional studies, also allowing for reduction of animals per

group considering the 3R principles of animal welfare [26,27]. However, when applying

longitudinal *in vivo* micro-CT, the effect of repeated anesthesia, handling and radiation

associated with the scans on the study's main outcome parameters and the general well-

being of the animals have to be considered.

In this study, isoflurane was used as anesthetic agent for the *in vivo* micro-CT scans, which is

the most commonly used inhalation anesthetic in longitudinal imaging studies with a fast on-

and offset of anesthesia and low metabolism rate [13]. Despite its general use, isoflurane has

been associated with several adverse effects, such as hepatic degenerative changes [28],

immunomodulation [29], oxidative DNA damage [30] as well as alterations in expression profiles

of oncogene and tumor suppressor genes in the bone marrow [31], which might potentially

also affect fracture healing and the general well-being of the animals. A recent study by

Hohlbaum et al. (2017) [13], which assessed the impact of repeated isoflurane anesthesia (6

times for 45 min at an interval of 3–4 days) and the associated handling on the well-being of

adult C57BL/6JRj mice, categorized the degree of distress as mild according to the EU

Directive 2010/63 with only short-term impairment of well-being, mainly in the immediate

post-anesthetic period. Therefore, isoflurane-based inhalation anesthesia and the associated

handling of the animal is suggested for longitudinal studies with multiple anesthetic sessions

per animal.

In this study, the applied micro-CT settings (55 kVp, 145 µA, 350 ms integration time, 500

projections per 180°, 21 mm field of view (FOV), scan duration ca. 15 min) were adapted

from well-established protocols used for longitudinal monitoring of bone adaptation and

implant integration in tail vertebrae in mice [10,32]. These settings are similar to protocols used

in 2 of the 5 longitudinal fracture healing studies published so far [3,21], while the 3 other

studies did not specify the CT settings [2,22,23]. The radiation dosage (CT dose index, CTDI)

associated with each scan in the current study was previously estimated to be 0.67 Gy per

scan [32,33]. A radiation control experiment using similar settings did not show detrimental

effects of 5 weekly *in vivo* micro-CT scans on bone development in tail vertebrae of mice [33].

However, some studies in non-fractured bone indicate cumulative radiation-associated

effects on bone properties particularly when using high radiation doses (>2G; [15]). The

radiation effects also depend on animal age [15,34] and ovariectomy (OVX) [7], with younger and

ovariectomized animals being more susceptible to radiation. So far, only one study assessed

radiation-associated effects of longitudinal *in vivo* micro-CT on bone healing in a uni-cortical

tibia burr-hole defect model, and did not find significant radiation effects on bone properties

[1]. In this study, similar CT settings (45kVp, 133 µA, 200 ms integration time, 1000 projections

per 180°) were used; however, due to the smaller dimensions of the defect and resulting

shorter scanning time, the radiation dosage (0.36 Gy) was lower compared to our protocol

(0.64 Gy).

In the current study, we assessed the combined imaging-associated impact (anesthesia,

handling, radiation; 1x/week for 6 weeks) on callus formation and remodeling in a

diaphyseal femur defect model in adult female C57BL/6J mice. We did not see any significant

imaging-associated changes in bone volume and turnover in the fracture callus during the

remodeling phase (post-operative week 5 and 6). Furthermore, no significant differences in

callus mineralization between scanned (d0, week1-6) and control animals (d0, week 5+6) in

any of the assessed callus volumes (TOT, DC, DP, FP) were observed. The distribution of the

osseous callus volume into the three sub-volumes was also not significantly different

between groups, indicating similar healing patterns. In addition, when separately assessing

the cortical fragments between the inner pins (FC), also no significant differences in any of

the assessed parameters were detected between groups. In respect to clinical healing

outcome, the percentage of unions in animals above the sub-critical defect length was 67%

in both groups. Defects without cortical bridging only showed small cartilage residuals as

visualized by Safranin-O staining, indicating manifestation of non-union formation. In the

defect regions of unions, no cartilage was seen, indicating progression towards the end of

the healing period. As adverse effects of radiation on bone properties have been shown to

be mediated via Sclerostin inhibition of osteoanabolic Wnt-signaling [24], we also visualized

Sclerostin expression in the fracture callus by immunohistochemistry, but did not see

differences in number of stained osteocytes, staining intensity and pattern between groups

in any of the assessed callus regions.

In summary, we did not see significant differences in a total of 7 assessed CT and histological

parameters in the total volume (TOT) between the inner pins of the fixator as well as in the

different callus sub-volumes (DC, DP, FP) and the adjacent cortical fragments (FC) between

the scan and the control group. The sub-volume specific analysis also allowed capturing of

distinct features only present in one region (e.g. significant post-operative increase in bone

resorption in FC), that showed contrary progression compared to the overall picture

(significant post-operative increase in bone formation in DC, DP and FP). This also shows the

importance of VOI selection. The other longitudinal fracture healing studies either assessed a

VOI consisting of the defect as well as adjacent bone fragments, or then only the callus in the

defect region including the last slice of intact cortex [2,3], or the callus in the defect and

periosteal region combined [21] was evaluated. An overall VOI including the defect region and

cortical fragments can give an overview of callus characteristics, but sub-volume-specific differences might be missed. Therefore, analysis of different endosteal and periosteal callus regions as well as the adjacent cortical volume is favorable.

The longitudinal *in vivo* micro-CT based approach established in this study allows monitoring of the different healing phases in mouse femur defect models without significant anesthesia-, handling- and radiation-associated effects on callus properties. By registering consecutive scans of the defect region for each animal, data on bone turnover can be obtained with distinct characteristics of the different fracture healing phases. Importantly, repeated anesthesia, handling and radiation associated with the scans did not impair callus formation and remodeling. Therefore, this study supports the application of longitudinal *in vivo* micro-CT for healing-phase specific monitoring of fracture repair in mice. Further studies should evaluate the potential of this micro-CT-based monitoring approach for healing phase-specific discrimination of normal and impaired healing conditions.

Methods

Study design

A micro-CT based approach for longitudinal *in vivo* monitoring of fracture healing was established for a mouse femur defect model. All mice received a femur defect and post-operative micro-CT scans (vivaCT 40, ScancoMedical, Brüttisellen, Switzerland) were performed. The scan group then received weekly scans of the defect area (weeks 1-6). To assess the combined effect of radiation, anesthesia and handling associated with weekly micro-CT measurements, controls were only scanned after 5 weeks. Control animals then

received another scan in post-operative week 6 to enable the assessment of both, static and

dynamic bone parameters during the final remodeling phase of fracture healing.

Animals

All animal procedures were approved by the authorities (licence number: 36/2014;

Kantonales Veterinäramt Zürich, Zurich, Switzerland). Female 12 week-old C57BL/6J mice

were purchased from Janvier (Saint Berthevin Cedex, France) and housed in the animal

facility of the ETH Phenomics Center (EPIC; 12h:12h light-dark cycle, maintenance feed

(3437, KLIBA NAFAG, Kaiseraugst, Switzerland), 5 animals/cage). At an age of 20 weeks, all

animals received a femur defect by performing an osteotomy (group 1: control group, defect

length - 1.45mm ± 0.16, n=8; group 2: scan group, defect length -1.47mm ± 0.16, n=11;

housing after surgery: 2-3 animals/cage). Perioperative analgesia (25 mg/L, Tramal®,

Gruenenthal GmbH, Aachen, Germany) was provided via the drinking water two days before

surgery until the third postoperative day. For surgery and micro-CT scans, animals were

anesthetized with isoflurane (induction/maintenance: 5%/1-2% isoflurane/oxygen).

Femur osteotomy

An external fixator (Mouse ExFix, RISystem, Davos, Switzerland; stiffness: 24N/mm [35]) was

positioned at the craniolateral aspect of the right femur and attached using four mounting

pins. First, the most distal pin was inserted approximately 2mm proximal to the growth

plate, followed by placement of the most proximal and the inner pins. Subsequently, a femur

defect was created using 2 Gigli wire saws.

 Time-lapsed *in vivo* micro-CT

Immediate post-surgery correct positioning of the fixator and the defect was visualized using

a vivaCT 40 (Scanco Medical AG, Brüttisellen, Switzerland) (isotropic nominal resolution: 10.5

µm; 2 stacks of 211 slices; 55 kVp, 145 µA, 350 ms integration time, 500 projections per

180°, 21 mm field of view (FOV), scan duration ca. 15 min). Subsequently, the fracture callus

and the adjacent bone between the inner pins of the fixator were scanned weekly using the

same settings. Scans were registered consecutively and morphometric indices (bone volume

- BV, bone volume/total volume – BV/TV, bone formation rate – BFR, bone resorption rate -

BRR) were computed (threshold: 395mg HA/cm^3; [4,11]). To assess mineralization progression,

a second threshold (645mg HA/cm^3) was applied and the ratio between highly and lowly

mineralized tissue (BV_{645}/BV_{395}) was calculated. According to the standard clinical evaluation

of X-rays, the number of bridged cortices per callus was evaluated in two perpendicular

planes (UCT Evaluation V6.5-1, Scanco Medical AG, Brüttisellen, Switzerland). A ''healed

fracture'' was considered as having a minimum of at least three bridged cortices per callus.

For evaluation, four volumes of interest (VOIs) were defined, which were created

automatically from the post-operative measurement (Fig. 1): defect center (DC), defect

periphery (DP), cortical fragment center (FC), and fragment periphery (FP). Data were

normalised to the central VOIs: DC/DC, DP/DC, FC/FC, FP/FC.

Histology

Histological analyses were performed in a sub-set of animals (n=6/group). On day 42, femora

were excised, the femoral head was removed and the samples were placed in 4% neutrally

buffered formalin for 24 hours and subsequently decalcified in 12.5% EDTA for 10-14 days.

The samples were embedded in paraffin and 4.5 µm longitudinal sections were stained with

Safranin-O/Fast Green: Weigert's iron haematoxylin solution (HT1079, Sigma-Aldrich, St.

Louis, MO) - 4min, 1:10 HCl-acidified 70% ethanol - 10s, tap water - 5min, 0.02% Fast Green

(F7258, Sigma-Aldrich, St. Louis, MO) - 3min, 1% acetic acid - 10s, 0.1% Safranin-O (84120,

Fluka, St. Louis, MO) - 5min. Images were taken with Slide Scanner Pannoramic 250 (3D

Histech, Budapest, Hungary) at 20x magnification.

For immunohistochemical staining of Sclerostin, nonspecific sites were blocked (1% BSA/PBS

+ 1% rabbit serum) for 60 min at room temperature. Subsequently, the sections were

incubated with the primary antibody against Sclerostin (AF1589, R&D Systems, Minneapolis,

MN; 1:150 in 1%BSA/PBS + 0.2% rabbit serum) overnight at 4°C. To detect the primary

antibody, a secondary biotinylated rabbit anti-goat-IgG antibody (BAF017, R&D Systems,

Minneapolis, MN) was added for 1 h at room temperature. For signal amplification, the

slides were incubated with avidin-biotin complex (PK-6100 Vector Laboratories, Burlingame,

CA) for 30 min. Diaminobenzidine (Metal Enhanced DAB Substrate Kit, 34065 ThermoFisher

Scientific, Waltham, MA) was used as detection substrate. Counterstaining was performed

with FastGreen (F7258, Sigma-Aldrich, St. Louis, MO). Species-specific IgG was used as

isotype control. Images were taken with Slide Scanner Pannoramic 250 (3D Histech,

Budapest, Hungary) at 40x magnification.

Statistics

CT analysis: Data were tested for normal distribution (Shapiro-Wilk-Test) and homogeneity

of variance (Levene-Test). Depending on the test outcome, group comparisons (scan versus

control group) of data derived at single time points were performed by two-tailed Student's

t-test or Mann-Whitney U-test (IBM SPSS Statistics Version 23). For statistical evaluation of

repeated measurements (scan group) dependent on results from normality and variance

tests, either one-way analysis of variance (ANOVA) with Bonferroni correction or Friedman

test with Dunn correction for multiple comparisons (GraphPad Prism 7) were performed.

Two-way ANOVA was used for longitudinal comparison of the body weight between the

control and the scan group. The level of significance was set at $p < 0.05$.

Acknowledgements

The authors gratefully acknowledge support from the EU (BIODESIGN FP7-NMP-2012-

262948 and ERC Advanced MechAGE ERC-2016-ADG-741883). E. Wehrle received funding

from the ETH Postdoctoral Fellowship Program (MSCA-COFUND, FEL-25_15-1).

Author Contributions Statement

The study was designed by E.W., G.A.K., S.H. and R.M.. The experiments were performed

by E.W., G.A.K. and A.C.S.. Data analyses were performed by E.W. and D.C.B.. The

manuscript was written by E.W. and reviewed and approved by all authors.

Data availability

All necessary data generated or analyzed during the present study are included in this

published article and its Supplementary Information files (preprint available on BioRxiv

(BIORXIV/2019/692343). Additional information related to this paper may be requested

from the authors.

Competing Interests

429 The authors declare no competing interests.

References

1   Taiani, J. T. *et al.* Embryonic stem cell therapy improves bone quality in a model of impaired
fracture healing in the mouse; tracked temporally using in vivo micro-CT. *Bone* **64**, 263-272,
doi:10.1016/j.bone.2014.04.019 (2014).
2   Locher, R. J. *et al.* Traumatic brain injury and bone healing: radiographic and biomechanical
analyses of bone formation and stability in a combined murine trauma model. *J. Musculoskel.*
Neuron. **15**, 309-315 (2015).
3   Graef, F. *et al.* Impaired fracture healing with high non-union rates remains irreversible after
traumatic brain injury in leptin-deficient mice. *J. Musculoskel. Neuron.* **17**, 78-85 (2017).
4   Betts, D., Wehrle, E., Kuhn, G. A., Hofmann, S. & Muller, R. Quantification of Mechanical
Stimuli and Bone Formation in Fracture Healing Using In Vivo Time-Lapsed Imaging. *Tissue*
Eng. Pt. A **21**, S56-S56 (2015).
5   Wehrle, E., Kuhn, G. A., Betts, D. C., Scheuren, A. & Muller, R. Influence of Longitudinal In
Vivo Micro-Computed Tomography on Fracture Healing in a Femoral Defect Model. *Tissue*
Eng. Pt. A **21**, S406-S406 (2015).
6   Boyd, S. K., Davison, P., Muller, R. & Gasser, J. A. Monitoring individual morphological
changes over time in ovariectomized rats by in vivo micro-computed tomography. *Bone* **39**,
854-862, doi:10.1016/j.bone.2006.04.017 (2006).
7   Klinck, R. J., Campbell, G. M. & Boyd, S. K. Radiation effects on bone architecture in mice and
rats resulting from in vivo micro-computed tomography scanning. *Med. Eng. Phys.* **30**, 888-
895, doi:10.1016/j.medengphy.2007.11.004 (2008).
8   Lambers, F. M., Kuhn, G., Schulte, F. A., Koch, K. & Muller, R. Longitudinal Assessment of In
Vivo Bone Dynamics in a Mouse Tail Model of Postmenopausal Osteoporosis. *Calcified Tissue*
Int. **90**, 108-119, doi:10.1007/s00223-011-9553-6 (2012).
9   David, V. *et al.* Noninvasive in vivo monitoring of bone architecture alterations in hindlimb-
unloaded female rats using novel three-dimensional microcomputed tomography. *J. Bone*
Miner. Res. **18**, 1622-1631, doi:10.1359/jbmr.2003.18.9.1622 (2003).
10   Lambers, F. M., Schulte, F. A., Kuhn, G., Webster, D. J. & Muller, R. Mouse tail vertebrae
adapt to cyclic mechanical loading by increasing bone formation rate and decreasing bone
resorption rate as shown by time-lapsed in vivo imaging of dynamic bone morphometry.
Bone **49**, 1340-1350, doi:10.1016/j.bone.2011.08.035 (2011).

11   Schulte, F. A., Lambers, F. M., Kuhn, G. & Muller, R. In vivo micro-computed tomography
allows direct three-dimensional quantification of both bone formation and bone resorption
parameters using time-lapsed imaging. *Bone* **48**, 433-442, doi:10.1016/j.bone.2010.10.007
(2011).
12   Kuhn, G., Weigt, C., Lambers, F. M., Schulte, F. A. & Muller, R. Bisphosphonate treatment
does not impair load adaptation in osteopenic mouse bones. *Bone* **50**, S156-S156,
doi:10.1016/j.bone.2012.02.487 (2012).
13   Hohlbaum, K. *et al.* Severity classification of repeated isoflurane anesthesia in C57BL/6JRj
mice-Assessing   the   degree   of   distress.   *Plos   One*   **12**,   doi:ARTN
e017958810.1371/journal.pone.0179588 (2017).
14   Tremoleda, J. L. *et al.* Imaging technologies for preclinical models of bone and joint disorders.
Ejnmmi Res. **1**, doi:Artn 1110.1186/2191-219x-1-11 (2011).
15   Laperre, K. *et al.* Development of micro-CT protocols for in vivo follow-up of mouse bone
architecture   without   major   radiation   side   effects.   *Bone*   **49**,   613-622,
doi:10.1016/j.bone.2011.06.031 (2011).
16   Hildebrandt, I. J., Su, H. & Weber, W. A. Anesthesia and other considerations for in vivo
imaging of small animals. *ILAR J.* **49**, 17-26 (2008).
17   Dudziak, M. E. *et al.* The effects of ionizing radiation on osteoblast-like cells in vitro. *Plast.*
Reconstr. Surg. **106**, 1049-1061 (2000).
18   Matsumura, S., Jikko, A., Hiranuma, H., Deguchi, A. & Fuchihata, H. Effect of X-ray irradiation
on proliferation and differentiation of osteoblast. *Calcified Tissue Int.* **59**, 307-308 (1996).
19   Scheven, B. A. A., Burger, E. H., Kawilarangdehaas, E. W. M., Wassenaar, A. M. & Nijweide, P.
J. Effects of Ionizing Irradiation on Formation and Resorbing Activity of Osteoclasts In vitro.
Lab. Invest. **53**, 72-79 (1985).
20   Brouwers, J. E. M., Van Rietbergen, B. & Huiskes, R. No effects of in vivo micro-CT radiation
on structural parameters and bone marrow cells in proximal tibia of Wistar rats detected
after eight weekly scans. *J. Orthop. Res.* **25**, 1325-1332, doi:10.1002/jor.20439 (2007).
21   Mehta, M., Checa, S., Lienau, J., Hutmacher, D. & Duda, G. N. In vivo tracking of segmental
bone defect healing reveals that callus patterning is related to early mechanical stimuli. *Eur.*
Cells Mater. **24**, 358-371; discussion 371 (2012).
22   Tsitsilonis, S. *et al.* The effect of traumatic brain injury on bone healing: an experimental
study in a novel in vivo animal model. *Injury* **46**, 661-665, doi:10.1016/j.injury.2015.01.044
(2015).
23   Wang, A. Y. *et al.* Effect of cervus and cucumis peptides on osteoblast activity and fracture
healing   in   osteoporotic   bone.   *Evid.   Based   Compl.   Alt.*   **2014**,   958908,
doi:10.1155/2014/958908 (2014).
24   Chandra, A. *et al.* Suppression of Sclerostin Alleviates Radiation-Induced Bone Loss by
Protecting Bone-Forming Cells and Their Progenitors Through Distinct Mechanisms. *J. Bone*
Miner. Res. **32**, 360-372, doi:10.1002/jbmr.2996 (2017).
25   Zwingenberger, S. *et al.* Establishment of a femoral critical-size bone defect model in
immunodeficient mice. *J. Surg. Res.* **181**, e7-e14, doi:10.1016/j.jss.2012.06.039 (2013).
26   Russel, W. M. S. & Burch, R. L. *The Principles of Humane Experimental Technique* (1959).
27   Tannenbaum, J. & Bennett, B. T. Russell and Burch's 3Rs Then and Now: The Need for Clarity
in Definition and Purpose. *J. Am. Assoc. Lab. Anim.* **54**, 120-132 (2015).
28   Stevens, W. C. *et al.* Comparative Toxicities of Halothane, Isoflurane, and Diethyl-Ether at
Subanesthetic Concentrations in Laboratory-Animals. *Anesthesiology* **42**, 408-419, doi:Doi
10.1097/00000542-197504000-00008 (1975).
29   Stollings, L. M. *et al.* Immune Modulation by Volatile Anesthetics. *Anesthesiology* **125**, 399-
411, doi:10.1097/Aln.0000000000001195 (2016).

30   Kim, H. *et al.* Oxidative damages in the DNA, lipids, and proteins of rats exposed to
isofluranes and alcohols. *Toxicology* **220**, 169-178, doi:10.1016/j.tox.2005.12.010 (2006).
31   Kadar, B. *et al.* Effects of isoflurane exposure on oncogene and tumour suppressor gene
expressions in vital organs of CBA/CA mice. *In Vivo* **21**, 861-865 (2007).
32   Li, Z. H., Kuhn, G., Schirmer, M., Muller, R. & Ruffoni, D. Impaired bone formation in
ovariectomized mice reduces implant integration as indicated by longitudinal in vivo micro-
computed tomography. *Plos One* **12**, doi:ARTN e018483510.1371/journal.pone.0184835
(2017).
33   Lambers, F. M. *et al.* Bone adaptation to cyclic loading in murine caudal vertebrae is
maintained with age and directly correlated to the local micromechanical environment. *J.*
Biomech. **48**, 1179-1187, doi:10.1016/j.jbiomech.2014.11.020 (2015).
34   Willie, B. M. *et al.* Diminished response to in vivo mechanical loading in trabecular and not
cortical bone in adulthood of female C57Bl/6 mice coincides with a reduction in deformation
to load. *Bone* **55**, 335-346, doi:10.1016/j.bone.2013.04.023 (2013).
35   Paul, G. R., Cherbuin, M., Wehrle, E. & Müller, R. Minimising external fixator stiffness
variation via computational analysis. *Annual Meeting, Swiss Soc. Biomed. Eng*, 45 (2017).

<u>Figure Legends</u>

Figure 1. Volumes of interest (VOIs) for micro-CT evaluation of callus and adjacent bone:

Defect center (DC - red), defect periphery (DP - yellow), fragment center (FC - blue),

fragment periphery (FP - orange).

Figure 2. Representative images (full image and cut; threshold: 645 mg HA/cm^3) of the

defect region from animals of the scan group (week 0-6): union defect (top panel), non-

union defect (bottom panel). Visualization of bone formation (orange) and resorption (blue)

via registration of micro-CT scans from week 5 and 6.

Figure 3. Micro-CT based evaluation of bone parameters in the scan group using different

VOIS: total VOI (TOT) - defect and adjacent bone fragments (top), defect VOI (DC+DP) –

defect center and periphery (bottom). a+d. Bone formation rate (solid) and bone resorption

rate (dashed line) in the femur defect (TV) given in percent per day. b+e: Bone volume (BV)

normalized to TV (DC+FC for TOT and DC for DC+DP). c+f: Degree of bone mineralization

given as ratio of bone volume with a density ≥645 mg HA/cm^3 to the total osseous volume in

the defect (threshold ≥395 mg HA/cm^3). n=10; * indicates $p < 0.05$ between consecutive

weeks determined by Friedman test with Dunn correction for multiple comparisons (a-e)/

repeated measurements ANOVA with Bonferroni correction (f).

Figure 4. Micro-CT based evaluation of bone parameters in the scan (red) and control group

(blue) using different VOIS: total VOI (TOT) - defect and adjacent bone fragments (top),

defect VOI (DC+DP) – defect center and periphery (bottom). a+d. Formed (solid) and

resorbed (empty) bone volume (BV) in the femur defect (TV) given in percent per day. b+e:

Bone volume (BV) normalized to TV (DC+FC for TOT and DC for DC+DP). c+f: Degree of bone

mineralization given as ratio of bone volume with a density ≥645 mg HA/cm^3 to the total

osseous volume in the defect (threshold ≥395 mg HA/cm^3). Control group: n=8, scan group:

n=10. * indicates $p < 0.05$ determined by two-tailed Student's t-test (a-e; f: week 6)/Mann-

Whitney U-test (f: week 5).

Figure 5. Representative longitudinal sections of fractured femora 6 weeks after defect

surgery of unions and non-unions in the scan and control group. Top panel: Safranin-O

staining - overview images, scale bar = 2 mm; mid panel: Safranin-O staining – area between

inner pins of fixator, scale bar = 500 µm; bottom panel: Sclerostin staining – area centered

between inner pins of fixator, scale bar = 100 µm.

Tables

Table 1. Micro-CT based evaluation of bone parameters in the scan group using different VOIS

		Post-operative week								Post-operative week							
		0-1	1-2	2-3	3-4	4-5	5-6	p-value #		0	1	2	3	4	5	6	p-value #
DC	BFR [%/day]	0.23 ± 0.10*	2.24 ± 1.20	2.40 ± 1.15	1.30 ± 0.49	0.63 ±0.14	0.41 ± 0.09	< 0.0001	BV/TV [%]	1 ± 0	3 ± 1	17 ± 9*	28 ± 15	30 ± 14	29 ± 12	28 ± 10	< 0.0001
	BRR [%/day]	0.06 ± 0.02	0.14 ± 0.04*	0.88 ± 0.36	1.06 ± 0.85	0.75 ± 0.43	0.52 ± 0.25	< 0.0001	BV_{645}/BV_{395} [%]	1 ± 1	34 ± 10	26 ± 6	49 ± 7	67 ± 6	76 ± 6	80 ± 6	< 0.0001
DP	BFR [%/day]	0.03 ± 0.02*	0.38 ± 0.16	1.07 ± 0.88	0.68 ± 0.52	0.28 ± 0.20	0.18 ± 0.11	< 0.0001	BV/TV [%]	0 ± 0	0 ± 0	3 ± 1	9 ± 7	11 ± 9	11 ± 9	11 ± 8	< 0.0001
	BRR [%/day]	0.01 ± 0.0	0.03 ± 0.02	0.19 ± 0.08	0.34 ± 0.31	0.30 ± 0.23	0.18 ± 0.16	< 0.0001	BV_{645}/BV_{395} [%]	2 ± 3	24 ± 11	8 ± 2	37 ± 9	61 ± 10	71 ± 7	74 ± 9	< 0.0001
FC	BFR [%/day]	0.22 ± 0.08	0.26 ± 0.15	0.31 ± 0.12	0.44 ± 0.19	0.45 ± 0.11	0.40 ± 0.12	0.0019	BV/TV [%]	57 ± 7	57 ± 6	53 + 6	48 ± 5	44 ± 5	41 ± 6	40 ± 7	< 0.0001
	BRR [%/day]	0.24 ± 0.17*	0.78 ± 0.31	1.13 ± 0.30	1.01 ± 0.30	0.80 ± 0.25	0.60 ± 0.21	< 0.0001	BV_{645}/BV_{395} [%]	95 ± 1	92 ± 1	89 ±2	87 ± 1	86 ± 2	86 ± 2	86 ± 3	< 0.0001
FP	BFR [%/day]	0.43 ± 0.09*	1.81 ± 0.58	1.77 ± 0.76	0.85 ± 0.41	0.51 ± 0.28	0.33 ± 0.20	< 0.0001	BV/TV [%]	5 ± 1	6 ± 1	16 ± 4	21 ± 7	20 ± 8	20 ± 9	20 ± 9	< 0.0001
	BRR [%/day]	0.20 ± 0.09	0.40 ± 0.13*	1.16 ± 0.30	0.90 ± 0.38	0.57 ± 0.22	0.35 ± 0.14	< 0.0001	BV_{645}/BV_{395} [%]	0 ± 0	34 ± 9	23 ± 7	52 ± 7	69 ± 5	75 ± 4	78 ± 3	< 0.0001

DC = defect center, DP = defect periphery, FC = cortical fragment center, FP = cortical fragment periphery; BFR – bone formation rate; BRR – bone resorption rate; BV/TV - bone volume normalized to DC for DC, DP and FC for FC, FP; BV645/BV395 - degree of bone mineralization given as ratio of bone volume with a density ≥645 mg HA/cm^3 to the total osseous volume (threshold: 395mg HA/cm^3); *, significant difference compared to subsequent week (p < 0.05); #, p-value over time as tested with repeated measurements ANOVA with Bonferroni correction/Friedman test with Dunn correction for multiple comparisons, n=10

VOI	Parameter	Evaluation time point/period	Control group	Scan group
DC	BFR [%/day]	week 5-6	0.59 ± 0.24	0.41 ± 0.09
	BRR [%/day]	week 5-6	0.62 ± 0.28	0.52 ± 0.25
	BV/TV [%]	week 5	35 ± 11	29 ± 12
		week 6	35 ± 11	28 ± 10
	BV_{645}/BV_{395} [%]	week 5	76 ± 2	76 ± 6
		week 6	80 ± 3	80 ± 6
DP	BFR [%/day]	week 5-6	0.18 ± 0.12	0.18 ± 0.11
	BRR [%/day]	week 5-6	0.15 ± 0.11	0.18 ± 0.16
	BV/TV [%]	week 5	11 ± 8	11 ± 9
		week 6	11 ± 8	11 ± 8
	BV_{645}/BV_{395} [%]	week 5	71 ± 7	71 ± 7
		week 6	75 ± 6	74 ± 9
FC	BFR [%/day]	week 5-6	0.44 ± 0.19	0.40 ± 0.12
	BRR [%/day]	week 5-6	0.63 ± 0.19	0.60 ± 0.21
	BV/TV [%]	week 5	42 ± 6	41 ± 6
		week 6	41 ± 6	40 ± 7
	BV_{645}/BV_{395} [%]	week 5	84 ± 3	86 ± 2
		week 6	84 ± 3	86 ± 3
FP	BFR [%/day]	week 5-6	0.32 ± 0.09	0.33 ± 0.20
	BRR [%/day]	week 5-6	0.30 ± 0.09	0.35 ± 0.14
	BV/TV [%]	week 5	21 ± 8	20 ± 9
		week 6	21 ± 8	20 ± 9
	BV_{645}/BV_{395} [%]	week 5	76 ± 5	75 + 4
		week 6	79 ± 4	78 + 3

DC - defect center; DP - defect periphery; FC – fragment center; FP – fragment periphery; BFR – bone
formation rate; BRR – bone resorption rate; BV/TV - bone volume normalized to DC for DC, DP and FC
for FC, FP. BV_{645}/BV_{395} - degree of bone mineralization given as ratio of bone volume with a density
≥645 mg HA/cm^3 to the total osseous volume (threshold: 395mg HA/cm^3). Control/scan group -
n=8/10; *, significant difference compared to control group, $p < 0.05$, assessed by two-tailed
Student's t-test or Mann-Whitney U-test.

Table 3. Number of bridged cortices per callus evaluated in two perpendicular planes and
number of mice with successful fracture healing (≥ 3 bridged cortices) evaluated 5 and 6
weeks after the defect surgery.

	Number of bridged cortices					Fracture healing outcome	
Group	0	1	2	3	4	Not healed	Healed
Scan							
all animals	3	0	1	0	7	4	7
defects ≥ 1.5mm	3	0	1	0	2	4	2
Control							
all animals	2	0	0	0	6	2	6
defects ≥ 1.5mm	2	0	0	0	1	2	1

Scan		Control	
Union	**Non-Union**	**Union**	**Non-Union**