Spatial arrangement drastically changes the neural representation of multiple visual stimuli that compete in more than one feature domain

Steven Wiesner, Ian W. Baumgart, Xin Huang

Department of Neuroscience, School of Medicine and Public Health
Physiology Graduate Training Program
McPherson Eye Research Institute
University of Wisconsin - Madison, WI 53705, U.S.A.

Correspondence should be addressed to:
Xin Huang, Department of Neuroscience, School of Medicine and Public Health, University of Wisconsin, Madison, WI 53705, USA. Email: *Xin.Huang@wisc.edu*

Abbreviated title: Neural representation of multiple competing stimuli

Number of Pages: 38

Number of Figures: 9

Number of Tables: 2

Word Counts: Abstract (250), Significance (120), Introduction (641), Discussion (1483)

Conflict of interest: The authors declare no competing financial interests.

Acknowledgment

   We thank Jianbo Xiao for assistance on electrophysiological recordings and data analysis and Bryce Arseneau for technical support. This work was supported by National Institutes of Health Grant R01EY022443, and in part by the Office of the Director, National Institutes of Health Grant P51OD011106 to the Wisconsin National Primate Research Center, University of Wisconsin-Madison. This research was conducted at a facility constructed with support from Research Facilities Improvement Program grant numbers RR15459-01 and RR020141-01.

ABSTRACT

Natural scenes often contain multiple objects and surfaces. However, how neurons in the
visual cortex represent multiple visual stimuli is not well understood. Previous studies have shown
that, when multiple stimuli compete in one feature domain, the evoked neuronal response is biased
toward the stimulus that has a stronger signal strength. Here we investigate how neurons in the
middle temporal (MT) cortex of macaques represent multiple stimuli that compete in more than
one feature domain. Visual stimuli were two random-dot patches moving in different directions.
One stimulus had low luminance contrast and moved with high coherence, whereas the other had
high contrast and moved with low coherence. We found that how MT neurons represent multiple
stimuli depended on the spatial arrangement of the stimuli. When two stimuli were overlapping,
MT responses were dominated by the stimulus component that had high contrast. When two
stimuli were spatially separated within the receptive fields, the contrast dominance was abolished.
We found the same results when using contrast to compete with motion speed. Our neural data and
computer simulations using a V1-MT model suggest that the contrast dominance found with
overlapping stimuli is due to normalization occurring at an input stage fed to MT, and MT neurons
cannot overturn this bias based on their own feature selectivity. The interaction between spatially
separated stimuli can largely be explained by normalization within MT. Our results revealed new
rules on stimulus competition and highlighted the impact of hierarchical processing on
representing multiple stimuli in the visual cortex.

SIGNIFICANCE STATEMENT

19
20
Previous studies have shown that the neural representation of multiple visual stimuli can
be accounted for by a divisive normalization model. By using multiple stimuli that compete in
more than one feature domain, we found that luminance contrast has a dominant effect in
determining competition between multiple stimuli when they were overlapping but not spatially
separated. Our results revealed that neuronal responses to multiple stimuli in a given cortical area
cannot be simply predicted by the population neural responses elicited in that area by the individual
stimulus components. To understand the neural representation of multiple stimuli, rather than
considering response normalization only within the area of interest, one must consider the
computations including normalization occurring along the hierarchical visual pathway.

Introduction

        In natural scenes, multiple visual stimuli are often present in a local spatial region. While it is generally well understood how neurons in the visual cortex encode a single stimulus, how neurons encode multiple visual stimuli within their receptive fields (RFs) remains to be elucidated. Because visual perception depends critically on the integration and segregation of multiple visual stimuli (Braddick, 1993), understanding the neural representation of multiple stimuli is of significant importance.

        The middle temporal (MT) cortex is an extrastriate brain area that is important for visual motion processing (Britten, 2003; Born and Bradley, 2005; Park and Tadin, 2018). Neurons in area MT receive feedforward inputs from direction-selective neurons in V1 (Movshon and Newsome, 1996) and have RFs about ten times larger in size than those of V1 neurons at the same eccentricities (Gattass and Gross, 1981; Albright and Desimone, 1987). Previous studies have shown that neuronal responses in area MT elicited by multiple moving stimuli follow a sub-linear summation of the responses elicited by the individual stimulus components (Snowden et al., 1991; Qian and Andersen, 1994; Recanzone et al., 1997; Ferera and Lisberger, 1997; Britten and Heuer, 1999; Heuer and Britten, 2002; McDonald et al., 2014), consistent with a model of divisive normalization (Simoncelli and Heeger, 1998; Britten and Heuer, 1999; Carandini and Heeger, 2011).

        Work in our laboratory has shown that the direction tuning curves of MT neurons to overlapping random-dot stimuli moving transparently in different directions can also be described as a weighted sum of the responses elicited by the individual stimulus components (Xiao et al., 2014; Xiao and Huang, 2015). When two stimulus components have different signal strengths in one feature domain, defined either by motion coherence or luminance contrast, MT neurons pool the stimulus component that has a stronger signal strength with greater weight (Xiao et al., 2014). The response bias in MT toward the stimulus component that has a stronger signal strength can be accounted for by a descriptive model of divisive normalization (Xiao et al., 2014), similar to the contrast normalization model used to describe neuronal responses in V1 (Carandini et al., 1997; Busse et al. 2009).

However, natural scenes contain multiple visual stimuli that often differ in more than one
feature domain. For example, one stimulus may have a stronger signal strength in feature A but a
weaker signal strength in feature B, whereas another stimulus may have a weaker signal strength
in feature A but a stronger signal strength in feature B. In this case, it is unclear which stimulus
has an overall stronger signal strength and, more generally, how visual stimuli with multiple
competing features interact within neurons' RFs.

One possibility is that, to neurons in a given brain area, the overall signal strength of a
visual stimulus is reflected in the evoked responses of a population of neurons in that area. Due to
divisive normalization within that area, a neuron may weigh a visual stimulus more strongly if the
population neural response elicited by that stimulus is greater than the population response elicited
by a competing stimulus. Alternatively, how neurons in a given brain area weigh multiple
competing stimuli may be the result of neural computations occurring in multiple stages along the
hierarchical visual pathway and may not be explained by simply considering the population neural
responses elicited by the individual stimulus components in the area of interest.

Here, we investigate the rule by with neurons in area MT encode multiple moving stimuli
that compete in more than one feature domain. We found that MT responses to multiple stimuli
changed drastically when the spatial arrangement of the visual stimuli was varied. Our results
revealed how visual stimuli that differ in multiple feature domains interact within neurons' RFs
and shed light on how the neuronal responses in a given cortical area are shaped by neural
processing along the hierarchical visual pathway.

84
85
86   **Materials and Methods**
87
Two male adult rhesus monkeys (*Macaca mulatta*) were used in the neurophysiological
experiments. Experimental protocols were approved by the Institutional Animal Care and Use
Committee of UW-Madison and conform to U.S. Department of Agriculture regulations and to the
National Institutes of Health guidelines for the care and use of laboratory animals. Procedures for
surgical preparation and electrophysiological recordings were routine and similar to those
described previously (Xiao et al., 2015). A head post and a recording cylinder were implanted

during sterile surgery with the animal under isoflurane anesthesia. For electrophysiological

recordings from neurons in area MT, we took a vertical approach and used tungsten electrodes (1-

3 MΩ, FHC). We identified area MT by its characteristically large portion of directionally selective

neurons, small RFs relative to those of neighboring medial superior temporal cortex (area MST),

its location at the posterior bank of the superior temporal sulcus, and visual topography of the RFs

(Gattass and Gross, 1981). Electrical signals were amplified and single units were identified with

a real-time template-matching system and an offline spike sorter (Plexon). Eye position was

monitored using a video-based eye tracker (EyeLink, SR Research) with a rate of 1000 Hz.

Visual stimuli and experimental procedure

Stimulus presentation and data acquisition were controlled by a real-time data acquisition

program "Maestro" (https://sites.google.com/a/srscicomp.com/maestro/home). Visual stimuli

were presented on a 25-inch CRT monitor at a viewing distance of 63 cm. Monitor resolution was

1024×768 pixels, with a refresh rate of 100 Hz. Stimuli were generated by a Linux workstation

using an OpenGL application that communicated with an experimental control computer. The

luminance of the video monitor was measured with a photometer (LS-110, Minolta) and was

gamma-corrected.

Visual stimuli were achromatic random-dot patches presented within a circular aperture

with a diameter of 3°. Individual dots were squares of 2 pixels extending 0.08° on each side, and

each random-dot patch had a dot density of 2.7 dots/deg^2. The dots had a luminance of either 79

or 22 cd/m^2, presented on a uniform background with a luminance of 10 cd/m^2, which gives rise

to a Michelson contrast of either 77.5% or 37.5%. Random dots in each patch moved within the

stationary aperture in a specified direction. The motion coherence of each random-dot patch was

set to either 100% or 60%. To generate a random-dot patch moving at N% of motion coherence

(after Newsome and Pare 1988; Britten et al. 1992), N% of the "signal" dots were selected to move

coherently, while the rest of the dots referred to as the "noise" dots were repositioned randomly

within the aperture. Random selections of the "signal" and "noise" dots occurred at each monitor

frame. Therefore, a given dot would switch back and forth between a signal dot and a noise dot.

The lifetime of each dot was as long as the motion duration.

In each experimental trial, the monkey maintained fixation within a $1° \times 1°$ electronic

window around a small fixation point. After a neuron was isolated, we first characterized its

direction selectivity by interleaving trials of a $30° \times 27°$ random-dot patch, moving in different

directions at a step of $45°$ and at a speed of $10°/s$. The direction selectivity and preferred direction

(PD) were determined on-line using MATLAB (MathWorks). We then characterized the speed

tuning of the neuron using a random-dot patch moving at different speeds (1, 2, 4, 8, 16, 32, or

$64°/s$) in the neuron's PD. Using a cubic spline, the preferred speed (PS) of the neuron was taken

as the speed that evoked the highest firing rate in the fitted speed tuning curve. Next, we used a

series of $5° \times 5°$ random-dot patches moving in the PD and at the PS of the neuron to map the

neuron's RF. The location of the patch was randomized and the screen was tiled in $5°$ steps. The

RF map was interpolated at $0.5°$ intervals, and the location giving rise to the highest firing rate

was taken as the center of the RF.

In the main experiments, the visual stimuli appeared after the monkey maintained fixation

for 200 ms. To separate the neuronal responses to the stimulus motion from those due to the

stimulus onset, the visual stimuli were first turned on and remained stationary for 200 ms before

they started to move for 500 ms. The visual stimuli were then turned off. The monkeys maintained

fixation for an additional 200 ms after the stimulus offset. In some stimulus trials, two random-dot

patches that moved in different directions, referred to as two stimulus components, were presented

simultaneously. The direction separation between two stimulus components was fixed at $90°$. We

varied the vector averaged (VA) direction of the bi-directional stimulus around $360°$ to

characterize the response tuning curve. The two stimulus components were either overlapping in

one of two locations (site a or b) within the RF, or they were spatially separated within the RF, one

centered at site a and the other at site b, with at least $1°$ gap between the borders of the two random-

dot patches (illustrated in Fig. 1). In other trials, only one stimulus component was presented at

either site a or site b and the direction was varied to characterize the tuning curve to the stimulus

component. For the majority of the experiments, the VA and component directions were varied in

a step of $15°$. In a small set of experiments, the directions were varied in a step of $30°$. The trials

presenting bi-directional stimuli and individual stimulus components were randomly interleaved.

In the first experiment, one random-dot patch, referred to as the "low contrast & high coherence" component, had a luminance contrast of 37.5% and a motion coherence of 100%. The other random-dot patch, referred to as the "high contrast & low coherence" component, had a luminance contrast of 77.5% and a motion coherence of 60%. Both stimulus components moved at the same speed, which was set at the neuron's PS if it was below 10°/s, or at 10°/s if the PS was at or greater than 10°/s. Note that when a random-dot patch moved at 60% coherence in a given direction, the visual stimulus was different from a situation where 60% of the dots always moved coherently and the rest of the 40% of dots always moved randomly. Because the random selection of signal and noise dots occurred at each monitor frame in our stimuli, a noise dot at one frame may turn into a signal dot in the next frame and move in the coherent direction. Perceptually, it is difficult to segregate the noise dots from the signal dots of the same stimulus component. The noise dots of the "high contrast & low coherence component" are not an independent entity and do not appear to interfere with the coherence of the "low contrast & high coherence" component perceptually.

In the second experiment, we set the motion coherence of both random-dot patches to 100% but used different speeds for the two stimulus components. One random-dot patch, referred to as the "low contrast & faster speed" component, had a luminance contrast of 37.5% and moved at 10°/s. The other random-dot patch, referred to as the "high contrast & slower speed" component, had a luminance contrast of 77.5% and moved at 2.5°/s.

Data analysis

Response firing rate was calculated during the period of 500-ms stimulus motion and averaged across repeated trials. We fitted the raw direction tuning curves for the bi-directional stimuli and the individual stimulus components using splines at a resolution of 1°. We then rotated the spline-fitted tuning curve to the bi-directional stimuli so that the VA direction of 0° was aligned with the PD of each neuron. In the first experiment, the responses of each neuron to the bi-directional stimuli and individual stimulus components were normalized by the maximum response to the "low contrast & high coherence" component. In the second experiment, the responses of each neuron were normalized by the maximum response to the faster speed

component. We averaged the rotated and normalized tuning curves across neurons to obtain

population-averaged tuning curves.

To quantify the relationship between the responses elicited by the bi-directional stimuli

and those elicited by the individual stimulus components, we fitted the direction tuning curves

using a summation plus nonlinear interaction (SNL) model (Eq. 1), which has been shown to

provide a better fit of MT responses elicited by bi-directional stimuli than a linear weighted

summation model (Xiao et al., 2014).

$$R_{pred}(\theta_1, \theta_2) = w_1 R_1(\theta_1) + w_2 R_2(\theta_2) + b R_1(\theta_1) R_2(\theta_2), \qquad (1)$$

where R_{pred} is the response to the bi-directional stimuli predicted by the model; θ_1 and θ_2 are the

two component directions; R_1 and R_2 are the measured component responses elicited by the two

stimulus components when presented alone; w_1 and w_2 are the response weights for R_1 and R_2,

respectively; and b is the coefficient of multiplicative interaction between the component

responses. To determine whether the response elicited by the bi-directional stimuli showed a

significant bias toward one of the two stimulus components, we compared the response weights

w_1 and w_2 using either a paired t-test or a Wilcoxon signed-rank test.

We also fitted the response tuning curves to the bi-directional stimuli using a few variants

of a divisive normalization model (Carandini and Heeger, 2011) (see Results). The model fits were

obtained using the constrained minimization tool 'fmincon' (MATLAB) to minimize the sum of

squared error.

To evaluate the goodness of fit of a model for the response tuning curve to the bi-directional

stimuli, we calculated the percentage of variance (PV) accounted for by the model as:

$$PV = 100 \times \left(1 - \frac{SSE}{SST}\right), \qquad (2)$$

where SSE is the sum of squared errors between the model fit and the neuronal data, and SST is

the sum of squared differences between the data and the mean of the data (Morgan et al., 2008).

    V1-MT Model

We adapted a computational model proposed by Simoncelli and Heeger (1998)

(http://www.cns.nyu.edu/~lcv/MTmodel/) to reconstruct our visual stimuli and to simulate the

neuronal response tuning to the bi-directional stimuli that were either overlapping or spatially

separated. The model contained several consecutive stages, which can be interpreted as V1 simple,

V1 complex, and MT (Simoncelli and Heeger, 1996; Rust et al., 2006). Based on the dimensions

of video monitor and viewing distance in our neurophysiological experiments, 1° of visual angle

corresponds to 21 pixels. The random-dot patch in our model simulations had a circular aperture

with a diameter of 63 pixels (i.e. 3°) and the same dot density as used in our experiments. Each

dot had a size of 2×2 pixels.

We set the RFs of model neurons by Gaussian convolutional filters (Table 1). We estimated

the size of the RF for each neuron type by summing the lengths of the incorporated filters. For the

spatially-separated stimuli, we set a blank gap between the two stimulus components as the RF

size of the V1 complex neuron, which is 1.2°, to ensure that no V1 neuron would be driven by

both stimulus components. We generated direction-selective neuron populations that

approximately tiled a sphere in the frequency domain. We tuned the contrast response functions

by adjusting C_{50} values for V1 and MT neurons. These C_{50} values were represented in the model

as σ^2 in the normalization equation (Eq. 3), which was applied to both V1 complex cell and MT

stages of the model (adapted from Simoncelli and Heeger 1998 and Rust et al., 2006).

240
$$R'_n(t) \;=\; \frac{\lfloor R_n(t) \rfloor}{K \sum_m \lfloor w_m \cdot R_m(t) \rfloor + \sigma^2} \;,\qquad\qquad (3)$$

where $R_n(t)$ represents the *nth* neuron's linear filter response; $R'_n(t)$ represents the normalized

response of either V1 complex cell or MT neuron; $\lfloor\;\rfloor$ denotes half-wave rectification; K represents

the strength of normalization, which was set as $1-\sigma^2$; m represents the *nth* neuron's normalization

pool; w represents the Gaussian spatial weighting profile of the normalization pool, with a standard

deviation of SD_{norm}. The model parameters for V1 and MT stages are defined in Table 1. We fitted

the model contrast response functions to neural data from V1 and MT as described in Sclar et al.

(1990). Similarly, we tuned coherence responses by varying the spatial scale of the normalization

pool (*m*), the weighting profile within the pool (*w*), and the size of the V1 linear RF. The MT

coherence response function was fitted to data replotted from Figure 1C in Britten and Newsome

(1998). We are not aware of published neural data on V1 coherence response function. So the

parameters for V1 model neurons were varied to simulate our MT responses to bi-directional

stimuli without a constraint on V1 coherence response function. The same model parameters were

used for the overlapping and spatially separated conditions.

Table 1. Model parameters for V1 and MT neurons

Model parameters	V1 stage	MT stage
RF size (pixels)	15 (simple cell) 25 (complex cell)	211
Standard Deviation (SD) of Gaussian RF profile (pixels)	0.73/2.19/6.57 (3 scales for simple cells) 7 (complex cell)	53
σ^2 (C_{50})	0.0016	0.000049
Size of normalization pool *m* (pixels)	38	153
SD_{norm} (pixels)	5	52
Baseline activity	0	0.1

258
259

We explored several variants of the model architecture. The model parameters were fitted

after each architectural manipulation. The following changes enabled the model to better capture

the trends of the stimulus competition found in our neural data. First, we used area-normalized

Gaussian functions to set the weights for the spatial pooling and local population normalization.

Second, multiple frequency scales for V1 simple cells were computed by tripling the standard

deviation of the underlying 3rd order derivative Gaussian, similar to the doubling suggested in

Simoncelli and Heeger (1998) – this change was made after spectral analysis of stimuli showed

that a wider range of scales was necessary to capture motion at lower coherence. Third, V1 afferent

weights were not adjusted to zero mean, allowing MT neurons to have variable proportions of

positive and negative inputs. Finally and importantly, rectification and static nonlinearity were

applied to the MT stage after spatial pooling and before normalization, which is physiologically

plausible and provides a better fit of our neural data.

272
273
274
275

Results

We asked the question of how neurons in extrastriate area MT represent multiple visual stimuli that compete in more than one feature domain. To address this question, we conducted neurophysiological experiments and computer simulations. We recorded from isolated single neurons in area MT of two macaque monkeys while they performed a fixation task. Visual stimuli were two random-dot patches moving simultaneously in different directions within the RFs. In the first experiment, we used luminance contrast and motion coherence as two competing features. One stimulus had high contrast but moved with low coherence, whereas the other stimulus had low contrast but moved with high coherence (see Methods). We manipulated the spatial arrangement of the visual stimuli to investigate the contributions of earlier visual areas and area MT in mediating the competition between multiple stimuli. In a second experiment, we used luminance contrast and motion speed as two competing features. We first present the results from the neurophysiological experiments and then computer simulations.

Neurophysiological experiments

We measured the direction tuning curves of MT neurons in response to two stimuli that had competing visual features and moved simultaneously in different directions. Our dataset includes recordings from 76 MT neurons, 43 from monkey G and 33 from monkey B. We set the angular separation between the motion directions of two individual stimuli, referred to as the stimulus components, at 90° and varied the VA direction of the stimuli. In the first experiment, one stimulus component had a low contrast of 37.5% and moved at a high motion coherence of 100%. The other component had a high contrast of 77.5% and moved at a low coherence of 60%. Figure 1 shows the direction tuning curves of two representative neurons. The red curve shows the neuronal response elicited when both stimulus components were present, as a function of the VA direction of the two stimulus components. The green and blue curves show the neuronal responses elicited by the individual stimulus components when presented alone. The tuning curves of the component responses are arranged such that, at each VA direction, the data points on the green and blue curves correspond to the responses elicited by the individual stimulus components of that VA direction (note the color-coded abscissas for the component directions in Fig. 1A2).

For the two example neurons, the peak response of the direction tuning curve to the "low

contrast & high coherence" component alone (shown in blue) was greater than that to the "high

contrast & low coherence" component (shown in green) (Fig. 1). This is expected since MT

neurons are sensitive to motion coherence within a large coherence range (Britten et al., 1993),

whereas their contrast response function saturates at a low luminance contrast (Sclar et al., 1990).

Consequently, the average of the response tuning curves to the two stimulus components (shown

in gray) was biased toward the "low contrast & high coherence" component. Surprisingly, we

found that when the two stimulus components were overlapping, the neuronal responses elicited

by the bi-directional stimuli were strongly biased toward the "high contrast & low coherence"

component (Fig. 1A). This response bias was robust and occurred when we placed the overlapping

stimuli at a different site within the RF (Fig. 1B).

Two overlapping visual stimuli could stimulate not only the RFs of single MT neurons but

also the RFs of single V1 neurons. The response bias toward the "high contrast & low coherence"

component may be caused by the neural processes within area MT, or alternatively inherited from

earlier visual areas such as V1. To determine the contribution of earlier visual areas to the response

bias, we placed two stimulus components at different locations within the RF of a given MT

neuron. The two stimulus components were separated by a gap of at least 1° (illustrated in Fig.

1C). With this spatial arrangement, the RF of a single V1 neuron could only be stimulated by one

of the two stimulus components, whereas the RF of an MT neuron could still be stimulated by both

components. We found that the response tuning to the bi-directional stimuli changed drastically

when stimulus components were spatially separated. MT responses elicited by the bi-directional

stimuli no longer showed a bias toward the "high contrast & low coherence" component, but

roughly followed a scaled average of the component responses (Fig. 1C).

Figure 2 shows the tuning curves averaged across 70 MT neurons. The population-

averaged response elicited by the "low contrast & high coherence" component moving in the PD

of each neuron, aligned to 0°, was significantly greater than that elicited by the "high contrast &

low coherence" component moving in the PD (one-tailed paired t-test, p = 4.1×10^{-7}). However,

when the two stimuli were overlapping, the population response elicited by the bi-directional

stimuli was almost completely biased toward the weaker "high contrast & low coherence"

component, regardless of the spatial location within the RF (Fig. 2A and 2B). The bias toward the

"high contrast & low coherence" component at a given VA direction was in a manner of "higher-

contrast-take-all". For example, at a VA direction of 45° where the "low contrast & high

coherence" component moved in the PD (0°) and the "high contrast & low coherence" component

moved in 90° (indicated by a dotted line in Fig. 2A), the bi-directional response closely followed

the much weaker response elicited by the "high contrast & low coherence" component. When the

two stimulus components were spatially separated within the RF, the strong bias toward the "high

contrast & low coherence" component was abolished (Fig. 2C). The population response to the

bi-directional stimuli now showed roughly equal weighting of the responses elicited by the

individual stimulus components.

The SNL model (see Eq. 1 in Methods) provided an excellent fit of the MT responses

elicited by the bi-directional stimuli, illustrated by the black curves in Figure 1. Across our neuron

population the model fit accounted for, on average, 83% of the response variance (see Methods).

Figure 3 compares the response weights for the two stimulus components obtained from the SNL

model fits. In the overlapping condition, the mean response weight w_2 for the "high contrast & low

coherence" component was significantly greater than the weight w_1 for the "low contrast & high

coherence" component (one-tailed paired t-test, p = 1.9×10^{-45} for *site a*, p = 2.5×10^{-28} for *site b*)

(Fig. 3A). Nearly all data points, each representing the result from one neuron, were below the

unity line. The mean response weight for the "high contrast & low coherence" component was

0.97 (std = 0.24), whereas the mean weight for the "low contrast & high coherence" component

was 0.23 (std = 0.25), indicating a dominant effect of the "high contrast & low coherence"

component in determining the neuronal response to the bi-directional stimuli.

When the two stimulus components were spatially separated within the RF, the response

weights changed significantly, becoming symmetrically distributed relative to the unity line (Fig.

3B). The spread of weights in the spatially-separated condition is larger than that in the overlapping

condition. The mean weight for the "high contrast & low coherence" component decreased to 0.66

(std = 0.32), whereas the mean weight for the "low contrast & high coherence" component

increased to 0.68 (std = 0.43). The mean weights for the two components were no longer different

(paired t-test, p = 0.8) but were significantly greater than 0.5 of response averaging (t-test, p <

0.001).

To quantify the response bias toward an individual stimulus component, we calculated a

bias index (BI) :

375                         $BI = (w_2 - w_1)/(w_2 + w_1)$ (4)

A positive value of the index indicates a bias toward the "high contrast & low coherence"

component. Figure 3C shows how this bias index changes with the spatial arrangement of the

visual stimuli. In the overlapping condition, the mean BI is 0.73 (std = 0.23), which is significantly

greater than 0 (one-tailed t-test, p = 7.5×10^{-35}). In the spatially-separated condition, the mean BI is

-0.01 (std = 0.95), which is not significantly different from 0 (p = 0.7). The mean BI obtained in

the overlapping condition is significantly greater than that in the spatially-separated condition

(one-tailed paired t-test, p = 4.7×10^{-9}), indicating a change of the response bias when the spatial

arrangement of the visual stimuli is altered.

We previously found that the tuning curves of some MT neurons to overlapping bi-

directional stimuli can show a directional "side-bias" toward one of the two direction components

(Xiao and Huang, 2015). A subgroup of neurons prefers the stimulus component at the clockwise

side of two motion directions, whereas another group prefers the component direction at the

counter-clockwise side. These response biases can occur even when both stimulus components

have the same contrast and coherence. In the experiment shown in Figures 1-3, the "high contrast

& low coherence" component always moved at the counter-clockwise side direction (Fig. 2A, 2B).

Could the strong bias toward the "high contrast & low coherence" component in the overlapping

condition be due to a biased neuron sample that happened to have a strong bias toward the direction

component at the counter-clockwise side? To address this concern, we arranged the direction

components differently.

Figure 4A and B show the averaged direction tuning curves of 15 MT neurons when the

direction of the "high contrast & low coherence" component was placed at the counter-clockwise

side under the overlapping and spatially separated conditions, as in Figure 2. When the "high

contrast & low coherence" component was placed at the clockwise side of the two component

directions, the responses of the same 15 neurons to the bi-directional stimuli still showed a strong

bias toward the "high contrast & low coherence" component under the overlapping condition (Fig.

4C), and showed roughly equal weighting of the two components under the spatially-separated

condition (Fig. 4D). Placing the "high contrast & low coherence" component at the clockwise or

counter-clockwise side of the two component directions had no effect on the response bias, as

measured by the bias index under the overlapping and spatially-separated conditions (Wilcoxon

rank-sum test, $p = 0.6$).

To shed light on the neural mechanisms underlying the response bias, we examined the

timecourse of the neuronal responses in the overlapping and spatially separated conditions. Figure

5 shows the PSTHs calculated using a 10-ms time bin when either the "high contrast & low

coherence" component or the "low contrast & high coherence" component moved in the PD.

When stimuli were overlapping, as soon as MT neurons started to respond to the onset of the static

stimuli (see Methods), the response elicited by both stimulus components already closely followed

the "high contrast & low coherence" component, even before the onset of the stimulus motion

(Fig. 5A, B). After the onset of the motion response, the neuronal response to the bi-directional

stimuli continued to follow the response elicited by the "high contrast & low coherence"

component throughout the motion period, regardless of whether the component moved in the PD

and elicited a strong response (Fig. 5A), or 90° away from the PD and elicited a weak response

(Fig. 5B). Since the strong bias towards the "high contrast & low coherence" component in the

overlapping condition occurred at the very beginning of the stimulus onset, it is unlikely that the

bias was due to selective attention (see Discussion).

When stimuli were spatially separated, MT neurons also followed the "high contrast & low

coherence" component in response to the onset of the static stimuli (Fig. 5C, D). After the motion

onset, when the "high contrast & low coherence" component moved in the PD, the motion response

elicited by the bi-directional stimuli initially followed the "high contrast & low coherence"

component for ~30 ms, and was then "pulled down" by the non-PD component (see the arrow in

Fig. 5C). When the "high contrast & low coherence" component moved in the non-PD, the motion

response elicited by the bi-directional stimuli followed the "high contrast & low coherence"
component for ~10 ms after the onset of the motion response to the PD component, and was then
"pulled up" by the PD component (see the arrow in Fig. 5D). These results suggest that response
normalization under the spatially separated condition takes 10~30 ms to occur.

When two stimulus components overlap, the random dots from each component constitute
only half of the total number of dots of the two moving surfaces. Could the strong response bias
toward the "high contrast & low coherence" component be due to a reduction of the motion
coherence of the "low contrast & high coherence" component when the stimuli overlapped? We
think this is an unlikely explanation because overlapping reduces the percentage of the signal dots
relative to the total number of dots for both stimulus components. In addition, our stimuli moved
in two directions separated by 90º. Human observers can reliably segregate the two stimulus
components at this angle separation and the "low contrast & high coherence" component still
appears to move coherently. Overlapping does not change the relative coherence levels nor the
perceived coherence of the two stimulus components. When overlapping random-dot stimuli have
the same luminance contrast but move at different motion coherences, macaque MT response to
both stimulus components is biased toward the high coherence component (Xiao et al., 2014),
indicating that stimulus overlapping does not prevent the response bias toward the high coherence
component given equal contrast.

To determine whether the dominance by the high-contrast component on MT responses
elicited by overlapping stimuli occurs only when luminance contrast and motion coherence
compete with each other, we conducted a second experiment using visual stimuli that differ in
luminance contrast and motion speed. We previously found that when two overlapping random-
dot patches moved in the same direction at different speeds, within a range of low to intermediate
speeds, the responses of MT neurons elicited by the bi-speed stimuli was biased toward the faster
speed component (X. Huang et al., unpublished data). Motivated by this finding, we used motion
speed to compete with luminance contrast. As in the main experiment, the visual stimuli contained
two random-dot patches moving in two directions separated by 90° and we varied the VA direction
to measure the direction tuning curves. One stimulus component had a high luminance contrast of
77.5% and moved at a slower speed of 2.5°/s. The other stimulus component had a low luminance

contrast of 37.5% and moved at a faster speed of 10°/s. Both stimulus components moved at 100%

coherence and were either overlapping or spatially-separated within the RF of a given MT neuron

as in the first experiment. We also measured the direction tuning curves when the two stimulus

components both had high luminance contrast (77.5%) and moved at 2.5°/s and 10°/s, respectively,

at 100% coherence.

We recorded from 13 MT neurons using these visual stimuli. Figure 6 shows the

population-averaged tuning curves. When both stimulus components had high contrast, the peak

response elicited by the faster (10°/s) stimulus component moving in the PD (i.e. 0°) was greater

than that elicited by the slower (2.5°/s) component moving in the PD. The component responses

are shown in green and purple in Figure 6A. When the two stimulus components were overlapping,

the tuning curve elicited by both stimulus components (shown in red) is biased toward the faster

stimulus component, more than what is predicted by the average of the component responses

(shown in gray) (Fig. 6A). We fitted the direction tuning curves using the SNL model for each

neuron (Eq. 1). The median response weight obtained by the model fit for the faster stimulus

component (0.88) was significantly greater than the median weight (0.41) for the slower

component (Wilcoxon signed-rank test, p = 7.3 x 10^{-4}). This result extended our previous finding

of the response bias toward the faster stimulus component for stimuli moving in the same direction

(unpublished results) to stimuli moving in different directions.

When the overlapping stimuli moving at different speeds had different luminance contrasts,

the responses elicited by both stimulus components showed a strong bias toward the "high contrast

& slower speed" component, even though the peak response to this component alone was

significantly weaker than that to the "low contrast & faster speed" component (Fig. 6B). We found

the same result when the two stimulus components overlapped at a different site within the RF

(Fig. 6C). Under the overlapping condition, the median response weight for the "high contrast &

slower speed" component was 0.81, which was significantly greater than the median weight for

the "low contrast & faster speed" component (0.17) (Wilcoxon signed-rank test, p = 2.4 x 10^{-4}).

Separating the two stimulus components spatially within the RF abolished the bias toward the

"high contrast & slower speed" component (Fig. 6D). As the spatial arrangement of the stimulus

components changed from overlapping to spatially separated, the median bias index (Eq. 4)

decreased significantly from 0.65 to -0.08 (Wilcoxon signed-rank test, p = 0.0012). These results

confirmed that luminance contrast has a dominant effect on MT responses elicited by overlapping

stimuli, which is not unique to the competition between contrast and motion coherence. The spatial

arrangement of visual stimuli can substantially change the competition between multiple stimuli

within the RF.

Fitting response tuning curve using the normalization model

Previous studies have shown that neuronal responses elicited by multiple stimuli in many

brain areas can be described by a divisive normalization model (Carandini and Heeger, 2011). We

asked whether our results could also be accounted for by response normalization. We first fitted

the data using the following equation:

$$R_{pred}(\theta_1, \theta_2) \ = \frac{S_1^n}{S_1^n + S_2^n + \sigma} R_1(\theta_1) + \frac{S_2^n}{S_1^n + S_2^n + \sigma} R_2(\theta_2) + c \ , \qquad (5)$$

where R_1 and R_2 are the evoked direction tuning curves to the two stimulus components 1 and 2,

respectively. θ_1 and θ_2 are the component directions. S_1 and S_2 represent the signal strengths of the

"low contrast & high coherence" component and the "high contrast & low coherence" component,

respectively. R_{pred} is the model-predicted response elicited by both stimulus components presented

simultaneously. n, σ, and c are model parameters with the constraints of $n \geqslant 1$ and $c > 0$. Equations

of the similar form have been used previously to describe normalization involving contrast, in

which case the signal strength is simply the luminance contrast (Carandini et al., 1997; Busse et

al., 2009; Xiao et al., 2014; Bao and Tsao, 2018). Since our visual stimuli competed in more than

one feature domain, it was not obvious which stimulus component had an overall stronger signal

strength. Because the brain has to make an inference of the signal strength based on the elicited

neural responses, we assumed that the signal strength of a stimulus component, in the "eye" of MT

neurons, is reflected in the neural responses elicited by that stimulus component moving in a fixed

direction summed across a population of MT neurons that have different PDs evenly spanning

360º. This summed population response is invariant to the direction of the stimulus component,

which is suitable for representing signal strength. Equivalently, the summed population neural

response in MT can be approximated by summing the responses of each neuron elicited by

stimulus component i moving in different directions spanning 360º and averaged across neurons

in our data sample, e.g. to sum the population-averaged component responses across directions in

Figure 2. This was how we calculated S_i, (i =1, 2).

This normalization model (Eq. 5) failed to capture the response tuning to overlapping bi-

directional stimuli, accounting for only 33% of the response variance (34% for site a, 32% for site

b). The model performed better when stimuli were separated, accounting for 66% of the variance.

We found similar results when using this model to fit the data from our second experiment, in

which luminance contrast competed with motion speed. The model accounted for an average of

44% of the response variance (38% for site a, 50% for site b) when stimuli were overlapping, and

77% of the variance when stimuli were separated (Table 2).

It has been suggested that response normalization can be tuned, such that individual

stimulus components contribute differently to normalization (Ni et al., 2012; Rust et al., 2006; also

see Carandini et al., 1997). We therefore fitted our data using a tuned normalization equation:

$$R_{pred}(\theta_1, \theta_2) \; = \frac{S_1^n}{S_1^n + \alpha S_2^n + \sigma} R_1(\theta_1) + \frac{S_2^n}{S_1^n + \alpha S_2^n + \sigma} R_2(\theta_2) + c \, , \qquad (6)$$

where α is a positive parameter that scales the contribution of S_2 with respect to S_1 to normalization.

We found that introducing tuned normalization did not improve the model performance at all when

stimuli were overlapping, accounting for an average of 33% of the response variance (34% for site

a, 32% for site b). When stimuli were separated, the tuned normalization model accounted for 68%

of the variance. We found the same results when fitting the data collected when contrast competed

with speed (Table 2).

The poor fit of the responses under the overlapping condition by the standard normalization

model (Eq. 5) can be understood because MT neurons showed a very strong bias toward the high

contrast component, whereas S_1 and S_2 were similar. The tuned normalization was not able to

improve the fit because, although it changed the relative contributions of the stimulus components

to the normalization pool in the denominator, it kept the numerators in Equation 6 unchanged.

Hence the relative weights for the two stimulus components did not change. To capture the strong

bias toward the high contrast component in the overlapping condition, a weighting parameter is

needed in the numerator. Accordingly, we fitted our results using the following equation:

557
$$R_{pred}(\theta_1, \theta_2) = \frac{S_1^n}{S_1^n + \beta S_2^n + \sigma} R_1(\theta_1) + \frac{\beta S_2^n}{S_1^n + \beta S_2^n + \sigma} R_2(\theta_2) + c, \qquad (7)$$

where β is a positive parameter and appears in both the numerator and the denominator. This

parameter allows the relative response weights for the two stimulus components to vary. When β

is greater than one, the response weight for the high contrast component (R_2) is greater than that

for the low contrast component (R_1). As expected, this equation fitted the data well, accounting for

>80% of the response variance for both the overlapping and spatially separated conditions (Table

2). However, the normalization model itself does not provide an explanation for why the response

weight is greater for the high contrast component in the overlapping condition but not in the

spatially separated condition.

Table 2. Fitting the direction tuning curves using the normalization model

Visual Stimuli	S_1	S_2	Percentage of Variance Accounted for (mean $\pm$ std)		
Contrast vs. Coherence (N = 70)	Low contrast & high coherence	High contrast & low coherence	Normalization (Eq. 5)	Tuned Normalization (Eq. 6)	Normalization with weighted Numerators (Eq. 7)
Overlapping (site a)	122.3	126.0	34 ± 18	34 ± 18	86 ± 16
Overlapping (site b)	128.3	130.9	32 ± 19	32 ± 19	81 ± 19
Spatially Separated	130.4	130.3	66 ± 24	68 ± 25	83 ± 17
Contrast vs. Speed (N = 13)	Low contrast & faster speed	High contrast & slower speed			
Overlapping (site a)	128.1	83.6	38 ± 21	39 ± 21	88 ± 14
Overlapping (site b)	113.3	81.3	49 ± 20	49 ± 20	84 ± 15
Spatially Separated	128.1	81.3	77 ± 20	77 ± 20	90 ± 5

570
571
Computer simulations using a V1-MT model

Our spatially separated visual stimuli fall inside the RFs of single MT neurons, whereas

only one of the stimulus components would fall inside the RFs of single V1 neurons. Hence, our

spatially-separated visual stimuli can interact within the RFs of MT neurons but not V1 neurons.

In contrast, the overlapping stimuli can interact within the RFs of both MT and V1 neurons. To

explore the neural mechanisms underlying our physiological findings, we conducted computer

simulations using a hierarchical feedforward model adapted from Simoncelli and Heeger (1998).

This model consists of two processing stages corresponding to areas V1 and MT. Each stage carries

out a series of computations including spatiotemporal filtering, spatial pooling, rectification, and

divisive normalization. At the V1 stage, simple cells receive input directly from the visual stimulus

and complex cells pool inputs from rectified and divisively normalized responses of V1 simple

cells. At the MT stage, MT neurons pool inputs from V1 complex cells, followed by rectification

and divisive normalization (Simoncelli and Heeger, 1998; Rust et al., 2006).

We generated random-dot visual stimuli that are similar to those used in our physiological

experiments and simulated the neuronal responses in areas MT and V1. The visual stimuli and a

simplified architecture of the model are illustrated in Figure 7. The diameter of each random-dot

patch was 3°, extending 63 pixels. The RF sizes of model V1 and MT neurons, set by the sizes of

the convolution filters, were 1.2° and 10° in diameter, respectively (see Methods). The populations

of model neurons in V1 and MT stages approximately tiled a sphere in the spatiotemporal

frequency domain, as in Simoncelli and Heeger's model (1998). The RFs of V1 and MT neuron

populations covered a region of the visual field that was 17.3° x 17.3°. In the overlapping

condition, the apertures of two random-dot patches overlapped within the RFs (Fig. 7A). In the

spatially-separated condition, the two random-dot patches were placed side by side, separated by

a blank gap that was 1.2° wide, within the RFs of single MT neurons (Fig. 7B). In the overlapping

condition, the V1 neurons whose RFs covered *site a* were activated by both stimulus components

(Fig. 7A). In the spatially-separated condition, V1 neurons were activated by only one stimulus

component, either at site *a* or site *b* (Fig. 7B).

We tuned the model parameters (see Methods) to match the experimentally measured

contrast response functions of V1 and MT neurons (Sclar et al., 1990) and the coherence response

function of MT neurons (Britten and Newsome, 1998). The simulated contrast response functions

of V1 and MT neurons fitted the experimental data almost perfectly, and the simulated coherence

response function of MT neurons also matched the data well (Fig. 8A-C). As far as we know, an

experimentally measured coherence response function of V1 neurons has not been described
previously. Our simulations show that V1 responses increased monotonically with the coherence
level of moving random-dot stimuli (Fig. 8D). The model V1 neurons had lightly higher firing
rates in response to low coherence stimuli and more trial-to-trial variability in comparison with the
model MT neurons (Fig. 8C and D).

The MT responses elicited by our visual stimuli that competed between luminance contrast
and motion coherence were well captured by the model. Consistent with our experimental data
(Fig. 2), the tuning curve of model MT neurons to the "low contrast & high coherence" component
had a greater peak response than that of the "high contrast & low coherence" component (Fig. 9A,
B). In the overlapping condition, the simulated MT response elicited by the bi-directional stimuli
was nearly completely biased toward the weaker "high contrast & low coherence" component (Fig.
9A), as found in the neural data. The model also captured the change of MT response tuning when
visual stimuli were rearranged spatially. In the spatially-separated condition, the tuning curve of
model MT neurons elicited by the bi-directional stimuli was no longer dominated by the "high
contrast & low coherence" component (Fig. 9B).

At the V1 stage of the model, the tuning curves of V1 complex cells showed a slightly
greater mean peak response to the "high contrast & low coherence" component than to the "low
contrast & high coherence" component (Fig. 9C). In the overlapping condition, the simulated V1
response elicited by the bi-directional stimuli was strongly biased toward the "high contrast & low
coherence" component (Fig. 9C), to the extent similar to that found in model MT neuron (Fig. 9A),
as measured by the weights for the component responses using the SNL model fits. The bias index
(Eq. 4) for the V1 model neuron was 0.90 and that for the MT model neuron was 0.93. These
simulation results suggest that the strong bias toward the "high contrast & low coherence"
component found in MT is inherited from V1.

In the spatially-separated condition, the V1 response elicited by the bi-directional stimuli
was the same as that elicited by the single stimulus component placed within the RFs of V1 neurons
(Fig. 9D, E). Although the V1 peak response elicited by the "high contrast & low coherence"
component at site *a* was slightly stronger than that elicited by the "low contrast & high coherence"

component at site *b*, the MT response elicited by the bi-directional stimuli was skewed toward the

"low contrast & high coherence" component, consistent with the average of the component

responses (Fig. 9B). These simulation results suggest that MT response elicited by the bi-

directional stimuli in the spatially-separated condition (Fig. 9B) may be due to feature competition

within MT.

The response tuning curves of single MT neurons measured by varying the VA direction

of the bi-directional stimuli can be mapped to the responses of a population of MT neurons that

have different PDs, elicited by the bi-directional stimuli moving in a given VA direction. Figure 7

summarizes the changes of the response distributions across neuron populations at V1 and MT

stages, under the overlapping and spatially-separated conditions. These results reveal the

importance of neural processing at different stages of the visual hierarchy on determining how

multiple visual stimuli compete within neurons' RFs in a given brain area.

651
652
653   **Discussion**
654

We have shown that how MT neurons represent multiple stimuli competing in more than

one feature domain depends on the spatial arrangement of the visual stimuli. When two stimuli are

overlapping, MT responses are dominated by the stimulus component that has high contrast. When

two stimuli are spatially separated, the contrast dominance is abolished. Our neural data and model

simulations suggest that the contrast dominance found with overlapping stimuli is due to

normalization occurring at an input stage fed to MT, and MT neurons cannot overturn this contrast

dominance based on their own feature selectivity. The interaction between spatially separated

stimuli can largely be explained by normalization within area MT. By using multiple visual stimuli

competing in more than one features domain, our study revealed how neural processing along the

hierarchical visual pathway shapes neural representation of multiple visual stimuli in extrastriate

cortex.

Consideration of the effect of attention

Attention can bias neuronal responses elicited by multiple stimuli in the RF in favor of the

attended stimulus (Reynolds et al., 1999; Li and Basso, 2005; Treue and Maunsell, 1996; Ferrera

and Lisberger, 1997; Treue and Martinez-Trujillo, 1999; Recanzone and Wurtz, 2000; Lee and

Maunsell, 2010). Although in this study the animals performed a fixation task without the need to

engage goal-directed attention, could the high contrast component capture stimulus-driven

attention (Corbetta and Shulman, 2002) and bias the neuronal response elicited by the overlapping

stimuli?　Several considerations argue against this possibility. While an abrupt stimulus onset

captures attention (Yantis and Jonides, 1984), a visual stimulus that is brighter than other

distractors does not automatically capture attention (Jonides and Yantis, 1988). The two stimulus

components of our overlapping stimuli were turned on and started to move at the same time. The

stimulus onset may automatically draw attention toward the spatial location of the overlapping

stimuli, but it is unlikely to draw attention toward only the high contrast component. Furthermore,

stimulus-driven attention occurs with a time delay (Nakayama and Mackeben, 1989) and its effect

on neuronal responses in MT is transient, lasting for about 70 ms (Busse et al., 2008). In contrast,

we found that the response bias toward the high contrast component is present in the very

beginning of the neuronal responses following the onset of the static stimuli, and the bias is

persistent throughout the motion period (Fig. 5). In addition, Wannig and colleagues (2007) have

shown that attention directed to one of two overlapping surfaces can alter the responses of MT

neurons. However, attention led to a response magnitude modulation of about 20% in MT between

conditions when attention was directed to two different surfaces (Wannig et al., 2007). Even if, for

some reason, the animals were consistently attending to the high contrast component throughout

the stimulus presentation period in our study, the effect of attention would be insufficient to

account for the nearly complete dominance by the high contrast component.

Mechanisms underlying stimulus interactions

The primate visual system is hierarchically organized (Maunsell and van Essen, 1983;

Felleman and Van Essen, 1991). The response properties of neurons in a visual area are shaped by

feedforward input, as well as intra-areal and feedback processes. To understand the mechanisms

underlying neural encoding of multiple stimuli, it is important to determine how these processes

contribute to the RF properties in a given visual area. However, it is often difficult to disentangle

the contribution of feedforward input from other neural processes. We have previously found that,

in response to overlapping stimuli, MT neurons show a bias toward the stimulus component that

has a higher signal strength, defined by either luminance contrast or motion coherence (Xiao et

al., 2014). The response bias can be described by a model of divisive normalization. Because neurons in V1 also show a bias toward the stimulus component that has a higher contrast (Busse et al., 2009; MacEvoy et al., 2009) and divisive normalization may occur in both V1 and MT (Simoncelli and Heeger, 1998; Heuer and Britten, 2002), it was unclear how the feedforward input from V1 contributed to the response bias found in MT.

In this study, we are able to differentiate the impact of feedforward input from other neural processes on the response properties of MT neurons. Our results suggest that neurons in V1 may respond more strongly to the "high contrast & low coherence" component than to the "low contrast & high coherence" component used in our experiment, due to V1 neurons' sensitivities to contrast and coherence. When two stimuli overlap, the responses of V1 neurons elicited by both stimulus components may already show a strong bias toward the "high contrast & low coherence" component due to divisive normalization in V1 (Fig. 9C). MT neurons are no longer able to remix the stimulus components according to their own sensitivities to contrast and coherence. In other words, MT neurons inherit the response bias toward the high contrast component from their input. When two visual stimuli are spatially separated, MT neurons receive inputs from two different pools of V1 neurons and each neuron pool responds to only one stimulus component (Fig. 7B). The neuronal responses elicited by the two stimulus components remain separated in V1. MT neurons can mix the responses elicited by the two stimulus components via spatial and directional pooling and divisive normalization within MT. As a result, the mixing in MT may well reflect the sensitivities of MT neurons to different stimulus features. Our model simulations make predictions regarding how V1 neurons respond to multiple competing stimuli (e.g. as shown in Fig. 9C), which can be tested in future physiological study.

Implications on normalization and encoding of multiple visual stimuli

Our finding that the response weighting for competing stimuli depends on the spatial arrangement provides a new perspective on the well-established normalization model (Carandini and Heeger, 2011). The basic form of normalization equations (Eqs. 5-6) predicts that the response weight for a stimulus component increases with its signal strength, but does not consider the spatial arrangement of the visual stimuli. We made a surprising finding that MT response to overlapping stimuli cannot be predicted by the population neural responses in MT elicited by the individual

stimulus components. One must consider the neural computations occurring along the hierarchical

visual pathway.

Majaj, Carandini, and Movshon (2007) showed that pattern-direction selective neurons in

MT characterized by overlapping drifting gratings (i.e. plaid) do not integrate the directions of the

component gratings when they were spatially separated within the RF, suggesting that the

computation underlying pattern-direction selectivity in MT is local. Different from the plaid, the

overlapping random-dot stimuli used in our study elicit the percept of motion transparency. We

showed that changing the spatial arrangement of visual stimuli can have a substantial impact not

only on motion integration but also on the competition between multiple stimuli. Our results

revealed that contrast has a dominant effect in determining stimulus competition within a local

spatial region when multiple stimuli differ in more than one feature domain. When visual stimuli

are spatially separated, the effect of contrast is substantially reduced.

A seminal model involving MT neurons pooling inputs from V1 and divisive normalization

in both V1 and MT has been successful in explaining a range of experimental results of MT

responses (Simoncelli and Heeger, 1998; Rust et al., 2006). However, the model in its original

form does not specify how features are spatially integrated and it does not differentiate overlapping

and spatially separated stimuli (Majaj et al., 2007). In our study, we adapted this model to simulate

both overlapping and spatially separated conditions and showed that the framework can explain

our main physiological findings. Also using this model, Busse, Wade, and Carandini (2009)

previously demonstrated the impact of response normalization in V1 on neural response in MT.

They showed that, by making the contrasts of two drifting gratings of a plaid to be unequal, the

response of a model MT neuron changed from representing the pattern motion of the plaid to

mostly representing the higher-contrast grating component, likely due to contrast normalization in

V1 (Busse et al., 2009). However, the MT response elicited by the higher-contrast grating alone

could also be greater than that elicited by the lower-contrast grating. The model-predicted response

bias toward the higher-contrast component in MT may also be contributed by response

normalization within MT, akin to our experimental result obtained using random-dot stimuli with

unequal contrasts (Xiao et al., 2014). In comparison, our current study provides unequivocal new

evidence on how responses in MT are shaped by the hierarchical network. By using two stimuli

competing in more than one feature domain, we demonstrated neurophysiologically and

computationally the substantial impact of stimulus competition in the input stage on the neuronal

responses in MT and how that impact changes with the spatial arrangement of visual stimuli. Our

finding may also apply to other visual areas in the hierarchical network, including those in the

ventral visual stream where response normalization has been well documented.

References

768 **References**
769
Albright TD, Desimone R. 1987. Local precision of visuotopic organization in the middle
temporal area (MT) of the macaque. *Exp Brain Res* 65: 582-92
Bao P, Tsao DY. 2018. Representation of multiple objects in macaque category-selective areas.
Nat Commun 9: 1774
Born RT, Bradley DC. 2005. Structure and function of visual area MT. *Annu Rev Neurosci* 28:
157-89
Braddick O. 1993. Segmentation versus integration in visual motion processing. *Trends Neurosci*
16: 263-8
Britten KH. (2003). The middle temporal area: motion processing and the link to perception In
The visual neurosciences, ed. LM Chalupa, JS Werner, pp. 1203-16. London: The MIT
press
Britten KH, Heuer HW. 1999. Spatial summation in the receptive fields of MT neurons. *J
Neurosci* 19: 5074-84
Britten KH, Newsome WT. 1998. Tuning bandwidths for near-threshold stimuli in area MT. *J
Neurophysiol* 80: 762-70
Britten KH, Shadlen MN, Newsome WT, Movshon JA. 1992. The analysis of visual motion: a
comparison of neuronal and psychophysical performance. *J Neurosci* 12: 4745-65
Britten KH, Shadlen MN, Newsome WT, Movshon JA. 1993. Responses of neurons in macaque
MT to stochastic motion signals. *Vis Neurosci* 10: 1157-69
Busse L, Katzner S, Treue S. 2008. Temporal dynamics of neuronal modulation during
exogenous and endogenous shifts of visual attention in macaque area MT. *Proc Natl
Acad Sci U S A* 105: 16380-5
Busse L, Wade AR, Carandini M. 2009. Representation of concurrent stimuli by population
activity in visual cortex. *Neuron* 64: 931-42
Carandini M, Heeger DJ. 2011. Normalization as a canonical neural computation. *Nat Rev
Neurosci* 13: 51-62
Carandini M, Heeger DJ, Movshon JA. 1997. Linearity and normalization in simple cells of the
macaque primary visual cortex. *J Neurosci* 17: 8621-44
Corbetta M, Shulman GL. 2002. Control of goal-directed and stimulus-driven attention in the
brain. *Nat Rev Neurosci* 3: 201-15
Ferrera VP, Lisberger SG. 1997. Neuronal responses in visual areas MT and MST during smooth
pursuit target selection. *J Neurophysiol* 78: 1433-46
Gattass R, Gross CG. 1981. Visual topography of striate projection zone (MT) in posterior
superior temporal sulcus of the macaque. *J Neurophysiol* 46: 621-38
Heuer HW, Britten KH. 2002. Contrast dependence of response normalization in area MT of the
rhesus macaque. *J Neurophysiol* 88: 3398-408
Jonides J, Yantis S. 1988. Uniqueness of abrupt visual onset in capturing attention. *Percept
Psychophys* 43: 346-54
Lee J, Maunsell JH. 2010. Attentional modulation of MT neurons with single or multiple stimuli
in their receptive fields. *J Neurosci* 30: 3058-66
Li X, Basso MA. 2005. Competitive stimulus interactions within single response fields of
superior colliculus neurons. *J Neurosci* 25: 11357-73
MacEvoy SP, Tucker TR, Fitzpatrick D. 2009. A precise form of divisive suppression supports

population coding in the primary visual cortex. *Nat Neurosci* 12: 637-45

Majaj NJ, Carandini M, Movshon JA. 2007. Motion integration by neurons in macaque MT is

local, not global. *J Neurosci* 27: 366-70

Maunsell JH, van Essen DC. 1983. The connections of the middle temporal visual area (MT) and

their relationship to a cortical hierarchy in the macaque monkey. *J Neurosci* 3: 2563-86

Morgan ML, Deangelis GC, Angelaki DE. 2008. Multisensory integration in macaque visual

cortex depends on cue reliability. *Neuron* 59: 662-73

Movshon JA, Newsome WT. 1996. Visual response properties of striate cortical neurons

projecting to area MT in macaque monkeys. *J Neurosci* 16: 7733-41

Nakayama K, Mackeben M. 1989. Sustained and transient components of focal visual attention.

Vision Res 29: 1631-47

Newsome WT, Pare EB. 1988. A selective impairment of motion perception following lesions of

the middle temporal visual area (MT). *J Neurosci* 8: 2201-11

Ni AM, Ray S, Maunsell JH. 2012. Tuned normalization explains the size of attention

modulations. *Neuron* 73: 803-13

Park, W. J. and Tadin, D. (2019). Motion Perception. In Stevens' Handbook of Experimental

Psychology and Cognitive Neuroscience, J. T. Wixted (Ed)

Qian N, Andersen RA. 1994. Transparent motion perception as detection of unbalanced motion

signals. II. Physiology. *J Neurosci* 14: 7367-80

Recanzone GH, Wurtz RH. 2000. Effects of attention on MT and MST neuronal activity during

pursuit initiation. *J Neurophysiol* 83: 777-90

Recanzone GH, Wurtz RH, Schwarz U. 1997. Responses of MT and MST neurons to one and

two moving objects in the receptive field. *J Neurophysiol* 78: 2904-15

Reynolds JH, Chelazzi L, Desimone R. 1999. Competitive mechanisms subserve attention in

macaque areas V2 and V4. *J Neurosci* 19: 1736-53

Rust NC, Mante V, Simoncelli EP, Movshon JA. 2006. How MT cells analyze the motion of

visual patterns. *Nat Neurosci* 9: 1421-31

Sclar G, Maunsell JH, Lennie P. 1990. Coding of image contrast in central visual pathways of

the macaque monkey. *Vision Res* 30: 1-10

Simoncelli EP, Heeger DJ. 1998. A model of neuronal responses in visual area MT. *Vision Res*

38: 743-61

Snowden RJ, Treue S, Erickson RG, Andersen RA. 1991. The response of area MT and V1

neurons to transparent motion. *J Neurosci* 11: 2768-85

Treue S, Martinez Trujillo JC. 1999. Feature-based attention influences motion processing gain

in macaque visual cortex. *Nature* 399: 575-9

Treue S, Maunsell JH. 1996. Attentional modulation of visual motion processing in cortical areas

MT and MST. *Nature* 382: 539-41

Wannig A, Rodriguez V, Freiwald WA. 2007. Attention to surfaces modulates motion

processing in extrastriate area MT. *Neuron* 54: 639-51

Xiao J, Huang X. 2015. Distributed and Dynamic Neural Encoding of Multiple Motion

Directions of Transparently Moving Stimuli in Cortical Area MT. *J Neurosci* 35: 16180-

98

Xiao J, Niu YQ, Wiesner S, Huang X. 2014. Normalization of neuronal responses in cortical area

MT across signal strengths and motion directions. *J Neurophysiol* 112: 1291-306

Yantis S, Jonides J. 1984. Abrupt visual onsets and selective attention: evidence from visual

search. *J Exp Psychol Hum Percept Perform* 10: 601-21

Figures and Figure Legends

Figure 1. The response tuning curves of two example MT neurons to overlapping (A, B) and spatially-separated stimuli (C). Visual stimuli were achromatic random-dot patches moving in two directions separated by 90°. The "low contrast & high coherence" component (shown in blue arrow) moved at the clockwise side of the two component directions, whereas the "high contrast & low coherence" component (shown in green arrow) moved in the direction at the counter-clockwise side. The X-axis labeled in black indicates the vector average direction of the bi-directional stimuli. The X-axes labeled in blue and green (**A2**) indicate the direction of the "low contrast & high coherence" component (Dir. 1) and the direction of "high contrast & low coherence" component (Dir. 2), respectively. The three X-axes are aligned such that the component directions shown in blue and green correspond to the directions of the two stimulus components at each vector average direction. **A1-C1:** Response tuning curves from one neuron. **A2-C2:** Response tuning curves from another neuron. The responses elicited by the bi-directional stimuli are shown in red (R12). The SNL model fits of R12 are shown in black. Error bars represent standard errors.

Figure 2. Population-averaged tuning curves to the bi-directional stimuli (red) and the unidirectional stimulus components (blue and green). The vector average direction of the bi-directional stimuli and the directions of individual stimulus components are labeled in the corresponding X-axes (**A**), following the same convention as in Figure 1. The direction of 0° was aligned with each neuron's PD before the tuning curves were averaged across neurons. The stimulus components were overlapping at site *a* (**A**) or site *b* (**B**), or spatially separated (**C**) within the RFs. The width of each tuning curve represents the standard error. The average of the responses to the two stimulus components is shown in gray.

Figure 3. The effect of the spatial arrangement of the bi-directional stimuli on the response weights for the stimulus components. Each dot represents the result from one neuron. Comparing the response weights for the "low contrast & high coherence" component (ordinate) with the "high contrast & low coherence" component (abscissa) under the overlapping (**A**) and the spatially separated (**B**) conditions. **C.** Comparing the bias indices between the spatially separated (ordinate) and overlapping (abscissa) conditions. The histograms in C show the distributions of the bias index for the overlapping (top) and spatially separated (right) conditions.

Figure 4. Control for the directional arrangement of the two stimulus components. **A, B.** Response tuning curves averaged across 15 MT neurons to the bi-directional stimuli and the stimulus components when the direction of the "high contrast & low coherence" component was placed at the counter-clockwise side of the two component directions, as in Figures 1 and 2. **C, D.** Response tuning curves averaged across the same 15 neurons when the direction of the "high contrast & low coherence" component was placed at the clockwise side of the two component directions. **A, C.** Overlapping condition. **B, D.** Spatially separated condition. Notice the switch of the values in the X-axes of the component directions, shown in blue and green, between A and C, as well as between B and D.

Figure 5. Timecourse of the neuronal responses to the bi-directional stimuli and the stimulus components. Peristimulus time histograms (PSTHs) were calculated using a 10-ms time bin and averaged across 70 neurons. **A, B.** The two stimulus components overlapped (at site *a*) within the RF. **C, D.** The two stimulus components were spatially separated within the RF. The dashed vertical lines at -200 ms indicate the onset of the static stimuli. The solid vertical lines at time 0 indicate motion onset. The solid horizontal bars shown in A and C indicate the stimulus motion period. The "high contrast & low coherence" component moved in the PD in A and C, and moved in a non-PD in B, D.

Figure 6. Averaged response tuning curves to two stimulus components that moved in different directions and at different speeds. Both stimulus components moved at 100% coherence. The response tuning to both stimulus components presented simultaneously is shown in red. The width of each tuning curve represents the standard error. The average of the component responses elicited by the individual stimulus components is shown in gray. **A.** Both stimulus components had high luminance contrast and were overlapping. **B-D.** The two stimulus components competed in luminance contrast and motion speed. The faster speed component had low luminance contrast, whereas the slower speed component had high luminance contrast. The stimulus components were overlapping at site *a* (**B**) or site *b* (**C**), or were spatially separated (**D**) within the RFs. The vector average direction of the bi-directional stimuli and the directions of individual stimulus components are labeled in the corresponding X-axes (**A, B**), following the same convention as in Figure 2.

Figure 7. Illustration of a simplified architecture for the V1-MT model. Each MT neuron receives feedforward inputs from multiple neurons at the V1 stage. Responses are divisively normalized by the sum of local population activity at both V1 and MT stages. Each small circle represents a neuron and the black arrow inside the circle indicates the PD. The color of each circle indicates the response magnitude of the neuron. Yellow means maximum response and red means minimum response. Visual stimuli are illustrated below neural circuit as the input to the V1 stage. The green and blue arrows represent the "high contrast & low coherence" component and the "low contrast & high coherence" component, respectively. Two pools of neurons at the V1 stage that respond only to site *a* or site *b* respectively are illustrated. The RFs of the MT neurons are illustrated by the dotted ellipse and cover both site *a* and site *b*. **A.** Overlapping condition. **B.** Spatially separated condition.

Figure 8. Contrast and coherence response functions of model V1 (**B, D**) and MT (**A, C**) neurons.
A, B. Fitted contrast response functions to sinusoidal gratings for model neurons. Green curves
are experimental data replotted from Sclar et al. (1990). **C**. Fitted coherence response function to
high contrast random-dots for model MT neurons. Green dots are experimental data replotted from
Britten and Newsome (1998). The green curve is the spline fit of the experimental data points. ***D***:
Coherence response to high contrast random-dots for model V1 complex cells. The widths of the
blue curves in C and D represent the standard deviation. N indicates the number of repeats for
simulations. The stimulus dots were regenerated randomly for each simulation in C and D.

Figure 9. Computer simulations of direction tuning curves of MT and V1 neurons to the bi-directional stimuli used in the main physiological experiment. The visual stimuli are either overlapping (**A, C**) or spatially separated (**B, D, E**) within the RFs of model MT neurons. The two stimulus components compete in luminance contrast and motion coherence. The simulated responses to the "low contrast & high coherence" component and the "high contrast & low coherence" component are shown in blue and green, respectively. The responses to the bi-directional stimuli are shown in red. The vector average direction and the directions of individual stimulus components are labeled in the corresponding X-axes (**A**), following the same convention as in Figure 2. **A, B.** Simulated responses of model MT neurons. **C-E.** Simulated responses of model V1 complex cells. Widths of the tuning curves in A and B and the error bars in C-E represent standard deviations.