Diverse dynamics of glutamatergic input from sensory neurons underlie heterogeneous responses of olfactory bulb outputs *in vivo*

Andrew K. Moran[1,2], Thomas P. Eiting[2], Matt Wachowiak[2]

[1]Interdepartmental Program in Neuroscience, [2]Department of Neurobiology and Anatomy, University of Utah School of Medicine, Salt Lake City, UT

Correspondence: Matt Wachowiak, Department of Neurobiology and Anatomy, University of Utah, Salt Lake City, UT, USA. email: matt.wachowiak@utah.edu

ABSTRACT

Mitral/tufted (MT) cells of the olfactory bulb (OB) show diverse temporal responses to odorant stimulation that are thought to encode odor information. To understand the role of sensory input dynamics versus OB circuit mechanisms in generating this diversity, we imaged glutamate signaling onto MT cell dendrites in anesthetized and awake mice. We found surprising diversity in the dynamics of these signals, including excitatory, suppressive, and biphasic responses as well as nonlinear changes in glutamate signaling across inhalations. Simultaneous imaging of glutamate and calcium signals from MT cell dendrites revealed highly correlated responses for both signals. Glutamate responses were only weakly impacted by blockade of postsynaptic activity, implicating sensory neurons as a primary source of glutamate signaling onto MT cells. Thus, the dynamics of sensory input alone, rather than emergent features of OB circuits, may account for much of the diversity in MT cell responses that underlies OB odor representations.

Keywords: imaging, glutamate, calcium, temporal coding, active sensing

INTRODUCTION

The neural activity underlying odor perception is inherently dynamic. In vertebrates, olfactory sensory input arrives at the brain as bursts of activity driven by each inhalation, and patterns of inhalation are actively controlled by an animal as it samples its environment. At the level of the olfactory bulb (OB), olfactory sensory neurons (OSNs) expressing a single type of odorant receptor converge onto glomeruli[1] where they make glutamatergic synapses onto targets including juxtaglomerular interneurons and the principal OB output neurons, mitral and tufted (MT) cells[2]. Odorant responses among MT cells can be temporally complex, displaying sequences of excitation and suppression that occur relative to a single inhalation and across multiple inhalations[3-6]. Olfactory information is likely represented both by the identity of activated MT cells as well as by their temporal dynamics[4,7-9].

Much of the complexity in MT cell response patterns has been attributed to processing by inhibitory OB circuits[10-13]. However, excitatory synaptic inputs to MT cells, which occur solely on the MT cell apical tuft in each glomerulus, can also contribute to generating temporally complex and diverse patterns of MT cell spiking. OSNs themselves respond to odorant inhalation with different latencies and show nonlinear changes in their activation patterns as a function of odorant concentration, sampling time and sampling frequency[14,15]. Additional diversity in excitatory synaptic input to MT cells may arise from multisynaptic pathways – for example, via external tufted (ET) cells[12,16-18]. However, the contribution of glutamatergic input dynamics to determining MT cell odorant response patterns in vivo remains poorly understood.

Here, we imaged odorant-evoked and inhalation-linked glutamate signals in glomeruli of the mouse OB using the genetically-encoded glutamate sensor iGluSnFR and its second-generation variants[19,20], expressed selectively in MT cells. Glomerular glutamate transients reported inhalation-linked dynamics with high fidelity, allowing temporally precise monitoring of glutamatergic signaling across the dorsal OB in both the anesthetized and awake mouse. We observed a striking degree of diversity in the temporal dynamics of glutamatergic input onto MT cell dendrites in a glomerulus, over time scales involving a single inhalation and across multiple inhalations of odorant. We found a high correspondence between the dynamics of glutamatergic input and that of

MT cell output across glomeruli. Glutamate responses were only weakly impacted by

blockade of postsynaptic activity, pointing to OSNs as the primary source of odorant-

evoked MT cell excitation and a subtle role for multisynaptic excitation. These results

reveal remarkable diversity in the dynamics of glutamatergic drive onto MT cells and

suggest that the complex spatiotemporal activity patterns thought to play a critical role in

coding odor information may arise largely from diverse patterns of feedforward

excitation driven directly by sensory afferents.

RESULTS

Diverse inhalation-linked glutamate transients imaged from MT cell dendrites

74   The timing of odorant- and inhalation-evoked MT cell spiking varies across MT

cells and across odorants, and has long been hypothesized to play a role in coding odor

information[4,7,8]. To assess the degree to which such variability is present at the level of

glutamatergic input to MT cells, we targeted iGluSnFR and its second-generation

variants to MT cells using Cre-dependent viral vectors and MT cell-specific Cre driver

lines[21,22] (Fig. 1a). We have previously shown, using the first-generation iGluSnFR, that

each inhalation of odorant drives a transient of glutamate release onto juxtaglomerular

interneuron processes in the glomerular neuropil[23]**.** We observed similar glutamate

signals with iGluSnFRs expressed in MT cells, with different glomeruli showing

glutamate transients with varying onset latencies and durations, and with distinct

changes in the glutamate signal evoked by successive inhalations across a longer

odorant presentation (Fig. 1b). In the superficial OB layers, glutamate signals appeared

restricted to MT cell apical tufts within glomeruli, with negligible signals outside of the

glomerulus or on MT cell primary dendrites or somata (Fig. 1c, d). Odorant-evoked

glutamate signals were also detected along MT cell lateral dendrites in the external

plexiform layer (Supplementary Fig. 1a, b).

90   We analyzed inhalation-linked dynamics of the glutamate signal across

glomerulus-odor pairs by generating 'inhalation-triggered average' responses (ITAs)

from multiple inhalations at low frequencies (0.25 - 0.5 Hz)[3,24]. Most ITA transients

consisting of a rapid rise and a slower decay to baseline. Odorant-evoked ITA dynamics

varied substantially across glomeruli and odorants: glomeruli activated by the same

odorant could respond with different onset latencies, rise-times and durations, and the

same glomerulus could show distinct ITA dynamics for different odorants (Fig. 2a, b).

Inhalation occasionally elicited fluorescence decreases in a glomerulus, suggesting a

phasic decrease in ongoing glutamate release (Fig. 2b, see ROI 6), possibly due to the

inhalation of clean air into the nasal cavity[24].

We compared ITAs across glomerulus-odor pairs using the three SF-iGluSnFR

variants, A184S, A184V and S72A, which have higher, medium and lower affinities for

glutamate, respectively[20]. ITA dynamics varied only slightly between the SF-iGluSnFR

variants (Supplementary Fig. 1c). We made the most measurements with the medium-

and high-affinity variants due to their higher signal-to-noise ratios; across all glomeruli

imaged with A184S, ITA onset latencies ranged from 113 to 253 ms (10^{th} - 90^{th}

percentiles), time to ITA peak ranged from 300 to 533 ms, and ITA durations ranged

from 360 to 867 ms (Fig. 2c). Responses using the A184V variant were similar, with

onset latencies ranging from 153 to 313 ms (10^{th} - 90^{th} percentiles, 113 pairs from 3

mice), time to peak ranging from 313 to 727 ms, and ITA durations ranging from 260 to

1100 ms. This overlap suggests that variation in the dynamics of the SF-iGluSnFR

signal imaged from MT cell tufts largely reflects the variable dynamics of glutamate

signaling in the glomerulus, as opposed to kinetic differences of the reporter. We

compared the variability seen in the glutamate ITA with that from OSNs expressing the

genetically-encoded reporter GCaMP6f, in response to the same odorants in separate

mice (Fig. 2d). Despite the slower kinetics of GCaMP6f compared to iGluSnFR[25], the

variability in onset latencies (relative to median latency) was similar for GCaMP6f and

iGluSnFR signals (Fig. 2d). These results suggest that variability in the onset times of

glutamate transients onto MT cells after each inhalation largely reflect differences in the

timing of spike bursts arriving at OSN presynaptic terminals.

We next asked whether the dynamics of glutamatergic input differ across mitral

and tufted cell subpopulations. Mitral cells and superficial tufted cells (sTCs) have

distinct odorant response properties[26,27]. Many sTCs express the peptide

neurotransmitter cholecystokinin (CCK)[28]; these neurons are strongly driven by

monosynaptic input from OSNs[29], and we have previously shown that they have faster-

onset and less diverse odorant-evoked responses than the general MT cell

population[6,24,30]. We compared glutamate dynamics onto CCK+ sTCs and piriform

cortex-projecting MT cells (pcMTs), using Flex.AAV9.iGluSnFR.A184S virus injection

into the superficial EPL of CCK-IRES-Cre mice or retrograde viral expression via

injection into anterior piriform cortex of Tbet-Cre mice (Fig. 2e). The latter approach

biases expression in mitral and deep tufted cells, with relatively little expression in

sTCs[31]. We found no difference in the dynamics of inhalation-linked glutamate

transients onto CCK+ sTCs compared to pcMT cells, with both cell types showing a

similar distribution of onset latencies, peak times and durations of the iGluSnFR ITA

(Fig. 2f). These results indicate that reported differences in odorant-evoked response

patterns in sTCs and pcMT cells do not arise from differences in the dynamics of their

excitatory input.

Diversity of glutamatergic signaling across multiple inhalations.

To more fully investigate the temporal diversity in patterns of glutamate signaling

onto MT cells, we imaged glomerular responses to 23 odorants in single sessions (see

Table 1). Odorants were delivered at concentrations such that each odorant evoked

relatively sparse responses across the imaging field (Fig. 3a). 16.5% of all glomerulus-

odor pairs (748/4520; 7 fields of view, 3 mice) showed responses that were significant

according to a conservative criterion of ± 7 SD deviation from baseline. Responsive

glomeruli were narrowly tuned across the 23-odorant panel, with high values of lifetime

sparseness for excited glomeruli (S = 0.91 ± 0.05, mean ± SD).

Temporal patterns of odorant-evoked glutamate signals were diverse but robust,

occurring consistently over repeated presentations (Fig. 3b). The most common

response pattern consisted of inhalation-linked transients that persisted over the 2-sec

presentation and returned rapidly to baseline (Fig. 3c, e.g., isopropyl tiglate). Other

response patterns included transient increases that occurred immediately after odorant

onset, prolonged increases that returned slowly to baseline after odorant offset, and

slow, 'facilitating' rises in glutamate (Fig. 3c, e.g. hexanoic acid). Response patterns

varied for different glomeruli activated by the same odorant and for responses of the

same glomerulus to different odorants (Fig. 3c), indicating that response patterns were

neither a function of SF-iGluSnFR kinetics or expression levels in a particular

glomerulus, nor a function of any particular odorant. Odorants also could evoke

responses with a pronounced suppressive component (Fig. 3b, c), although these were

relatively rare, with pure suppression seen in 7.2% (54/748) of glomerulus-odor pairs

and biphasic (excitatory and suppressed) responses seen in 3.5% (26/748) of pairs

(Fig. 3d, e).

A consequence of this diversity was that relative glutamate levels in different

glomeruli varied over the odor presentation. To quantify this, we generated vectors

consisting of time-binned glutamate signals (bin width, 387 ms) across all glomeruli in a

field of view and correlated the vector in the first time bin with that from each successive

time bin over the 2 sec odor presentation. Among all odorant-fields of view (n = 111),

there was a substantial decorrelation beginning in the second bin (time bin 2 vs. time

bin 1: $\Delta r = 0.52 \pm 0.32$) and continuing to the last bin (time bin 5 vs. time bin 1: $\Delta r = 0.39$

± 0.37), although the degree of decorrelation varied greatly for different odorants (Fig.

3f; see also Fig. 3c insets). This decorrelation is qualitatively similar to that observed in

patterns of MT cell spiking over repeated inhalations[3,5,6], suggesting that time-

dependent decorrelation can arise at the level of glutamatergic input to MT cells.

Glomerular glutamate signaling shows diverse temporal dynamics in awake mice

We next imaged iGluSnFR signals in awake, head-fixed mice while monitoring

nasal airflow. Respiration frequencies were typically between 3 and 6 Hz (mean

frequency per session, 4.4 ± 0.34 Hz, measured from 5 sessions), with occasional

pauses and bouts of higher-frequency sniffing. Glutamate transients in awake mice

were temporally diverse, similar to anesthetized mice. For many glomeruli, each

inhalation at 'resting' frequencies (defined as below 5 Hz) elicited a distinct glutamate

transient (Fig. 4a, 4b). We measured inhalation-linked dynamics for glomerulus-odorant

pairs showing significant respiratory modulation by constructing ITA waveforms as

before, but using an external sensor to determine inhalation timing. ITA waveforms

generated from these data were approximately sinusoidal (Fig. 4a, lower right), allowing

for straightforward estimates of onset latency and time-to-peak of the glutamate signal

relative to inhalation. Latency differences between glomeruli were apparent in single

trials and were consistent across repeated inhalations of odorant (Fig. 4a). Overall, ITA

onset latencies, compiled from four mice, varied over a range of 127 ms relative to the

median latency across all glomerulus-odor ITAs in a session (SD = 25 ms, 13 odorants,

7 fields of view, 108 glomerulus-odorant pairs); ITA peak latencies spanned a range of

187 ms (SD = 34 ms) (Fig. 4b). As seen with artificial inhalation, these dynamics could

vary across glomeruli and odorants.

For many glomerulus-odorant pairs, glutamate signals were not modulated by

respiration but instead consisted of a tonic glutamate increase. The same glomerulus

could show strong coupling to respiration for one odorant and a lack of respiratory

coupling for another odorant (Fig. 4c). Similarly, different glomeruli responsive to the

same odorant could show either strong or no respiratory coupling (Fig. 4d, indicating

that the lack of coupling was not due to differences in respiration from trial to trial, nor

was it due to differences in odorant stimulus profile. Odorants could also elicit

decreases in the glutamate signal (Fig. 4d, top trace). Respiration-coupled and tonic

glutamate signals were apparent to a similar degree in mice expressing SF-

iGluSnFR.A184V or SF-iGluSnFR.A184S (2 mice each).

Glutamate signals in awake mice also showed diverse odorant responses over

multiple inhalations, including slowly facilitating and adapting response patterns (Fig. 4c,

d). We also observed concentration-dependent changes in glutamate dynamics, with

increasing odorant concentrations leading to stronger adaptation of glutamate signal for

strongly-responsive glomeruli and recruitment of slowly-increasing responses in weakly-

responsive glomeruli (Fig. 4e). Consequently, the relative patterns of glutamate signals

across glomeruli changed from their initial pattern over repeated sniffs, and glutamate

response patterns decorrelated more substantially for some odorants than others (Fig.

4f). Overall, these data indicate that the diversity in dynamics of glutamatergic inputs to

MT cells observed in awake mice is qualitatively similar to that imaged with artificial

inhalation.

Correspondence between dynamics of glutamatergic inputs and MT cell postsynaptic

activity.

While glutamatergic signaling in the glomerulus is the sole source of excitatory

input to MT cells, inhibitory circuits and MT cell intrinsic properties may also shape

patterns of MT cell excitation[2,10,11,32,33]. To assess these relative contributions, we

compared glutamate signals with calcium signals measured from MT cell apical tufts of

the same glomerulus, imaged simultaneously. Because MT cell spikes back-propagate

along the MT cell primary dendrite and invade the apical tuft, calcium signals imaged

from the tuft provide a proxy for patterns of MT cell spike output from each glomerulus[34-

36].

We co-expressed SF-iGluSnFR.A184S and the red calcium reporter

jRGECO1a[37] in MT cells using viral co-injection into the OB of Tbet-Cre mice (Fig. 5a).

In response to odorant stimulation, jRGECO1a signals were apparent both in an

activated glomerulus as well as in the primary dendrites of the MT cells innervating it,

while SF-iGluSnFR signals were not apparent outside the glomerular neuropile (Fig. 5b,

e). As in our earlier report[30], we saw very high correspondence between odorant-

evoked GCaMP6f signals imaged from the soma or primary dendrite of MT cells and

those imaged from the neuropil of their parent glomerulus (Fig. 5b; Supplementary Fig.

3). In anesthetized mice, inhalation-linked transients were apparent in both signals at 1

Hz inhalation, although the jRGECO1a calcium signal decayed more slowly (Fig.5b).

Despite these kinetic differences, there was a close correspondence in the inhalation-

linked dynamics of the two signals, with different glomeruli showing the same relative

differences in the onset latency and time to peak of the SF-iGluSnFR and jRGECO1a

ITA waveform (Fig. 5c, d). We also observed a close - though not perfect -

correspondence in the occurrence of spontaneous transients in the SF-iGluSnFR and

jRGECO1a signals (Supplementary Fig. 2a). In awake mice, jRGECO1a signals

showed little or no inhalation-linked modulation despite strong inhalation-linked

transients in the SF-iGluSnFR signal (Fig. 5f).

To compare odorant-evoked response patterns over a time-scale involving

multiple inhalations, we first performed dual-color imaging in anesthetized mice using

the same 23 odorant panel as for the single-color imaging (5 fields of view from 3 mice).

Overall, there was a striking degree of correspondence in response patterns of the two

signals, with only rare exceptions (Fig. 6a; Supplementary Fig. 2b). There was high

concordance in glomeruli showing purely excitatory responses as measured with SF-

iGluSnFR and jRGECO1a (using the conservative criterion of ± 7 SD above baseline):

70% (273/387) of glomerulus-odor pairs with an excitatory SF-iGluSnFR signal also

showed a significant jRGECO1a response, and 79% (273/346) of glomeruli with

significant jRGECO1a increases showed excitatory SF-iGluSnFR signals

(Supplementary Fig. 2c). In awake mice, using a smaller odorant panel, only 46%

(57/123) of glomerulus-odor pairs with a significant excitatory SF-iGluSnFR signal also

showed jRGECO1a responses; however, 93% (57/61) of glomerulus-odor pairs with

excitatory jRGECO1a responses also showed SF-iGluSnFR responses. Temporal

response patterns were also highly correlated in awake mice (Fig. 6b; Supplementary

Fig. 2d); we quantified this using an index, T_2-T_1/T_{max}, that distinguished sustained,

rapidly adapting, and facilitating responses. T_2-T_1/T_{max} values were highly correlated for

glutamate and Ca^{2+} signals measured for the same glomerulus-odor pair, both in

anesthetized and awake mice (Fig. 6c). These results suggest that the evolution of MT

cell activity patterns across multiple inhalations largely follows that of glutamatergic

input to these cells.

To infer how inhibitory circuits might shape MT cell spike output from glomeruli,

we compared the prevalence of suppressive response components in the glutamate

and Ca^{2+} signals. In anesthetized mice, suppressive responses were rare, and their

prevalence may be underestimated from the relatively few repeat trials given (3 - 4 per

odorant) and the strict significance criteria we used. Nonetheless, jRGECO1a

responses with suppressive components were detected significantly more frequently

than SF-iGluSnFR responses, being present in 1.8% (72/3840) versus 0.7% (27/3840)

of all glomerulus-odor pairs, and 17% versus 7% of all significant responses (Chi-

Square statistic = 20.45, p = 6.0 e^{-6}). In awake mice, suppressive responses were more

prevalent (Supplementary Fig. 3d), with a similar prevalence of suppressive

components in both signals, being present in 21% (jRGECO1a: 92/446) versus 17%

(SF-iGluSnFR: 78/446) of all glomerulus-odor pairs (Chi-Square statistic = 1.15, p =

0.28). Notably, suppressive jRGECO1a responses were very rarely seen in glomeruli

showing excitatory SF-iGluSnFR signals, detected in only 2 of 72 suppressive

jRGECO1a responses in anesthetized mice (2.7%) and 5 of 92 responses (5.4%) in

awake mice; this prevalence is approaching the false positive rate of our measure of

response significance. This result is surprising given predictions from slice experiments

that weak excitatory input can lead to MT cell suppression by driving feedforward

inhibition[11,17,38]. However, these results are consistent with a model in which

interglomerular inhibition suppresses MT cell spiking in a small fraction of

glomeruli[30,39,40].

Limited contribution of multisynaptic pathways to odorant-evoked glutamate signaling

onto MT cells.

Sensory-evoked glutamatergic input onto MT cells can arise from OSNs or from

multisynaptic excitation involving dendritic glutamate release from ET cells and,

possibly, sTCs[16,29,41-43]. In addition, inhibitory circuits can modulate multisynaptic

glutamate signaling[11,12,17]. Glomerular glutamate signals may also arise from glutamate

release by MT cell dendrites[44,45], and thus at least partially reflect MT cell excitation

itself. To isolate the direct contribution of OSN inputs to MT cell excitation dynamics, we

compared evoked glutamate dynamics before and after pharmacological blockade of

postsynaptic activity with ionotropic glutamate receptor antagonists[46-48]. Because

iGluSnFRs are insensitive to these antagonists[19,20], we reasoned that this approach

would largely prevent OSN-driven excitation of postsynaptic neurons and allow us to

image evoked MT cell glutamate signals arising solely from OSN inputs.

To confirm postsynaptic blockade, we tested the ability of APV+NBQX to block

activation of CCK-expressing sTCs, as these, like ET cells, are strongly driven by

monosynaptic OSN input[29]. We imaged activation of CCK+ OB neurons in CCK-IRES-

Cre mice crossed to a Cre-dependent GCaMP6f reporter line. We first applied the

GABA$_B$ receptor antagonist CGP35348 (1 mM) to the dorsal OB to further enhance

transmitter release from OSNs by removing presynaptic inhibition, as done

previously[23,49]. As expected, CGP35348 led to a modest (20%) increase in the peak

CCK+ GCaMP6f response (Fig. 7b). Subsequent application of APV+NBQX (1 mM/0.5

mM) had mixed effects in different glomeruli, with responses almost completely

eliminated in 8 of 14 glomeruli (96% mean reduction in peak response, 3 experiments;

see Fig. 7c legend for summary statistics) but only partially reduced in the remaining 6

(46% mean reduction) (Fig. 7b - d). We performed additional experiments using a 5x

higher concentration of antagonist (5 mM APV/2.5 mM NBQX) and without

preapplication of CGP35348. In these, APV+NBQX completely blocked postsynaptic

activation in 11 of 13 glomeruli (from 3 experiments), with strongly reduced responses in

the remaining two (Fig. 7e).

Importantly, in glomeruli with persistent (but weakened) responses after

APV+NBQX application, inhalation-linked glutamate transients were significantly

delayed compared to the predrug condition (Fig. 7c, k), with a median increase in onset

latency of 285 ms (range: 164 – 667 ms) and median increase in time to peak of 247 ms

(range: 211 – 567 ms). An explanation for this substantial delay is that, in glomeruli

receiving the strongest sensory input, glutamate concentrations are eventually able to

overcome APV+NBQX blockade sufficiently to trigger spike bursts in sTCs. In additional

experiments imaging GCaMP6f signals from the general population of Tbet-expressing

MT cells, APV+NBQX eliminated inhalation-linked transients in glomeruli, with only a

small-amplitude tonic glutamate signal remaining (Supplementary Fig. 4 a,b), possibly

mediated by metabotropic glutamate receptors. These results demonstrate that

APV+NBQX blocks postsynaptic activation in glomeruli receiving all but the strongest

OSN inputs, and even in those glomeruli, substantially weakens and delays activation of

monosynaptically-driven TCs.

We next used APV+NBQX blockade to test the contribution of mono- versus di-

or polysynaptic glutamate signaling to inhalation-linked glutamate transients onto pcMT

cells. We first tested effects of CGP35348 and subsequent application of APV+NBQX (1

mM/0.5 mM) on glutamate transients using the medium-affinity SF-iGluSnFR.A184V.

Inhalation-linked glutamate transients were increased by CGP35348, confirming earlier

results[23]. Surprisingly, subsequent application of APV+NBQX had no impact on

response amplitudes (Fig. 7f, g). Importantly, GABA$_B$ receptor or iGluR blockade had

little to no effect on inhalation-linked glutamate dynamics (Fig. 7f, k). We obtained

similar results in epifluorescence experiments imaging iGluSnFR signals from a Tbet-

expressing MT cell population across a larger area of the dorsal OB, using

epifluorescence imaging (Supplementary Fig. 4d-f).

Multisynaptic glutamatergic excitation may preferentially drive MT cell responses

to weak inputs[17,43,50], and the odorant concentrations used in the preceding experiments

were relatively high (20 – 50 ppm see Table 1). Removal of presynaptic inhibition with

CGP35348 may also enhance OSN input strength to unnaturally high levels. Thus, we
next tested the impact of glutamate receptor blockade on inhalation-linked transients in
pcMT cell responses to odorants presented at 10 – 100x lower concentrations (0.1 – 5
ppm), using a higher concentration of APV+NBQX (5 mM/2.5 mM) and omitting
CGP35348. Here, iGluR blockade was weakly effective at reducing the magnitude of
inhalation-linked glutamate transients, causing a significant reduction in two of four
experiments and a mean reduction of 32 ± 12% overall (mean ± s.e.m. of median ratios
across 4 experiments, n = 8 – 16 glomerulus-odor pairs per experiment) (Fig. 7i, j).
However, APV+NBQX had no impact on the dynamics of inhalation-linked glutamate
responses, with no change in peak times in any experiment (Fig. 7k).
Finally, to examine the role of direct versus multisynaptic excitation in shaping
diverse temporal patterns of glutamatergic input to MT cells, we used a 12-odorant
panel (delivered concentrations ranging from 0.03 - 15 ppm, see Table 1) and 3 Hz
artificial inhalation, comparing response patterns before and after application of
APV+NBQX (1 mM/0.5 mM). This approach yielded significant odorant responses in 83
glomerulus-odorant pairs across 3 mice, and included diverse excitatory response
patterns (Fig. 8a). Consistent with the earlier experiments, APV+NBQX had little overall
impact on the magnitude of evoked iGluSnFR signals: while there was some variability
in the effect for particular glomerulus-odor pairs, neither mean nor peak excitatory
response magnitudes showed a significant overall change after APV+NBQX application
(paired t-tests, p = 0.147 for mean amplitude, p = 0.605 for peak amplitude, n = 83) (Fig.
8b).
APV+NBQX did, however, impact the temporal patterns of the glutamate signal
for a subset of responses. In particular, APV+NBQX tended to reduce adaptation that
occurred following the initial inhalation of odorant or enhance longer-lasting, tonic-type
responses (Fig. 8a). This was reflected as an increase in T_2-T_1/T_{max} values by
APV+NBQX, with significant increases seen in each of three mice (Fig. 8c). Taken
together, these results suggest that multisynaptic glutamatergic signaling contributes
rather subtly to odorant-evoked excitation of MT cells, slightly enhancing weak
inhalation-linked glutamate transients but also contributing to adaptation of glutamate
signals during repeated odorant sampling.

DISCUSSION

Temporally complex patterns of excitation and inhibition among principal cells

(MT cells) of the OB are hypothesized to play important roles in coding olfactory

information, and have been attributed to emergent properties of excitatory and inhibitory

circuit interactions[4-6,30,33]. By directly imaging glutamate signaling onto MT cell dendrites

of OB glomeruli, we observed a striking degree of heterogeneity in this signaling both on

the time-scale of a single respiratory cycle and over a slower time scale involving

repeated samples of odorant. Simultaneous imaging of presynaptic glutamate and

postsynaptic calcium from MT cells in the same glomerulus showed a high

correspondence in the dynamics of glutamatergic input MT output. Pharmacological

blockade of postsynaptic activity had little impact on glutamate dynamics, implicating

direct inputs from OSNs as a primary source of this diversity. These results suggest a

model of OB circuit function in which the dynamics of sensory input to glomeruli can

account for much of the diversity of MT cell responses that underlies odor

representations in vivo.

The iGluSnFR sensors are well-suited to characterizing glutamatergic signaling

in the OB in vivo, with intrinsic kinetics (rise and decay times of <10 to ~ 20 ms,

respectively[19,51,52]) that are much faster than the dynamics of odorant-evoked excitatory

input to MT cells (EPSP rise- and decay times of ~100 ms and 200 ms, respectively[53]).

iGluSnFR expression can prolong the decay of glutamate transients by competing with

glutamate transporters[54], and decay rates are slower (~140 ms) for the high-affinity SF-

iGluSnFR.A184S [20,55]. While we did find that decay times skewed longer for ITA

responses measured with the high-affinity variant, their distribution largely overlapped

with that of the medium-affinity A184V. Furthermore, the degree of respiratory

modulation as well as overall diversity in slow iGluSnFR signal dynamics appeared

similar for the two variants. Thus, the SF-iGluSnFR signal imaged from MT cell apical

tufts is a reliable reporter of the variable dynamics of glutamate signaling in the

glomerulus.

Another potential confound to interpreting the iGluSnFR signal is dendritic

release of glutamate from MT cells themselves[44,45]. However, near-complete

pharmacological blockade of MT cell activation did not alter ITA waveforms, suggesting
that glutamate signals imaged from OB glomeruli reflect glutamatergic input to, as
opposed to glutamate release from, MT cells.

OSN inputs to the OB respond with different latencies relative to inhalation in a
glomerulus- and odorant-specific manner[15,24,56], and we found a close match in the
range of response latencies for SF-iGluSnFR signals and those measured from OSN
presynaptic terminals using Ca^{2+} reporters. We also saw variation in other aspects of
inhalation-linked glutamate dynamics that are not predicted from presynaptic Ca^{2+}
imaging, with the duration of inhalation-linked glutamate transients ranging from 200 ms
to over 1.5 sec. Glutamatergic signaling onto MT cells also showed complex dynamics
over a slower time scale spanning multiple inhalations that could include adaptation,
facilitation, or biphasic response patterns. This diversity likely has substantial impacts
on the temporal dynamics of MT cell spiking, and suggests that MT cell excitation is not
a simple linear combination of responses to a single inhalation[57].

Several results emerged from imaging glutamate dynamics in awake mice that
inform models of how olfactory information is represented during natural odor sampling.
First, the relative magnitudes of glutamate signal across different glomeruli changed
over the course of an odor presentation, suggesting that odorant representations evolve
rapidly during repeated sampling of odorant. This result is consistent with recent
findings of dynamic odor representations on a similar time scale at the level of MT cell
spiking[3,5]. Second, we found that a substantial fraction of odorant-evoked glutamate
signals showed minimal or no respiratory patterning, instead showing tonic increases in
glutamate when odorant was present. Assuming a similar fraction of MT cells fail to
show respiratory patterning in their spike timing in awake mice, this result has
implications for theories of odor identity or intensity coding that are based on the timing
of MT cell spikes within the breathing cycle[4,7-9]. Third, in both awake and in anesthetized
mice, we found that the temporal evolution of glutamate responses as well as the
degree of inhalation-linked patterning varies with odorant identity and concentration as
well as glomerular identity. Comparable richness in response kinetics have been well-
characterized in Drosophila OSNs, where such features are attributed to odorant
receptor identity[58]. In mice, further experiments are necessary to understand the

determinants of these glutamate dynamics - for example, whether they are determined

by the affinity and intensity of an odorant relative to the dynamic range of a particular

glomerulus-odorant pair, or instead by properties of odorant sorption in the olfactory

epithelium.

Finally, we observed a very close (but not perfect) correspondence between the

dynamics of glutamatergic input to MT cells and those of postsynaptic Ca^{2+} signals

reflecting MT cell spiking, imaged simultaneously. These results suggest that the

dynamics of MT cell spiking patterns closely follow those of glutamatergic input to their

glomerulus. Interestingly, we saw no evidence for broadly-organized inhibition

surrounding glomeruli receiving excitatory inputs, as predicted by models of center-

surround or global inhibition[18,59]. We also saw little evidence for suppression of MT cell

output in glomeruli receiving weak excitatory inputs, as predicted from models of

feedforward inhibition[11,17,38]. Instead, suppressive MT cell responses were sparse and

often co-occurred with a reduction in the glutamate signal to below baseline levels,

indicating that MT cell suppression commonly reflects a reduction in excitatory drive.

This reduction could be could be due to adaptation of OSN inputs[48], inhibition at the

receptor level[60,61], or inhibition of ongoing glutamatergic drive from ET cells[11,12]. Overall,

the high correspondence between glutamatergic input and MT cell output patterns

suggests that transformations of odor representations by OB circuitry- at least in the

spatial and temporal domains - may be more limited than previously thought. Instead,

inhibitory circuits may function to shape response features such as sensitivity and

response gain by acting coincident with excitation. Such a model would be more similar

to the function of many inhibitory circuits in cortex, which are reliant on and closely

coupled to excitation[62].

Another surprising aspect of our results is the relatively small impact of blocking

postsynaptic activity on glutamate signaling onto MT cells. Studies from OB slices

suggest that ET cells provide a large fraction of excitatory drive to MT cells via

glutamate spillover from the ET- to - periglomerular cell synapse[16,17,44,50]. Extrapolating

to OB circuit function in vivo is difficult, however – for example, glutamate spillover may

contribute less to MT cell drive in vivo than in OB slices due to differences in glutamate

transporter efficacy or in the dynamics of odorant-evoked glutamate release from OSN

inputs. We found that a near-complete blockade of postsynaptic activation, as confirmed

with GCaMP reporters expressed in monosynaptically-driven tufted cells, had no impact

on the dynamics of inhalation-linked glutamate transients and only modestly impacted

the amplitude of transients in response to low odorant concentrations, as well having

rather subtle effects on responses across inhalations.

Our results are consistent with some models of ET cell-mediated excitation of MT

cells, which propose that the ET cell pathway is most important in regimes of weak OSN

input (i.e., low odorant concentrations)[17,43,50], or that direct OSN inputs are shunted by

gap junctions between MT cells[41], which would not be reflected in our iGluSnFR

recordings. Nonetheless, the overall persistence of glutamate signals after postsynaptic

blockade, as well as the strong correspondence between glutamate dynamics and MT

cell activation, suggests that monosynaptic drive from OSNs is the primary determinant

of MT cell response patterns across a large regime of odorant stimulation and natural

odor sampling. An alternate possibility is that the disynaptic circuit provides tonic

glutamatergic drive to MT cells via ET cell bursting in vivo, allowing for modulation of

MT cell excitability by inhibitory circuits. Further use of iGluSnFRs and other transmitter-

specific reporters could help disentangle the contributions of excitation and inhibition to

shaping dynamic odor representations in vivo.

MATERIALS AND METHODS

Animals

Experiments were performed on male and female mice expressing Cre recombinase

(Cre) in defined neural populations. Mouse strains used were: Pcdh21-Cre (Tg(Pcdh21-

cre)BYoko), Gensat Stock #030952-UCD; OMP-Cre (Tg(Omp-tm4-Cre)Mom), JAX

Stock #006668, Tbet-Cre (Tg(Tbx21-cre)1Dlc), JAX Stock #024507, and CCK-IRES-

Cre (Tg(CCK-IRES-Cre)Zjh), JAX Stock #012706[22]. Mice ranged from 3-8 months in

age. Mice were housed up to 4/cage and kept on a 12/12 h light/dark cycle with food

and water available ad libitum. All procedures were carried out following the National

Institutes of Health Guide for the Care and Use of Laboratory Animals and were

approved by the University of Utah Institutional Animal Care and Use Committee.

Viral vector expression

Viral vectors were obtained from the University of Pennsylvania Vector Core (AAV1 or 5 serotype, AAV.hSynap-FLEX.iGluSnFR and AAV.hSynap-FLEX.jRGECO1a), Addgene (AAV1 serotype, pAAV.hSynap-FLEX.SF-iGluSnFR.S72A, #106182), HHMI Janelia Campus or Vigene (AAV1 or 5 serotype, pAAV.hSynap-FLEX.SF-iGluSnFR.A184V, pAAV.hSynap-FLEX.SF-iGluSnFR.A184S). Virus injection was done using pressure injections and beveled glass pipettes, as described previously[24,31,63]. For coinjection of jRGECO1a and SF-iGluSnFR.A184S, virus was either diluted 1:10 and injected in separate pipettes through the same craniotomy, or mixed and coinjected via the same pipette. After injection, mice were given carprofen (Rimadyl, S.C., 5 mg/kg; Pfizer) as an analgesic and enrofloxacin (Baytril, I.M., 3 mg/kg; Bayer) as an antibiotic immediately before and 24 hours after surgery. Mice were singly housed after surgery on ventilated racks and used 21-35 days after virus injection. In some mice, viral expression was characterized with post-hoc histology using native fluorescence.

In vivo two photon imaging

Two-photon imaging in anesthetized mice was performed as described previously[30,63]. Mice were initially anesthetized with pentobarbital (50 - 90 mg/kg) then maintained under isoflurane (0.5 – 1% in O_2) for data collection. Body temperature and heart rate were maintained at 37 $^{\circ}$C and ~ 400 beats per minute. Mice were double tracheotomized and isoflurane was delivered passively via the tracheotomy tube without contaminating the nasal cavity[64]. Two-photon imaging occurred after removal of the bone overlying the dorsal olfactory bulb and stabilizing the brain surface with agarose and a glass coverslip.

Imaging in awake, head-fixed mice was performed through a chronic imaging window implanted over one dorsal OB. The imaging window consisted of a custom double coverslip with minimum diameter of 1.5 mm. Virus (500-750 nL) was injected at a depth of 250 µm at the time of window implant. Animals were acclimated to the imaging rig for at least one 30-minute session the day prior to imaging. Imaging sessions lasted from 30-60 minutes per day, over the course of several days. The behavior and imaging apparatus has been described previously[30]. Mice were naïve to

the odors at their first imaging session. Respiration (sniff) signals were obtained with an

external flow sensor (FBAM200DU, First Sensor, AG, Berlin) in front of the animal's

right nostril[65], or by using a thermistor (MEAS-G22K7MCD419, Measurement

Specialties; Hampton, VA) implanted in the right nasal bone[66].

Imaging was carried out with a two-photon microscope (Sutter Instruments or

Neurolabware) coupled to a pulsed Ti:Sapphire laser (Mai Tai HP, Spectra-Physics; or

Chameleon Ultra, Coherent) at 920-940 nm and controlled by either Scanimage (Vidrio)

or Scanbox (Neurolabware) software. Imaging was performed through a 16X, 0.8 N.A.

objective (Nikon) and emitted light detected with GaAsP photomultiplier tubes

(Hamamatsu). Fluorescence images were acquired using unidirectional resonance

scanning at 15.2 or 15.5 Hz. For SF-iGluSnFR.S72A imaging, bidirectional scanning at

30Hz was utilized to capture faster responses. For dual-color imaging, a second laser

(Fidelity-2; Coherent) was utilized to optimally excite jRGECO1a (at 1070nm) and

emitted red fluorescence collected with a second PMT, as described previously[24].

In vivo pharmacology

In vivo pharmacology was carried out after removing the bone and dura overlying

the dorsal olfactory bulb, using protocols described previously[23,46]. Drug solutions (1mM

CGP35348 and 0.5 mM NBQX / 1 mM APV) were dissolved in Ringers solution, pre-

warmed on a heating block, and applied in bulk to the dorsal OB without the use of

agarose or coverslip. For higher concentration experiments, we utilized 2.5 mM NBQX /

5 mM APV dissolved in Ringers. We waited at least 10 minutes after drug application to

allow for absorption and temperature equilibration across the tissue. NBQX + APV was

applied immediately after CGP35348 without an intervening control wash; due to its

extremely slow washout in vivo, we considered CGP35348 to still be present during the

subsequent NBQX + APV application. For statistical analysis of pharmacology

experiments, all comparisons were done within-animal unless otherwise noted.

Odorant stimulation

In some experiments, odorants were presented as precise dilutions from

saturated vapor (s.v.) in cleaned, desiccated air using a custom olfactometer under

computer control, as described previously[30,67]. Odorants were presented for durations

ranging from 2 - 8 sec. Clean air was passed across the nostrils in between trials to

avoid contribution from extraneous odorants in the environment. Odorants were pre-

diluted in solvent (1:10 or 1:25 in mineral oil or medium chain triglyceride oil) to allow for

lower final concentrations and then diluted to concentrations ranging from .3% to 1%

s.v. Relative increases in concentration were confirmed with a photoionization detector

(miniRAE Lite, PGM-7300, RAE Systems) 3 cm away from the flow dilution port.

Estimated final concentrations of odorants used ranged from 0.01 – 300 ppm,

depending on vapor pressure and s.v. dilution (Table 1). For experiments testing a

larger panel of odorants (e.g., Figs. 3 and 5, see Table 1), we used a novel, recently

described olfactometer design that allowed for rapid switching between odorants with

minimal cross-contamination[68]. Here, odorants were presented for 2 or 3 seconds, in

random order from among a bank of 12 odorant cartridges, using 10 second

interstimulus intervals. Odorant presentation was as described in a previous

publication[68], using an eductor nozzle for optimal mixing in a carrier stream of filtered

air. The end of the eductor was placed 5-7 cm away from the nose. With the

configuration used, estimated dilutions of odorant were approximately 1.5% s.v.;

odorants were prediluted to achieve relatively sparse activation of dorsal glomeruli[68].

Estimated final concentrations ranged from 0.01 – 16 ppm (Table 1).

Data Analysis

Image analysis was performed using custom software written in MATLAB

(Mathworks). For display, odorant response maps were displayed using ΔF values

rather than ΔF/F to minimize noise from nonfluorescent regions. Activity maps were

scaled as indicated in the Fig. and were kept to their original resolution (512 x 512

pixels) and smoothed using a Gaussian kernel with σ of 1 pixel. For time-series

analysis, regions of interest (ROIs) were chosen manually based on the mean

fluorescence image and were further refined based on odorant response ΔF maps, then

all pixels averaged within an ROI. All signals were upsampled to 150 Hz for analysis

using the MATLAB pchip function. Time-series were typically computed and displayed

as ΔF/F, with F defined as the mean fluorescence in the 1-2 sec prior to odorant onset.

591   For analysis of inhalation-linked dynamics, inhalation-triggered average (ITA)

592 responses were generated by averaging each inhalation (delivered at 0.25 Hz) over a

593 70 sec odorant presentation (17 inhalations averaged in total). Onset latencies, peak

594 responses, and ITA durations (full-width at half-maximum) were defined as previously[24].

595 To calculate decay time-constants (Fig. 1), a single exponential was fitted from the peak

596 of the unfiltered ITA trace and extending to the end of the trace. Time-to-peak values

597 were calculated from unfiltered ITAs as the time from the inhalation start to the peak of

598 the response.

599   For analysis of odorant-evoked dynamics over multiple sniffs, responses to 3 - 8

600 presentations of odorant were averaged before analysis. Changes in response

601 amplitude over time, or T_2-T_1/T_{max} (Fig. 6) were calculated as the difference in amplitude

602 between the peak ΔF following the first (T_1) and the second-to-last (T_2) inhalation during

603 odorant presentation, divided by the maximum ΔF during the 4 sec odor presentation.

604   To analyze response patterns across the 23-odorant panel (i.e., Fig. 3),

605 responses were averaged across 3 - 4 randomized presentations of each odorant.

606 Responses were classified as having significant excitatory and/or suppressive

607 components as follows: First, each averaged response for an ROI-odor pair was low-

608 pass Gaussian filtered at 2 Hz (to test for excitatory responses) or 0.5 Hz (for

609 suppressive responses) and z-scored using a baseline from 0.5 to 2 sec prior to odorant

610 onset, with z defined as the SD of the baseline period concatenated for all 23 odorant

611 responses for each ROI. Peak excitation was measured as 95% of the maximum signal

612 during the 3- or 4-sec odorant presentation; suppressive responses were measured as

613 the 15th percentile of all values in a time-window from odorant onset to 500 ms after

614 odorant offset. We then used a very conservative criterion for significance of z = $\pm$7 SD

615 for identifying significant excitatory or suppressive response components. This cutoff

616 was chosen to yield a false positive rate of approximately 1% (based on visual

617 inspection of a subset of traces).

618   Glomerular tuning was characterized from response matrices thresholded

619 according to these significance values, using a measure of lifetime sparseness[69]

620 $S = \{1 - [(\sum N_{j=1} r_j / N)^2 / \sum N_{j=1} (r_j^2 / N)]\} / [1 - (1/N)]$

Where N is number of odors and r_j is the thresholded z-scored value for a given

glomerulus of the response to odor j. Sparseness was calculated separately for

excitatory and suppressive response components.

Time-dependent decorrelation of odorant response patterns across glomeruli

(Figs. 3 and 4) was calculated as previously described [6], using all glomeruli in a field of

view without thresholding. Each odorant response was binned into five time bins (each

of 387 ms) spanning the length of odorant presentation but omitting the first 65 ms to

account for delays in odorant onset. Pearson's correlation coefficient (*r*) was calculated

using the response amplitudes across all ROIs in the first time bin as a vector,

compared to the same vector taken from each subsequent bin. We calculated an overall

correlation time series by computing the mean and s.e.m. for all odorant responses,

pooled across all fields of view.

For analysis of inhalation-linked timing of glutamate signals in awake mice, we

restricted analysis to inhalations occurring at a minimum period of 200 ms (5 Hz

respiration rate). Inhalation timing was detected from the external flow sensor using

peak/trough detection and maximal slopes of the flow signal. Inhalation peak was

defined as the trough of the external flow signal; inhalation onset was defined as the

maximal slope of the transition from exhalation to inhalation. Because inhalation peak

timing was a more robust measure, we generated inhalation-triggered average

iGluSnFR signals based on inhalation peak. ITAs were generated by averaging the

unfiltered optical signal taken from -0.3 sec before to 0.4 sec after inhalation. To avoid

confounds of the airflow waveform or measurement method (airflow versus thermistor)

in determining absolute ITA response latencies, we report the range of latencies relative

to the median of a given set of measurements. Statistical tests were performed either in

Origin (OriginLab Corp.), MATLAB (Mathworks) or R (version 3.3.2). Nonparametric

tests were used in most cases; parametric tests were used only on datasets determined

to be normally distributed. Summary statistics are reported as mean ± standard

deviation (SD), unless otherwise stated. All measurement of response parameters was

done using analysis code that was independent of treatment or comparison condition.

ACKNOWLEDGEMENTS

We thank L. Looger and J. Marvin and the GENIE Project from HHMI Janelia Research

Campus for sharing SF-iGluSnFR virus. We also thank M. Economo, D. Brunert, Y.

Tsuno and F. Fang for initial characterization of iGluSnFRs, G. Vasquez-Opazo, R.

Kummer and J. Ball for technical assistance, T. Rust for data analysis software, as well

as S. Burton, S. Short, I. Youngstrom, and K. Podgorski for providing helpful feedback

and discussion. This work was supported by the NINDS T32NS076067 (to A.K.M) and

NIDCD R01DC006441 (to M.W.), R01DC006441-13S1 (to A.K.M), and F32DC015389

(to T.P.E).

COMPETING INTERESTS

These authors declare no competing financial interests.

DATA AVAILABILITY

Data underlying the findings of this study are available from the corresponding author

upon request.

CODE AVAILABILITY

Custom code (Matlab) used in data analysis are available from the corresponding

author upon request.

CONTRIBUTIONS

A.K.M and M.W. designed research. A.K.M, T.P.E, and M.W. performed

research, A.K.M., T.P.E., and M.W. analyzed data, A.K.M. and M.W. wrote the paper.

REFERENCES

1. Vassar, R. *et al.* Topographic organization of sensory projections to the olfactory bulb. *Cell* **79**, 981-991 (1994).
2. Wachowiak, M. & Shipley, M.T. Coding and synaptic processing of sensory information in the glomerular layer of the olfactory bulb. *Semin Cell Dev Biol* **17**, 411-423 (2006).
3. Diaz-Quesada, M. *et al.* Inhalation Frequency Controls Reformatting of Mitral/Tufted Cell Odor Representations in the Olfactory Bulb. *J Neurosci* **38**, 2189-2206 (2018).
4. Chaput, M.A. Respiratory-phase-related coding of olfactory information in the olfactory bulb of awake freely-breathing rabbits. *Physiol Behav* **36**, 319-324 (1986).
5. Patterson, M.A., Lagier, S. & Carleton, A. Odor representations in the olfactory bulb evolve after the first breath and persist as an odor afterimage. *Proc Natl Acad Sci U S A* **110**, E3340-3349 (2013).
6. Eiting, T.P. & Wachowiak, M. Differential Impacts of Repeated Sampling on Odor Representations by Genetically-Defined Mitral and Tufted Cell Subpopulations in the Mouse Olfactory Bulb. *J Neurosci* **40**, 6177-6188 (2020).
7. Schaefer, A.T. & Margrie, T.W. Spatiotemporal representations in the olfactory system. *Trends Neurosci* **30**, 92-100 (2007).
8. Wilson, C.D., Serrano, G.O., Koulakov, A.A. & Rinberg, D. A primacy code for odor identity. *Nat Commun* **8**, 1477 (2017).
9. Li, A., Rao, X., Zhou, Y. & Restrepo, D. Complex neural representation of odour information in the olfactory bulb. *Acta physiologica*, e13333 (2019).
10. Schoppa, N.E. & Urban, N.N. Dendritic processing within olfactory bulb circuits. *Trends Neurosci* **26**, 501-506 (2003).
11. Gire, D.H. & Schoppa, N.E. Control of on/off glomerular signaling by a local GABAergic microcircuit in the olfactory bulb. *J Neurosci* **29**, 13454-13464 (2009).

12.   Shao, Z., Puche, A.C., Liu, S. & Shipley, M.T. Intraglomerular inhibition Shapes
the Strength and Temporal Structure of Glomerular Output. *J Neurophysiol*
(2012).

13.   Geramita, M. & Urban, N.N. Differences in Glomerular-Layer-Mediated
Feedforward Inhibition onto Mitral and Tufted Cells Lead to Distinct Modes of
Intensity Coding. *J Neurosci* **37**, 1428-1438 (2017).

14.   Verhagen, J.V., Wesson, D.W., Netoff, T.I., White, J.A. & Wachowiak, M. Sniffing
controls an adaptive filter of sensory input to the olfactory bulb. *Nat Neurosci* **10**,
631-639 (2007).

15.   Spors, H., Wachowiak, M., Cohen, L.B. & Friedrich, R.W. Temporal dynamics
and latency patterns of receptor neuron input to the olfactory bulb. *J Neurosci* **26**,
1247-1259 (2006).

16.   De Saint Jan, D., Hirnet, D., Westbrook, G.L. & Charpak, S. External Tufted Cells
Drive the Output of Olfactory Bulb Glomeruli. *J. Neurosci.* **29**, 2043-2052 (2009).

17.   Gire, D.H., Zak, J.D., Bourne, J.N., Goodson, N.B. & Schoppa, N.E. Balancing
Extrasynaptic Excitation and Synaptic Inhibition within Olfactory Bulb Glomeruli.
eNeuro **6** (2019).

18.   Banerjee, A. *et al.* An Interglomerular Circuit Gates Glomerular Output and
Implements Gain Control in the Mouse Olfactory Bulb. *Neuron* **87**, 193-207
(2015).

19.   Marvin, J.S. *et al.* An optimized fluorescent probe for visualizing glutamate
neurotransmission. *Nat Meth* **10**, 162-170 (2013).

20.   Marvin, J.S. *et al.* Stability, affinity, and chromatic variants of the glutamate
sensor iGluSnFR. *Nat Methods* **15**, 936-939 (2018).

21.   Nagai, Y., Sano, H. & Yokoi, M. Transgenic expression of Cre recombinase in
mitral/tufted cells of the olfactory bulb. *genesis* **43**, 12-16 (2005).

22.   Haddad, R. *et al.* Olfactory cortical neurons read out a relative time code in the
olfactory bulb. *Nat Neurosci* **16**, 949-957 (2013).

23.   Brunert, D., Tsuno, Y., Rothermel, M., Shipley, M.T. & Wachowiak, M. Cell-Type-
Specific Modulation of Sensory Responses in Olfactory Bulb Circuits by

Serotonergic Projections from the Raphe Nuclei. *J Neurosci* **36**, 6820-6835
(2016).

24. Short, S.M. & Wachowiak, M. Temporal Dynamics of Inhalation-Linked Activity
across Defined Subpopulations of Mouse Olfactory Bulb Neurons Imaged In
Vivo. *eNeuro* **6**, ENEURO.0189-0119.2019 (2019).

25. Richter, F.G., Fendl, S., Haag, J., Drews, M.S. & Borst, A. Glutamate Signaling in
the Fly Visual System. *iScience* **7**, 85-95 (2018).

26. Igarashi, K.M. *et al.* Parallel Mitral and Tufted Cell Pathways Route Distinct Odor
Information to Different Targets in the Olfactory Cortex. *The Journal of*
Neuroscience **32**, 7970-7985 (2012).

27. Fukunaga, I., Berning, M., Kollo, M., Schmaltz, A. & Schaefer, Andreas T. Two
Distinct Channels of Olfactory Bulb Output. *Neuron* **75**, 320-329 (2012).

28. Seroogy, K.B., Brecha, N. & Gall, C. Distribution of cholecystokinin-like
immunoreactivity in the rat main olfactory bulb. *J Comp Neurol* **239**, 373-383
(1985).

29. Sun, X., Liu, X., Starr, E.R. & Liu, S. CCKergic Tufted Cells Differentially Drive
Two Anatomically Segregated Inhibitory Circuits in the Mouse Olfactory Bulb. *J*
Neurosci **40**, 6189-6206 (2020).

30. Economo, M.N., Hansen, K.R. & Wachowiak, M. Control of Mitral/Tufted Cell
Output by Selective Inhibition among Olfactory Bulb Glomeruli. *Neuron* (2016).

31. Rothermel, M., Brunert, D., Zabawa, C., Díaz-Quesada, M. & Wachowiak, M.
Transgene Expression in Target-Defined Neuron Populations Mediated by
Retrograde Infection with Adeno-Associated Viral Vectors. *The Journal of*
Neuroscience **33**, 15195-15206 (2013).

32. Burton, S.D. Inhibitory circuits of the mammalian main olfactory bulb. *J*
Neurophysiol **118**, 2034-2051 (2017).

33. Fukunaga, I., Herb, J.T., Kollo, M., Boyden, E.S. & Schaefer, A.T. Independent
control of gamma and theta activity by distinct interneuron networks in the
olfactory bulb. *Nat Neurosci* **17**, 1208-1216 (2014).

34. Charpak, S., Mertz, J., Beaurepaire, E., Moreaux, L. & Delaney, K. Odor-evoked
calcium signals in dendrites of rat mitral cells. *Proc Natl Acad Sci U S A* **98**,
1230-1234 (2001).
35. Bischofberger, J. & Jonas, P. Action potential propagation into the presynaptic
dendrites of rat mitral cells. *J Physiol* **504 (Pt 2)**, 359-365 (1997).
36. Chen, W.R., Midtgaard, J. & Shepherd, G.M. Forward and backward propagation
of dendritic impulses and their synaptic control in mitral cells. *Science* **278**, 463-
467 (1997).
37. Dana, H. *et al.* Sensitive red protein calcium indicators for imaging neural activity.
eLife **5**, e12727 (2016).
38. Cleland, T.A. & Linster, C. On-Center/Inhibitory-Surround Decorrelation via
Intraglomerular Inhibition in the Olfactory Bulb Glomerular Layer. *Front Integr*
Neurosci **6**, 5 (2012).
39. Whitesell, J.D., Sorensen, K.A., Jarvie, B.C., Hentges, S.T. & Schoppa, N.E.
Interglomerular lateral inhibition targeted on external tufted cells in the olfactory
bulb. *J Neurosci* **33**, 1552-1563 (2013).
40. Liu, S., Puche, A.C. & Shipley, M.T. The Interglomerular Circuit Potently Inhibits
Olfactory Bulb Output Neurons by Both Direct and Indirect Pathways. *J Neurosci*
36, 9604-9617 (2016).
41. Gire, D.H. *et al.* Mitral Cells in the Olfactory Bulb Are Mainly Excited through a
Multistep Signaling Path. *The Journal of Neuroscience* **32**, 2964-2975 (2012).
42. Hayar, A., Karnup, S., Ennis, M. & Shipley, M.T. External tufted cells: A major
excitatory element that coordinates glomerular activity. *J. Neurosci.* **24**, 6676-
6685 (2004).
43. Vaaga, C.E. & Westbrook, G.L. Parallel processing of afferent olfactory sensory
information. *J Physiol* **594**, 6715-6732 (2016).
44. Isaacson, J.S. Glutamate spillover mediates excitatory transmission in the rat
olfactory bulb. *Neuron* **23**, 377-384 (1999).
45. Najac, M. *et al.* Intraglomerular lateral inhibition promotes spike timing variability
in principal neurons of the olfactory bulb. *J Neurosci* **35**, 4319-4331 (2015).

46. Pírez, N. & Wachowiak, M. In vivo modulation of sensory input to the olfactory
bulb by tonic and activity-dependent presynaptic inhibition of receptor neurons. *J*
Neurosci **28**, 6360-6371 (2008).
47. Gurden, H., Uchida, N. & Mainen, Z.F. Sensory-evoked intrinsic optical signals in
the olfactory bulb are coupled to glutamate release and uptake. *Neuron* **52**, 335-
345 (2006).
48. Lecoq, J., Tiret, P. & Charpak, S. Peripheral adaptation codes for high odor
concentration in glomeruli. *J Neurosci* **29**, 3067-3072 (2009).
49. Wachowiak, M. *et al.* Inhibition of olfactory receptor neuron input to olfactory bulb
glomeruli mediated by suppression of presynaptic calcium influx. *J Neurophysiol*
94, 2700-2712 (2005).
50. Najac, M., De Saint Jan, D., Reguero, L., Grandes, P. & Charpak, S.
Monosynaptic and Polysynaptic Feed-Forward Inputs to Mitral Cells from
Olfactory Sensory Neurons. *The Journal of Neuroscience* **31**, 8722-8729 (2011).
51. Armbruster, M., Hanson, E. & Dulla, C.G. Glutamate Clearance Is Locally
Modulated by Presynaptic Neuronal Activity in the Cerebral Cortex. *The Journal*
of Neuroscience **36**, 10404-10415 (2016).
52. Pinky, N.F., Wilkie, C.M., Barnes, J.R. & Parsons, M.P. Region- and Activity-
Dependent Regulation of Extracellular Glutamate. *The Journal of Neuroscience*
38, 5351-5366 (2018).
53. Cang, J. & Isaacson, J.S. In Vivo Whole-Cell Recording of Odor-Evoked Synaptic
Transmission in the Rat Olfactory Bulb. *J. Neurosci.* **23**, 4108-4116 (2003).
54. Armbruster, M., Dulla, C.G. & Diamond, J.S. Effects of fluorescent glutamate
indicators on neurotransmitter diffusion and uptake. *Elife* **9** (2020).
55. Jensen, T.P. *et al.* Multiplex imaging relates quantal glutamate release to
presynaptic Ca2+ homeostasis at multiple synapses in situ. *Nature*
Communications **10**, 1414 (2019).
56. Iwata, R., Kiyonari, H. & Imai, T. Mechanosensory-Based Phase Coding of Odor
Identity in the Olfactory Bulb. *Neuron* **96**, 1139-1152 e1137 (2017).

57. Gupta, P., Albeanu, D.F. & Bhalla, U.S. Olfactory bulb coding of odors, mixtures
and sniffs is a linear sum of odor time profiles. *Nat Neurosci* **advance online**
publication (2015).

58. Mathew, D. *et al.* Functional diversity among sensory receptors in a Drosophila
olfactory circuit. *Proc Natl Acad Sci U S A* **110**, E2134-2143 (2013).

59. Luo, M. & Katz, L.C. Response correlation maps of neurons in the mammalian
olfactory bulb. *Neuron* **32**, 1165-1179 (2001).

60. Zak, J.D., Reddy, G., Vergassola, M. & Murthy, V.N. Antagonistic odor
interactions in olfactory sensory neurons are widespread in freely breathing mice.
Nature Communications **11**, 3350 (2020).

61. Inagaki, S., Iwata, R., Iwamoto, M. & Imai, T. Widespread Inhibition, Antagonism,
and Synergy in Mouse Olfactory Sensory Neurons In Vivo. *Cell Reports* **31**,
107814 (2020).

62. Isaacson, J.S. & Scanziani, M. How inhibition shapes cortical activity. *Neuron* **72**,
231-243 (2011).

63. Wachowiak, M. *et al.* Optical Dissection of Odor Information Processing In Vivo
Using GCaMPs Expressed in Specified Cell Types of the Olfactory Bulb. *The*
Journal of Neuroscience **33**, 5285-5300 (2013).

64. Eiting, T.P. & Wachowiak, M. Artificial Inhalation Protocol in Adult Mice. *Bio*
Protoc **8** (2018).

65. Jordan, R., Fukunaga, I., Kollo, M. & Schaefer, A.T. Active Sampling State
Dynamically Enhances Olfactory Bulb Odor Representation. *Neuron* **98**, 1214-
1228 e1215 (2018).

66. McAfee, S.S. *et al.* Minimally invasive highly precise monitoring of respiratory
rhythm in the mouse using an epithelial temperature probe. *J Neurosci Methods*
263, 89-94 (2016).

67. Bozza, T., McGann, J.P., Mombaerts, P. & Wachowiak, M. In vivo imaging of
neuronal activity by targeted expression of a genetically encoded probe in the
mouse. *Neuron* **42**, 9-21 (2004).

68. Burton, S.D., Wipfel, M., Guo, M., Eiting, T.P. & Wachowiak, M. A Novel
Olfactometer for Efficient and Flexible Odorant Delivery. *Chem Senses* **44**, 173-
188 (2019).
69. Schlief, M.L. & Wilson, R.I. Olfactory processing and behavior downstream from
highly selective receptor neurons. *Nat Neurosci* (2007).

FIGURE LEGENDS

Fig. 1. Characterization of iGluSnFRs as reporters of glutamatergic signaling onto MT cells.

a. Expression of SF-iGluSnFR in MT cells of the OB. Image shows confocal stack with expression in MT cells after AAV2.1.hSynap.Flex.SF-iGluSnFR.A184V in the OB of a Tbet-Cre mouse. White arrows indicate a tufted cell with expression in the soma and primary dendrite extending to the glomerular layer (GL). EPL, external plexiform layer, MCL, mitral cell layer.

b. Top left: Mean fluorescence image taken in vivo, showing iGluSnFR expression in glomeruli after injection of AAV2.1.hSynap.Flex.SF-iGluSnFR into the OB a Pcdh21-Cre mouse. Top right: ΔF image showing responses to ethyl butyrate (mean of 8 presentations, 2 Hz inhalation). Bottom: Traces showing odorant-evoked iGluSnFR signal in three glomeruli, with dashed region expanded at right. Note distinct temporal responses across successive inhalations for each glomerulus.

c. Examples of SF-iGluSnFR.A184S responses imaged at high zoom from different glomeruli. Images show mean F (left) and ΔF response maps (right) to ethyl butyrate, showing localization of iGluSnFR signal to the glomerular neuropil (glom.), with a lack of signal on primary dendrites (pr. dend.) or MT cell somata (somas).

d. Traces showing responses to ethyl butyrate taken from the glomerular neuropil, primary dendrite, and MT somata (locations indicated in (d)), indicating negligible signal on the apical dendrite or somata.

Fig. 2. Glomerular glutamate signals show diverse inhalation-driven dynamics.

a. Left: Mean fluorescence image of SF-iGluSnFR.A184V expression imaged in vivo. Right: Inhalation-triggered ΔF response maps for three odorants.

b. ITA response traces for the glomeruli indicated in (A). Traces are averages across 17 inhalations. Left: ITAs from one glomerulus (ROI 5) responsive to 3 odorants, illustrating odorant-specific ITA dynamics. Right: ITAs from different glomeruli responsive to each of the three odorants, illustrating glomerulus-specific dynamics.

c. Plot of ITA onset latency, peak latency, and duration (FWHM, full-width half-max) for all responsive glomerulus-odor pairs using SF-iGluSnFR.A184S. Notch box plots (left)

of latency values (right). Notch: median, square: mean, box edges: 25th and 75th

percentile, n = 431 glomerulus-odor pairs, 5 mice.

d. Cumulative fraction of median-subtracted, ITA onset latencies for SF-

iGluSnFR.A184S signals, compared to those measured from GCaMP6f expressed in

OSN axon terminals (OMP-Cre x Rosa-GCaMP6f mice). Median-subtracted onset

latencies were similar: OSNs/GCaMP6f: 0.03 ± 0.11 sec, mean ± SD, n = 91 glomerulus

odor pairs, 6 mice; MTs/iGluSnFR.A184S: 0.002 ± 0.06 sec, n = 438 glomerulus-odor

pairs, 7 mice, Mann-Whitney Test, p = 0.1.

e. Preferential iGluSnFR expression in sTCs versus mitral cells. Left: Confocal image of

tissue section after Flex.AAV.SF-iGluSnFR injection into the superficial EPL of a CCK-

IRES-Cre mouse. Note strongest expression in superficial EPL. Right: Similar image

after Flex.AAV.iGluSnFR injection into anterior piriform cortex in a Tbet-Cre mouse.

f. Identical distributions of onset latencies, times-to-peak, and durations (FWHM) of

inhalation-triggered average waveforms imaged from CCK+ sTC and pcMT cell

populations. Range of onset latencies (10th - 90th percentiles): 195 – 451 ms (CCK+)

vs.192 to 447 ms (pcMTs); times to ITA peak: 373 – 938 ms (CCK+) vs. 367 to 828 ms

(pcMTs); FWHM: 313 – 1058 ms (CCK+) vs. 326 to 1007 ms (pcMTs). n = 174 and 160

glomerulus-odor pairs, respectively, from 3 mice each. All statistical comparisons,

p>0.05, Kolmogorov-Smirnov test.

Fig. 3. Glomerular glutamate signals show diverse temporal patterns across

repeated inhalations of odorant.

a. Mean fluorescence (left) and ΔF response maps for 5 odorants imaged with SF-

iGluSnFR.A184S (right). Odor numbers on the left side of the odorant map indicate

odorant identification code for listed odor name (for reference to (B)). Colored text

indicates odorants whose responses are shown in panels B, C and F.

b. Time-series of SF-iGluSnFR.A184S signal from 4 glomeruli, showing continuous

signal during 3 repetitions of 12 odorants, presented in random order. Shaded rectangle

indicates time of odorant presentations (2 sec duration). Signal is low-pass filtered at

0.5 Hz for display.

c. Average time-course of odorant-evoked responses in different glomeruli responsive to 4 of the 12 odorants from (B). Insets show change in response pattern across the field of view from (A) over the duration of odorant presentation, expressed as Pearson's *r* relative to the initial response in successive time bins of 387 ms (see Text). Dotted line indicates r = 0.

d. 'Waterfall' plot showing time-course of odorant-evoked SF-iGluSnFR.A184S signal in all significantly-responding glomerulus-odor pairs (one pair per row), normalized to the peak $\Delta F/F$ for each pair. Suppressive responses are low-pass filtered at 0.5 Hz and excitatory responses are low pass filtered at 2 Hz.

e. Proportion of excitatory (Exc.), suppressive (Sup) and biphasic (Exc. & Sup.) responses across the responsive population (right), as well as the entire population of glomeruli (left).

f. Change in glutamate response patterns during odorant presentation, summarized over all presentations. Black plot shows mean correlation over time, averaged across all odorant presentations. Shaded area indicates s.e.m. Dot plot at right shows correlation coefficients at the final time bin for all presentations.

Fig. 4. Diversity of glutamate dynamics imaged from MT cells in the awake mouse.

a. Top: mean SF-iGluSnFR.A184V fluorescence and Bottom: ΔF response map (ethyl butyrate) from the dorsal OB of an awake, head-fixed mouse. Right: Glutamate signals from 3 glomeruli (shown in (A)) during a single presentation of odorant (ethyl butyrate). Top trace shows respiration/sniffing as measured with an external flow sensor, with inhalation oriented downward. Lower left, snippet of signals from each glomerulus, illustrating temporal lag between different glomeruli and consistency of dynamics with each sniff. Signals are unfiltered. Lower right shows ITA waveforms for each glomerulus, generated across 137 sniffs over 6 presentations of ethyl butyrate. Vertical line indicates time of inhalation onset.

b. Left: Range of ITA onset (black) and peak (red) latencies, relative to median values, compiled across 108 glomerulus-odorant pairs (14 odorants, 7 fields of view). Notch: median, Box edges: 25th and 75th percentiles, square: mean values. Right: Spread of

latency values seen across multiple responsive glomeruli imaged in the same field of

view for the same odorant (as defined as the difference between maximum (lat_{max}) and

minimum (lat_{min}) latencies), for 23 unique odorant-FOVs.

c. SF-iGluSnFR.A184V signals imaged from 4 glomeruli from the same FOV showing

responses to each of two odorants, showing distinct temporal response patterns across

glomeruli and odorants. Traces are unfiltered and from a single presentation. Top trace

shows respiration/sniffing for each trial; vertical dotted lines indicate inhalation peak for

reference.

d. SF-iGluSnFR.A184V signals imaged from 3 glomeruli, taken from a separate field of

view in the same mouse, showing distinct response patterns to a third odorant (3-Octen-

2-one). Note weak suppressive response in one glomerulus (top trace).

e. Increasing odorant concentration elicits changes in glutamate signal dynamics in

awake mice. Traces show averaged responses (8 presentations) from 4 glomeruli in the

same field of view to a low (1:1000 dilution) versus high (1:10) concentration of ethyl

tiglate. Note that the response in the most sensitive glomerulus shows increased

adaptation at the higher concentration, while other glomeruli show facilitating

responses.

f. Changes in glutamate response patterns over the course of odorant presentation,

calculated as described for Fig. 3 in 4 awake mice, 33 odorant-FOVs (2 mice, SF-

iGluSnFR.A184V; 2 mice, SF-iGluSnFR.A184S). Plots of individual odorant responses

are shown in grey, thick black line shows mean, shaded region is quartiles (25[th] and 75[th]

percentiles). Note high variability in degree of decorrelation over time.

Fig. 5. Dual-color imaging reveals high correspondence between pre- and

postsynaptic signals in MT cells of the same glomerulus.

a. Post-hoc confocal image showing coexpression of jRGECO1a (magenta) and SF-

iGluSnFR.A184S (A184S, green) in MT cells of a Tbet-Cre mouse.

b. Dual-color two-photon imaging of SF.iGluSnFR.A184S and jRGECO1a signals from

MT cells. Left: odorant-evoked ΔF response maps for SF-iGluSnFR and jRGECO1a

signals (magenta imaged simultaneously. Note jRGECO1a signal in dendrites of several

MT cells exiting the central glomerulus, with SF-iGluSnFR signal confined to the

glomerular neuropil. Right: traces showing time-course of the fluorescence signal in
each channel (1 Hz inhalation). Top traces are from glomerular neuropil; lower traces
are from dendrites outside of the glomerulus.

c. Pseudocolor odorant-evoked ITA response maps across the green (A184S) and red
(jRGECO1a) channels. Arrows indicate ROIs with traces plotted in (d).

d. ITA traces taken from different glomeruli activated by the odorant in (c), with different
onset latencies, times to peak and durations in different glomeruli. Left traces: SF-
iGluSnFR.A184S; right traces: jRGECO1a; traces from the same glomerulus are shown
with the same color in each set. Each ITA trace is scaled to the same maximum.
Vertical lines indicate peak time for the signal in each glomerulus. The relative order of
peak times is the same for both signals.

e. Dual-color imaging of SF.iGluSnFR.A184S and jRGECO1a signals from MT cells in
the awake mouse. Left: Composite dual-color ΔF response map showing SF-iGluSnFR
(green) and jRGECO1a (magenta) signals evoked by cyclohexylamine. Right: Traces
showing fluorescence signal taken from the glomerular neuropile (top) and soma
(bottom) of an innervating TC. Traces are average of 8 presentations.

f. Trial-averaged ΔF response maps and traces for SF-iGluSnFR.A184S and

jRGECO1a signals imaged from a single presentation of odorant in an awake mouse.
Top trace shows respiration measured via external flow sensor. The SF-iGluSnFR
signal clearly follows each inhalation in only one of the three glomeruli shown, while the
jRGECO1a signal does not follow individual inhalations well in any glomeruli.

1003

Fig. 6. High correspondence between odorant-evoked temporal patterns of
glutamate signaling and calcium activity in MT cells of the same glomerulus.
a. ΔF response maps (left) and trial-averaged SF-iGluSnFR.A184S and jRGECO1a
signals imaged from select glomeruli for three odorants, illustrating high
correspondence in slow response dynamics. Traces are average of 4 presentations in
the same anesthetized mouse. Numbers indicate different glomeruli. Traces for
cyclohexylamine and 3-octen-2-one are scaled to the same maximum.
b. ΔF response maps and trial-averaged SF-iGluSnFR.A184S and jRGECO1a
responses taken from four glomeruli in response to three odorants in an awake mouse.

Traces are average of 16 presentations. Note distinct response patterns for the same

glomerulus in response to different odorants, but, with a few exceptions, similar

response patterns for SF-iGluSnFR and jRGECO1a signals.

c. High correlation between $T2-T1/T_{max}$ values measured for excitatory SF-iGluSnFR

and jRGECO1a responses in the same glomerulus in anesthetized (top) or awake mice

(bottom). Correlation coefficients for $T2-T1/T_{max}$ values between SF-iGluSnFR and

jRGECO1a responses were r = 0.87 in anesthetized mice (273 pairs,3 mice) and r =

0.90 in awake mice (57 pairs, 2 mice).

1021

Fig. 7. Effects of pharmacological blockade of polysynaptic glutamate signaling

onto MT cells.

a. Odorant response maps of GCaMP6f signals imaged from CCK+ sTCs, taken across

predrug, CGP35348 and APV+NBQX conditions, scaled to predrug response levels.

b. Response traces for the two glomeruli indicated in (k), showing enhanced response

amplitudes after application of CGP35348 (magenta) and either elimination (top) or

strong reduction (bottom) in responses after subsequent application of APV+NBQX

(green). In the latter case, the remaining response after APV+NBQX is delayed.

c. Example from a second experiment of a glomerulus with persistent responses after

APV+NBQX application. Traces (below) show expansion of response to first two

inhalations of odorant, showing delayed GCaMP6f response after APV+NBQX.

d. Scatter plot comparing CCK+ GCaMP6f responses (peak amplitude, first inhalation)

across CGP35348 and subsequent APV+NBQX conditions, imaged from 14 glomeruli in

3 experiments (2 mice). Summary statistics: ratio of post:pre APV+NBQX amplitudes,

0.27 ± 0.27, Wilcoxon signed ranks Z = 3.26, p = 0.001, n = 14 glomeruli). Shaded

regions indicate two glomeruli with near-totally blocked responses or with persistent but

weakened responses after APV+NBQX. Median post:pre APV+NBQX ratios, persistent:

0.54 ± 0.16 (n = 6), blocked: 0.06 ± 0.03 (n = 8).

e. Scatter plot of CCK+ GCaMP6f responses before and after APV+NBQX (5 mM/2.5

mM), without prior application of CGP35348.

f. Traces showing iGluSnFR.A184V signals imaged from pcMT cells in an aPC-injected Tbet-Cre mouse, comparing responses to the same odorant across Predrug, CGP35348, and APV+NBQX conditions. Inhalation frequency, 1 Hz.

g. Comparison of peak iGluSnFR responses in pcMT cells across CGP35348 and APV+NBQX conditions for first inhalation of odorant. Points indicate glomeruli from two mice (red and black). APV+NBQX/CGP35348 ratios, mouse 1 median: 1.03 (range, 0.89 - 1.11, n = 6), mouse 2: 0.86 (range, 0.79 - 1.05, n = 6). Combined comparison: Wilcoxon signed ranks, Z = 0.59, p = 0.55, n = 12. CGP35348/predrug ratios for the two mice (not shown), mouse 1 median: 1.32 (range, 1.28 - 1.52), mouse 2 median: 1.80 (range, 1.59 - 2.33); Combined comparison: Wilcoxon signed ranks, Z = -3.02, p = 0.003, n =12.

h. Top: Mean fluorescence and ΔF response maps showing odorant-evoked iGluSnFR signals in several glomeruli after SF-iGluSnFR.A184S expression in piriform-projecting MT cells. Bottom: Inhalation-triggered average responses from four of the glomeruli, imaged before and after application of APV+NBQX (5 mM / 2.5 mM). Response amplitudes in each glomerulus are modestly reduced, but relative latencies and durations are unchanged.

i. ITA response amplitudes before and after APV+NBQX (5 mM / 2.5 mM) application in pcMT cells from four experiments (47 total glomeruli). In each experiment, the same odorant was presented at two concentrations varying by a factor of 2.5 – 10; responses are plotted normalized to the maximal pre-drug response in each experiment. Summary statistics, paired Wilcoxon signed ranks per experiment, Exp. 1: n = 13, z = 0.21, p = 0.83; Exp. 2: n=16, z=0.18, p=0.86; Exp. 3: n = 8, z=2.5, p=0.008; Exp. 4: n = 14, z=3.3, p=1.2e-4.

j. Same data as in (i), showing ratio of post- to pre-drug ITA amplitude as a function of pre-drug response. Scatter indicates lack of consistent effect of APV + NBQX as a function of initial response amplitude.

k. Comparison of change in onset latencies and peak times of inhalation-linked responses after APV+NBQX application, measured from CCK+ GCaMP6f datasets (persistent glomeruli, (**d, e**)), pcMT iGluSnFR with 1 mM/0.5 mM APV+NBQX (data in (**g**)), and pcMT iGluSnFR with 5 mM/2.5 mM APV+NBQX (data in (**i**)). Note no change

in latencies for iGluSnFR signals. Boxes indicate 25th and 75th percentiles, line indicates

median, and whiskers denote outliers with a coefficient of 1.5. Summary statistics: 1)

CCK+ GCaMP6f, APV+NBQX vs. CGP35348, median Δ onset latency: 285 ms (range,

165 – 667 ms, n = 9 glomeruli), Z = -2.6 p = 0.004, Δ peak latency: 247 ms (range, 211

– 567 ms), Z = -2.6, p = 0.004. 2) iGluSnFR pcMTs, APV+NBQX vs. CGP35348 Δ onset

latency: mouse 1: 9.9 ms (range, -9.1 – 34.7 ms, n = 6), mouse 2: 9.8 ms (range, -19.4

– 18.6 ms, n = 6); Z = -1.47 p = 0.14 (same for both mice); Δ peak latency: mouse 1:

40.05 ms (range, 25.4 – 62.6 ms, n = 6), mouse 2: 14.5 ms (range, 1.8 – 38.8 ms, n =

6); Z = -1.89, -2.10, p = 0.06, 0.04); 3) iGluSnFR pcMTs, pre- vs. post- APV+NBQX,

median Δ peak latency, -6.7 ms; paired Wilcoxon signed ranks: expt. 1, n = 13, z=1.64,

p=0.1; expt. 2: n = 14, z=-0.55, p=0.58; expt. 3: n = 6, z=00, p=1; expt. 4: n = 14,

z=1.09, p=0.27.

1085

Fig. 8. Polysynaptic glutamate signaling contributes to responses to low-

concentration odorants and to slow temporal dynamics across inhalations.

a. Left: ΔF response maps showing responses to two odorants before application of

APV+NBQX (1 mM/0.5 mM). Signals are from SF-iGluSnFR.A184S injected into the OB

of a Tbet-Cre mouse. Right: Traces (mean of 4 presentations) from two glomeruli (top,

bottom) showing diverse temporal responses to different odorants, before and after

APV+NBQX application. Response to cyclohexylamine (bottom) shows a loss of the

initial transient and replacement by tonic signal. Response to vinyl butyrate (top) shows

increase in initial response magnitude and apparent loss of adaptation. Temporally

distinct patterns of response patterns to other odorants persist after drug application.

b. Mean and peak normalized response amplitudes after application of APV+NBQX,

with distributions indicated as previously.

c. Scatter plot of T_2-T_1/T_{max} values before and after APV+NBQX for all responsive

glomerulus-odor pairs across 3 mice (each mouse indicated by color), showing

decrease in adaptation for many pairs after application of APV+NBQX. Summary

statistics: paired t-tests, mouse 1, n= 7 glomerulus-odor pairs, p=0.001; mouse 2: n =

34, p=0.001; mouse 3: n = 42, p=5.45 x 10^{-6}.

Table 1. Odorants and concentration used.

Odor Name	Odor Number	Final Estimated ppm (by Experiment Type)				
		ITA	Diversity	Awake	Pharm. (hi)	Pharm. (low)
Butyric acid	1		0.08			
2-methylbutyric acid	2		0.03	0.03		
Valeric acid	3		0.03			
Hexanoic acid	5	0.2	0.01	0.01		
Isovaleric acid	6		0.04			
2-methylbuteraldehyde	9		2			
trans-2-methyl-2-butenal	10		0.7	0.7		
Heptanal	11		0.7			
2-methylvaleraldehyde	13		3	3		
Butyl acetate	18				17	
Ethyl butyrate	19	10			20	1 - 3
Vinyl butyrate	20	9	2	2	1	
Methyl valerate	22	8		5	40	2 - 5
Ethyl tiglate	22	2		0.6		0.07 - 0.7
Hexyl tiglate	23		1	1		
s-Methyl thiobutanoate	24		5			
Hexyl acetate	25		3			
Isopropyl tiglate	27		5	5		
Isoamylacetate	28		1			
2-hexanone	29	10		12	48	
Menthone	32	0.01				
2-butanone	33		16			
1-Hexanol	40		2			
Cyclohexylamine	43		15	15		
n-Methyl piperidine	45		1			
2-methoxy-3-methylpyrazine	51			1		
2-Isopropyl-3-methoxypyrazine	56		0.2			
Benzaldehyde	57	0.1				
Methyl benzoate	60	0.3				
Isobutylthiazole	81			0.07		
3-Octen-2-one	102		4			
trans-2-dodecanal	107		0.2			
Furfuyl mercaptan	110		0.7			
Isobutyraldehyde	136		5			

Each row represents a different odorant used in each experimental manipulation. Final Estimated ppm (by Experiment type) is the estimated vapor concentration of each odorant delivered to the animal, estimated from liquid dilution ratios used, reported vapor pressures, and calibration of the odor delivery device used. Pharmacology (Pharm.) estimated ppm shows the average of ppms across a range of presentations for

1111   ethyl butyrate, methyl valerate, and 2-hexanone. Odor number represents the internal
1112   reference number associated with each odorant, mostly used in Figure 3.

1113

1114

1115

Figure 1

Figure 2

Figure 3

a

mean F   valeric acid (3)   hexanoic acid (5)   heptanal (11)   isopropyl tiglate (27)   hexyl acetate (25)

SF-iGluSnFR.A184S

% max ΔF
95
-10

b

odor #: 136 102 2 43 20 25 3 27 56 5 11 10 20 5 10 25 136 56 102 2 3 11 27 43 2 56 136 11 3 25 10 20 102 27 5

3 Hz inhal.

20% ΔF/F
10 sec

c

valeric acid   hexanoic acid   heptanal   isopropyl tiglate

20% ΔF/F 1 s   3 Hz inhal.   20% ΔF/F 1 s   20% ΔF/F 1 s   20% ΔF/F 1 s

r: 1.0 / -0.4, time (sec) 0.25 – 1.8

d

odor

Glom - odor pair: 100, 200, 300, 400, 500, 600, 700

max
0 ΔF/F
-max

Time relative to odor onset (sec): -1 0 1 3 7 9 11

e

% of all Glom - odor pairs: 0 – 100

Unresponsive   Exc (89.3%)   Sup. (7.2%)   Exc. & Sup. (3.5%)

f

Correlation versus initial response

r (bin 1 vs. bin x): 1.0, 0.5, 0.0, -0.5

time (sec): 0.25 – 1.8

All odorant responses

Figure 4

Figure 5

a A184S jRGECO 100 µm 50 µm

b SF-iGluSnFR.A184S isobutylthiazole jRGECO1a
0 % max ΔF 90   0 % max ΔF 90
glom. SF-iGluSnFR jRGECO1a
50% ΔF/F 1s
dend.
odor 1 Hz inhal.

c SF-iGluSnFR.A184S jRGECO1a
95 % max ΔF -10
ethyl tiglate

d scaled ITAs
scaled to max inhal. SF-iGluSnFR.A184S inhal. jRGECO1a

e ΔF map (awake) cyclohexylamine
0 % max ΔF 95
glom. 2 sec
soma 20% ΔF/F
odor

f Awake, single-trial methyl benzoate
ΔF, iGluSnFR A184S 95 % max ΔF -10
ΔF, jRGECO1a
inhal. exhal.
1s

Figure 6

Figure 7

Figure 8

a

SF-iGluSnFR.A184S

vinyl butyrate

% max ΔF

-10

cyclohexylamine

Predrug

APV+NBQX

vinyl butyrate
ethyl butyrate
t-2M2B
2-MBA
isopropyl tiglate

20%
ΔF/F

1 sec

cyclohexylamine

3 Hz inhal.

odor

odor

b

APV+NBQX

Normalized amplitudes (Predrug)

1.5

1.0

0.5

0.0

mean   peak

c

APV+NBQX
T2-T1/Tmax

1.0

0.5

0.0

-0.5

-1.0

facilitation

adaptation

Exp. 1
Exp. 2
Exp. 3

-1.0   -0.5   0.0   0.5   1.0

Predrug
T2-T1/Tmax