Parallel Mosaic Speciation via Mutation-order and Ecological Divergence

Maddie E. James[1,2*†], Maria C. Melo[1†], Federico Roda[1,3], Diana Bernal-Franco[1,4], Melanie J. Wilkinson[1,2,5], Greg M. Walter[1,6], Huanle Liu[1,7], Jan Engelstädter[1], and Daniel Ortiz-Barrientos[1,2*]

[1]The University of Queensland, School of the Environment, St Lucia, Queensland, Australia.

[2]Australian Research Council Centre of Excellence for Plant Success in Nature and Agriculture, The University of Queensland, St Lucia, Queensland, Australia.

[3]Current Address: Universidad Nacional de Colombia, Departamento de Biología, Bogotá, Colombia.

[4]Current Address: Los Andes University, Bogotá, Colombia.

[5]Current Address: Queensland Alliance for Agriculture and Food Innovation, St Lucia, Queensland, Australia.

[6]Current Address: University of Tasmania, School of Natural Sciences, Hobart, Australia.

[7]Current Address: Centre for Genomic Regulation, Barcelona, Catalonia, Spain.

[*]Corresponding should be addressed to d.ortizbarrientos@uq.edu.au and m.james4@uq.edu.au
[†]Co-first authors

Short title: Parallel speciation in *Senecio*

Keywords: Parallel speciation, mutation-order, ecological speciation, natural selection, reproductive isolation, ecotypes

ABSTRACT

Natural selection shapes how new species arise, yet the mechanisms that generate reproductive barriers remain debated. Although ecological divergence in contrasting environments and mutation-order processes in similar environments are often viewed as distinct speciation mechanisms, we show they act together as part of a continuum we call 'parallel mosaic speciation.' In the *Senecio lautus* species complex, Dune and Headland ecotypes have evolved repeatedly along the Australian coastline. Through crossing experiments and field studies, we find that divergent natural selection promotes strong reproductive isolation between the Dune and Headland ecotypes. While uniform selection maintains reproductive compatibility among ecologically similar Dune populations, Headland populations have evolved reproductive barriers despite their convergent prostrate phenotypes, likely driven by adaptation to heterogeneous environments. To understand how habitat heterogeneity contributes to patterns of reproductive isolation, we extend previous theoretical work on the accumulation of hybrid incompatibilities to account for environmental gradients and polygenic adaptation. We show that the probability of reproductive isolation depends on three factors: how similar the environments are, how complex the genetic architecture is, and how selection coefficients are distributed among beneficial mutations. These theoretical findings explain how reproductive isolation arises in systems like *Senecio*, where multiple forms of selection jointly drive parallel speciation.

INTRODUCTION

Speciation, the process by which new species arise, is a fundamental driver of biodiversity. However, demonstrating the direct role of natural selection in speciation remains a challenge. Systems exhibiting parallel evolution—where populations independently acquire similar phenotypes under comparable selective pressures—provide a powerful framework for investigating the role of selection in speciation (Schluter 2001; Langerhans and Riesch 2013). These natural replicates of the evolutionary process (Schluter and Nagel 1995; Schluter 2000) enable a thorough examination of the mechanisms that drive reproductive isolation. Current theoretical frameworks distinguish two primary modes of speciation by natural selection: ecological speciation, driven by deterministic adaptation to contrasting environments, and mutation-order speciation, where stochastic processes generate reproductive barriers between populations adapting to similar conditions (Schluter 2009; Schluter and Conte 2009; Nosil and Flaxman 2011; Nosil 2012; Crespi and Nosil 2013, but see Sobel et al. 2010; Langerhans and Riesch 2013).

Parallel ecological speciation occurs when divergent natural selection repeatedly generates reproductive isolation between populations adapting to contrasting environments, while maintaining reproductive compatibility among populations experiencing similar selective pressures (Figure 1A; Schluter and Nagel 1995; Ostevik et al. 2012). The resulting reproductive barriers can manifest through environment-dependent isolation, where immigrants and hybrids show reduced fitness in parental habitats, and environment-independent isolation, characterized by intrinsic hybrid incompatibilities regardless of ecological context (Rundle and Nosil 2005; Nosil 2012). The threespine stickleback (*Gasterosteus aculeatus*) system exemplifies this process, where the repeated colonization of freshwater habitats by marine ancestors has driven parallel evolution of adaptive traits, including reduced body armor and modified feeding morphology (Colosimo et al. 2005; Wund et al. 2008). Reproductive isolation between marine and freshwater forms has evolved primarily through extrinsic mechanisms, while geographically separated freshwater populations maintain reproductive compatibility (Rundle et al. 2000; McKinnon and Rundle 2002; McKinnon et al. 2004). This pattern—where ecological divergence predicts reproductive isolation more strongly than geographical separation—provides compelling evidence for the role of natural selection in speciation. Similar patterns of parallel ecological speciation have been documented in other animal taxa, including *Littorina* snails (Johannesson et al. 2010, 2024) and *Timema* walking stick insects (Nosil et al. 2002; Soria-Carrasco et al. 2014), although comparable evidence in plants remains limited (see Ostevik et al. 2012; James et al. 2023).

Mutation-order speciation is an alternative mechanism distinct from ecological divergence that can

generate reproductive isolation. Under this process, uniform natural selection favors the same

beneficial alleles across populations adapting to the same selective pressures (Figure 1B). However,

reproductive incompatibilities arise as different beneficial alleles randomly arise and fix in each

population. These alleles are incompatible when combined in hybrid genomes due to negative

epistatic interactions, and as a result, can cause reproductive isolation between taxa (Mani and

Clarke 1990; Schluter 2009; Nosil and Flaxman 2011; Nosil 2012). The mechanistic basis of

mutation-order speciation differs fundamentally from ecological speciation: rather than arising from

maladaptation to parental environments, reproductive barriers emerge from deleterious interactions

between independently acquired beneficial mutations. Because these mutations experience positive

selection, their fixation and subsequent incompatibilities can accumulate more rapidly than under

neutral divergence through genetic drift (Mani and Clarke 1990).

Experimental studies provide compelling evidence for mutation-order speciation, demonstrating the

evolution of reproductive barriers between populations adapting to identical environments such as

in *Drosophila* (Hsu et al. 2024) and *Saccharomyces* (Ono et al. 2017). While the strictest definitions

of mutation-order speciation require populations to inhabit identical environments (Mani and Clarke

1990), some definitions acknowledge it can also occur when populations adapt to similar but non-

identical habitats (e.g., Schluter 2009; Schluter and Conte 2009; Nosil and Flaxman 2011). This

makes documenting mutation-order speciation in nature challenging: determining whether

reproductive barriers arose from the stochastic fixation of beneficial mutations or subtle ecological

differences can blur the distinction between mutation-order and ecological processes. Addressing

this challenge requires systematic investigation of speciation mechanisms across diverse taxa with

broad geographic ranges (Anderson and Weir 2022), particularly in understudied groups such as

plants.

The *Senecio lautus* species complex (Ali 1964; Radford et al. 2004; Thompson 2005) provides an

exceptional system for examining the relationship between natural selection and speciation. Two

phenotypically distinct ecotypes have repeatedly evolved in the species complex: an ancestral erect

Dune ecotype occupying sheltered sand dunes and a derived prostrate Headland ecotype inhabiting

exposed rocky outcrops (Figure 2A). These ecotypes form multiple Dune-Headland parapatric pairs

along the Australian coastline (Figure 2B), which can be considered natural replicates of the

evolutionary process (Roda et al. 2013a; James et al. 2021a). Each ecotype is locally adapted to a

distinct habitat type (Roda et al. 2013b; Walter et al. 2016, 2018a; James et al. 2021b), yet

Headland sites exhibit fine-scale heterogeneity in soil chemistry, microclimate, and biotic

interactions, which generates a complex mosaic of selective pressures (Roda et al. 2013b; Walter et

al. 2016). This environmental variation has significant implications for the genetic architecture of

adaptation: while Headland populations exhibit convergent prostrate phenotypes, they achieve this

convergence through distinct combinations of small-effect alleles, revealing the polygenic nature of

adaptive evolution in the system (Roda et al. 2013b; James et al. 2021b, 2023a; Kaur 2024).

Research in *Senecio* has revealed complex interactions between environment-dependent and

intrinsic reproductive barriers. Reciprocal transplant experiments demonstrate extrinsic isolation

between Dune and Headland ecotypes, indicating that divergent selection maintains reproductive

boundaries through adaptation to contrasting environments (Melo et al. 2014; Richards and Ortiz-

Barrientos 2016; Richards et al. 2016; Walter et al. 2016, 2018b; Wilkinson et al. 2021). While F_1

hybrids between eastern coastal populations are largely reproductively compatible (Melo et al.

2014; Richards and Ortiz-Barrientos 2016; Walter et al. 2016), F_2 and subsequent generations

display strong forms of reproductive isolation (Walter et al. 2020; Wilkinson et al. 2021). Finally,

the identification of a shared genetic basis between gravitropism—a key ecological trait

differentiating the ecotypes—and hybrid sterility in *Senecio* demonstrates a direct pathway through

which natural selection on ecological traits promotes speciation (Wilkinson et al. 2021).

In this work, we use empirical analyses to test fundamental predictions of parallel ecological and

mutation-order speciation in the *Senecio lautus* species complex. We investigate how

environmental heterogeneity shapes the accumulation of reproductive barriers and create a

mathematical framework to explicitly incorporate environmental gradients and polygenic

architectures, providing novel insights into speciation across heterogeneous landscapes. Our work

shows that deterministic and stochastic processes interact in complex ways to generate reproductive

isolation during parallel speciation.

METHODS

Phylogenetic Independence of Replicate Populations

Samples

We studied parallel speciation in six parapatric Dune-Headland *Senecio lautus* population pairs

from the eastern and southern coasts of Australia (Figure 2A, Table S1). For each population, we

used previously collected leaf samples from 12 individuals (Roda et al. 2013a). To strengthen our

phylogenetic analysis, we included two additional ecotypes (two Inland and two Alpine

populations; $n_{pop} = 11$), and the closely related African species *S. madagascariensis* (see Roda et al.

2013) as an outgroup ($n = 4$) (Figure 2A, Table S1). We extracted DNA from each sample using a

modified CTAB protocol (Clarke 2009), purified samples using Promega's Wizard® SV Gel and

PCR Clean-Up System, and standardized each sample to 30ng/μL. We undertook targeted re-

sequencing of nuclear genomic regions, resulting in 13 neutral markers across all populations (see

Supplementary Material 1 for details on primers, library preparation, sequencing, bioinformatics

and neutrality tests; Table S2, S3). For a detailed taxonomy of the *S. lautus* species complex, please

see supplementary tables in Roda et al. (2013a).

Phylogeny

Previous research in *S. lautus* demonstrated phylogenetic independence of five out of the six Dune-

Headland population pairs in the current study (James et al. 2021a). To confirm the independence of

all six pairs, we conducted a Bayesian phylogenetic analysis using the 13 neutral loci with *BEAST

v1.7.5 (Heled and Drummond 2010), an extension of BEAST (Drummond et al. 2012) designed for

multilocus and multi-individual species tree estimation. We generated the XML file for *BEAST

using BEAUTi v1.7.5 (Drummond et al. 2012), and ran the analysis with a chain length of

300,000,000 under a strict molecular clock. Using ITS as our reference locus (mutation rate =

4.13×10^{-9} subs/site/year for herbaceous plants; Kay et al. 2006), we estimated relative mutation

rates for all other loci. The HKY model (Hasegawa et al. 1985) best fit our 13 loci based on the

Bayesian Information Criterion implemented in jModelTest (Posada and Crandall 1998). We used a

Yules speciation process for species tree estimation, which assumes that lineages split at a constant

rate. We generated the maximum clade credibility tree in TreeAnnotator v1.7.5 (Drummond et al.

2012) with a burn-in of 10,000 steps, and visualized the tree in FigTree v1.4.4 using *S.*

madagascariensis as the outgroup.

Population Structure

We assessed population structure by identifying the number of genetic clusters (K) across all

populations using STRUCTURE v2.3.4, a Bayesian Markov Chain Monte Carlo (MCMC) approach

(Pritchard et al. 2000). We performed analyses with the variable sites of the 13 neutral loci using an

admixture model and the correlated allele frequency model (Falush et al. 2003). Following

guidelines from Gilbert et al. (2012) and Janes et al. (2017), we tested K values from 1 to 16,

running 20 iterations per K with a burn-in of 100,000 and MCMC run length of 100,000. We

confirmed convergence of model parameters through visual inspection of MCMC summary

statistics. The most likely K value was evaluated using methods from Pritchard et al. (2000) and

Evanno et al. (2005), implemented in STRUCTURE HARVESTER v0.6.93 (Earl and vonHoldt

2012). Since both methods tend to overestimate K, and higher K values provided no additional

clustering information, we selected the smallest K that captured the major population structure in

the data. We aligned cluster assignments across iterations using CLUMPP v1.1 (Jakobsson and

Rosenberg 2007) with the complete search algorithm, and visualized results using DISTRUCT v1.1

(Rosenberg 2004).

Intrinsic Reproductive Isolation Between and Within Ecotypes

Samples

To investigate patterns of reproductive isolation, we selected four Dune-Headland population pairs:

two from the eastern clade (LH and CH) and two from the southeastern clade (VC and SA) of the

phylogeny (Figure 2; Table S1). From each population, we collected seeds from 30 individuals in

the field spaced at least ten meters apart. We stored seeds in dry conditions at 4°C at The University

of Queensland. To create seed stocks for each population as well as to eliminate maternal effects

(Bischoff and Müller-Schärer 2010), we first germinated and grew plants for one generation. To

induce germination, we scarified seeds by trimming 1 mm off the micropyle side and placed them

on moist filter paper in Petri dishes in a controlled growth room at 25°C. We kept seeds in darkness

for three days to promote root elongation, and then transferred to a 12h:12h light:dark cycle for

seven days to encourage vegetative growth. We then transplanted seedlings into 0.25L pots

containing a soil mix (70% pine bark, 30% coco peat) supplemented with slow-release Osmocote

fertilizer (5 kg/m³) and Suscon Maxi insecticide (830 g/m³). After two months, we conducted

controlled crosses by repeatedly rubbing mature flower heads together over three to five days. This

ensured bi-directional pollen transfer and maximum fertilization opportunity. The resulting seeds

were stored at 4°C.

Reproductive Isolation Measures

We quantified two components of intrinsic reproductive isolation: F_1 seed set and F_1 viability

(germination). To assess these barriers, we compared the success of crosses between populations of

the *same* ecotype (D × D and H × H population crosses) with those of *different* ecotypes (D × H).

This was undertaken both *within* and *between* the eastern and southern phylogenetic clades to

account for phylogenetic divergence time (Coyne and Orr 1997). We also conducted intra-

population crosses as controls. Using the protocol outlined above, we grew up to 14 families from

each of the eight populations from the eastern and southeastern clades of the phylogeny (Table S1).

When the plants flowered, we performed intra- and inter-population crosses twice daily, completing

260 crosses in total (see Table S4).

To measure reproductive isolation at the F_1 seed set stage, we counted the proportion of fertilized

seeds per flower head, identifying unfertilized seeds by their thin size and pale color. We quantified

the strength of reproductive isolation (RI) for F_1 seed set between populations as:

$$RI_{seed\ set} = 1 - \frac{proportion\ of\ fertilized\ seeds\ from\ the\ interspecific\ crosses}{average\ proportion\ of\ fertilized\ seeds\ from\ the\ two\ parental\ intraspecific\ crosses}$$

201 To assess reproductive isolation at the F_1 viability stage, we measured germination rates from the

202 crosses defined above. For each of the 260 crosses, we germinated five seeds by placing them onto

203 moist filter paper in Petri dishes (one cross per Petri dish). We used the same germination

204 conditions as above but without scarification to test the intrinsic ability of embryos to emerge from

205 the seed coat. The strength of reproductive isolation for F_1 viability was calculated as:

$$RI_{viability} = 1 - \frac{proportion\ of\ germinated\ seeds\ from\ the\ interspecific\ crosses}{average\ proportion\ of\ germinated\ seeds\ from\ the\ two\ parental\ intraspecific\ crosses}$$

Statistical Analyses

We analyzed patterns of intrinsic reproductive isolation using linear mixed effect models in R

v4.1.0 (R Core Team 2021) with the *lmerTest* package (Kuznetsova et al. 2017):

RI = Clade + Ecotype + (1 | CrossType)

where *RI* is the reproductive isolation values for either F_1 seed set or F_1 viability, *Clade* indicates

whether crosses occurred within or between phylogenetic clades, *Ecotype* denotes whether crosses

were between the same or different ecotypes, and *CrossType* is a random effect with a random

intercept model, representing the population comparison. For both F_1 seed set and viability models,

we initially tested for an interaction between *Clade* and *Ecotype* but removed it due to non-

significance. To examine specific cross-type effects, we ran additional linear models:

RI = Clade + EcotypeCross

where *EcotypeCross* specifies the exact cross comparison (D × D, H × H, D × H, or H × D). Again,

we removed the interaction terms as they were non-significant. Finally, we used one-sided t-tests to

determine whether reproductive isolation values for D × D and H × H crosses (both within and

between clades) were significantly greater than zero.

Extrinsic Reproductive Isolation Within Ecotypes

 Samples

We tested for mutation-order effects on speciation by examining extrinsic reproductive isolation

within ecotypes. If this type of speciation is occurring in *Senecio*, populations of the same ecotype

should exhibit similar performance at other localities within the same environment. This was

assessed by reanalyzing data from a previous field transplant experiment by Walter et al. (2016).

Briefly, in this study seeds of six populations (three Dune-Headland replicate pairs) were generated

under common garden conditions and then transplanted (n = 150-180 seeds per population in each

environment; Table S1) into the dune and headland environments of the Lennox Head (LH)

population pair. Seeds were planted in fully randomized grids with six replicate blocks in each

environment, seedling survival was monitored over time, and seedling establishment was recorded

when plants had grown ten true leaves. For further details, see Walter et al. (2016).

Statistical Analyses

We compared local population performance (LH Dune seeds planted in the LH dune habitat; LH

Headland seeds planted in LH headland habitat) against non-local populations (i.e., Stradbroke

Island and Cabarita Beach Dune and Headland seeds in the LH dune and headland environments,

respectively). Our analyses examined two fitness components: survival and establishment.

Using the R package *coxme* (Therneau 2024), we modeled survival separately for dune and

headland environments:

Survival = Population + (1 | Block)

where *Survival* is days survived (censored at day 320), *Population* indicates seed source location,

and *Block* is a random effect for replicate experimental blocks within each environment.

We then tested whether local populations showed similar probabilities of reaching seedling

establishment compared to non-local populations using a generalized linear mixed-effects model in

the R package *lme4* (Bates et al. 2015). The model used the binomial family with a logit link

function to model the probability of establishment success. We analyzed dune and headland

environments separately, with the model:

Establishment = Population + (1 | Block)

where *Establishment* is a binary variable indicating whether a plant reached ten leaves and *Block* is

a random effect for replicate experimental blocks within each environment.

In both analyses, we tested whether non-local populations performed similarly to the local

population. To do this, we explicitly set the local population as the reference level (intercept) for the

Population factor. This parameterization allows coefficient estimates to represent deviations of non-

local populations from local population performance, where significant negative coefficients

indicate local adaptation.

Connecting Ecology and Phenotype with Intrinsic Reproductive Isolation

To further assess the role of mutation order in driving parallel speciation in *Senecio*, we examined

the relationship between intrinsic reproductive isolation and environmental or phenotypic distance.

For mutation-order to hold true, we expect no associations because populations are expected to fix

advantageous mutations randomly rather than in response to ecological or phenotypic differences

(Schluter 2009).

Intrinsic Reproductive Isolation and Environmental Distance

To test the role of ecological differences to intrinsic isolation, we explored the association between

intrinsic reproductive isolation (F_1 seed set and F_1 viability, as measured above) and environmental

variance among Headland populations. We used previously published soil data from Roda et al.

(2013b) to quantify environmental variance, where 38 soil variables (nutrients, salts and metals)

were measured for each population. We used R for statistical analyses and scaled each soil variable

to a mean of zero and standard deviation of one. We used the *vegan* package (Blanchet et al. 2018)

in R to create a distance matrix of the soil variables for each population. We performed

multidimensional scaling to calculate the Euclidian distance between populations in environmental

space using the first five principal component axes, which explained more than 95% of the

variance. To estimate genetic distance, we used the *ape* package (Paradis et al. 2004) in R to

calculate pairwise distances between populations based on branch lengths from the phylogeny

constructed above.

To assess the relationship between reproductive isolation and environmental distances, we

performed linear models:

RI = GeneticDistance + EnvironmentalDistance

where *RI* is the reproductive isolation value for either F_1 seed set or F_1 viability.

We examined the association between intrinsic reproductive isolation (F_1 seed set and F_1 viability)

with phenotypic variance between Headland populations. We used a combination of previously

published phenotypic data from Walter et al. (2018a), and unpublished data for additional

populations measured at the same time, where phenotypes were measured in controlled glasshouse

conditions. This dataset consists of four plant architecture traits (vegetative height, stem

length/width, number of branches, and stem diameter) and six leaf traits (area, $perimeter^2/area^2$,

circularity, number of indents, indent width, and indent depth), measured for up to 17 individuals

per population (Table S1), see Walter et al. (2018a) for specific details on growing conditions and

trait measurements. As in the environmental analysis, we scaled each phenotypic variable to have a

mean of zero and standard deviation of one, calculated the distance matrix of the phenotypic traits

for each population, and performed multidimensional scaling to calculate the Euclidian distance

between populations in environmental space using the first seven principal component axes, which

explained more than 95% of the variance. We used the genetic distances calculated above in *ape*.

We performed linear models to assess the relationship between reproductive isolation and

phenotypic distances, again averaging the reciprocal crosses:

RI = GeneticDistance + PhenotypicDistance

where *RI* is the reproductive isolation value for either the F_1 seed set or F_1 viability for each

population comparison.

Mathematical Analysis of Parallel Speciation

To further understand how various forms of natural selection drive patterns of reproductive

isolation in *Senecio*, we created a mathematical framework to explore how both ecological and

mutation-order processes can jointly underlie patterns of reproductive isolation. Our framework

begins with the classic Dobzhansky-Muller model, which establishes how hybrid fitness decreases

through epistatic interactions between incompatible alleles (Orr 1995). Next, we build a model

based on the Unckless-Orr framework (Unckless and Orr 2009), which quantifies the probability of

Dobzhansky-Muller Incompatibility (DMI) formation between populations adapting to the same

environments – representing the strictest form of mutation-order speciation, where environmental

conditions are identical between populations. To capture the dynamics of divergent natural

selection, we extend the model to incorporate environmental heterogeneity, where DMIs

accumulate across a gradient of environmental similarity. This gradient ranges from identical

environments where mutation-order processes dominate, to contrasting environments where

ecological divergence occurs. Additionally, we integrate polygenic adaptation into the framework,

providing a more nuanced understanding of how incompatibilities evolve with increasing genetic

complexity.

Foundations

The foundation of our analysis builds upon the classic Unckless-Orr model examining DMI

formation between two populations adapting to identical environments (Unckless and Orr 2009).

This model considers two interacting loci A and B, each with ancestral (A_0, B_0) and derived

(A_1, B_1) alleles. The ancestral genotype (A_0, B_0) has a fitness of 1, while genotypes with single

derived alleles $A_1 B_0$ and $A_0 B_1$ have fitness values of $1 + s_A$ and $1 + s_B$ respectively. The

combination of both derived alleles $(A_1 B_1)$ creates an incompatibility with fitness $1 - t$, where t

quantifies the severity of the DMI (Orr 1995). When $t = 1$, the hybrid combination is completely

inviable/sterile, while smaller values of t represent partial reproductive isolation. The inclusion of t

in the model helps capture the concept that while individual-derived alleles can be beneficial in

their genetic backgrounds, their combination in hybrids can be deleterious $(t > 0)$. This is a key

feature of the Dobzhansky-Muller model of speciation—incompatibilities arise not from individual

mutations being deleterious, but from their negative epistatic interactions when brought together in

hybrids.

Under strong selection $(Ns >> 1)$ and weak mutation $(N\mu << 1)$, where adaptive trajectories

retain stochastic elements despite directional selection, the probability of DMI formation depends

on selection coefficients through the Unckless-Orr model:

$$P_{DM,U} = \frac{2 s_A s_B}{(s_A + s_B)^2} \qquad (1)$$

We use the subscript *U* to refer to the Unckless-Orr model. This equation reveals that during

adaptation to identical environments, DMIs are most probable when selection coefficients are

similar $(s_A \approx s_B)$, as populations experience maximum stochasticity in the order of mutation

fixation. When selection coefficients are equal, either beneficial mutation is equally likely to fix

first in each population, maximizing the probability that populations will fix different alleles.

However, when selection coefficients differ substantially, both populations are likely to fix the

mutation with the stronger selective advantage first, reducing the probability of DMI formation.

Extensions

We extend this framework by modeling how environmental differences create asymmetric selection

on alleles between populations. For any locus, we assume its selection coefficient in its "home"

environment (s_i) is reduced by a factor φ when it occurs in the alternative environment (φs_i). This

parameter $\varphi \in [0,1]$ captures the degree to which selection pressures transfer between

environments, with $\varphi = 1$ indicating identical selection and $\varphi = 0$ indicating complete asymmetry

where alleles that are beneficial in one environment are neutral in alternative environments.

Under these assumptions, the probability of DMI formation becomes:

$$P_{DM,E} = \frac{s_A s_B (1 + \varphi^2)}{(s_A + \varphi s_B)(\varphi s_A + s_B)} \qquad (2)$$

When environments are identical ($\varphi = 1$), the equation reduces to the classic Unckless-Orr model.

As environmental differences increase (decreasing φ), selection becomes increasingly asymmetric

between populations, making it more likely they will fix different alleles. In the extreme case of

complete asymmetry ($\varphi = 0$), DMI formation becomes inevitable because alleles strongly favored

in one environment are neutral in the other.

For polygenic traits involving n interacting loci with selection coefficients s_i, the probability of at

least one DMI forming is:

$$P_{DM,P} = 1 - \frac{\varphi}{\sigma_1 \sigma_2} \sum_{i=1}^{n} s_i^2 \qquad (3)$$

353 where σ_1 and σ_2 represent the sums of selection coefficients in each population. When

354 environments are identical ($\varphi = 1$), this reduces to an n-locus extension of the two-locus

355 Unckless and Orr (2010) model, while complete asymmetry ($\varphi = 0$) guarantees DMI formation. For

356 equal selection coefficients ($s_i = s$), the formula simplifies to:

$$P_{DM,Ps(i=s)} = 1 - \frac{4\varphi}{n(1 + \varphi)^2} \qquad (4)$$

This shows how genetic complexity amplifies the effects of selection asymmetry through

combinatorial growth in potential incompatibilities (see *Supplementary Material 2* for detailed

derivations of all equations).

Implementation

We implemented these models using Python v3.10.9 and NumPy v1.23.5 (Harris et al. 2020) for

numerical calculations, and Matplotlib v3.9.2 (Hunter 2007) for visualization. Our analysis

explored selection coefficient space (s_A vs s_B), environmental similarity (φ), and the impact of

interacting loci number (n) on DMI probability, providing a theoretical explanation for the

observed patterns of reproductive isolation in *Senecio*.

RESULTS AND DISCUSSION

Empirical Patterns of Parallel Speciation

Phylogenetic Independence Reveals Multiple Origins of Coastal Ecotypes

To investigate parallel speciation in *Senecio*, we first confirmed the independent origins of Dune

and Headland populations in this study (also see Roda et al. 2013a; James et al. 2021a). Using 13

neutral markers, we found that populations cluster into two main clades (eastern and southeastern;

Figure 2C, 2D), that align with their coastal geographic distribution (Figure 2B). As expected, the

phylogenetic relationships support parallel evolution: Dune-Headland population pairs form sister

taxa, with both ecotypes present within each clade (Figure 2C). This result confirms previous

independent datasets in the *Senecio* system (Roda et al. 2013a; James et al. 2021a). Although gene

flow can create false signals of parallel evolution (Endler 1977; Barton and Hewitt 1985; Coyne and

Orr 2004; Bierne et al. 2013), previous work in *Senecio* showed that gene flow among *Senecio*

populations, both within and between ecotypes, is negligible (James et al. 2021a). These findings

indicate that the observed phylogenetic patterns reflect the independent adaptive divergence of

Dune-Headland pairs along the Australian coastline.

Divergent Selection Drives Strong Reproductive Isolation Between Ecotypes

Parallel ecological speciation generates specific predictions about patterns of reproductive isolation:

populations adapting to contrasting environments should develop reproductive incompatibilities,

while those inhabiting similar environments should maintain reproductive compatibility. In

contrast, mutation-order processes predict that populations evolve reproductive barriers even when

adapting to similar environments through the independent fixation of different beneficial alleles

(Schluter and Nagel 1995; Ostevik et al. 2012). To evaluate these contrasting theoretical predictions

in *Senecio*, we quantified intrinsic reproductive isolation between populations from the same

environments (within Dune or within Headland ecotypes) and different environments (between

Dune and Headland ecotypes). Our analysis focused on two quantitative components of

reproductive isolation, measured relative to parental averages: F_1 hybrid seed production and F_1

hybrid viability during germination.

Consistent with the role of divergent selection, populations from similar environments showed

higher reproductive compatibility than those from different environments. This pattern held for both

seed set ($F_{1,15.73} = 5.93, P = 0.023, R^2 = 0.58$ after taking into account the effect of clade

$F_{1,15.61} = 13.49, P = 0.002$) and viability ($F_{1,14.48} = 7.25, P = 0.017, R^2 = 0.36$ after taking into

account the impact of clade $F_{1,14.48} = 7.21, P = 0.018$) (Figure 3A). Crosses within clades were

generally viable, with crosses between the same ecotypes showing some hybrid vigor (reproductive

isolation values < 0), suggesting complementary genetic interactions. The most substantial (almost

complete) reproductive isolation occurred in Dune-Headland crosses between the two clades

(Figure 3A). These results indicate that natural selection drives speciation in *Senecio*, with

reproductive isolation increasing as genetic divergence accumulates between ecotypes. This pattern

mirrors findings in other systems, including stickleback fish (Hendry et al. 2009; Stuart et al. 2017),

Littorina snails (Johannesson et al. 2024), and cichlid fishes (Weber et al. 2021).

Unexpected Reproductive Barriers Emerge Among Convergent Headland Populations

To establish whether the observed patterns of intrinsic reproductive isolation represent parallel

ecological speciation in *Senecio*, we must demonstrate reproductive compatibility within each

ecotype, for both hybrid viability and fertility. Cross type (D × D, H × H, D × H, or H × D)

significantly affected reproductive isolation for both F₁ seed set ($F_{3,29} = 3.44, P = 0.03, R^2 = 0.46$

after taking into account the effect of clade $F_{1,29} = 14.44, P = 0.0007$) and F₁ viability ($F_{3,29} = $

$3.52, P = 0.03, R^2 = 0.34$ after taking into account the effect of clade $F_{1,29} = 4.51, P = 0.042$)

(Figure 3B). Dune populations showed the expected pattern of parallel ecological speciation,

maintaining reproductive compatibility within and between clades. Their reproductive isolation

values were not significantly greater than zero for either F₁ seed set (within clades $t_5 = -1.85, P = $

0.84, between clades $t_5 = -0.44, P = 0.66$), or F₁ viability (within clades $t_5 = -4.4, P = 0.93$,

between clades $t_5 = -1.51, P = 0.9$). Headland populations were similarly compatible within

clades for both F₁ seed set ($t_3 = -1.96, P = 0.93$) and F₁ viability ($t_3 = -1.02, P = 0.81$), and

between clades for viability ($t_5 = -0.95, P = 0.19$).

Our analyses revealed an unexpected pattern of reproductive incompatibility among Headland

populations from different clades, manifested as significantly reduced F₁ seed set ($t_5 = 8.12, P = $

0.0002) (Figure 3B). While this pattern contradicts predictions of ecological speciation, it aligns

with the expectation that mutation-order processes contribute to speciation. The reproductive

isolation between Headland populations suggests parallel evolution of prostrate phenotypes

occurred through distinct adaptive trajectories, resulting in different combinations of beneficial

alleles in each population that create intrinsically unfit hybrids between populations. This

interpretation is supported by previous genetic analyses in *Senecio*, which demonstrated that

convergent Headland morphology evolved through the selection of distinct alleles and genes across

populations (Roda et al. 2013b; James et al. 2021b). To further explore the role of mutation-order

effects on speciation in *Senecio*, we tested additional predictions of speciation theory.

Ecological Differences do not Explain Reproductive Isolation in Headland Populations

Previous work in *Senecio* revealed contrasting patterns of environmental variation. Each ecotype

occupies a distinct soil type, where variation within ecotypes is lower than between ecotypes (Roda

et al. 2013b; Walter et al. 2016). The soils of different Headland populations share fundamental

characteristics of being shallow and nutrient-rich but vary in specific mineral composition based on

local geology. This habitat heterogeneity contrasts with the Dune populations, which occupy

remarkably similar habitats even when separated by thousands of kilometers. To determine whether

ecological variance reflect fitness variation within each ecotype, we reanalyzed reciprocal

transplant experiments from Walter et al. (2016). In these experiments, three eastern-coast Dune

populations were transplanted into the dune habitat at Lennox Head, while three eastern-coast

Headland populations were transplanted into the rocky headland. Dune populations showed no

signal of local adaptation: non-local individuals performed equally well or better than local Lennox

Head individuals in both survival (Figure 4A) and establishment (Figure 4B). In contrast, the local

Lennox Head Headland population consistently outperformed non-local populations in both fitness

components (Figure 4).

Given the fitness differences among Headland populations at Lennox Head, we tested whether

environmental variation (measured by soil ecology) explains patterns of intrinsic reproductive

isolation across geography. If ecological divergence in soil composition drives reproductive

barriers, we expect to find a correlation between measured environmental differences and intrinsic

reproductive isolation across Headland populations. After controlling for genetic distance, we found

no significant relationship between soil-based ecological divergence and reproductive isolation for

F_1 seed set $\left(F_{1,2} = 3.00, P = 0.228\right)$, or F_1 viability $\left(F_{1,2} = 0.16, P = 0.729\right)$. The absence of an

environment-reproductive isolation correlation for soil composition is consistent with mutation-

order effects on speciation, where populations fix advantageous mutations randomly rather than in

response to ecological differences (Schluter 2009). However, we cannot disregard the potential role

of ecological differences in generating reproductive incompatibilities for two main reasons: other

non-measured ecological variables may be the main selective force driving the association, and the

relatively small sample size of soil data reduces our power to detect an effect.

Phenotypic Divergence does not Predict Reproductive Isolation in Headland Populations

We next asked whether subtle phenotypic variation among Headland populations could explain

their reproductive isolation patterns. If divergent rather than uniform selection drives intrinsic

reproductive isolation between Headlands, populations with greater phenotypic differences should

show stronger reproductive barriers. However, after controlling for genetic distance, we found no

significant relationship between phenotypic divergence and reproductive isolation for either F_1 seed

set ($F_{1,2} = 12.88, P = 0.070$) or F_1 viability ($F_{1,2} = 4.45, P = 0.170$). The absence of a

correlation between phenotypic divergence and intrinsic reproductive isolation further supports

mutation-order effects on speciation, where genetic incompatibilities independently arise and

contribute to the same adaptive phenotypes across populations (Schluter 2009). However, we again

cannot disregard a possible association between phenotypic differences and reproductive

incompatibilities due to potential unmeasured ecological variables and relatively small sample sizes

for this analysis.

Parallel Mosaic Speciation in *Senecio*

The patterns of reproductive isolation in the *Senecio lautus* species complex present an evolutionary

paradox that challenges our traditional understanding of parallel speciation (Figure 6). While

divergent natural selection in contrasting coastal environments has repeatedly driven the evolution

of reproductive barriers between Dune and Headland ecotypes (Melo et al. 2014; Richards and

Ortiz-Barrientos 2016; Richards et al. 2016; Walter et al. 2016, 2018b; Wilkinson et al. 2021), the

pattern within each ecotype reveals an unexpected asymmetry. In Dunes, populations experience

relatively uniform selection due to consistent environmental pressures (Roda et al. 2013b; Walter et

al. 2016), allowing for reproductive compatibility despite geographic separation of thousands of

kilometers and ancient divergence. Crosses between distant Dune populations even exhibit

enhanced fitness through hybrid vigor. In contrast, Headland populations have evolved reproductive

barriers in crosses between the phylogenetic clades, despite convergence on similar prostrate

phenotypes. This suggests that the repeated evolution of similar phenotypes may obscure profound

differences in how populations traverse adaptive landscapes. These incompatibilities between

Headlands do not correlate with environmental or morphological differences, suggesting that

mutation-order processes are at play. However, environmental heterogeneity (Roda et al. 2013b;

Walter et al. 2016) and localized selective pressures among Headland populations indicate that

patterns of reproductive isolation do not strictly align with mutation-order speciation.

In *Senecio*, this combination of divergent, uniform and heterogeneous selection forms what we call

'parallel mosaic speciation'—a continuum of speciation patterns where reproductive isolation arises

through multiple mechanisms (also see Langerhans and Riesch 2013). While theoretical

frameworks have traditionally treated mutation-order and ecological speciation as distinct processes

(Schluter 2009; Nosil and Flaxman 2011), empirical evidence suggests natural systems operate

along a continuum of selective pressures (Johannesson et al. 2024). This complexity is particularly

evident in plants, such as the *Senecio* system, where fine-scale environmental heterogeneity creates

intricate selective mosaics that challenge the binary classification of uniform versus divergent

selection (Lowry et al. 2019; James et al. 2023b). Here our parallel mosaic speciation framework

explicitly incorporates within-ecotypic environmental heterogeneity to illustrate how various forms

of natural selection shape patterns of reproductive isolation across heterogeneous landscapes.

Parallel mosaic speciation creates several theoretical challenges: How do systems of parallel

evolution evolve different forms of reproductive isolation within and between ecotypes? How can

populations that evolved nearly identical phenotypes accumulate reproductive barriers? And what is

the effect of genetic architecture and environmental variance in creating patterns of reproductive

isolation during parallel speciation? Addressing these questions requires an integrated theoretical

framework to explain parallel speciation under mixed selection regimes and complex genetic

architectures. Although theory has addressed components of these problems (e.g., Orr 1995; Barton

2001; Orr and Turelli 2001; Porter and Johnson 2002; Gavrilets 2004; Palmer and Feldman 2009;

Fierst and Hansen 2010; Nosil and Flaxman 2011; Bank et al. 2012; Chevin et al. 2014; Fraïsse et

al. 2014; Yamaguchi and Otto 2020; Thompson et al. 2024), many models oversimplify the genetic

architectures underlying adaptation. Additionally, current frameworks inadequately account for how

genetic architecture interacts with environmental heterogeneity to drive the accumulation of

reproductive barriers. By integrating these factors, we can better understand how speciation unfolds

across heterogeneous landscapes, moving us beyond the simple dichotomy of ecological vs.

mutation-order speciation.

Environmental Symmetry Determines the Probability of Incompatibility Formation

We developed a mathematical framework demonstrating how environmental symmetry and genetic

architecture shape reproductive isolation during parallel mosaic speciation (Table 1). We found that

the probability of Dobzhansky-Muller Incompatibility (DMI) formation is governed by the

interaction of three key parameters: the distribution of selection coefficients (s), environmental

symmetry (φ), and the genetic architecture (n) of the interacting loci. The classic Unckless-Orr

model ($P_{DM,U}$) demonstrates that under identical environments, the probability of DMI formation

reaches a maximum when selection coefficients are equal (Table 2). This upper bound reflects

maximum stochasticity in adaptive trajectories: when mutations confer equal fitness benefits ($s_A =$

s_B), either allele is equally likely to fix first in separate populations, maximizing the probability of

incompatible combinations.

 The Role of the Environment in Mosaic Speciation

We extended the Unckless-Orr model by incorporating environmental symmetry (φ), to reveal how

environmental heterogeneity systematically modifies DMI formation probability through its effects

on relative selection pressures (Figure 5A). This environmental similarity parameter enables us to

model speciation scenarios beyond the binary categorisation of identical versus contrasting

environments, offering a more nuanced perspective on how environmental gradients drive

reproductive isolation. This extended model (Equation 2, $P_{DM,E}$) implies the following. First, when

the environments are identical between populations $(\varphi = 1)$, $P_{DM,E}$ reduces to $P_{DM,U}$, recovering

classic predictions about the relationship between selection coefficients and reproductive isolation.

This is akin to the *Senecio* Dune populations, where populations remain reproductively compatible

due to nearly identical selective pressures. Second, as environments become increasingly different

between populations $(\varphi \rightarrow 0)$, the probability of DMI formation approaches unity regardless of

selection coefficient ratios, demonstrating how environmental differences can override

deterministic selection effects, as seen between the Dune and Headland ecotypes. Third, for

intermediate symmetry $(0 < \varphi < 1)$, the probability of DMI formation increases monotonically

with decreasing φ while becoming progressively less dependent on the ratio s_B/s_A. This is similar to

the Headland populations, which experience some degree of environmental heterogeneity.

When $\varphi \neq 1$, our model predicts that stochastic fixation of different alleles can also lead to DMIs,

even under partial environmental overlap. This result parallels the findings of Yamaguchi and Otto

(2020), which reveal the speed and direction of environmental change are pivotal in promoting

speciation, particularly in cases where populations experience similar but shifting selection

pressures. Yamaguchi and Otto (2020) further demonstrate that periods of rapid environmental

change increase the likelihood of large-effect mutations becoming fixed, thereby enhancing the

probability of DMIs between populations. In the context of parallel evolution, loci controlling

convergent traits are predicted to exhibit parallel molecular evolution when selection coefficients

are highly asymmetric $(s_A \gg s_B)$ and environmental symmetry is minimal $(\varphi \rightarrow 1)$. This pattern

is supported by systems such as the parallel loss of armor plates in marine sticklebacks (Jones et al.

2012) and mimicry rings in butterflies (Van Belleghem et al. 2017).

Complex Genetic Architectures Enhance the Probability of Reproductive Isolation

Our extension of the model to polygenic traits (Equation 3, $P_{DM,P}$) reveals that genetic complexity

amplifies the probability of reproductive isolation. $P_{DM,P}$ captures how combinatorial growth in

potential interactions amplifies the effects of selection asymmetry (Figure 5B). Specifically, we

show that polygenic adaptation amplifies the effects of selection asymmetry, particularly when

multiple loci with varying selection coefficients contribute to adaptation. Our framework explicitly

accounts for the combinatorial growth of potential incompatibilities with increasing numbers of

loci, explaining why polygenic traits readily accumulate DMIs despite weak selection asymmetry.

These results align with Yamaguchi and Otto (2020), who show how pleiotropic mutations, which

affect multiple traits, are likely to generate incompatibilities under conditions of rapid adaptation.

The *Senecio* system exemplifies these findings: Headland populations achieve phenotypic

convergence through distinct combinations of alleles at gravitropism and architectural loci,

facilitated by conditional neutrality and asymmetric selection ($\varphi < 1$; James et al. 2023a; Kaur

2024).

Our polygenic framework also aligns with the work of Schneemann et al. (2020), who explore how

the geometry of population divergence in a multi-trait space influences hybrid fitness,

distinguishing between intrinsic and extrinsic components of reproductive isolation. Schneemann et

al.'s observation that complex trait architectures can either enhance or mitigate hybrid

incompatibilities, depending on the distribution of adaptive alleles, closely mirrors our findings of

how polygenic architectures and adaptive divergence influence the likelihood and strength of hybrid

incompatibilities. Furthermore, their distinction between intrinsic and extrinsic isolation supports

our observation that mosaic speciation arises from a dynamic interplay between local adaptation

and stochastic divergence. However, as with any proposed mechanistic model, loci contributing to

adaptation will not contribute to intrinsic reproductive isolation unless they allow the evolution of

DMI systems that require conditional neutrality to create a genetic correlation between adaptation

and speciation. See *Supplementary Material 3* for a mechanistic model that connects patterns of

reproductive compatibility to parallel evolution in *Senecio*.

Our mathematical framework provides insights into how populations move along the adaptive

landscape. When φ approaches 1, populations traverse similar adaptive landscapes, minimizing

opportunities for incompatibility evolution. As φ decreases, increasing landscape complexity

creates multiple possible adaptive trajectories, facilitating the accumulation of reproductive barriers

even between phenotypically convergent populations, or arriving to different adaptive peaks. These

theoretical predictions align with empirical observations in several systems, including Caribbean

Anolis lizards (Mahler et al. 2013), where fine-scale habitat differences create multiple adaptive

solutions despite broad-scale ecological similarities, or in bacteria (Nahum et al. 2015) where

environmental heterogeneity can drive populations toward distinct adaptive peaks despite

experiencing similar selective pressures.

Together, our mathematical framework demonstrates how uniform, similar and divergent selection

can simultaneously shape patterns of reproductive isolation during parallel speciation. The

framework unifies ecological and mutation-order mechanisms through a single parameter (φ) while

revealing how genetic architecture modulates the translation of selection asymmetry into

reproductive barriers. These results provide a quantitative foundation for understanding the

emergence of biodiversity across heterogeneous landscapes such as in systems like *Senecio*, where

environmental variation creates complex mosaics of selection pressures, and therefore of speciation

during parallel evolution. Future research should extend the model to include additional

complexities, such as recombination dynamics, diploid populations, dominance effects and

temporal fluctuations in selection pressures.

Synthesis and Broader Implications

Mechanisms Generating Genetic Incompatibilities During Parallel Speciation

During parallel evolution, reproductive isolation can emerge through several key processes. As

shown by our parallel mosaic speciation framework, reproductive incompatibilities can arise via the

stochastic fixation of alternative beneficial alleles under similar selective pressures (as with

mutation-order speciation), or the deterministic fixation of alternative alleles under divergent

selective pressures (as with ecological speciation), or a combination of both (parallel mosaic

speciation). These forms of speciation, where reproductive isolation accumulates as a byproduct of

natural selection, likely underlie the patterns observed in *Senecio*.

Our theoretical framework demonstrates that genetic incompatibilities may arise through the

divergence of pleiotropic developmental pathways that affect both adaptive traits and reproductive

compatibility. This mechanism generates several predictions: reproductive barriers should involve

genes with dual roles in development and reproduction, when selection coefficients are similar

across interacting developmental components, populations can reach similar phenotypic outcomes

through different genetic architectures, and hybrids should show a genetic correlation between

adaptive and reproductive isolation traits. The *Senecio* system is consistent with this mechanism.

Previous work in *Senecio* has found that hormone signaling pathways, particularly auxin regulation,

appear to coordinate adaptive traits (gravitropism, architecture; Wilkinson et al. 2021; James et al.

2023a; Broad et al. 2024) and reproductive processes (fertilization success; Wilkinson et al. 2021).

Because hormone pathways involve multiple interacting components (McGlothlin and Ketterson

2008) with potentially similar selection coefficients, they provide opportunities for selection to

generate different genetic solutions to the same challenges, as seen in other systems of replicated

evolution (e.g., Rokas and Carroll 2008; Reid et al. 2016; Birkeland et al. 2020). This integration of

deterministic and stochastic processes might explain how *Senecio* populations achieve similar

phenotypes through different genetic architectures while accumulating reproductive barriers.

Reproductive isolation can also accumulate between populations via neutral process (True and

Haag 2001; Gavrilets 2004; Palmer and Feldman 2009). For instance, system drift is a form of

genetic drift where populations traverse a "fitness ridge" of functionally equivalent genetic

networks. As populations move along this ridge via stochastic processes, they may develop genetic

incompatibilities while maintaining their phenotype and fitness (True and Haag 2001; Schiffman

and Ralph 2021; James et al. 2022). Under this process, we expect three key patterns: no systematic

relationship between reproductive isolation and ecological divergence, stronger isolation in

populations with smaller effective sizes, and random accumulation of incompatibilities with respect

to adaptive traits. Our data provide limited support for this mechanism in *Senecio*. Reproductive

isolation shows consistent ecotype-dependence rather than random accumulation, effective

population sizes remain large across ecotypes (James et al. 2021a), and Headland populations

exhibit clear signatures of local adaptation rather than drift. Furthermore, most candidate speciation

genes in plants (Rieseberg and Blackman 2010) do not support system drift as a primary driver.

Instead, they reflect deterministic evolution due to selection on ecological and reproductive traits.

However, specific cases, such as certain S-RNase self-incompatibility genes controlling unilateral

incompatibility (Murfett et al. 1996; Hancock et al. 2003) and rare cytoplasmic male sterility loci

(Bentolila et al. 2002; Wang et al. 2006), could represent drift-driven incompatibilities, particularly

in species or populations with small effective sizes. Even so, these examples are exceptions rather

than the rule, suggesting that drift plays a limited role in the evolution of reproductive isolation in

plant systems.

Genetic conflict and sexual selection are key mechanisms known to generate intrinsic

incompatibilities in plants (Moore and Pannell 2011; Crespi and Nosil 2013; Tonnabel et al. 2021;

Coughlan 2023). This can arise through divergent evolution of mating systems (e.g., Martin and

Willis 2007), parent-of-origin effects on hybrid fitness (e.g., Coughlan et al. 2020), or antagonistic

coevolution between male and female reproductive traits (e.g., Arnqvist et al. 2000). Recent work

on conspecific pollen precedence in *Senecio* revealed that local adaptation influences competitive

gametic interactions controlled by females (Arenas-Castro 2022), suggesting sexual selection might

contribute to reproductive isolation. However, we found no significant asymmetries in hybrid

viability or fertility that would indicate strong parent-of-origin effects. The absence of pronounced

reciprocal cross differences suggests that while sexual selection may fine-tune reproductive

barriers, it is unlikely to be the primary driver of isolation in this system.

Release from Trade-offs and the Evolution of Reproductive Isolation

Trade☐offs arise when a set of alleles or traits that confer advantages in one environment impose

costs in another (Stearns 1989). During the colonization of a new habitat, these constraints can be

relaxed or entirely lifted if the selective context shifts, thereby freeing previously suboptimal or

deleterious alleles to persist or even become beneficial (Lahti et al. 2009). In the extended

Unckless–Orr model, this release is captured mathematically by allowing the parameter φ to deviate

from 1. When φ is significantly less than 1, ancestral constraints no longer fully operate in the new

habitat, enabling novel sets of mutations to fix without incurring previous fitness penalties. This

release from ancestral trade☐offs can accelerate adaptation because it expands the genetic

"toolbox" populations have at their disposal. As a result, different populations can fix different

beneficial alleles for what appears to be the same ecological task, a phenomenon seen particularly

in Headland environments of *Senecio*. Despite convergent prostrate morphologies, Headland

populations have accumulated genetic differences that reduce hybrid fitness. These apparently

paradoxical patterns arise when the new habitat only partially resembles the ancestral one, making

φ an intermediate value rather than strictly 1 or 0. Such intermediate environmental similarity

provides enough shared selective pressures for similar phenotypes to evolve while simultaneously

allowing distinct adaptive trajectories, ultimately yielding a mosaic of reproductive barriers across

the range. In this way, the relaxation of trade☐offs not only promotes phenotypic innovation but

can also generate the conditions necessary for the formation of DMIs under parallel mosaic

speciation.

Effect of Gene Flow on the Likelihood of Parallel Speciation

Population connectivity further complicates the dynamics of speciation because gene flow between

populations homogenizes allele frequencies and reduces the likelihood of independent fixation of

alternative beneficial alleles (Bank et al. 2012). For example, in populations adapted to similar

environments, mutation-order speciation is challenging under moderate gene flow (Schluter 2009;

Nosil and Flaxman 2011). This is because advantageous mutations are likely to spread and fix in all

populations, thereby preventing the accumulation of incompatibilities. Consequently, mutation-

order speciation typically requires very low or absent gene flow (Nosil and Flaxman 2011; Nosil

2012). In *Senecio*, the negligible gene flow in the system (James et al. 2021a) has created ideal

conditions for intrinsic incompatibilities to accumulate among Headland populations.

During ecological speciation, where alternative alleles are favored in different environments, there

is an antagonism between gene flow and divergent natural selection (Felsenstein 1981). As with

mutation-order speciation, gene flow constrains the evolution of reproductive barriers during

adaptation, yet speciation remains possible if mechanisms arise to resolve this conflict (also see

(Porter and Johnson 2002). If locally adaptive alleles reside in genomic regions of reduced

recombination, such as chromosomal inversions, population divergence and speciation can proceed

despite ongoing gene flow (Noor et al. 2001; Rieseberg 2001; Butlin 2005; Kirkpatrick and Barton

2006; Ortiz-Barrientos et al. 2016). The role of chromosomal rearrangements in mitigating the

antagonism between gene flow and selection has been demonstrated across numerous systems

including sticklebacks (Samuk et al. 2017), Atlantic cod (Berg et al. 2017) *Heliconius* butterflies

(Martin et al. 2019), *Mimulus* monkeyflowers (Lowry and Willis 2010), *Littorina* snails (Le Moan

et al. 2024) and *Drosophila* (Poikela et al. 2024). In the *Senecio* system, the low levels of

contemporary gene flow between the Dune and Headland ecotypes likely facilitated the evolution of

incompatibilities, yet ancient gene flow may also have shaped their divergence (James et al. 2021a).

Future work will explore the role of chromosomal inversions during parallel evolution in *Senecio*,

shedding light on their contributions to speciation dynamics.

Patterns of Intrinsic Reproductive Isolation across Traits and Development

Reproductive isolation can manifest differently across traits and developmental stages due to

varying selective pressures and genetic complexities (Cutter 2023). For instance, while natural

selection may act divergently at the population level, individual traits may experience selection that

ranges from uniform to highly divergent (Langerhans and Riesch 2013). Additionally, the type and

strength of selection upon traits may change over the course of development of the organism, such

as when trade-offs exist where traits favoured early in development are not beneficial in later life

(Stearns 1989; Garland et al. 2022). Finally, shared genetic architectures among traits can impose

pleiotropic constraints, shaping how reproductive barriers evolve and influencing the potential for

independent adaptation of specific traits (Mauro and Ghalambor 2020; Zhang 2023).

In *Senecio*, we found that reproductive barriers manifest differently in seed set versus viability.

Crosses between Headland populations from different clades show reduced F_1 seed set, supporting

our theoretical predictions about DMI formation under similar selection pressures. However, hybrid

viability does not follow this pattern, suggesting that different components of reproductive isolation

may evolve through distinct genetic architectures. This asymmetry in reproductive barriers aligns

with our mathematical framework: the probability of parallel DMI formation critically depends on

both the number of interacting loci and the distribution of selection coefficients. Seed set may be

governed by a small number of loci with large effect sizes, such as those involved in specific

pollen-pistil recognition and hormone signaling pathways, making parallel selection more likely to

generate consistent reproductive barriers. In contrast, hybrid viability likely involves a greater

number of loci with smaller, more diffuse effects across genetic networks with an extensive

repertoire to disrupt development, increasing the likelihood of different outcomes across the

population pairs. This distinction between reproductive barriers requires further exploration to help

explain if different components of isolation can evolve through distinct mechanistic pathways even

under similar selective pressures.

A Continuous Framework for Speciation

Our work expands our understanding of how incompatibilities arise during uniform selection,

proposing a broader concept of mutation-order speciation reframed as stochastic speciation. This

reframing incorporates some degree of environmental heterogeneity as well as the dynamics of

conditional neutrality, emphasizing not only the sequence in which beneficial mutations fix but also

the potential for different beneficial alleles to achieve similar adaptive outcomes in comparable

environments – an aspect often overlooked in traditional theories of mutation-order speciation.

While pure mutation-order speciation can theoretically occur under identical environments, it

becomes increasingly improbable with increasing genetic complexity. Most apparent cases of

speciation during adaptation to "identical" environments likely involve interactions among multiple

mechanisms, including alternative adaptive solutions and subtle environmental heterogeneity. This

may explain why the most convincing cases of mutation-order speciation arise in experimental

systems (Ono et al. 2017; Hsu et al. 2024), where environmental conditions can be controlled to

ensure they are effectively identical. To fully understand the relative contributions of stochastic

fixation versus ecological adaptation in natural systems, identifying the specific loci underlying

reproductive barriers and their link to local adaptation is crucial (Wu and Ting 2004; Lowry et al.

2008; Nosil and Schluter 2011). Only by characterizing the molecular basis of incompatibilities can

we determine whether they arose through the random fixation of alternative beneficial alleles or

through adaptation to subtle ecological differences. This will help clarify the extent to which

apparent cases of mutation-order speciation in nature truly reflect stochastic processes versus

environmental heterogeneity.

The parallel mosaic speciation framework we propose challenges traditional distinctions between

ecological and mutation-order speciation (Schluter and Nagel 1995; Ostevik et al. 2012), positing

that these modes represent endpoints of a continuous spectrum rather than discrete processes

(Langerhans and Riesch 2013). This aligns with recent work showing how speciation dynamics can

exhibit "tipping points" where small changes trigger major transitions, suggesting a continuous but

potentially nonlinear process (Nosil et al. 2017). Furthermore, local adaptation inevitably involves

some degree of reproductive isolation, further supporting this continuous view of speciation (Butlin

and Faria 2024). Our framework illustrates how reproductive isolation progresses across gradients

of environmental similarity: from deterministic adaptation in identical environments, to adaptation

driven by subtle environmental heterogeneity, to divergent adaptation in contrasting environments,

where reproductive isolation becomes nearly inevitable. By incorporating environmental symmetry

and genetic architecture, our mathematic models help predict speciation outcomes in systems with

partial reproductive isolation or varying degrees of environmental heterogeneity. Explicitly treating

these factors as continuous variables moves beyond the simple dichotomies of parallel versus

divergent evolution towards a more nuanced understanding of speciation mechanisms.

Recent theoretical advances, such as the "speciation hypercube" framework (Bolnick et al. 2023;

Johannesson et al. 2024), emphasize the multidimensionality of speciation, integrating genetic,

phenotypic, ecological, and temporal factors. In contrast to the traditional "speciation continuum"

model (Seehausen et al. 2014; Stankowski and Ravinet 2021), our mathematical analysis

complements these multidimensional perspectives by highlighting how parameter-level

interactions—including environmental symmetry, selection coefficients, and genetic architecture—

generate quantitatively predictable patterns of reproductive isolation. This mechanistic approach

bridges descriptive speciation models and evolutionary processes, offering insights into the relative

contributions of deterministic versus stochastic processes in driving divergence. We urge

researchers to move beyond the categorical classification of ecotypes and instead incorporate

within-ecotype environmental variation to better understand the role of environmental heterogeneity

in the evolution of reproductive incompatibilities.

CONCLUSIONS

Our study demonstrates how multiple forms of selection jointly drive the parallel evolution of

reproductive isolation in *Senecio lautus*. By integrating field experiments, genetic analyses, and an

updated theoretical framework, we reveal that divergent selection between contrasting

environments, uniform selection in identical dune environments and similar selection in headland

environments together shape speciation dynamics. These findings of parallel mosaic speciation

challenge the traditional dichotomy between ecological and mutation-order speciation, highlighting

their complementary roles in generating biodiversity.

Incorporating polygenic adaptation and environmental symmetry into our mathematic framework

demonstrates that genetic complexity increases the probability of speciation, even without stark

ecological contrasts. These findings underscore the importance of genetic architecture in shaping

evolutionary outcomes and reveals how phenotypically convergent populations can obscure

profound genetic divergence. Future work integrating genomic and ecological data across broader

landscapes will further illuminate how divergent and stochastic processes jointly drive the origins of

species. Our findings contribute to a growing understanding of how biodiversity emerges and
persists across heterogeneous environments.

AUTHOR CONTRIBUTIONS

Conceptualization: Daniel Ortiz-Barrientos. Formal Analysis: Maddie James, Maria Melo, and Greg
Walter. Mathematical Model: Daniel Ortiz-Barrientos and Jan Engelstädter. Experiments: Maria
Melo, Maddie James, Federico Roda, Diana Bernal-Franco, Melanie Wilkinson, and Huanle Liu.
Data Curation: Maddie James, Maria Melo. Visualization: Maddie James, Greg Walter, and Daniel
Ortiz-Barrientos. Writing: Maddie James, and Daniel Ortiz-Barrientos, with contributions from
Greg Walter, Melanie Wilkinson, Maria Melo, Federico Roda, Diana Bernal-Franco, Huanle Liu,
and Jan Engelstädter. Supervision: Daniel Ortiz-Barrientos. Funding Acquisition: Daniel Ortiz-
Barrientos and Jan Engelstädter.

ACKNOWLEDGEMENTS

The authors acknowledge the Turrbal and Yuggera, Ningy Ningy, Barunggam, Bundjalung,
Gumbaynggirr, Gungarri, Gunaikurnai, Gunditjmara, Wirangu, Paredarerme, and
Tyerrernotepanner peoples, the traditional owners of the land on which this work was undertaken,
and pay their respects to Elders past, present, and emerging. We thank Mohamed Noor, Kieran
Samuk, Kate Ostevik, Loren Rieseberg, Eric Baack, Matthew Hahn, and Thomas Richards for their
critical discussions on the empirical foundations of this work. We acknowledge the use of ChatGPT
v4 Turbo and Claude v3.5 Sonnet assist with minor grammatical edits to the main text, and code
development for visualization of the theoretical framework. This research was supported by grants
from the Australian Research Council to Daniel Ortiz-Barrientos (DP120104559), Daniel Ortiz-
Barrientos and Jan Engelstädter (DP190103039), and Daniel Ortiz-Barrientos through The
Australian Research Council Centre of Excellence for Plant Success in Nature and Agriculture
(CE200100015).

CONFLICTS OF INTEREST

815 The authors declare no conflicts of interest.

DATA ACCESSIBILITY

817 All data sets and code will be uploaded to GitHub upon acceptance of the manuscript.

REFERENCES

Ali, S. 1964. *Senecio lautus* complex in Australia. I. Taxonomic considerations and discussion of
some of the related taxa from New Zealand. Aust. J. Bot. 12:282–291.
Anderson, S. A. S., and J. T. Weir. 2022. The role of divergent ecological adaptation during
allopatric speciation in vertebrates. Science 378:1214–1218.
Arenas-Castro, H. 2022. The evolution of gametic interactions during speciation. The University of
Queensland. https://doi.org/10.14264/1627e2b.
Arnqvist, G., M. Edvardsson, U. Friberg, and T. Nilsson. 2000. Sexual conflict promotes speciation
in insects. Proceedings of the National Academy of Sciences 97:10460–10464.
Bank, C., R. Bürger, and J. Hermisson. 2012. The limits to parapatric speciation: Dobzhansky–
Muller Incompatibilities in a continent–island model. Genetics 191:845–863.
Barton, N. H. 2001. The role of hybridization in evolution. Molecular Ecology 10:551–568.
Barton, N. H., and G. M. Hewitt. 1985. Analysis of hybrid zones. Annu. Rev. Ecol. Syst. 16:113–
148.
Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using
lme4. J. Stat. Softw. 67:1–48.
Bentolila, S., A. A. Alfonso, and M. R. Hanson. 2002. A pentatricopeptide repeat-containing gene
restores fertility to cytoplasmic male-sterile plants. Proc Natl Acad Sci U S A 99:10887–
10892.
Berg, P. R., B. Star, C. Pampoulie, I. R. Bradbury, P. Bentzen, J. A. Hutchings, S. Jentoft, and K. S.
Jakobsen. 2017. Trans-oceanic genomic divergence of Atlantic cod ecotypes is associated
with large inversions. Heredity 119:418–428.
Bierne, N., P. A. Gagnaire, and P. David. 2013. The geography of introgression in a patchy
environment and the thorn in the side of ecological speciation. Curr. Zool. 59:72–86.
Birkeland, S., A. L. S. Gustafsson, A. K. Brysting, C. Brochmann, and M. D. Nowak. 2020.
Multiple genetic trajectories to extreme abiotic stress adaptation in Arctic Brassicaceae.
Molecular Biology and Evolution 37:2052–2068.
Bischoff, A., and H. Müller-Schärer. 2010. Testing population differentiation in plant species – how
important are environmental maternal effects. Oikos 119:445–454.
Blanchet, G., M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R. B. O'Hara, G. L.
Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner. 2018. vegan:
Community Ecology Package. R package version 2.5-2. https://CRAN.R-
project.org/package=vegan.
Bolnick, D. I., A. K. Hund, P. Nosil, F. Peng, M. Ravinet, S. Stankowski, S. Subramanian, J. B. W.
Wolf, and R. Yukilevich. 2023. A multivariate view of the speciation continuum. Evolution
77:318–328.
Broad, Z., J. Lefevre, M. J. Wilkinson, S. Barton, F. Barbier, H. Jung, D. Donovan, and D. Ortiz-
Barrientos. 2024. Gravitropic gene expression divergence associated with adaptation to
contrasting environments in an Australian wildflower. Mol Ecol e17543.
Butlin, R. K. 2005. Recombination and speciation. Mol. Ecol. 14:2621–2635.
Butlin, R. K., and R. Faria. 2024. Local adaptation and reproductive isolation: when does speciation
start? Evolutionary Journal of the Linnean Society 3:kzae003.
Chevin, L.-M., G. Decorzent, and T. Lenormand. 2014. Niche dimensionality and the genetics of
ecological speciation. Evolution 68:1244–1256.
Clarke, J. D. 2009. Cetyltrimethyl Ammonium Bromide (CTAB) DNA miniprep for plant DNA
isolation. Cold Spring Harb. Protoc. 2009:Pdb.prot5177.

Colosimo, P. F., K. E. Hosemann, S. Balabhadra, G. Villarreal, M. Dickson, J. Grimwood, J.
Schmutz, R. M. Myers, D. Schluter, and D. M. Kingsley. 2005. Widespread parallel
evolution in sticklebacks by repeated fixation of Ectodysplasin alleles. Science 307:1928–
1933.

Coughlan, J. M. 2023. The role of conflict in shaping plant biodiversity. New Phytologist
240:2210–2217.

Coughlan, J. M., M. Wilson Brown, and J. H. Willis. 2020. Patterns of hybrid seed inviability in the
Mimulus guttatus sp. complex reveal a potential role of parental conflict in reproductive
isolation. Current Biology 30:83-93.e5.

Coyne, J. A., and H. A. Orr. 1997. "Patterns of speciation in *Drosophila*" revisited. Evolution
51:295–303.

Coyne, J. A., and H. A. Orr. 2004. Speciation. Sunderland: Sinauer Associates.

Crespi, B., and P. Nosil. 2013. Conflictual speciation: species formation via genomic conflict.
TREE 28:48–57.

Cutter, A. D. 2023. Speciation and development. Evolution & Development 25:289–327.

Drummond, A. J., M. A. Suchard, D. Xie, and A. Rambaut. 2012. Bayesian phylogenetics with
BEAUti and the BEAST 1.7. Mol. Biol. Evol. 29:1969–1973.

Earl, D. A., and B. M. vonHoldt. 2012. STRUCTURE HARVESTER: a website and program for
visualizing STRUCTURE output and implementing the Evanno method. Conserv. Genet.
Resour. 4:359–361.

Endler, J. 1977. Geographic variation, speciation, and clines. Princeton: Princeton University Press.

Evanno, G., S. Regnaut, and J. Goudet. 2005. Detecting the number of clusters of individuals using
the software structure: a simulation study. Mol. Ecol. 14:2611–2620.

Falush, D., M. Stephens, and J. K. Pritchard. 2003. Inference of population structure using
multilocus genotype data: linked loci and correlated allele frequencies. Genetics 164:1567–
1587.

Felsenstein, J. 1981. Skepticism towards Santa Rosalia, or why are there so few kinds of animals?
Evolution 35:124–138.

Fierst, J. L., and T. F. Hansen. 2010. Genetic architecture and postzygotic reproductive isolation:
Evolution of Bateson-Dobzhansky-Muller incompatibilities in a polygenic model. Evolution
64:675–693.

Fraïsse, C., J. a. D. Elderfield, and J. J. Welch. 2014. The genetics of speciation: are complex
incompatibilities easier to evolve? Journal of Evolutionary Biology 27:688–699.

Garland, T., C. J. Downs, and A. R. Ives. 2022. Trade-offs (and constraints) in organismal biology.
Physiological and Biochemical Zoology 95:82–112.

Garner, A. G., A. M. Kenney, L. Fishman, and A. L. Sweigart. 2016. Genetic loci with parent-of-
origin effects cause hybrid seed lethality in crosses between *Mimulus* species. New Phytol.
211:319–331.

Gavrilets, S. 2004. Fitness landscapes and the origin of species. Princeton University Press.

Gilbert, K. J., R. L. Andrew, D. G. Bock, M. T. Franklin, N. C. Kane, J.-S. Moore, B. T. Moyers, S.
Renaut, D. J. Rennison, T. Veen, and T. H. Vines. 2012. Recommendations for utilizing and
reporting population genetic analyses: the reproducibility of genetic clustering using the
program STRUCTURE. Mol. Ecol. 21:4925–4930.

Grant, P. R., and B. R. Grant. 2008. How and why species multiply: The radiation of Darwin's
finches. Princeton University Press.

Hancock, C. N., K. Kondo, B. Beecher, and B. McClure. 2003. The S-locus and unilateral
incompatibility. Philos Trans R Soc Lond B Biol Sci 358:1133–1140.
Harris, C. R., K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau, E.
Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk,
M. Brett, A. Haldane, J. F. del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K.
Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, and T. E. Oliphant. 2020. Array
programming with NumPy. Nature 585:357–362.
Hasegawa, M., H. Kishino, and T. Yano. 1985. Dating of the human-ape splitting by a molecular
clock of mitochondrial DNA. J. Mol. Evol. 22:160–174.
Heled, J., and A. J. Drummond. 2010. Bayesian inference of species trees from multilocus data.
Mol. Biol. Evol. 27:570–580.
Hendry, A. P., D. I. Bolnick, D. Berner, and C. L. Peichel. 2009. Along the speciation continuum in
sticklebacks. Journal of Fish Biology 75:2000–2036.
Hsu, S.-K., W.-Y. Lai, J. Novak, F. Lehner, A. M. Jakšić, E. Versace, and C. Schlötterer. 2024.
Reproductive isolation arises during laboratory adaptation to a novel hot environment.
Genome Biology 25:141.
Hunter, J. D. 2007. Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering
9:90–95.
Jakobsson, M., and N. A. Rosenberg. 2007. CLUMPP: a cluster matching and permutation program
for dealing with label switching and multimodality in analysis of population structure.
Bioinformatics 23:1801–1806.
James, M. E., R. N. Allsopp, J. S. Groh, A. Kaur, M. J. Wilkinson, and D. Ortiz-Barrientos. 2023a.
Uncovering the genetic architecture of parallel evolution. Molecular Ecology 32:5575–5589.
James, M. E., H. Arenas-Castro, J. S. Groh, S. L. Allen, J. Engelstädter, and D. Ortiz-Barrientos.
2021a. Highly replicated evolution of parapatric ecotypes. Mol. Biol. Evol. 38:4805–4821.
James, M. E., T. Brodribb, I. J. Wright, L. H. Rieseberg, and D. Ortiz-Barrientos. 2023b. Replicated
evolution in plants. Annu. Rev. Plant Biol. 74:697–725.
James, M. E., N. L. V. O'Brien, and A. Bukkuri. 2022. Digest: Stable phenotypes, fluid genotypes:
how stochasticity impacts network evolution and speciation. Evolution evo.14452.
James, M. E., M. J. Wilkinson, D. M. Bernal, H. Liu, H. L. North, J. Engelstädter, and D. Ortiz-
Barrientos. 2021b. Phenotypic and genotypic parallel evolution in parapatric ecotypes of
Senecio. Evolution 75:3115–3131.
Janes, J. K., J. M. Miller, J. R. Dupuis, R. M. Malenfant, J. C. Gorrell, C. I. Cullingham, and R. L.
Andrew. 2017. The $K = 2$ conundrum. Mol. Ecol. 26:3594–3602.
Johannesson, K., R. Faria, A. L. Moan, M. Rafajlović, A. M. Westram, R. K. Butlin, and S.
Stankowski. 2024. Diverse pathways to speciation revealed by marine snails. TiG 40:337–
351.
Johannesson, K., M. Panova, P. Kemppainen, C. André, E. Rolán-Alvarez, and R. K. Butlin. 2010.
Repeated evolution of reproductive isolation in a marine snail: unveiling mechanisms of
speciation. Philos. Trans. R. Soc. B 365:1735–1747.
Jones, F. C., Y. F. Chan, P. Russell, E. Mauceli, J. Johnson, R. Swofford, M. Pirun, M. C. Zody, S.
White, E. Birney, S. Searle, J. Schmutz, J. Grimwood, M. C. Dickson, R. M. Myers, C. T.
Miller, B. R. Summers, A. K. Knecht, S. D. Brady, H. Zhang, A. A. Pollen, T. Howes, C.
Amemiya, E. S. Lander, F. Di Palma, K. Lindblad-Toh, and D. M. Kingsley. 2012. The
genomic basis of adaptive evolution in threespine sticklebacks. Nature 484:55–61.

Kaur, A. 2024. The polygenic architecture of local adaptation in *Senecio lautus*. The University of Queensland. https://doi.org/10.14264/4cef722.

Kay, K. M., J. B. Whittall, and S. A. Hodges. 2006. A survey of nuclear ribosomal internal transcribed spacer substitution rates across angiosperms: An approximate molecular clock with life history effects. BMC Evol. Biol. 6:36.

Kirkpatrick, M., and N. Barton. 2006. Chromosome inversions, local adaptation and speciation. Genetics 173:419–434.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests in linear mixed effects models. J. Stat. Softw. 82:1–26.

Lahti, D. C., N. A. Johnson, B. C. Ajie, S. P. Otto, A. P. Hendry, D. T. Blumstein, R. G. Coss, K. Donohue, and S. A. Foster. 2009. Relaxed selection in the wild. Trends in Ecology & Evolution 24:487–496.

Langerhans, R. B., and R. Riesch. 2013. Speciation by selection: A framework for understanding ecology's role in speciation. Curr. Zool. 59:31–52.

Le Moan, A., S. Stankowski, M. Rafajlović, O. Ortega-Martinez, R. Faria, R. K. Butlin, and K. Johannesson. 2024. Coupling of twelve putative chromosomal inversions maintains a strong barrier to gene flow between snail ecotypes. Evolution Letters 8:575–586.

Lenormand, T., D. Roze, and F. Rousset. 2009. Stochasticity in evolution. Trends Ecol. Evol. 24:157–165.

Losos, J. B. 2011. Convergence, adaptation, and constraint. Evolution 65:1827–1840.

Lowry, D. B., J. T. Lovell, L. Zhang, J. Bonnette, P. A. Fay, R. B. Mitchell, J. Lloyd-Reilley, A. R. Boe, Y. Wu, F. M. Rouquette, R. L. Wynia, X. Weng, K. D. Behrman, A. Healey, K. Barry, A. Lipzen, D. Bauer, A. Sharma, J. Jenkins, J. Schmutz, F. B. Fritschi, and T. E. Juenger. 2019. QTL × environment interactions underlie adaptive divergence in switchgrass across a large latitudinal gradient. Proceedings of the National Academy of Sciences 116:12933–12941.

Lowry, D. B., J. L. Modliszewski, K. M. Wright, C. A. Wu, and J. H. Willis. 2008. The strength and genetic basis of reproductive isolating barriers in flowering plants. Philos T R Soc B 363:3009–3021.

Lowry, D. B., and J. H. Willis. 2010. A widespread chromosomal inversion polymorphism contributes to a major life-history transition, local adaptation, and reproductive isolation. PLoS Biol. 8:e1000500.

Mahler, D. L., T. Ingram, L. J. Revell, and J. B. Losos. 2013. Exceptional convergence on the macroevolutionary landscape in island lizard radiations. Science 341:292–295.

Mani, G. S., and B. C. Clarke. 1990. Mutational order: a major stochastic process in evolution. Proc. R. Soc. Lond. B 240:29–37.

Martin, N. H., and J. H. Willis. 2007. Ecological divergence associated with mating system causes nearly complete reproductive isolation between sympatric *Mimulus* species. Evolution 61:68–82.

Martin, S. H., J. W. Davey, C. Salazar, and C. D. Jiggins. 2019. Recombination rate variation shapes barriers to introgression across butterfly genomes. PLOS Biology 17:e2006288.

Mauro, A. A., and C. K. Ghalambor. 2020. Trade-offs, pleiotropy, and shared molecular pathways: A unified view of constraints on adaptation. Integrative and Comparative Biology 60:332–347.

McGlothlin, J. W., and E. D. Ketterson. 2008. Hormone-mediated suites as adaptations and evolutionary constraints. Philos Trans R Soc Lond B Biol Sci 363:1611–1620.

McKinnon, J. S., S. Mori, B. K. Blackman, L. David, D. M. Kingsley, L. Jamieson, J. Chou, and D. Schluter. 2004. Evidence for ecology's role in speciation. Nature 429:294–298.

McKinnon, J. S., and H. D. Rundle. 2002. Speciation in nature: the threespine stickleback model systems. TREE 17:480–488.

Melo, M. C., A. Grealy, B. Brittain, G. M. Walter, and D. Ortiz-Barrientos. 2014. Strong extrinsic reproductive isolation between parapatric populations of an Australian groundsel. New Phytol. 203:323–334.

Moore, J. C., and J. R. Pannell. 2011. Sexual selection in plants. Current Biology 21:R176–R182.

Murfett, J., T. J. Strabala, D. M. Zurek, B. Mou, B. Beecher, and B. A. McClure. 1996. S RNase and interspecific pollen rejection in the genus *Nicotiana*: multiple pollen-rejection pathways contribute to unilateral incompatibility between self-incompatible and self-compatible species. Plant Cell 8:943–958.

Nahum, J. R., P. Godfrey-Smith, B. N. Harding, J. H. Marcus, J. Carlson-Stevermer, and B. Kerr. 2015. A tortoise–hare pattern seen in adapting structured and unstructured populations suggests a rugged fitness landscape in bacteria. Proceedings of the National Academy of Sciences 112:7530–7535.

Noor, M. A. F., K. L. Grams, L. A. Bertucci, and J. Reiland. 2001. Chromosomal inversions and the reproductive isolation of species. Proceedings of the National Academy of Sciences 98:12084–12088.

Nosil, P. 2012. Ecological speciation. Oxford University Press.

Nosil, P., B. J. Crespi, and C. P. Sandoval. 2002. Host-plant adaptation drives the parallel evolution of reproductive isolation. Nature 417:440–443.

Nosil, P., J. L. Feder, S. M. Flaxman, and Z. Gompert. 2017. Tipping points in the dynamics of speciation. Nat Ecol Evol 1:1.

Nosil, P., and S. M. Flaxman. 2011. Conditions for mutation-order speciation. Proceedings of the Royal Society B: Biological Sciences 278:399–407.

Nosil, P., and D. Schluter. 2011. The genes underlying the process of speciation. Trends Ecol. Evol. 26:160–167.

Nosil, P., T. H. Vines, and D. J. Funk. 2005. Perspective: Reproductive isolation caused by natural selection against immigrants from divergent habitats. Evolution 59:705–719.

Ono, J., A. C. Gerstein, and S. P. Otto. 2017. Widespread genetic incompatibilities between first-step mutations during parallel adaptation of *Saccharomyces cerevisiae* to a common environment. PLoS Biology 15:e1002591.

Orr, H. A. 1995. The population genetics of speciation: The evolution of hybrid incompatibilities. Genetics 139:1805–1813.

Orr, H. A., and M. Turelli. 2001. The evolution of postzygotic isolation: accumulating Dobzhansky-Muller incompatibilities. Evolution 55:1085–1094.

Ortiz-Barrientos, D., J. Engelstädter, and L. H. Rieseberg. 2016. Recombination rate evolution and the origin of species. Trends Ecol. Evol. 31:226–236.

Ostevik, K. L., B. T. Moyers, G. L. Owens, and L. H. Rieseberg. 2012. Parallel ecological speciation in plants? Int. J. Ecol. 2012:939862.

Palmer, M. E., and M. W. Feldman. 2009. Dynamics of hybrid incompatibility in gene networks in a constant environment. Evolution 63:418–431.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: Analyses of phylogenetics and evolution in R language. Bioinformatics 20:289–290.

Poikela, N., D. R. Laetsch, V. Hoikkala, K. Lohse, and M. Kankare. 2024. Chromosomal inversions
and the demography of speciation in *Drosophila montana* and *Drosophila flavomontana*.
Genome Biology and Evolution 16:evae024.

Porter, A. H., and N. A. Johnson. 2002. Speciation despite gene flow when developmental
pathways evolve. Evolution 56:2103–2111.

Posada, D., and K. A. Crandall. 1998. MODELTEST: testing the model of DNA substitution.
Bioinformatics 14:817–818.

Presgraves, D. C. 2010. Darwin and the origin of interspecific genetic incompatibilities. Am. Nat.
176:S45–S60.

Pritchard, J. K., M. Stephens, and P. Donnelly. 2000. Inference of population structure using
multilocus genotype data. Genetics 155:945–959.

R Core Team. 2021. R: A Language and Environment for Statistical Computing. R Foundation for
Statistical Computing, Vienna, Austria.

Radford, I. J., R. D. Cousens, and P. W. Michael. 2004. Morphological and genetic variation in the
Senecio pinnatifolius complex: are variants worthy of taxonomic recognition? Aust. Syst.
Bot. 17:29–48.

Reid, N. M., D. A. Proestou, B. W. Clark, W. C. Warren, J. K. Colbourne, J. R. Shaw, S. I.
Karchner, M. E. Hahn, D. Nacci, M. F. Oleksiak, D. L. Crawford, and A. Whitehead. 2016.
The genomic landscape of rapid repeated evolutionary adaptation to toxic pollution in wild
fish. Science 354:1305–1308.

Richards, T. J., and D. Ortiz-Barrientos. 2016. Immigrant inviability produces a strong barrier to
gene flow between parapatric ecotypes of *Senecio lautus*. Evolution 70:1239–1248.

Richards, T. J., G. M. Walter, K. McGuigan, and D. Ortiz-Barrientos. 2016. Divergent natural
selection drives the evolution of reproductive isolation in an Australian wildflower.
Evolution 70:1993–2003.

Rieseberg, L. H. 2001. Chromosomal rearrangements and speciation. Trends in Ecology &
Evolution 16:351–358.

Rieseberg, L. H., and B. K. Blackman. 2010. Speciation genes in plants. Ann. Bot. 106:439–455.

Roda, F., L. Ambrose, G. M. Walter, H. L. Liu, A. Schaul, A. Lowe, P. B. Pelser, P. Prentis, L. H.
Rieseberg, and D. Ortiz-Barrientos. 2013a. Genomic evidence for the parallel evolution of
coastal forms in the *Senecio lautus* complex. Mol. Ecol. 22:2941–2952.

Roda, F., H. Liu, M. J. Wilkinson, G. M. Walter, M. E. James, D. M. Bernal, M. C. Melo, A. Lowe,
L. H. Rieseberg, P. Prentis, and D. Ortiz-Barrientos. 2013b. Convergence and divergence
during the adaptation to similar environments by an Australian groundsel. Evolution
67:2515–2529.

Rokas, A., and S. B. Carroll. 2008. Frequent and Widespread Parallel Evolution of Protein
Sequences. Molecular Biology and Evolution 25:1943–1953.

Rosenberg, N. A. 2004. DISTRUCT: A program for the graphical display of population structure.
Mol. Ecol. Notes 4:137–138.

Rundle, H. D., L. M. Nagel, J. W. Boughman, and D. Schluter. 2000. Natural selection and parallel
speciation in sympatric sticklebacks. Science 287:306–308.

Rundle, H. D., and P. Nosil. 2005. Ecological speciation. Ecol. Lett. 8:336–352.

Samuk, K., G. L. Owens, K. E. Delmore, S. E. Miller, D. J. Rennison, and D. Schluter. 2017. Gene
flow and selection interact to promote adaptive divergence in regions of low recombination.
Mol. Ecol. 26:4378–4390.

Schiffman, J. S., and P. L. Ralph. 2021. System drift and speciation. Evolution evo.14356.

Schluter, D. 2001. Ecology and the origin of species. Trends in Ecology & Evolution 16:372–380.

Schluter, D. 2009. Evidence for ecological speciation and its alternative. Science 323:737–741.

Schluter, D. 2000. The ecology of adaptive radiation. Oxford: Oxford University Press.

Schluter, D., and G. L. Conte. 2009. Genetics and ecological speciation. PNAS 106:9955–9962.

Schluter, D., and L. M. Nagel. 1995. Parallel speciation by natural selection. Am. Nat. 146:292–
301.

Schneemann, H., B. De Sanctis, D. Roze, N. Bierne, and J. J. Welch. 2020. The geometry and
genetics of hybridization. Evolution 74:2575–2590.

Seehausen, O., R. K. Butlin, I. Keller, C. E. Wagner, J. W. Boughman, P. A. Hohenlohe, C. L.
Peichel, G.-P. Saetre, C. Bank, Å. Brännström, A. Brelsford, C. S. Clarkson, F.
Eroukhmanoff, J. L. Feder, M. C. Fischer, A. D. Foote, P. Franchini, C. D. Jiggins, F. C.
Jones, A. K. Lindholm, K. Lucek, M. E. Maan, D. A. Marques, S. H. Martin, B. Matthews,
J. I. Meier, M. Möst, M. W. Nachman, E. Nonaka, D. J. Rennison, J. Schwarzer, E. T.
Watson, A. M. Westram, and A. Widmer. 2014. Genomics and the origin of species. Nat.
Rev. Genet. 15:176–192.

Sobel, J. M., G. F. Chen, L. R. Watt, and D. W. Schemske. 2010. The biology of speciation.
Evolution 64:295–315.

Soria-Carrasco, V., Z. Gompert, A. A. Comeault, T. E. Farkas, T. L. Parchman, J. S. Johnston, C.
A. Buerkle, J. L. Feder, J. Bast, T. Schwander, S. P. Egan, B. J. Crespi, and P. Nosil. 2014.
Stick insect genomes reveal natural selection's role in parallel speciation. Science 344:738–
742.

Stankowski, S., and M. Ravinet. 2021. Defining the speciation continuum. Evolution 75:1256–
1273.

Stearns, S. C. 1989. Trade-offs in life-history evolution. Functional Ecology 3:259–268.

Stuart, Y. E., T. Veen, J. N. Weber, D. Hanson, M. Ravinet, B. K. Lohman, C. J. Thompson, T.
Tasneem, A. Doggett, R. Izen, N. Ahmed, R. D. H. Barrett, A. P. Hendry, C. L. Peichel, and
D. I. Bolnick. 2017. Contrasting effects of environment and genetics generate a continuum
of parallel evolution. Nat. Ecol. Evol. 1:0158.

Therneau, T. M. 2024. coxme: Mixed Effects Cox Models. R package version 2.2-22.

Thompson, I. R. 2005. Taxonomic studies of Australian *Senecio* (Asteraceae): 5. The *S.*
pinnatifolius/S. lautus complex. Muelleria 21:23–76.

Thompson, K. A., Y. Brandvain, J. M. Coughlan, K. E. Delmore, H. Justen, C. R. Linnen, D. Ortiz-
Barrientos, C. A. Rushworth, H. Schneemann, M. Schumer, and R. Stelkens. 2024. The
ecology of hybrid incompatibilities. Cold Spring Harb Perspect Biol 16:a041440.

Tonnabel, J., P. David, T. Janicke, A. Lehner, J.-C. Mollet, J. R. Pannell, and M. Dufay. 2021. The
scope for postmating sexual selection in plants. Trends in Ecology & Evolution 36:556–567.

True, J. R., and E. S. Haag. 2001. Developmental system drift and flexibility in evolutionary
trajectories. Evolution & Development 3:109–119.

Unckless, R. L., and H. A. Orr. 2009. Dobzhansky–Muller incompatibilities and adaptation to a
shared environment. Heredity 102:214–217.

Van Belleghem, S. M., P. Rastas, A. Papanicolaou, S. H. Martin, C. F. Arias, M. A. Supple, J. J.
Hanly, J. Mallet, J. J. Lewis, H. M. Hines, M. Ruiz, C. Salazar, M. Linares, G. R. P.
Moreira, C. D. Jiggins, B. A. Counterman, W. O. McMillan, and R. Papa. 2017. Complex
modular architecture around a simple toolkit of wing pattern genes. Nat Ecol Evol 1:52.

Walter, G. M., J. D. Aguirre, M. W. Blows, and D. Ortiz-Barrientos. 2018a. Evolution of genetic
variance during adaptive radiation. Am. Nat. 191:E108–E128.

Walter, G. M., T. J. Richards, M. J. Wilkinson, M. W. Blows, J. D. Aguirre, and D. Ortiz-Barrientos. 2020. Loss of ecologically important genetic variation in late generation hybrids reveals links between adaptation and speciation. Evol. Lett. 4:302–316.

Walter, G. M., M. J. Wilkinson, J. D. Aguirre, M. W. Blows, and D. Ortiz-Barrientos. 2018b. Environmentally induced development costs underlie fitness tradeoffs. Ecology 99:1391–1401.

Walter, G. M., M. J. Wilkinson, M. E. James, T. J. Richards, J. D. Aguirre, and D. Ortiz-Barrientos. 2016. Diversification across a heterogeneous landscape. Evolution 70:1979–1992.

Wang, Z., Y. Zou, X. Li, Q. Zhang, L. Chen, H. Wu, D. Su, Y. Chen, J. Guo, D. Luo, Y. Long, Y. Zhong, and Y.-G. Liu. 2006. Cytoplasmic male sterility of rice with boro II cytoplasm is caused by a cytotoxic peptide and is restored by two related PPR motif genes via distinct modes of mRNA silencing. Plant Cell 18:676–687.

Weber, A. A.-T., J. Rajkov, K. Smailus, B. Egger, and W. Salzburger. 2021. Speciation dynamics and extent of parallel evolution along a lake-stream environmental contrast in African cichlid fishes. Science Advances 7:eabg5391.

Wilkinson, M. J., F. Roda, G. M. Walter, M. E. James, R. Nipper, J. Walsh, H. L. North, C. A. Beveridge, and D. Ortiz-Barrientos. 2021. Adaptive divergence in shoot gravitropism creates hybrid sterility in an Australian wildflower. PNAS 118:e2004901118.

Wu, C.-I., and C.-T. Ting. 2004. Genes and speciation. Nat Rev Genet 5:114–122.

Wund, M. A., J. A. Baker, B. Clancy, J. L. Golub, and S. A. Foster. 2008. A test of the "Flexible Stem" model of evolution: Ancestral plasticity, genetic accommodation, and morphological divergence in the threespine stickleback radiation. The American Naturalist 172:449–462.

Yamaguchi, R., and S. P. Otto. 2020. Insights from Fisher's geometric model on the likelihood of speciation under different histories of environmental change. Evolution 74:1603–1619.

Zhang, J. 2023. Patterns and evolutionary consequences of pleiotropy. Annual Review of Ecology, Evolution, and Systematics 54:1–19.

TABLES

Table 1. Theoretical framework for parallel mosaic speciation. Dobzhansky-Muller Incompatibilities (DMIs) accumulate between populations through asymmetric selection effects and genetic architecture. The framework extends classic models to incorporate environmental heterogeneity and polygenic adaptation, providing a quantitative foundation for understanding parallel mosaic speciation.

Component	Mathematical Form	Key Parameters
Classic Unckless-Orr Model	$P_{DM,U} = (2s_A s_B)/(s_A + s_B)^2$	s_A, s_B: selection coefficients
Environmental Effects Model	$P_{DM,E} = \dfrac{s_A s_B (1 + \varphi^2)}{(s_A + \varphi s_B)(\varphi s_A + s_B)}$	φ: environmental symmetry (0 to 1)
Polygenic Architecture Model	$P_{DM,P} = 1 - \dfrac{\varphi}{\sigma_1 \sigma_2} \sum_{i=1}^{n} s_i^2$	n: number of interacting loci

Table 2. Unification of speciation models through a parallel mosaic framework. Environmental symmetry (φ) and selection coefficients (s) interact to produce diverse patterns of reproductive isolation across heterogeneous landscapes. Classic parallel ecological and mutation-order speciation represent extreme cases of parallel mosaic speciation where environmental heterogeneity and genetic architecture jointly determine the probability of speciation. φ refers to how selection coefficients transfer between environments. For instance, high symmetry means an allele experiences similar selection pressure in both populations. Also see Table 1.

Model Component	Environmental Symmetry (φ)	Selection Coefficients	Population Structure	Reproductive Isolation	Example Systems
Parallel Ecological Speciation	$\varphi = 0$ between environments $\varphi = 1$ within environments	$s_A \gg s_B$ (highly asymmetric)	Discrete habitat transitions	Strong between environments; absent within environments	Threespine stickleback marine-freshwater pairs (Rundle et al. 2000)
Mutation-Order Speciation	$\varphi = 1$ (symmetric selection)	$s_A \approx s_B$ (similar magnitudes)	Independent populations in similar habitats	Present despite selection symmetry	Experimental *Drosophila* lineages (Hsu et al. 2024)
Parallel Mosaic Speciation	Continuous: $0 \leq \varphi \leq 1$	Variable relationship between s_A and s_B	Spatially structured populations across environmental mosaics	Varies with Φ and selection coefficient relationships	*Senecio lautus* complex

FIGURE CAPTIONS

Figure 1. Schematic diagram representing the differences between parallel ecological speciation and mutation-order speciation. Circles represent populations, whereas colors represent different environmental conditions. Ancestral populations are in grey. A) In ecological speciation, intrinsic and extrinsic reproductive isolation (RI) evolves during the adaptation of populations to different environments (i.e., during divergent natural selection selection). B) In contrast, during mutation-order speciation, intrinsic reproductive isolation evolves during the adaptation of populations to similar environments (i.e., during uniform natural selection).

Figure 2. Parallel evolution of *Senecio lautus* Dune and Headland populations. A) Illustrations of the Dune and Headland environments and ecotypes. B) Geographic distribution in Australia of the 16 populations in the study (blue Dune ecotype, orange Headland ecotype, pink Alpine ecotype, green Inland ecotype). C) Bayesian phylogeny constructed with 13 neutral markers implemented in *BEAST. Numbers on nodes are credible posterior probabilities. D) Bayesian assignment of individuals to genetic clusters within STRUCTURE for $K = 2$. Each individual is depicted as a bar, with colors representing ancestry proportions to each K cluster.

Figure 3. Strength of intrinsic reproductive isolation (RI) of coastal Dune (D) and Headland (H) *Senecio lautus* populations for F_1 seed set (left graphs) and F_1 viability. Data points represent the mean of multiple crosses for each population comparison (F_1 seed set: $n_{total} = 260$ crosses; F_1 viability: $n_{total} = 259$ crosses). Crosses were undertaken within and between the two clades defined in the phylogeny of Figure 1. Reproductive isolation was examined (A) between the same ecotype (D $\times$ D and H x H) or between different ecotypes (D $\times$ H and H $\times$ D), as well as (B) for each separate cross type (D $\times$ D, H $\times$ H, D $\times$ H and H $\times$ D). Positive values imply that hybrids perform worse than parents, and negative values that hybrids perform better than parents.

Figure 4. Strength of extrinsic reproductive isolation of coastal Dune and Headland *Senecio lautus* populations. Fitness of three replicate populations (LH – Lennox Head, SI – Stradbroke Island; CB – Cabarita Beach) from the Dune and Headland ecotypes transplanted into the dune and headland environments, respectively, at Lennox Head (Dune: $n_{total} = 479$ individuals; Headland: $n_{total} = 480$ crosses). Fitness was measured by A) survival curves, where the local population is denoted in color and dashed lines are 95% confidence intervals, and B) the probability of seedling establishment (production of 10 leaves), where asterisks denote significant differences from the local population.

Figure 5. Environmental and polygenic effects on Dobzhansky-Muller incompatibility (DMI) formation probability. A) Environmental effects model ($P_{DM,E}$), showing how the probability of incompatibility formation varies with environmental similarity φ under different selection ratios s_A/s_B. Lower φ (greater environmental difference) increases DMI probability, particularly when selection is more asymmetric (larger s_A/s_B). In the boundary case $\varphi = 1$ and $s_A = s_B$, this single‐locus model reduces to the Unckless–Orr result of $P_{DM,U} = 0.5$ (purple line). B) Polygenic architecture model ($P_{DM,P}$), illustrating how the probability of DMI increases with the number of contributing loci n. Genetic complexity

amplifies the impact of selection on reproductive isolation, especially under strongly asymmetric selection regimes.

Figure 6. The dynamics of parallel mosaic speciation in *Senecio lautus*. A) Patterns of intrinsic reproductive isolation in *Senecio*. Strong intrinsic reproductive isolation has evolved between ecotypes. In contrast, weaker intrinsic reproductive isolation has evolved between Headland (H) populations but not the Dune (D) populations, as shown by the thickness of the reproductive isolation lines. These patterns of intrinsic isolation are apparent for the most divergent crosses (i.e., crosses between clades). Reproductive isolation values are means for the crosses for F_1 seed set. B) Schematic diagram representing fitness landscapes of the Dune (blue) and Headland (orange) ecotypes. Dune populations share one optimum, driven by uniform selective pressures, whereas the heterogeneous headland environments likely lead to several optima for the Headland ecotype.

A)

Parallel ecological speciation

Geographic location 1

Geographic location 2

Time

Intrinsic and extrinsic RI

Divergent natural selection

Intrinsic and extrinsic RI

Divergent natural selection

Ancestral environment   Environment 1   Environment 2

B)

Parallel mutation-order speciation

Geographic location 1

Geographic location 2

Time

Intrinsic RI

Uniform natural selection

Intrinsic RI

Uniform natural selection

Ancestral environment   Environment 1

A)

Intrinsic reproductive isolation

Dune environment
Headland environment

Clade 1   Clade 2

D   H   D   H   D   H

Reproductive isolation

-0.10
-0.26
-0.48

Crosses within clades

-0.08
0.41
0.8

Crosses between clades

B)

Fitness landscapes

Dune Ecotype
One optimum
Unique path few adaptive alleles

Headland Ecotype
Several optima
Multiple paths many adaptive alleles

Both Ecotypes
Dune optimum
Headland optima