Evidence for mutation-order speciation between convergent ecotypes

Maria C. Melo[¶], Maddie E. James[¶], Federico Roda, Diana Bernal-Franco, Melanie J.

Wilkinson, Huanle Liu, Greg M. Walter, and Daniel Ortiz-Barrientos[*]

[¶]These authors contributed equally to this work.

The University of Queensland, School of Biological Sciences, St. Lucia, Queensland

4072, Australia.

[*]Corresponding author:

Daniel Ortiz-Barrientos

The University of Queensland

School of Biological Sciences

St. Lucia, Queensland 4072, Australia

E-mail: d.ortizbarrientos@uq.edu.au

Phone: 617-3365-1767

Fax: 617-3365-1655

Keywords: Parallel Speciation, Natural Selection, Reproductive Isolation, Gene Flow,

Senecio

In a number of animal species, divergent natural selection repeatedly and

independently drove the evolution of reproductive isolation between populations

adapted to contrasting, but not to similar environments[1]. This process is known as

parallel ecological speciation, and examples in plants are enigmatically rare[2]. Here,

we perform a comprehensive test of the ecological speciation hypothesis in an

Australian wildflower where parapatric populations found in coastal sand dunes

(Dune ecotype) and headlands (Headland ecotype) have repeatedly and

independently diverged in growth habit. Consistent with a role for divergent natural selection driving the evolution of reproductive isolation, we found that Dune populations with erect growth habit were easy to transplant across sand dunes, were largely interfertile despite half-a-million years of divergence, and were reproductively isolated from equally divergent Headland populations with prostrate growth habit. However, we unexpectedly discovered that both extrinsic and intrinsic reproductive isolation evolved between prostrate Headland populations, suggesting that populations evolving convergent phenotypes can also become new species rapidly. Mutation-order speciation[2], where the random accumulation of adaptive alleles create genetic incompatibilities between populations inhabiting similar habitats, provides a compelling explanation for these complex patterns of reproductive isolation, and is consistent with the increasingly common observation that there are multiple biochemical avenues to create convergent phenotypes. Our results suggest that natural selection can drive speciation effectively, but environmental and genetic complexity might make parallel ecological speciation uncommon in plants despite strong morphological convergence.

Convergent and divergent phenotypes can evolve when populations adapt repeatedly to contrasting environments. For instance, stickleback fishes adapted to freshwater lakes from the northern hemisphere lack armour and are similar in size when compared to their armoured and smaller relatives from the ocean[3]. This pattern of convergence and divergence is also seen in cave fishes[4], stick insects[5], and marine snails[6], eucalypts[7], and dandelions[8,9], and is thought to be driven by the repeated and independent action of

divergent natural selection between multiple population pairs occupying contrasting

environments.

Senecio lautus is a species complex from Australia and the pacific islands, generally

known as the common groundsel, that has repeatedly evolved two coastal ecotypes

adapted to contrasting environments[10]. The Dune (D) ecotype occupies the sheltered

areas behind the sand dunes of the beach, whereas the Headland (H) ecotype lives in

exposed and salty rocky outcrops and cliff tops that interrupt the beach. These ecotypes

have contrasting growth habits, where Dunes grow erect and Headlands grow prostrate

and form mats on top of the ground[10,11] (Figure 1A). Multiple transplant experiments

have shown that Dune and Headland populations are exposed to strong divergent

natural selection[12-14], which creates strong ecologically-dependent reproductive

isolation between them[12,14,15]. However, it remains unknown whether intrinsic

reproductive isolation has evolved as a by-product of adaptation to contrasting or

similar environments. Here we test the fundamental predictions of the parallel

speciation hypothesis for the evolution of both extrinsic and intrinsic reproductive

isolation in *S. lautus* by using a range of glasshouse, field, and laboratory experiments.

We first confirmed if divergent natural selection drove the independent and repeated

evolution of these two coastal ecotypes (Figure 1A). As we observed before[10], Dune and

Headland populations that live next to each other (parapatric DH pairs) are

phylogenetically more related to each other than to any other Dune or Headland

population, respectively (Figure 1B, Table S1). Using 13 neutral nuclear loci common

to sixteen populations of *S. lautus* we show that DH pairs are sister taxa. DH pairs

sorted into two main clades (South-Eastern, and Southern), which we also detected in a

population genetics analysis of the same loci using the STRUCTURE algorithm (Supp.

Figure 1). Adjacent populations showed little genetic differentiation, possibly

suggesting that high levels of ancestral polymorphism are still present in these

population pairs, or that gene flow after secondary contact could have homogenised

them (Supp. Figure 1 and 2). To disentangle these scenarios we used coalescent models

of isolation with migration[16] to estimate migration rates in each coastal pair. If gene

flow after secondary contact were responsible for the phylogenetic pattern predicted by

parallel speciation, we expect to detect high levels of gene flow in DH pairs.

Alternatively, if DH pairs originated multiple times independently, we expect low to nil

levels of gene flow in each pair, indicating that genetic similarities between populations

reflect segregation of ancestral polymorphism. We failed to detect migration in each DH

pair (Table S1), except in pair (D03H02), which showed asymmetric gene flow

($Nm_{H02>D03}$ = 23.01, 95% credible interval 12.30-34.98, Table S2). The phylogenetic

pattern detected in coastal *Senecio* and the lack of gene flow between parapatric pairs

strongly suggests that Dune and Headland morphologies have evolved independently

multiple times.

Next, we asked if the independent evolution of Dune and Headland populations lead to

the evolution of reproductive isolation between individuals occupying contrasting

environments (DxH crosses or transplants), and between individuals found in similar

environments (DxD and HxH crosses or transplants). We performed crosses in the

glasshouse both within and between the Southern (S) and South-Eastern clades (SE),

thus correcting for divergence time, and reanalysed previous transplant experiments in

the field across multiple Dune and Headland habitats[13]. In each glasshouse cross, we

measured the ability to produce hybrid seed as well as ability to germinate relative to

the average of the parents. In transplant experiments in the field, we previously

measured the viability component of fitness by tracking individual mortality since

germination. Consistent with parallel speciation driven by divergent natural selection,

we found that populations that inhabit similar environments were on average

reproductively and ecologically compatible, while populations inhabiting different

environments were reproductively and ecologically isolated. We asked if reproductive

and ecological compatibility was similar between Dune populations or between

Headland populations. Surprisingly, we found that only Dune populations were easily

crossed or transplanted into each other's habitat, whereas Headland populations were

difficult to cross and transplant between headland sites (Figure 2, 3 Supplementary

Table S3 and S4). Altogether, it appears that costal parapatric pairs of *S. lautus* have

evolved by divergent natural selection, but Headland populations have done it uniquely

every time, thus creating further reproductive isolation among them.

Contrary to expectations from the parallel speciation hypothesis, our results indicate

that the evolution of the same morphology does not guarantee reproductive

compatibility between populations. A straightforward explanation is adaptive polygenic

divergence between populations inhabiting similar environments. Here, populations

evolving similar values of a complex trait accumulate adaptive differences in random

order[1,2] some of which incidentally cause intrinsic reproductive isolation. This

speciation mode is referred to as mutation-order speciation, and it has been used as an

alternative view to speciation driven by divergent natural selection. Our findings

suggest that the two processes might have more in common than previously anticipated.

Previous results in *S. lautus* have suggested that DH pairs are divergent in similar

signalling pathways even if the differentiated genes are not common across populations

pairs[17]. One of these pathways controls auxin movement, which is known to affect

variation in growth habit in Arabidopsis and other model plant systems. Re-use of a

genetic pathway might be an easy solution for creating convergent phenotypes, yet it

might incidentally lead to the origins of new species. This might explain why parallel

ecological speciation might be rare in some organisms, such as plants.

Methods

Study populations and sample preparation

Leaf samples were collected from six parapatric *Senecio lautus* Dune and Headland

populations (*S. pinnatifiolius* var. *pinnatifolius*, *S. pinnatifolius* var. *maritimus*) as well

as two Inland and two Alpine populations (*S. pinnatifolius* var *serratus* and *S.*

pinnatifolius var. *alpinus*) (Table S1 and map in Figure 1, and supplementary

materials[10] for more details). *Senecio madagascariensis*, a closely related species to *S.*

lautus from Africa was included as an outgroup[10]. We extracted DNA with a modified

CTAB protocol and samples were standardised to 30ng/μL[10].

Library construction and sequencing

We undertook targeted re-sequencing in the Fluidigm Access Array system[18], which

enables the simultaneous amplification of 48 primers across 48 individuals, whilst

barcoding individuals for next generation sequencing. Individuals were barcoded using

the 4-primer PCR process with 454 barcodes[18]. Individuals were randomised and pooled

in equimolar quantities into two pools, and sent for sequencing at the Beijing Genomics

Institute. Pools were cleaned with an Agencourt AMPure XP purification kit, and then

sequenced using the emPCR (Lib-A) kit for bi-directional sequencing on two lanes of

the Roche GS FLX Titanium platform.

SNP calling

After trimming barcodes with TagCleaner[19], reads that had more than 40% low quality

bases (<Q20), had Ns higher than 2%, or were shorter than 50bp were removed.

PRGmatic [20] was used for read alignment. The PRGmatic pipeline aligns reads into loci

and constructs haplotypes for each individual using dependent programs including

CAP3[21], BWA[22], SAMtools[23], and VarScan[24]. We used default parameters and an

overlap of 100bp for read alignment, as suggested for 454 data[20]. A locus was only

considered for further analyses if haplotypes were assigned in six or more individuals

per population. BLAT[25] was used to map loci from PRGmatic to the expected

amplicons. 26 loci were selected and aligned in MUSCLE[26] and subsequently used in

phylogenetic ananylses and neutrality tests. See Tables S5-S7 for genes and primer

sequences.

DNA sequence polymorphism analyses

Deviations from neutrality[27] for each gene within each population were tested using

HKA[28], Tajima's D[29] and Fu and Li's tests[30] using DNAsp. Loci selected for

phylogenetic and population genetics analyses did not reject neutrality via the HKA test,

or via the other two tests if HKA was impossible to perform because a locus did not

amplify in the outgroup species. As a consequence, 26 loci were used in estimating gene

flow between different pairwise comparisons between populations, and 13 neutral loci

across populations were used for the phylogeny (Table S5-S7).

Phylogenetic analysis and population differentiation for neutral markers

A Bayesian phylogenetic analysis was performed using *Beast for species tree

estimation[31]. Analysis was performed with a chain 300,000,000 long using a strict

molecular clock. ITS was used as the reference locus, with a mutation rate for

herbaceous plants of $4.13*10^{-9}$ subs/site/year [32], from which the rate of the other genes

were estimated. According to Bayesian Information Criterion values found using

jModeltest[33], the best substitution model for 17 out of the 26 genes was the HKY

model[34]. We used a *Yules* species process for species tree estimation (this assumes that

lineages split at a constant rate).

We explored population structure across the species complex by inferring the number of

genetic clusters (K) using STRUCTURE v2.3.4[35]. This was undertaken at two levels: *i)*

including all populations, and *ii)* by pairs (six parapatric and two allopatric).

STRUCTURE was run using an admixture model and the correlated allele frequency

model[36]. Parameters used for *i)* K=1-16, 20 iterations per K, burnin 100,000, MCMC

100,000; and for *ii)* K=1-9, 20 iterations per K, burnin 100,000, MCMC 100,000, as

suggested by *Gilbert et al.* (2012) . To choose the most likely K value, both methods by

Pritchard *et al.* (2000) and Evanno *et al.* (2005) were examined. Because both methods

tend to overestimate K, and high K values did not add any additional clustering

information compared to smaller Ks, the smallest K that captured the major structure in

the data was chosen, ensuring that all summary statistics converged.

Gene flow

Gene flow estimates were derived from genes that did not reject neutrality tests in most

populations and were non-recombining. Intra-locus recombination was assessed with

the four-gamete test[39] in DNAsp[40]. The longest non-recombining DNA segment per

locus was included in the analysis, as sequences including recombination events could

lead to a false signature of gene flow[41]. Estimates of gene flow were assessed using

IMa2[16] with pairwise comparisons between the six parapatric pairs and two allopatric

comparisons. Gene flow between proximate populations inhabiting the same habitat was

also estimated (between Dunes and between Headlands). We used a 100,000 burnin for

a minimum of 100,000 genealogies. Runs involved 150 to 180 independent chains,

effective sample size (ESS) among all parameters ranged from several hundred to

>10,000. Upper bounds of the prior distributions for each parameter were set based on

the results of a preliminary run. Mutation rates for each locus were obtained using the

relationship D=2uT, where D is the average number of substitutions separating the

Senecio lautus complex from *S. madagascariensis,* and T is the age of the *S. lautus*

clade[17] (Roda *et al.* 2013a). Gene flow estimates did not change if we ignored mutation

rates (data not shown). We ran three independent IMa2 analyses varying the seed

number to ensure parameter estimation was consistent.

Estimates of reproductive isolation

Seeds from 30 individuals from four population pairs, two from the eastern coast (D01-

H01 and D04-H05) and two from the southern coast (D32-H12 and D23-H21), were

collected and stored in dry conditions at 4°C in the Ortiz-Barrientos Laboratory at The

University of Queensland, Australia. Seeds were scarified (1mm trimmed at the

micropyle side) and germinated on moist filter paper in petri dishes. Seeds were kept in

dark and controlled conditions for three days to induce root elongation, and

subsequently placed under the light for 7 days to induce vegetative growth. Seedlings

were then transplanted into 0.25L pots filled with standard potting mix and transferred

to a glasshouse with constant temperature (25°C) and 12h:12h light:dark cycle. After

two months, flowering individuals were crossed by rubbing flower heads over a period

of three to five days to create seed stocks for each population.

F1 hybrid seed set

15 families for each population were germinated and grown under glasshouse

conditions (described above). Intra- and inter-population crosses were performed twice

a day by gently rubbing flower heads (capitula): each flower head was crossed at least

three times to saturate the number of fertilised florets. Due to logistic constraints,

crosses were blind with respect to research assistants but not with respect to MCM who

supervised the experiments. We calculated seed set by estimating the proportion of

fertilised (filled seeds)[14] seeds in flower heads. We divided the number of fertilised

seeds in an inter-population cross by the average number of seeds produced in parental

intra-population crosses; we subtracted this fraction from one to calculate postmating

prezygotic reproductive isolation in the system[17].

F1 viability at early stages of the life cycle

We tested for the viability of F1 hybrids between multiple inter-population crosses (see

above) by counting the proportion of seeds that germinated in relation to the parental

germination proportions. A total of 260 families equally distributed across all parental

and hybrid cross types were germinated, placing five seeds of each family into moist

filter paper in petri dishes (one family per petri dish). We placed petri dishes into trays

(20 per tray) while randomising the position of all families. Germination conditions are

the same as described above, except for tray position on the shelves, which in this case

was switched daily. We did not scarify seeds to investigate the intrinsic ability of

embryos to germinate. Data collection was blind, where cells in a tray were assigned a

number that was cross-referenced with the original genotype and family ID at the end of

the experiment. We calculated F1 inviability by estimating the proportion of germinated

seeds in a cross, divided by the average number of seeds that germinated of the two

parental populations.

Strength of reproductive isolation

Estimates of reproductive isolation were done for two reproductive isolating barriers in

the following way: *Hybrid seed set (I)* in the glasshouse or whether the proportion of

fertilised seeds in a flower head from an inter-population cross differed from an intra-

population cross as $RI_{seed\ set}$=1-(Pf$_{inter}$/Pf$_{intra}$) were Pf stands for proportion fertilised.

Hybrid viability (I), or whether hybrid seedlings germinated equally well as their

parents in the glasshouse, as H_{hf}=1-(v$_{F1}$/v$_{parents}$), where v$_{F1}$ is the average germination of

F1 hybrids, and v$_{parents}$ is the average germination of the two parents.

Estimates of local and non-local field performance

To identify whether local adaptation was stronger among populations for the Headland

versus the Dune ecotype, we reanalysed field transplant experiments that used seeds

from three populations of each ecotype (n=150-180 seeds/population/transplant

environment). Methods of the transplants into the two natural environments at Lennox

Head (NSW), including the local population are described in detail in Walter et al.

2016, but briefly, we fixed each seed to a toothpick using non-drip supa glue. The

toothpicks were then placed in a fully randomised grid under natural conditions in both

the sand dunes and rocky headland at Lennox Head. We used six blocks to sample each

environment. Shadecloth (50%) was suspended 15cm above the grids and seeds watered

for three weeks to replicate ideal germination conditions. We recorded seedling

emergence as a binary trait, we then tracked seedling survival and recorded whether

plants reached 10 leaves (as a measure of seedling establishment).

To understand whether local populations performed better than non-local populations

(from the same habitat), we first implemented a survival analysis within the R package

'coxme'[43] where survival was censored at day 320, and we included population as a

fixed effect, and environmental block as a random effect. Next, to test whether local

populations had a higher probability of reaching seedling establishment, we

implemented a generalised linear mixed effects model within the R package 'lme4'[44].

To do so, we included seedling establishment as a binary response variable, with

population as a fixed effect, and environmental block as a random effect. In both

analyses we wanted to test whether the two non-local populations performed more

poorly than the local population. Thus, for each analysis we designated the local

population as the intercept, testing each non-local population relative to the intercept.

References
270
271   1   Schluter, D. Evidence for ecological speciation and its alternative. *Science* **323**,
272   737 (2009).
273   2   Ostevik, K. L., Moyers, B. T., Owens, G. L. & Rieseberg, L. H. Parallel
274   ecological speciation in plants? *International Journal of Ecology* **2012** (2012).
275   3   Boughman, J. W., Rundle, H. D. & Schluter, D. Parallel evolution of sexual
276   isolation in sticklebacks. *Evolution* **59**, 361-373, doi:10.1111/j.0014-
277   3820.2005.tb00995.x (2005).
278   4   Strecker, U., Hausdorf, B. & Wilkens, H. Parallel speciation in *Astyanax* cave
279   fish (Teleostei) in Northern Mexico. *Molecular Phylogenetics and Evolution* **62**,
280   62-70 (2012).
281   5   Nosil, P., Crespi, B. J. & Sandoval, C. P. Host-plant adaptation drives the
282   parallel evolution of reproductive isolation. *Nature* **417**, 440-443 (2002).
283   6   Johannesson, K. *et al.* Repeated evolution of reproductive isolation in a marine
284   snail: unveiling mechanisms of speciation. *Philosophical Transactions of the*

Royal Society B: Biological Sciences **365**, 1735-1747,
doi:10.1098/rstb.2009.0256 (2010).
7   Foster, S. A., McKinnon, G. E., Steane, D. A., Potts, B. M. & Vaillancourt, R.
E. Parallel evolution of dwarf ecotypes in the forest tree *Eucalyptus globulus*.
New Phytologist **175**, 370-380 (2007).
8   Vijverberg, K., Kuperus, P., Breeuwer, J. & Bachmann, K. Incipient adaptive
radiation of New Zealand and Australian Microseris (Asteraceae): an amplified
fragment length polymorphism (AFLP) study. *Journal of Evolutionary Biology*
13, 997-1008 (2000).
9   Vijverberg, K., Mes, T. H. & Bachmann, K. Chloroplast DNA evidence for the
evolution of *Microseris* (Asteraceae) in Australia and New Zealand after long-
distance dispersal from western North America. *American Journal of Botany* **86**,
1448-1463 (1999).
10   Roda, F. *et al.* Genomic evidence for the parallel evolution of coastal forms in
the *Senecio lautus* complex. *Mol Ecol* **22**, 2941-2952, doi:10.1111/mec.12311
(2013).
11   Thompson, I. Taxonomic studies of Australian *Senecio* (Asteraceae): 5. The *S.*
pinnatifolius/S. lautus complex. *Muelleria* **21**, 23-76 (2005).
12   Richards, T. J., Walter, G. M., McGuigan, K. & Ortiz-Barrientos, D. Divergent
natural selection drives the evolution of reproductive isolation in an Australian
wildflower. *Evolution; international journal of organic evolution* **70**, 1993-
2003, doi:10.1111/evo.12994 (2016).
13   Walter, G. M., Wilkinson, M. J., Aguirre, J. D., Blows, M. W. & Ortiz-
Barrientos, D. Environmentally induced development costs underlie fitness
tradeoffs. *Ecology* **99**, 1391-1401, doi:doi:10.1002/ecy.2234 (2018).
14   Melo, M. C., Grealy, A., Brittain, B., Walter, G. M. & Ortiz-Barrientos, D.
Strong extrinsic reproductive isolation between parapatric populations of an
Australian groundsell. *New Phytologist* **doi: 10.1111/nph.12779** (2014).
15   Richards, T. J. & Ortiz-Barrientos, D. Immigrant inviability produces a strong
barrier to gene flow between parapatric ecotypes of Senecio lautus. *Evolution;*
international journal of organic evolution **70**, 1239-1248,
doi:10.1111/evo.12936 (2016).
16   Hey, J. Isolation with migration models for more than two populations.
Molecular Biology and Evolution **27**, 921-933 (2010).
17   Roda, F. *et al.* Convergence and divergence during the adaptation to similar
environments by an Australian groundsel. *Evolution* **67**, 2515-2529,
doi:10.1111/evo.12136 (2013).
18   Moonsamy, P. V. *et al.* 201-P Use of the Fluidigm® access Array™ system
provides simplified amplicon library preparation in next generation sequencing
for high throughput HLA genotyping. *Human Immunology* **72**, S142 (2011).
19   Schmieder, R., Lim, Y., Rohwer, F. & Edwards, R. TagCleaner: Identification
and removal of tag sequences from genomic and metagenomic datasets. *BMC*
Bioinformatics **11**, 341 (2010).
20   Hird, S. M., Brumfield, R. T. & Carstens, B. C. PRGmatic: an efficient pipeline
for collating genome‐enriched second‐generation sequencing data using a '
provisional‐reference genome'. *Molecular Ecology Resources* **11**, 743-748
(2011).

21   Huang, X. & Madan, A. CAP3: A DNA sequence assembly program. *Genome*
Research **9**, 868-877 (1999).
22   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–
Wheeler transform. *Bioinformatics* **25**, 1754-1760,
doi:10.1093/bioinformatics/btp324 (2009).
23   Li, H. *et al.* The sequence alignment/map format and SAMtools. *Bioinformatics*
25, 2078-2079, doi:10.1093/bioinformatics/btp352 (2009).
24   Koboldt, D. C. *et al.* VarScan: variant detection in massively parallel sequencing
of individual and pooled samples. *Bioinformatics* **25**, 2283-2285,
doi:10.1093/bioinformatics/btp373 (2009).
25   Kent, W. J. BLAT—the BLAST-like alignment tool. *Genome Research* **12**, 656-
664 (2002).
26   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and
high throughput. *Nucleic Acids Research* **32**, 1792-1797 (2004).
27   Kimura, M. *The neutral theory of molecular evolution.* (Cambridge University
Press, 1985).
28   Hudson, R. R., Kreitman, M. & Aguadé, M. A test of neutral molecular
evolution based on nucleotide data. *Genetics* **116**, 153-159 (1987).
29   Tajima, F. Statistical method for testing the neutral mutation hypothesis by
DNA polymorphism. *Genetics* **123**, 585-595 (1989).
30   Fu, Y.-X. & Li, W.-H. Statistical tests of neutrality of mutations. *Genetics* **133**,
693-709 (1993).
31   Heled, J. & Drummond, A. J. Bayesian inference of species trees from
multilocus data. *Molecular biology and evolution* **27**, 570-580 (2010).
32   Kay, K., Whittall, J. & Hodges, S. A survey of nuclear ribosomal internal
transcribed spacer substitution rates across angiosperms: an approximate
molecular clock with life history effects. *BMC Evolutionary Biology* **6**, 36
(2006).
33   Posada, D. & Crandall, K. A. MODELTEST: testing the model of DNA
substitution. *Bioinformatics* **14**, 817-818, doi:10.1093/bioinformatics/14.9.817
(1998).
34   Hasegawa, M., Kishino, H. & Yano, T.-a. Dating of the human-ape splitting by
a molecular clock of mitochondrial DNA. *Journal of molecular evolution* **22**,
160-174 (1985).
35   Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure
using multilocus genotype data. *Genetics* **155**, 945-959 (2000).
36   Falush, D., Stephens, M. & Pritchard, J. K. Inference of population structure
using multilocus genotype data: linked loci and correlated allele frequencies.
Genetics **164**, 1567-1587 (2003).
37   Gilbert, K. J. *et al.* Recommendations for utilizing and reporting population
genetic analyses: the reproducibility of genetic clustering using the program
structure. *Molecular Ecology* **21**, 4925-4930, doi:10.1111/j.1365-
294X.2012.05754.x (2012).
38   Evanno, G., Regnaut, S. & Goudet, J. Detecting the number of clusters of
individuals using the software structure: a simulation study. *Molecular Ecology*
14, 2611-2620, doi:10.1111/j.1365-294X.2005.02553.x (2005).

39   Hudson, R. R. & Kaplan, N. L. Statistical properties of the number of
379   recombination events in the history of a sample of DNA sequences. *Genetics*
380   **111**, 147-164 (1985).
381   40   Librado, P. & Rozas, J. DnaSP v5: A software for comprehensive analysis of
382   DNA polymorphism data. *Bioinformatics* **25** (2009).
383   41   Hey, J. & Nielsen, R. Multilocus methods for estimating population sizes,
384   migration rates and divergence time, with applications to the divergence of
385   *Drosophila pseudoobscura* and *D. persimilis*. *Genetics* **167**, 747-760,
386   doi:10.1534/genetics.103.024182 (2004).
387   42   Ramsey, J., Bradshaw Jr, H. & Schemske, D. Components of reproductive
388   isolation between the monkeyflowers *Mimulus lewisii* and *M. cardinalis*
389   (Phrymaceae). *Evolution* **57**, 1520-1534 (2003).
390   43   Huebner, M. & Therneau, T. Mixed Effects Cox Models For Gene Set Analysis
391   In Lung Cancer. *Genet Epidemiol* **36**, 157-157 (2012).
392   44   Bates, D., Machler, M., Bolker, B. M. & Walker, S. C. Fitting Linear Mixed-
393   Effects Models Using lme4. *J Stat Softw* **67**, 1-48 (2015).
394

Figure Legends

Figure 1. A) Morphological differences between the two ecotypes: Dunes grow erect and have big leaves, while Headlands grow prostrate and have smaller succulent leaves. Modified from[10], with permission from the authors. **B)** Geographic distribution of 16 populations in the study and phylogeny based on 13 neutral markers using Bayesian inference of the populations in the study. S and SE respectively refer to the Southern and South-Eastern clades observed in the phylogeny. Numbers above nodes are estimates of divergence time obtained in IMa2, whereas numbers below nodes are credible posterior probabilities obtained in BEAST*.

Figure 2. F1 hybrid seed set and F1 hybrid inviability in crosses between coastal populations in *Senecio lautus*. Each panel represents a combination of crosses within ecotype (DxD, with Dune mother; HxH, with Headland mother) or between ecotypes (DxH, with Dune mother; HxD, with Headland mother) from the same or from a different clade. Positive values of Strength of RI imply that hybrids perform worse than parents, and negative values that hybrids perform better than parents.

Figure 3. Field performance of three replicate populations from each ecotype in their native habitat at Lennox Head (NSW) in the form of Estimated S (survival). Coloured populations represent the local population and error bars represent 95% confidence intervals. (**A**) Survival was similar for the three Dune populations, with only a significant difference between the local population (Lennox Head) and the Cabarita Beach population. The local Headland population survived much better than the two non-local populations. Tables inset describe tests of each non-local population relative

to the local population. (**B**) Probability of reaching seedling establishment was similar for all Dune populations, but was much higher for the local Headland population (asterisks denote significant difference to local population).

Figure 4. Observed pattern of extrinsic and intrinsic reproductive isolation in *Senecio lautus*. Rectangles represent sets of reproductively or ecologically compatible populations. For instance, all Dune (orange) and Headland (green) populations are reproductively compatible within clades (top right panel). However, only Dune populations are ecologically compatible within clades as they cannot be transplanted easily into rocky headland sites, and Headland populations perform best in their own rocky site (top left panel). This pattern is also found when examining intrinsic reproductive isolation between clades and ecotypes (bottom right panel).

Figure 1.

(A)

	Stradbroke Island	Cabarita Beach	Lennox Head	Coffs Harbour	South East	Cape Bridgewater
H ●						
D ●						
	D00-H00	D03-H02	D01-H01	D04-H05	D12-H14	D15-H12

(B)

Figure 2

Figure 3

Figure 4