Stable and widespread structural heteroplasmy in chloroplast genomes revealed by a new long-read quantification method

Weiwen Wang [a], Robert Lanfear [a]

[a] Research School of Biology, the Australian National University, Canberra, ACT, Australia, 2601

Corresponding Author:

Weiwen Wang, wei.wang@anu.edu.au

Robert Lanfear, rob.lanfear@anu.edu.au, +61 2 6125 2536

Keywords: Single copy inversion, flip-flop recombination, chloroplast genome structural heteroplasmy

Abstract

The chloroplast genome usually has a quadripartite structure consisting of a large single copy region and a small single copy region separated by two long inverted repeats. It has been known for some time that a single cell may contain at least two structural haplotypes of this structure, which differ in the relative orientation of the single copy regions. However, the methods required to detect and measure the abundance of the structural haplotypes are labour-intensive, and this phenomenon remains understudied. Here we develop a new method, Cp-hap, to detect all possible structural haplotypes of chloroplast genomes of quadripartite structure using long-read sequencing data. We use this method to conduct a systematic analysis and quantification of chloroplast structural haplotypes in 61 land plant species across 19 orders of Angiosperms, Gymnosperms and Pteridophytes. Our results show that there are two chloroplast structural haplotypes which occur with equal frequency in most land plant individuals. Nevertheless, species whose chloroplast genomes lack inverted repeats or have short inverted repeats have just a single structural haplotype. We also show that the relative abundance of the two structural haplotypes remains constant across multiple samples from a single individual plant, suggesting that the process which maintains equal frequency of the two haplotypes operates rapidly, consistent with the hypothesis that flip-flop recombination mediates chloroplast structural heteroplasmy. Our results suggest that previous claims of differences in chloroplast genome structure between species may need to be revisited.

Significance Statement

Chloroplast genome consists of a large single copy region, a small single copy region, and two inverted repeats. Some decades ago, a discovery showed that there are two types chloroplast genome in some plants, which differ the way that the four regions are put together. However, this phenomenon has been shown in just a small number of species, and many open questions remain. Here, we develop a fast method to measure the chloroplast genome structures, based on long-reads. We show that almost all plants we analysed contain two possible genome structures, while a few plants contain only one structure. Our findings hint at the causes of the phenomenon, and provide a convenient new method with which to make rapid progress.

Introduction

Chloroplasts are organelles which are vital for photosynthesis. Most land plant chloroplast genomes are 120 – 160 kb in size (1, 2), and have a quadripartite structure consisting of a pair of identical rRNA-containing inverted repeats (hereafter referred to as IR) of ~10-30kb divided by a large single copy (LSC) region of ~80 -90 kb and a small single copy (SSC) region of ~10-20 kb.

Surprisingly, chloroplast genomes can exist two structural haplotypes differing in the orientation of single copy regions (3). The presence of this structural heteroplasmy has been confirmed in some land plants (3-6) and algae (7, 8), but its cause remains

unknown. One hypothesis to explain the presence of structural heteroplasmy is known

as flip-flop recombination (4). This hypothesis suggests that the large IRs could

mediate frequent intramolecular recombination, resulting in the maintenance of

roughly equal amounts of the two haplotypes differing only in the orientation of their

single copy regions.

A better understanding of chloroplast genome structural heteroplasmy is important for

a number of reasons. First, some recent papers have suggested that the orientation of

the single copy regions differ between species (9-12), but these studies seem to have

overlooked the possibility that both orientations may coexist in a single individual.

Second, the relationship between the structure of the chloroplast genome and the

existence or otherwise of structural heteroplasmy remains poorly understood. For

example, if flip-flop recombination causes chloroplast structural heteroplasmy, then it

seems likely that the presence of two long IRs may be a pre-requisite for the existence

of heteroplasmy. Consistent with this, no heteroplasmy was observed in a chloroplast

genome with highly reduced IRs (13), but the generality of this observation remains to

be tested. Third, if heteroplasmy is the norm rather than the exception, then this may

represent a challenge for the assembly of chloroplast genomes, which may be easily

overcome by simply allowing for the existence of two structural haplotypes during the

assembly process. A large-scale analysis across many different species has the

potential to provide a more complete picture of chloroplast heteroplasmy, further

elucidating this fascinating phenomenon and potentially improving genome assembly

and inference. In this study, we perform this large-scale analysis using a new method

which we developed to quickly and conveniently quantify structural heteroplasmy in

chloroplast genomes from long-read sequencing data.

Currently there are two methods to detect different structural heteroplasmy in

chloroplast genomes: Bacterial Artificial Chromosomes (BAC)-End-Sequence (BES)

(5), and restriction digests (4). For the BES method, one first constructs BAC libraries

in which large pieces of chloroplast DNA are inserted into bacterial chromosomes.

One then uses the BAC libraries to sequence short fragments of both ends of many

such pieces of chloroplast DNA to ascertain which chloroplast structures exist in a

given sample. BES reads can only provide information on chloroplast genome

structure when a single read covers an entire IR region, with one end in LSC region

and another end in the SSC region. Although a very useful method, BES reads may be

limited for detecting highly atypical chloroplast genome structures, because only a

short fragment from each end of a read is actually sequenced. For the restriction

digestion method, one uses restriction enzymes to digest the chloroplast genome, and

then decodes the chloroplast genome structure by studying the distribution of the

resulting fragment lengths using agarose gel electrophoresis. Using this approach,

Stein *et al* (4) found that the IR region was present in four different fragments which

could be separated into two groups for which the sum of lengths was equal. They

therefore concluded that chloroplast genome contained two equimolar isomers, which

differed in their single copy orientation. The restriction digest method has provided

much of the existing information on chloroplast genome heteroplasmy, but it is

limited by the availability of suitable restriction sites (which may be unknown in

some species), and requires a labour-intensive hybridisation step to infer chloroplast

genome heteroplasmy. Other methods have also been proposed, such as the use of

PCR to attempt to amplify diagnostic regions of the chloroplast genome. However, it

is not easy to generate PCR fragments which are longer than the IR region (usually

10-30 kb), so while PCR-based methods could only work well with short IRs (e.g. of

a few hundred bp) (13), their general application is likely to remain limited.

Furthermore, PCR-mediated recombination could result in false positives when using

PCR-based methods (14, 15). In short, current methods for detecting structural

heteroplasmy in chloroplast genomes are labour- and time-intensive, which makes

them difficult to employ for broad-scale studies. Current methods have been applied

in a relatively small number of land plant species (*Beta vulgaris* (6), *Musa acuminata*

(5), *Phaseolus vulgaris* (3), *Osmunda cIaytoniana*, *Osmunda cinnamomea* and

Osmunda regalis (4), and some species in the Pinaceae (13)), and as a result our

understanding of the patterns and causes of structural heteroplasmy in chloroplast

genomes remains somewhat limited.

In this study, we develop and apply a fast, simple, and cheap method to detect and

quantify structural heteroplasmy in chloroplast genomes using long-read sequencing.

This method requires just a single DNA extraction and sequencing run, followed by a

simple bioinformatic analysis for which we provide free and open-source code

(https://github.com/asdcid/Cp-hap). Our approach, which we call Cp-hap, relies on

sequencing individual DNA molecules longer than the length of the IR region

(roughly 10-30 kb in most species). Sequencing reads of this length are commonly

obtained in great abundance when sequencing whole-plant DNA extractions on the

Oxford Nanopore MinION device (16, 17). Our method sifts through these long

sequencing reads to find those that uniquely match one of the 32 uniquely-identifiable

structural haplotypes (Fig. S1), and then uses the abundance of these reads to estimate

the relative abundance of each haplotype (details see Materials and Methods). Since

many new plant samples are being sequenced with long-read sequencing technology,

this method has the potential to provide many insights in the coming years, simply as

a by-product of existing sequencing efforts.

We use Cp-hap to quantify the abundance of chloroplast genome structural haplotypes

in 61 plant species. Our results suggest that most land plants contain the two

commonly-found chloroplast genome haplotypes in equal abundance whenever the

chloroplast genomes contain a pair of large IRs. However, when the IR is small (e.g.

only a few hundred bp in size) or repeat regions are not inverted but positioned, we

find only a single haplotype. Finally, we show that multiple samples taken from a

single individual all contain the same two haplotypes in equal abundance, suggesting

that the process which maintains equimolar concentrations of the two haplotypes

operates relatively rapidly, and consistent with the suggestion that flip-flop

recombination is the underlying cause.

Results

Data collection

We searched the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) database, and selected 90 land plants species which met our length and data amount requirement (see Materials and Methods) to analysis (Table S1). Two of these species (*Herrania umbratica* and *Siraitia grosvenorii*) lacked suitable chloroplast reference genomes, so we therefore assembled these two chloroplast genomes *de-novo* (see Materials and Methods). The chloroplast genomes of *Herrania umbratica* and *Siraitia grosvenorii* were 158,475 bp and 158,757 bp in size, consisting of a LSC (88,369 bp and 87,625 bp), a SSC (19,040 bp and 18,556 bp), and two IRs (25,533 bp and 26,288 bp), and encoding 134 and 133 genes respectively (Fig. S2).

Chloroplast structural haplotypes across the plant tree of life

For these 90 species, we obtained adequate long-read sequencing data to assess chloroplast genome structural heteroplasmy for 61 land plants species. Across all 61 species, we find just three structural haplotypes (Fig. 1, Table 1). All 58 of the Angiosperms in our dataset are heteroplasmic, and contain roughly equal frequencies of haplotypes A and B (binomial test for departure from equal frequency, p value >0.05 in all cases). The remaining three species are not heteroplasmic (binomial test for departure from equal frequency $p = 2.2 \times 10^{-16}$ in all cases). The two Gymnosperms in our dataset both contain only haplotype B, and one of the

Pteridophytes in our dataset (*Selaginella tamariscina*) contains only haplotye C. Fig.

2 shows clearly that, as expected, there is a trend for datasets with larger sample sizes

to show relative haplotype abundances closer to exactly 50/50.

Across 8 different samples taken from a single large *Eucaltyptus pauciflora* plant

(Table S2), we detected roughly equal frequencies of haplotypes A and B in all 8

samples (binomial p-values > 0.05 in all cases), no evidence for differences in relative

frequencies among samples (chi-square p-value > 0.05), and no evidence for

phylogenetic signal in the any of the relative abundances (p-value > 0.05 in all tests),

where the underlying topology and branch lengths are taken to be the physical

structure of the tree itself (see Materials and Methods).

Discussion

In this study, we develop a new method to detect and quantify structural heteroplasmies in the chloroplast genome. By applying this method to a broad range of plant species including representatives from Angiosperms, Gymnosperms and Pteridophytes, we show that of the large number of possible structural haplotypes, only a very small number are observed in nature. For example, species with IRs almost universally contain the same two structural haplotypes in ratios that do not depart significantly from 50/50. The only exceptions to this are species in which the repeats are not inverted (e.g. one of the Pteridophytes in our dataset), or in which the repeats are inverted but much reduced in length (e.g. both of the Gymnosperms in our dataset). In these species, we detect just one haplotype. Our results demonstrate not only that the relative abundance of the two most common haplotypes stable across the Angiosperm tree of life (i.e. all 58 Angiosperms in our dataset had a frequency that was not different from 50/50), but that it also stable within an individual plant (i.e. within a single large plant, all sampled leaves contained both haplotypes at a ratio that was not different from 50/50). This suggests that the process which maintains the equal frequencies of the two commonly-observed haplotypes operates relatively rapidly.

The most parsimonious interpretation of our results suggests that structural heteroplamsy should be extremely common across all angiosperms. In light of this, it may be necessary to re-examine many previous suggestions of structural differences

in the chloroplast genomes of angiosperms *between* species (9-12). For example,

Ibrahim *et al* (9) suggested that the SSC of *Gossypium barbadense* was inverted when

compared to *Gossypium hirsutum.* However, our study shows that both *Gossypium*

barbadense and *Gossypium hirsutum* contain two structural haplotypes (haplotypes A

and B (Table 1)), thus showing that there are no structural differences between these

two species. Future studies may benefit from assuming at the outset that it is highly

likely that the chloroplast genome of any angiosperm species, particularly if it

contains two long inverted repeats, is most likely to exist in two roughly equimolar

haplotypes, haplotypes A and B shown in Fig. 1.

Of the 61 species for which we had sufficient data, we find just three species in which

there is no evidence for structural heteroplasmy in the chloroplast genome:

Selaginella tamariscina, *Pinus taeda* and *Picea sitchensis*. *Selaginella tamariscina* is

a pteridophyte in which the repeats in the chloroplast genome are positioned in-line

rather than inverted (18). The existence of a single structural haplotype in this species

is therefore consistent with the hypothesis that the process which generates

chloroplast structural heteroplasmy is mediated by long IRs. In the gymnosperms

Pinus taeda and *Picea sitchensis*, the repeats are inverted but their lengths are highly

reduced from the typical 10-30 kb in most species to just 399 bp and 440 bp,

respectively. The existence of a single structural haplotype in these species is also

consistent with the flip-flop recombination theory, because it is feasible that the

highly reduced IRs preclude the formation of the dumbbell-like structure (4) which is

necessary to activate flip-flop recombination. Whether or not flip-flop recombination

is the underlying cause, our results confirm that chloroplast structural heteroplasmy

appears to require the existence of two long IRs in the chloroplast genome, and

therefore that groups which lack this trait would be expected to lack chloroplast

structural heteroplasmy. For example, chloroplast genomes in the Pinaceae and

Cupressophytes (two major groups of gymnosperms) usually lack one IR or have IRs

that are highly reduced in size (13, 19-24), and we therefore expect that these species

would lack chloroplast structural heteroplasmy.

In this study, we focussed on examining heteroplasmies in the quadripartite structure

of the chloroplast genome. However, other smaller structural heteroplasmies have

been observed in some chloroplast genomes, and their existence challenges the

conclusion that long inverted repeats are a pre-requisite for the existence of structural

heteroplasmy in the more general sense. Although flip-flop recombination may be

absent when the IR regions are small (see above), some studies have suggested that

homologous recombination could induce structural heterplasmy from other much

smaller inverted repeats in chloroplast genomes (23, 24). For instance, the *trnQ-UUG*

gene (~150 bp) is duplicated in the LSC region in some *Juniperus* species, and this

duplication is associated with a 36 kb inversion resulting in the coexistence of two

isomeric chloroplast genome structures, although their abundance is unequal (0.8% -

5.0%) (23). Qu *et al* (24) reported a similar finding in other Cupressoideae species,

and suggest that even a pair of very small inverted repeats (11 bp) could induce a 34

kb inversion and resulting structural heteroplasmy in *Calocedrus macrolepis*. In light

of these results, it remains unclear why the highly reduced IRs of *Pinus taeda* and

Picea sitchensis studied here fail to induce structural heteroplasmy, while much

shorter inverted repeats in regions of the chloroplast genome outside the IR region

apparently do induce structural heteroplasmy. One possibility is that different

processes mediate the formation of structural heteroplasmies in these cases (23, 24).

Another possibility is that the frequency of the alternative haplotype is too low for us

to detect in *Pinus taeda* and *Picea sitchensis*. However, if this were the case the

frequency of haplotype A in these species would have to be extremely low: for

example in *Picea sitchensis* we detected 2845 long reads supporting haplotype B and

no reads supporting haplotype A.

The Cp-hap pipeline we present here provides a convenient method for quantifying

structural heteroplasmy in chloroplast genomes. It adds to suite of existing methods

such as restriction digests (3, 4) and BES (5), but differs from these methods in that it

uses long sequencing reads to provide direct evidence for uniquely-identifiable

chloroplast structures. It is of note that the two available long-read sequencing

methods, PacBio and Oxford Nanopore sequencing, differ in their utility for detecting

chloroplast structural heteroplasmy. For example, only ~63% (46 out of 73 species) of

the PacBio datasets we used in this study contained enough long chloroplast reads to

adequately measure chloroplast structural heteroplasmy. In contrast, ~88% (15 out of

17 species) of Oxford Nanopore datasets provided enough long-read to measure the

chloroplast genome heteroplasmy. Given the increasingly popularity of Oxford

Nanopore sequencing, and ongoing improvements in the read-lengths available from

PacBio sequencers, we hope that the simplicity of the Cp-hap pipeline will accelerate

further work on chloroplast structural heteroplasmy.

Materials and Methods

Cp-hap, a new method for quantifying chloroplast structural

haplotypes using long-read sequencing data

New long-read sequencing methods such as those developed by Oxford Nanopore can

routinely sequence single DNA molecules of 10s of kb in length. These molecules can

be used to provide direct evidence for the existence and relative abundance of a large

range of chloroplast genome haplotypes. Furthermore, long-reads from the chloroplast

genome are often highly abundant in plant genome sequencing projects, because the

chloroplast genome is typically present in copy numbers at least two orders of

magnitude higher than the nuclear genome (25, 26). Thus, most whole genome

sequencing projects of plant provide abundant information for quantifying chloroplast

structural haplotypes. Here, we describe a simple approach that uses this long-read

sequencing data to estimate and quantify chloroplast structural haplotypes. We call

this approach the Cp-hap pipeline, and provide open source code and detailed

instructions for running this pipeline on GitHub at https://github.com/asdcid/Cp-hap.

The Cp-hap pipeline consists of three steps: (i) create a fasta file that contains all

haplotypes of interest; (ii) map the long-reads to the potential structures in step (i); (iii)

use the results of step (ii) to quantify the relative abundance of all possible haplotypes.

Below, we describe each of these steps in more detail.

Step (i) requires us to create a fasta file containing all of the structural haplotypes of

interest. To create a list of all possible haplotypes from the LSC, SSC, and IR regions,

we first consider that each of these regions can have four different orientations:

original, reversed, complement, and reverse-complement (Fig. S3A). This results in

256 possible structural haplotypes (256 = 4x4x4x4) in which the regions retain the

ordering conserved across plant chloroplast genomes of LSC-IRA-SSC-IRB. These

256 haplotypes can be grouped into 128 identifiable structures, since it is not possible

to distinguish between one structure and its direct complement given the two-stranded

nature of the DNA molecule.

To uniquely identify one of these 128 structures, a single sequencing read would need

to cover at least some parts of all four regions (LSC, SSC and the two IR regions), for

which the read would need to be at least 30-50 kb. This is because to cover all four

regions, at a minimum a read must entirely cover the SSC (~20 kb) region and one IR

region (10-30 kb) and at least partially cover the LSC region and the other IR region.

But the abundance of reads of this length tends to be relatively low in most datasets,

meaning that in most cases it is almost impossible to uniquely identify one of the 128

possible structures by simply mapping a read to each structure.

To reduce the length of long-reads required to uniquely identify chloroplast structural

haplotypes, the Cp-hap pipeline assumes by default that the two large repeat regions

are always inverted. When assuming that the IR regions are always inverted, there are

only 32 uniquely identifiable chloroplast genome structural haplotypes (Fig. S1). In

this situation, a read only needs to entirely cover one IR region and partially cover the

two adjacent LSC and SSC regions to provide evidence to uniquely identify one of the

32 structures (Fig. S3B). Because of this, this approach can provide direct evidence

for one of the 32 possible structures using reads that are just 10-30 kb in size (i.e. at

least as long as the length of a single IR region). Reads of this length are often highly

abundant in long-read datasets, meaning that the default Cp-hap method is readily

applicable to many long-read datasets being produced today.

In Step (i), the default Cp-hap pipeline creates a fasta file of the 32 possible structural

haplotypes when assuming that the IRs are inverted by simply combining the LSC,

SSC and IR region sequences with different orientations. After creating each

structural haplotype, we then duplicate and concatenate it, such that single reads

which span the point at which the genome was linearised will still align successfully

to the relevant haplotype in the fasta file. This creates a fasta file with 32 sequences,

each of which represents one of the 32 uniquely identifiable chloroplast structural

haplotypes, and each of which contains a sequence that is twice as long as the

chloroplast genome of interest.

In step (ii) of the pipeline, we align our long-reads to our reference set of 32 possible

structural haplotypes with Minimap2 (27), using "–secondary=no" setting to ensure

that only the best alignment for each read is retained. After aligning all long-reads to

the set of 32 possible structural haplotypes, we examine each mapped read, and retain

only those that completely cover a whole IR region and at least 1 kb of the adjacent

LSC and SSC regions. We then consider that these reads, which we term valid reads,

uniquely identify one of the 32 possible structural haplotypes.

In step (iii) of the Cp-hap pipeline, we calculate the relative proportion of each of the

32 possible haplotypes. To do this, we simply divide the number of valid reads that

aligned to each of the haplotypes by total number of reads that aligned to all

haplotypes. These proportions provide direct estimates of the relative abundance of all

32 uniquely identifiable chloroplast genome structural haplotypes.

However, we acknowledge that the IR regions are not always the case. For example,

the large repeat regions are positioned in-line instead of inverted in *Selaginella*

tamariscina chloroplast genome (18). In cases such as this, Cp-hap pipeline is also

able to figure out the structure. Here, we provide an example that how Cp-hap

pipeline confirms a chloroplast genome with atypical structure, such as a paired of

in-line repeats (Fig. S3C). Since Cp-hap duplicates and concatenates all default 32

structures (with IRs), reads from chloroplast genome with in-line repeats are able to

map to Block A and/or C region of LSC_IR_SSC_IRrc structure (one of the 32 default

structure in Cp-hap pipeline), or Block B region of LSC_IRrc_SSC_IR structure (the

other default structure in Cp-hap pipeline). Those reads are impossible to map to

Block B region of LSC_IR_SSC_IRrc, or Block A or C region of LSC_IRrc_SSC_IR.

Therefore, if some reads only map to Block A and/or C region of LSC_IR_SSC_IRrc

and Block B region of LSC_IRrc_SSC_IR, it suggests that the chloroplast genome

acquires in-line repeats.

In sum, Cp-hap pipeline can provide direct evidence to identify one of the 32

chloroplast structural haplotypes with a pair of IRs, and provide indirect evidence to

identify other 96 chloroplast genome structural haplotypes with a pair of non-inverted

repeats.

Assessing chloroplast structural heteroplasmy across the plant tree of

life

We used the Cp-hap pipeline to assess chloroplast genome structural heteroplasmy

across the plant tree of life. To do this we searched for publicly available long-read

data from land plants in the NCBI SRA database (https://www.ncbi.nlm.nih.gov). We

only selected datasets in which the average read length was >9 kb and for which the

total amount of sequence data was >5Gb. 90 of the 164 potential species with datasets

met this requirement in April, 2019.

For the 90 species, 19 of them do not have available chloroplast reference genomes.

Due to the relative conservation of land plant chloroplast genomes, we used the

chloroplast genome in the same genus as the reference genome for 17 species (Table

S1). For the final two species, *Herrania umbratica* and *Siraitia grosvenorii*, we

assembled and annotated chloroplast genomes *de novo* using long and short-reads

(Table S1) following the pipeline described in (28). The chloroplast genome of

Herrania umbratica and *Siraitia grosvenorii* have been depostied in GitHub

(https://github.com/asdcid/Cp-hap/tree/master/Chloroplast%20genome).

Full details of the reads, accession numbers, and chloroplast genomes used for each of

the 90 species in our dataset are given in Table S1.

Chloroplast structural haplotype quantification

For the 90 species for which we have long-read data, we ran the Cp-hap pipeline

using default settings to generate the 32 possible haplotypes from the LSC, SSC, and

IR regions. We then ran the rest of the Cp-hap pipeline with default settings for all

species, resulting in measurements for each species of the number of valid reads that

mapped to each haplotype, and the relative abundance of each haplotype. We

calculated the relative abundance of each haplotype only if the species had more than

five valid reads in total. This filter led to us excluding 29 datasets that did not have

more than five valid chloroplast reads, resulting in a final dataset of 61 species for

which we have sufficient long-read data to estimate relative haplotype abundance.

For the *Eucalyptus pauciflora* dataset, we quantified the chloroplast structural

haplotypes not only for a single individual, but also for eight separate samples taken

from different branches of the same individual. This was possible because this data

was sequenced from leaves collected from eight different branch tips from a single

individual plant, and each sample contained sufficient long chloroplast reads to

provide reliable haplotype quantification. This provides a unique opportunity to

understand whether the relative abundance of chloroplast structural haplotypes varies

or remains constant within an individual, potentially providing insights into the

mechanisms which lead to the existence of multiple haplotypes.

Statistical analyses

Although our method can detect a large number of structural haplotypes, across all of

the species we examined with inverted repeats, we observed three haplotypes,

haplotypes A, B, and C (Fig. 1).

Because no species contained more than two haplotypes, we used a binomial test to

ask whether the abundances of haplotype A and B differ from a 1:1 ratio in each

species, or in each individual sample in the case of *Eucalyptus pauciflora*. A

significant result from a binomial test suggests that the observed haplotype

abundances are unlikely to be explained by an underlying 1:1 ratio of chloroplast

structural haplotypes.

In addition, since we have data for eight different branch tips of a single *Eucalyptus*

pauciflora individual, we used two additional approaches to test for differences

between samples. First, we used a chi-square test to ask whether the abundances of

the two haplotypes differ among samples. Second, we tested for phylogenetic signal

among the relative abundance of haplotypes observed in the eight branch tips. To do

this, we used the structure of the physical tree (with branch lengths in units of

centimetres) and the phylosignal package (29) in R (30). This package implements

five commonly-used tests of phylogenetic signal: (i) Abouheif's Cmean (31); (ii & iii)

Blomberg's K and K* (32); (iv) Moran's I (33); (v) Pagel's Lambda (34). These tests

use different approaches to ask whether the physical structure of the individual tree

helps to explain the observed variation in the relative haplotype abundance of the

eight samples. For example, if the process that generates different haplotypes operates

relatively slowly with respect to the age of this individual, which is roughly 40 years,

then we might expect haplotype frequencies to vary due to genetic drift, and thus for

the haplotype frequencies in neighbouring parts of the tree to be more similar than

would be expected by chance. This similarity would be revealed by the detection of

significant phylogenetic signal in the relative abundance data. However, if the process

that generates the different haplotypes operates very quickly with respect to the age of

this individual, then we would expect all haplotype frequencies to be roughly equal,

and thus we would expect to observe no phylogenetic signal in the haplotype

abundance data.

Acknowledgments

We thank the help of Chen Han in the statistical analysis. We also thank the people below for providing data: Xuzhi Chao (for *Selaginella tamariscina* data), Hongyan Du (for *Eucommia ulmoides* data). This work was supported by grants from the Australian Research Council Future Fellowship, FT140100843.

References

1. S. Wicke, G. M. Schneeweiss, C. W. dePamphilis, K. F. Muller, D. Quandt, The evolution of the plastid chromosome in land plants: gene content, gene order, gene function. *Plant Mol Biol* **76**, 273-297 (2011).

2. X. M. Zheng *et al.*, Inferring the evolutionary mechanism of the chloroplast genome size by comparing whole-chloroplast genome sequences in seed plants. *Scientific Reports* **7**, 1555 (2017).

3. J. D. Palmer, Chloroplast DNA exists in two orientations. *Nature* **301**, 92 (1983).

4. D. B. Stein, J. D. Palmer, W. F. Thompson, Structural evolution and flip-flop recombination of chloroplast DNA in the fern genus Osmunda. *Current Genetics* **10**, 835-841 (1986).

5. G. Martin, F. C. Baurens, C. Cardi, J. M. Aury, A. D'Hont, The complete chloroplast genome of banana (Musa acuminata, Zingiberales): insight into plastid monocotyledon evolution. *PLoS One* **8**, e67350 (2013).

6. T. Brears, C. L. Schardl, D. M. Lonsdale, Chloroplast genome organisation in sugar

beet and maize. *Plant Mol Biol* **6**, 171-177 (1986).

7.   H. J. Bohnert, W. Löffelhardt, Cyanelle DNA from Cyanophora paradoxa exists in two

forms due to intramolecular recombination. *FEBS Letters* **150**, 403-406 (1982).

8.   J. Aldrich, B. Cherney, E. Merlin, C. Williams, L. Mets, Recombination within the

inverted repeat sequences of the Chlamydomonas reinhardii chloroplast genome

produces two orientation isomers. *Current Genetics* **9**, 233-238 (1985).

9.   R. I. Ibrahim, J. Azuma, M. Sakamoto, Complete nucleotide sequence of the cotton

(Gossypium barbadense L.) chloroplast genome with a comparative analysis of

sequences among 9 dicot plants. *Genes Genet Syst* **81**, 311-321 (2006).

10.   M. Yang *et al.*, The complete chloroplast genome sequence of date palm (Phoenix

dactylifera L.). *PLoS One* **5**, e12762 (2010).

11.   Y. Liu *et al.*, Complete chloroplast genome sequences of Mongolia medicine Artemisia

frigida and phylogenetic relationships with other plants. *PLoS One* **8**, e57533 (2013).

12.   Y. Zhang, L. Li, T. L. Yan, Q. Liu, Complete chloroplast genome sequences of Praxelis

(Eupatorium catarium Veldkamp), an important invasive species. *Gene* **549**, 58-69

(2014).

13.   C. S. Wu, C. P. Lin, C. Y. Hsu, R. J. Wang, S. M. Chaw, Comparative chloroplast

genomes of pinaceae: insights into the mechanism of diversified genomic

organizations. *Genome Biol Evol* **3**, 309-319 (2011).

14.   R. Cronn, M. Cedroni, T. Haselkorn, C. Grover, J. F. Wendel, PCR-mediated

recombination in amplification products derived from polyploid cotton. *Theor Appl*

Genet **104**, 482-489 (2002).

15. D. J. Lahr, L. A. Katz, Reducing the impact of PCR-mediated recombination in

molecular evolution and environmental studies using a new-generation high-fidelity

DNA polymerase. *Biotechniques* **47**, 857-866 (2009).

16. C. Belser *et al.*, Chromosome-scale assemblies of plant genomes using nanopore

long reads and optical maps. *Nature Plants* **4**, 879-887 (2018).

17. S. Deschamps *et al.*, A chromosome-scale assembly of the sorghum genome using

nanopore sequencing and optical mapping. *Nature Communications* **9**, 4844 (2018).

18. Z. Xu *et al.*, Genome Analysis of the Ancient Tracheophyte Selaginella tamariscina

Reveals Evolutionary Features Relevant to the Acquisition of Desiccation Tolerance.

Mol Plant **11**, 983-994 (2018).

19. C. S. Wu, Y. N. Wang, C. Y. Hsu, C. P. Lin, S. M. Chaw, Loss of different inverted

repeat copies from the chloroplast genomes of Pinaceae and cupressophytes and

influence of heterotachy on the evaluation of gymnosperm phylogeny. *Genome Biol*

Evol **3**, 1284-1295 (2011).

20. C. S. Wu, S. M. Chaw, Large-Scale Comparative Analysis Reveals the Mechanisms

Driving Plastomic Compaction, Reduction, and Inversions in Conifers II

(Cupressophytes). *Genome Biol Evol* **8**, 3740-3750 (2016).

21. L. A. Raubeson, R. K. Jansen, A rare chloroplast-DNA structural mutation is shared by

all conifers. *Biochemical Systematics and Ecology* **20**, 17-24 (1992).

22. X. Yi, L. Gao, B. Wang, Y. J. Su, T. Wang, The complete chloroplast genome

sequence of Cephalotaxus oliveri (Cephalotaxaceae): evolutionary comparison of

cephalotaxus chloroplast DNAs and insights into the loss of inverted repeat copies in

498   gymnosperms. *Genome Biol Evol* **5**, 688-698 (2013).

499 23. W. Guo *et al.*, Predominant and substoichiometric isomers of the plastid genome

500   coexist within Juniperus plants and have shifted multiple times during cupressophyte

501   evolution. *Genome Biol Evol* **6**, 580-590 (2014).

502 24. X.-J. Qu, C.-S. Wu, S.-M. Chaw, T.-S. Yi, Insights into the Existence of Isomeric

503   Plastomes in Cupressoideae (Cupressaceae). *Genome Biology and Evolution* **9**,

504   1110-1119 (2017).

505 25. S. A. Morley, B. L. Nielsen, Chloroplast DNA Copy Number Changes during Plant

506   Development in Organelle DNA Polymerase Mutants. *Front Plant Sci* **7**, 57 (2016).

507 26. H. Daniell, C. S. Lin, M. Yu, W. J. Chang, Chloroplast genomes: diversity, evolution,

508   and applications in genetic engineering. *Genome Biol* **17**, 134 (2016).

509 27. H. Li, Minimap and miniasm: fast mapping and de novo assembly for noisy long

510   sequences. *Bioinformatics* **32**, 2103-2110 (2016).

511 28. W. Wang *et al.*, Assembly of chloroplast genomes with long- and short-read data: a

512   comparison of approaches using Eucalyptus pauciflora as a test case. *BMC*

513   *Genomics* **19**, 977 (2018).

514 29. F. Keck, F. Rimet, A. Bouchez, A. Franc, phylosignal: an R package to measure, test,

515   and explore the phylogenetic signal. *Ecol Evol* **6**, 2774-2780 (2016).

516 30. R Core Team, R: A language and environment for statistical computing. *R Foundation*

517   *for Statistical Computing* (2013).

518 31. E. Abouheif, A method for testing the assumption of phylogenetic independence in

519   comparative data. *Evolutionary Ecology Research* **1**, 895-909 (1999).

32. S. P. Blomberg, T. Garland, Jr., A. R. Ives, Testing for phylogenetic signal in

comparative data: behavioral traits are more labile. *Evolution* **57**, 717-745 (2003).

33. P. A. Moran, Notes on continuous stochastic phenomena. *Biometrika* **37**, 17-23

(1950).

34. M. Pagel, Inferring the historical patterns of biological evolution. *Nature* **401**, 877-884

(1999).

Figure Legends

528
529
Fig. 1. The three different structural haplotypes of chloroplast genomes detected in
this study. The green region is the Long Single Copy (LSC) region and blue region is
the Short Single Copy (SSC) region. The two black regions are the Inverted Repeat
(IR) regions. The arrow denotes 5'-3' orientation. *psbA* is in the minus strand of LSC
region, whereas *rrn23* is in the plus strand of IR regions. *ndhF* is in the minus strand
of SSC region, while *ccsA* is in the plus strand of SSC region. For ease of
communication, we use the relative order of three genes (*psbA* in LSC, and *ndhF* and
ccsA in the SSC) to label these two haplotypes 'A' and 'B'. In haplotype A, these
genes are ordered *psbA-ndhF-ccsA*. In haplotype B these genes are ordered
psbA-ccsA-ndhF. For haplotype C, these three genes are ordered the same as
haplotype A, but the repeat regions are in-line rather than inverted.

541
542
543
Fig. 2. The relationship between haplotype counts and haplotype frequencies. Each
point is one species. A blue dot means binomial p-value > 0.05, whereas the orange
dot means binomial p-value ≤ 0.05. Two species, *Pinus taeda*, *Picea sitchensis* only
showed evidence of haplotype B, therefore the frequencies of them were 0. For the
remaining 58 species, they show higher level of equal frequency between haplotype A
and B with the increase of sample size. One species (*Selaginella tamariscina*) is
omitted from this figure because it contained only a third haplotype (haplotype C,
Figure 1).

 Table 1. The frequency of existing chloroplast genome haplotypes from 61 species.

Division	Order	Species	Size (bp)	LSC (bp)	SSC (bp)	IR (bp)	HA	HB	Frequency	p-value
Angiospermae	Poales	*Andropogon gerardii*	139,712	81,548	12,592	22,786	31	31	0.50	1.00
		Brachypodium†mexicanum	~135,199	~79,447	~12,668	~21,542	403	373	0.52	0.30
		Eleusine coracana†	~135,151	~80,667	~12,646	~20,919	14	11	0.56	0.69
		Hordeum vulgare	136,462	81,671	12,701	21,045	347	329	0.51	0.51
		Oropetium thomaeum†	~133,880	~79,181	~12,591	~21,054	27	28	0.49	1.00
		Oryza coarctata	134,750	80,816	12,334	20,800	40	38	0.51	0.91
		Oryza longistaminata	134,567	80,596	12,357	20,807	8	4	0.67	0.39
		Oryza meyeriana	136,133	81,832	12,499	20,901	5	5	0.50	1.00
		Panicum miliaceum	140,048	81,841	9,603	24,302	37	22	0.63	0.07
		Saccharum officinarum	141,176	83,046	12,540	22,795	51	56	0.48	0.70
		Saccharum spontaneum	141,168	83,046	12,544	22,789	25	28	0.47	0.78
		Sorghum bicolor	140,754	83,733	12,503	22,259	94	71	0.57	0.09
		Zea mays	140,384	82,352	12,536	22,748	564	555	0.50	0.30
		Zoysia japonica	135,854	81,348	12,582	20,962	35	54	0.39	0.06
	Zingiberales	*Musa schizocarpa†*	~169,503	~87,828	~11,487	~35,094	192	210	0.48	0.40
	Asparagales	*Yucca aloifolia†*	~157,785	~86135	~18,292	~26,679	87	105	0.45	0.22
		Yucca filamentosa	157,785	86,135	18,292	26,679	200	202	0.50	0.96
	Ranunculales	*Papaver somniferum*	152,931	83,029	17,920	25,991	12	13	0.48	1.00
	Ericales	*Actinidia eriantha*	156,964	88,759	20,541	23,832	122	94	0.56	0.11
	Lamiales	*Osmanthus fragrans*	155,326	86,564	17,370	25,696	65	63	0.51	0.93
		Salvia splendens†	~153,995	~84,573	~17,590	~25,916	15	11	0.58	0.56
	Solanales	*Cuscuta campestris*	86,749	50,978	7,063	14,354	5	8	0.38	0.58

	Nicotiana attenuata	155,886	86,602	18,518	25,383	3	6	0.33	0.51
Asterales	*Lactuca sativa*	152,765	84,103	18,596	25,033	39	31	0.56	0.40
Cucurbitales	*Siraitia grosvenorii*	158,757	87,625	18,556	26,288	5	4	0.56	1.00
Fabales	*Ammopiptanthus nanus*	154,084	84,070	18,014	26,000	31	43	0.42	0.24
	Arachis hypogaea	156,391	85,946	18,797	25,824	17	17	0.50	1.00
	Vigna unguiculata	152,415	81,822	17,425	26,584	4	3	0.43	1.00
Malpighiales	*Populus trichocarpa*	157,033	85,129	16,600	27,652	58	78	0.39	0.40
	Salix purpurea	155,590	84,454	16,220	27,458	41	47	0.47	0.59
Rosales	*Fragaria vesca*	155,691	85,606	18,173	25,956	31	25	0.55	0.51
	Fragaria ananassa	155,549	85,532	18,145	25,936	4	2	0.67	0.69
	Prunus avium†	~157,859	~85,977	~19,120	~26,381	114	128	0.47	0.40
	Rosa chinensis	156,546	85,727	18,754	26,030	5	4	0.56	1.00
	Rubus occidentalis†	~155,760	~85,430	~18,768	~25,781	9	7	0.56	0.80
Fagales	*Betula pendula*	161,147	89,428	19,607	26,056	22	16	0.58	0.42
Brassicales	*Arabidopsis thaliana*	154,478	84,170	17,780	26,264	135	150	0.47	0.41
	Brassica napus	152,860	83,030	17,760	26,035	51	38	0.57	0.20
	Brassica oleracea	153,364	83,136	17,834	26,197	1,714	1,773	0.49	0.33
	Brassica rapa	153,483	83,282	17,775	26,213	1,097	1,077	0.51	0.68
Malvales	*Bombax ceiba*	158,997	89,022	21,111	24,432	71	96	0.43	0.06
	Gossypium arboreum	160,230	88,722	20,274	25,617	38	47	0.45	0.39
	Gossypium barbadense	160,317	88,841	20,294	25,591	11	14	0.44	0.69
	Gossypium bickii	159,422	88,073	20,183	25,583	333	330	0.50	0.94
	Gossypium darwinii	160,378	88,906	20,266	25,603	93	95	0.49	0.94
	Gossypium hirsutum	160,301	88,817	20,280	25,602	623	586	0.52	0.30
	Gossypium longicalyx	160,241	88,667	20,278	25,648	205	213	0.49	0.73
	Gossypium mustelinum	160,313	88,824	20,271	25,609	50	52	0.49	0.92

	Order	Species	Size	LSC	SSC	IR	HA	HB	Frequency	p-value
		Gossypium raimondii	160,161	88,654	20,205	25,651	48	38	0.56	0.33
		Gossypium tomentosum	160,433	88,932	20,273	25,614	42	61	0.41	0.08
		Theobroma cacao	160,619	89,333	20,194	25,546	86	77	0.52	0.53
	Myrtales	*Eucalyptus albens†*	~160,076	~88,828	~18,476	~26,386	20	25	0.44	0.55
		Eucalyptus marginata	160,076	88,828	18,476	26,386	13	16	0.45	0.71
		Eucalyptus melliodora	160,386	89,073	18,557	26,378	371	373	0.50	0.97
		Eucalyptus pauciflora	159,942	88,787	18,421	26,367	1,431	1,459	0.50	0.62
	Sapindales	*Acer yangbiense†*	~156,262	~86,018	~18,072	~26,086	104	94	0.53	0.52
		Citrus maxima	160,133	87,739	18,395	26,999	2	6	0.25	0.29
		Xanthoceras sorbifolium	161,231	85,299	18,692	28,620	110	119	0.48	0.60
Gymnosperm	Pinales	*Pinus taeda*	121,530	66,444	54,288	399	0	898	0.00	2.2e-16*
		Picea sitchensis	124,049	67,411	55,758	440	0	2,845	0.00	2.2e-16*
Pteridophytes	Selaginellales	*Selaginella tamariscina ‡*	126,399	53,176	47,573	12,825	0	0	-	-

*: p-value ≤ 0.05

†: The long-reads were mapped to the chloroplast genome from the same genius species due to the lack of chloroplast genome of this species, so

the lengths of LSC, SSC and IR are around (~).

‡: *Selaginella tamariscina* only present one haplotype, haplotype C, with 174 counts, and p-value is 2.2e-16.

Size: the chloroplast genome size

HA/B: Haplotype A/B, the number of supported reads of LSC and SSC having identical/opposite orientation.

Frequency: the proportion of the count of haplotype A in the count of haplotype A+B