**Evolutionary dynamics of abundant 7 bp satellites in the genome of *Drosophila virilis***

Jullien M. Flynn[1], Manyuan Long[2], Rod A. Wing[3], Andrew G. Clark[1]

[1]Department of Molecular Biology and Genetics, Cornell University, Ithaca, USA

[2]Department of Ecology and Evolution, University of Chicago, Chicago, USA

[3]Arizona Genomics Institute, School of Plant Sciences, University of Arizona, Tucson, Arizona, USA

Corresponding author:

Jullien M. Flynn

jmf422@cornell.edu

**Abstract**

The factors that drive the rapid changes in satellite DNA genomic composition we see in eukaryotes

are not well understood. *Drosophila virilis* has one of the highest relative amounts of simple

satellites of any organism that has been studied, with an estimated >40% of its genome composed

of a few related 7 bp satellites. Here we use *D. virilis* as a model to understand technical biases

affecting satellite sequencing and the evolutionary processes that drive satellite composition. By

analyzing sequencing data from Illumina, PacBio, and Nanopore platforms, we identify platform-

specific biases and suggest best practices for accurate characterization of satellites by sequencing.

We use comparative genomics and cytogenetics to demonstrate that the highly abundant satellite

family arose from a related satellite in the branch leading to the virilis phylad 4.5 - 11 million years

ago before exploding in abundance in some species of the clade. The most abundant satellite is

conserved in sequence and location in the pericentromeric region but has diverged widely in

abundance among species, whereas the satellites nearest the centromere are rapidly turning over

in sequence composition. By analyzing multiple strains of *D. virilis*, we saw that one centromere-

proximal satellite is increasing in abundance along a geographical gradient while the other is

contracting in an anti-correlated manner, suggesting ongoing conflicts at the centromere. In

conclusion, we illuminate several key attributes of satellite evolutionary dynamics that we

hypothesize to be driven by processes like selection, meiotic drive, and constraints on satellite

sequence and abundance.

**Introduction**

Repetitive DNA is abundant in most eukaryotic genomes, and is now understood to be correlated

with the manifold variation in genome size across the tree of life (Elliott and Gregory 2015). For

most species, transposable elements (TEs) dominate the repeat landscape, including in humans,

plants, and *Drosophila melanogaster*. Satellite DNA, which is characterized by tandem repeats

spanning long arrays, very rarely has dominated a genome to a similar extent as TEs. An

unprecedented case is that of *Drosophila virilis*, the Drosophila species with the largest estimated

genome size (up to 389 Mb) (Bosco *et al.* 2007), where some 40% of the genome is comprised of

just three simple 7-mer satellites: AAACTAC, AAACTAT, and AAATTAC (Gall *et al.* 1971; Gall and

Atherton 1974). Since the 1970s, there has been no follow-up to validate the amount of 7-mers

with modern techniques, or evolutionary studies to understand how and why these satellite repeats

expanded so explosively. The genomic composition of simple satellites in *D. virilis* provides an

excellent model for an investigation of the evolutionary dynamics involved in their expansion in the

genome as well as the technical challenges facing simple satellite analysis.

Satellites are rapidly evolving in sequence and copy number, and there is a high level of

variation in satellite content among and within species (Wei *et al.* 2014, 2018). The reasons for such

dramatic variation is not well understood, and cannot be fully explained by current models.

Satellites have been long hypothesized to be slightly deleterious and therefore governed primarily

by the strength of negative selection (Ohno 1972). However, the amount of satellite in the genome

that causes negative effects that could be selected against depends on many factors and cannot be

easily predicted (Charlesworth *et al.* 1994; Gregory 2001). The fact that most organisms have

satellite repeats in or near centromeres suggests that they are important for centromere function.

Satellite repeats can also be important for maintenance of the chromocenter and packaging of

chromosomes in the nucleus (Jagannathan *et al.* 2018, 2019), and the transcripts of some satellites

may be essential for fertility (Mills *et al*. 2019). In heterozygotes with alleles that differ in

pericentromeric satellite sequence or abundance, one allele may assemble a stronger kinetochore

during female meiosis I, increasing its probability of transmission into the egg (rather than polar

bodies). This transmission advantage, known as centromere drive, allows satellites to rapidly

change in composition in the population, regardless of their whole-organism fitness effects

(Henikoff et al. 2001). If satellite DNA is an essential component of genomes or is only a burden (i.e.

is selfish), it is still not clear why some species have almost no pericentromeric satellite DNA while

others, like *D. virilis*, possess pericentromeric satellites that make up almost half of the genome.

64            Comparing the satellites of *D. virilis* to those of its sister species can elucidate when the

abundant satellites arose, and how rapidly their copy numbers and sequences evolved. *D. virilis* is

4.5 MY diverged from its sister species *D. novamexicana* and *D. americana*, which are both

restricted to North America, unlike globally-distributed *D. virilis* (Caletka and McAllister 2004).  *D.*

*novamexicana* and *D. americana* have a smaller estimated genome size than *D. virilis* (~250 Mb vs.

389 Mb), suggesting these species may have less satellite content (Bosco *et al.* 2007). Additionally,

using intra-species comparisons across global populations can give indications about factors that

may be influencing satellite dynamics. For example, in *D. melanogaster*, patterns of abundance of

the *Prodsat* satellite closely mirror the migration patterns of species, suggesting an ongoing

expansion of this satellite (Wei *et al.* 2014). Genetic drift or meiotic drive may contribute to

patterns of geographical gradients of satellite abundance. We can also use intra-species data to

pose hypotheses about non-neutral processes that may be driving satellite content. Previous work

has shown evidence for conflicts or trade-offs between satellites within the genome, and these

constraints can be illuminated by analyzing satellites in several strains (Flynn *et al.* 2017, 2018).

Genome-wide characterization of satellites has taken off since high-throughput sequencing

has become widely available. We have learned from several informative studies about the

sequences and relative abundances of satellites in various species (Pavlek *et al.* 2015; Flynn *et al.*

2017; de Lima *et al.* 2017; Wei *et al.* 2018), but technical challenges may prevent accurate

quantitative estimates. Satellites may be more prone to errors or biases in the sequencing process

that do not affect the better studied regions of the genome. Satellites are difficult to assemble even

with long-read sequencing (Chang and Larracuente 2019). The genome assembly of *D. virilis* is

approximately half its estimated genome size by flow cytometry (~200 Mb vs 389 Mb) (Bosco *et al.*

2007), and it is likely that much of what is missing is simple satellite DNA. However, even using

alignment-free raw read methods have not produced satellite DNA estimates that approach the

amount that is missing from the genome assembly and was estimated from early work (Gall *et al.*

1971; Gall and Atherton 1974; Wei *et al.* 2018). Now, as long read sequencing is also being

exploited to study satellites, we must evaluate satellite DNA abundance estimates to assess if there

are platform-specific biases that may affect evolutionary analysis of satellite DNA.

The purpose of this paper is two-fold; first to explore the technical biases preventing

accurate characterization and quantification of simple satellites, and second to use a comparative

approach to understand the evolutionary dynamics of the extremely abundant 7mers in the *D. virilis*

group. First, we characterize satellites in *D. virilis* sequencing data from different platforms and

assess biases that affect accurate satellite characterization. We then use comparative genomics and

cytogenetics in *D. virilis* and its sister species to understand the composition and changes in the

highly abundant simple satellites. Finally we sequence multiple strains of *D. virilis* and sister species

to estimate polymorphism in satellite abundance and infer processes that may be influencing their

evolution. From this we infer that there are likely a variety of understudied processes affecting

satellite DNA in this organism, including positive selection, meiotic drive, and constraints and trade-

offs between satellites.

**RESULTS**

**Technical biases in characterizing simple satellites from sequencing**

*Long-read genome assemblies have an under-representation of simple satellites*

Long-read sequencing technologies have an advantage because of their long reads, but a

disadvantage due to their high error rate, prompting a need for extensive alignments for error-

correction and assembly. First we asked whether assemblies from long read technologies can better

assemble simple satellite reads than the previous Sanger assembly. We compared the amount of

simple 7-mer satellites (AAACTAC, AAACTAT, AAATTAC, AAACAAC) in three *D. virilis* genome

assemblies: the CAF1 assembly produced from Sanger sequencing (Drosophila 12 Genomes

Consortium *et al.* 2007), a PacBio assembly produced by our group by ~100x coverage (available at

https://www.ncbi.nlm.nih.gov/bioproject/?term=txid7214[Organism:noexp]), and a Nanopore

assembly produced from ~20x sequencing coverage (Miller *et al.* 2018). All assemblies were

approximately the same size at ~200 Mb. The PacBio and Nanopore assemblies contained a

similarly low amount of simple 7-mer satellites, 29 and 28 kb, respectively. The CAF1 assembly,

however contained 7.36 Mb of these satellites. This discrepancy is likely largely due to the

difference in assembly algorithms used for short read and long read data. Long reads must be

aligned and corrected to be incorporated into the assembly because of their high error rate,

whereas this is not necessary for Sanger-based assemblies. Use of modified methods can improve

assemblies of repetitive regions (Chang and Larracuente 2019), but for highly homogeneous simple

satellites, whose arrays span 10-100x longer than the current maximum read length, it is practically

impossible to produce a continuous assembly.

*Simulations to assess simple repeat quantification from long read sequencing data*

Due to assembly issues of simple satellites, they must be quantified from raw unassembled reads.

Long read sequencing data poses a significant challenge because of the high error rate including a

high indel rate in the raw reads. We therefore used two different approaches along with

simulations to assess their accuracy. The first approach used k-Seek (Wei et al. 2014) to select

repeat-rich reads and then Phobos ([https://www.ruhr-uni-bochum.de/ecoevo/cm/cm_phobos.htm)](https://www.ruhr-uni-bochum.de/ecoevo/cm/cm_phobos.htm)

to quantify satellites. This approach allows for *de novo* discovery of satellite sequences. We used

Noise-Cancelling Repeat Finder (NCRF, Harris *et al.* 2019) for our second approach, providing our

target satellites. Both methods are relatively sensitive to imperfect repeats, which we expect with

the high error rate of long-read sequencing.

To evaluate our approaches, we created a mock *D. virilis*-like genome containing

pericentromeric and centromeric repeats on each of five chromosomes (See Materials and

Methods). We then simulated 10x PacBio reads from this genome, and then quantified satellites

using both approaches. NCRF works by doing alignments of target satellites to the reads and

allowing up to a user-specified maximum divergence. To determine the most appropriate maximum

divergence, we simulated a range of values for this parameter from 18-30% and chose the lowest

asymptotic value - which was 25% in this case (Figure S1). NCRF found almost the same amount of

satellites that truly existed in the mock genome whereas the k-Seek + Phobos method only found

about 20% (Figure 1A).

*The amounts and biases in simple 7mer repeats differ between Nanopore and PacBio sequencing*

*reads*

Next, we quantified simple satellites in the long-read data generated from our 100x PacBio

sequencing and 20x Nanopore sequencing using the two approaches mentioned above. Unlike in

the simulations, both approaches produced very similar (but lower than expected) estimates at 8.8-

10.9 Mb for the PacBio data (Figure 1B). The Nanopore data contained almost 3 times the 7mer

satellites compared to PacBio, with 23.4 - 28.2 Mb (Figure 1C). This may represent a platform-

specific difference in the ability to sequence long arrays of simple tandem repeats. Both the PacBio

reads and the Nanopore reads contained a greater amount of simple satellites than data produced

in our lab previously with Illumina HiSeq sequencing (Wei et al. 2018), however did not approach

the estimated >100 Mb in the genome.

Both the PacBio and Nanopore reads contained large amounts of what we expect to be

artefactual repeats, which were found with the k-Seek + Phobos approach, and validated with

NCRF. NCRF found 4.4 Mb (normalized to 1x genome coverage) of AAACGAC in the PacBio reads.

This satellite was not found in the Nanopore data or Illumina data (this and previous studies) or in

previous studies that characterized the most abundant satellites in *D. virilis*. Manual inspection

proved that the AAACGAC satellite was the true consensus found in long arrays in the reads and did

not represent an error in our approaches' characterization of satellites. Similarly, AAATCAC,

AGCCTAT, ACAGGCT, and AATGG were found in megabase quantities (after normalization) in the

Nanopore data - whereas these satellites were not found in Illumina or PacBio data. We suggest

these satellites are also technical artifacts introduced at the base-calling level.

**Fig1**: Issues in quantifying simple satellites in sequencing data (all data shown is *D. virilis*). (**A**) Cumulative stacked barplot comparing the performance of the two tested approaches on PacBio data simulated with PBSim from a mock genome. (**B**) Comparing the results of the two approaches on the real PacBio data; "other" refers to additional satellites in the family, including suspected artefactual ones (AAAGCAC for PacBio and AAATCAC + AGCCTAT for Nanopore). (**C**) Same as B for Nanopore data. (**D**) Strand biases in the sequenced satellites in long read sequencing data. Satellites with asterisks are suspected artefactual ones. N refers to the number of satellite regions of reads used for the calculation. (E) Amount of satellites quantified in different datasets: imaginal discs (pure diploid), compared to flies (some polyteny), and fly data that has been quality filtered.

In the PacBio data, the relative amounts of 7mer satellites (AAACTAC, AAACTAT, and

AAATTAC) were lower than expected. This additional evidence led us to hypothesize that there

were context-specific errors in our PacBio data affecting our particular satellites. If the sequencing

were unbiased, we would expect to have an equal amount of satellites being detected on reads

coming from both DNA strands. We evaluated the strand bias in the simulated and real long-read

data for the three most abundant true satellites, as well as some artefactual satellites. We

arbitrarily label the positive strand as AAACTAC and the negative strand as GTAGTTT, etc.  In the

simulated data, the positive and negative strands of satellites were detected in equal amounts

(Figure 1D). However, there was a strong strand bias for all satellites in both the PacBio and

Nanopore data (Figure 1D). For PacBio, the real satellites AAACTAC, AAACTAT, AAATTAC had a

positive strand bias, whereas the artefactual satellite had a negative strand bias: 98% of the reads

with this satellite were from the negative strand. Based on communication with PacBio

representatives, this issue seemed to be caused by context-specific issues with base calling

algorithms used for this sequencing run. As base calling algorithms improve, these issues will likely

begin to be remedied. In fact, we received PacBio Circular Consensus Sequencing or "HiFi" data for

a closely related species, *D. americana*, and the base-calling issue was remedied. In the Nanopore

data, strand biases were even more extreme: the negative strand was sequenced almost exclusively

for real satellites AAACTAC and AAATTAC and suspect satellite AAATCAC. However, the AAACTAT

real satellite was sequenced almost exclusively on the positive strand. In this case, strand biases

may be caused by unsequenceable secondary structures developing more frequently on one strand

of the satellite DNA than the other. We analyzed Illumina NextSeq reads for *D. virilis*, and no such

strand bias was found.

*D. virilis whole-flies have 40% less pericentromeric satellites than non-polytene tissue*

Polyteny occurs in all differentiated tissues of Dipterans, and is characterized by multiple

rounds of local DNA replication within the same nucleus and without cell division, a process known

as endoreduplication (Smith and Orr-Weaver 1991; Kim *et al.* 2011). However, the pericentromeric

heterochromatin, where most satellite DNA is located, is under-replicated (Belyaeva *et al.* 1998). It

has never been tested if the level of polyteny in an adult fly makes a difference in the estimate of

satellites per genome. Thus, we sequenced adult male flies (which have multiple polytene tissues)

and imaginal discs (which are diploid) from male larvae and compared the amount of simple

satellites in these datasets. We used Illumina sequencing and PCR-free library preparations to

reduce known PCR bias (Wei *et al.* 2018). We found that for each of the four most abundant 7mer

satellites in the *D. virilis* genome, there was approximately 40% less in the flies compared to the

imaginal discs (Figure 1E). This pattern is not observed for microsatellites which are known to

localize outside of pericentromeric heterochromatin (Figure S2A). We also analyzed publicly

available *D. melanogaster* data, including flies, imaginal discs, and salivary glands (which are the

most extreme in polyteny), and observed this same pattern of under-replication of satellite repeats

in polytene tissues (Figure S2B and S2C).

*Reads with tandem repeats had lower quality scores in Illumina data*

Upon inspection with FastQC of our data from the polyteny analysis, we found a bimodal

distribution of quality scores, with one peak at 22 and another at 37 (Figure S3A). After filtering low

quality reads, the majority of the reads with simple satellites were removed (Figure S3). The

quantity of satellites was reduced by ~15 x after quality filtering (Figure 1E). It is apparent that in

our dataset, simple satellite-containing reads were highly enriched for low quality scores. We

examined other published *D. virilis* Illumina datasets to evaluate if this issue existed in other

sequencing runs. Two other datasets were available and the one that was produced on the Illumina

NextSeq platform like our data (Miller *et al.* 2018) showed the same pattern of biased quality scores

in repetitive reads (Figure S4). The dataset produced on the HiSeq platform (Wei *et al.* 2018) did not

show this pattern. It should be noted however that the amounts of 7mer satellites sequenced in the

NextSeq datasets were higher than the HiSeq dataset. Our libraries were multiplexed with other

non-*D. virilis* group samples from unrelated projects and only represented ~20% of the total

sequenced lane so that we would not have issues related to low complexity. We also noticed this

pattern (but less dramatically) in our Illumina sequencing of multiple strains.

**Related species have similar but fewer simple repeats**

*D. novamexicana* and *americana* which are 0.38 MY diverged from each other, are sister species of

*D. virilis*, which is approximately 4.5 MY diverged (Caletka and McAllister 2004) (Figure 2A). We

sequenced these species with high coverage PacBio runs and characterized and quantified satellites.

We emphasize the comparison of relative satellite amounts since all are likely under-represented.

*D. americana* was sequenced with PacBio HiFi reads, which eliminated artefactual satellites, but

make quantitative comparisons difficult since different chemistries have different efficiencies of

sequencing satellites. Nevertheless, we also found a high enrichment of 7bp satellites in *D.*

*novamexicana* and *D. americana* (Fig 2B). Interestingly, we found the most abundant satellite in *D.*

*virilis*, AAACTAC, is also the most abundant in *D. novamexicana* and *D. americana*, albeit with about

half the total amount. The second and third most abundant repeats, AAACTAT and AAATTAC,

however were not present in long tandem arrays in *D. novamexicana*. The second most abundant

satellite in *D. novamexicana* and *americana* was AAACAAC, whereas in *D. virilis* there is only a few

kilobases.

By analyzing sequencing data in more diverged species, we can infer when the AAACTAC

satellite family arose. *D. hydei* is approximately 26 MY diverged from *D. virilis* (Izumitani *et al.* 2016),

and we had PacBio long read data for this species. Here 7 bp satellites are again the most enriched

(Fig 2B), but the sequences are unrelated to those in *D. virilis* (ACCCATG, AAAGGTC from PacBio

data). We analyzed Illumina data for *D. montana*, another member of the virilis group that is 7-11

MY diverged from *D. virilis* (Ostrega and Thompson 1986; Spicer and Bell 2002) (Figure 2A). This

species does not have any AAACTAC family satellites, and in fact no enrichment of 7 bp satellites.

The most abundant satellite in *D. montana* is AAAC. From these data, we infer that the AAACTAC

family of satellites arose in the clade leading to the *D. virilis* phylad 4.5-11 MYA. We also analyzed

Illumina sequencing data for *D. lummei*, which is 3 MY diverged from *D. novamexicana/americana*

(Fig 2A). AAACTAC is conserved in *D. lummei,* but it is the only enriched 7 bp satellite in this species

and its relative estimated abundance is lower than the other three *D. virilis* phylad species.

**Complex satellites are also abundant in *D. virilis* group genomes**

We searched the high-quality genome assemblies for complex satellites (defined here as unit

lengths greater than 20 bp). In *D. virilis,* we found a 36-bp satellite

AAAACGACATAACTCCGCGCGGAGATATGACGTTCC making up ~800 kb of the assembly. This satellite

was found in previous studies and is thought to be associated with the possibly mobile element pDv

(Zelentsova et al 1986, Heikkinen et al. 1995). In *D. novamexicana*, we found a 32 bp satellite

AAAAGCTGATTGCTATATGTGCAATAGCTGAC along with a related 29 bp satellite. The 32 bp satellite

spanned over 1.1 Mb on a single 3 Mb contig in the *D. novamexicana* assembly. The non-satellite

portion of the contig had similarity to chromosome 6 (dot chromosome/Muller element F) (Figure

S5). In *D. americana*, we found this identical 32 bp satellite, but in total its span was only ~150 kb. In

all *D. virilis* group species, we also found a series of similar satellites varying in size (150-500 bp)

related to the previously described helitron central repeat that has expanded to tandem repeats in

the virilis group (Dias *et al.* 2015).

**Fluorescence *in situ* hybridization reveals evolutionary dynamics of 7 bp repeats**

The location of the 7 bp satellites on metaphase chromosomes has never been shown in the *D.*

*virilis* group. From our sequencing data, we know that the AAACTAC satellite is conserved between

*D. virilis, D. novamexicana,* and *D. americana*, but the abundance varies by approximately two-fold.

The second most abundant satellites have turned over between *D. virilis* and

*novamexicana/americana*. We used FISH of the most abundant 7mers (AAACTAC, AAACTAT,

AAATTAC, AAACAAC) in these three sister species. *D. virilis* and *D. novamexicana* have the same

karyotype with five acrocentric chromosomes plus the very small F element or "dot chromosome".

The strain of *D. americana* we used has centromere-centromere fusions between the X and 4[th]

chromosomes and the 2[nd] and 3[rd] chromosomes.

**Fig 2**. Comparative analysis of simple satellites in the *D. virilis* group. (A) Phylogeny demonstrating when satellites arose (+) and were lost (-). Dashed box: virilis phylad; solid box: virilis group. (B) Total amount of different unit lengths (k-mers) of satellites across four related species. The (*) for *D. americana* indicates that it was sequenced with PacBio HiFi reads, whereas the other species were sequenced with chemistry version 2.0. (C) DNA-FISH image of *D. virilis* male mitotic cells (D) DNA-FISH image of *D. virilis* female mitotic cells (E) DNA-FISH image of *D. novamexicana* male mitotic cells (F) DNA-FISH image of *D. americana*. Up to three different fluorescent probes were used each time.

281    FISH results in *D. virilis* show that the most abundant satellite determined by sequencing,

282  AAACTAC, is clearly the most abundant and occurs in approximately equal amounts in the

pericentromeric region on the five pairs of large chromosomes. The Y chromosome appears to have

slightly less AAACTAC satellite. The second and third most abundant satellites, AAATTAC and

AAACTAT, are localized more proximally at or near the centromere. There are five single

chromosomes having each of these satellite, indicating that one chromosome pair has different

satellite content - which we hypothesized to be the X and Y.  Based on differences between male

and female FISH results (Figure 2C and 2D), we suggest the Y chromosome has AAACTAT at both

distal ends of the chromosome and AAACTAC only flanking one end, whereas the X chromosome

has the other centromeric repeat AAATTAC. We were also able to visualize the dot chromosomes in

*D. virilis*, which we find is mostly composed of AAACTAT. The AAACAAC satellite is present in small

amounts in *D. virilis*, very likely on a single chromosome (Figure S6).

We estimated that *D. novamexicana* has approximately half the AAACTAC as *D. virilis*, and

visualizing it with FISH reveals a pattern that suggests aspects of its evolution. Its pericentromeric

localization is conserved. One chromosome pair has the same amount of AAACTAC as *D. virilis*,

whereas all other chromosomes have a very small amount (Figure 2E). Based on the FISH images, it

appears that it is the 5th chromosome in *D. novamexicana* that has the greatest amount of

pericentromeric AAACTAC conserved. The centromeric repeat on all major chromosomes is

AAACAAC in *D. novamexicana* and *D. americana.* Our images illustrate clearly the centromere-

centromere fusion between chromosome X-4 and 2-3 in *D. americana* with the satellites being

maintained on both sides of the fusion (Figure 2F)*.* None of the four simple satellite probes bound

to the Y chromosome of *D. novamexicana* or *D. americana*. Based on the images we suggest that *D.*

*americana* has an intermediate amount of pericentromeric AAACTAC satellite compared to *D. virilis*

and *novamexicana*.

**Some satellite-containing reads are linked to TEs**

We used RepeatMasker to detect if any of the reads containing satellites also contain transposable elements. TEs might be located in islands within the simple repeats at the centromere as in *D. melanogaster* (Chang *et al.* 2019) (Figure 3A), in more distal regions flanking the pericentromeric heterochromatin (Figure 3B), or some TEs may have inserted into long pericentromeric satellite arrays (Figure 3C). For AAACTAC (and its artefactual counterpart AAACGAC), ~3.5% of reads (2473/75,364) also contained at least 500 bp of a TE insertion. In the satellite reads that also contain TE sequences, TEs were enriched at the beginning and ends of reads, concordant with the hypothesis that a high proportion of the TEs we found are flanking the long arrays of AAACTAC distally (Figure 3B). In order to understand how many reads would be expected to contain both satellites and TEs if TEs flanked this satellite and were not interspersed, we simulated a situation where a large satellite block flanked a large TE block. This simulation revealed a much smaller amount of reads containing both satellites and TEs (0.06 %). This result suggests that not only are TEs flanking the pericentromeric satellite AAACTAC, but there have likely been TE insertions into the satellite arrays. Likely flanking the proximal end of the pericentromeric satellite are the centromeric satellites AAACTAT or AAATTAC. 144 reads contained both AAACTAC and AAACTAT repeats (0.19% of AAACTAC reads) and 94 reads contained both AAACTAC and AAATTAC repeats (0.12% of AAACTAC reads). Based on our simulations, these proportions of overlapping reads are consistent with our expectation based on our FISH results that the pericentromeric and centromeric satellites have relatively clean boundaries and they directly flank each other.

**Fig 3**. Transposable elements are in close proximity to satellite arrays. (**A**) Satellites near the centromere may be linked to potential islands of retroelements (Chang et al. 2019) and telomeric TEs. (**B**) Heterochromatic TEs flank satellite arrays. Histograms show the start and end position of transposable element in the satellite-rich read. (**C**) Transposable elements may have inserted into the satellite arrays.

Many of the TE insertions into satellites seemed to be very recent. 2080/2473 TE-containing

AAACTAC reads had a TE insertion with less than 15% divergence from the Repbase consensus,

which is the expected error rate of PacBio reads. These insertions included the superfamilies: DNA

elements, LINE/CR1, LINE/I-Jockey, LINE/Penelope, LINE/R1, LTR/Copia, LTR/Gypsy, LTR/Pao, and

Helitrons. We acknowledge however, that there may be a detection bias for insertions that are less

divergent from the consensus. We also remind readers that there were likely fewer satellite reads

sequenced than expected, and that this may have biased these results if satellite-only regions were

sequenced less efficiently than satellite-TE regions. For centromeric satellites AAATTAC and

AAACTAT, the results are more difficult to interpret since these were more strongly under-

represented. However, 300/715 reads of AAACTAT and 49/385 reads AAATTAC contained TEs. Like

the AAACTAC pericentromeric satellite, most TE insertions were low divergence from the Repbase

consensus in the centromere-proximal satellite reads. 288/300 and 46/49 TE containing reads had a

TE insertion > 500 bp with < 15% divergence for AAACTAT and AAATTAC, respectively.

There were differences in the TE composition of reads with different satellites. For the

pericentromeric satellite AAACTAC, Gypsy-10_Dvi was the most enriched, followed by Helitrons

(Helitron-1N1_DVir, Helitron-1_DVir, Helitron-2N1_DVir, Helitron-2_DVir). For the AAACTAT

centromeric satellite, Gypsy-10_Dvi was again the most enriched, followed by Penelope. For the

AAATTAC centromeric satellite, Gypsy-2_DVir was the most enriched followed by Penelope. In both

AAACTAC and AAACTAT reads, CR1-1_DVi was the second or third most abundant TE. Interestingly,

R1 was present in relatively high amounts in AAACTAC. In 110/132 of these R1-AAACTAC reads,

rDNA sequences were not also linked. This suggests that some R1 elements, which are generally

localized to rDNA loci, have jumped into or near satellite arrays. This is concordant with findings

that some R1 elements are located outside rDNA loci in Drosophila (Stage and Eikbush 2009). All

centromeres in *D. virilis* are acrocentric, meaning that the telomeric TEs Het-A and TART

(Casacuberta and Pardue 2003) are likely near the centromere satellites. We found 12 reads linked

to AAATTAC that contained matches to TART. Only two reads linked to any satellite contained a

sequence matching HeT-A. We also used BLAST to detect matches between the genome assembly

(masked from the 7mer satellites) and the 7mer satellite reads. We could not detect any unique

regions of the genome that matched non-satellite sequence on the reads because they had low

quality matches to hundreds of places in the genome each.

**Variation in *D. virilis* group global strains**

*D. virilis* is globally distributed while its sister species are localized to North America, with *D.*

*novamexicana* more restricted than *D. americana.* Patterns of variation in satellites may reveal

potential mechanisms that can be hypothesized to be driving satellite evolution. Additionally, *D.*

*americana* has a polymorphic fusion between the X and 4th chromosomes, so we may be able to

identify differences in satellite composition associated with the fusion. This fusion has been shown

to be currently undergoing meiotic drive, potentially mediated by a larger total centromere or

pericentromere size in the fused strains compared to the non-fused strains (Stewart et al. 2019). On

the other hand, chromosome fusions are often caused by Robertsonian translocations with loss of

some non-essential DNA, which might include pericentromeric satellites (Schubert and Lysak 2011).

We used Illumina sequencing with PCR-free library preparation and k-Seek to estimate the

abundance of 7mer satellites across 12 worldwide strains of *D. virilis*, eight strains of *D. americana*

(including four strains that have the X-4 fusion and four that do not), and five strains of *D.*

*novamexicana* (Table S1). All sequenced strains were male except a female of the *D. virilis* inbred

strain 87 as a comparison. A PCA using only the four most abundant 7mers shows clustering of the

three species, but the separation is much more dramatic in the PCA using the 20 most abundant

simple satellites (Figure S7). Overall, *D. virilis* had the highest AAACTAC satellite content as well as

the highest variation, with *D. americana* intermediate between *D. virilis* and *D. novamexicana*

(Figure 4A). Using different normalization procedures including mapping and GC correction (see

Materials and Methods), produced the same relative ranking of satellite abundances between

species. In all cases, the inbred strain from which the genome sequence was produced had the

lowest abundance of AAACTAC. In the case of *D. virilis*, this difference was very high. This was not

due to a normalization bias as we did mapping-free normalization.

Satellite abundances in *D. virilis* displayed a pattern that appeared to be correlated to the

geographic location from which strains were collected. For the centromeric satellite AAATTAC,

there was a linear decrease in abundance from West to East then South following probable

migration from Beringia (Throckmorton 1982) beginning in China (Figure 4C). For the centromeric

satellite AAACTAT, the pattern was the opposite; a linear increase in abundance from West to East

then South (Figure 4D). We also analyzed sequence variation in the satellite repeats across strains

and species to determine if there were any interesting patterns. On average, the centromeric

satellite arrays were very homogeneous (average above 99% sequence identity in Illumina reads).

However, AAACTAT had a slightly higher sequence identity than AAATTAC (Figure 4 C and D). There

was no pattern in average sequence identity with respect to geography for any of the satellites

(Figure 4 C and D). The pericentromeric satellite AAACTAC has almost identical sequence divergence

across the three species (~98.5%). When comparing between a male and female of the same strain,

the male had lower sequence identity. In *D. americana* and *D. novamexicana*, the AAACAAC satellite

had lower average sequence identity at 97.5%.

**Fig 4**. Variation in satellites across species and strains. (**A**) AAACTAC total abundance across the three species. (**B**) AAACTAC, (**C**) AAACTAC, and (**D**) AAATTAC, abundance across *D. virilis* strains originating from different localities (x axis). The strain Dvir is strain 87, the inbred strain used for genome assemblies.

There were also several other satellites besides the most abundant four that varied

between the *D. virilis* group species. AAACAAT was either absent or in low abundance in *D. virilis*

and *novamexicana*, but present between 100,000 - 250,000 copies in *D. americana,* indicating a

very recent expansion of this satellite. AAAAAC was present in ~100,000 copies in *D. americana* and

*D. novamexicana*, whereas it is almost absent in *D. virilis*. To differentiate Y-specific satellites, we

included a female of strain 87, the reference genome strain, in our Illumina sequencing run in

addition to a male. There were several Y-specific satellites in *D. virilis* found by Wei et al. 2018 and

validated by our data, which varied between species in the group (AATAATAG, AATAGATT, and

ACATAT). All had different patterns of relative abundance between species

(https://github.com/jmf422/D_virilis_satellites/blob/master/Intra_inter_species_sequencing/virilis

_group_intra-inter_species.html).  There was no detectable difference in centromeric or

pericentromeric satellite abundance in *D. americana* strains with vs. without the polymorphic

centromere-centromere fusion. We conclude that molecular events surrounding the fusion did not

produce any changes in satellite abundance (Figure S8).

Amplified repeats of the DINE-1 helitron have been found on the *D. virilis* 5th and Y

chromosomes (Dias *et al.* 2015). We also examined variation in this satellite abundance by counting

reads that mapped to this family of satellites. There was no striking pattern with respect to

geography, however the strain with the highest AAACTAC content had the lowest DINE-1 content,

besides the inbred strain 87, which had lower satellite content all-around (Figure S9). We expect

AAACTAC and DINE-1 to be both in the pericentromeric region of Chr5 in *D. virilis* based on our FISH

results and the previous work. The male contained 2.6x more DINEs than the female of the same

strain, indicating that ~70% of the DINE-1 repeats in the genome are located on the Y chromosome

in *D. virilis*.

**Discussion**

Here, we used the satellite DNA-rich genome of *D. virilis* to highlight three previously

uncharacterized mechanisms for biases that occur in sequencing and analyzing satellite DNA. We

emphasize that comparing satellite DNA amounts between different platforms (e.g. Illumina,

PacBio, Nanopore, and even different versions of each) should be done with caution as each

technology has its own biases. We have found that issues arise when long arrays of simple satellite

DNA are attempted to be sequenced by long-read platforms. In the case of PacBio, systematic

errors in base calling may be introduced when sequencing through long arrays of satellites. This

issue is not specific to our satellites, as a recent study has also found systematic errors and strand

biases in shorter arrays of human satellites in both PacBio and Nanopore reads (Mitsuhashi *et al.*

2019). Circular consensus sequencing (CCS) or "HiFi", a type of sequencing offered by PacBio which

allows an accurate consensus to be produced after multiple rounds of sequencing the same

molecule, may be more appropriate for sequencing analysis of satellite DNA. No systematic errors

in satellite sequences resulted with the new CCS platform after collaboration with PacBio

representatives. In the case of Nanopore, it is possible that a similar satellite-specific base calling

errors exists, or that there is a strand-specific difference in secondary or tertiary structures that

occur in long strands of simple satellite DNA. We caution readers in interpreting simple satellite

DNA from long-read sequencing data and suggest validation with satellites of known sequence and

abundance if available or Illumina reads (without quality filtering). Long read platforms are already

improving their chemistry and software for better satellite characterization. Because long reads are

likely to cross boundaries of different repetitive regions, long read sequencing proved useful in

understanding the length of the satellite arrays and TE insertions into them. Moreover, we

demonstrate that the abundance of satellites in pericentromeric heterochromatin are

underestimated when sequencing flies compared to pure diploid tissue because of polyteny. We

caution readers in performing quality filtering before simple satellite analysis, as satellite containing

reads may be enriched for lower quality scores.

From comparative analysis of satellites, we found the abundant AAACTAC family of satellites

arose in the branch leading to the *virilis* phylad 4.5-11 MYA (Figure 2A). Interestingly, the most

abundant satellite in *D. montana*, 7-11 MY diverged, is AAAC. The AAACTAC and AAAC satellites

were likely derived from a common ancestor satellite (Fig 2A). From both FISH and sequencing

analysis, we found that *D. virilis* has the highest total amount of AAACTAC family satellites, *D.*

*novamexicana* has about half of *D. virilis*, and *D. americana* intermediate between the two species.

*D. lummei* has the lowest relative satellite content, and its only high-abundance simple satellite is

AAACTAC. Unlike the pericentromeric satellite, the centromere-proximal satellite sequence has

turned over between *D. virilis* and *D. americana/novamexicana*. The AAACAAC satellite likely

evolved in the branch leading to the virilis phylad since it is present in three of the four species

studied (Fig 2A). AAACAAC is present in *D. virilis* in relatively low amounts, whereas it became the

centromeric satellite on almost all chromosomes in *D. americana* and *novamexicana*. The AAACTAT

and AAATTAC satellites are unique to *D. virilis* and occupy the centromeric region. The emerging

pattern is that the centromere-proximal satellites have turned over more rapidly than the

pericentromeric satellite. This is likely due to satellites participating in conflicts at centromeres

(Bayes and Malik 2006, and discussed below). Although sequencing quantified only up to 30 Mb of

the AAACTAC family of satellites, FISH confirmed that these satellites are extremely abundant in *D.*

*virilis* and the 40% of the genome estimate seems realistic.

We can make hypotheses about how and why the satellites expanded in *D. virilis*. We know

that mutation rates for changes in copy number of satellite DNA are high, and potentially have a

tendency to expand rather than contract in the absence of selection (Flynn *et al.* 2017, 2018). High

rates of mutation must be accompanied by a regime that would allow a satellite copy number

increase to sweep the population - which could be mediated by positive selection if there is a

benefit of the satellite increase, or centromere drive if the phenomenon is at play. Alternatively, in

a situation where satellites are slightly deleterious, small effective population sizes in isolated

populations or continued bottlenecks could allow satellites to expand in the genome without being

removed by selection. However, *D. novamexicana* has the lowest effective population size of the

virilis phylad and yet it has the lowest amount of satellite DNA. We already know that the

centromere-to-centromere fusions in *D. americana* have undergone meiotic drive hypothesized to

be mediated by the increase in centromere total size with the fusion (Stewart et al. 2019). The

mechanism allowing drive in *D. americana* may have been at play in the branch leading to *D. virilis*

or may be currently occurring. Why have satellites not expanded to this extent in the other species?

*D. virilis* might have some attributes about its biology that made the satellite expansion favorable or

allowable. For example, genome size is positively correlated with development time in

Drosophilidae (Gregory and Johnston 2008). *D. virilis* has a slow development time, and this may

have evolved in concert with the expansion in satellite abundance in its genome.

We can use data from multiple strains to make hypotheses about factors driving satellite

DNA evolution in *D. virilis*. Ancestrally, *D. virilis* had a relatively small effective population size in an

isolated range in Asia, and has undergone a recent population and range expansion (Mirol et al.

2008). The amount of the most abundant pericentromeric satellite AAACTAC, does not show a

geographical pattern across the global strains. Assuming we sequenced a strain from the ancestral

range, this suggests that population bottlenecks were not what allowed AAACTAC to expand, and

the satellite expansion likely occurred before the population expansion.

Our observation of rapid evolution and enrichment of AAACTAC in *D. virilis* in a short

evolutionary time period (a few million years) is consistent with the centromere-drive model to

account for the evolution of centromere complexity in genetic conflict (Malik and Bayes, 2006). In

this model, the asymmetric female meiosis can cause competition between the centromeres with

or without newly formed satellites or with more or less satellites, to be included into the oocyte to

pass to next generation. A consequence of the competition would be runway expansions of

centromeric satellites, and rapid replacements by novel satellites. We hypothesize that the pattern

of the centromere-proximal satellite AAACTAT increasing on a geographical gradient while AAATTAC

decreases along the same gradient is driven by centromeric conflicts. AAACTAT may be starting to

occupy centromeres that AAATTAC occupied, benefitting from a transmission advantage

(centromere drive), while the AAATTAC satellite may be decreasing in parallel because of selection

"pushing back", for example because of a maximum limit on satellite amount in the centromeric

region. Another line of evidence that centromere related conflicts are playing a role is the rapid rate

of turnover of the centromere-proximal satellites compared to the pericentromeric satellite.

Interestingly, in *D. novamexicana*, AAACTAC was greatly reduced in the pericentromeric

regions on all chromosome pairs except one. Based on the FISH images in *D. novamexicana* and

*americana*, we hypothesize that it is the 5th chromosome that has the high amount of AAACTAC

satellite. This is interesting because previous work has shown that the 5th chromosome contains a

high amount of DINE-1 helitron satellite in *D. virilis* but not in *D. americana* (Dias et al. 2015). This

may be evidence of past and ongoing competition and trade-offs between the DINE-1 satellite and

AAACTAC.  We found that all chromosomes including Chr5 contain a large amount of AAACTAC in *D.*

*virilis*. DINE-1 had a relatively consistent amount across different *D. virilis* strains, however the

strain with the highest AAACTAC amount is an outlier with a lower DINE-1 amount (Figure S9). This

may indicate a maximum threshold of satellites was reached on this chromosome, and one satellite

had to reduce its abundance. We have seen evidence for this trade-off, or appearance of

competitive exclusion, being invoked under selection in our previous studies (Flynn *et al.* 2017,

2018). There may have been a similar conflict on Chr5 of *D. novamexicana*, where AAACTAC

retained a high copy number to prevent DINE-1 from expanding. Interestingly, the opposite has

occurred on the *D. novamexicana* and *D. americana* Y chromosome, where AAACTAC family

satellites are absent but DINE repeats are abundant. A potential mechanism mediating apparent

stabilizing selection on total satellite abundance is that satellites can act as a sink for

heterochromatin factors, with their abundance affecting chromatin state (Lemos *et al*. 2010).

The AAACTAC satellite has remained conserved in sequence and location in the virilis

phylad. It has also maintained high levels of sequence identity that is equal in the three species we

sequenced (98.5% based on Illumina reads). The conservation may reflect a constraint due to

selection or a pervasive mechanism of concerted evolution. The periodicity of the sequence may

stabilize the DNA helix wrapping around nucleosomes, or it may be constrained by coevolution of

an important satellite DNA binding protein (Maio *et al.* 1977; Jagannathan *et al.* 2018). Additionally,

within the AAACTAC family, the position and identity of the four A-nucleotides are conserved in all

four satellites (A̲A̲A̲CTA̲C, A̲A̲A̲TTA̲C, A̲A̲A̲CTA̲T, A̲A̲A̲CA̲A̲C) - which may indicate constraint based on

the above mechanisms. Conservation of particular satellite unit lengths and "AA" periodicities have

been found in other divergent species (Lowman and Bina 1990). Concerted evolution of satellites

could be achieved by repeated recycling of units by copy number changes associated with

replication slippage or unequal recombination or gene conversion (Walsh 1987; Elder and Turner

1995). However, recombination in the pericentromeric heterochromatin has never been detected

in wild-type flies (Mehrotra and McKim 2006; Hughes *et al.* 2018). On the other hand, if

recombination were occurring, satellite arrays will eventually be lost unless they are conserved by

selection (Charlesworth *et al.* 1986). Clearly, we are still lacking in understanding how and why long

simple satellite arrays maintain their homogeneity, and whether recombination plays a role in their

dynamics. Concordant with the hypothesis that recombination is playing a role, males have lower

average sequence identity in the 7 bp satellites than females, which could indicate increased decay

on the Y chromosome where there no homologous recombination (Figure 4A).

Moreover, our results suggest that transposable elements flank the AAACTAC satellite array

(likely more distally to the centromere) and some TEs have inserted within the array. Our analysis

suggests that most TE insertions are recent, but because of the under-representation of satellite

containing reads, we cannot estimate the number of TE insertions that have occurred into the

satellite arrays. Our analysis is also not precise enough to determine if there are islands of TEs at

the centromere in *D. virilis* as has been demonstrated in *D. melanogaster* (Chang et al. 2019).

We found no difference in centromeric and pericentromeric satellites abundances between

*D. americana* strains that differ in their X-4 fusion status. This suggests that the fusion event did not

result in a large loss of satellites, making the total centromere and pericentromere size is indeed

larger on the X-4 fused chromosome than the single unfused chromosome, concordant with the

hypothesis that a larger centromeric region results in centromere drive (Stewart et al. 2019).

Another interesting observation from the sequencing of multiple strains of the three species was

that in all cases, the inbred strain that the reference genome was made from had the lowest

amount of AAACTAC. For *D. virilis*, this difference was extreme. It is tempting to speculate that the

process of inbreeding and/or long periods in the lab may have driven the reduction in

pericentromeric satellite abundance.

In conclusion, our results show very rapid dynamics in the abundant satellites of the *D. virilis*

group that are likely explained by various cellular and population-level forces that are not yet

understood. Further studies can test if there is a species-specific upper limit to satellite amount per

genome or per chromosome upon which negative fitness effects occur, which may result in trade-

offs or competition between satellites. Centromere drive may be an important process affecting

satellite evolution in this species group, and might partially explain why the satellites expanded 4.5-

11 MYA, why satellite sequences at the centromere turned over more rapidly, and why there is a

gradient of increasing satellite content related to geographical distribution of strains. A more

extensive study to determine if inbreeding or extended periods in the lab drives a reduction in

satellite abundance will help illuminate the processes that are important for maintaining satellite

content. Determining the frequency of recombination in the large pericentromeric heterochromatin

blocks in species like *D. virilis* will be challenging but important for understanding how the satellites

maintain homogeneity in their sequence. To understand the role of satellites and the importance of

their sequence, unit length, and abundance, researchers can strive to develop methods to engineer

satellites by modifying specific bases and their abundances.

**MATERIALS AND METHODS**

All scripts for analyzing the data and to produce the results we show are here:

https://github.com/jmf422/D_virilis_satellites. Illumina sequencing reads generated for this study

are deposited in NCBI SRA under accession PRJNA548201. Raw PacBio reads will be deposited under

the same accession. Both will be released upon publication.

**Characterizing satellite DNA from genome assemblies**

All scripts and R markdown files used for this analysis are provided in

https://github.com/jmf422/D_virilis_satellites/tree/master/Genome_assembly_analysis.

We used genome assemblies produced by the PacBio sequencing project

(https://www.ncbi.nlm.nih.gov/bioproject/?term=txid7214[Organism:noexp]) of *D. virilis*, *D.*

*novamexicana*, and *D. americana*. We also downloaded the *D. virilis* genome produced by

Nanopore sequencing from (Miller *et al.* 2018), and the CAF1 assembly from (Drosophila 12

Genomes Consortium *et al.* 2007). We used Phobos (https://www.ruhr-uni-

bochum.de/spezzoo/cm/cm_phobos.htm) and Tandem repeats finder (Benson 1999) to

characterize simple and complex satellites in these genome assemblies. To identify the

chromosomal linkage of complex satellites in the genome assembly, we produced a dotplot with D-

GENIES (Cabanettes and Klopp 2018).

**Characterizing satellite DNA from raw long reads**

Characterizing and quantifying satellites from long reads is a challenge because of the sequencing

high error rate. We used two approaches to characterize satellites from raw long reads. The first

approach, we call k-Seek + Phobos, in which we first broke the reads into 100 bp subreads and ran

k-Seek on them. k-Seek is very efficient for analyzing many reads, however is not very sensitive for

reads with a high error rate since it was designed for Illumina reads (Wei *et al.* 2014). If k-Seek

found satellites on at least one subread, we would run the complete parent read through Phobos.

Phobos is more sensitive to imperfect repeats and error rates, but cannot handle huge quantities of

data; thus why we only ran the portion of reads identified by k-Seek to have tandem repeats. This

approach allowed us to characterize satellites *de novo* and quantify them. All scripts for the analysis

of long reads with the k-Seek + Phobos approach are located here:

https://github.com/jmf422/D_virilis_satellites/tree/master/LongRead_kseek_Phobos.  The second

approach we used is Noise-Cancelling Repeat Finder (NCRF, (Harris *et al.* 2019) ). This program was

designed to quantify satellites from long reads with high error rates. However, it cannot identify

satellites *de novo* and requires specific satellite sequences to search for. NCRF also requires a "max

divergence allowed" parameter, which we tuned with simulations (see below). Scripts used for the

NCRF approach are located here:

https://github.com/jmf422/D_virilis_satellites/tree/master/LongRead_NCRF.

We did simulations to assess both approaches:

https://github.com/jmf422/D_virilis_satellites/tree/master/Simulations. First, we created a

simplified mock *D. virilis* genome with a satellite DNA composition estimated from our FISH results.

We could not use the genome assembly because it contained very little satellite DNA. Specifically,

each chromosome had a centromeric satellite either AAATTAC or AAACTAC followed by the

pericentromeric satellite AAACTAC, combined taking up 40% of the genome. The non-satellite DNA

portion of the genome was generated randomly with a 40% GC content. We then used PBSim (Ono

*et al.* 2013) to simulate PacBio reads and we used these simulated reads for multiple analyses. First,

we used them to tune the max divergence parameter of NCRF by running NCRF repeatedly with

max divergence parameters ranging 18-30%. We found that the amount of satellites found,

particularly the most abundant one, levelled off at 25% max divergence. This is the parameter value

we used moving forward. We also used these simulated reads to quantify satellites with both

approaches and compare them. Finally, we used these simulated reads to assess strand biases in

long read sequencing data (see below).

**Identification of biases in simple satellites in long read data**

We suspected that there were biases in the satellite DNA found in the *D. virilis* group PacBio (and

Nanopore) data because we found high abundance satellites that had never been found before with

other types of data, and so we suspected they were artifactual. These artifactual satellites were

found with both kSeek + Phobos and NCRF approaches, but were not found in the simulated data.

We tried testing for a strand bias in reads that contained satellite DNA. Using both the summarized

output from NCRF and validated with a custom script

(LongRead_NCRF/which_strand_pacbio_script.sh), we counted the satellite DNA stretches that

originated from each the positive and negative strand. The positive strand is defined as the one that

contains the satellite AAACTAC and derivatives (more As than Ts), and the negative strand is the

one that contains the reverse complement (e.g. GTAGTTT, more Ts than As). We did this for the

three satellites used in the simulated data and real and artefactual satellites found in the PacBio

and Nanopore data. Detailed analysis and visualization of the biases is shown here:

LongRead_kseek_Phobos/longread_analysis.html

**Sequencing of polytene and non-polytene tissue**

To acquire *D. virilis* pure diploid tissue, we dissected male 3$^{rd}$ instar larvae and collected imaginal

discs including the eye-antennal disc and wing discs. Approximately 100 larvae were required to get

enough DNA (>1 ug). We also collected ~5 adult flies for fly libraries. We used the inbred genome

assembly strain 87 for these libraries. DNA was extracted with Qiagen DNeasy blood and tissue kit

and PCR-free libraries were prepared. Libraries were run on an Illumina NextSeq with 1 x 150 bp

reads, and each sample took up approximately 7% of the flowcell. The other libraries run on this

flowcell were from an unrelated project including RNAseq from other species. Reads were analyzed

with k-Seek both before and after filtering with Trimmomatic (Bolger et al. 2014). FastQC was run to

evaluate the quality of the reads. Scripts are here:

https://github.com/jmf422/D_virilis_satellites/tree/master/Polyteny. We also analyzed publicly

available *D. melanogaster* data from the same strain and same sequencing platform of embryos

(non-polytene), salivary glands (extreme polyteny) from (Yarosh and Spradling 2014), and flies

(varied levels of polyteny) from (Gutzwiller *et al.* 2015).

**Fluorescence in situ hybridization of satellite DNAs**

We followed the protocol of (Larracuente and Ferree 2015) for satellite DNA FISH. We ordered the

following probes from IDT with 5' modifications: $(AAACTAC)_6$ with alexa-488 fluorophore,

$(AAACTAT)_6$ with Cyanine5 fluorophore, $(AAATTAC)_6$ with Cyanine3 fluorophore, $(AAACAAC)_6$ with

Cyanine3 fluorophore, and $(AAACGAC)_6$ with Cyanine5 fluorophore. We hybridized three probes at

a time, to allow for similar probes to compete to result in specific hybridization with the rationale

shown in (Beliveau *et al.* 2015). Hybridization temperature was 32°C. We imaged on an Olympus

fluorescent microscope and Metamorph capture system at the Cornell Imaging Facility. Composite

images were produced with ImageJ.

**Characterizing TEs linked to satellites and satellites anchored to the genome assembly**

We extracted the reads identified to have the 7 bp satellites on them from NCRF results, and then

we ran RepeatMasker (http://www.repeatmasker.org) on these reads using parameters: "-nolow"

and "-species Drosophila". All reads had at least 500 bp of tandem satellite on them according to

NCRF default parameters, and to avoid spurious identification of TEs from semi-repetitive

fragments, we described only TEs in reads that also had at least 500 bp of a TE identified from

RepeatMasker. We also BLASTed the same satellite reads to the genome assembly to evaluate if

satellite reads could be anchored to the genome assembly. Analysis scripts are here:

https://github.com/jmf422/D_virilis_satellites/tree/master/TEs_satellites.

**Sequencing of multiple *D. virilis* group strains**

We obtained as many strains of *D. virilis* that have information about where they were collected as

possible. This included 12 strains as live stocks we obtained either from stocks in our lab or from the

Drosophila species stock center (Table S1). We also prepared a female library for strain 87 for when

we wanted to differentiate Y-specific satellite patterns. We also obtained five strains of *D.*

*novamexicana* and eight strains of *D. americana*. All were obtained from live stocks and the inbred

genome strains were included for both species as well (strain 14 and G96, respectfully). For *D.*

*americana*, we included four strains that have the chromosome X-4 fusion and four strains that do

not have it, based on communication with the Bryant McAllister lab. DNA was extracted as above

from five flies each and samples were prepared identically as above and sequenced on 50% of 3

flowcells of Illumina NextSeq 1 x 150 bp reads. We dispersed the samples from each species

between multiple flowcells. Our samples took up only half the flowcell with the other half being

occupied by a RNAseq libraries from an unrelated project.

All scripts used to analyze these data are located here:

https://github.com/jmf422/D_virilis_satellites/tree/master/Intra_inter_species_sequencing.

Reads were evaluated with FastQC and not filtered for quality based on the potential bias of

Illumina quality scores on satellites. We used k-Seek to quantify satellites. We tried several

normalization strategies but decided the most appropriate was a mapping-free normalization. We

estimated average depth by dividing the total number of bases sequenced by the estimated

genome size by flow cytometry (Bosco *et al.* 2007). We believe this was the best option in this case

because: 1) we were concerned about a mapping bias because for each species the strain that the

genome assembly was made from may have more reads map to it; 2) after masking the genome

from the 7mer satellites and also excluding the X and Y contigs (because we had male and female

strains, and the Y chromosome contained more low GC regions) - there was little difference in

coverage based on GC content. We include results when we used a mapping based GC

normalization in the sub-directory "AlternativeNormalization".

We used NCRF with modified parameters (minlength=100, maxdiv=10) to characterize the

average sequence identity of satellite arrays from the Illumina data. To quantify DINE-1 satellites

across *D. virilis* strains, we produced a library of DINE-1 satellite variants based on our PacBio

genome analysis. We then mapped Illumina reads to this library and normalized the number of

reads that mapped to any sequence in the library by the estimated depth. We also analyzed

Illumina DNA sequencing reads of *D. montana* (Parker et al. 2018) and *D. lummei* (Ahmed-Braimah

*et al.* 2017) with k-Seek to identify the most abundant satellites and whether or not the AAACTAC

satellite family was present.

**Acknowledgments**

We thank Yasir Ahmed-Braimah for helpful discussions and advice for some analyses. We also thank

Bryant McAllister for providing *D. americana* strains along with their fusion status. We are grateful

for Elissa Cosgrove's help with some computational trouble-shooting and Asha Jain's help in

preparing DNA sequencing libraries. We also thank Danny Miller for useful discussions and for

providing the raw Nanopore reads from his study. Sarah Kingan, Jane Landolin, and Greg Young

from Pacific Biosciences were very helpful in exploring the causes for the artefactual repeats and in

producing the HiFi data. We thank Amanda Larracuente for advice on FISH protocols and the Cornell

Imaging Facility for use of their microscope. This project was funded by NIH grant number:

GM116113 to RAW, ML, and AGC and GM119125 to AGC and Daniel Barbash. JMF was supported

by an NSERC PGS D fellowship.

**References**

Ahmed-Braimah Y. H., R. L. Unckless, Andrew G. Clark, 2017 Evolutionary Dynamics of Male

Reproductive Genes in the *Drosophila virilis* Subgroup. G3. 7:3145-3155.

Beliveau B. J., A. N. Boettiger, M. S. Avendaño, R. Jungmann, R. B. McCole, *et al.*, 2015 Single-

molecule super-resolution imaging of chromosomes and in situ haplotype visualization using

Oligopaint FISH probes. Nat. Commun. 6: 7147.

Belyaeva E. S., I. F. Zhimulev, E. I. Volkova, A. A. Alekseyenko, Y. M. Moshkin, *et al.*, 1998 Su(UR)ES:

a gene suppressing DNA underreplication in intercalary and pericentric heterochromatin of

*Drosophila melanogaster* polytene chromosomes. Proc. Natl. Acad. Sci. U. S. A. 95: 7532–7537.

Benson G., 1999 Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids

Research 27: 573–580.

Bolger A. M., M. Lohse, B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics 30:2114-2120.

Bosco G., P. Campbell, J. T. Leiva-Neto, and T. A. Markow, 2007 Analysis of Drosophila species

genome size and satellite DNA content reveals significant differences among strains as well as

between species. Genetics 177: 1277–1290.

Cabanettes F., and C. Klopp, 2018 D-GENIES: dot plot large genomes in an interactive, efficient and

simple way. PeerJ 6: e4958.

Caletka B. C., and B. F. McAllister, 2004 A genealogical view of chromosomal evolution and species

delimitation in the *Drosophila virilis* species subgroup. Mol. Phylogenet. Evol. 33: 664–670.

Casacuberta E., and M.-L. Pardue, 2003 HeT-A elements in *Drosophila virilis*: retrotransposon

telomeres are conserved across the Drosophila genus. Proc. Natl. Acad. Sci. U. S. A. 100:

14091–14096.

Chang C.-H., and A. M. Larracuente, 2019 Heterochromatin-Enriched Assemblies Reveal the

Sequence and Organization of the *Drosophila melanogaster* Y Chromosome. Genetics 211:

333–348.

Chang C.-H., A. Chavan, J. Palladino, X. Wei, N. M. C. Martins, *et al.*, 2019  Islands of retroelements

are the major components of Drosophila centromeres. PLoS Biology 17(5):e3000241.

Charlesworth B., C. H. Langley, and W. Stephan, 1986 The evolution of restricted recombination and

the accumulation of repeated DNA sequences. Genetics 112: 947–962.

Charlesworth B., P. Sniegowski, and W. Stephan, 1994 The evolutionary dynamics of repetitive DNA

in eukaryotes. Nature 371: 215–220.

Dias G. B., P. Heringer, M. Svartman, and G. C. S. Kuhn, 2015 Helitrons shaping the genomic

architecture of Drosophila: enrichment of DINE-TR1 in α- and β-heterochromatin, satellite DNA

emergence, and piRNA expression. Chromosome Res. 23: 597–613.

Drosophila 12 Genomes Consortium, A. G. Clark, M. B. Eisen, D. R. Smith, C. M. Bergman, *et al.*,

2007 Evolution of genes and genomes on the Drosophila phylogeny. Nature 450: 203–218.

Elder J. F. Jr, and B. J. Turner, 1995 Concerted evolution of repetitive DNA sequences in eukaryotes.

Q. Rev. Biol. 70: 297–320.

Elliott T. A., and T. Ryan Gregory, 2015 What's in a genome? The C-value enigma and the evolution

of eukaryotic genome content. Philos. Trans. R. Soc. Lond. B Biol. Sci. 370: 20140331.

Flynn J. M., I. Caldas, M. E. Cristescu, and A. G. Clark, 2017 Selection Constrains High Rates of

Tandem Repetitive DNA Mutation in *Daphnia pulex*. Genetics genetics.300146.2017.

Flynn J. M., S. E. Lower, D. A. Barbash, and A. G. Clark, 2018 Rates and Patterns of Mutation in

Tandem Repetitive DNA in Six Independent Lineages of *Chlamydomonas reinhardtii*. Genome

Biol. Evol. 10: 1673–1686.

Gall J., E. Cohen, and M. Polan, 1971 Repetitive DNA sequences in Drosophila. Chromosoma 33.

https://doi.org/10.1007/bf00284948

Gall J. G., and D. D. Atherton, 1974 Satellite DNA sequences in Drosophila virilis. J. Mol. Biol. 85:

633–664.

Gregory T. R., 2001 Coincidence, coevolution, or causation? DNA content, cell size, and the C-value

enigma. Biol. Rev. Camb. Philos. Soc. 76: 65–101.

Gregory T. R., and J. S. Johnston, 2008 Genome size diversity in the family Drosophilidae. Heredity

101: 228–238.

Gutzwiller F., C. R. Carmo, D. E. Miller, D. W. Rice, I. L. G. Newton, *et al.*, 2015 Dynamics of

*Wolbachia pipientis* Gene Expression Across the Drosophila melanogaster Life Cycle. G3 5:

2843–2856.

Harris R. S., M. Cechova, K.D. Makova. 2019 Noise-cancelling repeat finder: uncovering tandem

repeats in error-prone long-read sequencing data. Bioinformatics: btz484

Heikkinen E., V. Launonen, E. Muller, L. Bachmann 1995 The pvB370 *BamHI* Satellite DNA Family of

the *Drosophila virilis* Group and Its Evolutionary Relation to Mobile Dispersed Genetic pDv

Elements. J Mol Evol 41:604-614.

Henikoff S., K. Ahmad, H.S. Malik. 2001 The Centromere Paradox: Stable Inheritance with Rapidly

Evolving DNA. Science 293: 1098–1102.

Hughes S. E., D. E. Miller, A. L. Miller, and R. S. Hawley, 2018 Female Meiosis: Synapsis,

Recombination, and Segregation in. Genetics 208: 875–908.

Izumitani H. F., Y. Kusaka, S. Koshikawa, M. J. Toda, and T. Katoh, 2016 Phylogeography of the

Subgenus Drosophila (Diptera: Drosophilidae): Evolutionary History of Faunal Divergence

between the Old and the New Worlds. PLoS One 11: e0160051.

Jagannathan M., R. Cummings, and Y. M. Yamashita, 2018 A conserved function for pericentromeric

satellite DNA. Elife 7. https://doi.org/10.7554/eLife.34122

Jagannathan M., R. Cummings, and Y. M. Yamashita, 2019 The modular mechanism of

chromocenter formation in. Elife 8. https://doi.org/10.7554/eLife.43938

Kim J. C., J. Nordman, F. Xie, H. Kashevsky, T. Eng, *et al.*, 2011 Integrative analysis of gene

amplification in Drosophila follicle cells: parameters of origin activation and repression. Genes

Dev. 25: 1384–1398.

Larracuente A. M., and P. M. Ferree, 2015 Simple Method for Fluorescence DNA In Situ

Hybridization to Squashed Chromosomes. Journal of Visualized Experiments. 95: e52288.

Lemos B., A. T. Branco, D. L. Hartl, 2010 Epigenetic effects of polymorphic Y chromosomes modulate

chromatin components, immune response, and sexual conflict. PNAS 107:15826-15831.

Lima L. G. de, M. Svartman, and G. C. S. Kuhn, 2017 Dissecting the Satellite DNA Landscape in Three

Cactophilic Sequenced Genomes. G3 7: 2831–2843.

Lowman H., and M. Bina, 1990 Correlation between dinucleotide periodicities and nucleosome

positioning on mouse satellite DNA. Biopolymers 30: 861–876.

Malik H.S.  and J.J. Bayes, 2006. Genetic conflicts during meiosis and the evolutionary origins of

centromere complexity.  Biochem Soc Trans.  34: 569-573.

Maio J. J., F. L. Brown, and P. R. Musich, 1977 Subunit structure of chromatin and the organization

of eukaryotic highly repetitive DNA: recurrent periodicities and models for the evolutionary

origins of repetitive DNA. J. Mol. Biol. 117: 637–655.

Mehrotra S., and K. S. McKim, 2006 Temporal Analysis of Meiotic DNA Double-Strand Break

Formation and Repair in Drosophila Females. PLoS Genetics 2: e200.

Miller D. E., C. Staber, J. Zeitlinger, and R. Scott Hawley, 2018 Highly Contiguous Genome

Assemblies of 15 Drosophila Species Generated Using Nanopore Sequencing. G3:

Genes|Genomes|Genetics 8: 3131–3141.

Mills W. K., Y. C. G. Lee, A. M. Kochendoerfer, E. M. Dunleavy, G. H. Karpen. 2019 RNA transcribed

from heterochromatic simple-tandem repeats are required for male fertility and histone-

protamine exchange in *D. melanogaster*. Biorxiv doi: https://doi.org/10.1101/617175.

Mitsuhashi S., M. C. Frith, T. Mizuguchi, S. Miyatake, T. Toyota, *et al.*, 2019 Tandem-genotypes:

Robust detection of tandem repeat expansions from long DNA reads. Genome Biology 20:58.

Ohno S., 1972 So much "junk" DNA in our genome. Brookhaven Symp. Biol. 23: 366–370.

Ono Y., K. Asai, and M. Hamada, 2013 PBSIM: PacBio reads simulator—toward accurate genome

assembly. Bioinformatics 29: 119–121.

Ostrega M. S., and V. Thompson, 1986 Mitochondrial DNA Restriction site polymorphism in

*Drosophila montana* and *Drosophila virilis*. Biochemical Systematics and Ecology 14: 515–519.

Parker D. J., R. A. W. Wiberg, U. Trivedi, V. I. Tyukmaeva, K. Gharbi, *et al.*, 2018 Inter and

Intraspecific Genomic Divergence in *Drosophila montana* Shows Evidence for Cold Adaptation.

Genome Biol. Evol. 10: 2086–2101.

Pavlek M., Y. Gelfand, M. Plohl, and N. Meštrović, 2015 Genome-wide analysis of tandem repeats in

*Tribolium castaneum* genome reveals abundant and highly dynamic tandem repeat families

with satellite DNA features in euchromatic chromosomal arms. DNA Res. 22: 387–401.

Schubert I., and M. A. Lysak, 2011 Interpretation of karyotype evolution should consider

chromosome structural constraints. Trends Genet. 27: 207–216.

Smith A. V., and T. L. Orr-Weaver, 1991 The regulation of the cell cycle during Drosophila

embryogenesis: the transition to polyteny. Development 112: 997–1008.

Spicer G. S., and C. D. Bell, 2002 Molecular Phylogeny of the *Drosophila virilis* Species Group

(Diptera: Drosophilidae) Inferred from Mitochondrial 12S and 16S Ribosomal RNA Genes.

Annals of the Entomological Society of America 95: 156–161.

Stage D. E., T. H. Eickbush, 2009 Origin of nascent lineages and the mechanisms used to prime

second-strand DNA synthesis in the R1 and R2 retrotransposons of Drosophila. Genome

Biology 10:R49.

Stewart B. S., Y. H. Ahmed-Braimah, D. G. Cerne, B. F. McAllister, Female meiotic drive preferentially

segregates derived metacentric chromosomes in Drosophila.

Throckmorton LH: *The Genetics and Biology of Drosophila Volume 3b*. Academic Press: New York;

1982.

Walsh J. B., 1987 Persistence of tandem arrays: implications for satellite and simple-sequence

DNAs. Genetics 115: 553–567.

Wei K. H.-C., J. K. Grenier, D. A. Barbash, and A. G. Clark, 2014 Correlated variation and population

differentiation in satellite DNA abundance among lines of *Drosophila melanogaster*. Proc. Natl.

Acad. Sci. U. S. A. 111: 18793–18798.

Wei K. H.-C., S. E. Lower, I. V. Caldas, T. J. S. Sless, D. A. Barbash, *et al.*, 2018 Variable Rates of

Simple Satellite Gains across the Drosophila Phylogeny. Mol. Biol. Evol. 35: 925–941.

Yarosh W., and A. C. Spradling, 2014 Incomplete replication generates somatic DNA alterations

within Drosophila polytene salivary gland cells. Genes & Development 28: 1840–1855.

Zelentsova E. S., R. P. Vashakidze, A. S Krayev, M. B. Evgen'ev 1986 Dispersed repeats in *Drosophila*

*virilis*: Elements mobilized by interspecific hybridization. Chromosoma 93: 469-476.