Evolutionary dynamics of abundant 7 bp satellites in the genome of *Drosophila virilis*

Jullien M. Flynn[1], Manyuan Long[2], Rod A. Wing[3], Andrew G. Clark[1]

[1]Department of Molecular Biology and Genetics, Cornell University, Ithaca, USA

[2]Department of Ecology and Evolution, University of Chicago, Chicago, USA

[3]Arizona Genomics Institute, School of Plant Sciences, University of Arizona, Tucson, Arizona,

USA

Abstract

The factors that drive the rapid changes in satellite DNA genomic composition we see in

eukaryotes are not well understood. *Drosophila virilis* has one of the highest relative amounts

of simple satellites of any organism that has been studied, with an estimated 40+% of its

genome composed of a few related 7 bp satellites. Here we use *D. virilis* as a model to

understand technical biases affecting satellite sequencing and the evolutionary processes that

drive satellite composition. By analyzing sequencing data from Illumina, PacBio, and Nanopore

platforms, we identify platform-specific biases and suggest best practices for accurate

characterization of satellites by sequencing. We use comparative genomics and cytogenetics to

demonstrate that the satellite family arose in the branch leading to the virilis phylad 4.5 - 11

million years ago. The most abundant satellite is conserved in sequence and location in the

pericentromeric region but has diverged widely in abundance among species, whereas the

satellites nearest the centromere are rapidly turning over in sequence composition. The

amount of satellite DNA is lowest in the species with the smallest effective population size in

the *D. virilis* phylad, contrary to what would be expected if satellites were only evolving under

weak negative selection. We found evidence for transposable elements flanking the satellite

arrays as well as recent insertions into the arrays. By analyzing multiple strains of *D. virilis*, we

saw that one centromere-proximal satellite is undergoing an expansion, while the other is

contracting in a correlated manner, suggesting possible trade-offs or competition between

satellites. In conclusion, we illuminate several key attributes of satellite evolutionary dynamics

that we hypothesize to be driven by processes like positive selection, meiotic drive, and

constraints on satellite sequence and abundance.

Author summary

Tandem arrays of sequence repeats, or satellite DNA, makes up large and variable proportions

of eukaryote genomes. Satellites may be an important component of genomes, but their

evolution is not well understood because they are subject to a variety of different evolutionary

forces due to their repetitive nature and proximity to centromeres, and they are difficult to

sequence. Here we use a fruit fly *Drosophila virilis* – whose genome experienced a recent

explosion in satellite DNA to make up 40+% of its genome – to understand the evolutionary

processes that drive satellite composition. We found that the satellite emerged and expanded

rapidly, and is variable among related species. The satellites have different patterns of

evolution depending on their proximity to the centromere. We show evidence that the

satellites may act selfishly and result in trade-offs on other satellites due to the push-back of

selection.

Introduction

Repetitive DNA is abundant in most eukaryotic genomes, and is now understood to be

correlated with the manifold variation in genome size across the tree of life (Elliott and Ryan

Gregory 2015). For most species, transposable elements (TEs) dominate the repeat landscape,

including in humans, plants, and *Drosophila melanogaster*. Satellite DNA, which is characterized

by tandem repeats spanning long arrays, very rarely has dominated a genome to a similar

extent as TEs. An unprecedented case is that of *Drosophila virilis*, the Drosophila species with

the largest estimated genome size (up to 389 Mb) (Bosco *et al.* 2007), where some 40% of the

genome is comprised of just three simple 7-mer satellites: AAACTAC, AAACTAT, and AAATTAC

(Gall *et al.* 1971; Gall and Atherton 1974). Since the 1970s, there has been no follow-up to

validate the amount of 7-mers with modern techniques, or evolutionary studies to understand

how and why these satellite repeats expanded so explosively. The genomic composition of

simple satellites in *D. virilis* provides an excellent model for an investigation of the evolutionary

dynamics involved in their expansion in the genome as well as the technical challenges facing

simple satellite analysis.

Satellites are rapidly evolving in sequence and copy number, and there is a high level of

variation in satellite content among and within species (Wei *et al.* 2014, 2018). The reasons for

such dramatic variation is not well understood, and cannot be fully explained by current

models. Satellites have been long hypothesized to be slightly deleterious and therefore

governed primarily by the strength of negative selection (Ohno 1972). However, the amount of

satellite in the genome that causes negative effects that could be selected against depends on

many factors and cannot be easily predicted (Charlesworth *et al.* 1994; Gregory 2001). The fact

that most organisms have satellite repeats in or near centromeres suggests that they are

important for centromere function. Satellite repeats can also be important for maintenance of

the chromocenter and packaging of chromosomes in the nucleus (Jagannathan *et al.* 2018,

2019), and the transcripts of some satellites may be essential for fertility (Mills *et al*. 2019). In

heterozygotes with alleles that differ in pericentromeric satellite sequence or abundance, one

allele may assemble a stronger kinetochore during female meiosis I, increasing its probability of

transmission into the egg (rather than polar bodies). This transmission advantage, known as

centromere drive, allows satellites to rapidly change in composition in the population,

regardless of their whole-organism fitness effects (Henikoff 2001). If satellite DNA is an

essential component of genomes or is only a burden (i.e. is selfish), it is still not clear why some

species have almost no pericentromeric satellite DNA while others, like *D. virilis*, possess

pericentromeric satellites that make up almost half of the genome.

Comparing the satellites of *D. virilis* to those of its sister species can elucidate when the

abundant satellites arose, and how rapidly their copy numbers and sequences evolved. *D. virilis*

is 4.5 MY diverged from its sister species *D. novamexicana* and *D. americana*, which are both

restricted to North America, unlike globally-distributed *D. virilis* (Caletka and McAllister 2004).

D. novamexicana and *D. americana* have a smaller estimated genome size than *D. virilis* (~250

Mb vs. 389 Mb), suggesting these species may have less satellite content (Bosco *et al.* 2007).

Additionally, using intra-species comparisons across global populations can give indications

about factors that may be influencing satellite dynamics. For example, in *D. melanogaster*,

patterns of abundance of the *Prodsat* satellite closely mirror the migration patterns of species,

suggesting an ongoing expansion of this satellite (Wei *et al.* 2014). Genetic drift or meiotic drive

may contribute to patterns of geographical gradients of satellite abundance. We can also use

intra-species data to pose hypotheses about non-neutral processes that may be driving satellite

content. Previous work has shown evidence for conflicts or trade-offs between satellites within

the genome, and these constraints can be illuminated by analyzing satellites in several strains

(Flynn *et al.* 2017, 2018).

Genome-wide characterization of satellites has taken off since high-throughput

sequencing has become widely available. We have learned from several informative studies

about the sequences and relative abundances of satellites in various species (Pavlek *et al.* 2015;

Flynn *et al.* 2017; de Lima *et al.* 2017; Wei *et al.* 2018), but technical challenges may be

preventing accurate quantitative estimates in some cases. Satellites may be more prone to

errors or biases in the sequencing process that do not affect the better studied regions of the

genome. Satellites are difficult to assemble even with long-read sequencing (Chang and

Larracuente 2019). The genome assembly of *D. virilis* is approximately half its estimated

genome size by flow cytometry (~200 Mb vs 389 Mb) (Bosco *et al.* 2007), and it is likely that

much of what is missing is simple satellite DNA. However, even using alignment-free raw read

methods have not produced satellite DNA estimates that approach the amount that is missing

from the genome assembly and was estimated from early work (Gall *et al.* 1971; Gall and

Atherton 1974; Wei *et al.* 2018). Now, as long read sequencing is also being exploited to study

satellites, we must evaluate satellite DNA abundance estimates to assess if there are platform-

specific biases that may affect evolutionary analysis of satellite DNA.

The purpose of this paper is two-fold; first to explore the technical biases preventing

accurate characterization and quantification of simple satellites, and second to use a

comparative approach to understand the evolutionary dynamics of the extremely abundant

7mers in the *D. virilis* group. First, we characterize satellites in *D. virilis* sequencing data from

different platforms and assess biases that affect accurate satellite characterization. We then

use comparative genomics and cytogenetics in *D. virilis* and its sister species to understand the

composition and changes in the highly abundant simple satellites. Finally we sequence multiple

strains of *D. virilis* and sister species to estimate polymorphism in satellite abundance and infer

processes that may be influencing their evolution. From this we infer that there are likely a

variety of understudied processes affecting satellite DNA in this organism, including positive

selection, meiotic drive, and constraints and trade-offs between satellites.

RESULTS

Technical biases in characterizing simple satellites from sequencing

Long-read genome assemblies have an under-representation of simple satellites

Long-read sequencing technologies have an advantage because of their long reads, but a

disadvantage due to their high error rate, prompting a need for extensive alignments for error-

correction and assembly. First we asked whether assemblies from long read technologies can

better assemble simple satellite reads than the previous Sanger assembly. We compared the

amount of simple 7-mer satellites (AAACTAC, AAACTAT, AAATTAC, AAACAAC) in three *D. virilis*

genome assemblies: the CAF1 assembly produced from Sanger sequencing (Drosophila 12

Genomes Consortium *et al.* 2007), a PacBio assembly produced by our group by ~100x coverage

(available at https://www.ncbi.nlm.nih.gov/bioproject/?term=txid7214[Organism:noexp]), and a

Nanopore assembly produced from ~20x sequencing coverage (Miller *et al.* 2018). All

assemblies were approximately the same size at ~200 Mb. The PacBio and Nanopore

assemblies contained a similarly low amount of simple 7-mer satellites, 29 and 28 kb,

respectively. The CAF1 assembly, however contained 7.36 Mb of these satellites. This

discrepancy is likely largely due to the difference in assembly algorithms used for short read

and long read data. Long reads must be aligned and corrected to be incorporated into the

assembly because of their high error rate, whereas this is not necessary for Sanger-based

assemblies. Use of modified methods can improve assemblies of repetitive regions (Chang and

Larracuente 2019), but for highly homogeneous simple satellites, whose arrays span 10-100x

longer than the current maximum read length, it is practically impossible to produce a

continuous assembly.

Simulations to assess simple repeat quantification from long read sequencing data

Due to assembly issues of simple satellites, they must be quantified from raw unassembled

reads. Long read sequencing data poses a significant challenge because of the high error rate

including a high indel rate in the raw reads. We therefore used two different approaches along

with simulations to assess their accuracy. The first approach used k-Seek to select repeat-rich

reads and then Phobos to quantify satellites. This approach allows for *de novo* discovery of

satellite sequences. We used Noise-Cancelling Repeat Finder (NCRF, Harris *et al.* 2018) for our

second approach, providing our target satellites. Both methods are relatively sensitive to

imperfect repeats, which we expect with the high error rate of long-read sequencing.

To evaluate our approaches, we created a mock *D. virilis*-like genome containing

pericentromeric and centromeric repeats on each of 5 chromosomes (See Materials and

Methods). We then simulated 10x PacBio reads from this genome, and then quantified

satellites using both approaches. NCRF works by doing alignments of target satellites to the

reads and allowing up to a user-specified maximum divergence. To determine the most

appropriate maximum divergence, we simulated a range of values for this parameter from 18-

30% and chose the lowest asymptotic value - which was 25% in this case (Figure S1). NCRF

found almost the same amount of satellites that truly existed in the mock genome whereas the

k-Seek + Phobos method only found about 20% (Figure 1A).

The amounts and biases in simple 7mer repeats differ between Nanopore and PacBio

sequencing reads

Next, we quantified simple satellites in the long-read data generated from our 100x PacBio

sequencing and 20x Nanopore sequencing using the two approaches mentioned above. Unlike

in the simulations, both approaches produced very similar (but lower than expected) estimates

at 8.8-10.9 Mb for the PacBio data (Figure 1B). The Nanopore data contained almost 3 times

the 7mer satellites compared to PacBio, with 23.4 - 28.2 Mb (Figure 1C). This may represent a

platform-specific difference in the ability to sequence long arrays of simple tandem repeats.

Both the PacBio reads and the Nanopore reads contained a greater amount of simple satellites

than data produced in our lab previously with Illumina HiSeq sequencing (Wei et al. 2018).

Both the PacBio and Nanopore reads contained large amounts of what we expect to be

artifactual repeats, which were found with the k-Seek + Phobos approach, and validated with

NCRF. NCRF found 4.4 Mb (normalized to 1x genome coverage) of AAACGAC in the PacBio

reads. This satellite was not found in the Nanopore data or Illumina data (this and previous

studies) or in previous studies that characterized the most abundant satellites in *D. virilis*.

Manual inspection proved that the AAACGAC satellite was the true consensus found in long

arrays in the reads and did not represent an error in our approaches' characterization of

satellites. Similarly, AAATCAC, AGCCTAT, ACAGGCT, and AATGG were found in megabase

quantities (after normalization) in the Nanopore data - whereas these satellites were not found

in Illumina or PacBio data. We suggest these satellites are also technical artifacts introduced at

the base-calling level.

Fig1: Issues in quantifying simple satellites in sequencing data (all data shown is *D. virilis*). (**A**) Cumulative stacked barplot comparing the performance of the two tested approaches on PacBio data simulated with PBSim from a mock genome. (**B**) Comparing the results of the two approaches on the real PacBio data; "other" refers to additional satellites in the family, including suspected artefactual ones (AAAGCAC for PacBio and AAATCAC + AGCCTAT for Nanopore). (**C**) Same as B for Nanopore data. (**D**) Strand biases in the sequenced satellites in long read sequencing data. Satellites with asterisks are suspected artefactual ones. N refers to the number of satellite regions of reads used for the calculation. (E) Amount of satellites quantified in different datasets: imaginal discs (pure diploid), compared to flies (some polyteny), and fly data that has been quality filtered.

In the PacBio data, the relative amounts of 7mer satellites (AAACTAC, AAACTAT, and

AAATTAC) were lower than expected. This additional evidence led us to hypothesize that there

were context-specific errors in our PacBio data affecting our particular satellites.

If the sequencing were unbiased, we would expect to have an equal amount of satellites being

detected on reads coming from both DNA strands. We evaluated the strand bias in the

simulated and real long-read data for the three most abundant true satellites, as well as some

artifactual satellites. We arbitrarily label the positive strand as AAACTAC and the negative

strand as GTAGTTT, etc. In the simulated data, the positive and negative strands of satellites

were detected in equal amounts (Figure 1D). However, there was a strong strand bias for all

satellites in both the PacBio and Nanopore data (Figure 1D). For PacBio, the real satellites

AAACTAC, AAACTAT, AAATTAC had a positive strand bias, whereas the artifactual satellite had a

negative strand bias: 98% of the reads with this satellite were from the negative strand. Based

on communication with PacBio representatives, this issue seemed to be caused by context-

specific issues with base calling algorithms used for this sequencing run. As base calling

algorithms improve, these issues will likely begin to be remedied. In the Nanopore data, strand

biases were even more extreme: the negative strand was sequenced almost exclusively for real

satellites AAACTAC and AAATTAC and suspect satellite AAATCAC. However, the AAACTAT real

satellite was sequenced almost exclusively on the positive strand. In this case, strand biases

may be caused by unsequenceable secondary structures developing more frequently on one

strand of the satellite DNA than the other. We analyzed Illumina NextSeq reads for *D. virilis*, and

no such strand bias was found.

D. virilis whole-flies have 40% less pericentromeric satellites than non-polytene tissue

210
Polyteny occurs in all differentiated tissues of Dipterans, and is characterized by

multiple rounds of local DNA replication within the same nucleus and without cell division, a

process known as endoreduplication (Smith and Orr-Weaver 1991; Kim *et al.* 2011). However,

the pericentromeric heterochromatin, where most satellite DNA is located, is under-replicated

(Belyaeva *et al.* 1998). It has never been tested if the level of polyteny in an adult fly makes a

difference in the estimate of satellites per genome. Thus, we sequenced adult male flies (which

have multiple polytene tissues) and imaginal discs (which are diploid) from male larvae and

compared the amount of simple satellites in these datasets. We used Illumina sequencing and

PCR-free library preparations to reduce known PCR bias (Wei *et al.* 2018). We found that for

each of the four most abundant 7mer satellites in the *D. virilis* genome, there was

approximately 40% less in the flies compared to the imaginal discs (Figure 1E). This pattern is

not observed for microsatellites which are known to localize outside of pericentromeric

heterochromatin (Figure S2A). We also analyzed publicly available *D. melanogaster* data,

including flies, imaginal discs, and salivary glands (which are the most extreme in polyteny), and

observed this same pattern of under-replication of satellite repeats in polytene tissues (Figure

S2B and S2C).

Reads with tandem repeats had lower quality scores in Illumina data

Upon inspection with FastQC of our data from the polyteny analysis, we found a bimodal

distribution of quality scores, with one peak at 22 and another at 37 (Figure S3A). After filtering

low quality reads, the majority of the reads with simple satellites were removed (Figure S3). The

amount of satellites were reduced by ~15 x after quality filtering (Figure 1E). It is apparent that

in our dataset, simple satellite-containing reads were highly enriched for low quality scores. We

examined other published *D. virilis* Illumina datasets to evaluate if this issue existed in other

sequencing runs. Two other datasets were available and the one that was produced on the

Illumina NextSeq platform like our data (Miller *et al.* 2018) showed the same pattern of biased

quality scores in repetitive reads (Figure S4). The dataset produced on the HiSeq platform (Wei

et al. 2018) did not show this pattern. It should be noted however that the amounts of 7mer

satellites sequenced in the NextSeq datasets were higher than the HiSeq dataset. Our libraries

were multiplexed with other non-*D. virilis* group samples from unrelated projects and only

represented ~20% of the total sequenced lane so that we would not have issues related to low

complexity. We also noticed this pattern (but less dramatically) in our Illumina sequencing of

multiple strains.

Related species have similar but fewer simple repeats

D. novamexicana and *americana* which are 0.38 MY diverged from each other, are sister

species of *D. virilis*, which is approximately 4.5 MY diverged (Caletka and McAllister 2004)

(Figure 2A). We sequenced these species with high coverage PacBio runs and characterized and

quantified satellites. We emphasize the comparison of relative satellite amounts since all are

likely under-represented. Interestingly, we found the most abundant satellite in *D. virilis*,

AAACTAC, is also the most abundant in *D. novamexicana*, albeit with about half the total

amount. The second and third most abundant repeats, AAACTAT and AAATTAC, however were

barely present in *D. novamexicana*. The second most abundant satellite in *D. novamexicana*

was AAACAAC, whereas in *D. virilis* there is only a small amount. *D. americana* has a similar

satellite composition as *D. novamexicana*, but because of issues with a different PacBio

sequencing chemistry used, we do not report the data here.

259   By looking at sequencing data in more diverged species, we can infer when the

AAACTAC satellite family arose. *D. montana*, another member of the virilis group that is 7-11

MY diverged from *D. virilis* (Ostrega and Thompson 1986; Spicer and Bell 2002) (Figure 2A),

does not have any AAACTAC family satellites, and in fact no enrichment of 7 bp satellites. The

most abundant satellite is AAAC, which may be related to the AAACTAC family. *D. hydei* is

approximately 26 MY diverged from *D. virilis* (Izumitani *et al.* 2016), and here 7 bp satellites are

again the most enriched, but the sequences are unrelated to those in *D. virilis* (ACCCATG,

AAAGGTC from PacBio data). From these data, we infer that the AAACTAC family of satellites

arose in the clade leading to the *D. virilis* phylad 4.5-11 MYA. We also analyzed Illumina

sequencing data for *D. lummei*, which is 3 MY diverged from *D. novamexicana/americana*.

AAACTAC is conserved in *D. lummei,* but it is the only enriched 7mer satellite in this species and

its estimated abundance is much lower than the other three *D. virilis* phylad species.

Complex satellites are also abundant in *D. virilis* group genomes

We searched the high-quality genome assemblies for complex satellites (defined here as unit

lengths greater than 20 bp), and we applied NCRF to search in the raw reads. In *D. virilis,* we

found a previously uncharacterized 36-bp satellite

AAAACGACATAACTCCGCGCGGAGATATGACGTTCC making up ~800 kb of our assembly. In *D.*

novamexicana, we found a previously uncharacterized 32 bp satellite

AAAAGCTGATTGCTATATGTGCAATAGCTGAC along with a related 29 bp satellite that spanned

over 1.1 Mb on a single contig. The 3 Mb contig was approximately half satellite and half non-

satellite. The non-satellite half had similarity to chromosome 6 (dot chromosome/Muller

element F) (Figure S5). In *D. americana*, we found this identical 32 bp satellite, but in total its

span was only ~150 kb. In all *D. virilis* group species, we also found a series of similar satellites

varying in size (150-500 bp) related to the previously described helitron central repeat that has

expanded to tandem repeats in the virilis group (Dias *et al.* 2015).

Fluorescence in situ hybridization reveals evolutionary dynamics of 7mer repeats

The location of the 7mer satellites on metaphase chromosomes has never been shown in the *D.*

virilis group. The AAACTAC satellite is conserved between all three species included in this

study: *D. virilis, D. novamexicana,* and *D. americana*, but the abundance varies by

approximately two-fold. The second most abundant satellites have turned over between *D.*

virilis and *novamexicana/americana*. We used FISH of the most abundant 7mers (AAACTAC,

AAACTAT, AAATTAC, AAACAAC) in these three sister species. *D. virilis* and *D. novamexicana*

have the same karyotype with five acrocentric chromosomes plus the very small F element or

"dot chromosome". The strain of *D. americana* we used has centromere-centromere fusions

between the X and 4th chromosomes and the 2nd and 3rd chromosomes.

Fig 2. Comparative analysis of simple satellites in the *D. virilis* group. (A) Phylogeny demonstrating when satellites evolved. Dashed box: virilis phylad; solid box: virilis group. (B) Total amount of different unit lengths (k-mers) of satellites across four related species. (C) DNA-FISH image of *D. virilis* male mitotic cells (D) DNA-FISH image of *D. virilis* female mitotic cells (E) DNA-FISH image of *D. novamexicana* male mitotic cells (F) DNA-FISH image of *D. americana*. Up to three different fluorescent probes were used each time.

297
298
299   FISH results in *D. virilis* show that the most abundant satellite determined by

300 sequencing, AAACTAC, is clearly the most abundant and occurs in approximately equal amounts

301 in the pericentromeric region on the five pairs of large chromosomes. The Y chromosome

302 appears to have slightly less AAACTAC satellite. The second and third most abundant satellites,

303 AAATTAC and AAACTAT, are localized beyond the pericentromeric repeat at or near the

304 centromere. There are five single chromosomes having each of these repeats, indicating that

305 one chromosome pair has different satellite content - which we hypothesized to be the X and Y.

306 Based on differences between male and female FISH results (Figure 2C and 2D), we suggest the

307 Y chromosome has AAACTAT at both distal ends of the chromosome and AAACTAC only flanking

308 one end, whereas the X chromosome has the other centromeric repeat AAATTAC. We were also

able to visualize the dot chromosomes in *D. virilis*, which we find is mostly composed of

AAACTAT. The AAACAAC satellite is present in small amounts in *D. virilis*, very likely on a single

chromosome (Figure S6).

We estimated that *D. novamexicana* has approximately half the AAACTAC as *D. virilis*,

and visualizing it with FISH reveals a pattern that suggests aspects of its evolution. Its

pericentromeric localization is conserved. One chromosome pair has the same amount of

AAACTAC as *D. virilis*, whereas all other chromosomes have a very small amount (Figure 2E).

Based on the FISH images, it appears that it is the 5th chromosome in *D. novamexicana* that has

the greatest amount of pericentromeric AAACTAC conserved. The centromeric repeat on all

major chromosomes is AAACAAC in *D. novamexicana* and *D. americana.* Our images illustrate

clearly the centromere-centromere fusion between chromosome X-4 and 2-3 in *D. americana*

with the satellites being maintained on both sides of the fusion (Figure 2F). None of the four

simple satellite probes bound to what we hypothesize is the Y chromosome of *D. novamexicana*

or *D. americana*. Based on the images we suggest that *D. americana* has an intermediate

amount of pericentromeric AAACTAC satellite compared to *D. virilis* and *novamexicana*.

Some satellite-containing reads are linked to TEs

We used RepeatMasker to detect if any of the reads containing satellites also contain

transposable elements. TEs might be located in islands within the simple repeats at the

centromere as in *D. melanogaster* (Chang *et al.* 2019) (Figure 3A), in more distal regions

flanking the pericentromeric heterochromatin (Figure 3B), or some TEs may have inserted into

long pericentromeric satellite arrays (Figure 3C). For AAACTAC (and its artifactual counterpart

AAACGAC), ~3.5% of reads (2473/75,364) also contained at least 500 bp of a TE insertion. In the

reads that contain TE sequences, satellite repeats were enriched at the beginning and ends of

reads, concordant with the hypothesis that a high proportion of the TEs we found are actually

flanking the long arrays of AAACTAC distally (Figure 3B). In order to understand how many

reads would be expected to contain both satellites and TEs if TEs flanked this satellite and were

not interspersed, we simulated a situation where a large satellite block flanked a large TE block.

This simulation revealed a much smaller amount of reads containing both satellites and TEs

(0.06 %). This result suggests that not only are TEs flanking the pericentromeric satellite

AAACTAC, but there have likely been TE insertions into the satellite arrays. Likely flanking the

proximal end of the pericentromeric satellite are the centromeric satellites AAACTAT or

AAATTAC. 144 reads contained both AAACTAC and AAACTAT repeats (0.19% of AAACTAC reads)

and 94 reads contained both AAACTAC and AAATTAC repeats (0.12% of AAACTAC reads). Based

on our simulations, these proportions of overlapping reads are consistent with our expectation

based on our FISH results that the pericentromeric and centromeric satellites have relatively

clean boundaries and they directly flank each other.

Fig 3. Transposable elements are in close proximity to satellite arrays. (**A**) Satellites near the centromere may be linked to potential islands of retroelements and telomeric TEs. (**B**) Heterochromatic TEs flank satellite arrays. Histograms show the start and end position of transposable element in the satellite-rich read. (**C**) Transposable elements may have inserted into the satellite arrays.

Many of the TE insertions into satellites seemed to be very young. 2080/2473 TE-

containing AAACTAC reads had a TE insertion with less than 15% divergence from the Repbase

consensus, which is the expected error rate of PacBio reads. These insertions included the

superfamilies: DNA elements, LINE/CR1, LINE/I-Jockey, LINE/Penelope, LINE/R1, LTR/Copia,

LTR/Gypsy, LTR/Pao, and Helitrons. We acknowledge however, that there may be a detection

bias for insertions that are less divergent from the consensus. We also remind readers that

there were likely fewer satellite reads sequenced than expected, and that this may have biased

these results if satellite-only regions were sequenced less efficiently than satellite-TE regions.

For centromeric satellites AAATTAC and AAACTAT, the results are more difficult to interpret

since these were more strongly under-represented. However, 300/715 reads of AAACTAT and

49/385 reads AAATTAC contained TEs. Like the AAACTAC pericentromeric satellite, most TE

insertions were low divergence from the Repbase consensus in the centromeric satellite reads.

288/300 and 46/49 TE containing reads had a TE insertion > 500 bp with < 15% divergence for

AAACTAT and AAATTAC, respectively.

There were differences in the TE composition of reads with different satellites. For the

pericentromeric satellite AAACTAC, Gypsy-10_Dvi was the most enriched, followed by Helitrons

(Helitron-1N1_DVir, Helitron-1_DVir, Helitron-2N1_DVir, Helitron-2_DVir). For the AAACTAT

centromeric satellite, Gypsy-10_Dvi was again the most enriched, followed by Penelope. For

the AAATTAC centromeric satellite, Gypsy-2_DVir was the most enriched followed by Penelope.

In both AAACTAC and AAACTAT reads, CR1-1_DVi was the second or third most abundant TE.

Interestingly, R1 was present in relatively high amounts in AAACTAC. In 110/132 of these R1-

AAACTAC reads, rDNA sequences were not also linked. This suggests that some R1 elements,

which are generally localized to rDNA loci, have jumped into or near satellite arrays. This is

concordant with findings that some R1 elements are located outside rDNA loci in Drosophila

(Stage and Eikbush 2009). All centromeres in *D. virilis* are acrocentric, meaning that the

telomeric TEs Het-A and TART (Casacuberta and Pardue 2003) are likely near the centromere

satellites. We found 12 reads linked to AAATTAC that contained matches to TART. Only two

reads linked to any satellite contained a sequence matching HeT-A. We also used BLAST to

detect matches between the genome assembly (masked from the 7mer satellites) and the 7mer

satellite reads. We could not detect any unique regions of the genome that matched non-

satellite sequence on the reads. Non-satellite parts of satellite reads matched hundreds of

places in the genome each.

Variation in *D. virilis* group global strains

D. virilis is globally distributed while its sister species are localized to North America, with *D.*

novamexicana more restricted than *D. americana.* Patterns of variation in satellites may reveal

potential mechanisms that can be hypothesized to be driving satellite evolution. Additionally, *D.*

americana has a polymorphic fusion between the X and 4th chromosomes, so we can identify

differences in satellite composition associated with the fusion. This fusion has been shown to

be currently undergoing meiotic drive, potentially mediated by a larger total centromere or

pericentromere size in the fused strains compared to the non-fused strains (Stewart et al.). On

the other hand, chromosome fusions are often caused by Robertsonian translocations with loss

of some non-essential DNA, which might include pericentromeric satellites (Schubert and Lysak

2011).

395   We used Illumina sequencing with PCR-free library preparation and k-Seek to estimate

the abundance of 7mer satellites across 12 worldwide strains of *D. virilis*, eight strains of *D.*

americana (including four strains that have the X-4 fusion and four that do not), and five strains

of *D. novamexicana* (Table S1). All sequenced strains were male except a female of the *D. virilis*

inbred strain 87 as a comparison. A PCA using only the four most abundant 7mers shows

clustering of the three species, but the separation is much more dramatic in the PCA using the

20 most abundant simple satellites (Figure S7). Overall, *D. virilis* had the highest AAACTAC

satellite content as well as the highest variation, with *D. americana* intermediate between *D.*

virilis and *D. novamexicana* (Figure 4A). Using different normalization procedures including

mapping and GC correction (see Materials and Methods), produced the same relative ranking

of satellite abundances between species. In all cases, the inbred strain from which the genome

sequence was produced had the lowest abundance of AAACTAC. In the case of *D. virilis*, this

difference was very high. This was not due to a normalization bias as we did mapping-free

normalization.

Satellite abundances in *D. virilis* displayed a pattern that appeared to be correlated to

the geographic location strains were collected from. For the centromeric satellite AAATTAC,

there was a linear decrease in abundance from West to East then South following migration

from Beringia (Throckmorton 1982) beginning in China (Figure 4C). For the centromeric satellite

AAACTAT, the pattern was the opposite; a linear increase in abundance from West to East then

South (Figure 4D). We analyzed variation in the satellite repeats to determine if the apparent

linear expansion in AAATTAC also correlated with newer and hence more homogeneous

sequences. On average, the centromeric satellite arrays were very homogeneous (average

above 99% sequence identity in Illumina reads). However, AAACTAT had a slightly higher

sequence identity than AAATTAC (Figure 4 C and D). There was no pattern in average sequence

identity with respect to geography for any of the satellites (Figure 4 C and D). The

pericentromeric satellite AAACTAC has almost identical sequence divergence across the three

species (~98.5%). When comparing between a male and female of the same strain, the male

had lower sequence identity. In *D. americana* and *D. novamexicana*, the AAACAAC satellite had

lower average sequence identity at 97.5%.

Fig 4. Variation in satellites across species and strains. (**A**) AAACTAC abundance across the three species. (**B**) AAACTAC abundance across strains originating from different localities. (**C**) AAACTAC (centromere satellite) abundance across strains originating from different localities. (**D**) AAATTAC abundance across strains originating from different localities.

There were also several other satellites besides the most abundant four that varied

between the *D. virilis* group species. AAACAAT was either absent or in low abundance in *D.*

virilis and *novamexicana*, but present between 100,000 - 250,000 copies in *D. americana,*

indicating a very recent expansion of this satellite. AAAAAC was present in ~100,000 copies in

D. americana and *D. novamexicana*, whereas it is almost absent in *D. virilis*. To differentiate Y-

specific satellites, we included a female of strain 87, the reference genome strain, in our

Illumina sequencing run in addition to a male. There were several Y-specific satellites in *D. virilis*

found by Wei et al. 2018 and validated by our data, which varied between species in the group

(AATAATAG, AATAGATT, and ACATAT). All had different relative patterns of abundance

between species

(https://github.com/jmf422/D_virilis_satellites/blob/master/Intra_inter_species_sequencing/vi

rilis_group_intra-inter_species.html). There was no detectable difference in centromeric or

pericentromeric satellite abundance in *D. americana* strains with and without the polymorphic

centromere-centromere fusion that might be explained by a change in satellite abundance

resulting from molecular events surrounding the fusion (Figure S8).

Amplified repeats of the DINE-1 helitron have been found on the *D. virilis* 5th and Y

chromosomes (Dias *et al.* 2015). We also examined variation in this satellite abundance by

counting reads that mapped to this family of satellites. There was no striking pattern with

respect to geography, however the strain with the highest AAACTAC content had the lowest

DINE-1 content, besides the inbred strain 87, which had lower satellite content all-around

(Figure S9). We expect AAACTAC and DINE-1 to be both in the pericentromeric region of Chr5 in

D. virilis based on our FISH results and the previous work. The male contained 2.6x more DINEs

than the female of the same strain, indicating that ~70% of the DINE-1 repeats in the genome

are located on the Y chromosome in *D. virilis*.

Discussion

Here, we used the satellite DNA-rich genome of *D. virilis* to highlight three previously

uncharacterized mechanisms for biases that occur in sequencing and analyzing satellite DNA.

We emphasize that comparing satellite DNA amounts between different platforms (e.g.

Illumina, PacBio, Nanopore, and even different versions of each) should be done with caution

as each technology has its own biases. We have found that issues arise when multiple kilobase

molecules of simple satellite DNA are attempted to be sequenced by long-read platforms. In

the case of PacBio, systematic errors in base calling may be introduced when sequencing

through long arrays of satellites. This issue is not specific to our satellites, as a recent study has

also found systematic errors and strand biases in shorter arrays of human satellites in both

PacBio and Nanopore reads (Mitsuhashi *et al.*). Circular consensus sequencing (CCS) or "HiFi", a

type of sequencing offered by PacBio which allows an accurate consensus to be produced after

multiple rounds of sequencing the same molecule, may be more appropriate for sequencing

analysis of satellite DNA. Preliminary results based on collaborative efforts with PacBio

representatives show no systematic errors in satellite sequences with the new CCS platform. In

the case of Nanopore, it is possible that a similar satellite-specific base calling errors exists, or

that there is a strand-specific difference in secondary or tertiary structures that occur in long

strands of simple satellite DNA. We caution readers in interpreting simple satellite DNA from

long-read sequencing data and suggest validation with satellites of known sequence and

abundance if available or Illumina reads (without quality filtering). Long read platforms are

already improving their chemistry and software for better satellite characterization. Because

long reads are likely to cross boundaries of different repetitive regions, long read sequencing

proved useful in understanding the length of the satellite arrays and TE insertions into them.

Moreover, we demonstrate that the abundance of satellites in pericentromeric

heterochromatin are underestimated when sequencing flies compared to pure diploid tissue

because of polyteny. We caution readers in performing quality filtering before simple satellite

analysis, as satellite containing reads may be enriched for lower quality scores.

From comparative analysis of satellites, we found the abundant AAACTAC family of

satellites arose in the branch leading to the *virilis* phylad 4.5-11 MYA (Figure 2A). Interestingly,

the most abundant satellite in *D. montana*, 7-11 MY diverged, is AAAC. The AAACTAC and AAAC

satellites may have been derived from one another. However, an enrichment of different 7 bp

satellites does exist in the even more distantly related *D. hydei.* From both FISH and sequencing

analysis, we found that *D. virilis* has the highest total amount of AAACTAC family satellites, *D.*

novamexicana has the lowest, and *D. americana* intermediate. Unlike the pericentromeric

satellite, the centromere-proximal satellite sequence has turned over between *D. virilis* and *D.*

americana/novamexicana. The AAACAAC satellite is present in *D. virilis* in relatively low

amounts, likely on a single chromosome. This satellite then became the centromeric satellite in

D. americana and *novamexicana*, replacing the AAACTAT and AAATTAC satellites. Interestingly,

D. lummei has lower satellite content than the other species and its only high-abundance

simple satellite is AAACTAC. Although sequencing quantified only up to 30 Mb of the AAACTAC

family of satellites, FISH confirmed that these satellites are extremely abundant in *D. virilis* and

the 40% of the genome estimate seems realistic.

We can make hypotheses about how and why the satellites expanded in *D. virilis*. We

know that mutation rates for changes in copy number of satellite DNA are high, and potentially

have a tendency to expand rather than contract in the absence of selection (Flynn *et al.* 2017,

2018). High rates of mutation must be accompanied by a regime that would allow a satellite

copy number increase to sweep the population - which could be mediated by positive selection

if there is a benefit of the satellite increase, or centromere drive if the phenomenon is at play in

this group. Alternatively, in a situation where satellites are slightly deleterious, small effective

population sizes in isolated populations or continued bottlenecks could allow satellites to

expand in the genome without being removed by selection. However, *D. novamexicana* has the

lowest effective population size of the virilis pylad and yet it has the lowest amount of satellite

DNA. We already know that the centromere-to-centromere fusions in *D. americana* have

undergone meiotic drive hypothesized to be mediated by the increase in centromere total size

with the fusion (Stewart et al.). The mechanism allowing drive in *D. americana* may have been

at play in the branch leading to *D. virilis* or may be currently occurring. Why have satellites not

expanded to this extent in the other species? *D. virilis* might have some attributes about its

biology that made the satellite expansion favorable or allowable. For example, genome size is

positively correlated with development time in Drosophilidae (Gregory and Johnston 2008). *D.*

virilis has a slow development time, and this may have evolved in concert with the expansion in

satellite abundance in its genome.

We can also use data from multiple strains to make hypotheses about factors driving

satellite DNA evolution in *D. virilis*. Ancestrally, *D. virilis* had a relatively small effective

population size in an isolated range in Asia, and has undergone a recent population and range

expansion (Mirol et al. 2008). The amount of the most abundant pericentromeric satellite,

AAACTAC, does not vary greatly across strains from different localities. Assuming we sequenced

a strain from the ancestral range, this suggests that population bottlenecks did not allow

AAACTAC to expand, and the satellite expansion likely occurred before the population

expansion. However, the abundance of the centromere-proximal satellites does show a pattern

with respect to where samples were collected. There is increasing AAACTAT abundance on a

gradient from west to east, correlated with decreasing AAATTAC on the same gradient.

AAACTAT may be experiencing an expansion or starting to occupy additional centromeres,

while the AAATTAC satellite may be decreasing in parallel because of a trade-off or maximum

limit on satellite amount. AAACTAT is located on the Y, two other chromosomes and the dot,

and AAATTAC is located on the X and two other chromosomes. AAACTAT has a slightly higher

average sequence identity than AAATTAC, concordant with the hypothesis that it has recently

increased its copy number.

Our observation of rapid evolution and enrichment of AAACTAC in *D. virilis* in a short

evolutionary time period (a few million years) is consistent with the centromere-drive model to

account for the evolution of centromere complexity in genetic conflict (Malik and Bayes, 2006).

In this model, the asymmetric female meiosis can cause competition between the centromeres

with or without newly formed satellites (or with more or less satellites) to be included into the

oocyte to pass to next generation. A consequence of the competition would be runway

expansions of centromeric satellites, and rapid replacements by novel satellites. According to

this model, our observation of increasing AAACTAT abundance in a west-east gradient may

reflect an ongoing centromere drive combined with selection "pushing back".

Interestingly, in *D. novamexicana*, AAACTAC was greatly reduced in the pericentromeric

regions on all chromosome pairs except one. Based on the FISH images in *D. novamexicana* and

americana, we hypothesize that it is the 5th chromosome that has the high amount of

AAACTAC satellite. This is interesting because previous work has shown that the 5th

chromosome contains a high amount of DINE-1 helitron satellite in *D. virilis* but not in *D.*

americana (Dias et al. 2015). We found that all chromosomes including Chr5 contain a large

amount of AAACTAC in *D. virilis*. The DINE-1 repeat and the pericentromeric satellite may both

be present (and possibly in conflict or competing) on the *D. virilis* 5th chromosome, but in the

branch leading to *D. novamexicana*, AAACTAC took over. DINE-1 had a relatively consistent

amount across different *D. virilis* strains, however the strain with the highest AAACTAC amount

is an outlier with a lower DINE-1 amount (Figure S9). This may indicate a maximum threshold of

satellites was reached on this chromosome, and one satellite had to reduce its abundance. We

have seen evidence for this trade-off, or appearance of competitive exclusion, being invoked

under selection in our previous studies (Flynn *et al.* 2017, 2018). There may have been a similar

conflict on Chr5 of *D. novamexicana*, where AAACTAC retained a high copy number to prevent

DINE-1 from expanding. Interestingly, the opposite has occurred on the *D. novamexicana* and

D. americana Y chromosome, where AAACTAC family satellites are absent but DINE repeats are

abundant. A potential mechanism mediating apparent stabilizing selection on total satellite

abundance is that satellites can act as a sink for heterochromatin factors, with their abundance

affecting chromatin state (Lemos *et al*. 2010).

The AAACTAC satellite has remained conserved in sequence and location in the virilis

phylad. It has also maintained high levels of sequence identity that is equal in the three species

we sequenced (98.5% based on Illumina reads). The conservation may reflect a constraint due

to selection or a pervasive mechanism of concerted evolution. The periodicity of the sequence

may stabilize the DNA helix wrapping around nucleosomes, or it may be constrained by

coevolution of an important satellite DNA binding protein (Maio *et al.* 1977; Jagannathan *et al.*

2018). Additionally, within the AAACTAC family, the position and identity of the four A-

nucleotides are conserved in all four satellites (AAACTAC , AAATTAC, AAACTAT, AAACAAC) -

which may indicate constraint based on the above mechanisms. Conservation of particular

satellite unit lengths and "AA" periodicities have been found in other divergent species

(Lowman and Bina 1990). Concerted evolution of satellites could be achieved by repeated

recycling of units by copy number changes associated with replication slippage or unequal

recombination or gene conversion (Walsh 1987; Elder and Turner 1995). However,

recombination in the pericentromeric heterochromatin has never been detected in wild-type

flies (Mehrotra and McKim 2006; Hughes *et al.* 2018). On the other hand, if recombination were

occurring, satellite arrays will eventually be lost unless they are conserved by selection

(Charlesworth *et al.* 1986). Clearly, we are still lacking in understanding how and why long

simple satellite arrays maintain their homogeneity, and whether recombination plays a role in

their dynamics. Concordant with the hypothesis that recombination is playing a role, males

have lower average sequence identity in the 7mer satellites than females, which could indicate

increased decay on the Y chromosome where there no homologous recombination (Figure 4A).

Moreover, our results suggest that transposable elements flank the AAACTAC satellite

array (likely more distally to the centromere) and some TEs have inserted within the array. Our

analysis suggests that most TE insertions are recent, but because of the under-representation

of satellite containing reads, we cannot estimate the number of TE insertions that have

occurred into the satellite arrays. Our analysis is also not precise enough to determine if there

are islands of TEs at the centromere in *D. virilis* as has been demonstrated in *D. melanogaster*

(Chang et al. 2019).

We found no difference in centromeric and pericentromeric satellites abundances

between *D. americana* strains that differ in their X-4 fusion status. This suggests that the fusion

event did not result in a large loss of satellites, making the total centromere and

pericentromere size is indeed larger on the X-4 fused chromosome than the single unfused

chromosome, concordant with the hypothesis that a larger centromeric region results in

centromere drive (Stewart et al.). Another interesting observation from the sequencing of

multiple strains of the three species was that in all cases, the inbred strain that the reference

genome was made from had the lowest amount of AAACTAC. For *D. virilis*, this difference was

extreme. It is tempting to speculate that the process of inbreeding and/or long periods in the

lab may have driven the reduction in pericentromeric satellite abundance.

In conclusion, our results show very rapid dynamics in the abundant satellites of the *D.*

virilis group that are likely explained by various cellular and population-level forces that are not

yet understood. Further studies can test if there is a species-specific upper limit to satellite

amount per genome or per chromosome upon which negative fitness effects occur, which may

result in trade-offs or competition between satellites. Centromere drive may be an important

process affecting satellite evolution in this species group, and might partially explain why the

satellites expanded 4.5-11 MYA, why satellite sequences at the centromere turned over more

rapidly, and why there is a gradient of increasing satellite content related to geographical

distribution of strains. A more extensive study to determine if inbreeding or extended periods

in the lab drives a reduction in satellite abundance will help illuminate the processes that are

important for maintaining satellite content. Determining the frequency of recombination in the

large pericentromeric heterochromatin blocks in species like *D. virilis* will be challenging but

important for understanding how the satellites maintain homogeneity in their sequence. To

understand the role of satellites and the importance of their sequence, unit length, and

abundance, researchers can strive to develop methods to engineer satellites by modifying

specific bases and their abundances.

MATERIALS AND METHODS

All scripts for analyzing the data and to produce the results we show are here:

https://github.com/jmf422/D_virilis_satellites. Illumina sequencing reads generated for this

study are deposited in NCBI SRA under accession PRJNA548201. Raw PacBio reads will be

deposited under the same accession. Both will be released upon publication.

Characterizing satellite DNA from genome assemblies

All scripts and R markdown files used for this analysis are provided in

https://github.com/jmf422/D_virilis_satellites/tree/master/Genome_assembly_analysis.

We used genome assemblies produced by the PacBio sequencing project

(https://www.ncbi.nlm.nih.gov/bioproject/?term=txid7214[Organism:noexp]) of *D. virilis, D.*

novamexicana, and *D. americana*. We also downloaded the *D. virilis* genome produced by

Nanopore sequencing from (Miller *et al.* 2018), and the CAF1 assembly from (Drosophila 12

Genomes Consortium *et al.* 2007). We used Phobos (https://www.ruhr-uni-

bochum.de/spezzoo/cm/cm_phobos.htm) and Tandem repeats finder (Benson 1999) to

characterize simple and complex satellites in these genome assemblies. To identify the

chromosomal linkage of complex satellites in the genome assembly, we produced a dotplot

with D-GENIES (Cabanettes and Klopp 2018).

Characterizing satellite DNA from raw long reads

Characterizing and quantifying satellites from long reads is a challenge because of the

sequencing high error rate. We used two approaches to characterize satellites from raw long

reads. The first approach, we call k-Seek + Phobos, in which we first broke the reads into 100 bp

subreads and ran k-Seek on them. k-Seek is very efficient for analyzing many reads, however is

not very sensitive for reads with a high error rate since it was designed for Illumina reads (Wei

et al. 2014). If k-Seek found satellites on at least one subread, we would run the complete

parent read through Phobos. Phobos is more sensitive to imperfect repeats and error rates, but

cannot handle huge quantities of data; thus why we only ran the portion of reads identified by

k-Seek to have tandem repeats. This approach allowed us to characterize satellites *de novo* and

quantify them. All scripts for the analysis of long reads with the k-Seek + Phobos approach are

located here:

https://github.com/jmf422/D_virilis_satellites/tree/master/LongRead_kseek_Phobos. The

second approach we used is Noise-Cancelling Repeat Finder (NCRF, (Harris *et al.*)). This

program was designed to quantify satellites from long reads with high error rates. However, it

cannot identify satellites *de novo* and requires specific satellite sequences to search for. NCRF

also requires a "max divergence allowed" parameter, which we tuned with simulations (see

below). Scripts used for the NCRF approach are located here:

https://github.com/jmf422/D_virilis_satellites/tree/master/LongRead_NCRF.

We did simulations to assess both approaches:

https://github.com/jmf422/D_virilis_satellites/tree/master/Simulations. First, we created a

simplified mock *D. virilis* genome with a satellite DNA composition estimated from our FISH

results. We could not use the genome assembly because it contained very little satellite DNA.

Specifically, each chromosome had a centromeric satellite either AAATTAC or AAACTAC

followed by the pericentromeric satellite AAACTAC, combined taking up 40% of the genome.

The non-satellite DNA portion of the genome was generated randomly with a 40% GC content.

We then used PBSim (Ono *et al.* 2013) to simulate PacBio reads and we used these simulated

reads for multiple analyses. First, we used them to tune the max divergence parameter of NCRF

by running NCRF repeatedly with max divergence parameters ranging 18-30%. We found that

the amount of satellites found, particularly the most abundant one, levelled off at 25% max

divergence. This is the parameter value we used moving forward. We also used these simulated

reads to quantify satellites with both approaches and compare them. Finally, we used these

simulated reads to assess strand biases in long read sequencing data (see below).

Identification of biases in simple satellites in long read data

We suspected that there were biases in the satellite DNA found in the *D. virilis* group PacBio

(and Nanopore) data because we found high abundance satellites that had never been found

before with other types of data, and so we suspected they were artifactual. These artifactual

satellites were found with both kSeek + Phobos and NCRF approaches, but were not found in

the simulated data. We tried testing for a strand bias in reads that contained satellite DNA.

Using both the summarized output from NCRF and validated with a custom script

(LongRead_NCRF/which_strand_pacbio_script.sh), we counted the satellite DNA stretches that

originated from each the positive and negative strand. The positive strand is defined as the one

that contains the satellite AAACTAC and derivatives (more As than Ts), and the negative strand

is the one that contains the reverse complement (e.g. GTAGTTT, more Ts than As). We did this

for the three satellites used in the simulated data and real and artefactual satellites found in

the PacBio and Nanopore data. Detailed analysis and visualization of the biases is shown here:

LongRead_kseek_Phobos/longread_analysis.html

Sequencing of polytene and non-polytene tissue

To acquire *D. virilis* pure diploid tissue, we dissected male 3rd instar larvae and collected

imaginal discs including the eye-antennal disc and wing discs. Approximately 100 larvae were

required to get enough DNA (>1 ug). We also collected ~5 adult flies for fly libraries. We used

the inbred genome assembly strain 87 for these libraries. DNA was extracted with Qiagen

DNeasy blood and tissue kit and PCR-free libraries were prepared. Libraries were run on an

Illumina NextSeq with 1 x 150 bp reads, and each sample took up approximately 7% of the

flowcell. The other libraries run on this flowcell were from an unrelated project including

RNAseq from other species. Reads were analyzed with k-Seek both before and after filtering

with Trimmomatic (Bolger et al. 2014). FastQC was run to evaluate the quality of the reads.

Scripts are here: https://github.com/jmf422/D_virilis_satellites/tree/master/Polyteny. We also

analyzed publicly available *D. melanogaster* data from the same strain and same sequencing

platform of embryos (non-polytene), salivary glands (extreme polyteny) from (Yarosh and

Spradling 2014), and flies (varied levels of polyteny) from (Gutzwiller *et al.* 2015).

Fluorescence in situ hybridization of satellite DNAs

We followed the protocol of (Larracuente and Ferree 2015) for satellite DNA FISH. We ordered

the following probes from IDT with 5' modifications: $(AAACTAC)_6$ with alexa-488 fluorophore,

$(AAACTAT)_6$ with Cyanine5 fluorophore, $(AAATTAC)_6$ with Cyanine3 fluorophore, $(AAACAAC)_6$

with Cyanine3 fluorophore, and $(AAACGAC)_6$ with Cyanine5 fluorophore. We hybridized three

probes at a time, to allow for similar probes to compete to result in specific hybridization with

the rationale shown in (Beliveau *et al.* 2015). Hybridization temperature was 32°C. We imaged

on an Olympus fluorescent microscope and Metamorph capture system at the Cornell Imaging

Facility. Composite images were produced with ImageJ.

Characterizing TEs linked to satellites and satellites anchored to the genome assembly

We extracted the reads identified to have the 7 bp satellites on them from NCRF results, and

then we ran RepeatMasker (http://www.repeatmasker.org) on these reads using parameters:

"-nolow" and "-species Drosophila". All reads had at least 500 bp of tandem satellite on them

according to NCRF default parameters, and to avoid spurious identification of TEs from semi-

repetitive fragments, we described only TEs in reads that also had at least 500 bp of a TE

identified from RepeatMasker. We also BLASTed the same satellite reads to the genome

assembly to evaluate if satellite reads could be anchored to the genome assembly. Analysis

scripts are here: https://github.com/jmf422/D_virilis_satellites/tree/master/TEs_satellites.

Sequencing of multiple *D. virilis* group strains

We obtained as many strains of *D. virilis* that have information about where they were

collected as possible. This included 12 strains as live stocks we obtained either from stocks in

our lab or from the Drosophila species stock center (Table S1). We also prepared a female

library for strain 87 for when we wanted to differentiate Y-specific satellite patterns. We also

obtained five strains of *D. novamexicana* and eight strains of *D. americana*. All were obtained

from live stocks and the inbred genome strains were included for both species as well (strain 14

and G96, respectfully). For *D. americana*, we included four strains that have the chromosome X-

4 fusion and four strains that do not have it, based on communication with the Bryant

McAllister lab. DNA was extracted as above from five flies each and samples were prepared

identically as above and sequenced on 50% of 3 flowcells of Illumina NextSeq 1 x 150 bp reads.

We dispersed the samples from each species between multiple flowcells. Our samples took up

only half the flowcell with the other half being occupied by a RNAseq libraries from an

unrelated project.

All scripts used to analyze these data are located here:

https://github.com/jmf422/D_virilis_satellites/tree/master/Intra_inter_species_sequencing.

Reads were evaluated with FastQC and not filtered for quality based on the potential bias of

Illumina quality scores on satellites. We used k-Seek to quantify satellites. We tried several

normalization strategies but decided the most appropriate was a mapping-free normalization.

We estimated average depth by dividing the total number of bases sequenced by the estimated

genome size by flow cytometry (Bosco *et al.* 2007). We believe this was the best option in this

case because: 1) we were concerned about a mapping bias because for each species the strain

that the genome assembly was made from may have more reads map to it; 2) after masking the

genome from the 7mer satellites and also discluding the X and Y contigs (because we had male

and female strains, and the Y chromosome contained more low GC regions) - there was little

difference in coverage based on GC content. We include results when we used a mapping based

GC normalization in the sub-directory "AlternativeNormalization".

We used NCRF with modified parameters (minlength=100, maxdiv=10) to characterize

the average sequence identity of satellite arrays from the Illumina data. To quantify DINE-1

satellites across *D. virilis* strains, we produced a library of DINE-1 satellite variants based on our

PacBio genome analysis. We then mapped Illumina reads to this library and normalized the

number of reads that mapped to any sequence in the library by the estimated depth. We also

analyzed Illumina DNA sequencing reads of *D. montana* (Parker et al. 2018) and *D. lummei*

(Ahmed-Braimah *et al.* 2017) with k-Seek to identify the most abundant satellites and whether

or not the AAACTAC satellite family was present.

Acknowledgments

We thank Yasir Ahmed-Braimah for helpful discussions and advice for some analyses. We also

thank Bryant McAllister for providing *D. americana* strains along with their fusion status. We

are grateful for Elissa Cosgrove's help with some computational trouble-shooting and Asha

Jain's help in preparing DNA sequencing libraries. We also thank Danny Miller for useful

discussions and for providing the raw Nanopore reads from his study. Sarah Kingan and Jane

Landolin, from PacBio, were very helpful in exploring the causes for the artefactual repeats. We

thank Amanda Larracuente for advice on FISH protocols and the Cornell Imaging Facility for use

of their microscope. This project was funded by NIH grant number: GM116113 to RAW, ML, and

AGC. JMF was supported by an NSERC PGS D fellowship.

775
776
References

Ahmed-Braimah Y. H., R. L. Unckless, Andrew G. Clark, 2017 Evolutionary Dynamics of Male

Reproductive Genes in the *Drosophila virilis* Subgroup. G3. 7:3145-3155.

Beliveau B. J., A. N. Boettiger, M. S. Avendaño, R. Jungmann, R. B. McCole, *et al.*, 2015 Single-molecule

super-resolution imaging of chromosomes and in situ haplotype visualization using Oligopaint FISH

probes. Nat. Commun. 6: 7147.

Belyaeva E. S., I. F. Zhimulev, E. I. Volkova, A. A. Alekseyenko, Y. M. Moshkin, *et al.*, 1998 Su(UR)ES: a

gene suppressing DNA underreplication in intercalary and pericentric heterochromatin of

Drosophila melanogaster polytene chromosomes. Proc. Natl. Acad. Sci. U. S. A. 95: 7532–7537.

Benson G., 1999 Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Research

27: 573–580.

Bolger A. M., M. Lohse, B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics 30:2114-2120.

Bosco G., P. Campbell, J. T. Leiva-Neto, and T. A. Markow, 2007 Analysis of Drosophila species genome

size and satellite DNA content reveals significant differences among strains as well as between

species. Genetics 177: 1277–1290.

Cabanettes F., and C. Klopp, 2018 D-GENIES: dot plot large genomes in an interactive, efficient and

simple way. PeerJ 6: e4958.

Caletka B. C., and B. F. McAllister, 2004 A genealogical view of chromosomal evolution and species

delimitation in the Drosophila virilis species subgroup. Mol. Phylogenet. Evol. 33: 664–670.

Casacuberta E., and M.-L. Pardue, 2003 HeT-A elements in Drosophila virilis: retrotransposon telomeres

are conserved across the Drosophila genus. Proc. Natl. Acad. Sci. U. S. A. 100: 14091–14096.

Chang C.-H., and A. M. Larracuente, 2019 Heterochromatin-Enriched Assemblies Reveal the Sequence

and Organization of the Drosophila melanogaster Y Chromosome. Genetics 211: 333–348.

Chang C.-H., A. Chavan, J. Palladino, X. Wei, N. M. C. Martins, *et al.*, 2019 Islands of retroelements are

the major components of Drosophila centromeres. PLoS Biology 17(5):e3000241

Charlesworth B., C. H. Langley, and W. Stephan, 1986 The evolution of restricted recombination and the

accumulation of repeated DNA sequences. Genetics 112: 947–962.

Charlesworth B., P. Sniegowski, and W. Stephan, 1994 The evolutionary dynamics of repetitive DNA in

eukaryotes. Nature 371: 215–220.

Dias G. B., P. Heringer, M. Svartman, and G. C. S. Kuhn, 2015 Helitrons shaping the genomic architecture

of Drosophila: enrichment of DINE-TR1 in α- and β-heterochromatin, satellite DNA emergence, and

piRNA expression. Chromosome Res. 23: 597–613.

Drosophila 12 Genomes Consortium, A. G. Clark, M. B. Eisen, D. R. Smith, C. M. Bergman, *et al.*, 2007

Evolution of genes and genomes on the Drosophila phylogeny. Nature 450: 203–218.

Elder J. F. Jr, and B. J. Turner, 1995 Concerted evolution of repetitive DNA sequences in eukaryotes. Q.

Rev. Biol. 70: 297–320.

Elliott T. A., and T. Ryan Gregory, 2015 What's in a genome? The C-value enigma and the evolution of

eukaryotic genome content. Philos. Trans. R. Soc. Lond. B Biol. Sci. 370: 20140331.

Flynn J. M., I. Caldas, M. E. Cristescu, and A. G. Clark, 2017 Selection Constrains High Rates of Tandem

Repetitive DNA Mutation in Daphnia pulex. Genetics genetics.300146.2017.

Flynn J. M., S. E. Lower, D. A. Barbash, and A. G. Clark, 2018 Rates and Patterns of Mutation in Tandem

Repetitive DNA in Six Independent Lineages of Chlamydomonas reinhardtii. Genome Biol. Evol. 10:

1673–1686.

Gall J., E. Cohen, and M. Polan, 1971 Repetitive DNA sequences in Drosophila. Chromosoma 33.

https://doi.org/10.1007/bf00284948

Gall J. G., and D. D. Atherton, 1974 Satellite DNA sequences in Drosophila virilis. J. Mol. Biol. 85: 633–

664.

Gregory T. R., 2001 Coincidence, coevolution, or causation? DNA content, cell size, and the C-value

enigma. Biol. Rev. Camb. Philos. Soc. 76: 65–101.

Gregory T. R., and J. S. Johnston, 2008 Genome size diversity in the family Drosophilidae. Heredity 101:

228–238.

Gutzwiller F., C. R. Carmo, D. E. Miller, D. W. Rice, I. L. G. Newton, *et al.*, 2015 Dynamics of Wolbachia

pipientis Gene Expression Across the Drosophila melanogaster Life Cycle. G3 5: 2843–2856.

Harris R. S., M. Cechova, and K. Makova, Noise-Cancelling Repeat Finder: Uncovering tandem repeats in

error-prone long-read sequencing data:

Henikoff S., 2001 The Centromere Paradox: Stable Inheritance with Rapidly Evolving DNA. Science 293:

1098–1102.

Hughes S. E., D. E. Miller, A. L. Miller, and R. S. Hawley, 2018 Female Meiosis: Synapsis, Recombination,

and Segregation in. Genetics 208: 875–908.

Izumitani H. F., Y. Kusaka, S. Koshikawa, M. J. Toda, and T. Katoh, 2016 Phylogeography of the Subgenus

Drosophila (Diptera: Drosophilidae): Evolutionary History of Faunal Divergence between the Old

and the New Worlds. PLoS One 11: e0160051.

Jagannathan M., R. Cummings, and Y. M. Yamashita, 2018 A conserved function for pericentromeric

satellite DNA. Elife 7. https://doi.org/10.7554/eLife.34122

Jagannathan M., R. Cummings, and Y. M. Yamashita, 2019 The modular mechanism of chromocenter

formation in. Elife 8. https://doi.org/10.7554/eLife.43938

Kim J. C., J. Nordman, F. Xie, H. Kashevsky, T. Eng, *et al.*, 2011 Integrative analysis of gene amplification

in Drosophila follicle cells: parameters of origin activation and repression. Genes Dev. 25: 1384–

1398.

Larracuente A. M., and P. M. Ferree, 2015 Simple Method for Fluorescence DNA In Situ Hybridization to

Squashed Chromosomes. Journal of Visualized Experiments.

Lemos B., A. T. Branco, D. L. Hartl, 2010 Epigenetic effects of polymorphic Y chromosomes modulate

chromatin components, immune response, and sexual conflict. PNAS 107:15826-15831.

Lima L. G. de, M. Svartman, and G. C. S. Kuhn, 2017 Dissecting the Satellite DNA Landscape in Three

Cactophilic Sequenced Genomes. G3 7: 2831–2843.

Lowman H., and M. Bina, 1990 Correlation between dinucleotide periodicities and nucleosome

positioning on mouse satellite DNA. Biopolymers 30: 861–876.

Malik H.S. and J.J. Bayes, 2006. Genetic conflicts during meiosis and the evolutionary origins of

centromere complexity. Biochem Soc Trans. 34: 569-573.

Maio J. J., F. L. Brown, and P. R. Musich, 1977 Subunit structure of chromatin and the organization of

eukaryotic highly repetitive DNA: recurrent periodicities and models for the evolutionary origins of

repetitive DNA. J. Mol. Biol. 117: 637–655.

Mehrotra S., and K. S. McKim, 2006 Temporal Analysis of Meiotic DNA Double-Strand Break Formation

and Repair in Drosophila Females. PLoS Genetics 2: e200.

Miller D. E., C. Staber, J. Zeitlinger, and R. Scott Hawley, 2018 Highly Contiguous Genome Assemblies of

15 Drosophila Species Generated Using Nanopore Sequencing. G3: Genes|Genomes|Genetics 8:

3131–3141.

Mills W. K., Y. C. G. Lee, A. M. Kochendoerfer, E. M. Dunleavy, G. H. Karpen. 2019 RNA transcribed from

heterochromatic simple-tandem repeats are required for male fertility and histone-protamine

exchange in *D. melanogaster*. Biorxiv doi: https://doi.org/10.1101/617175.

Mitsuhashi S., M. C. Frith, T. Mizuguchi, S. Miyatake, T. Toyota, *et al.*, 2019 Tandem-genotypes: Robust

detection of tandem repeat expansions from long DNA reads. Genome Biology 20:58.

Ohno S., 1972 So much "junk" DNA in our genome. Brookhaven Symp. Biol. 23: 366–370.

Ono Y., K. Asai, and M. Hamada, 2013 PBSIM: PacBio reads simulator—toward accurate genome

assembly. Bioinformatics 29: 119–121.

Ostrega M. S., and V. Thompson, 1986 Mitochondrial DNA Restriction site polymorphism in Drosophila

montana and drosophila virilis. Biochemical Systematics and Ecology 14: 515–519.

Parker D. J., R. A. W. Wiberg, U. Trivedi, V. I. Tyukmaeva, K. Gharbi, *et al.*, 2018 Inter and Intraspecific

Genomic Divergence in Drosophila montana Shows Evidence for Cold Adaptation. Genome Biol.

Evol. 10: 2086–2101.

Pavlek M., Y. Gelfand, M. Plohl, and N. Meštrović, 2015 Genome-wide analysis of tandem repeats in

Tribolium castaneum genome reveals abundant and highly dynamic tandem repeat families with

satellite DNA features in euchromatic chromosomal arms. DNA Res. 22: 387–401.

Schubert I., and M. A. Lysak, 2011 Interpretation of karyotype evolution should consider chromosome

structural constraints. Trends Genet. 27: 207–216.

Smith A. V., and T. L. Orr-Weaver, 1991 The regulation of the cell cycle during Drosophila

embryogenesis: the transition to polyteny. Development 112: 997–1008.

Spicer G. S., and C. D. Bell, 2002 Molecular Phylogeny of the Drosophila virilis Species Group (Diptera:

Drosophilidae) Inferred from Mitochondrial 12S and 16S Ribosomal RNA Genes. Annals of the

Entomological Society of America 95: 156–161.

Stage D. E., T. H. Eickbush, 2009 Origin of nascent lineages and the mechanisms used to prime second-

strand DNA synthesis in the R1 and R2 retrotransposons of Drosophila. Genome Biology 10:R49.

Stewart B. S., Y. H. Ahmed-Braimah, D. G. Cerne, B. F. McAllister, Female meiotic drive preferentially

segregates derived metacentric chromosomes in Drosophila.

Throckmorton LH: *The Genetics and Biology of Drosophila Volume 3b*. Academic Press: New York; 1982.

Walsh J. B., 1987 Persistence of tandem arrays: implications for satellite and simple-sequence DNAs.

Genetics 115: 553–567.

Wei K. H.-C., J. K. Grenier, D. A. Barbash, and A. G. Clark, 2014 Correlated variation and population

differentiation in satellite DNA abundance among lines of Drosophila melanogaster. Proc. Natl.

Acad. Sci. U. S. A. 111: 18793–18798.

Wei K. H.-C., S. E. Lower, I. V. Caldas, T. J. S. Sless, D. A. Barbash, *et al.*, 2018 Variable Rates of Simple

Satellite Gains across the Drosophila Phylogeny. Mol. Biol. Evol. 35: 925–941.

Yarosh W., and A. C. Spradling, 2014 Incomplete replication generates somatic DNA alterations within

Drosophila polytene salivary gland cells. Genes & Development 28: 1840–1855.

Supporting Information Captions

Figure S1. NCRF Simulations results. Running NCRF with different max error allowed on reads from

PBSIM off a mock genome composed of perfect repeats.

Figure S2. Polyteny vs. diploid analysis. (A) Microsatellites expected to not be exclusive in

pericentromeric heterochromatin (control for Fig 2E). (B) Analysis of *D. melanogaster* sequencing

data from embryos, flies, and salivary glands. (C) Microsatellites in *D. melanogaster* data, control

for Fig S2B.

Fig S3. Satellite containing reads are enriched for low quality scores in Illumina data. A) Quality score

distribution in the raw reads. (B) Quality score distribution after quality filtering. (C) GC distribution

of raw reads. (D) GC distribution of reads after quality filtering.

Fig S4. Satellite containing reads are enriched for low quality reads in one Illumina dataset (A) produced

on a NextSeq, but not on a second dataset (B) produced on a HiSeq.

Figure S5. Dotplot of D. novamexicana contig 000018F, which contained a high amount of the 32 bp

satellite (from 0-1.5 M) and its similarity to a large region on *D. virilis* chromosome 6 (dot).

Fig S6. FISH image of D. virilis male. The AAACAAC satellite appears on a single chromosome pair. This is

clear from the metaphase chromosomes (middle) as well as the interphase cells where two distinct

foci are localized. The distinct foci in interphase cells is in contrast to the AAACTAC satellite, which

takes up a large portion of the nucleus.

Fig S7. PCA of satellite DNA copy number of D. virilis group strains. (A) Using only the abundances of the

satellites AAACTAC, AAATTAC, AAACTAT, AAACAAC. (B) Using the abundances of the 20 most

abundant simple satellites.

Fig S8. Comparison of D. americana strains with and without the X-4 chromosomal fusion. (A) PCA using

the top 20 simple satellites. (B) Boxplot of AAACTAC copy number. (C) Boxplot of AAACAAC copy

number.

Fig S9. DINE amount vs. AAACTAC copy number. DINE amount is the number of reads mapped to the

DINE repeat / estimated depth.