The little caterpillar that could – Tobacco Hornworm (*Manduca sexta*) caterpillars as a
novel host model for the study of fungal virulence and drug efficacy

Naomi Lyons[1], Isabel Softley[2], Andrew Balfour[2], Carolyn Williamson[2], Heath O'Brien[3], and
Stephanie Diezmann[2, 4]

[1] School of Molecular Cell Biology and Biotechnology, Tel Aviv University, Tel Aviv, Israel
[2] Department of Biology & Biochemistry, University of Bath, Bath, United Kingdom
[3] LivingDNA, Frome, United Kingdom
[4] School of Cellular and Molecular Medicine, University of Bristol, Bristol, United Kingdom

Corresponding author: Stephanie Diezmann
14   s.diezmann@bristol.ac.uk

16 Running title: Caterpillars as fungal virulence host model

18 Keywords: *Manduca sexta*,
19   caterpillar,
20   fungal virulence,
21   *Candida*,
22   *Cryptococcus*,
23   *Saccharomyces*,
24   *Metschnikowia*,
25   antifungal drug,
26   host model,
27   fungal burden

Abstract

Pathogenic yeast species can cause life-threatening infections in humans. The two leading

yeast pathogens, *Candida albicans* and *Cryptococcus neoformans*, cause systemic infections

in >1.4 million patients world-wide with mortality rates approaching 75%. It is thus

imperative to study fungal virulence mechanisms, stress response pathways, and the efficacy

of antifungal drugs. This is commonly done using mammalian models. To address ethical and

practical concerns, invertebrate models, such as wax moth larvae, nematodes, or flies, have

been introduced over the last two decades. To address short-comings in existing invertebrate

host models, we developed fifth instar caterpillars of the Tobacco Hornworm moth *Manduca*

sexta as a novel host model for the study of fungal virulence and drug efficacy. These

caterpillars can be raised at standardised conditions, maintained at 37˚C, can be injected with

defined amounts of yeast cells, and are susceptible to the most threatening yeast pathogens,

including *C. albicans*, *C. neoformans*, *C. auris*, and *C. glabrata*. Infected caterpillars can be

rescued by treatment with commonly deployed antifungal drugs and importantly, fungal

burden can be assessed daily throughout the course of infection in a single caterpillar's faeces

and hemolymph. Notably, these animals are large enough so that weight provides a reliable

and reproducible measure of fungal virulence. This model combines a suite of parameters that

recommend it for the study of fungal virulence.

Introduction

Fungal infections pose a serious threat to human health and well-being world-wide.

Each year, as many, if not more patients, die of fungal infections than of malaria or

tuberculosis[1]. The leading yeast pathogens, *Candida albicans* and *Cryptococcus neoformans*,

together account for >1,400,000 life-threatening infections world-wide with mortality rates

approaching 75%[1]. Candidemia, most commonly caused by *C. albicans*, is the fourth most

common cause of nosocomial blood stream infections, only surpassed by infections with

Staphylococci and *Enterococcus* spp. Disturbingly, candidemia incidence rates are on the

rise. Within less than ten years, incidence rates increased by 36%[2]. Although cryptococcosis

incidence rates are on the decline in North America, this AIDS-defining illness is responsible

for 15% of all AIDS-related deaths world-wide[2,3]. This already dire situation is further

confounded by the emergence of drug resistant yeast species. Patients at risk of developing

invasive candidemia are often prophylactically treated with fluconazole, while the

echinocandins are considered a first line defence strategy[4]. Yet, *C. glabrata*, the most

common non- *albicans Candida* species associated with nosocomial blood stream infections[5],

is intrinsically less susceptible to azole drugs and acquires resistance to echinocandins

rapidly[6]. The rapid global spread of multi-drug resistant *C. auris* has further exacerbated the

threat posed by fungi. *C. auris* was first reported in 2009 in Japan[7]. In 2015, *C. auris* arrived

in Europe causing an outbreak involving 72 patients in a cardio-thoracic hospital in London[8].

C. auris outbreaks have been reported from South Korea, India, Spain, Columbia,

Switzerland, Germany, Israel, Kuwait, and Oman[9]. Most concerningly, up to 25% of *C. auris*

isolates are multi-drug resistant, with some strains being resistant to three of the four drug

classes available for the treatment of systemic candidemia. In addition to the unacceptably

high burden on human health, fungal infections substantially increase health care costs.

Treatment requires extended hospitalisation, resulting in additional costs of up to $45,000 in

adult patients or up to $119,000 in paediatric patients in the case of candidemia[10].

It is thus imperative to investigate fungal virulence and host response mechanisms.

This is traditionally done in mammalian models. The most frequently employed models

include the mouse tail vain infection model for systemic candidemia, the mouse

gastrointestinal infection model of candidemia, the mouse *Candida* vaginitis model[11], the

mouse inhalation model of cryptococcosis[12], the rabbit chronic cryptococcal meningitis

model[13], and the rabbit *Candida* keratitis model[14]. While mammalian models combine a

number of features that make them particularly amenable for the study of fungal diseases,

such as susceptibility, availability of knock-out mutants, and comparable histology to human

disease, using mammals is ethically controversial, economically challenging, and requires

extensive board certifications and documentations. Furthermore, specific applications, such

as the screening of large-scale fungal mutant libraries are not sustainable in mammalian

models.

In an effort to reduce the usage of mammals as model hosts, alternative invertebrate

models have been developed and used in fungal virulence research over the past two decades.

The most commonly employed invertebrate species include the nematode *Caenorhabditis*

elegans, the fly *Drosophila melanogaster*, and larvae of the Greater Wax moth *Galleria*

mellonella. All three species can be easily maintained in the laboratory at a much lower cost

than mice or rabbits and have been successfully used for the study of diverse yeast pathogens,

such as *C. neoformans*[15,16], *C. albicans*[16-18], *C. parapsilosis*[18-20], *C. glabrata*[19,21]. Of note,

invertebrate models differ in their applicability and the best suitable model should be

carefully selected[22]. Unlike mammalian host models, these invertebrates do not have adaptive

immunity but all have components of the innate immune system[23,24], some of which are

conserved with mammals. This includes the Toll-like receptors found in the fly[25] and the

homolog of the MKK3/6 kinase in the nematode[26]. Ironically, it is the Toll-like receptors that

protect flies from infections with *C. neoformans*[27], *C. albicans*[28], and *C. glabrata*[29] and the

MKK3/6 homolog SEK-1 protects the nematode from bacterial invaders[26]. Thus, to increase

susceptibility of flies and nematodes to fungal pathogens, Toll and *sek-1*[30] mutants need to be

used. A key limitation for the study of human pathogens, is the inability of the nematode and

the fly to survive human body temperature. Only *Galleria* can withstand $37°C$[16]. The *Galleria*

genome has been announced very recently[31], yet a detailed analysis, including annotations, is

still missing. Due to their long-standing history as eukaryotic models, well-curated genomes

and genome databases exist for the nematode and the fly. Yet, neither organism allows for the

delivery of an exact inoculum of fungal cells, only *Galleria* can be directly injected with a

defined cell number. Until recently, *Galleria* larvae for research had to be purchased from

fishing shops. Now, UK-based TruLarv is selling research grade larvae while extensive stock

collections exist for the fly and nematode.

An insect model with a long history in research, the Tobacco Hornworm *Manduca*

sexta, has yielded important insights into flight mechanisms, nicotine resistance, hormonal

regulation of development, metamorphosis, antimicrobial defences, and bacterial

pathogenesis. *M. sexta* laboratory stocks have been derived from animals collected in North

Carolina, USA[32] and been maintained in laboratories on both sides of the Atlantic for several

decades. In the wild, *M. sexta* is most commonly encountered in the southern United States,

where it feeds on solanaceous plants and is thus considered a plant pest. *M. sexta*'s research

portfolio includes innate immunity[33], a genome sequence that has been complemented with

tissue-specific transriptomic analyses[34], numerous successful applications of RNAi[35-39], and

protocols for the efficient extraction of hemocytes for down-stream analyses[40]. But despite its

versatility and prominent role in bacterial pathogenesis research, *M. sexta* has yet to be

explored for its suitability as a host model for fungal infections.

Here, we aimed to establish *M. sexta* as a novel model host for the study of fungal

virulence. Inbred animals from the University of Bath's research colony, that has been

established in the 1980s, were tested for their ability to live at $37°C$, their susceptibility to

different yeast species, and the reproducibility of *C. albicans* mutant phenotypes obtained in

mice virulence studies. Indeed, the caterpillars grow at 37˚C while maintaining susceptibility,

specific *C. albicans* mutants are just as attenuated in their virulence in *M. sexta* as they are in

mice, and notably, *M. sexta* are susceptible to the leading yeast pathogens *C. albicans*, *C.*

neoformans, as well as the emerging *C. auris*. To expand *M. sexta's* applicability as a host

model, we developed an infection protocol that permits screening of fungal burden

throughout the course of infection in a single animal and uses weight as a proxy measure for

virulence in addition to survival. *M. sexta* can furthermore be used to test efficacy of common

antifungal drugs. Our results define *M. sexta* characteristics that recommend the caterpillars

as a non-mammalian host model for the study of fungal virulence.

Materials and Methods

Origin of the Bath colony of *Manduca sexta*

The colony has been in continuous culture since 1978 without the addition of animals

from elsewhere. Bath's genetic stock was derived from animals from the Truman-Riddiford

laboratories at the University of Washington in Seattle, USA. Their animals date back to the

ones originally collected in North Carolina in 1976[32].

.

Caterpillar and yeast culture conditions

M. sexta caterpillars were reared to fifth instar under standardised conditions. They

were maintained in 125 ml disposable cups (Sarstedt Ltd., Cat. No. 75.1335), on a wheat

germ-based diet (Appendix 1), at a constant temperature of 25˚C with 50% humidity, and 12

hours of light and dark cycles. Three days prior to infections with fungi, animals were shifted

to a formaldehyde-free diet as the compound is toxic to non-methylotrophic yeast.

For infection assays, yeasts were grown overnight in 50 ml YPD (1% yeast extract,

2% peptone, 2% dextrose) and cells harvested by centrifugation for 3 minutes at 3,000 rpm.

The cell pellet was washed twice with 1x phosphate buffered saline (PBS) and suspended in 5

ml 1x PBS. Cells were counted and numbers adjusted as indicated. *C. albicans* YSD85

(Table 1) cells were heat-inactivated by incubation at 65˚C for 20 minutes. For long-term

storage, yeast isolates were maintained at -80°C in 25% glycerol.

Yeast infections and measurements of fungal burden and drug efficacy

100 µl of washed and number-adjusted yeast suspension were injected into each

caterpillar with a 30G1/2" needle (BD Microlance) and a 1 ml NORM-JECT syringe.

Animals were injected through their distal left proleg. Following injection, each animal's

weight was recorded. Animals were scored for survival and weight once daily for three to

four days post infection. During the course of the experiment, animals were kept on a 12 hour

light and dark cycle at the temperature indicated and on their regular diet.

To measure fungal burden in caterpillar faeces and hemolymph, six animals were

injected with either 1x PBS or 10^6 cells of the wild type YSD89 or the *hog1* mutant strain

YSD883 and kept at 37˚C. On day 1, two animals were selected from each group. These

animals were weighted and their hemolymph and faeces collected daily throughout the course

of infection. To collect hemolymph, animals were first kept on ice for 15 minutes. The 'horn'

was then surface sterilised with 70% ethanol and its top 1-2 mm clipped with a pair of micro

scissors. Hemolymph was collected in a pre-chilled 1.5 ml Eppendorf tube and cooled

immediately to reduce polymerisation and melanisation. One faecal pellet was collected daily

with sterile forceps, weighted and suspended in 500 µl 1x PBS. Prior to diluting, the mixture

was thoroughly vortexed for 10 seconds, and centrifuged for 5 seconds using a table top

centrifuge to separate faecal matter. To quantify fungal burden, hemolymph and faecal

samples were plated either directly onto YPD-agar with Kanamycin 50 µg/ml or in ten-fold

serial dilutions. Agar plates were incubated at 30˚C for 48 hours and colonies counted.

To assess the efficacy of commonly used antifungal drugs, animals were infected with

10^7 cells of YSD85 or PBS and treated with increasing doses of fluconazole and caspofungin

(Sigma Aldrich, Inc.) as indicated. Drugs were injected with an ethanol-sterilized Hamilton

syringe in a total volume of 10 µl per animal, 30 minutes post-infection. Caterpillars were

weighted and scored for survival on the day of injection and the following three days.

Statistical analyses

Survival plots were made using the **survminer** R package (https://CRAN.R-

project.org/package=survminer), and differences were evaluated using the Kaplan-Meier

method. Weight and fungal burden were plotted using ggplot2[41] and weight differences were

evaluated using linear models with day post-inoculation and the interaction between

treatment and dpi as fixed effects and individual as a random effect using nlme

(https://CRAN.R-project.org/package=nlme). All analyses were done using RStudio version

1.1.442.

Results

We first aimed to determine if *M. sexta* fifth instar caterpillars, reared and maintained

at standard conditions (Fig. 1a), are susceptible to *Candida albicans*. To do so, groups of ten

animals were infected with increasing doses of the widely used *C. albicans* laboratory strains

SC5314 and SN95[42]. Animals were scored daily for survival for three consecutive days while

being maintained at 25˚C. Dead animals differ from live ones in that their bodies go limp and

turn grey-green in colour, which is in stark contrast to the vivid turquoise of live animals

(Fig. 1b). Indeed, caterpillars that were infected with *C. albicans* succumbed to the yeast in a

dose-dependent manner. Both *C. albicans* strains killed *M. sexta* caterpillars efficiently at

inocula of 10^6 or 10^7 cells per animal (Fig. 1c). To determine if survival measures in

caterpillars are comparable to those obtained in the current gold standard, the murine model

of systemic candidemia, we tested *C. albicans* mutants with published phenotypes of either

attenuated virulence, such as the *hog1Δ/Δ*[43] and *ahr1Δ/Δ*[44] mutants, or wild-type levels of

virulence, such as *cka2Δ/Δ*[45]. Cross-species virulence levels are comparable for Hog1, which

is as essential for virulence in caterpillars as it is in mice. Cka2 is not required to establish

systemic infections in mammals but is in caterpillars. Ahr1, while required for virulence in

mammals, appears to be dispensable for virulence in caterpillars (Fig. 1d).

Given the importance of temperature for fungal virulence, we aimed to determine if

M. sexta retained their susceptibility to *C. albicans* at human body temperature of 37˚C.

Temperature itself does not affect caterpillar survival or development (Fig. S1) but animals

are ten times more susceptible to infections with *C. albicans* at 37˚C than they are at 25˚C

(Fig. 2a). At 37˚C, 10^6 *C. albicans* cells per animal lead to 100% mortality on day 4, while

10^7 cells are required for the same outcome at 25˚C (Fig. 1b). To exclude the possibility that

mortality is due to starvation rather than the outcome of a host-pathogen interaction, we

infected caterpillars with live and heat-killed *C. albicans* wild-type cells at 37˚. Only live

cells, but not heat-killed *Candida* cells, kill caterpillars suggesting that killing is not due to

nutritional limitations (Fig. S2). Demonstrating susceptibility of *M. sexta* caterpillars to

Candida albicans supports their suitability as an alternative host model for the study of

fungal virulence but also highlights the need for additional measures of fungal virulence. To

add granularity to fungal virulence data collected from *M. sexta*, we complemented measures

of survival with quantifications of weight and fungal burden throughout the course of

infection. To collect weight data, caterpillars were weighted prior to infection and then daily

throughout the course of infection. Weight gain in animals infected with a low dose of 10^4

cells did not significantly differ from those injected with 1x PBS but caterpillars infected

with10^5 cells per animal exhibited significant weight loss. Too few animals survived

infection with 10^6 cells to allow for a meaningful comparison (Fig. 2b). After establishing

susceptibility of *M. sexta* to *C. albicans* at 37˚C, we aimed to validate the attenuated

virulence phenotype of the *hog1Δ/Δ* mutant strain. To test this, caterpillars were infected with

10^6 cells per animal with the wild type strain, the *hog1Δ/Δ* mutant, and the complemented

strain *hog1/hog1::HOG1* and compared to the control group injected with 1x PBS. The

hog1Δ/Δ mutant is less virulent than the wild type (Fig. 2c). Animals infected with the

hog1Δ/Δ mutant strain JC 50 gained significantly more weight than those infected with the

wild type RM1000, while infection with the complemented strain JC52 leads to a comparable

lack in weight gain (Fig. 2d).

To further expand the applicability of *M. sexta* caterpillars as a host model for fungal

infections, fungal burden in the hemolymph and faeces was quantified daily throughout the

course of infection. Since neither the collection of hemolymph nor that of faeces necessitates

killing of the animal, data could be collected daily throughout the infection for the same

caterpillar. Animals infected with the wild type and the *hog1Δ/Δ* mutant strain YSD883 were

compared to control animals injected with 1x PBS only. Analysing fungal burden by counting

colony-forming units in faeces and hemolymph in two animals per group, revealed an

increase in CFU counts in the animals infected with the wild type but not the *hog1Δ/Δ* mutant

or the PBS-control group (Figs. 3a, b). While no yeasts were detected in either group one day

post infection (p.i), CFUs in faeces and hemolymph are detectable two days p.i. in animals

injected with the wild type. Three days p.i., the CFU count in the wild-type group increase

while CFUs are detected for the first time in the *hog1Δ/Δ* group. In keeping with the increase

in fungal burden, weight gain in the wild-type infected group was reduced, while *hog1Δ/Δ*

infected animals gained weight at a similar rate to the control animals (Fig. 3c). Thus,

measurements of fungal burden, that can be obtained from the same animal throughout the

course of infection, provide a valuable parameter for the study of fungal virulence.

The currently available armamentarium of antifungal drugs is limited and often

lacking in efficacy. Studying drug efficacy and drug mode of action in the host are thus

pertinent to the much needed development of novel antifungal drugs and further

characterisation of existing compounds. To establish suitability of *M. sexta* as a model for

drug efficacy testing, we recorded survival and weight of animals infected with the *C.*

albicans wild-type strain SC5314 and treated with increasing doses of two common

antifungals, fluconazole and caspofungin (Fig.4). Treatment with fluconazole or caspofungin

resulted in overall improved survival and weight gain. Pair-wise comparisons between

different drug concentrations with the untreated control animals yielded no statistical

significance for the fluconazole-treated group of caterpillars but those being treated with 2

mg/kg and 4 mg/kg of caspofungin survived significantly better than those without treatment

or those that only received 1 mg/kg of caspofungin (Table 2).

While *C. albicans* undoubtedly is a leading fungal pathogen of humans, clinical

manifestations of fungal infections are not limited to *C. albicans*. To broaden *M. sexta*'s

applicability, we sought to determine the caterpillars' susceptibility to other fungal

pathogens. To this end, animals were infected with *C. neoformans*, *C. auris*, and *C. glabrata*

in addition to *C. albicans*. Type strains of *Saccharomyces cerevisiae*, the baker's or brewer's

yeast and *Metschnikowia pulcherrima*, a yeast inhabiting fruits and flowers[46] served as

reference points for attenuated virulence infections. Animals were infected with increasing

doses of yeast cells starting at 10^5 cells per animal and up to 10^9 cells per animal. Groups of

ten animals per yeast cell dose and species were then screened for survival and weight daily

(Figs. 5 and 6). Notably, only infections with pathogenic yeast species caused reduced

survival of caterpillars (Fig. 5). Infections with *S. cerevisiae* or *M. pulcherrima* did not affect

survival of caterpillars. Comparing survival amongst the pathogenic yeast species revealed *C.*

albicans to be the most severe. 10^7 *C. albicans* cells per animal result in 100% killing within

in 24 hours. 10^9 *C. auris* cells were required to kill all animals within four days and the same

dose of *C. glabrata* resulted in 75% killing. 10^8 *C. neoformans* cells were required to achieve

50% killing within four days, comparable to survival of animals infected with *C. glabrata*. It

should be noted that due to high viscosity of the cell suspension, we could not test higher

concentrations than the ones stated. In addition to assessing survival, each surviving

caterpillar's weight was recorded daily. Interestingly, while infections with *S. cerevisiae* or

M. pulcherrima did not kill caterpillars, infected animals displayed reduced weight gain (Fig.

6). Infections with *C. albicans* resulted in severely reduced weight gain, even at the lowest

yeast dose tested. Animals infected with *C. auris* and *C. glabrata* responded in a dose-

dependent manner, the higher the yeast dose, the more dramatic the weight loss. Notably,

animals infected with the lowest dose of *C. neoformans* grew better than PBS-injected control

animals. Increasing the yeast dosage, however, resulted in significantly reduced weight gain.

This bi-phasic pattern, resembling hormesis[47] in which a low dose of an environmental agent

is beneficial, while a high dose is toxic, could be due to *C. neoformans'* immunogenic

capsule[48]. Caterpillars displaying variable degrees of susceptibility to different yeast species,

with weight providing an additional measure of host damage, suggests that this host model is

broadly applicable for the study of fungal virulence.

Discussion

M. sexta caterpillars are naturally susceptible to the leading human yeast pathogens.

Unlike other host models, they permit assessment of fungal burden daily throughout the

course of infection in a single animal by either collecting faeces or hemolymph. *M. sexta* can

be maintained at 37˚C, is large enough to be injected with a specified yeast inoculum and for

weight to be a reliable measure of virulence. *C. albicans* mutant virulence phenotypes found

in mice can be replicated and yeast inocula required to elicit a response in caterpillars are

comparable to those used in the murine model. Additionally, the caterpillars permit study of

antifungal drug efficacy. These parameters commend this invertebrate as a novel host model

for the study of fungal virulence.

This new model system allows for fungal burden to be monitored throughout the

course of infection in a single animal via CFU count. This is unlike any other experimental

system, where fungal burden is either an endpoint measure in the mouse kidney, in

homogenised was moth larvae[49],nematodes[50], flies[51], or requires genetically modified

fluorescent yeast strains for microscopic imaging and analyses[52]. While we did detect very

low CFU counts in the PBS control 3 days post infection, we consider this spurious finding

due to cross contamination as preliminary experiments of plating contents of hemolymph and

faeces of naïve animals did not detect any yeast growth (data not shown). As a consequence,

we amended to protocol to include changing gloves when handling animals of different

treatment groups.

When reviewing the weight data collected as part of our study, we noticed two

interesting aspects. First, while *S. cerevisiae* and *M. pulcherrima* do not affect caterpillar

survival, infections with either species led to reduced weight gain. The dichotomy between

survival and weight observed here, further emphasizes that fungal virulence comprises more

than a measure of survival. It appears that *M. sexta* would allow to discrimination between

disease (weight) and death (survival) quantitatively adding further granularity to measuring

fungal virulence. Secondly, infections with *C. neoformans* resulted in increased caterpillar

weight gain at a low yeast dose but reduced weight gain at higher doses. This pattern

resembles the concept of hormesis often deployed by toxicologists to describe the response to

toxins, where exposure to a low dose is beneficial while a higher dose results toxicity due to

overcompensation in response to disruption of homeostasis[53]. While tissue-specific hormetic

responses have been described in flies, where a virus-acquired cytokine relays ageing[54], and

examples of abiotic factors or signalling molecules affecting peas and aphid infestation[55],

immunity in plants[56,57], life-span in malaria-transmitting *Anopheles*[58], and larval development

in Black Cutworm[59], this example here could be the first involving a eukaryotic host-

pathogen relationship. The underlying factor remains to be elucidated but *C. neoformans'*

highly immunogenic capsule seems to be an excellent candidate[48].

Heat-killed *C. albicans* cells appear to be entirely non-pathogenic in *M. sexta*

caterpillars as neither survival nor weight are affected by innoculation with heat-killed yeast

cells. The lack of mortality in response to innoculation with heat-killed yeast cells indicates

that yeast viability and proliferation are required for pathogenesis and excludes the possibility

of death due to an allergic reaction in response to a large number of fungal cells. This appears

to differ from the responses of other host models to fungal pathogens. While susceptibility

was reduced, but still measurable, in *G. mellonella*[60] and the two-spotted cricket[61], heat-killed

C. albicans cells elicited 100% mortality in a sepsis-like murine model[62]. In mice, serum

levels of β-(1,3)-glucan levels were elevated in animals injected with heat-killed yeast cells

when compared to those infected with live cells. Indeed, heat inactivation leads to increased

exposure of β-(1,3)-glucan on the *C. albicans* cell surface[63] and β-(1,3)-glucan activates the

innate immune response in invertebrates and mammals[64]. Given the complexity of receptors

involved in recognition of fungal invaders[65] and the lack of a response in *M. sexta* and other

invertebrates indicates that they may not host all or identical receptors as mammals.

We showed that *Manduca sexta* caterpillars expand the repertoire of invertebrate

models for the study of fungal disease. It combines the unique advantage of allowing

measurement of fungal burden with standard quantifications of survival and weight.

Although, *M. sexta* genomic and transcriptomic analyses are currently still in their infancy,

we would expect that the Tobacco Hornworm's long history of being an invaluable model for

diverse facets of biology will lead to reliable tools in combination with genetic tractability

and protocols establishing the yeasts' fate inside the caterpillar.

Acknowledgements

We would like to thank Ewan Basterfield and Chris Apark for expert advice and

technical assistance in preparing *M. sexta* caterpillars. Thanks to all the laboratories who

kindly shared their strains with us. This work was supported by an ERC Marie Curie Career

Integration Grant to SD and undergraduate and postgraduate research funding from the

University of Bath.

Disclosure of interest

The authors report no interest of conflict.

References

1. Brown GD, Denning DW, Gow NAR, Levitz SM, Netea MG, White TC. Hidden killers: Human fungal infections. Sci Transl Med 2012; 4:165rv13.

2. Pfaller MA, Diekema DJ. Epidemiology of invasive mycoses in North America. Crit Rev Microbiol 2010; 36:1–53.

3. Nyazika TK, Tatuene JK, Kenfak-Foguena A, Verweij PE, Meis JF, Robertson VJ, Hagen F. Epidemiology and aetiologies of cryptococcal meningitis in Africa, 1950–2017: Protocol for a systematic review. BMJ Open 2018; 8:e020654–4.

4. Pappas PG, Kauffman CA, Andes D, Benjamin DK Jr, Calandra TF, Edwards JE Jr, Filler SG, Fisher JF, Kullberg B-J, Ostrosky Zeichner L, et al. Clinical practice guidelines for the management of candidiasis: 2009 update by the Infectious Diseases Society of America. Clin Inf Dis 2009; 48:503–35.

5. Pfaller MA, Diekema DJ. Epidemiology of invasive candidiasis: A persistent public health problem. Clin Microbiol Rev 2007; 20:133–63.

6. Jensen RH, Johansen HK, Søes LM, Lemming LE, Rosenvinge FS, Nielsen L, Olesen B, Kristensen L, Dzajic E, Astvad KMT, et al. Posttreatment antifungal resistance among colonizing *Candida* isolates in candidemia patients: Results from a systematic multicenter study. Antimicrob Agents Chemother 2016; 60:1500–8.

7. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. Microbiol Immunol 2009; 53:41–4.

8. Schelenz S, Hagen F, Rhodes JL, Abdolrasouli A, Chowdhary A, Hall A, Ryan L, Shackleton J, Trimlett R, Meis JF, et al. First hospital outbreak of the globally emerging *Candida auris* in a European hospital. Antimicrob Resist Infect Control 2016; 5:1–7.

9. Bidaud AL, Chowdhary A, Dannaoui E. *Candida auris*: An emerging drug resistant yeast – A mini-review. J Mycol Med 2018; 28:568–73.

10. Zaoutis TE, Argon J, Chu J, Berlin JA, Walsh TJ, Feudtner C. The epidemiology and attributable outcomes of candidemia in adults and children hospitalized in the United States: A propensity analysis. Clin Infect Dis 2005; 41:1232–9.

11. Segal E, Frenkel M. Experimental *in vivo* models of candidiasis. JoF 2018; 4:21–10.

12. Zaragoza O, Alvarez M, Telzak A, Rivera J, Casadevall A. The relative susceptibility of mouse strains to pulmonary *Cryptococcus neoformans* infection is associated with pleiotropic differences in the immune response. Infect Immun 2007; 75:2729–39.

13. Perfect JR, Lang SD, Durack DT. Chronic cryptococcal meningitis: A new

395   experimental model in rabbits. Am J Pathol 1980; 101:177–94.

396   14.   Ray WA, O'Day DM, Head WS, Robinson RD. Variability in isolate recovery
397   rates from multiple and single breeds of outbred pigmented rabbits in an
398   experimental model of *Candida* keratitis. Curr Eye Res 2009; 3:949–53.

399   15.   Mylonakis E, Ausubel FM, Perfect JR, Heitman J, Calderwood SB. Killing of
400   *Caenorhabditis elegans* by *Cryptococcus neoformans* as a model of yeast
401   pathogenesis. PNAS 2002; 99:15675–80.

402   16.   Fuchs BB, O'Brien E, Khoury El JB, Mylonakis E. Methods for using *Galleria*
403   *mellonella* as a model host to study fungal pathogenesis. Virulence 2014; 1:475–
404   82.

405   17.   Pukkila-Worley R, Ausubel FM, Mylonakis E. *Candida albicans* infection of
406   *Caenorhabditis elegans* induces antifungal immune defenses. PLoS Pathog
407   2011; 7:e1002074.

408   18.   Chamilos G, Lionakis MS, Lewis RE, Lopez-Ribot JL, Saville SP, Albert ND,
409   Halder G, Kontoyiannis DP. *Drosophila melanogaster* as a facile model for
410   large-scale studies of virulence mechanisms and antifungal drug efficacy in
411   *Candida* species. J Infect Dis 2006; 193:1014–22.

412   19.   Ortega-Riveros M, De-la-Pinta I, Marcos-Arias C, Ezpeleta G, Quindós G, Eraso
413   E. Usefulness of the non-conventional *Caenorhabditis elegans* model to assess
414   *Candida* virulence. Mycopathologia 2017; 182:785–95.

415   20.   Gago S, García-Rodas R, Cuesta I, Mellado E, Alastruey-Izquierdo A. *Candida*
416   *parapsilosis*, *Candida orthopsilosis*, and *Candida metapsilosis* virulence in the
417   non-conventional host *Galleria mellonella*. Virulence 2014; 5:278–85.

418   21.   Ames L, Duxbury S, Pawlowska B, Ho H-L, Haynes K, Bates S. *Galleria*
419   *mellonella* as a host model to study *Candida glabrata* virulence and antifungal
420   efficacy. Virulence 2017; 8:1–9.

421   22.   Desalermos A, Fuchs BB, Mylonakis E. Selecting an invertebrate model host for
422   the study of fungal pathogenesis. PLoS Pathog 2012; 8:e1002451.

423   23.   Mylonakis E, Aballay A. Worms and flies as genetically tractable animal models
424   to study host-pathogen interactions. Infect Immun 2005; 73:3833–41.

425   24.   Vogel H, Altincicek B, Glöckner G, Vilcinskas A. A comprehensive
426   transcriptome and immune- gene repertoire of the lepidopteran model host
427   *Galleria mellonella*. BMC Genomics 2011; 12:308.

428   25.   Lemaitre B, Nicolas E, Michaut L, Reichhart JM, Hoffmann JA. The
429   dorsoventral regulatory gene cassette spätzle/Toll/cactus controls the potent
430   antifungal response in *Drosophila* adults. Cell 1996; 86:973–83.

431   26.   Kim DH, Feinbaum R, Alloing G, Emerson FE, Garsin DA, Inoue H, Tanaka-
432   Hino M, Hisamoto N, Matsumoto K, Tan M-W, et al. A conserved p38 MAP
433   kinase pathway in *Caenorhabditis elegans* innate immunity. Science 2002;

434   297:623–6.

27.   Apidianakis Y, Rahme LG, Heitman J, Ausubel FM, Calderwood SB, Mylonakis E. Challenge of *Drosophila melanogaster* with *Cryptococcus neoformans* and role of the innate immune response. Eukaryot Cell 2004; 3:413–9.

28.   Alarco A-M, Marcil A, Chen J, Suter B, Thomas D, Whiteway M. Immune-deficient *Drosophila melanogaster*: A model for the innate immune response to human fungal pathogens. J Immunol 2004; 172:5622–8.

29.   Quintin J, Asmar J, Matskevich AA, Lafarge MC, Ferrandon D. The *Drosophila* Toll pathway controls but does not clear *Candida glabrata* infections. J Immunol 2013; 190:2818–27.

30.   Breger J, Fuchs BB, Aperis G, Moy TI, Ausubel FM, Mylonakis E. Antifungal chemical compounds identified using a *C. elegans* pathogenicity assay. PLoS Pathog 2007; 3:e18.

31.   Lange A, Beier S, Huson DH, Parusel R, Iglauer F, Frick J-S. Genome sequence of *Galleria mellonella* (Greater Wax Moth). Genome Announc 2018; 6:61–2.

32.   Flowers RW, Entomologist RYF, 1982. Feeding on non-host plants by the Tobacco Hornworm (*Manduca sexta* (Lepidoptera: Sphingidae). JSTOR 65:523–30.

33.   Kanost MR, Jiang H, Yu X-Q. Innate immune responses of a lepidopteran insect, *Manduca sexta*. Immunol Rev 2004; 198:97–105.

34.   Kanost MR, Arrese EL, Cao X, Chen Y-R, Chellapilla S, Goldsmith MR, Grosse-Wilde E, Heckel DG, Herndon N, Jiang H, et al. Multifaceted biological insights from a draft genome sequence of the Tobacco Hornworm moth, *Manduca sexta*. Insect Biochem Mol Biol 2016; 76:118–47.

35.   Flores-Escobar B, Rodríguez-Magadan H, Bravo A, Soberón M, Gómez I. Differential role of *Manduca sexta* aminopeptidase-N and alkaline phosphatase in the mode of action of Cry1Aa, Cry1Ab, and Cry1Ac toxins from *Bacillus thuringiensis*. Appl Environ Microbiol 2013; 79:4543–50.

36.   Burke WG, Kaplanoglu E, Kolotilin I, Menassa R, Donly C. RNA interference in the Tobacco Hornworm, *Manduca sexta*, using plastid-encoded long double-stranded RNA. Front Plant Sci 2019; 10:313.

37.   Eleftherianos I, Millichap PJ, ffrench-Constant RH, Reynolds SE. RNAi suppression of recognition protein mediated immune responses in the tobacco hornworm *Manduca sexta* causes increased susceptibility to the insect pathogen *Photorhabdus*. Dev Comp Immunol 2006; 30:1099–107.

38.   Eleftherianos I, Gökçen F, Felföldi G, Millichap PJ, Trenczek TE, ffrench-Constant RH, Reynolds SE. The immunoglobulin family protein Hemolin mediates cellular immune responses to bacteria in the insect *Manduca sexta*. Cell Microbiol 2007; 9:1137–47.

39. Kumar P, Pandit SS, Baldwin IT. Tobacco Rattle Virus vector: A rapid and
transient means of silencing *Manduca sexta* genes by plant mediated RNA
interference. PLoS ONE 2012; 7:e31347–10.

40. Stoepler TM, Castillo JC, Lill JT, Eleftherianos I. A simple protocol for
extracting hemocytes from wild caterpillars. JoVE 2012; :1–6.

41. Wickham H. ggplot2. New York, NY: Springer-Verlag New York; 2009.

42. Noble SM, Johnson AD. Strains and strategies for large-scale gene deletion
studies of the diploid human fungal pathogen *Candida albicans*. Eukaryot Cell
2005; 4:298–309.

43. Alonso-Monge R, Navarro-García F, Molero G, Diez-Orejas R, Gustin M, Pla J,
Sánchez M, Nombela C. Role of the mitogen-activated protein kinase Hog1p in
morphogenesis and virulence of *Candida albicans*. J Bacteriol 1999; 181:3058–
68.

44. Askew C, Sellam A, Epp E, Mallick J, Hogues H, Mullick A, Nantel A,
Whiteway M. The zinc cluster transcription factor Ahr1p directs Mcm1p
regulation of *Candida albicans* adhesion. Mol Microbiol 2010; 79:940–53.

45. Chiang LY, Sheppard DC, Bruno VM, Edwards JE, Filler SG. *Candida albicans*
protein kinase CK2 governs virulence during oropharyngeal candidiasis. Cell
Microbiol 2007; 9:233–45.

46. Vadkertiová R, Molnárová J, Vránová D, Sláviková E. Yeasts and yeast-like
organisms associated with fruits and blossoms of different fruit trees. Can J
Microbiol 2012; 58:1344–52.

47. Mattson MP. Hormesis defined. Ageing Res Rev 2008; 7:1–7.

48. Casadevall A, Coelho C, Cordero RJB, Dragotakes Q, Jung E, Vij R, P W. The
capsule of *Cryptococcus neoformans*. Virulence 2018; 0:1–10.

49. Lu M, Yu C, Cui X, Shi J, Yuan L, Sun S. Gentamicin synergises with azoles
against drug-resistant *Candida albicans*. Int J Antimicrob Ag 2018; 51:107–14.

50. Sun L, Liao K, Wang D. Effects of magnolol and honokiol on adhesion, yeast-
hyphal transition, and formation of biofilm by *Candida albicans*. PLoS ONE
2015; 10:e0117695–20.

51. Wurster S, Bandi A, Beyda ND, Albert ND, Raman NM, Raad II, Kontoyiannis
DP. *Drosophila melanogaster* as a model to study virulence and azole treatment
of the emerging pathogen *Candida auris*. J Antimicrob Chemother 2019;
74:1904–10.

52. Sun L, Zhi L, Shakoor S, Liao K, Wang D. microRNAs involved in the control
of innate immunity in *Candida* infected *Caenorhabditis elegans*. Nature 2016;
:1–13.

53. Calabrese1 EJ, Baldwin LA. Hormesis: A generalizable and unifying hypothesis.

Cr Rev Toxicol 2008; 31:353–424.

54.   Mlih M, Khericha M, Birdwell C, West AP, Karpac J. A virus-acquired host
cytokine controls systemic aging by antagonizing apoptosis. PLoS Biol 2018;
16:e2005796–30.

55.   Woźniak A, Drzewiecka K, Kęsy J, Marczak Ł, Narożna D, Grobela M, Motała
R, Bocianowski J, Morkunas I. The influence of lead on generation of signalling
molecules and accumulation of flavonoids in pea seedlings in response to pea
aphid infestation. Molecules 2017; 22:1404–30.

56.   Bektas Y, Rodriguez-Salus M, Schroeder M, Gomez A, Kaloshian I, Eulgem T.
The synthetic elicitor DPMP (2,4-dichloro-6-{(E)-[(3-
methoxyphenyl)imino]methyl}phenol) triggers strong immunity in *Arabidopsis*
thaliana and tomato. Nature 2016; :1–16.

57.   Rodriguez-Salus M, Bektas Y, Schroeder M, Knoth C, Vu T, Roberts P,
Kaloshian I, Eulgem T. The synthetic elicitor 2-(5-bromo-2-hydroxy-phenyl)-
thiazolidine-4-carboxylic acid links plant immunity to hormesis. Plant Physiol
2015; 170:444–58.

58.   Drexler AL, Pietri JE, Pakpour N, Hauck E, Wang B, Glennon EKK, Georgis M,
Riehle MA, Luckhart S. Human IGF1 regulates midgut oxidative stress and
epithelial homeostasis to balance lifespan and *Plasmodium falciparum* resistance
in *Anopheles stephensi*. PLoS Pathog 2014; 10:e1004231–14.

59.   Kullik SA, Sears MK, Schaafsma AW. Sublethal effects of Cry 1F Bt corn and
clothianidin on Black Cutworm (Lepidoptera: Noctuidae) larval development.
Jnl Econ Entom 2011; 104:484–93.

60.   Cotter G, Doyle S, Kavanagh K. Development of an insect model for the *in vivo*
pathogenicity testing of yeasts. FEMS Immunol Med Microbiol 2000; 27:163–9.

61.   Kochi Y, Matsumoto Y, Sekimizu K, Kaito C. Two-spotted cricket as an animal
infection model of human pathogenic fungi. DD&T 2017; 11:259–66.

62.   Panpetch W, Somboonna N, Bulan DE, Issara-Amphorn J, Finkelman M,
Worasilchai N, Chindamporn A, Palaga T, Tumwasorn S, Leelahavanichkul A.
Oral administration of live- or heat-killed *Candida albicans* worsened cecal
ligation and puncture sepsis in a murine model possibly due to an increased
serum $(1{\rightarrow}3)$-β-D-glucan. PLoS ONE 2017; 12:e0181439–15.

63.   Wellington M, Dolan K, Krysan DJ. Live *Candida albicans* suppresses
production of reactive oxygen species in phagocytes. Infect Immun 2008;
77:405–13.

64.   Means TK, Mylonakis E, Tampakakis E, Colvin RA, Seung E, Puckett L, Tai
MF, Stewart CR, Pukkila-Worley R, Hickman SE, et al. Evolutionarily
conserved recognition and innate immunity to fungal pathogens by the scavenger
receptors SCARF1 and CD36. J Exp Med 2009; 206:637–53.

65.   Netea MG, Brown GD, Kullberg B-J, Gow NAR. An integrated model of the

recognition of *Candida albicans* by the innate immune system. Nat Rev Micro
2008; 6:67–78.

66. Blankenship JR, Heitman J. Calcineurin is required for *Candida albicans* to
survive calcium stress in serum. Infect Immun 2005; 73:5767–74.

67. Bruno VM. Regulation of azole drug susceptibility by *Candida albicans* protein
kinase CK2. Mol Microbiol 2005; 56:559–73.

68. Smith DA, Nicholls S, Morgan BA, Brown AJP, Quinn J. A conserved stress-
activated protein kinase regulates a core stress response in the human pathogen
Candida albicans. Mol Biol Cell 2004; 15:4179–90.

69. Sniegowski P, Dombrowski P. *Saccharomyces cerevisiae* and *Saccharomyces
paradoxus* coexist in a natural woodland site in North America and display
different levels of reproductive isolation. FEMS Yeast Res 2002; 1:299–306.

70. Pitt JI, Mycologia MM, 1968. Sporulation in *Candida pulcherrima, Candida
reukaufii* and *Chlamydozyma* species: Their relationships with *Metschnikowia*.
Mycologia 1968; 60:663–685.

71. Dujon B, Sherman D, Fischer G, Durrens P, Casaregola S, Lafontaine I, De
Montigny J, Marck C, Neuvéglise C, Talla E, et al. Genome evolution in yeasts.
Nature 2004; 430:35–44.

72. Ben-Ami R, Berman J, Novikov A, Bash E, Shachor-Meyouhas Y, Zakin S,
Maor Y, Tarabia J, Schechner V, Adler A, et al. Multidrug-Resistant *Candida
haemulonii* and *C. auris*, Tel Aviv, Israel. Emerg Infect Dis 2017; 23:1–9.

73. Wong B, Perfect JR, Beggs S, Wright KA. Production of the hexitol D-mannitol
by *Cryptococcus neoformans in vitro* and in rabbits with experimental
meningitis. Infect Immun 1990; 58:1664–70.

575
576

577 **Appendices**

578 **Appendix 1 – Food preparation**

Premix		Diet cake	
Amount	Ingredient	Amount	Ingredient
2,700 g	Wheatgerm	336 g	Premix
1,260 g	Casein	1,770 ml	Distilled water
1,080 g	Sucrose	22.5 g	Agar
540 g	Dried active yeast	4 ml	Corn oil
360 g	Wesson's Salt	4 ml	Linseed oil
36 g	Choline chloride	8 ml	4% Formaldehyde
72 g	Cholesterol	0.2 g	Chlorotetracycline
36 g	Methyl paraben	0.2 g	Vanderssant vitamins
54 g	Sorbic acid	8 g	Ascorbic acid

For the premix, deactivate the yeast by microwaving for 5 minutes on low power
before mixing all components thoroughly. Store in a cool, dry place.

For the diet cake, heat up 650 ml of water on a hot plate while melting the agar in 1 l
of water by microwaving. Combine the agar with the pre-warmed water, 336 g of premix,
formaldehyde and oils and mix thoroughly using a stand mixer. Dissolve vitamins, antibiotic,
and ascorbic acid in the remaining 30 ml of water and add to mixing bowl once the content
has cooled below 50°C to prevent inactivation of vitamins and antibiotic. Line a large ice
cube tray with sterile aluminium foil (sterilise by spraying with 70% ethanol) and pour
mixture into tray. Let the diet mix set for about 1.5 hours, wrap tightly in aluminium foil and
store at 4°C. Keeps for 3 weeks.

Tables

Table 1: Yeast strains used in this study.

Strain ID	Strain Name	Species	Source
YSD89	SN95	*C. albicans*	
YSD85	SC5314	*C. albicans*	
YSD883	hog1Δ/Δ	*C. albicans*	This study
YSD87	BWP17	*C. albicans*	[66]
YSD190	VIC84 (cka2Δ/Δ)	*C. albicans*	[67]
YSD192	VIC93 (cka2/cka2::CKA2)	*C. albicans*	[67]
YSD302	CAS12 (ahr1Δ/Δ)	*C. albicans*	[44]
YSD303	CAS13 (ahr1/ahr1::AHR1)	*C. albicans*	[44]
YSD304	RM1000	*C. albicans*	[68]
YSD305	JC52 (hog1/hog1::HOG1)	*C. albicans*	[68]
YSD306	JC50 (hog1Δ/Δ)	*C. albicans*	[68]
YSD790	YPS143	*S. cerevisiae*	[69]
YSD1448	NCYC2580	*M. pulcherrima*	[70]
YSD465	2001	*C. glabrata*	[71]
YSD1454	TA004-14	*C. auris*	[72]
YSD1028	H99	*C. neoformans*	[73]

595 **Table 2:** Statistical significance values of survival of animals treated with antifungal drugs
596 compared to those not receiving treatment.
597

Fluconazole		Caspofungin	
Concentration	p-value	Concentration	p-value
1 mg/kg	0.54	1 mg/kg	0.8
4 mg/kg	0.63	2 mg/kg	0.015
16 mg/kg	0.68	4 mg/kg	0.0022
32 mg/kg	0.27		
64 mg/kg	0.13		

598
599

Figure Captions

Figure 1: *M. sexta* caterpillars are susceptible to infections with *C. albicans*. a) *M. sexta* fifth instar caterpillar prior to injection weighing ~2 g. **b)** 24 hours post injection, the dead animal on the left has lost colour and rigor compared to the live animal on the right. **c)** Survival curves of animals infected with *C. albicans* SN95 or SC5314. Killing occurs in a dose-dependent manner at 10^6 and 10^7 yeast cells per animal. **d)** Survival curves of animals infected with *C. albicans* mutants with attenuated virulence phenotypes in mice or epithelial cell models. The *hog1Δ/Δ* and *cka2Δ/Δ* mutants exhibit attenuated virulence, while virulence of the *ahr1Δ/Δ* mutant is comparable to the wild type. The *hog1Δ/Δ* mutant differs significantly from the wild type (p=0.0001). The complemented strain *hog1/hog1::HOG1* kills *M. sexta* at a level comparable to that of the wild type strain (p=0.21). The *cka2Δ/Δ* mutant is significantly less virulent than the wild type (p=0.00021), while the complemented strain *cka2/cka2::CKA2* is not (p=0.054). Virulence of the *ahr1Δ/Δ* mutant and the complemented strain *ahr1/ahr1::AHR1* does not differ significantly from the wild type (*ahr1Δ/Δ* p=0.72; *ahr1/ahr1::AHR1* p=0.38).

Figure 2: Elevated temperatures increase susceptibility of *M. sexta* to *C. albicans*. a) *M. sexta* caterpillars succumb to infection with the laboratory strain SC5314 in a dose-dependent manner at 37˚C but less inoculum is required than for infections at 25˚C. **b)** The weight of caterpillars infected with 10^4 cells per animal was comparable to that of caterpillars injected with 1x PBS, while animals infected with 10^5 cells showed a significant reduction in weight. **c)** Attenuated virulence of the *hog1Δ/Δ* mutant is retained at 37˚C. **d)** Animals infected with the mutant strain gain significantly more weight than those infected with the wild type strain. Infection with the complemented strain results in a comparable lack of weight gain.

Figure 3: Fungal burden in faeces and hemolymph increases throughout the course of infection. Colony forming units per gram faeces **(a)** and 100 μl hemolymph **(b)** in the *hog1Δ/Δ* mutant (YSD883), the wild type (SN95) and the control animals. Counts increase over time in the wild-type infected animals but not the ones injected with the *hog1Δ/Δ* mutant or PBS. **c)** Weight increases in animals infected with *hog1Δ/Δ* comparable to those injected with PBS, while animals injected with the wild type *C. albicans* strain experience reduced weight gain. The peak and drop in weight observed in the *hog1Δ/Δ* and PBS groups, marked with a ⊙ is coinciding with the onset of pupation. This 'pupation drop' is due to the animals refraining from food upon entering the early stages of pupation.

Figure 4: Antifungal efficacy testing of fluconazole and caspofungin. Groups of ten *C. albicans* infected caterpillars were treated with increasing doses of antifungal drug. a) Survival improves upon treatment with fluconazole, while weight remains largely stagnant in surviving animals (b). c) Caspofungin treatment has a positive effect on *M. sexta* survival and weight (d). Weight data were not analysed for significance due to the lack of animals in the no treatment group.

Figure 5: Caterpillars are susceptible to common yeast pathogens. Groups of ten animals were infected with increasing doses of yeasts and survival was recorded daily. Caterpillars are not susceptible to *S. cerevisiae* or *M. pulcherrima* but infections with *C. neoformans*, *C. glabrata*, *C. auris* and *C. albicans* result in significantly reduced survival rates.

Figure 6: Caterpillar weight as a measure of virulence. Recording caterpillar weight daily revealed that yeast infections affect weight regardless of mortality rates. Daily weight measures were compared to that of PBS-injected animals and statistical significance assessed as follows: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure S1: Caterpillar growth is comparable between standard colony temperature and host body temperature. Groups of ten animals were kept on standard diet at 25˚C and 30˚C **(a)** or 25˚C and 37˚C **(b)** for four days. Animals were scored daily for survival and weight as a measure of development. Within these parameters, caterpillar development is comparable across all three temperatures.

Figure S2: Heat-killed *C. albicans* cells do not affect caterpillar survival or weight gain. Groups of ten animals were injected with either live *C. albicans* wild-type YSD85 cells, heat-killed cells, or PBS. a) Only live yeast cells kill caterpillars. b) The weight gain in animals infected with heat-killed *Candida* cells is comparable to that of animals injected with PBS.

Figure 1 - Lyons et al.

Figure 2 - Lyons et al.

Figure 3 - Lyons et al.

Figure 4 - Lyons et al.

a)

b)

p < 0.0001

Treatment: 0 mg/kg, 1 mg/kg, 4 mg/kg, 16 mg/kg, 32 mg/kg, 64 mg/kg, PBS

c)

d)

p < 0.0001

Treatment: 0 mg/kg, 1 mg/kg, 2 mg/kg, 4 mg/kg, PBS

Figure 5 - Lyons et al.

Figure 6 - Lyons et al.

Figure S1 - Lyons et al.

Figure S2 - Lyons et al.

a)

b)