Tobacco Hornworm (*Manduca sexta*) caterpillars as a novel host model for the study of fungal virulence and drug efficacy

Naomi Lyons[1, 2], Isabel Softley[2], Andrew Balfour[2], Carolyn Williamson[2], Heath E. O'Brien[3], Amol C. Shetty[4], Vincent M. Bruno[4], and Stephanie Diezmann[2, 5]

[1] School of Molecular Cell Biology and Biotechnology, Tel Aviv University, Tel Aviv, Israel

[2] Department of Biology & Biochemistry, University of Bath, Bath, United Kingdom

[3] LivingDNA, Frome, United Kingdom

[4] Institute for Genome Sciences, University of Maryland School of Medicine, Baltimore, MD, USA

[5] School of Cellular and Molecular Medicine, University of Bristol, Bristol, United Kingdom

Corresponding author:　　　Stephanie Diezmann

　　　　　　　　　　　　　s.diezmann@bristol.ac.uk

Running title: Caterpillars as fungal virulence host model

Abstract

The two leading yeast pathogens of humans, *Candida albicans* and *Cryptococcus neoformans*, cause systemic infections in >1.4 million patients world-wide with mortality rates approaching 75%. It is thus imperative to study fungal virulence mechanisms, efficacy of antifungal drugs, and host response pathways. While this is commonly done in mammalian models, which are afflicted by ethical and practical concerns, invertebrate models, such as wax moth larvae and nematodes have been introduced over the last two decades. To complement existing invertebrate host models, we developed fifth instar caterpillars of the Tobacco Hornworm moth *Manduca sexta* as a novel host model. These caterpillars can be maintained at 37˚C, are suitable for injections with defined amounts of yeast cells, and are susceptible to the most threatening yeast pathogens, including *C. albicans*, *C. neoformans*, *C. auris*, and *C. glabrata*. Importantly, fungal burden can be assessed daily throughout the course of infection in a single caterpillar's faeces and haemolymph. Infected caterpillars can be rescued by treatment with antifungal drugs. Notably, these animals are large enough for weight to provide a reliable and reproducible measure of fungal disease. *M. sexta* caterpillars combine a suite of parameters that make them suitable for the study of fungal virulence.

Keywords: *Manduca sexta*,

caterpillar,

fungal virulence,

Candida,

Cryptococcus,

Saccharomyces,

Metschnikowia,

antifungal drug,

host model,

fungal burden

Introduction

Fungal infections pose a serious threat to human health and well-being world-wide. Each year, as many, if not more patients, die of fungal infections than of malaria or tuberculosis[1]. The leading yeast pathogens, *Candida albicans* and *Cryptococcus neoformans*, account for ~1,400,000 life-threatening infections world-wide with mortality rates of up to 70%[1,2]. Candidaemia, most frequently caused by *C. albicans*, is the fourth most common cause of nosocomial blood stream infections, only surpassed by infections with Staphylococci and *Enterococcus* spp. Disturbingly, candidaemia incidence rates are on the rise. Within less than ten years, they increased by 36%[3]. Although cryptococcosis incidence rates are on the decline in North America, cryptococcosis as an AIDS-defining illness is responsible for 15% of all AIDS-related deaths world-wide[3,4]. This dire situation is further confounded by the emergence of drug resistant yeast species, such as *C. glabrata* and *C. auris*. Patients at risk of developing invasive candidaemia are often prophylactically treated with fluconazole and echinocandins as a first line defence strategy[5]. Yet, *C. glabrata*, the most common non-*albicans Candida* species associated with nosocomial blood stream infections[6], is intrinsically less susceptible to azole drugs and acquires resistance to echinocandins rapidly[7]. The rapid global spread of multi-drug resistant *C. auris* has further exacerbated the threat posed by fungal infections. *C. auris* was first reported in 2009 in Japan[8]. In 2015, *C. auris* arrived in Europe causing an outbreak involving 72 patients in a cardio-thoracic hospital in London[9]. *C. auris* outbreaks have been reported from South Korea, India, Spain, Columbia, Switzerland, Germany, Israel, Kuwait, and Oman[10]. Most concerningly, up to 25% of *C. auris* isolates are multi-drug resistant, with some strains being resistant to three of the four drug classes available for the treatment of systemic candidaemia. In addition to the unacceptably high burden on human health, fungal infections substantially increase health care costs. Treatment of candidaemia requires extended hospitalisation, resulting in additional costs of up to $45,000 in adult patients or up to $119,000 in paediatric patients[11].

It is thus imperative to investigate fungal virulence and host response mechanisms. This is traditionally done in mammalian models. The most frequently deployed models include the mouse tail vein infection model for systemic candidaemia, the mouse gastrointestinal infection model of candidaemia, the mouse *Candida* vaginitis model[12], the mouse inhalation model of cryptococcosis[13], the rabbit chronic cryptococcal meningitis model[14], and the rabbit *Candida* keratitis model[15]. While mammalian models combine a

number of features that make them particularly amenable for the study of fungal diseases,

such as susceptibility, availability of knock-out mutants, and comparable histology to human

disease, using mammals is ethically controversial, economically challenging, and requires

extensive board certifications and documentations.

In an effort to reduce the usage of mammalian hosts, alternative invertebrate models

have been developed and successfully used over the past two decades. The most commonly

employed invertebrate species include the nematode *Caenorhabditis elegans*, the fly

Drosophila melanogaster, and larvae of the Greater Wax moth *Galleria mellonella*. All three

can be easily maintained in the laboratory, are much less expensive than mice or rabbits, and

have been used for the study of diverse yeast pathogens, such as *C. neoformans*[16,17], *C.*

albicans[17-19], *C. parapsilosis*[19-21], *C. glabrata*[20,22]. Of note, invertebrate models differ in their

applicability and the best suitable model should be carefully selected[23]. Unlike mammalian

models, these invertebrates do not have adaptive immunity but all share components of the

innate immune system[24,25], some of which are conserved with mammals. This includes the

Toll-like receptors found in the fly[26] and the homolog of the MKK3/6 kinase in the

nematode[27]. Ironically, it is the Toll-like receptors that protect flies from infections with *C.*

neoformans[28], *C. albicans*[29], and *C. glabrata*[30] and the MKK3/6 homolog SEK-1 protects the

nematode from bacterial invaders[27]. Thus, to increase susceptibility of flies and nematodes to

fungal pathogens, Toll and *sek-1*[31] mutants are required. A key limitation for the study of

human pathogens, is the inability of nematodes and flies to survive human body temperature.

Only *Galleria* can withstand 37°C[17]. Due to their long-standing history as eukaryotic models,

well-curated genomes and genome databases exist for the nematode and the fly. Although the

Galleria genome has been announced very recently[32], detailed analyses and annotations are

still outstanding. Delivery of an exact inoculum of fungal cells is crucial when comparing

mutant and wild type yeast strains, neither fly nor nematode permit routine direct injection.

Galleria can be directly injected with a defined cell number. Extensive stock collections

provide fly and nematode strains, while until recently, *Galleria* larvae had to be purchased

from fishing shops. Now, UK-based TruLarv is selling research grade larvae.

Tobacco Hornworms are most commonly encountered in the southern United States,

where they feed on solanaceous plants and are thus considered a plant pest. As an insect

model with a long history in research, *Manduca sexta* has yielded important insights into

flight mechanisms, nicotine resistance, hormonal regulation of development, metamorphosis,

antimicrobial defences, and bacterial pathogenesis. *M. sexta* laboratory stocks have been

derived from animals collected in North Carolina, USA[33] and have been maintained in

laboratories on both sides of the Atlantic for several decades. *M. sexta*'s research portfolio

includes a draft genome sequence that has been complemented with tissue-specific

transcriptomic analyses[34], numerous successful applications of RNAi[35-39], and protocols for

the efficient extraction of hemocytes for down-stream analyses[40] of the animal's innate

immunity[41]. Despite its versatility, *M. sexta* has yet to be explored for its suitability as a host

model for fungal infections.

Here, we aimed to establish *M. sexta* as a novel model host for the study of fungal

virulence. Inbred animals from the University of Bath's research colony were tested for their

ability to live at 37˚C, their susceptibility to different yeast species, and the reproducibility of

C. albicans mutant phenotypes obtained in mice virulence studies. Indeed, the caterpillars

grow at 37˚C while maintaining susceptibility to the leading yeast pathogens *C. albicans*, *C.*

neoformans, and emergent *C. auris*. Specific *C. albicans* mutants are just as attenuated in

their virulence in *M. sexta* as they are in mice. To expand *M. sexta's* applicability as a host

model, we developed an infection protocol that permits screening of fungal burden

throughout the course of infection in a single animal and uses weight as a proxy measure for

virulence in addition to survival. *M. sexta* can furthermore be used to test efficacy of common

antifungal drugs and to query the host transcriptional response to systemic yeast infections.

Our results define *M. sexta* characteristics that recommend these caterpillars as a non-

mammalian host model for the study of fungal virulence with susceptibility to different yeast

species.

Materials and Methods

Origin of the Bath colony of *Manduca sexta*

The University of Bath colony has been in continuous culture since 1978 without the

addition of animals from elsewhere. Bath's genetic stock was derived from animals from the

Truman-Riddiford laboratories at the University of Washington in Seattle, USA. Their

animals date back to the ones originally collected in North Carolina in 1976[33].

Caterpillar maintenance and yeast culture conditions

M. sexta caterpillars were reared to fifth instar under standardised conditions. They

were maintained in 125 ml disposable cups (Sarstedt Ltd., Cat. No. 75.1335) on a wheat

germ-based diet (Appendix 'Food preparation') at a constant temperature of 25˚C with 50%

humidity and 12 hours light/dark cycles. Three days prior to fungal inoculations, animals

were shifted to a formaldehyde-free diet as the compound is toxic to non-methylotrophic

yeast.

For infection assays, yeasts were grown overnight in 50 ml YPD (1% yeast extract,

2% peptone, 2% dextrose) and cells harvested by centrifugation for 3 minutes at 3,000 rpm.

The cell pellet was washed twice with 1x phosphate buffered saline (PBS) and suspended in 5

ml 1x PBS. Cells were counted and numbers adjusted as indicated. *C. albicans* YSD85

(Table 1) cells were heat-inactivated by incubation at 65˚C for 20 minutes. For long-term

storage, yeast isolates were cryo-archived at -80˚C in 25% glycerol.

Yeast infections and measurements of fungal burden and drug efficacy

100 µl of washed and number-adjusted yeast suspension were injected into each

caterpillar's distal right proleg with a 30G1/2" needle (BD Microlance) and a 1 ml NORM-

JECT syringe. Following injection, each animal's weight was recorded. Animals were scored

for survival and weight once daily for three or four days post infection. During the course of

the experiment, animals were kept on a 12-hour light/dark cycle at the temperature indicated

and on their regular diet.

To measure fungal burden in caterpillar faeces and haemolymph, six animals were

injected with 100 µl of either 1x PBS or 10^6 cells of the wild type YSD89 or the *hog1* mutant

strain YSD883 and kept at 37˚C. On day 1, two animals were selected from each group.

These animals were weighted and their haemolymph and faeces collected daily throughout

the course of infection. To collect haemolymph, animals were first kept on ice for 15 minutes.

The 'horn' was then surface sterilised with 70% ethanol and its top 1-2 mm clipped with a

pair of micro scissors. Haemolymph was collected in a pre-chilled 1.5 ml Eppendorf tube and

cooled immediately to reduce polymerisation and melanisation. One faecal pellet was

collected daily with sterile forceps, weighted and suspended in 500 µl 1x PBS. Prior to

diluting, the suspension was thoroughly vortexed for 10 seconds, and centrifuged for 5

seconds using a table top centrifuge to separate faecal matter. To quantify fungal burden,

haemolymph and faecal samples were plated either directly onto YPD-agar with Kanamycin

50 µg/ml or in ten-fold serial dilutions. Agar plates were incubated at 30˚C for 48 hours and

colonies counted.

To assess the efficacy of commonly used antifungal drugs, animals were infected with

10^7 cells of YSD85 or 100 µl PBS and treated with increasing doses of fluconazole and

caspofungin (Sigma Aldrich, Inc.) as indicated. Drugs were injected 30 minutes post-

infection with an ethanol-sterilized Hamilton syringe in a total volume of 10 µl per animal.

Caterpillars were weighed and scored for survival on the day of injection and the following

three days.

Statistical analyses

Survival plots were made using the survminer R package (https://CRAN.R-

project.org/package=survminer), and differences were evaluated using the Kaplan-Meier

method. Weight and fungal burden were plotted using ggplot2[42] and weight differences were

evaluated using linear models with day post-inoculation and the interaction between

treatment and days post infection as fixed effects and individual as a random effect using

nlme (https://CRAN.R-project.org/package=nlme). Upset plots visualising gene expression

profiles were made using the UpSetR package[43]. All analyses were conducted in RStudio

version 1.1.442.

RNA sequencing and gene expression analysis

Five animals were injected with 100 µl 1x PBS or 10^6 yeast cells of the wild type

strain SN95 (YSD89) and maintained on a 12-hour light/dark cycle at 37˚C for 24 hours.

Three animals were then randomly selected for transcriptomic analyses of their midguts. In

preparation for midgut extraction, animals were surface sterilized and tissue extracted and

washed as described before[34]. Midgut RNA was extracted using TRIzol reagent (Invitrogen,

Inc.), quantified using the Agilent 2100 Bioanalyzer system and used to generate RNA-seq

libraries (strand-specific, paired end) using the TruSeq RNA sample prep kit (Illumina).

150 nucleotides of the sequence were determined from both ends of each cDNA

fragment using the HiSeq 2500 platform (Illumina). An average of 83.79 million read pairs

were obtained for each sample (range: 51.4 M to 109.8 M read pairs). Sequencing reads were

aligned to the reference *Manduca sexta* whole genome assembly v1.0

(https://i5k.nal.usda.gov/Manduca_sexta) using HISAT2[44] and alignment files were used to

generate read counts for each gene. Statistical analysis of differential gene expression was

performed using the DEseq2 package from Bioconductor[45]. A gene was considered

differentially expressed if the FDR-value for differential expression was less than 0.10.

In order to compare the transcriptional response to *C. albicans* infection by *M. sexta* to that of

a mouse, we generated a blast reference database using the peptide sequences for 67,960

proteins from the mouse genome (GRCm38) available from Ensembl

(https://useast.ensembl.org). We downloaded the peptide sequences for 27,403 proteins from

the *M. sexta* database (OGS v2.0) available from the National Agricultural Library

(https://i5k.nal.usda.gov/Manduca_sexta)[34] provided by the United States Department of

Agriculture. We computed peptide sequence similarity between the *M. sexta* proteins and the

mouse proteins using the blastp local alignment search tool (ncbi-blast+ v2.8.1)[46]. For each

M. sexta protein, we had multiple hits that were ranked based on the e-value computed by the

blastp search tool. For the list of differentially expressed *M. sexta* genes from each

comparison, we extracted the mouse orthologs with e-values smaller than e^{-20}. This list of

orthologs was then compared to differential expression lists based on RNA-seq analysis of

kidneys[47], tongues[48], and vaginas[49] from *C. albicans* infected mice. Gene lists were compared

using Venny 2.1.0 (https://bioinfogp.cnb.csic.es/tools/venny/index.html). RNAseq data are

available from NCBI's Short Read Archive (###).

Results

***M. sexta* caterpillars are susceptible to the leading fungal pathogen *C. albicans* at 37˚C**

We first aimed to determine if *M. sexta* fifth instar caterpillars (Fig. 1a), reared and

maintained at standard conditions, are indeed susceptible to *C. albicans*. To do so, groups of

ten animals were infected with increasing doses of the widely used *C. albicans* laboratory

strains SC5314 and SN95[50]. Animals were kept at 25˚C, their standard maintenance

temperature, and scored for survival on three consecutive days. Dead animals differ from live

ones in that their bodies go limp and turn brown-green in colour, which is in stark contrast to

the vivid turquoise of live animals (Fig. 1b). Caterpillars infected with *C. albicans*

succumbed in a dose-dependent manner. Both *C. albicans* strains killed *M. sexta* efficiently at

inocula of 10^6 or 10^7 cells per animal (Fig. 1c). To assess if survival measures in caterpillars

are comparable to those obtained in the current gold standard, the murine model of systemic

candidaemia, we tested *C. albicans* mutants with published phenotypes of either attenuated

virulence, such as the *hog1Δ/Δ*[51] and *ahr1Δ/Δ*[52] mutants, or wild-type levels of virulence,

such as *cka2Δ/Δ*[53]. Cross-species virulence levels are comparable for Hog1, which is just as

essential for virulence in caterpillars as it is in mice. Ahr1, while required for virulence in

mammals, appears to be dispensable for virulence in caterpillars. Cka2 is not required to

establish systemic infections in mammals but is in caterpillars (Fig. 1d).

Given the importance of temperature for fungal virulence, we queried if *M. sexta*

retained their susceptibility to *C. albicans* at human body temperature of 37˚C. Temperature

itself does not affect caterpillar survival or development (Fig. S1) but animals are ten times

more susceptible to infections with *C. albicans* at 37˚C than they are at 25˚C (Fig. 2a). At

37˚C, 10^6 *C. albicans* cells per animal lead to 100% mortality on day 4, while 10^7 cells are

required for the same outcome at 25˚C (Fig. 1b). To exclude the possibility that mortality is

due to starvation rather than the outcome of a host-pathogen interaction, we infected

caterpillars with live and heat-killed *C. albicans* wild-type cells and kept them at 37˚C. Only

live cells, but not heat-killed yeast cells, killed the caterpillars suggesting that killing is not

due to nutritional limitations (Fig. S2). Demonstrating susceptibility of *M. sexta* caterpillars

to *C. albicans* lent support to them as an alternative host model for the study of fungal

virulence and emphasized the need for additional measures of fungal virulence. To add

granularity to fungal virulence data collected from *M. sexta*, we complemented measures of

survival with quantifications of weight and fungal burden throughout the course of infection.

To collect weight data, caterpillars were weighed prior to infection and then daily throughout

the course of infection. Weight gain in animals infected with a low dose of 10^4 cells did not

significantly differ from those injected with 1x PBS but caterpillars infected with 10^5 cells per

animal exhibited significant weight loss. Too few animals survived infection with 10^6 cells to

allow for a meaningful comparison (Fig. 2b). After establishing susceptibility of *M. sexta* to

C. albicans at 37˚C, we aimed to validate the attenuated virulence phenotype of the *hog1Δ/Δ*

mutant strain. To test this, caterpillars were infected with 10^6 cells per animal with the wild

type strain, the *hog1Δ/Δ* deletion mutant, the *hog1/hog1::HOG1* complementation strain and

compared to the control group injected with 1x PBS. The *hog1Δ/Δ* mutant is less virulent

than the wild type (Fig. 2c). Animals infected with the *hog1Δ/Δ* mutant strain gained

significantly more weight than those infected with the wild type strain, while infection with

the complemented strain lead to a comparable lack in weight gain (Fig. 2d).

Measurements of fungal burden in faeces and haemolymph obtained from a single

animal throughout infection provides a useful measure of virulence

To further expand the applicability of *M. sexta* caterpillars as a host model for fungal

infections, fungal burden in the haemolymph and faeces was quantified daily throughout the

course of infection. Since the collection of haemolymph or faeces does not necessitate killing

the animal, data could be collected daily throughout the infection for the same caterpillar.

Animals infected with the wild type and the *hog1Δ/Δ* mutant were compared to control

animals injected with 1x PBS only and fungal burden measured as colony-forming units

(CFUs) in faeces and haemolymph in two animals per group. Yeasts were detected in the

faeces and haemolymph of animals infected with the wild type on day 2 and CFU counts

increased on day 3 (Fig. 3a, b). In animals infected with the *hog1Δ/Δ* mutant, we detected low

CFU counts on day 3 in faeces and haemolymph. Even fewer CFUs were detected in the

haemolymph of animals injected with PBS on day3. Consistent with the observed increase in

fungal burden in wild-type infected animals, indicative of a fungal infection, these animals

displayed reduced weight gain while *hog1Δ/Δ* infected animals gained weight at a similar rate

as the control animals (Fig. 3c). Thus, measurements of fungal burden, that can be obtained

from the same animal throughout the course of infection, provide a valuable parameter for the

study of fungal virulence.

Caterpillars are suitable for antifungal drug treatment studies

The currently available armamentarium of antifungal drugs is limited and drugs are

often lacking in efficacy. The ability to study drug efficacy and mode of action in a host

model are thus pertinent to the development of novel antifungal drugs. To determine *M.*

sexta's suitability for drug efficacy testing, we recorded survival and weight of animals

infected with the *C. albicans* wild-type strain SC5314 that were treated with increasing doses

of two common antifungals, fluconazole and caspofungin (Fig.4). Treatment with fluconazole

or caspofungin resulted in overall improved survival and weight gain. Animals treated with 2

mg/kg and 4 mg/kg of caspofungin survived significantly better than those without treatment

or those that only received 1 mg/kg of caspofungin. Pair-wise comparisons between different

drug concentrations and the untreated control animals yielded no statistical significance for

the fluconazole-treated group of caterpillars (Table 2).

***M. sexta* caterpillars are broadly susceptible to different yeast pathogens**

While *C. albicans* undoubtedly is a leading fungal pathogen of humans, clinical

manifestations of fungal infections are not limited to *C. albicans*. To broaden *M. sexta*'s

applicability, we sought to quantify the caterpillars' susceptibility to other fungal pathogens.

To this end, animals were also infected with wild type strains of *C. neoformans*, *C. auris*, and

C. glabrata. Type strains of *Saccharomyces cerevisiae*, the baker's or brewer's yeast and

Metschnikowia pulcherrima, a yeast inhabiting fruits and flowers[54] served as reference points

for attenuated virulence. Animals were infected with increasing doses of yeast cells starting at

10^5 cells per animal and up to 10^9 cells per animal. Groups of ten animals per yeast dose and

species were then screened for survival and weight daily (Figs. 5 and 6). Notably, only

infections with pathogenic yeast species affected survival of caterpillars (Fig. 5). Infections

with *S. cerevisiae* or *M. pulcherrima* did not significantly reduce caterpillar survival.

Comparing survival amongst the pathogenic yeast species revealed *C. albicans* to be the most

severe. 10^7 *C. albicans* cells per animal result in 100% killing within in 24 hours. 10^9 *C. auris*

cells were required to kill all animals within four days and the same dose of *C. glabrata*

resulted in 75% killing. 10^8 *C. neoformans* cells were required to achieve 50% killing within

four days, comparable to *C. glabrata*. It should be noted that due to the high viscosity of the

cell suspension, we could not test higher concentrations than the ones stated here. In addition

to assessing survival, each surviving caterpillar's weight was recorded daily. Interestingly,

while infections with *S. cerevisiae* or *M. pulcherrima* did not kill caterpillars, infected

animals did not gain weight as effectively as uninfected control animals (Fig. 6). Infections

with *C. albicans* resulted in severely reduced weight gain, even at the lowest yeast dose

tested. Animals infected with *C. auris* and *C. glabrata* responded in a dose-dependent

manner, the higher the yeast dose, the more dramatic the weight loss. Notably, animals

infected with the lowest dose of *C. neoformans* grew better than PBS-injected control

animals. Increasing the yeast dosage, however, resulted in significantly reduced weight gain.

This bi-phasic pattern, resembling hormesis[55] in which a low dose of an environmental agent

is beneficial while a high dose is toxic, could be due to *C. neoformans*' immunogenic

capsule[56]. Caterpillars displayed variable degrees of susceptibility to different yeast

pathogens and with weight providing an additional measure of host damage. Thus, *M. sexta*

caterpillars are broadly applicable for the study of fungal virulence.

M. sexta transcriptional response profiles differ between uninfected and infected

animals

As a proof-of-concept we recorded transcriptional profiles in three animals infected

with the wild-type strain and compared them to three uninfected control animals. To do so,

we collected the mid-gut, rinsed the tissue thoroughly, and extracted RNA, which was then

sequenced using Illumina' HiSeq 2500 platform. With the vast majority of reads mapping to

M. sexta rather than *C. albicans* (Fig. S3), animals infected with *Candida* displayed a

transcriptional profile different from that of uninfected animals (Fig. S4, Table S1). Infection

with the wild type strain elicited quantifiable transcriptional changes in the caterpillars. 157

M. sexta genes were up-regulated and 165 went down when compared to the PBS control

animals (Table S1).

Lastly, we compared our set of differentially expressed *M. sexta* genes to those

identified in three murine candidaemia studies. Here, tissue-specific responses in the

kidneys[47], tongues[48], and vaginas[49] were recorded for animals infected with *Candida* wild

type (Fig. 7). *M. sexta* genes up-regulated during infection and their homologs in all three

mouse tissues include serine peptidase inhibitors (Serpinb3c, Serpinb3b), a Ras-like protein

(Rit2), transmembrane proteases (Tmprss11a, Tmprss11f), a histidine decarboxylase (Hdc),

and a peptidoglycan recognition protein (Pglyrp3). *M. sexta* genes down-regulated in mouse

kidney and vagina include a muscle glycogen phosphorylase (Pygm), a bone morphogenetic

antagonist of the DAN family (Grem1), meprin 1 alpha (Mep1a), and calponin 1 (Cnn1).

Two of these genes are annotated as functioning in immune system processes (Grem1,

Pglyrp3) and three more are relevant for signalling and response to different stimuli (Rit2,

Pglyrp3, Pygm). Seven genes, however, are involved in protein metabolic processes

(Serpinb3c, Serpinb3b, Rit2, Tmprss11a, Tmprss11f, Grem1, Mep1a), suggesting this

fundamental process to be of importance in the defence against microbial attacks.

Discussion

Invertebrate host models have become a valuable alternative to mammalian hosts for

the study of fungal disease. When choosing the appropriate invertebrate model, care needs to

be taken and experimental parameters should be considered[57]. Here we developed *M. sexta*

caterpillars as a novel invertebrate model showing that they are naturally susceptible to

different human pathogenic yeast species, including the emergent *C. auris*. Unlike other host

models, these large caterpillars permit daily measures of fungal burden throughout the course

of infection in a single animal by either collecting faeces or haemolymph. *M. sexta* can be

maintained at 37˚C, is large enough to be injected with a specified yeast inoculum and for

weight to be a reliable measure of virulence. *C. albicans* mutant virulence phenotypes found

in mice can be replicated and yeast inocula required to elicit a response in caterpillars are

comparable to those used in the murine model. Additionally, the caterpillars permit study of

antifungal drug efficacy. These parameters commend this invertebrate species as a novel host

model for the study of fungal virulence.

This new model system allows for fungal burden to be monitored throughout the

course of infection in a single animal via CFU count. This is unlike any other experimental

system, where fungal burden is either an endpoint measure in the mouse kidney, in

homogenised wax moth larvae[58], nematodes[59], flies[60], or requires genetically modified

fluorescent yeast strains for microscopic imaging and analyses[61]. While we did detect very

low CFU counts in the PBS control 3 days post infection, we consider this a spurious finding

due to cross contamination as preliminary experiments of plating contents of haemolymph

and faeces of naïve animals did not detect any yeast growth (data not shown). As a

consequence, we amended the protocol to include changing gloves when handling animals of

different treatment groups.

When reviewing the weight data collected as part of our study, we noticed two

interesting aspects. First, while *S. cerevisiae* and *M. pulcherrima* do not affect caterpillar

survival, infections with either species led to reduced weight gain. The dichotomy between

survival and weight observed here, further emphasizes that fungal virulence comprises more

than a measure of survival. It appears that *M. sexta* would allow discrimination between

disease (weight) and death (survival) quantitatively adding further granularity to measuring

fungal virulence. Secondly, infections with *C. neoformans* resulted in increased caterpillar

weight gain at a low yeast dose but reduced weight gain at higher doses. This pattern

resembles the concept of hormesis often deployed by toxicologists to describe the response to

toxins, where exposure to a low dose is beneficial while a higher dose results in toxicity due

to overcompensation in response to disruption of homeostasis[62]. While tissue-specific

hormetic responses have been described in flies, where a virus-acquired cytokine delays

ageing[63], and examples of abiotic factors or signalling molecules affecting peas and aphid

infestation[64], immunity in plants[65,66], life-span in malaria-transmitting *Anopheles*[67], and larval

development in Black Cutworm[68], this example here could be the first involving a eukaryotic

host-pathogen relationship. The underlying factor remains to be elucidated but *C.*

neoformans' highly immunogenic capsule seems to be an excellent candidate[56].

Heat-killed *C. albicans* cells appear to be entirely non-pathogenic in *M. sexta*

caterpillars as neither survival nor weight are affected by inoculation with heat-killed yeast

cells. The lack of mortality in response to inoculation with heat-killed yeast cells indicates

that yeast viability and proliferation are required for pathogenesis and excludes the possibility

of death due to an allergic reaction in response to a large number of fungal cells. This appears

to differ from the responses of other host models to fungal pathogens. While susceptibility

was reduced, but still measurable, in *G. mellonella*[69] and the two-spotted cricket[70], heat-killed

C. albicans cells elicited 100% mortality in a sepsis-like murine model[71]. In mice, serum

levels of β-(1,3)-glucan levels were elevated in animals injected with heat-killed yeast cells

when compared to those infected with live cells. Indeed, heat inactivation leads to increased

exposure of β-(1,3)-glucan on the *C. albicans* cell surface[72] and β-(1,3)-glucan activates the

innate immune response in invertebrates and mammals[73]. Given the complexity of receptors

involved in recognition of fungal invaders[74] and the lack of a response in *M. sexta* and other

invertebrates indicates that they may not express all or identical receptors as mammals.

Measures of the host transcriptional response provide useful insights into the

relevance of specific defence mechanisms. Mouse transcriptional profiles in response to *C.*

albicans infections differ between different types of tissues[48,49,75], which allowed for the

identification of tissue-specific response mechanisms such as IL-17 signalling in oral

epithelia. Here, we detected numerous differentially expressed genes in the caterpillar midgut

when comparing animals infected with *C. albicans* to the PBS control. Given the size of *M.*

sexta caterpillars, which permits tissue-specific transcriptional analyses[34], this phenomenon

could be systematically investigated in caterpillars rather than mammalian hosts, who are

burdened by ethic concerns and economic challenges prohibiting large scale or time-course

studies. Furthermore, partial conservation of the transcriptional response to *C. albicans*

infections between insects and mammals offers the opportunity to identify and investigate

core eukaryotic response mechanisms.

In summary, *Manduca sexta* caterpillars expand the current repertoire of invertebrate

models for the study of fungal disease. They combine a suite of measures that commend it as

a new model system. Although, *M. sexta* genomic and transcriptomic analyses are currently

still in their infancy, we would expect that the Tobacco Hornworm's long history of being an

invaluable model for diverse facets of biology will lead to reliable tools in combination with

genetic tractability and protocols establishing the fungus' fate inside the caterpillar.

Acknowledgements

We would like to thank Ewan Basterfield and Chris Apark for expert advice and

technical assistance in preparing *M. sexta* caterpillars. Thanks to all the laboratories who

kindly shared their strains with us. This work was supported by an ERC Marie Curie Career

Integration Grant to SD as well as undergraduate and postgraduate research funding from the

University of Bath.

Disclosure of interest

The authors report no conflict of interest.

References

1. Brown GD, Denning DW, Gow NAR, Levitz SM, Netea MG, White TC. Hidden
killers: Human fungal infections. Sci Transl Med 2012; 4:165rv13.

2. Harsha MV, International SV, 2017. Emerging fungal pathogens-a major threat
to human life. IJPSR 8:1923–34.

3. Pfaller MA, Diekema DJ. Epidemiology of invasive mycoses in North America.
Crit Rev Microbiol 2010; 36:1–53.

4. Nyazika TK, Tatuene JK, Kenfak-Foguena A, Verweij PE, Meis JF, Robertson
VJ, Hagen F. Epidemiology and aetiologies of cryptococcal meningitis in Africa,
1950–2017: Protocol for a systematic review. BMJ Open 2018; 8:e020654–4.

5. Pappas PG, Kauffman CA, Andes D, Benjamin DK Jr, Calandra TF, Edwards JE
Jr, Filler SG, Fisher JF, Kullberg B-J, Ostrosky Zeichner L, et al. Clinical
practice guidelines for the management of candidiasis: 2009 update by the
Infectious Diseases Society of America. Clin Inf Dis 2009; 48:503–35.

6. Pfaller MA, Diekema DJ. Epidemiology of invasive candidiasis: A persistent
public health problem. Clin Microbiol Rev 2007; 20:133–63.

7. Jensen RH, Johansen HK, Søes LM, Lemming LE, Rosenvinge FS, Nielsen L,
Olesen B, Kristensen L, Dzajic E, Astvad KMT, et al. Posttreatment antifungal
resistance among colonizing *Candida* isolates in candidemia patients: Results
from a systematic multicenter study. Antimicrob Agents Chemother 2016;
60:1500–8.

8. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H.
Candida auris sp. nov., a novel ascomycetous yeast isolated from the external
ear canal of an inpatient in a Japanese hospital. Microbiol Immunol 2009; 53:41–
4.

9. Schelenz S, Hagen F, Rhodes JL, Abdolrasouli A, Chowdhary A, Hall A, Ryan
L, Shackleton J, Trimlett R, Meis JF, et al. First hospital outbreak of the globally
emerging *Candida auris* in a European hospital. Antimicrob Resist Infect
Control 2016; 5:1–7.

10. Bidaud AL, Chowdhary A, Dannaoui E. *Candida auris*: An emerging drug
resistant yeast – A mini-review. J Mycol Med 2018; 28:568–73.

11. Zaoutis TE, Argon J, Chu J, Berlin JA, Walsh TJ, Feudtner C. The epidemiology
and attributable outcomes of candidemia in adults and children hospitalized in
the United States: A propensity analysis. Clin Infect Dis 2005; 41:1232–9.

12. Segal E, Frenkel M. Experimental *in vivo* models of candidiasis. JoF 2018; 4:21–
10.

13. Zaragoza O, Alvarez M, Telzak A, Rivera J, Casadevall A. The relative
susceptibility of mouse strains to pulmonary *Cryptococcus neoformans* infection
is associated with pleiotropic differences in the immune response. Infect Immun

2007; 75:2729–39.

14. Perfect JR, Lang SD, Durack DT. Chronic cryptococcal meningitis: A new experimental model in rabbits. Am J Pathol 1980; 101:177–94.

15. Ray WA, O'Day DM, Head WS, Robinson RD. Variability in isolate recovery rates from multiple and single breeds of outbred pigmented rabbits in an experimental model of *Candida* keratitis. Curr Eye Res 2009; 3:949–53.

16. Mylonakis E, Ausubel FM, Perfect JR, Heitman J, Calderwood SB. Killing of *Caenorhabditis elegans* by *Cryptococcus neoformans* as a model of yeast pathogenesis. PNAS 2002; 99:15675–80.

17. Fuchs BB, O'Brien E, Khoury El JB, Mylonakis E. Methods for using *Galleria mellonella* as a model host to study fungal pathogenesis. Virulence 2014; 1:475–82.

18. Pukkila-Worley R, Ausubel FM, Mylonakis E. *Candida albicans* infection of *Caenorhabditis elegans* induces antifungal immune defenses. PLoS Pathog 2011; 7:e1002074.

19. Chamilos G, Lionakis MS, Lewis RE, Lopez-Ribot JL, Saville SP, Albert ND, Halder G, Kontoyiannis DP. *Drosophila melanogaster* as a facile model for large-scale studies of virulence mechanisms and antifungal drug efficacy in *Candida* species. J Infect Dis 2006; 193:1014–22.

20. Ortega-Riveros M, De-la-Pinta I, Marcos-Arias C, Ezpeleta G, Quindós G, Eraso E. Usefulness of the non-conventional *Caenorhabditis elegans* model to assess *Candida* virulence. Mycopathologia 2017; 182:785–95.

21. Gago S, García-Rodas R, Cuesta I, Mellado E, Alastruey-Izquierdo A. *Candida parapsilosis*, *Candida orthopsilosis*, and *Candida metapsilosis* virulence in the non-conventional host *Galleria mellonella*. Virulence 2014; 5:278–85.

22. Ames L, Duxbury S, Pawlowska B, Ho H-L, Haynes K, Bates S. *Galleria mellonella* as a host model to study *Candida glabrata* virulence and antifungal efficacy. Virulence 2017; 8:1–9.

23. Desalermos A, Fuchs BB, Mylonakis E. Selecting an invertebrate model host for the study of fungal pathogenesis. PLoS Pathog 2012; 8:e1002451.

24. Mylonakis E, Aballay A. Worms and flies as genetically tractable animal models to study host-pathogen interactions. Infect Immun 2005; 73:3833–41.

25. Vogel H, Altincicek B, Glöckner G, Vilcinskas A. A comprehensive transcriptome and immune- gene repertoire of the lepidopteran model host *Galleria mellonella*. BMC Genomics 2011; 12:308.

26. Lemaitre B, Nicolas E, Michaut L, Reichhart JM, Hoffmann JA. The dorsoventral regulatory gene cassette spätzle/Toll/cactus controls the potent antifungal response in *Drosophila* adults. Cell 1996; 86:973–83.

27. Kim DH, Feinbaum R, Alloing G, Emerson FE, Garsin DA, Inoue H, Tanaka-
Hino M, Hisamoto N, Matsumoto K, Tan M-W, et al. A conserved p38 MAP
kinase pathway in *Caenorhabditis elegans* innate immunity. Science 2002;
297:623–6.

28. Apidianakis Y, Rahme LG, Heitman J, Ausubel FM, Calderwood SB, Mylonakis
E. Challenge of *Drosophila melanogaster* with *Cryptococcus neoformans* and
role of the innate immune response. Eukaryot Cell 2004; 3:413–9.

29. Alarco A-M, Marcil A, Chen J, Suter B, Thomas D, Whiteway M. Immune-
deficient *Drosophila melanogaster*: A model for the innate immune response to
human fungal pathogens. J Immunol 2004; 172:5622–8.

30. Quintin J, Asmar J, Matskevich AA, Lafarge MC, Ferrandon D. The *Drosophila*
Toll pathway controls but does not clear *Candida glabrata* infections. J Immunol
2013; 190:2818–27.

31. Breger J, Fuchs BB, Aperis G, Moy TI, Ausubel FM, Mylonakis E. Antifungal
chemical compounds identified using a *C. elegans* pathogenicity assay. PLoS
Pathog 2007; 3:e18.

32. Lange A, Beier S, Huson DH, Parusel R, Iglauer F, Frick J-S. Genome sequence
of *Galleria mellonella* (Greater Wax Moth). Genome Announc 2018; 6:61–2.

33. Flowers RW, Entomologist RYF, 1982. Feeding on non-host plants by the
Tobacco Hornworm (*Manduca sexta* (Lepidoptera: Sphingidae). JSTOR 1982;
65:523–30.

34. Kanost MR, Arrese EL, Cao X, Chen Y-R, Chellapilla S, Goldsmith MR,
Grosse-Wilde E, Heckel DG, Herndon N, Jiang H, et al. Multifaceted biological
insights from a draft genome sequence of the Tobacco Hornworm moth,
Manduca sexta. Insect Biochem Mol Biol 2016; 76:118–47.

35. Flores-Escobar B, Rodríguez-Magadan H, Bravo A, Soberón M, Gómez I.
Differential role of *Manduca sexta* aminopeptidase-N and alkaline phosphatase
in the mode of action of Cry1Aa, Cry1Ab, and Cry1Ac toxins from *Bacillus*
thuringiensis. Appl Environ Microbiol 2013; 79:4543–50.

36. Burke WG, Kaplanoglu E, Kolotilin I, Menassa R, Donly C. RNA interference in
the Tobacco Hornworm, *Manduca sexta*, using plastid-encoded long double-
stranded RNA. Front Plant Sci 2019; 10:313.

37. Eleftherianos I, Millichap PJ, ffrench-Constant RH, Reynolds SE. RNAi
suppression of recognition protein mediated immune responses in the Tobacco
Hornworm *Manduca sexta* causes increased susceptibility to the insect pathogen
Photorhabdus. Dev Comp Immunol 2006; 30:1099–107.

38. Eleftherianos I, Gökçen F, Felföldi G, Millichap PJ, Trenczek TE, ffrench-
Constant RH, Reynolds SE. The immunoglobulin family protein Hemolin
mediates cellular immune responses to bacteria in the insect *Manduca sexta*. Cell
Microbiol 2007; 9:1137–47.

39.   Kumar P, Pandit SS, Baldwin IT. Tobacco Rattle Virus vector: A rapid and
transient means of silencing *Manduca sexta* genes by plant mediated RNA
interference. PLoS ONE 2012; 7:e31347–10.

40.   Stoepler TM, Castillo JC, Lill JT, Eleftherianos I. A simple protocol for
extracting hemocytes from wild caterpillars. JoVE 2012; :1–6.

41.   Kanost MR, Jiang H, Yu X-Q. Innate immune responses of a lepidopteran insect,
Manduca sexta. Immunol Rev 2004; 198:97–105.

42.   Wickham H. ggplot2. New York, NY: Springer-Verlag New York; 2009.

43.   Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. UpSet: Visualization
of intersecting sets. IEEE Trans Visual Comput Graphics 2016; 20:1983–92.

44.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol
2019; 37:907–15.

45.   Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol 2014; 15:31–21.

46.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden
TL. BLAST+: Architecture and applications. BMC Bioinformatics 2009;
10:421–9.

47.   Liu Y, Shetty AC, Schwartz JA, Bradford LL, Xu W, Phan QT, Kumari P,
Mahurkar A, Mitchell AP, Ravel J, et al. New signaling pathways govern the
host response to *C. albicans* infection in various niches. Genome Res 2015;
25:679–89.

48.   Conti HR, Bruno VM, Childs EE, Daugherty S, Hunter JP, Mengesha BG,
Saevig DL, Hendricks MR, Coleman BM, Brane L, et al. IL-17 receptor
signaling in oral epithelial cells is critical for protection against oropharyngeal
candidiasis. Cell Host Microbe 2016; 20:606–17.

49.   Bruno VM, Shetty AC, Yano J, Fidel PL, Noverr MC, Peters BM.
Transcriptomic analysis of vulvovaginal candidiasis identifies a role for the
NLRP3 inflammasome. mBio 2015; 6:3896–15.

50.   Noble SM, Johnson AD. Strains and strategies for large-scale gene deletion
studies of the diploid human fungal pathogen *Candida albicans*. Eukaryot Cell
2005; 4:298–309.

51.   Alonso-Monge R, Navarro-García F, Molero G, Diez-Orejas R, Gustin M, Pla J,
Sánchez M, Nombela C. Role of the mitogen-activated protein kinase Hog1p in
morphogenesis and virulence of *Candida albicans*. J Bacteriol 1999; 181:3058–
68.

52.   Askew C, Sellam A, Epp E, Mallick J, Hogues H, Mullick A, Nantel A,
Whiteway M. The zinc cluster transcription factor Ahr1p directs Mcm1p
regulation of *Candida albicans* adhesion. Mol Microbiol 2010; 79:940–53.

53. Chiang LY, Sheppard DC, Bruno VM, Edwards JE, Filler SG. *Candida albicans*
protein kinase CK2 governs virulence during oropharyngeal candidiasis. Cell
Microbiol 2007; 9:233–45.

54. Vadkertiová R, Molnárová J, Vránová D, Sláviková E. Yeasts and yeast-like
organisms associated with fruits and blossoms of different fruit trees. Can J
Microbiol 2012; 58:1344–52.

55. Mattson MP. Hormesis defined. Ageing Res Rev 2008; 7:1–7.

56. Casadevall A, Coelho C, Cordero RJB, Dragotakes Q, Jung E, Vij R, P W. The
capsule of *Cryptococcus neoformans*. Virulence 2018; 0:1–10.

57. Desalermos A, Fuchs BB, Mylonakis E. Selecting an invertebrate model host for
the study of fungal pathogenesis. PLoS Pathog 2012; 8:e1002451.

58. Lu M, Yu C, Cui X, Shi J, Yuan L, Sun S. Gentamicin synergises with azoles
against drug-resistant *Candida albicans*. Int J Antimicrob Ag 2018; 51:107–14.

59. Sun L, Liao K, Wang D. Effects of magnolol and honokiol on adhesion, yeast-
hyphal transition, and formation of biofilm by *Candida albicans*. PLoS ONE
2015; 10:e0117695–20.

60. Wurster S, Bandi A, Beyda ND, Albert ND, Raman NM, Raad II, Kontoyiannis
DP. *Drosophila melanogaster* as a model to study virulence and azole treatment
of the emerging pathogen *Candida auris*. J Antimicrob Chemother 2019;
74:1904–10.

61. Sun L, Zhi L, Shakoor S, Liao K, Wang D. microRNAs involved in the control
of innate immunity in *Candida* infected *Caenorhabditis elegans*. Nature 2016;
6:1–13.

62. Calabrese1 EJ, Baldwin LA. Hormesis: A generalizable and unifying hypothesis.
Cr Rev Toxicol 2008; 31:353–424.

63. Mlih M, Khericha M, Birdwell C, West AP, Karpac J. A virus-acquired host
cytokine controls systemic aging by antagonizing apoptosis. PLoS Biol 2018;
16:e2005796–30.

64. Woźniak A, Drzewiecka K, Kęsy J, Marczak Ł, Narożna D, Grobela M, Motała
R, Bocianowski J, Morkunas I. The influence of lead on generation of signaling
molecules and accumulation of flavonoids in pea seedlings in response to pea
aphid infestation. Molecules 2017; 22:1404–30.

65. Bektas Y, Rodriguez-Salus M, Schroeder M, Gomez A, Kaloshian I, Eulgem T.
The synthetic elicitor DPMP (2,4-dichloro-6-{(E)-[(3-
methoxyphenyl)imino]methyl}phenol) triggers strong immunity in *Arabidopsis
thaliana* and Tomato. Nature 2016; 6:1–16.

66. Rodriguez-Salus M, Bektas Y, Schroeder M, Knoth C, Vu T, Roberts P,
Kaloshian I, Eulgem T. The synthetic elicitor 2-(5-bromo-2-hydroxy-phenyl)-
thiazolidine-4-carboxylic acid links plant immunity to hormesis. Plant Physiol

2015; 170:444–58.

67.   Drexler AL, Pietri JE, Pakpour N, Hauck E, Wang B, Glennon EKK, Georgis M,
631   Riehle MA, Luckhart S. Human IGF1 regulates midgut oxidative stress and
632   epithelial homeostasis to balance lifespan and *Plasmodium falciparum* resistance
633   in *Anopheles stephensi*. PLoS Pathog 2014; 10:e1004231–14.

634   68.   Kullik SA, Sears MK, Schaafsma AW. Sublethal effects of Cry 1f Bt corn and
635   clothianidin on Black Cutworm (lepidoptera: noctuidae) larval development. Jnl
636   Econ Entom 2011; 104:484–93.

637   69.   Cotter G, Doyle S, Kavanagh K. Development of an insect model for the *in vivo*
638   pathogenicity testing of yeasts. FEMS Immunol Med Microbiol 2000; 27:163–9.

639   70.   Kochi Y, Matsumoto Y, Sekimizu K, Kaito C. Two-spotted cricket as an animal
640   infection model of human pathogenic fungi. DD&T 2017; 11:259–66.

641   71.   Panpetch W, Somboonna N, Bulan DE, Issara-Amphorn J, Finkelman M,
642   Worasilchai N, Chindamporn A, Palaga T, Tumwasorn S, Leelahavanichkul A.
643   Oral administration of live- or heat-killed *Candida albicans* worsened cecal
644   ligation and puncture sepsis in a murine model possibly due to an increased
645   serum (1→3)-β-D-glucan. PLoS ONE 2017; 12:e0181439–15.

646   72.   Wellington M, Dolan K, Krysan DJ. Live *Candida albicans* suppresses
647   production of reactive oxygen species in phagocytes. Infect Immun 2008;
648   77:405–13.

649   73.   Means TK, Mylonakis E, Tampakakis E, Colvin RA, Seung E, Puckett L, Tai
650   MF, Stewart CR, Pukkila-Worley R, Hickman SE, et al. Evolutionarily
651   conserved recognition and innate immunity to fungal pathogens by the scavenger
652   receptors SCARF1 and CD36. J Exp Med 2009; 206:637–53.

653   74.   Netea MG, Brown GD, Kullberg B-J, Gow NAR. An integrated model of the
654   recognition of *Candida albicans* by the innate immune system. Nat Rev Micro
655   2008; 6:67–78.

656   75.   Liu Y, Shetty AC, Schwartz JA, Bradford LL, Xu W, Phan QT, Kumari P,
657   Mahurkar A, Mitchell AP, Ravel J, et al. New signaling pathways govern the
658   host response to *C. albicans* infection in various niches. Genome Res 2015;
659   25:679–89.

660   76.   Blankenship JR, Heitman J. Calcineurin is required for *Candida albicans* to
661   survive calcium stress in serum. Infect Immun 2005; 73:5767–74.

662   77.   Bruno VM. Regulation of azole drug susceptibility by *Candida albicans* protein
663   kinase CK2. Mol Microbiol 2005; 56:559–73.

664   78.   Smith DA, Nicholls S, Morgan BA, Brown AJP, Quinn J. A conserved stress-
665   activated protein kinase regulates a core stress response in the human pathogen
666   *Candida albicans*. Mol Biol Cell 2004; 15:4179–90.

79.   Sniegowski P, Dombrowski P. *Saccharomyces cerevisiae* and *Saccharomyces*

paradoxus coexist in a natural woodland site in North America and display
different levels of reproductive isolation. FEMS Yeast Res 2002; 1:299–306.

80. Pitt JI, Mycologia MM, 1968. Sporulation in *Candida pulcherrima*, *Candida*
reukaufii and *Chlamydozyma* species: their relationships with *Metschnikowia*.
Mycologia 2018; 60:663–85.

81. Dujon B, Sherman D, Fischer G, Durrens P, Casaregola S, Lafontaine I, De
Montigny J, Marck C, Neuvéglise C, Talla E, et al. Genome evolution in yeasts.
Nature 2004; 430:35–44.

82. Ben-Ami R, Berman J, Novikov A, Bash E, Shachor-Meyouhas Y, Zakin S,
Maor Y, Tarabia J, Schechner V, Adler A, et al. Multidrug-resistant *Candida*
haemulonii and *C. auris*, Tel Aviv, Israel. Emerg Infect Dis 2017; 23:1–9.

83. Wong B, Perfect JR, Beggs S, Wright KA. Production of the hexitol D-mannitol
by *Cryptococcus neoformans in vitro* and in rabbits with experimental
meningitis. Infect Immun 1990; 58:1664–70.

Appendix – Food preparation

Premix		Diet cake	
Amount	Ingredient	Amount	Ingredient
2,700 g	Wheatgerm	336 g	Premix
1,260 g	Casein	1,770 ml	Distilled water
1,080 g	Sucrose	22.5 g	Agar
540 g	Dried active yeast	4 ml	Corn oil
360 g	Wesson's Salt	4 ml	Linseed oil
36 g	Choline chloride	8 ml	4% Formaldehyde
72 g	Cholesterol	0.2 g	Chlorotetracycline
36 g	Methyl paraben	0.2 g	Vanderssant vitamins
54 g	Sorbic acid	8 g	Ascorbic acid

For the premix, deactivate the yeast by microwaving for 5 minutes on low power
before mixing all components thoroughly. Store in a cool, dry place.

For the diet cake, heat up 650 ml of water on a hot plate while melting the agar in 1 l
of water by microwaving. Combine the agar with the pre-warmed water, 336 g of premix,
formaldehyde and oils and mix thoroughly using a stand mixer. Dissolve vitamins, antibiotic,
and ascorbic acid in the remaining 30 ml of water and add to mixing bowl once the content
has cooled below 50°C to prevent inactivation of vitamins and antibiotic. Line a large ice
cube tray with sterile aluminium foil (sterilise by spraying with 70% ethanol) and pour
mixture into tray. Let the diet mix set for about 1.5 hours, wrap tightly in aluminium foil and
store at 4°C. Keeps for 3 weeks.

Tables

Table 1: Yeast strains used in this study.

Strain ID	Strain Name/genotype	Species	Source
YSD89	SN95	*C. albicans*	
YSD85	SC5314	*C. albicans*	
YSD883	*hog1Δ/Δ*	*C. albicans*	This study
YSD87	BWP17	*C. albicans*	[76]
YSD190	VIC84 (*cka2Δ/Δ*)	*C. albicans*	[77]
YSD192	VIC93 (*cka2/cka2::CKA2*)	*C. albicans*	[77]
YSD302	CAS12 (*ahr1Δ/Δ*)	*C. albicans*	[52]
YSD303	CAS13 (*ahr1/ahr1::AHR1*)	*C. albicans*	[52]
YSD304	RM1000	*C. albicans*	[78]
YSD305	JC52 (*hog1/hog1::HOG1*)	*C. albicans*	[78]
YSD306	JC50 (*hog1Δ/Δ*)	*C. albicans*	[78]
YSD790	YPS143	*S. cerevisiae*	[79]
YSD1448	NCYC2580	*M. pulcherrima*	[80]
YSD465	2001	*C. glabrata*	[81]
YSD1454	TA004-14	*C. auris*	[82]
YSD1028	H99	*C. neoformans*	[83]

701 **Table 2:** Statistical significance values of survival of animals treated with antifungal drugs
702 compared to those not receiving treatment.
703

Fluconazole		Caspofungin	
Concentration	p-value	Concentration	p-value
1 mg/kg	0.54	1 mg/kg	0.8
4 mg/kg	0.63	2 mg/kg	0.015
16 mg/kg	0.68	4 mg/kg	0.0022
32 mg/kg	0.27		
64 mg/kg	0.13		

704
705

Figure Captions

Figure 1: *M. sexta* caterpillars are susceptible to infections with *C. albicans*. a) *M. sexta* fifth instar caterpillar prior to injection weighing ~2 g. **b)** 24 hours post injection, the dead animal on the left has lost colour and turgidity compared to the live animal on the right. **c)** Survival curves of animals infected with *C. albicans* SN95 or SC5314. Killing occurs in a dose-dependent manner at 10^6 and 10^7 yeast cells per animal. **d)** Survival curves of animals infected with *C. albicans* mutants with attenuated virulence phenotypes in mice or epithelial cell models. The *hog1Δ/Δ* and *cka2Δ/Δ* mutants exhibit attenuated virulence, while virulence of the *ahr1Δ/Δ* mutant is comparable to the wild type. The *hog1Δ/Δ* mutant differs significantly from the wild type (p=0.0001). The complemented strain *hog1/hog1::HOG1* kills *M. sexta* at a level comparable to that of the wild type strain (p=0.21). The *cka2Δ/Δ* mutant is significantly less virulent than the wild type (p=0.00021), while the complemented strain *cka2/cka2::CKA2* is not (p=0.054). Virulence of the *ahr1Δ/Δ* mutant and the complemented strain *ahr1/ahr1::AHR1* does not differ significantly from the wild type (*ahr1Δ/Δ* p=0.72; *ahr1/ahr1::AHR1* p=0.38).

Figure 2: Elevated temperatures increase susceptibility of *M. sexta* to *C. albicans*. a) *M. sexta* caterpillars succumb to infection with the laboratory strain SC5314 in a dose-dependent manner at 37˚C but less inoculum is required than for infections at 25˚C. **b)** The weight of caterpillars infected with 10^4 cells per animal was comparable to that of caterpillars injected with 1x PBS, while animals infected with 10^5 cells showed a significant reduction in weight. **c)** Attenuated virulence of the *hog1Δ/Δ* mutant is retained at 37˚C. **d)** Animals infected with the JC50 mutant strain gain significantly more weight than those infected with the wild type strain RM1000. Infection with the complemented strain JC52 resulted in a comparable lack of weight gain.

Figure 3: Fungal burden in faeces and haemolymph increases throughout the course of infection. Colony forming units per gram faeces **(a)** and 100 μl haemolymph **(b)** in the *hog1Δ/Δ* mutant (YSD883), the wild type (SN95) and the control animals. Counts increase over time in the wild-type infected animals but not the ones injected with the *hog1Δ/Δ* mutant or PBS. **c)** Weight increases in animals infected with *hog1Δ/Δ* comparable to those injected with PBS, while animals injected with the wild type *C. albicans* strain experience reduced weight gain. The peak and drop in weight observed in the *hog1Δ/Δ* and PBS groups, marked

with a ⊙ is coinciding with the onset of pupation. This 'pupation drop' is due to the animals
refraining from food upon entering the early stages of pupation.

Figure 4: Antifungal efficacy testing of fluconazole and caspofungin. Groups of ten *C.*
albicans infected caterpillars were treated with increasing doses of antifungal drug. **a)**
Survival improves upon treatment with fluconazole, while weight remains largely stagnant in
surviving animals **(b)**. **c)** Caspofungin treatment has a positive effect on *M. sexta* survival and
weight **(d)**. Weight data were not analysed for significance due to the lack of surviving
animals in the no treatment group.

Figure 5: Caterpillars are susceptible to common yeast pathogens. Groups of ten animals
were infected with increasing doses of yeasts and survival was recorded daily. Caterpillars
are not susceptible to *S. cerevisiae* or *M. pulcherrima* but infections with *C. neoformans*, *C.*
glabrata, *C. auris* and *C. albicans* result in significantly reduced survival rates.

Figure 6: Caterpillar weight as a measure of virulence. Recording caterpillar weight daily
revealed that yeast infections affect weight regardless of mortality rates. Daily weight
measures were compared to that of PBS-injected animals and statistical significance assessed
as follows: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure 7: Mouse homologs of *M. sexta* differentially expressed genes. Numbers of
Manduca genes and their murine homologs that were up- or down-regulated in mouse vagina,
kidney or tongue and the overlap between the different data sets as depicted in a Euler
diagram. Note, in some cases the same mouse gene was the best BLAST match for several *M.*
sexta genes.

Figure 1 - Lyons et al.

Figure 2 - Lyons et al.

Figure 3 - Lyons et al.

Figure 4 - Lyons et al.

a)

b)

c)

d)

Figure 5 - Lyons et al.

Figure 6 - Lyons et al.

Figure 7- Lyons et al.

up-regulated

Ces2f
Slc6a14
Ces2f
Ces2f
Ces2f
Ces2f

Serpinb3c
Serpinb3b
Rit2
Tmprss11a
Tmprss11f
Hdc
Serpinb3b
Pglyrp3
Tmprss11f

Hif3a
Adam8

Gria2
Kcnj3
Fut9
Cbl

Gm14295
Tuba1c

Cd53
Gpr84
Meltf
Galnt6

down-regulated

Hacd1
Eno3
Pgam2
Lipc
Abcb4

Pygm
Grem1
Mep1a
Cnn1

Ugt2b34
Fasn

Acsm1

Vagina   Kidney   Tongue

Supplemental Online Material

Figure S1 - Lyons et al.

a)   b)

Figure S1: Caterpillar growth is comparable between standard colony temperature and host body temperature. Groups of ten animals were kept on standard diet at 25˚C and 30˚C **(a)** or 25˚C and 37˚C **(b)** for four days. Animals were scored daily for survival and weight as a measure of development. Within these parameters, caterpillar development is comparable across all three temperatures.

Figure S2 - Lyons et al.

Figure S2: Heat-killed *C. albicans* cells do not affect caterpillar survival or weight gain.
Groups of ten animals were injected with either live *C. albicans* wild-type YSD85 cells, heat-killed cells, or PBS. **a)** Only live yeast cells kill caterpillars. **b)** The weight gain in animals infected with heat-killed *Candida* cells is comparable to that of animals injected with PBS.

Figure S3 - Lyons et al.

Figure S3: Mapping distribution of RNAseq reads. Between 17.8 M and 34.9 M read pairs per sample mapped either to the *M. sexta* or the *C. albicans*.

Figure S4 - Lyons et al.

Figure S4: *M. sexta* **genomic expression programs in response to infection with *C. albicans*.** Hierarchical clustering of global gene expression patterns recorded in samples from uninfected control animals (PBS, samples A4, D4, E4) and those infected with the *C. albicans* wild type (samples A5, B5, C5). Listed are 322 differentially expressed genes on the y-axis in corresponding order to Table S1. In the inset, count refers to the number of differentially expressed genes and value denotes their magnitude of difference.