Pupil responses to pitch deviants reflect predictability of melodic sequences.

Bianco, Roberta[1], Ptasczynski, Lena Esther[2], Omigie, Diana[2]

[1] Ear Institute, University College London, [2] Goldsmiths, University of London,

* Correspondence:
Roberta Bianco
r.bianco@ucl.ac.uk
7

Keywords: music, melodies, contextual entropy, deviants, pupillometry

ABSTRACT

Humans automatically detect events that, in deviating from their expectations, may signal prediction
failure and a need to reorient behaviour. The pupil dilation response (PDR) to violations has been
associated with subcortical signals of arousal and prediction resetting. However, whether and how PDR
to a deviant is modulated by the structure of the proximal context remains unexplored. Using
ecologically valid musical stimuli that we characterised using a computational model, we showed PDR
(to behaviourally irrelevant deviants) to be sensitive to contextual uncertainty (quantified as entropy),
whereby PDR was greater in low than high entropy contexts. PDR was also positively correlated with
the unexpectedness of deviants in high entropy contexts. No effects of music expertise were found,
suggesting a ceiling effect due to enculturation. These results show that the same sudden
environmental change can lead to differing arousal levels depending on contextual factors, providing
evidence for a sophistication of the PDR to the structure of the sensory environment.

Introduction

The experience of surprise is very common in the sensory realm, and often triggers automatic changes
in the arousal and attentional states that are fundamental to adaptive behaviours. Music is a
ubiquitous and ecological example of a situation where changes in listeners' arousal and attention are
intentionally manipulated. Composers may, for example, modulate the predictability of musical
passages in order to achieve differing levels of tension in a listener. A great deal of empirical work has
shown that surprising sounds are recognised by listeners in an effortless and automatic fashion
(Pearce, 2018). This process is thought to be supported by a mismatch between the current
unexpected input and the implicit expectations made possible by schematic and dynamic knowledge
of stimulus structure (Huron, 2006; Krumhansl, 2015; Tillmann, Bharucha, & Bigand, 2000; Vuust &
Witek, 2014). However, there is still rather little research examining mismatch responses under
different degrees of uncertainty during passive listening.

Evidence of listeners experiencing events in music as unexpected comes from studies investigating
behavioural (Marmel, Tillmann, & Delbé, 2010; Omigie, Pearce, & Stewart, 2012; Tillmann & Lebrun-
Guillaud, 2006) and brain responses to less regular musical events (Bianco, Novembre, Keller, Kim, et
al., 2016; Carrus, Pearce, & Bhattacharya, 2013; Koelsch, 2016; Koelsch, Gunter, et al., 2002; Maess,
Koelsch, Gunter, & Friederici, 2001; Miranda & Ullman, 2007; Omigie, Pearce, Williamson, & Stewart,
2013; M.T. Pearce, Ruiz, Kapasi, Wiggins, & Bhattacharya, 2010). With regard to the former, priming
paradigms have shown that a context allows perceivers to generate implicit expectations for future

events, leading to facilitated processing (i.e., priming) of expected events. With regard to the latter, violation paradigms have shown increased brain responses to deviant events (out of key notes, or harmonically incongruent chords) within structured contexts as well as events which are musically plausible but more improbable in the given context. For example, Omigie et al. (2013) tested brain responses to melodies whose notes were characterised in terms of their predictability by a model of auditory expectations (Pearce, 2005). They showed that surprising events (more improbable notes) within melodies elicited a mismatch response – often termed the mismatch negativity, MMN (Garrido, Kilner, Stephan, & Friston, 2009; Näätänen, Paavilainen, Rinne, & Alho, 2007). This response decreased in amplitude for progressively more predictable events, as estimated by a computational model of melodic expectation. A similar parametric sensitivity to note unexpectedness has since also been observed in subcortical regions like the anterior cingulate and insula (Omigie et al., 2019). Moreover, sensitivity to music structure violation seems to emerge in all members of the general population that have had sufficient exposure to a given musical system (Bigand & Poulin-Charronnat, 2006; Pearce, 2018; Rohrmeier, Rebuschat, & Cross, 2011), and this sensitivity is modulated by pre-existing schematic knowledge of music, as reflected in acquired levels of musical expertise (Fujioka, Trainor, Ross, Kakigi, & Pantev, 2004; Koelsch, Schmidt, & Kansok, 2002a; Tervaniemi, 2009; Vuust, Brattico, Seppänen, Näätänen, & Tervaniemi, 2012).

According to theoretical and empirical work framing perception in the context of predictive processing, the experience of surprise may be modulated by the predictability of a stimulus's structure as it unfolds (Clark, 2013; Friston, 2005; Ross & Hansen, 2016). Random or high entropic stimuli hinder the possibility of making accurate predictions about possible upcoming events, whilst stimuli characterized by familiarity or regular statistics will enable relatively precise predictions by permitting the assignment of high probability to a few possible continuations. Perceptually, it has been shown that listeners indeed experience high-entropic musical contexts with greater uncertainty compared to low entropy ones (Hansen & Pearce, 2014). Moreover, previous work has shown that neurophysiological signatures associated with auditory surprise display larger responses to a given deviant event when it is embedded in a low rather than high entropic context (Garrido, Sahani, & Dolan, 2013; Hsu, Le Bars, Hamalainen, & Waszak, 2015; Quiroga-Martinez, Hansen, Højlund, Pearce, Brattico, & Vuust, 2019; Rubin, Ulanovsky, Nelken, & Tishby, 2016; Southwell & Chait, 2018). Therefore, research suggests that to understand whether and how surprising events modulate arousal and re-orient behaviours, the statistics of the proximal context must be considered.

A vast literature has used pupil dilation response (PDR) as a general marker of arousal, selective attention, and surprise (Aston-Jones & Cohen, 2005; Sara, 2009). Pupil dilation is associated with the locus coeruleus- norepinephrine (LCN) system (Laeng, Sirois, & Gredeback, 2012; Widmann, Schröger, & Wetzel, 2018), the activation of which results in wide spread norepinephrine release in the brain. Increase of PDR has been extensively reported in response to violation of expectations or surprising/salient events even in distracted listeners (Damsma & van Rijn, 2017; Liao, Yoneya, Kidani, Kashino, & Furukawa, 2016; Wetzel, Buttelmann, Schieler, & Widmann, 2016; Zhao et al., 2018). Importantly, PDR has been specifically associated to violations of statistical regularities in the sensory input when these violations are behaviourally irrelevant (Alamia, VanRullen, Pasqualotto, Mouraux, & Zenon, 2019; Zhao et al., 2018). These results provide supporting evidence for a role of norepinephrine in the tracking of abrupt deviations from the current model of the world, and as a signal of prediction resetting that enables the discovery of new information (Dayan & Yu, 2006).

Music has the ability to play with our expectations, hence manipulating our arousal and emotions in an automatic fashion (Laeng, Eidet, Sulutvedt, & Panksepp, 2016; Meyer, 2001; Zatorre & Salimpoor, 2013). Abrupt changes in register, texture and tonality are all examples of instances where the listener may have to reset or potentially abandon current models about the unfolding music. Such changes may however appear less surprising if embedded in high entropy contexts. Here, we use music as an ecological setting with which to study how PDR to deviant musical events is modulated by structure of the stimulus context, specifically by its predictability. The growing field of computational musicology means that information in melodies can be statistically estimated. One particular model of auditory expectations – the Information Dynamics of Music, IDyOM (Pearce, 2005) – has been shown to model listeners' experience of surprise and uncertainty. This unsupervised Markov model learns and estimates the conditional probability of each subsequent note in a melody based on a corpus on which it has been trained (long-term sub-model; extra-opus-learning) and the given melody as it unfolds (short-term sub-model; intra-opus learning). The model outputs information content and entropy values, which, respectively, reflect the experienced unexpectedness of a certain note after its onset and the experienced uncertainty in precisely predicting a subsequent note based on the preceding pitch probabilities.

We created novel melodies (Figure 1) that adhered to the principles guiding western tonal melodic structure. We then created shuffled versions of these melodies to create stimuli that were higher in entropy albeit matched for pitch range and content. The information theoretic properties of all melodies (shuffled and original) were estimated using IDyOM (Pearce, 2005). Listeners were presented with these melodies either in their standard form or with a pitch deviant. PDR was measured and subjective ratings about the unexpectedness of each melody were collected at the end of each sample trial. We expected larger PDR to deviant notes that are embedded in predictable rather than unpredictable contexts, and that are higher in their unexpectedness – information content value – estimated by the computational model. Also, we expected entropy of the melodies to predict subjective ratings of stimulus unexpectedness (Hansen & Pearce, 2014). Differences due to musical expertise (Müllensiefen et al., 2014) were also investigated.

Methods

Participants

Forty-seven participants (age: M=26.19, SD=6.24, min.=20, max.=52, 68% female, representing 15 nationalities) took part in the study. The sample scored relatively high on the general musical sophistication index, GoldMSI (Müllensiefen et al., 2014), with M=87.40, SD=25.24, min.=32, max.=120. A big sample size was chosen based on a previous experiment using musical stimuli (Laeng et al., 2016) and to ensure statistical power. Two participants reported ophthalmologic concerns or surgery prior to the experiment but were not excluded from participation as the pupil dilates even in blindsight participants (Weiskrantz, Cowey, & Barbur, 1999). Technical problems occurred during the recording of four participants, who were therefore excluded from the analysis. One subject was further excluded as blink gaps were too large to be interpolated. In sum, forty-two participants' data were analysed.

Ethical approval for this study was granted by the Research Ethics Committee of Goldsmiths, University of London. Participants were instructed as to the purpose of the study, and consented to participate (written informed consent). Participation was remunerated with 5 pounds.

Stimuli

One-hundred-twenty melodies were used in this study (thirty originally composed, thirty matched
'shuffled' versions, and sixty corresponding versions with a deviant tone as the 13[th] note). The melodic
sequences were 5 seconds long, isochronous (with an inter-onset-interval of 250 ms, 4/4 bar with 240
bpm), and had constant intensity and timbre (MIDI generated piano timbre).

A chromatic scale was used to compose the original melodies, whereby harmony could be ambiguous.
Ambitus and tonal space varied across melodies. Interval size did not exceed a perfect fourth
(Narmour, 2015) between adjacent notes. Thus, the original melodies were characterized by a smooth
contour. To generate matched melodies that controlled for potential biases such as pitch class, range
or ambitus, high entropy melodies were created from the original melodies by randomizing the order
of constituent notes using midi processing tools (Eerola & Toiviainen, n.d.).

Deviant notes were inserted in all 60 melodies (original and shuffled version) at the onset of the 13[th]
note (3000ms on the salient onbeat) in order to create the corresponding set of *deviant melodies* (Fig
1B). The deviant note was integrated into the second half of the melody in order to allow establishment
of an expectation-forming context before its occurrence. Interval size of the deviant varied between
maj7 up/down, min9 up/down and aug11 up/down as those intervals sound particularly unusual
within a melodic progression. Larger interval size of the deviant was assigned to matched pairs with
lower entropy differences. This allowed for a variety of deviant interval sizes while ensuring salience
of deviants even in melodies showing relatively smaller entropy differences.

Mean entropy values for each melody were assigned by the IDyOM model to each note of the
melodies. The IDyOM model considered one pitch viewpoint, namely 'cpitch', whereby chromatic
notes count up and down from the middle pitch number (C=60) (Pearce, 2005). Through a process of
unsupervised learning, the model was trained on a corpus (903 Western tonal melodies) comprising
songs and ballads from Nova Scotia in Canada, German folk songs from the Essen Folk Song Collection
and chorale melodies harmonised by J.S. Bach. The probability of each note of the stimulus used here
were then estimated based on a combination of the training set's statistics and those of the given
melody at hand. The model outputs information content and entropy values. In mathematical terms,
information content is inversely proportional to the probability of an event xi, with $IC(x_i) = -\log_2 p(x_i)$
(MacKay, 2003), while the maximum entropy is reached when all potential events xi are equally
probable, with $p(x_i)=1/n$, where n equals the number of stimuli. In psychological terms, information
content represents how surprising each subsequent note is based on its fit to the prior context Pearce
& Wiggins, 2006). In contrast, entropy refers to the anticipatory difficulty in precisely predicting a
subsequent note. Mean entropy values, obtained by taking the mean entropy of all notes, were used
to characterise the predictability of each melody. These values were then used to predict subjective
inferred uncertainty about each melody.

Procedure

The experiment was presented using the Experiment Builder Software and pupil diameter was
recorded using EyeLink 1000 eye-tracker at a 250Hz sampling rate (SR Research, www.sr-
research.com/experiment-builder). Prior to the data acquisition, a three-dot calibration was
conducted to ensure adequate gaze measurements. Participants were further allowed to adjust the
sound volume to a comfortable level and were asked to reduce head movements to a minimum
throughout the recording session. As no differences were anticipated between the left and right pupil,
the left pupil was recorded in ten and the right pupil in 32 participants depending on the participants'

dominant eye. To reduce motion artefacts, the head was stabilized using the SR Research Head Support
chinrest placed 50 cm from the presentation screen.

During the experiment, the 120 melodies were presented binaurally through headphones in a
randomized order. Each trial was triggered by the experimenter on the control computer when the
fixation was stable at less than two arbitrary gaze units away from the fixation point. When recording
was enabled, a white fixation dot on the grey screen turned black to prepare participants for the onset
of the melody. Each trial was preceded by a 400 ms baseline period and followed by a 2400 ms post
stimulus offset.

Participants were instructed to carefully listen to the melody while fixating on the fixation point in the
centre of the screen throughout the recording period. After each trial, participants rated the final note
on a Likert-scale ranging from 1 (not at all unpredictable) to 7 (extremely unpredictable) in a forced-
choice task on the presentation screen. Data on the subjects' musical expertise and sociodemographic
background was collected at the end of each experiment using the GoldMSI (Müllensiefen et al., 2014).
The whole study lasted approximately one hour.

Data pre-processing

Blinks were identified and removed from the signal using MATLAB R2017b. These were characterized
by a rapid decline towards zero from blink onset, and a rapid rise back to the regular value at blink
offset. 100ms of signal was removed before and after the missing data points (Troncoso, Macknik, &
Martinez-Conde, 2008) and missing data were interpolated: four equally spaced time points were used
to generate a cubic spline fit to the missing time points between blink onset (t2) and blink offset (t3)
of the unsmoothed signal, with t1= t2-t3+t2 and t4= t3-t2+t3 (Mathôt, Fabius, Van Heusden, & Van der
Stigchel, 2018). Trials containing more than 15% missing data were excluded from the analysis (*M*=
0.3, *SD* = 0.9 trials across subjects). Data were cleaned of artefacts using Hampel filtering (median
filtered data (Pearson, Neuvo, Astola, & Gabbouj, 2016), and smoothened using a Savitzky-Golay filter
of polynomial order 1 over the entire trial epoch. Finally, data were z-scored, and then baseline-
corrected by subtracting the median pupil size of 400 ms baseline before melody onset (first analysis)
and before deviant onset (second analysis). The normalized pupil diameter was time-domain-averaged
across trials of each condition.

Experimental design and statistical analysis

We estimated a single time series for each sequence type: predictable or unpredictable with either
standard or deviant note type (sp/dp/su/du) by averaging across trials and participants. Statistical
analysis was performed using Fieldtrip's cluster-based permutation test (Maris & Oostenveld, 2007),
with a significance threshold at 5% to control family-wise error-rate (FWER). This analysis revealed the
time windows showing significant difference between the compared time series.

We first compared responses to deviant notes in the predictable and unpredictable contexts with the
corresponding standard notes using signals recorded across the entire melody duration (baselined 400
ms before melody onset). This ensured that differences between deviant and corresponding standards
were not due to differences in the immediately preceding note. Then, to determine how the deviant
PDR is affected by the entropy of the melodic context, we focussed on the time window from deviant
onset to the end of the melody (3000 to 5000 ms epochs baselined to 400 ms before deviant onset),
and we tested for an interaction of the deviant and context manipulation: (*dp - sp*) vs. (*du - su*). Further,

we compared the responses to standard tones in the two contexts (*sp - su*) to ensure that any
differences were not driven simply by the standard notes (the control conditions).

To assess a potential influence of expertise on PDR to deviants (baselined data between 3000 and 5000
ms), we computed the mean PDR to deviant trials as *dp – sp* for deviants in predictable context, and
du – su for deviants in unpredictable context. Participants were split into two groups of musical experts
and non-experts based on GoldMSI scores (*Mdn*=96). An ANOVA with within-subject factor context
(predictable/unpredictable) and expertise as between-subject factor (expert/non-expert) was
computed.

Results

Analyses were carried out to clarify the nature of all differences in information theoretic properties of
the different stimuli. Figure 1B shows that the predictable and unpredictable melodies were
characterized by significantly different mean entropy values regardless of whether they contained or
didn't contain a deviant note (dp: *M*=10.83, *SD*=5.31; du: *M*=11.41, *SD*=4.38; sp: *M*=4.32, *SD*=2.80; su:
M=4.85, *SD*=2.70). Indeed, an ANOVA with context (predictable / unpredictable) and note type
(deviant / standard) as between-group factors yielded a main effect of context [$F(1,116)$=15.56, *p* <
.001, *np2* = .12], a non-significant main effect of note type [$F(1,116)$=1.34, *p* = .249, *np2* = .01], and no
interaction between the two factors [$F(1,116)$=0.10, *p* = .757, *np2* < .01]- thus indicating higher entropy
in unpredictable than predictable melodies [$t(58)$ = 2.85, *p* = 0.006].

Figure 1C shows that the unexpectedness of deviant notes, as reflected by information content values,
was comparable between predictable and unpredictable melodies. An ANOVA with between group
factors context (predictable / unpredictable) and note type (deviant / standard) yielded a main effect
of note type [$F(1,116)$=82.97, *p* < .001, *np2* = .42], a non-significant main effect of context
[$F(1,116)$=0.68, *p* = .410, *np2* = .01], and no interaction between the two factors [$F(1,116)$=0.00, *p* =
.977, *np2* < .01]- thus indicating higher information content in deviant than standard notes regardless
of the context in which they were embedded [$t(58)$ = -8.069, *p* < 0.001].

Fig. 1. Experimental design including two factors: Contextual entropy (predictable/unpredictable) and Note type (standard/deviant). Subjective ratings about the unexpectedness of each melody were collected at the end of each sample trial. **A)** A high entropy melody example containing a pitch deviant note at the 13th note from melody onset. **B)** Mean entropy was larger for unpredictable than predictable melodies regardless of the presence of deviant (yellow) or standard (blue) tones. **C)** Mean information content of deviants (yellow) was larger than standard (blue) tones regardless of the predictability of the context.

Figure 2A shows the time course of the PDR across conditions (*sp* = standard predictable, *su* = standard unpredictable, *dp* = deviant predictable, *du* = deviant unpredictable) baselined 400 ms before melody onset. A comparison between *sp* and *su* showed no difference in PDR as a function of entropy levels of the melody, whilst the PDR to deviants (*dp* vs. *du*) was greater in predictable than unpredictable contexts (diverging at .56 s from deviant onset). The responses to deviants in the predictable contexts was greater than the relative standard condition (*dp* vs. *sp*: $p = .029$), significantly diverging from *sp* at 3.57 s after melody onset (0.57 s after deviant onset). Conversely, the responses to deviants in unpredictable contexts didn't differ from the relative standard condition (*du* vs. *su)*, despite their high information content (see figure 1C). Figure 2B shows the PDR to deviants baselined 400 ms before deviant onset. We show the conditions *dp* and *du* following subtraction of the relative standard conditions. The comparison between the two time-courses confirmed that *dp* evoked a larger response than *du* starting at .59 s from deviant onset ($p = .007$).

Figure 3B shows the relationship between the deviant-related PDR and deviant unexpectedness (information content) for predictable and unpredictable melodies. For each subject and for each melody containing deviants, the average PDR to deviants was computed relative to the deviant onset. These values showed a positive correlation with deviant information content values when embedded in unpredictable ($rho = 0.467$, $p = 0.009$), but not in predictable melodies ($rho = 0.264$, $p = 0.158$). Thus,

a dissociation between unexpectedness of deviants and entropy of the context was shown, whereby PDR is sensitive to a large range of unexpectedness levels in low entropic contexts, whilst in high entropic contexts it responds only to particularly larger deviants.

We further investigated potential influences of expertise on mean PDR to deviants (computed as *dp* – *sp* for deviants in predictable context, and *du* – *su* for deviants in unpredictable context). The ANOVA examining the effect of musical expertise on mean PDR to deviants yielded a main effect of context [$F(1,40)$ = 4.53, p = .040, $np2$ = .10]. The main effect of expertise was not significant [$F(1,40)$ = 0.59, p = .449, $np2$ = .01], and no interaction between the two factors was seen [$F(1,40)$ = 2.23, p = .143, $np2$ = .05]. This indicates that mean PDR to deviant was greater in the predictable than unpredictable contexts to an equal extent in both experts and non-experts.

Finally, we showed that the model reliably predicted subjective uncertainty levels (inferred entropy) of the melody progressions (Hansen & Pearce, 2014) (Figure 3A). This measure was collected after participants listened to each melody during the experimental session. They were asked to carefully listen to the melodies and rate the last note on a seven-step Likert scale – 1 equal 'not at all unpredictable' and 7 'extremely unpredictable'. The results of Fig. 3A showed that mean ratings for each melody strongly correlated with mean IDyOM entropy values.

Fig. 2. A) PDR for all conditions from melody onset (*sp* = standard predictable, *su* = standard unpredictable, *dp* = deviant predictable, *du* = deviant unpredictable). PDR to deviants compared to standard tones (*dp* vs. *sp*) in predictable contexts diverged at 3.57 s from melody onset (.57 s from deviant onset), but did not differ in unpredictable contexts. Also, the PDR to deviants was greater in predictable than unpredictable contexts (*dp* vs. *du*) (diverging at 3.056 s from melody onset), but there was no significant context-dependent difference between the standard tones (*sp* and *su*). **B)** Interaction effect of deviant and context predictability on PDR after deviant onset. The difference between PDR to deviant and standard tones was greater in the predictable than unpredictable contexts. This effect emerged .59 after deviant onset.

Fig. 3. A) The stimulus entropy was computed by the IDyOM model, and validated by participants' self-
reports about the overall unexpectedness of each melody. Each dot represents one of the 120
melodies. **B)** The unexpectedness of deviants were estimated by the IDyOM model and showed a
positive correlation (Spearman) with mean evoked PDR in the unpredictable *(upper panel)*, but not in
the predictable melodies *(lower panel)*. Each dot represents one of the 60 melodies containing a
deviant note.

Discussion

We report that pupil dilation response (PDR) to behaviourally irrelevant deviants occurs when deviants
are embedded in predictable rather than unpredictable melodies. We showed that the amplitude of
the response is predicted by the information content (or unexpectedness) of the musical deviants in
high but not low entropic contexts. We also replicate the previous finding that listeners' experience of
uncertainty is predicted by the entropy of the music (Hansen & Pearce, 2014). These results show that
the same sudden environmental change leads to differing levels of arousal depending on whether it
occurs in low or high states of uncertainty. Our results suggest that the more stable predictions formed
in predictable rather than unpredictable contexts may be more abruptly violated by surprising events,
possibly leading to greater changes in the listeners' arousal state.

The observed modulatory effect of context predictability on PDR is consistent with a body of
electrophysiological work showing context effects on mismatch like responses at the cortical level
(Garrido et al., 2013; Quiroga-Martinez et al., 2019; Southwell & Chait, 2018). Here we show a similar
pattern but in autonomic markers of arousal, as reflected by PDR. Pupil response is thought to be
driven by norepinephrine activity in the locus coeruleus (LC) (Joshi, Li, Kalwani, & Gold, 2016). This is a
key subcortical nucleus which widely connects to the brain (Sara, 2009) to signal unexpected and
abrupt contextual changes (Alamia et al., 2019; Damsma & van Rijn, 2017; Zhao et al., 2018). It has
been hypothesised (Zhao et al., 2018) that MMN-related brain systems (Garrido et al., 2009; Hsu et al.,
2015; Southwell & Chait, 2018) may trigger norepinephrine-mediated updating or interruption of
ongoing top-down expectations that are mediated by temporo-frontal networks. Being the temporo-
frontal network widely associated with MMN-like responses in music-violation paradigms (Bianco,
Novembre, Keller, Seung-Goo, et al., 2016; Koelsch, Gunter, et al., 2002; Tillmann, Janata, & Bharucha,
2003), our results are in line with a model resetting hypothesis, whereby stronger predictive models

(in the predictable melodies) supported by temporo-frontal cortical regions require a greater signal to
be interrupted.

Albeit indirectly, our results also establish a link between subcortical and cortical activity in response
to unexpected events under different states of uncertainty. Increased MMN response under low states
of uncertainty (Quiroga-Martinez et al., 2019; Southwell & Chait, 2018) are replicated in the pupil
response, reflecting a general increase in arousal. A possible interpretation, in line with the predictive
coding theory (Friston, 2005), is that more stable expectations, representative of a strong predictive
model, are reflected in precision-weighing of the prediction error, and hence a stronger response when
the input mismatches the current predictions. Conversely, in high entropic contexts predictive models
are weak, and the prediction error attenuated. This mirroring pattern between cortical (MMN) and
subcortical (as reflected by PDR) responses may have important behavioural advantages. Specifically,
strong predictive models can suddenly be abandoned when they are revealed to be erroneous, thus
allowing speedy reorienting behaviours and quicker engagement with new potentially relevant stimuli.
One limitation with regard to this possible interpretation is the nature of the deviant events used here,
whereby the deviating event was a single note that did not lead to any long lasting changes in the
statistics of the unfolding sequence. Further studies combining cortical and pupil response
measurements in continuously changing stimuli are necessary to corroborate our working hypothesis.

The absence of PDR differences between high and low entropic contexts suggests that the pupil is
relatively unresponsive to slowly unfolding stimulus structures when stimuli are behaviourally
irrelevant (Alamia et al., 2019; Zhao et al., 2018). Whilst slow cortical responses have been shown to
be sensitive to the statistics of the unfolding stimulus structure (Barascud, Pearce, Griffiths, Friston, &
Chait, 2016; Sohoglu & Chait, 2016; Southwell et al., 2017), subcortical responses to slowly evolving
structures may be indeed more vulnerable to stimulus properties and tasks demands (Zhao et al.,
2018).

A compelling finding was that the information content of the deviant predicted the size of the evoked
PDR only for deviants in the high entropy condition. This observation confirms the relative lack of
sensitivity listeners show for deviants in the unpredictable context. In other words, our results confirm
that in such high entropy contexts, only very large increases in information content are able to result
in changes in PDR. This is in contrast to low entropy environments where listener's expectations are
so precise as to lead to large PDR whenever these expectations are violated, regardless of the size of
the violation. Another interesting finding was the absence of a modulation of the PDR mismatch
response by degree of musical expertise. Larger MMN responses have been shown for musicians in a
range of studies examining electrophysiological correlates of expectancy violation in music (Koelsch,
Schmidt, & Kansok, 2002b; Oechslin, Van De Ville, Lazeyras, Hauert, & James, 2013; M. Tervaniemi,
Tupala, & Brattico, 2012). One possibility is that this reflects a ceiling effect whereby the rather salient
deviants used here were relatively easy to detect. Future studies examining PDR to more subtle
differences in musical structure may be expected to show similar expertise effects to those reported
in previous studies.

Finally, our results provide evidence of music's usefulness in investigating the neural mechanisms
underlying processing of stimuli statistical properties in ecological settings. While our work here
focused on pitch expectations, previous studies have shown that music-induced temporal expectations
are also tracked by the PDR (Damsma & van Rijn, 2017). Similarly, that music induced chills – high
arousal physiological responses associated with subjective pleasure – are associated with increased
PDR (Laeng et al., 2016) suggest a usefulness of music in examining the relationship between stimulus

information theoretic properties and reward processing. By showing that predictive uncertainty can be used to modulate prediction-error related arousal, our findings have implications for understanding the variety of forms listeners' aesthetic appreciation of music may take. However, considering, more generally, the tight coupling between the error-related norepinephrine system and the reward-seeking dopaminergic pathway (Laeng et al., 2016; Xing, Li, & Gao, 2016; Zatorre & Salimpoor, 2013), our results emphasize that measuring PDR may be useful for investigating the reward value of information across a range of modalities and domains.

In sum, we show that pupillometry in the auditory domain can reliably track the effect of context uncertainty on responses to sudden environmental change. Given the tight interplay between cortical and subcortical mechanisms involved in precision weighted anticipatory processing, a first milestone is set towards the non-invasive quantification of related arousal responses.

Competing Interests Statement

The author(s) declares no competing interests.

Author Contributions

RB conceived the experiments; analysed the bulk of the data; wrote the manuscript.

EP conceived and performed the experiments; contributed to analysis; wrote the manuscript; secured the funding.

DO conceived the experiments; contributed to analysis; wrote the manuscript; secured the funding.

Funding

This study was funded by a grant from the psychology department at Goldsmiths, University of London.

Acknowledgments

We are grateful to Prof. Maria Chait and Sijia Zhao for inspiring discussions.

Data Availability Statement

The datasets for this study can be found in the OSF repository (link: https://osf.io/u2qdr/?view_only=7a9fa6bacbb249b090282377c1542d29).

Informed Consent Statement

All participants provided written informed consent. The study was approved by the Ethics Committee at Goldsmiths, University of London.

References

Alamia, A., VanRullen, R., Pasqualotto, E., Mouraux, A., & Zenon, A. (2019). Pupil-linked arousal responds to unconscious surprisal. *The Journal of Neuroscience*, 3010–18.

Aston-Jones, G., & Cohen, J. D. (2005). An integrative theory of locus coeruleus-norepinephrine function: Adaptive Gain and Optimal Performance. *Annual Review of Neuroscience*, *28*(1), 403–450.

Barascud, N., Pearce, M., Griffiths, T., Friston, K., & Chait, M. (2016). Brain responses in humans

reveal ideal-observer-like sensitivity to complex acoustic patterns. *Proceedings of the National*
Academy of Sciences, *113*(5), E616-25.

Bianco, R., Novembre, G., Keller, P. E., Seung-Goo, K., Scharf, F., Friederici, A. D., … Sammler, D.
(2016). Neural networks for harmonic structure in music perception and action. *NeuroImage*,
142, 454–464.

Bigand, E., & Poulin-Charronnat, B. (2006). Are we "experienced listeners"? A review of the musical
capacities that do not depend on formal musical training. *Cognition*, *100*(1), 100–130.

Carrus, E., Pearce, M. T., & Bhattacharya, J. (2013). Melodic pitch expectation interacts with neural
responses to syntactic but not semantic violations. *Cortex*, *49*(8), 2186–200.

Clark, A. (2013). Whatever next? Predictive brains, situated agents, and the future of cognitive
science. *The Behavioral and Brain Sciences*, *36*(3), 181–204.

Damsma, A., & van Rijn, H. (2017). Pupillary response indexes the metrical hierarchy of unattended
rhythmic violations. *Brain and Cognition*, *111*, 95–103.

Dayan, P., & Yu, A. J. (2006). Phasic norepinephrine: A neural interrupt signal for unexpected events.
Network: Computation in Neural Systems, *17*(4), 335–350.

Eerola, T., & Toiviainen, P. (n.d.). *Midi toolbox for Matlab*.

Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal Society of*
London. Series B, Biological Sciences, *360*(1456), 815–36.

Fujioka, T., Trainor, L. J., Ross, B., Kakigi, R., & Pantev, C. (2004). Musical training enhances automatic
encoding of melodic contour and interval structure. *Journal of Cognitive Neuroscience*, *16*(6),
1010–21.

Garrido, M. I., Kilner, J. M., Stephan, K. E., & Friston, K. J. (2009). The mismatch negativity: a review of
underlying mechanisms. *Clinical Neurophysiology : Official Journal of the International*
Federation of Clinical Neurophysiology, *120*(3), 453–63.

Garrido, M. I., Sahani, M., & Dolan, R. J. (2013). Outlier Responses Reflect Sensitivity to Statistical
Structure in the Human Brain. *PLoS Computational Biology*, *9*(3).

Hansen, N. C., & Pearce, M. T. (2014). Predictive uncertainty in auditory sequence processing.
Frontiers in Psychology, *5*, 1052.

Hsu, Y.-F., Le Bars, S., Hamalainen, J. A., & Waszak, F. (2015). Distinctive Representation of
Mispredicted and Unpredicted Prediction Errors in Human Electroencephalography. *Journal of*
Neuroscience, *35*(43), 14653–14660.

Huron, D. (2006). *Sweet Anticipation : Music and the Psychology of Expectation by David Huron*. (M.
T. M. Press., Ed.), *Sweet Anticipation: Music and the Psychology of Expectation.* Cambridge.

Joshi, S., Li, Y., Kalwani, R. M., & Gold, J. I. (2016). Relationships between Pupil Diameter and
Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron*, *89*(1), 221–
234.

Koelsch, S. (2016). Under the hood of statistical learning: A statistical MMN reflects the magnitude of
transitional probabilities in auditory sequences. *Scientific Reports*, 1–11.

Koelsch, S., Gunter, T. C., v. Cramon, D. Y., Zysset, S., Lohmann, G., & Friederici, A. D. (2002). Bach
Speaks: A Cortical "Language-Network" Serves the Processing of Music. *NeuroImage*, *17*(2),
956–966.

Koelsch, S., Schmidt, B.-H., & Kansok, J. (2002a). Effects of musical expertise on the early right

anterior negativity: an event-related brain potential study. *Psychophysiology*, *39*(5), 657–63.

Krumhansl, C. L. (2015). Statistic, structures and style in music. *Music Perception*, *33*(1), 20–31.

Laeng, B., Eidet, L. M., Sulutvedt, U., & Panksepp, J. (2016). Music chills: The eye pupil as a mirror to
music's soul. *Consciousness and Cognition*, *44*, 161–178.

Laeng, B., Sirois, S., & Gredeback, G. (2012). Pupillometry: A Window to the Preconscious?
Perspectives on Psychological Science, *7*(1), 18–27.

Liao, H. I., Yoneya, M., Kidani, S., Kashino, M., & Furukawa, S. (2016). Human pupillary dilation
response to deviant auditory stimuli: Effects of stimulus properties and voluntary attention.
Frontiers in Neuroscience.

Maess, B., Koelsch, S., Gunter, T. C., & Friederici, A. D. (2001). Musical syntax is processed in Broca's
area: an MEG study. *Nature Neuroscience*, *4*(5), 540–545.

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of*
Neuroscience Methods, *164*, 177–190.

Marmel, F., Tillmann, B., & Delbé, C. (2010). Priming in melody perception: tracking down the
strength of cognitive expectations. *Journal of Experimental Psychology. Human Perception and*
Performance, *36*(4), 1016–1028.

Mathôt, S., Fabius, J., Van Heusden, E., & Van der Stigchel, S. (2018). Safe and sensible preprocessing
and baseline correction of pupil-size data. *Behavior Research Methods*, *50*(1), 94–106.

Meyer, L. B. (2001). Music and emotion: Distinction and uncertainties. In *Music and emotion: Theory*
and research. (pp. 341–360). Meyer, Leonard B.: Dept of Music, U Pennsylvania, 210 SO 34th St,
Philadelphia, PA, US, 19104: Oxford University Press.

Miranda, R. A., & Ullman, M. T. (2007). Double dissociation between rules and memory in music: An
event-related potential study. *NeuroImage*, *38*(2), 331–345.

Müllensiefen, D., Gingras, B., Musil, J., Stewart, L., Levitin, D., Hallam, S., … Winner, E. (2014). The
Musicality of Non-Musicians: An Index for Assessing Musical Sophistication in the General
Population. *PLoS ONE*, *9*(2), e89642.

Näätänen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity (MMN) in basic
research of central auditory processing: A review. *Clinical Neurophysiology*, *118*(12), 2544–
2590.

Narmour, E. (2015). Toward a unified theory of the I-R model (part 1): Parametrci scales and their
analogically isomorphic structures. *Music Perception*, *33*(1), 32–69.

Oechslin, M. S., Van De Ville, D., Lazeyras, F., Hauert, C.-A., & James, C. E. (2013). Degree of musical
expertise modulates higher order brain functioning. *Cerebral Cortex, 23*(9), 2213–24.

Omigie, D., Pearce, M., Lehongre, K., Hasboun, D., Navarro, V., Adam, C., & Samson, S. (2019).
Intracranial Recordings and Computational Modeling of Music Reveal the Time Course of
Prediction Error Signaling in Frontal and Temporal Cortices. *Journal of Cognitive Neuroscience*,
31(6), 855–873.

Omigie, D., Pearce, M. T., & Stewart, L. (2012). Tracking of pitch probabilities in congenital amusia.
Neuropsychologia, *50*(7), 1483–1493.

Omigie, D., Pearce, M. T., Williamson, V. J., & Stewart, L. (2013). Electrophysiological correlates of
melodic processing in congenital amusia. *Neuropsychologia*, *51*(9), 1749–1762.

Pearce, M. T. (2005). The Construction and Evaluation of Statistical Models of Melodic Structure in

Music Perception and Composition. *Dissertation*, (December), 267.

Pearce, M. T. (2018). Statistical learning and probabilistic prediction in music cognition: Mechanisms
of stylistic enculturation. *Annals of the New York Academy of Sciences*, *1423*, 378–395.

Pearce, M. T., Ruiz, M. H., Kapasi, S., Wiggins, G. A, & Bhattacharya, J. (2010). Unsupervised statistical
learning underpins computational, behavioural, and neural manifestations of musical
expectation. *NeuroImage*, *50*(1), 302–13.

Pearce, M. T., & Wiggins, G. A. (2006). Expectation in melody: the influence of context and learning.
Music Perception, *23*(45), 377–405.

Pearson, R. K., Neuvo, Y., Astola, J., & Gabbouj, M. (2016). Generalized Hampel Filters. *Eurasip
Journal on Advances in Signal Processing*.

Quiroga-Martinez, D. R., Hansen, N. C., Højlund, A., Pearce, M., Brattico, E., & Vuust, P. (2019).
Reduced prediction error responses in high-as compared to low-uncertainty musical contexts.
Cortex, https://doi.org/10.1016/j.cortex.2019.06.010.

Rohrmeier, M., Rebuschat, P., & Cross, I. (2011). Incidental and online learning of melodic structure.
Consciousness and Cognition, *20*(2), 214–222.

Ross, S., & Hansen, N. C. (2016). Dissociating Prediction Failure: Considerations from Music
Perception. *Journal of Neuroscience*, *36*(11), 3103–3105.

Rubin, J., Ulanovsky, N., Nelken, I., & Tishby, N. (2016). The Representation of Prediction Error in
Auditory Cortex. *PLOS Computational Biology*, *12*(8), e1005058.

Sara, S. J. (2009). The locus coeruleus and noradrenergic modulation of cognition. *Nature Reviews.
Neuroscience*, *10*(3), 211–223.

Sohoglu, E., & Chait, M. (2016). Detecting and representing predictable structure during auditory
scene analysis. *ELife*, *5*(Se), 1–17.

Southwell, R., Baumann, A., Gal, C., Barascud, N., Friston, K., & Chait, M. (2017). Is predictability
salient? A study of attentional capture by auditory patterns. *Philosophical Transactions of the
Royal Society B: Biological Sciences*, *372*(1714).

Southwell, R., & Chait, M. (2018). Enhanced deviant responses in patterned relative to random sound
sequences. *Cortex*, *109*, 92–103.

Tervaniemi, M. (2009). Musicians-Same or Different? *Annals of the New York Academy of Sciences*,
1169(1), 151–156.

Tervaniemi, M., Tupala, T., & Brattico, E. (2012). Expertise in folk music alters the brain processing of
Western harmony. *Annals of the New York Academy of Sciences*, *1252*(1), 147–151.

Tillmann, B., Bharucha, J. J., & Bigand, E. (2000). Implicit learning of tonality: A self-organizing
approach. *Psychological Review*, *107*(4), 885–913.

Tillmann, B., Janata, P., & Bharucha, J. J. (2003). Activation of the inferior frontal cortex in musical
priming. *Cognitive Brain Research*, *16*(2), 145–161.

Tillmann, B., & Lebrun-Guillaud, G. (2006). Influence of tonal and temporal expectations on chord
processing and on completion judgments of chord sequences. *Psychological Research*, *70*(5),
345–58.

Troncoso, X. G., Macknik, S. L., & Martinez-Conde, S. (2008). Microsaccades counteract perceptual
filling-in. *Journal of Vision*, *8*(14), 15–15.

Vuust, P., Brattico, E., Seppänen, M., Näätänen, R., & Tervaniemi, M. (2012). Practiced musical style
shapes auditory skills. *Annals of the New York Academy of Sciences*, *1252*(1), 139–146.

Vuust, P., & Witek, M. a. G. (2014). Rhythmic complexity and predictive coding: a novel approach to
modeling rhythm and meter perception in music. *Frontiers in Psychology*, *5*, 1111.

Weiskrantz, L., Cowey, A., & Barbur, J. L. (1999). Differential pupillary constriction and awareness in
the absence of striate cortex. *Brain*, *122*(8), 1533–1538.

Wetzel, N., Buttelmann, D., Schieler, A., & Widmann, A. (2016). Infant and adult pupil dilation in
response to unexpected sounds. *Developmental Psychobiology*, *58*(3), 382–392.

Widmann, A., Schröger, E., & Wetzel, N. (2018). Emotion lies in the eye of the listener: Emotional
arousal to novel sounds is reflected in the sympathetic contribution to the pupil dilation
response and the P3. *Biological Psychology*, *133*(January), 10–17.

Xing, B., Li, Y. C., & Gao, W. J. (2016). Norepinephrine versus dopamine and their interaction in
modulating synaptic function in the prefrontal cortex. *Brain Research*, *1641*, 217–233.

Zatorre, R. J., & Salimpoor, V. N. (2013). From perception to pleasure: music and its neural
substrates. *Proceedings of the National Academy of Sciences*, *110*, 10430–7.

Zhao, S., Chait, M., Dick, F., Dayan, P., Furukawa, S., & Liao, H.-I. (2018). Phasic norepinephrine is a
neural interrupt signal for unexpected events in rapidly unfolding sensory sequences - evidence
from pupillometry. *BioRxiv*, *42*, 466367.