HERV-K and HERV-W transcriptional activity in Myalgic Encephalomyelitis/ Chronic Fatigue Syndrome

Short title: Endogenous retroviruses in ME/CFS

Lucas S Rodrigues [1#], Luiz H da Silva Nali [1,2#], Cibele O D Leal [1], Ester C Sabino [1,3], Eliana M Lacerda[4], Caroline C Kingdon[4], Luis Nacul[4], Camila M Romano[1,5*]

1. Instituto de Medicina Tropical de São Paulo, Universidade de São Paulo, São Paulo, Brazil
2. Universidade Santo Amaro, Pós-Graduação em Ciências da Saúde
3. Departamento de Moléstias Infecciosas e Parasitárias, Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil
4. Department of Clinical Research, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom
5. Hospital das Clinicas HCFMUSP (LIM52), Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil

# These authors contributed equally to this manuscript

* Correspondence to: CMR, Laboratório de Virologia, Instituto de Medicina Tropical de São Paulo, Universidade de São Paulo, Rua Dr. Enéas de Carvalho Aguiar, 470, ZIP. 05403-000 São Paulo, SP, Brazil. phone: +55 11 3061-8668; cmromano@usp.br

Keywords: Endogenous retroviruses; HERV-W; HERV-K; myalgic encephalomyelitis; chronic fatigue

Abstract

Chronic Fatigue Syndrome / Myalgic Encephalomyelitis (CFS / MS) is an incapacitating chronic disease that dramatically compromise the life quality. The CFS/ME pathogenesis is multifactorial, and it is believed that immunological, metabolic and environmental factors play a role. It is well documented an increased activity of Human endogenous retroviruses (HERVs) from different families in autoimmune and neurological diseases, making these elements good candidates for biomarkers or even triggers for such diseases. Here the expression of Endogenous retroviruses K and W (HERV–K and HERV–W) was determined in blood from moderately and severely affected ME/CFS patients. HERV-K was overexpressed only in moderately affected individuals and HERV-W showed no difference. This is the first report about HERV-K differential expression in moderate ME/CFS.

Introduction

Myalgic Encephalomyelitis/ Chronic Fatigue Syndrome (ME/CFS) is a chronic and debilitating disease with unknown etiology [1]. Affected individuals have compromised motor and cognitive capacities. There is a wide variation in the symptoms of this disease, which include joint pains, mood disturbance, and malaise and worsening of symptoms following minimal physical or mental exertion. More severe symptoms can be also present including extreme exhaustion, severe joint pains with no apparent cause, non-restorative sleep and a range of immune and neurological symptoms. These symptoms may lead to depression and social isolation in the person with ME/CFS [1]. The pathophysiology of the ME/CFS is not understood and there is no diagnostic biomarker available. There is still controversy over the etiology of the disease; however, it is widely accepted that several immunological alterations are present in ME/CFS patients [2]. In addition, accumulated evidence for an association of ME/CFS with viral infections also exists and many patients report the onset of their symptoms during or right after a flu-like illness [3]. Thereafter, an unusual autoimmune response against the infection would be responsible for the perpetuation of the ME/CFS symptoms. Viral participation is finally supported by the evidences of clinical benefit of patients treated with valganciclovir [4]. Unfortunately, the absence of large cohort studies that investigate at the molecular level the participation of infectious agents on the ME/CFS pathogenesis impairs our understanding of this disease.

Human Endogenous Retroviruses (HERVs) are derived from exogenous retroviral infections, which occurred early in the evolution of vertebrates. Due to active replication and transposition events, HERVs are extensively distributed through the host

genome and constitute about 8% of the human genome [5]. Due to accumulated mutations over the primate and human evolution, most HERVs are non-functional, but intact open reading frames of some HERVs persist and can be reactivated in response to systemic and environmental factors such as hormones, stress, and infection by exogenous viruses including almost all human herpesviruses, HIV and others [6,7]. Given their potential pathogenic effects, which include molecular mimicry and immune deregulation, HERVs are often postulated as possible causes of autoimmune diseases. Among the more than 30 families, the K and W families are the most recently integrated, the most active, and have been frequently associated with neurological and autoimmune diseases such as multiple sclerosis, diabetes mellitus SLE, ALS and rheumatoid arthritis [8].

To our knowledge, only two studies have investigated the participation of endogenous retroviruses in ME/CFS with contrasting results [9,10].

Given the extensively described altered patterns of HERVs in several diseases and the gap in knowledge of its expression in ME/CFS, we investigated the expression of the HERVs K and W in patients diagnosed with ME/CFS.

Methods

Participants

 We used PBMC samples from a hundred patients diagnosed with ME/CFS and stored in the UK ME/CFS Biobank (UKMEB) at the London School of Hygiene and Tropical Medicine in this study. The UKMEB is among the few biorepositories worldwide with

advanced storage and linked research infrastructure dedicated to research into ME/CFS

[11]. Seventy five samples were requested from participants diagnosed with moderate

fatigue (ME/CFSm), and 25 from participants with severe fatigue (ME/CFSs). Samples

from 70 healthy controls (also provided by the UKMEB) were included. This study was

approved by LSHTM and University of São Paulo ethical committees [#EC.2017.02

and #2728254 respectively].

RNA extraction and Real Time PCR

RNA extraction from the PBMC samples was performed by the Trizol-chloroform

method, with 1ml Trizol and subsequent addition of chloroform to solubilize lipids

allowing its removal. The samples were centrifuged at 15,000 rpm for 15 minutes and

the upper phase containing the RNA was further used. The material was precipitated

with Isopropanol 100% and washed with 75% Ethanol. In both steps the material was

centrifuged at 15,000 RPM for 10 minutes at 4 °C. After this process, the pellets were

dried at room temperature for 10 minutes, and the RNA was eluted in 45μl of Nuclease-

Free H2O. The decontamination of remnant DNA was performed using two rounds

DNAse treatment (Turbo DNA-Free (Ambion) following the manufacturer's

instruction. The absence of DNA was confirmed by real time PCR without reverse

transcriptase using primers for HERV-K or HERV-W (see primers description bellow).

After this procedure, cDNA was synthesized using the High capacity cDNA Reverse

Transcription kit (Ambion, USA) according the manufacturer's instructions. Real-time

PCRs were performed for the HERV-W, - K and the endogenous gene using the primers

and conditions used previously by Nali et al [12] using the Sybr Green method. The

primers used are described in Table 1.

Table 1. Primers used in real-time PCR assays

Oligo	Sense	Antisense
HERV-W	CCAATGCATCAGGTGGGTAAC	GAGGTACCACAGACAAAAAATATTCCT
HERV-K	TCCCCTTGGAATACTCCTGTTTT	CATTCCTTGTGGTAAAACTTTCCA
GAPDH	ACCCACTCCTCCACCTTTGAC	TGTTGCTGTAGCCAAATTCGTT

The cycling conditions for both HERVs detection were: 50 °C for 2 minutes, 95 °C for

10 minutes, followed by 40 cycles of 95 °C for 15 seconds, 50 °C for 1 minute, 60 °C

for 1 minute. HERV activity was qualitatively (referred as presence/absence) and

quantitatively (level of expression) evaluated. As positive controls we used a plasmid

containing both HERV-W envelope and HERV-K polymerase fragments correspondent

to the region covered by the primers used. The level of expression was determined by

calculation of $2^{-\Delta\Delta Ct}$, and the results were represented as fold changes. Statistical analysis

was performed using the Wilcox test in the GraphPad Prism program v.6.04.

Samples were only considered positive for HERVs and included in the analyses if

expression of the endogenous control was also detected.

Results

General description of individuals included in the study is described in Table 2. As

expected, women were 4 times more prevalent than men. Therefore, we adjusted the

control group to the same gender prevalence.   HERV-K and W expression were

evaluated in ME/CFS patients and healthy controls; and some level of expression of

HERV-W was detected in all patients with severe fatigue and in 72/75 ME/CFSm

(96%). HERV-K was also detected in all severe cases but in 65/75 of moderate cases

(86.6%). The healthy control group was very similar to the moderate group, with 68/70

(97%) and 60/70 (85.7%) presenting expression of HERV-W and HERV-K respectively

(Table 2). Only one patient with moderate fatigue and one control individual had no

HERV activity at all. No relation was observed regarding HERV detection and duration

of disease.

Table 2. Main characteristics of individuals included in the study.

Participants (#)	Age median (max, min)	Gender (%)		Time of disease median (max, min)	Detection of HERV activity*	
		M	F		HK	HW
ME/CFSs (25)	43.3 (25-62)	24%	76%	16.8 (2.8-40)	100%	100%
ME/CFSm (75)	22.9 (18-64)	25,4%	74,6%	11.2 (0.2 – 33.7)	86.6%	96%
Controls (70)	42.8 (19-63)	25.7%	74.3%	-	85.7%	97%

* Qualitatively

Regarding to the level of expression (quantitative analysis), real time results revealed

that HERV-W did not present significant differences when the healthy controls (HCs)

or the two ME/CFS groups were compared between each other (Figure 1 A), i.e.

ME/CFSs vs HCs (p = 0.89), ME/CFSm vs HCs (p = 0.77), ME/CFSs vs ME/CFSm (p

= 0.95), all patients ME/CFS vs HCs (p = 0.78).

On the other hand, the HERV-K expression differed significantly between ME/CFSm

group and the HCs (p = 0.050). HERV-K activity was not distinct between the ME/CFS

groups: ME/CFSs vs ME/CFSm (p = 0.12), ME/CFS vs HCs (p = 0.17). ME/CFSs vs

HCs (p = 0.97) (Figure 1 B).

[FIGURE 1]

Discussion

The most recognized and widely-used case definitions (Fukuda [13] and Canadian

Consensus criteria [1]) are based on self-reported symptoms. Studies of energy

metabolism, oxidative stress and immunological alterations in ME/CFS have

demonstrated imbalance in all these pathways, but the use of such information for

diagnostic purposes is still far from reality.

Here, HERV-K and W transcripts were detected in all groups investigated, and we

found that HERV-K was overexpressed in moderate ME/CFS. It is possible that the

immunological, genic expression and metabolic alterations are different according to

disease severity.

The interplay between endogenous retroviruses and the immune system is complex.

ERVs are part of the host genome and in theory, they are supposed to be recognized as

self-antigens and an immune tolerance should be established during the early stages of

the organism development [14]. However, HERV products can interact with

components of the innate immune system leading to the activation of pro-inflammatory

pathways or, in some particular cases, their suppression [15]. The syncytin -2 protein

for example, is a product of the ERV-FRD Env gene that has an immunosuppressive

role by preventing maternal immune response against the fetus [16]. In a distinct

scenario, it was demonstrated using psoriasis model that a pro-inflammatory

environment could be able to suppress the expression of repetitive elements, including

HERVs [17]. It would be reasonable to suggest that the immunological enhancement

seen in more severe ME/CFS works by silencing the HERV transactivation that occurs

in moderate cases. Such transactivation could be caused by exogenous viral replication

or another as yet unknown factor. In line with this, Montoya and colleagues (2017)

found a cytokine signature of severity in people with ME/CFS [18]. They demonstrated

that from the 17 cytokines related to severity, 13 are pro-inflammatory, and (in addition

to the worsening of the symptoms) may cause the reversion of the HERV-K activity to

levels similar to those seen in healthy individuals. It may similarly occur with HERV-

W, which, despite not being at significant levels, there was a slight decrease in people

severely affected by ME/CFS when compared to those who are moderately affected.

Infection has often been considered as a trigger to ME/CFS. Many patients report that

the fatigue began during or short after an episode of infectious disease. A number of

pathogens including viruses have been associated with this disease [3]. And, due to its

life long persistence and broad cell tropism, the herpesviridae family, particularly HHV-

6 has been considered to be a possible trigger for ME/CFS for many, even though such

relationship has not been consistent [3,9]. Interestingly, HHV-6 as some other

herpesviruses, is also capable of transactivating HERVs, particularly, HERV-K [6].

Such transactivation may be either direct (through LTR activation by viral products) or

indirect (via transcriptional binding factors and cytokines produced by viral replication)

[3,6]. It is possible that as the disease progresses, whatever the exogenous infection that

would have act as the trigger factor is controlled, and consequently, the HERVs

transactivation decrease. Unfortunately, we did not perform serological or molecular

tests for exogenous viruses.

Two reports of HERV activity in ME/CFS were published some years ago but the

results were conflicting. In 2013 Oakes and his team found no difference on the

expression of HERV-K18 envelope in people with ME/CFS when compared with HCs

[9]. In the same year De Meirleir and colleagues, using immunohistochemical methods,

found immunoreactivity to HERV proteins (HERV-K, HERV-18, HERV-R and HERV-

FRD) in dendritic cells of the duodenum of individuals diagnosed with the syndrome

[10], suggesting that alterations in endogenous retroviruses expression pattern may

occur in ME/CFS. The differences between the results of Oakes and colleagues and ours

may be due to the methods used to detect HERV-K. While the present work used

generic primers for HERV-K that allow the detection of hundreds of elements from

most HML subfamilies the Oakes team searched for the HERV-K 18 envelope

transcripts only, using a method specific to this particular element, while neglecting all

the remaining proviruses from the K family. On the other hand, we are unable to

determine which K family proviruses are involved in the differential expression

observed.

The molecular method used here to detect HERV-W was also generic and was widely

used in several studies that found differential expression of this element in pathological

conditions, including in the blood, brain and CSF of multiple sclerosis (MS) patients

[19]. Therefore, despite the similarity of a number of symptoms and the strong

immunological component of ME/CFS and MS, the mechanisms responsible for HERV

reactivation in such diseases are likely distinct.

In conclusion, this is the first report that demonstrates increased expression of an endogenous retrovirus in the blood of individuals with moderate ME/CFS. While the increased expression of these retroelements can't be directly associated to the ME/CFS pathogeny, the observation of this phenomenon cannot be ignored.

Funding: This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) [grant number 15/05958-3]. LN, EL and CK have been funded by the National Institutes of Health (NIH USA) under award number [2R01AI103629], however the content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

References

1. Carruthers BM, Jain AK, De Meirleir KL, Peterson DL, Klimas NG, Lerner AM, et al. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. J Chronic Fatigue Syndr. 2003;11(1):7–115.

2. Lorusso L, Mikhaylova S V., Capelli E, Ferrari D, Ngonga GK, Ricevuti G. Immunological aspects of chronic fatigue syndrome. Autoimmun Rev. 2009;8(4):287–291.

3. Rasa S, Nora-Krukle Z, Henning N, Eliassen E, Shikova E, Harrer T, et al. Chronic viral infections in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). J Transl Med. 2018;16(1):268.

4. Watt T, Oberfoell S, Balise R, Lunn MR, Kar AK, Merrihew L, et al. Response to valganciclovir in chronic fatigue syndrome patients with human herpesvirus 6 and Epstein-Barr virus IgG antibody titers. J Med Virol. 2012;84(12):1967–1974.

5. Weiss RA. The discovery of endogenous retroviruses. Retrovirology. 2006;3(1):67.

6. Chen J, Foroozesh M, Qin Z. Transactivation of human endogenous retroviruses by tumor viruses and their functions in virus-associated malignancies. Oncogenesis. 2019;8(1):6.

7. Nellåker C, Yao Y, Jones-Brando L, Mallet F, Yolken RH, Karlsson H. Transactivation of elements in the human endogenous retrovirus W family by viral infection. Retrovirology. 2006;3(1):44.

8. Gröger V, Cynis H. Human Endogenous Retroviruses and Their Putative Role in the Development of Autoimmune Disorders Such as Multiple Sclerosis. Front Microbiol. 2018 ;9:265.

9. Oakes B, Hoagland-Henefield M, Komaroff AL, Erickson JL, Huber BT. Human Endogenous Retrovirus-K18 Superantigen Expression and Human Herpesvirus-6 and Human Herpesvirus-7 Viral Loads in Chronic Fatigue Patients. Clin Infect Dis. 2013;56(10):1394–1400.

10. De Meirleir KL, Khaiboullina SF, Frémont M, Hulstaert J, Rizvanov AA, Palotás A, et al. Plasmacytoid dendritic cells in the duodenum of individuals diagnosed with myalgic encephalomyelitis are uniquely immunoreactive to antibodies to

human endogenous retroviral proteins. In Vivo. 2013;27(2):177–187.

11.   Lacerda EM, Mudie K, Kingdon CC, Butterworth JD, O'Boyle S, Nacul L. The
UK ME/CFS Biobank: A Disease-Specific Biobank for Advancing Clinical
Research Into Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. Front
Neurol. 2018 Dec;9:1026.

12.   Nali LHS, Oliveira ACS, Alves DO, Caleiro GS, Nunes CF, Gerhardt D, et al.
Expression of human endogenous retrovirus K and W in babies. Arch Virol.
2017;162(3):857–861.

13.   Fukuda K, Straus SE, Hickie I, Sharpe MC, Dobbins JG, Komaroff A. The
chronic fatigue syndrome: a comprehensive approach to its definition and study.
International Chronic Fatigue Syndrome Study Group. Ann Intern Med.
1994;121(12):953–959.

14.   Tugnet N, Rylance P, Roden D, Trela M, Nelson P. Human Endogenous
Retroviruses (HERVs) and Autoimmune Rheumatic Disease: Is There a Link?
Open Rheumatol J. 2013;7:13–21.

15.   Magiorkinis G, Hurst TP. Activation of the innate immune response by
endogenous retroviruses. J Gen Virol. 2015;96(6):1207–1218.

16.   Mangeney M, Renard M, Schlecht-Louf G, Bouallaga I, Heidmann O, Letzelter
C, et al. Placental syncytins: Genetic disjunction between the fusogenic and
immunosuppressive activity of retroviral envelope proteins. Proc Natl Acad Sci.
2007;104(51):20534–20539.

17.   Lättekivi F, Kõks S, Keermann M, Reimann E, Prans E, Abram K, et al.
Transcriptional landscape of human endogenous retroviruses (HERVs) and other
repetitive elements in psoriatic skin. Sci Rep. 2018;8(1):4358.

18.   Montoya JG, Holmes TH, Anderson JN, Maecker HT, Rosenberg-Hasson Y,
Valencia IJ, et al. Cytokine signature associated with disease severity in chronic
fatigue syndrome patients. Proc Natl Acad Sci. 2017;114(34):E7150–8.

19.   Morandi E, Tanasescu R, Tarlinton RE, Constantinescu CS, Zhang W, Tench C,
et al. The association between human endogenous retroviruses and multiple
sclerosis: A systematic review and meta-analysis. PLoS One.
2017;12(2):e0172415.

327   Figure 1. Boxplot of expression levels (in fold change) of HERVs among the groups. A)

328   HERV-K and B) HERV-W. Significance between the groups (obtained by Wilcox test)

329   is evidenced by an asterisk.