Correcting for sparsity and non-independence in glycomic data through a systems biology framework

Bokan Bao[1,2,3,+], Benjamin P. Kellman[1,2,3,+], Austin W.T. Chiang[1,2], Austin K. York[1,2], Mahmoud A. Mohammad[4], Morey W. Haymond[4], Lars Bode[1], Nathan E. Lewis[1,2,5,*]

[1] Department of Pediatrics, University of California, San Diego, La Jolla, CA 92093, USA

[2] The Novo Nordisk Foundation Center for Biosustainability at the University of California, San Diego, La Jolla, CA 92093, USA

[3] Bioinformatics and Systems Biology Graduate Program, University of California, San Diego, La Jolla, CA 92093, USA

[4] Department of Pediatrics, Children's Nutrition Research Center, US Department of Agriculture/Agricultural Research Service, Baylor College of Medicine, Houston, Texas 77030, USA

[5] Department of Bioengineering, University of California, San Diego, La Jolla, CA 92093, USA

[+] These authors contributed equally to this work

[*] Corresponding author:

Name: Nathan E. Lewis

Address: 9500 Gilman Drive MC 0760, La Jolla, CA 92093

Phone: 858-246-1876

E-mail: nlewisres@ucsd.edu

Short running title: Comparative glycoprofiles analysis with glycan substructures

Abstract

Glycans are fundamental cellular building blocks, involved in many organismal functions. Advances in glycomics are elucidating the roles of glycans, but it remains challenging to properly analyze large glycomics datasets, since the data are sparse (each sample often has a minority of measured glycans) and detected glycans are non-independent, sharing many intermediate biosynthetic steps with other glycans. We address these challenges with GlyCompare, a glycomic data analysis approach that leverages shared biosynthetic pathway intermediates to correct for sparsity and non-independence in glycomics. Specifically, quantities of measured glycan are propagated to intermediate glycan substructures, which enable direct comparison of different glycoprofiles. Using this, we studied diverse N-glycan profiles from glycoengineered erythropoietin. We obtained biologically meaningful clustering of mutant cells and identified knockout-specific effects of fucosyltransferase mutants on tetra-antennary structures. We further analyzed human milk oligosaccharide profiles and identified novel impacts that the mother's secretor-status has on fucosylation and sialylation. Our GlyCompare substructure-oriented approach will enable researchers to take full advantage of the growing power and size of glycomics data.

Introduction

Glycosylation is a highly abundant and complex post-translational modification, decorating between one-fifth and one-half of eukaryotic proteins[1,2]. These diverse carbohydrates account for 12-25% of dry mass, and have important functional and pathological roles[3,4]. Despite their essentiality, glycans are difficult to study; thus, our understanding of their impact is limited. The complex structures of glycans arise from a non-template driven synthesis through a biosynthetic network. A simple change of a single intermediate glycan or glycosyltransferase will have cascading impacts on the final glycans obtained[5,6]. Unfortunately, glycoprofiling and glycomic data lack the necessary systems perspective to easily decode this potential for change. It is important to understand the network behind the glycoprofiles so that we can better understand the regulation of this process.

New tools that aid in the acquisition and aggregation of glycoprofiles are emerging, making large-scale comparisons of glycoprofiles possible. Advances in mass spectrometry now enable the rapid generation of large volumes of glycoprofiles with detailed glycan composition[7–10], exposing the complex and heterogeneous glycosylation patterns on lipids and proteins[11,12]. Large glycoprofile datasets and supporting databases are also emerging, including GlyTouCan[13], UnicarbDB[14], Glygen and UniCarbKB[15].

These new technologies and databases provide opportunities to examine global trends in glycan function and their association with disease. However, the rapid and accurate comparison of glycoprofiles can be challenging with the size, sparsity and heterogeneity of such datasets. Indeed in any one glycoprofiles, only a few glycans may be detected among the thousands of possible glycans. Thus, even in the same dataset, few glycans, if any, overlap between samples if there is a major perturbation to glycosylation. However, these non-overlapping glycans may only

differ in their synthesis by as few as one enzymatic step. Thus, it can be difficult to know which

glycans to compare. Furthermore, since glycans often share substantion portions of their

biosynthetic pathways with each other, statistical methods that assume independence (e.g., t-

tests, ANOVA, etc) are inappropriate for glycomics. Here we address these challenges by

proposing that glycan substructures, or intermediates, are useful functional units to enable

meaningful glycoprofile comparisons, since each substructure can capture one step in the

complex process of glycan synthesis. Thus, using substructures for comparison, we account for

the shared dependencies of all glycans.

Previous work has investigated the similarity across glycans using glycan motifs, such as

glycan fingerprinting to describe glycan diversity in glycan databases[16], alignment of glycan

structures[17], identification of glycan epitopes in glycoprofiles[18], graph-based deconvolution of

LC-MS to clarify glycan abundance[19], or comparison of glycans in glycoprofiles leveraging

simple structures[20]. These tools use information on glycan composition or epitopes. However,

accounting for the shared biosynthetic steps in the various glycans in glycoprofiles, provides

complete biosynthetic context to all glycan epitopes. That context includes connecting all

glycans to the enzymes involved in their synthesis, the order of the enzyme reactions, and

information on competition for glycan substrates. Thus, a generalized substructure approach can

facilitate the study of large numbers of glycoprofiles and connect them to the shared mechanisms

involved in making each glycan.

Here we present GlyCompare, a method enabling the rapid and scalable analysis and

comparison of any number of glycoprofiles while accounting for the biosynthetic similarities of

each glycan. This will facilitate the discovery of mechanisms underlying the changes among

glycoprofiles. We demonstrate the functionalities and performance of this approach with both

protein-conjugated and unconjugated glycomics, using recombinant erythropoietin (EPO) N-

glycosylation and human milk oligosaccharides (HMOs). Specifically, we analyzed sixteen

MALDI-TOF glycoprofiles of EPO, where each glycoprofile is of EPO produced in a different

glycoengineered CHO cell line[9,11]. We also analyzed forty-eight HMO glycoprofiles from six

mothers[21]. By analyzing these glycoprofiles with GlyCompare, we quantify the abundance of

important substructures, cluster the glycoprofiles of mutant cell lines, connect genotypes to

unexpected changes in glycoprofiles, and associate a phenotype of interest with substructure

abundance and synthesis network dynamics. The analysis of the EPO and HMO datasets

demonstrate that our novel framework presents a convenient and automated approach to

elucidate novel insights into complex patterns in glycobiology.

Results

GlyCompare decomposes glycoprofiles to facilitate glycoprofile comparison

Glycoprofiles can be decomposed into glycan substructure abundances. The resulting

substructure profile has richer information than whole glycans and enables more precise

comparison across conditions. However, the comparison does not need all substructures. Glycan

biosynthesis involves long pathways with multiple intermediates and much pathway redundancy.

Thus, pathways can be collapsed to obtain a subset of substructures while preserving the

information of all glycans in the dataset. We call this minimal set of substructures "glyco-

motifs". The GlyCompare workflow consists of several steps wherein glycoprofiles are

annotated and decomposed, glyco-motifs are prioritized, and each glyco-motif is quantified for

subsequent condition-specific comparisons (**Fig. 1**). The specific workflow is described as

follows.

First, to characterize each glycoprofile by substructures, all relevant substructures are defined

and quantified. Thus, a complete set of glycan substructures is obtained for all glycans in all

glycoprofiles being analyzed. For each glycoprofile, the abundance of each substructure is

calculated by summing the abundance of all glycans associated with the substructure. This

results in a substructure profile, which stores abundances for all glycan substructures (**Fig. 1a,b**)

in given glycoprofile.

Second, to identify the most informative substructures (glyco-motifs), substructures are

prioritized using the substructure network. The substructure network is built by connecting all

substructures into their biosynthetic pathways (**Fig. 1c**). To improve computational efficiency,

only substructures associated with glycans being analyzed are included in the substructure

network. Starting from the monosaccharides, each level of the network represents another

biosynthetic step, with one more monosaccharide than the previous level. The edges in the

network are the enzymatic reactions adding each monosaccharide. These edges are weighted by

the correlation between the abundances of the substrate and product substructures across all

samples. Redundant substructures can be easily identified since their parent-child substructure

abundances will be perfectly correlated. Substructure network reduction proceeds by collapsing

links with a perfect correlation between substrate and product substructures, and only retaining

the product substructure. The remaining substructures are termed glyco-motifs; they completely

describe the variance at the substructure level. The abundances of all glyco-motifs are then

represented with a glyco-motif profile, the minimal subset of meaningful substructure

abundances (**Fig. 1d**).

For larger datasets, summarizing the glyco-motifs becomes necessary. Since the glyco-motifs

are also clustered when clustering the glycoprofiles (**Fig. 1d**), representative substructures are

generated by summarizing the common structure in each cluster (**Fig. 1e**). The representative

substructure describes the glycan features that vary the most across samples. To extract the

common structure features, the average weight of each monosaccharide is calculated and the

monosaccharides with a weight larger than 51% are preserved, which illustrates the predominant

structure in the cluster. This allows one to quickly evaluate the distinguishing glycan features

that vary across samples in any given dataset.

The workflow we described here successfully connects all glycoprofiles in a data set by their

shared intermediate substructures, thus allowing robust analysis of the differences across

glycomics samples and the evaluation of the associated genetic bases.

GlyCompare accurately clusters glycoengineered EPO samples

In examining the EPO glycosylation data[9], we found the glycoprofiles failed to form robust

clusters that can be easily interpreted. Furthermore, glycoprofiles with low genetic similarity

clustered together (**Fig. 2a and Supplementary Fig. 1, 2**). This inconsistency and poor

clustering stems from the inherent sparseness of the glycoprofile datatype, i.e., each glycoprofile

only has a few glycans, making the matrix of all samples very sparse, unfit for standard

clustering approaches, and thus hard to interpret. Particularly problematic is that pairs of glycans

differing in a single monosaccharide are treated as two completely different glycans under

standard clustering approaches. Thus, we found that clustering is affected more by the presence

or absence of a glycan, rather than structural similarity.

GlyCompare addresses these problems by elucidating hidden similarities between glycans

after decomposing glycoprofiles to their composite substructures. The 52 glycans were

decomposed into their constituent glycan substructures, resulting in a substructure profile that

consists of 613 glycan substructures and a further simplified 120 glyco-motif profile

(**Supplementary Fig. 3**). The clustering automatically distinguished the samples based on the

structural patterns, and clustered the glyco-motif profiles into groups more consistently

associated with the extent of changes in the profile than the raw glycan-based clusters (**Fig. 2b**

and Supplementary Fig. 4, 5).

The sixteen glycoprofiles clustered into three groups with a few outliers (**Fig. 2b**), and the 120

glyco-motifs clustered into twenty-four groups, each summarized by representative substructures

(**Fig. 2c and Supplementary Fig. 5**). The clusters of glycoprofiles are consistent with the

genetic similarities among the host cells. Specifically, the major substructure patterns cluster

individual samples into groups of 'wild-type (WT)-like', 'mild', 'medium' and 'outlier'

categories. The WT-like category contains one group, WT and B4galt1/2/3/4/ knockouts, which

contains most of the substructures seen in WT cells. The mild group includes the Mgat4b/4a,

Mgat4b, and Mgat5 knockouts, which each lose the tetra-antennary structure, and an individual

St3gal4/6 knockout that loses the terminal sialylation. The medium category is a group that

contains knockouts of St3gal4/6 and Mgat4a/4b/5, knockouts of Mgat4a/4b/5 and B3gnt2,

knockouts of Mgat4a/4a/5 with a knock-in of human ST6GAL1, and knockouts of Mgat4a/4b/5

and St3gal4/6, which lose the tri-antennary structure. The 'outlier' category includes three

individual glycoprofiles with knockouts for Fut8, Mgat2, and Mgat1, each of which generate

many glycans not detected in the WT nor other three groups ('WT-like', 'mild' and 'medium').

While some glyco-motif clusters can be seen in the glycoprofile clusters, there are important

differences, and the glyco-motif clusters provide more information and improved cluster stability

(**Supplementary Fig. 2**). The results demonstrated the excellent performance of our

GlyCompare in assessing the structural similarity between different glycoprofiles.

GlyCompare identifies substructure changes across glycoprofiles

GlyCompare helps to more robustly group samples by accounting for the biosynthetic and

structural similarities of glycans. Further analysis of the representative structures provides

detailed insights into which structural features vary the most across samples. To accomplish this,

we standardized--using z-score normalization--the representative structure abundances. We then

identified significant changes in representative substructure abundances between mutant cells

and WT (**Fig. 2d**). This highlights the specific structural features of glycans that are impacted

when glycoengineering recombinant EPO.

   As expected, in the Mgat1 knockout glycoprofile, only high mannose N-glycans are seen.

Also, in the Mgat2 knockout, the glycan substructure of bi-antennary on one mannose linkage

significantly increases, and the unique structure of bi-antennary LacNac elongated in the N-

glycans emerges in the St3gal4/6 and Mgat4a/4b/5 knockouts. Along with expected changes in

α-1,6 fucosylation in the Fut8 knockout glycoprofile, we also observed an increase in the tetra-

antennary poly-LacNac elongated N-glycan without fucose, which has not been previously

reported (**Fig. 3a**). In the St3gal4/6 knockout (**Fig. 3a**), we observed the relative abundance of

structures with sialylation has significantly decreased, where the tetra-antennary and triantennary

poly-LacNAc elongated N-glycan substructure without sialylation significantly increased

(Rep13: $p<0.01$, Rep20: $p<0.001$). Finally, in the Mgat4b, Mgat4a/4b and Mgat5 knockout

glycoprofiles (**Fig. 3b**), tetra-antennary substructures (Rep16-Rep22) decreased significantly

($p<0.01$), while triantennary substructures with GlcNac elongation (Rep13-Rep14) increased

significantly ($p<0.01$). However, the poly-LacNac elongation structure decreased significantly

for the Mgat5 and Mgat4b knockout ($p<0.001$). Interestingly, while both the Mgat4b and Mgat5

knockouts significantly decreased in the triantennary poly-LacNac elongated N-glycan, the

Mgat4b knockout significantly increased in the tri-antennary LacNac elongated N-glycan. Thus,

through the use of GlyCompare, we identified the specific glycan features that are impacted not

only in individual glycoengineered cell lines, but also features shared by groups of related cell

lines.

GlyCompare reveals phenotype-associated substructures and trends invisible at the whole

glycan level

Many secreted and measured glycans are also precursors, or substructures, of larger glycans (**Fig.**

4a). Thus, the secreted and observed abundance of one glycan does not equal to the total amount

synthesized. GlyCompare can quantify the total abundance of a glycan by combining the glycan

abundance with the abundance of its products. To demonstrate this capability of GlyCompare,

we analyzed HMO abundance, and examined the impact of secretor status and days postpartum

on HMO abundance. We obtained forty-seven HMO glycoprofiles from 6 mothers (1, 2, 3, 4, 7,

14, 28 and 42 days postpartum (DPP)), 4 "secretor" mothers with functioning FUT2 (α-1,2

fucosyltransferase), and 2 "non-secretor" mothers with non-functional FUT2. With the non-

independence of HMOs addressed with GlyCompare, we were able to use powerful statistical

methods to study trends in HMO synthesis. Specifically, we used Generalized Estimating

Equations[22] to build regression models predicting secretor status and DPP from substructure

abundance.

We first checked both the glycan-level and substructure-level clustering of the glycoprofile.

Samples with same secretor status and days postpartum (DPP) were successfully grouped

(**Supplementary Fig. 6**). Further examination of the total amount of substructure synthesized

revealed phenotype-related trends invisible at the level of the raw glycoprofile. For example, the

LSTb substructure (X80) increased in secretor mothers (Wald p = 0.001) and decreased in non-

secretor mothers over time (Wald $p < 2.2x10^{-16}$; **Fig. 4b**). Yet, the same trend was weak or

inconsistent for all glycans containing the X80 substructure: LSTb, DSLNT and DSLNH. Like

X80, LSTb and DSLNT decreased in abundance over time in non-secretors (LSTb Wald

p=0.0046, DSLNT Wald p = 0.0045); however, unlike X80, neither LSTb nor DSLNT

significantly increased in secretor mothers over time (Wald p>0.15). Meanwhile, DSLNH, like

X80, increased over time in secretor mothers (Wald p=0.0062) (**Fig. 4e**). Rather than decreasing

like X80, non-secretor DSLNT abundance significantly increased over time (Wald $p=5.8x10^{-9}$)

(**Fig. 4d**). The secretor-specific trends in total LSTb are only visible examining the X80

substructure abundance (**Fig. 4c**). Thus, while secretor status is expected to impact HMO

fucosylation, GlyCompare reveals associations with non-fucosylated substructures. Viewing

substructure abundance as total substructure synthesized provides a new fundamental measure to

the study of glycoprofiles, it also creates an opportunity to explore trends in synthesis.

GlyCompare identifies condition-specific synthesis dynamics

We applied GlyCompare to explore changes in HMO synthesis over time. We estimate the fluxes

for each biosynthetic reaction by quantifying the ratio of product-reactant pairs (pairs of glycan

substructures that differ by one monosaccharide). By comparing the substructure abundance of

reactants and products, we can estimate how many reactant substructures are converted to

product substructure. With clearly defined glycan biosynthesis pathways, we explore synthesis

dynamics through the systematic estimation of reaction flux under various conditions.

We measured the product/reactant ratio for every reaction in the HMO substructure network

(**Supplementary Fig. 7**) to estimate flux for each reaction over time and across secretor status.

We found several reactions strongly associated with secretor status (**Supplementary Table 1**

and Supplementary Fig. 8), i.e. the ability of the mother to add an α-1,2 fucose to HMOs. As

expected, the reaction flux from the LNT substructure (X38) to the LNFPI substructure (X86),

was strongly associated with secretor status (Wald $p<2.2\times10^{-16}$) (**Fig. 4a**). In secretors, 36.2% of

X38 was converted to X86, compared to non-secretors, wherein only 5% of X38 was converted.

Eventhough secretor status is defined by the fucosyltransferase-2 genotype, not all secretor-

associated reactions were fucosylation reactions. We further explored the secretor-X80

association using the product/reactant ratio. Specifically, we examined the upstream reaction of

LNT (X38) to LSTb (X80) and the downstream reaction--LSTb (X80) to DSLNT (X103) (**Fig.**

4b, 5a). Using the product/reaction ratio of substructure abundances, we found secretor status

correlates with the reaction of LSTb to DSLNT but does not impact the conversion of LNT to

LSTb; any changes in sialylation is intriguing considering secretor status only directly impacts

fucosylation. We measured the upstream reaction of LNT converting to LSTb, using the

X80/X38 ratio over time; however, no significant change was observed with respect to secretor

status (Wald $p=0.44$). In the reaction of LSTb to DSLNT, we found a secretor-specific reaction

propensity. Specifically, we quantified the association of the X103/X80 ratio with secretor status.

The X103/X80 ratio was significantly higher (Wald $p=0.018$) in secretor mothers (**Fig. 5b)**. The

higher ratio suggests that the turnover from LSTb (X80) to DSLNT (X103) is significantly

higher in secretor mothers. In the average non-secretor mother, 52.3% of LSTb is converted to

DSLNT. Meanwhile in secretors, the average conversion rate is 81.8% (a 63.9% reduction in the

efficiency of the LSTb to DSLNT conversion). The LSTb to DSLNT conversion rate appears

higher in secretors while conversion from the LSTb precursor, LNT, appears unchanged.

Examining the product/reactant ratio has revealed a phenotype-specific reaction propensity thus

providing insight to the condition-specific synthesis.

Discussion

Glycosylation has generally been studied from the whole-glycan perspective using mass

spectrometry and other analytical methods. From this perspective, two glycans that differ by only

one monosaccharide are completely distinct and are not comparable. Thus, the comparative study

of glycoprofiles has been limited to changes between glycans shared by multiple glycoprofiles or

small manually curated glycan substructures[16]. Glycompare sheds light on the hidden

biosynthetic interdependencies between glycans by integrating the biosynthetic pathways into the

comparison. Glycoprofiles are converted to glyco-motif profiles, wherein each substructure

abundance represents the cumulative abundance of all glycans containing that substructure. This

enumeration and quantification of substructures can be easily scaled up to include many

glycoprofiles in large datasets. Additionally, since no prior information is required beyond

glycan identities and quantities, the method can even facilitate analysis of glycans with limited

characterization. Thus, it brings several advantages and new perspectives to enable the

systematic study of glycobiology.

First, the GlyCompare platform computes a glyco-motif profile (i.e., the abundances of the

minimal set of glycan substructures) that maintains the information of the original glycoprofiles,

while exposing the shared intermediates of measured glycans. These sample-specific glyco-motif

profiles more accurately quantify similarities across glycoprofiles. This is made possible since

glycans that share substructures also share many biosynthetic steps. If the glycan biosynthetic

network is perturbed, all glycans synthesized will be impacted and the nearest substructures will

directly highlight where the change occurred. For example, in EPO glycoprofiles studied here,

the tetra-antennary structure is depleted on the Mgat4A/4B/5 knockout group and the

downstream sialylated substructure depleted when St3gal4/6 were knocked out. Such structural

patterns emerge in GlyCompare since the tool exposes the substructures through clustering, thus

identifying common features in glycans measured across diverse samples.

Second, new trends in synthesis network dynamics become visible at the substructure level.

For example, in the HMO data set, multiple HMOs are made through a series of steps from LNT

to DSLNH (**Fig. 4a**). Only when the substructure abundances are computed are we able to

observe the secretor-dependent differences in the LSTb substructure, X80. This is particularly

interesting since secretor status is defined by changes in α-1,2 fucosylation, but we see here

additional secretor-dependent changes to sialylated structures with no fucose. These are the

systemic effects invisible without a systems-level perspective due to the interconnected nature of

glycan synthesis; this disparity underlines the power of this method.

Third, the sparse nature of glycobiology datasets and the inherent connections between

glycans make glycomic data unfit for many statistical analyses commonly used. However, the

translation of glycoprofiles into substructure abundance provides a framework for more powerful

and statistically rigorous analysis of glycomic datasets. Single sample perturbations, such as the

knockouts in the glycoengineered EPO, can be compared to wild-type; all substructure data can

be normalized and then rigorously distinguished from the control using a one sample t-test.

Conditions or phenotypes with many samples, such as the secretor status in the HMO dataset,

can be compared using regression models, like Generalized Estimating Equations (GEE)[22], to

evaluate the association between the phenotypes and glycosylation. For example, in HMO data,

the GEE revealed that the α-1,2 fucose substructure is enriched in secretor status, consistent with

the previous studies[23–25]. Additionally, the GEE discovered that the sialylation increases over

time in secretor mothers while decreasing in non-secretor. Finally, in combination with the

substructure network, we can systematically study glycan synthesis. The product/reactant ratio

provides a regular measure indicating the flux through the glycan biosynthetic pathways. Using

the HMO dataset, we demonstrate the power of this perspective by showing that more LSTb is

converted to DSLNT in the secretor mother using regression modeling. The perspectives made

available through GlyCompare are not limited to t-tests and regression models. We have reduced

the sparse nature of glycoprofiles, thereby making countless comparisons and new analyses

possible.

Conclusions

328   In conclusion, GlyCompare provides a novel paradigm for describing complex glycoprofiles,

329   thus enabling a wide range of analyses and facilitating the acquisition of detailed insights into the

330   molecular mechanisms controlling all types of glycosylation.

Methods

Data Availability and Implementation

333   **https://github.com/LewisLabUCSD/GlyCompare**

Glycoprofile preprocess procedures

There are three procedures required for preprocessing the studied glycoprofiles (**Fig. 1a**). First, glycoprofiles are parsed into glycans with abundance. In each glycoprofile, the glycans are manually drawn and exported with GlycoCT format using the Glytoucan Graphic Input tool[13]. The structure of glycan in GlycoCT format will be further loaded into python using "glypy" (a glycan analysis and glycoinformatics library for Python) for programmatic manipulation. Assuming we have a glycoprofile i, the corresponding abundance of each glycan j in glycoprofile i is represented by g_{ij}. For example, the relative m/z peak in the mass spectrum or the abundance value in an HPLC trace, will be rectified by the total abundances of glycans in this glycoprofile $g_{ij}/\Sigma g_{i*}$. Sometime, the glycans with ambiguous topology will be handled by assuming it could belong to every possible structure with the same probability, thereby creating all possible n structures but with $g_{ij}/n\Sigma g_{i*}$ abundance of each. Second, glycans are annotated with glycan substructure information and this information is transformed into the glycoprofile vector. The substructures within a glycan are exhaustively extracted by breaking down each linkage or a combination of linkages of the studied glycan. Note that this method cannot currently deal with glycans with ring topology. All substructures extracted will be merged into a substructure set S, in which the complexity degrees are sorted and the duplicated substructures are removed. Then, each glycan will match to the substructure set S resulting in a binary glycan substructure presence (1) or absence (0) vector, named x_{ij}. Lastly, a glycoprofile vector is calculated as $p_i = \Sigma x_{ij} g_{ij}/\Sigma g_{i*}$ representing the abundance of the substructures s in this glycoprofile, where $p_i = (s_{1i}, \ldots, s_{ni})$. Third, a substructure network is built based on the glycoprofile vectors. The substructure network is a directed acyclic graph and each node denotes a glycan substructure. Given the substructure set S, the root node starts from the monosaccharides or a defined root

core structure, and a child node is a substructure that has only one monosaccharide added to its

parent node. We note that one child node might have multiple parent nodes and vice versa. The

child node depends on its parent node(s) since it cannot exist alone without any parent node.

Generating the glyco-motif vector bases on the substructure abundance

A larger subset of the substructure network is necessary to uniquely describe a more diverse set

of glycoprofiles while fewer substructures are needed to sufficiently describe more similar

glycoprofiles. Comparisons become more focused when only examining these variable

substructures. By checking the substructure network, the substructures that have exactly the same

abundance can be merged without any information loss. In other words, after the substructure

network is generated, it is simplified by merging the substructure nodes. The merging criteria are

based on how child substructure node s_b depends on the parent substructure node s_a. The

dependency is the correlation of their abundance across all glycoprofiles, $corr(s_{a*}, s_{b*})$. If the

correlation is 1, we can conclude that the addition of the specific monosaccharide is not

perturbed across all glycoprofiles, which means they carry exactly the same information. Thus,

the parent node can be pruned without information loss. All remaining nodes, namely, the glyco-

motifs, are used to cluster the glycoprofiles.

Meanwhile, the weight of each node is initialized at a value of 1. When a node is removed, the

weight is equally divided and distributed to child nodes whose correlation with the removed node

is 1. Since this method happens from the root to leaves, the node that serves as checkpoint will

gain more weight. The weights W are used later for generating the representative substructures.

Procedures for glycoprofile clustering and identifying representative glycan substructures

The preprocessed glycoprofiles (see details in the "glycoprofile preprocess procedures") generate

the glycoprofile vectors to enable further clustering analysis. Here we used the Pearson

correlation and 'complete' distance to cluster the glycoprofiles. This procedure clusters the

glycoprofiles and substructures.

To identify the representative glycan substructures, a set of glycan substructures with weights

W are first aligned. Then, we calculate the sum of monosaccharide weights for each glycan

substructure. The representative substructure is thus defined as the glycan substructures with

their summed monosaccharide weights greater than 51% of the total weight of glycan

substructures. Lastly, the averaged abundances of the representative substructures are generated

to assess their differential expressions between different glycoprofiles.

Specification for HMO data

48 milk samples were collected from 6 human mothers (1, 2, 3, 4, 7, 14, 28 and 42 days

postpartum (DPP)). HPLC was used to measure the abundances of the 16 major glycans. The

relative abundance of each glycan in each milk sample is normalized by $w_i/\Sigma w_*$ for the given

sample. Then, we followed the substructure network was built as the procedure mentioned

above.

Testing the substructure-phenotype association

The General Equation Estimating (GEE from R::geepack[26,27]) with an exchangeable correlation

structure was used to measure associations between motif abundance, secretor status and DPP;

these regression models were trained to predict motif abundance from either secretor status or

DPP. A Wald Test was used to measure the significance of phenotype contribution.

Product/Reactant Ratio as a proxy for flux and estimating flux-phenotype associations

To further isolate motif-specific effects from biosynthetic biases, we explored methods to control

for the reactant-product relations. First, we isolated the relative abundance of motifs separated by

only one monosaccharide in the biosynthetic network; these are reactant-product relations like

LNT and LSTb. Motif abundance represents the total substructure synthesized therefore when

we examine the ratio of product/reactant we measure the total amount of the reactant

substructure converted to the product substructure in the sample. The Product/Reactant Ratio is

therefore a proxy for flux. Using the same GEE repression modeling approach from testing

substructure-phenotype associations, we can measure the association between flux between two

glycans and a phenotype of interest.

Acknowledgements

We thanks Joshua Klein for providing generous and timely guidance on using the glypy (https://github.com/mobiusklein/glypy). This work is a publication of the U.S. Department of Agriculture/Agricultural Research Service, Children's Nutrition Research Center, Department of Pediatrics, Baylor College of Medicine, Houston, Texas. The contents of this publication do not necessarily reflect the views or policies of the U.S. Department of Agriculture, nor does mention of trade names, commercial products, or organizations imply endorsement from the U.S. government. This work was conducted with support from the Novo Nordisk Foundation provided to the Center for Biosustainability at the Technical University of Denmark (NNF10CC1016517: A.W.T.C.), NIGMS (R35 GM119850: B.B., B.K.P., N.E.L.) , NICHD (R21 HD080682: L.B.) and USDA (USDA/ARS 6250-6001; M.W.H).

Author contributions

B.B, B.P.K designed the work. B.B, B.P.K, A.W.T.C., A.K.Y., and N.E.L. performed data analysis. The manuscript was written by B.B, B.P.K, A.W.T.C., L.B., and N.E.L.

Competing interests

The authors declare no competing financial interests.

References

1. Khoury, G. A., Baliban, R. C. & Floudas, C. A. Proteome-wide post-translational modification statistics: frequency analysis and curation of the swiss-prot database. *Sci. Rep.* **1**, 90 (2011).

2.   Apweiler, R., Hermjakob, H. & Sharon, N. On the frequency of protein glycosylation, as

deduced from analysis of the SWISS-PROT database. *Biochim. Biophys. Acta* **1473**, 4–8

(1999).

3.   RodrÍguez, E., Schetters, S. T. T. & van Kooyk, Y. The tumour glyco-code as a novel

immune checkpoint for immunotherapy. *Nat. Rev. Immunol.* **18**, 204–211 (2018).

4.   Gutierrez, J. M. *et al.* Genome-scale reconstructions of the mammalian secretory pathway

predict metabolic costs and limitations of protein secretion. Preprint at (*bioRxiv*)

https://www.biorxiv.org/10.1101/351387 (2018).

5.   Gabius, H. J., André, S., Kaltner, H. & Siebert, H. C. The sugar code: functional

lectinomics. *Biochimica et Biophysica Acta (BBA)* **1572**, 165–177 (2002).

6.   Spahn, P. N. & Lewis, N. E. Systems glycobiology for glycoengineering. *Curr. Opin.*

Biotechnol. **30**, 218–224 (2014).

7.   Holst, S. *et al.* N-glycosylation profiling of colorectal cancer cell lines reveals association of

fucosylation with differentiation and caudal type Homebox 1 (CDX1)/Villin mRNA

expression. *Mol. Cell. Proteomics* **15**, 124–140 (2016).

8.   Reiding, K. R., Blank, D., Kuijper, D. M., Deelder, A. M. & Wuhrer, M. High-throughput

profiling of protein N-glycosylation by MALDI-TOF-MS employing linkage-specific sialic

acid esterification. *Anal. Chem.* **86**, 5784–5793 (2014).

9.   Yang, Z. *et al.* Engineered CHO cells for production of diverse, homogeneous

glycoproteins. *Nat. Biotechnol.* **33**, 842–844 (2015).

10. Anugraham, M. *et al.* Specific glycosylation of membrane proteins in epithelial ovarian

cancer cell lines: glycan structures reflect gene expression and DNA methylation status.

Mol. Cell. Proteomics **13**, 2213–2232 (2014).

11. Čaval, T., Tian, W., Yang, Z., Clausen, H. & Heck, A. J. R. Direct quality control of

glycoengineered erythropoietin variants. *Nat. Commun.* **9**, 3342 (2018).

12. Riley, N. M., Hebert, A. S., Westphall, M. S. & Coon, J. J. Capturing site-specific

heterogeneity with large-scale N-glycoproteome analysis. *Nat. Commun.* **10**, 1311 (2019).

13. Aoki-Kinoshita, K. *et al.* GlyTouCan 1.0--The international glycan structure repository.

Nucleic Acids Res. **44**, D1237–42 (2016).

14. Campbell, M. P. *et al.* Validation of the curation pipeline of UniCarb-DB: building a global

glycan reference MS/MS repository. *Biochim. Biophys. Acta* **1844**, 108–116 (2014).

15. Campbell, M. P. *et al.* UniCarbKB: building a knowledge platform for glycoproteomics.

Nucleic Acids Res. **42**, D215–221 (2014).

16. Rademacher, C. & Paulson, J. C. Glycan fingerprints: calculating diversity in glycan

libraries. *ACS Chem. Biol.* **7**, 829–834 (2012).

17. Hosoda, M. *et al.* MCAW-DB: A glycan profile database capturing the ambiguity of glycan

recognition patterns. *Carbohydr. Res.* **464**, 44–56 (2018).

18. Alocci, D. *et al.* Understanding the glycome: an interactive view of glycosylation from

glycocompositions to glycoepitopes. *Glycobiology* **28**, 349–362 (2018).

19. Klein, J., Carvalho, L. & Zaia, J. Application of network smoothing to glycan LC-MS

profiling. *Bioinformatics* **34**, 3511–3518 (2018).

20. Sharapov, S. *et al.* Defining the genetic control of human blood plasma N-glycome using

genome-wide association study. Preprint at (*bioRxiv*)

https://www.biorxiv.org/10.1101/365486 (2018).

21. Mohammad, M. A., Hadsell, D. L. & Haymond, M. W. Gene regulation of UDP-galactose

synthesis and transport: potential rate-limiting processes in initiation of milk production in

humans. *Am. J. Physiol. Endocrinol. Metab.* **303**, E365–376 (2012).

22. Hanley, J. A., Negassa, A., Edwardes, M. D. D. & Forrester, J. E. Statistical analysis of

correlated data using generalized estimating equations: an orientation. *Am. J. Epidemiol.*

157, 364–375 (2003).

23. Koda, Y., Soejima, M., Liu, Y. & Kimura, H. Molecular basis for secretor type alpha(1,2)-

fucosyltransferase gene deficiency in a Japanese population: a fusion gene generated by

unequal crossover responsible for the enzyme deficiency. *Am. J. Hum. Genet.* **59**, 343–350

(1996).

24. Kudo, T. *et al.* Molecular genetic analysis of the human Lewis histo-blood group system. II.

Secretor gene inactivation by a novel single missense mutation A385T in Japanese

nonsecretor individuals. *J. Biol. Chem.* **271**, 9830–9837 (1996).

25. Viverge, D., Grimmonprez, L., Cassanas, G., Bardet, L. & Solere, M. Discriminant

carbohydrate components of human milk according to donor secretor types. *J. Pediatr.*

Gastroenterol. Nutr. **11**, 365–370 (1990).

26. Yan, J. & Fine, J. Estimating equations for association structures. *Stat. Med.* **23**, 859–874;

discussion 875–877, 879–880 (2004).

27. Halekoh, U., Højsgaard, S., Yan, J. The R package geepack for generalized estimating

equations. *J. Stat. Softw.* **15**, 1–11 (2006).

Figure Legends

Fig. 1 | The GlyCompare workflow for glycoprofile decomposition and comparison. a and b, A glycoprofile with annotated structure and relative abundance is obtained and the glycans are decomposed to a substructure vector S and the presence/absence matrix X is built. Each substructure vector is weighted by the glycan abundance, and the weighted substructure vectors are summed into a glycoprofile vector p. **c**, Seven example glycoprofiles are represented here with their glycoprofile vectors. The glycoprofile vectors are simplified to contain a minimal number of substructures, so a substructure network is constructed to identify the non-redundant glyco-motifs that change in abundance from their precursor substructures. **d**, The glycoprofiles can be re-clustered with simplified glyco-motif vectors for a clearer result. **e**, Clustered substructures can be analyzed to identify the most representative structure in the group. For example, four substructures with different relative abundance were aligned together and the monosaccharides' with weight over 51% was preserved.

Fig. 2 | Glycompare effectively clusters panels of distinct glycoprofiles and identifies differentially expressed representative substructures by resolving sparsity. a, Sixteen glycoprofiles from glycoengineered recombinant EPO cluster poorly when based solely on raw glycan abundance. **b**, GlyCompare was used to compute and cluster EPO glyco-motif vectors. This resulted in three dominant clusters of glycoprofiles and a few individuals that have very different structural pattern (distance threshold=0.5) and twenty-four clusters of glycan substructure (distance threshold=0.19). **c**, The representative substructure table contains representative substructures for each of the 24 substructure clusters. The color scale represents the averaged abundances of the substructures in each cluster. The substructures are sorted based

on the glycan structure complexity, followed by the number of branches, the degree of

galactosylation, sialylation, and fucosylation. **d**, The significantly differentially expressed glycan

substructures are illustrated by z-score normalized abundance of twenty-four glycan

substructures, compared with WT.

Fig. 3 | Changes in representative substructures are quantified and compared to WT with

the standardized abundance bar plot. a, Differential fucosylation is illustrated for the Fut8

knockout. The red (black) triangles represent the presence/absence of fucose in the representative

substructures. Differential sialylation is illustrated for the St3gal4/6 knockout. The purple/black

diamonds represent the presence/absence of the sialylation in the representative substructures. **b**,

Changes in branching are presented for the Mgat4A/4B/5 knockouts. The tetra-antennary

substructures (Rep16-Rep22) decreased considerably. The triantennary substructures with

elongated GlcNac (Rep13-Rep14) increase significantly (p-value < 0.01). However, the

elongated triantennary structure (Rep15) decreases significantly for the Mgat5 and Mgat4B

knockouts (p-value < 0.001). In the CHO dataset, the glycan substructure generated by

Mgat4A/4B and Mgat5 will be considered as the same topologically.

Fig. 4 | Analysis of intermediate substructures with GlyCompare elucidates associations

with secretor status over time, which are missed in the standard whole-glycan analysis. a,

The substructure intermediates for four connected HMOs are shown here. The synthesis of larger

HMOs must pass through intermediate substructures that are also observed HMOs, where the

substructures are as associated with measured HMOs as follow X38=LNT, X80=LSTb,

X103=DSLNT, X138=DSLNH. **b-e**, Over time (DPP), X80, LSTb, DSLNT, and DSLNH show

different trends for secretors and non-secretors. Furthermore, the abundance of aggregated X80

shows significant positive-correlation with secretor and negative-correlation with non-secretor.

Fig. 5 | Product-Reactant Ratio of two reactions. X38, the LNT substructure, is a precursor to

X80, the LSTb substructure, which is a precursor to X103, the DSLNT substructure. We estimate

the flux of these conversions from X38 to X80 and X80 to X103 by examining the

product/reactant ratio; the proportion of the total synthesize reactant converted to the product. **a**,

LSTb/LNT motif relative abundance ratios are not associated with secretor status while **b**,

DSLNT/LSTb ratios are. Each line is a simple linear regression leveraging sample-specific data.

Fig. 1 | The GlyCompare workflow for glycoprofile decomposition and comparison.

Fig. 2 | Glycompare effectively clusters panels of distinct glycoprofiles and identifies

differentially expressed representative substructures by resolving sparsity.

Fig. 3 | Changes in representative substructures can be quantified and compared to WT

567   **with the standardized abundance bar plot.**

Fig. 4 | Analysis of intermediate substructures with GlyCompare elucidates associations

 with secretor status over time, which are missed in the standard whole-glycan analysis.

574 **Fig. 5 | Product-Reactant Ratio of two reactions.**