1

Expression Profiling, Downstream Signalling and Inter-Subunit Interactions

of GPA2/GPB5 in the Adult Mosquito *Aedes aegypti*

Short Title: Characterization of GPA2/GPB5 Signalling in Mosquitoes

David A. Rocco[1*] and Jean-Paul V. Paluzzi[1*]

1. Department of Biology, York University, Toronto, Ontario, Canada, M3J1P3

*Corresponding Author Information:

1. David A. Rocco

    Department of Biology, York University, Toronto, Ontario, Canada, M3J1P3

    (416) 301- 4843

    davrocco@yorku.ca

    ORCID ID: 0000-0001-9545-7713

2. Jean-Paul Paluzzi

    Department of Biology, York University, Toronto, Ontario, Canada, M3J1P3

    (416) 736-2100 ext. 20999

    paluzzi@yorku.ca

    ORCID ID: 0000-0002-7761-0590

Abstract

GPA2/GPB5 and its receptor constitute a glycoprotein hormone-signalling system native to the genomes of most vertebrate and invertebrate organisms, including humans and mosquitoes. Unlike the well-studied gonadotropins and thyrotropin, the exact function of GPA2/GPB5 is unclear, and whether it elicits its functions as heterodimers, homodimers or as independent monomers remains unknown. Here, the glycoprotein hormone signalling system was investigated in adult mosquitoes, where GPA2 and GPB5 subunit transcripts co-localized to bilateral pairs of neuroendocrine cells within the first five abdominal ganglia of the central nervous system. Unlike human GPA2/GPB5 that demonstrated strong heterodimerization between subunits, the GPA2/GPB5 subunits in *A. aegypti* lacked evidence of heterodimerization when heterologously expressed. Interestingly, cross-linking analysis to determine subunit interactions revealed *A. aegypti* and *H. sapiens* GPA2 and GPB5 subunits form homodimers, and treatments with independent subunits did not activate *A. aegypti* LGR1 or *H. sapiens* TSH receptor, respectively. Since mosquito GPA2/GPB5 heterodimers were not evident by heterologous expression of independent subunits, a tethered construct was generated for expression of the subunits as a single polypeptide chain to improve heterodimer formation. Our findings revealed *A. aegypti* LGR1 elicited constitutive activity that elevated levels of cAMP as determined by increased cAMP-dependent luminescence. However, upon treatment with recombinant tethered GPA2/GPB5 heterodimers, an inhibitory G protein (Gi) signalling cascade is initiated and forskolin-induced cAMP production is inhibited. These results provide evidence towards the functional deorphanization of LGR1 and, moreover, further support the notion that GPA2/GPB5 heterodimerization is a requirement for glycoprotein hormone receptor activation.

44   Keywords: GPA2/GPB5, Mosquito, Glycoprotein hormone, Leucine-rich repeat-containing G

45   protein coupled-receptor 1, Homodimer, Heterodimer, Thyrostimulin

Significance Statement

The GPA2/GPB5 glycoprotein hormone signalling system is an evolutionarily ancient one that is involved in multiple physiological functions in diverse animal species, ranging from insects to worms to humans; however, its mode of signalling remains poorly characterized. To better understand how GPA2/GPB5 may exert its function, hormone subunit expression was localized in adult mosquitoes, and properties of subunit interaction along with receptor coupling for human and mosquito GPA2/GPB5 signalling was investigated. Our findings indicate certain aspects of GPA2/GPB5 signalling are unique from that of the classic glycoprotein hormones (i.e. the gonadotropins and thyrotropin) and could implicate independent physiological roles for heterodimers and homodimers in mosquitoes and humans.

Introduction

Members of the cystine knot growth factor (CKGF) superfamily, which are characterized with a CKGF domain as their primary structural feature, include (i) the glycoprotein hormones, (ii) the invertebrate bursicon hormone, (iii) the transforming growth factor beta (TGFβ) family, (iv) the bone morphogenetic protein (BMP) antagonist family, (v) the platelet-derived growth factor (PDGF) family and (vi) the nerve growth factor (NGF) family. Of the members of the CKGF superfamily, the glycoprotein hormones are of fundamental importance in the regulation of both vertebrate and invertebrate physiology.

In vertebrates, members of the glycoprotein hormone family include follicle-stimulating hormone (FSH), luteinizing hormone (LH), thyroid-stimulating hormone (TSH) as well as chorionic gonadotropin (CG), which are implicated in governing several aspects of physiology including reproduction, energy metabolism along with growth and development. Structurally, these hormones are formed by the heterodimerization of two cystine-knot glycoprotein subunits, an α subunit that is structurally identical for each hormone (GPA1), and a hormone-specific β subunit (GPB1-4) (1, 2).

Two novel glycoprotein hormone subunits were identified in the human genome, glycoprotein α2 (GPA2) and glycoprotein β5 (GPB5), and were found to heterodimerize (GPA2/GPB5) and act on the same receptor as TSH. As a result, GPA2/GPB5 was coined the name thyrostimulin to differentiate it from TSH in vertebrates (3, 4). Unlike other glycoprotein hormones which are restricted to the vertebrate lineage, homologous genes encoding GPA2/GPB5 subunits exist in all bilaterian organisms, where its function appears to be pleiotropic (3–5). In vertebrates, the function of GPA2/GPB5 has been implicated, or at least suggested, to be involved in reproduction (6), thyroxine production (3, 7), skeletal development

(8), immunoregulation (9, 10) and the proliferation of ovarian cancer cell lines (11). For

invertebrate species, GPA2/GPB5 has been implicated or demonstrated to function in

development (12–15), hydromineral balance (13–18) as well as in reproduction (18, 19).

Non-covalent interactions and heterodimerization between the subunits forming FSH,

LH, TSH and CG is required for their respective biological functions (1, 20). However, whether

heterodimerization is required for GPA2/GPB5 to activate its receptor and exert its physiological

role in vertebrates and invertebrates is debated. For FSH, LH, TSH and CG, the subunits are co-

expressed in the same cells and each hormone is released into circulation as heterodimers (21,

22). On the other hand, GPA2 and GPB5 subunit expression profiles, both for vertebrate and

invertebrate organisms, do not always occur in the same cells, and GPA2 is often expressed more

widely and abundantly than GPB5 in some tissues (7, 12, 23–25). Additionally, unlike the beta

subunits of the classic glycoprotein hormones, the structure of the GPB5 subunit lacks an extra

pair of cysteine residues that form an additional disulfide linkage, referred to as the 'seatbelt',

which strengthens and stabilizes its heterodimeric association with GPA2 (26).

Relative to other G protein-coupled receptors (GPCRs), members of the leucine-rich

repeat-containing G protein-coupled receptor (LGR) family are often characterized with a large

extracellular amino terminal domain responsible for the selective binding of their large hormone

ligands (27). Following its initial genomic and molecular characterization (28), an invertebrate

receptor for GPA2/GPB5, called LGR1, was functionally deorphanized in the fruit fly

Drosophila melanogaster, where GPA2/GPB5 heterodimers were found to activate LGR1 and

increased intracellular levels of cyclic AMP (cAMP) (29). Interestingly, stimulatory G protein

(Gs) coupling and signalling to elevate cAMP was also shown with GPA2/GPB5-TSH receptor

activation in humans (3, 6).

In the mosquito *Aedes aegypti*, genes encoding for GPA2, GPB5 and LGR1 were

identified and shown to be expressed in all developmental life stages, with expression enriched

in adults compared to juvenile stages (14). In adults, LGR1 was found localized to epithelia

throughout the gut where GPA2/GPB5 could regulate feeding-related processes and

hydromineral balance (18). Notably, LGR1 transcript expression was also observed in the

reproductive organs of males and females (19). In adult male mosquitoes, knockdown of LGR1

expression led to abnormal spermatogenesis with spermatozoa displaying malformations such as

shortened flagella and consequently, LGR1-knockdown males had 60% less spermatozoa as well

as significantly reduced fecundity relative to control mosquitoes (19).

With an interest in better understanding GPA2/GPB5 signalling in *A. aegypti* mosquitoes,

our work herein set out to characterize the tissue-specific and cellular distribution expression

profile of GPA2/GPB5 in mosquitoes. As well, we sought to determine GPA2/GPB5 interactions

and demonstrate GPA2/GPB5-LGR1 functional coupling using a heterologous system.

Combining various molecular techniques, we demonstrate GPA2/GPB5 cellular co-expression in

the central nervous system of adult mosquitoes, and that heterodimers are indeed required to

activate LGR1, which exhibits ligand-dependent G protein-coupling activity. Moreover, these

results provide evidence for human and mosquito GPA2 and GPB5 homodimers, which were

incapable of activating TSH receptor or LGR1, respectively. Overall, these findings appreciably

advance our understanding of GPA2/GPB5 signalling in mosquitoes and provide novel

directions to uncover the functions of homologous systems in other organisms.

Results

A. aegypti **GPA2 and GPB5 subunit expression localization**

To determine the distribution of GPA2 and GPB5 subunit expression in the central

nervous system and peripheral tissues, adult mosquito organs were analyzed using RT-qPCR.

GPA2 and GPB5 subunit transcript was detected in the central nervous system of adult male and

female mosquitoes, with significantly enriched expression in the abdominal ganglia relative to

other nervous and peripheral tissues (Fig. 1A-1B). Fluorescence *in situ* hybridization was

employed to localize cell-specific expression of the GPA2 and GPB5 transcripts in the

abdominal ganglia. GPA2 and GPB5 anti-sense RNA probes identified two bilateral pairs of

neuroendocrine cells (Fig. 1C-1D) within each of the first five abdominal ganglia in male and

female mosquitoes whereas the control sense probes did not detect cells in the nervous system

(Fig. 1E-1F). Within each of these first five abdominal ganglia, GPA2 transcript localized to

similar laterally-positioned cells as the GPB5 transcript (Fig. 1G-1H). To determine if cells

expressing GPA2 transcript were the same cells expressing GPB5 transcript, abdominal ganglia

were simultaneously treated with both GPA2 and GPB5 anti-sense RNA probes. Using this dual

probe approach, results confirmed the detection of only two bilateral pairs of cells (Fig. 1I-1J)

that displayed a greater staining intensity compared to the intensity of cells detected when using

either the GPA2 or GPB5 anti-sense probe alone (Fig. 1C-1D).

Using a custom antibody targeting *A. aegypti* GPB5, we next sought to immunolocalize

GPB5 protein in the abdominal ganglia. GPB5 immunoreactivity localized to two bilateral pairs

of cells (Fig. 2A) within the first five ganglia, which were in similar positions to cells expressing

GPA2 and GPB5 transcript (Fig. 1C-1D, 1G-J). In some preparations, three bilateral pairs of

cells immunolocalized to the first five abdominal ganglia of the ventral nerve cord (Fig. 2B),

however no cells were ever detected in the sixth terminal ganglion (Fig. 2C). Along the lateral

sides of each of the first five abdominal ganglia, GPB5 immunoreactive processes were observed

to interconnect and closely associate as a tract of axons emanating through the lateral nerve (Fig.

2B). Control treatments with GPB5 antibody preabsorbed with the GPB5 immunogenic antigen,

failed to detect any cells in ganglia (Fig. 2D)

Cross-linking analyses to determine *A. aegypti* GPA2/GPB5 dimerization patterns

A. aegypti GPA2 and GPB5 subunit protein interactions were studied using western blots

of recombinant proteins from HEK 293T cells expressing each subunit independently, or co-

expressing both subunits in the same cells using a dual promoter plasmid. Under control

conditions, GPA2 protein is represented as two bands at 16 kDa and 13 kDa, which correspond

to the glycosylated and non-glycosylated forms of *A. aegypti* GPA2, respectively (Fig. 3A).

Following deglycosylation with PNGase, the higher molecular weight band of GPA2 is

eliminated, and the non-glycosylated lower molecular weight band intensifies (Fig. 3A).

Interestingly, when the GPA2 subunit was tested to examine potential homodimerization, an

additional strong band at ~32 kDa was detected, which migrates to ~30 kDa when cross-linked

protein samples were deglycosylated using PNGase (Fig. 3A). Under control conditions, GPB5

protein is represented as a band size at 24 kDa and the migration pattern is not affected by

PNGase treatment (Fig. 3B). Upon treatment with DSS, a faint second band appears at 48 kDa,

which does not change in molecular weight following treatment with PNGase (Fig. 3B). Three

independent band sizes at 24 kDa (GPB5), 16 kDa (glycosylated GPA2) and 13 kDa (non-

glycosylated GPA2) were detected in lanes loaded with protein isolated from HEK 293T cells

co-expressing GPA2 and GPB5 subunits (Fig. 3C). After removing N-linked sugars, the 24 kDa

band is not affected but the 16 kDa band disappears and 13 kDa band intensifies (Fig. 3C), as

observed when assessing the GPA2 subunit independently (Fig. 3A). Cross-linked samples show

the addition of two higher molecular weight bands at ~48 kDa and ~32 kDa, the latter of which

migrates lower to ~30 kDa when subjected to PNGase treatment (Fig. 3C). Given the resolution

and intensity of the detected bands, molecular weight bands at ~48 kDa and ~32 kDa could

indicate GPA2/ GPB5 heterodimeric interactions (13 or 16 kDa + 24 kDa = 37-40 kDa).

Alternatively, these bands could reflect GPA2 (13 or 16 kDa + 13 or 16 kDa = 26-32 kDa) and

GPB5 (24 + 24kDa = 48 kDa) homodimers. As a result, to clarify whether *A. aegypti* GPA2 and

GPB5 subunits are heterodimeric partners, additional experiments were performed.

Heterodimerization of mosquito and human GPA2/ GPB5

Using yeast two-hybrid analyses and cross-linking experiments, it was previously shown

that human GPA2 (hGPA2) and GPB5 (hGPB5) subunits are capable of heterodimerization (3).

As a result, to verify whether *A. aegypti* GPA2 and GPB5 subunit proteins are heterodimeric

candidates, experiments were first performed alongside hGPA2/hGPB5 subunit proteins, using

the latter as an experimental control.

Initially, single-promoter expression constructs were designed to incorporate a FLAG-tag

and His-tag on the C-terminus of (human and mosquito) GPA2 and GPB5 subunits, respectively

(i.e. GPA2-FLAG and GPB5-His). When probed with an anti-His antibody, no bands were

detected in lanes containing only GPA2-FLAG (human and mosquito) protein (Fig. 4A-4B).

Similarly, no bands were detected when an anti-FLAG antibody was used to detect GPB5-His

protein (human and mosquito) (Fig. 4A-4B). Lanes loaded with cross-linked human GPB5-His

protein revealed two bands at 18 kDa (monomer) and 36 kDa (homodimer) (Fig. 4A). Similarly,

in lanes loaded with cross-linked human GPA2-FLAG protein, two bands at 20 kDa (monomer)

and 40 kDa (homodimer) were detected (Fig. 4A). To determine subunit heterodimerization,

GPA2-FLAG protein was combined with GPB5-His protein (i.e. produced separately in different

cell batches) and subsequently crosslinked. Results showed an intense 38 kDa band size that

correlated to the molecular weight of the hGPA2-FLAG/hGPB5-His heterodimers; detected with

both anti-His and anti-FLAG primary antibody solutions (Fig. 4A).

The same experiments were replicated using *A. aegypti* GPA2-FLAG/ GPB5-His protein

but using three-fold higher concentration of DSS cross-linker to help improve detection of inter-

subunit interactions. Results demonstrated that lanes loaded with DSS-treated GPB5-His protein

resulted in a 24 kDa monomer band (Fig. 4B), whereas lanes containing cross-linked GPA2-

FLAG detected a 16 kDa (glycosylated monomer), 30 kDa and 32 kDa (homodimers) band size

(Fig. 4B). As a result, unlike immunoblots containing hGPA2-FLAG and hGPB5-His protein

(Fig. 4A), lanes loaded with cross-linked *A. aegypti* GPA2-FLAG and GPB5-His failed to

provide evidence of bands correlating to the predicted molecular weight of an *A. aegypti*

GPA2/GPB5 heterodimer, which would be expected at 37-40 kDa (Fig. 4B).

***A. aegypti* GPA2/GPB5 unable to activate LGR1-mediated Gs and Gi/o signalling pathways**

Bioluminescent assays were employed to confirm LGR1 interaction with mosquito GPA2

and/or GPB5 subunits and elucidate downstream signalling pathways upon receptor activation.

Since human GPA2/GPB5 has previously been shown to bind and activate human thyrotropin

receptor (hTSHR) mediating a stimulatory G protein (Gs) signalling pathway (3), our

experimental design was first tested using hGPA2/GPB5 and hTSHR as a positive control.

Recombinant hGPA2 and hGPB5 subunit proteins were produced in HEK 293T cells with single

promoter expression constructs containing the hGPA2 or hGPB5 sequences. Conditioned culture

media containing secreted proteins were subsequently concentrated, and crude extracts

containing hGPA2, hGPB5 or a combination of hGPA2 and hGPB5 subunits were tested as

ligands on HEK 293T cells expressing the hTSHR and a mutant luciferase biosensor, which

produces bioluminescence upon interacting with cAMP. Control treatments involved the

incubation of hTSHR/ luciferase-expressing cells with concentrated media collected from

mCherry-transfected cells (negative control), or 250 nM forskolin (positive control) (Fig. 5A-

5B). Our findings indicate that incubation with extracts containing a combination of both hGPA2

and hGPB5 were required to stimulate a cAMP-mediated luminescent response from hTSHR-

expressing cells, but not when incubated with extracts containing individual subunits (Fig. 5A).

We next performed parallel experiments using *A. aegypti* GPA2/GPB5 and LGR1. Given the

availability of a dual promoter plasmid (pBud-CE), an additional treatment was performed

whereby GPA2/GPB5 subunits were co-expressed within the same cells and conditioned media

was concentrated as above. Unlike hGPA2/hGPB5 subunit homologs (Fig. 5A), no combination

of mosquito GPA2 and/or GPB5 subunit proteins led to an increase in the luminescent response

(Fig. 5B).

Using an *in silico* analysis to predict coupling specificity of *A. aegypti* LGR1 and human

TSHR to different families of G-proteins (30), we determined that *A. aegypti* LGR1 is strongly

predicted to couple to inhibitory (Gi) G proteins (Table S3). As a result, to determine whether *A.*

aegypti GPA2 and/or GPB5 activate a Gi/o signalling pathway, various combinations of GPA2

and GPB5 were tested for their ability to inhibit a forskolin-induced rise in cAMP measured by

changes in bioluminescence (Fig. 5C-5D). Results revealed that sole treatments of GPA2 and

GPB5 proteins alone significantly inhibited a forskolin-induced luminescent response, relative to

control treatments with mCherry proteins, when incubated with cells expressing LGR1 (Fig. 5C).

However, similar inhibitory effects of GPA2 and GPB5 proteins were also observed with cells

not expressing LGR1 (Fig. 5D).

Characterization of a tethered *A. aegypti* GPA2/GPB5 heterodimer

Activation of hTSHR was only observed when both human GPA2 and GPB5 subunits

were coapplied for receptor binding (Fig. 5A) and, unlike the heterodimerization of human

GPA2/GPB5 observed in our experiments, mosquito GPA2/GPB5 lacked evidence of

heterodimerization (Fig. 4). In light of these observations, it was hypothesized that the activation

of *A. aegypti* LGR1 also required subunit heterodimerization. To produce stable GPA2/GPB5

heterodimers using the heterologous expression system, both GPA2 and GPB5 mosquito

subunits were expressed as a tethered, single-chain polypeptide by fusing the C-terminus of the

GPB5 prepropeptide sequence with the N-terminus of the GPA2 propeptide sequence using a

tagged linker sequence composed of twelve amino acids, involving three glycine-serine repeats

and six histidine residues.

HEK 293T cells were transfected to transiently express a single promoter plasmid

construct containing the tethered GPA2/GPB5 sequence, or the red fluorescent protein

(mCherry) as a negative control. At 48-hrs post-transfection, cell lysates along with the

conditioned culture media, the latter of which contains secreted proteins, were collected for

immunoblot analysis. No bands were detected in lanes containing cell lysate or secreted protein

fractions of mCherry transfected cells (Fig. 6A). However, a strong band was detected at 32-40

kDa in the lysates of cells transfected to express tethered GPA2/GPB5 (Fig. 6A). Moreover, two

bands were detected at 37 kDa and 40 kDa in lanes loaded with secreted fractions of tethered

GPA2/GPB5-transfected cells, that matched the predicted molecular weights of *A. aegypti*

GPA2/GPB5 heterodimers corresponding to either non-glycosylated (13 kDa) or glycosylated

(16 kDa) GPA2 plus GPB5 (24 kDa) producing bands of 37 kDa or 40 kDa, respectively (Fig.

6A). After secreted protein extracts containing tethered GPA2/GPB5 proteins were treated with

PNGase, the higher 40 kDa molecular weight band disappears and the 37 kDa band size

intensifies, which confirms the observed molecular weight shift results from removal of N-linked

oligosaccharides (Fig. 6B).

***A. aegypti* GPA2/GPB5 heterodimers activate LGR1**

The effects of tethered GPA2/GPB5 heterodimers on LGR1 activity was examined. Cell

lysates or secreted protein fractions collected from mCherry- or tethered GPA2/GPB5-

transfected cells were incubated with HEK 293T cells co-expressing the cAMP luciferase

biosensor and either *A. aegypti* LGR1 or mCherry (i.e. not expressing LGR1). Whether tethered

GPA2/GPB5 proteins could elevate cAMP or inhibit a forskolin-induced rise in cAMP was

assessed and compared to control treatments with proteins harvested from mCherry-transfected

cells. Overall, the relative luminescent response was significantly greater in LGR1-transfected

cells compared to mCherry-transfected cells (Fig. 7). Unlike treatments with forskolin that

significantly increased cAMP-mediated luminescence, neither secreted protein fractions nor cell

lysates of tethered GPA2/GPB5-transfected cells elicited an increase in the cAMP-mediated

luminescent response relative to controls (Fig. 7A-7B). Secreted protein fractions containing

tethered GPA2/GPB5 protein had no effect on the forskolin-induced cAMP-mediated

luminescence, compared to control treatments with mCherry secreted proteins in LGR1-

expressing cells (Fig. 7C). Notably, however, treatments of LGR1-transfected cells, but not

mCherry-transfected cells or assay media, with cell lysates containing tethered GPA2/GPB5

protein significantly inhibited the forskolin-induced rise in cAMP relative to treatments with

mCherry-transfected cell lysates (Fig. 7D).

Discussion

Co-expression of *A. aegypti* GPA2/GPB5 in neuroendocrine cells of the abdominal ganglia

The central nervous system (CNS) of adult mosquitoes is comprised of a brain and a ventral nerve cord, consisting of three fused thoracic ganglia and six abdominal ganglia. Our findings demonstrate that the GPA2 and GPB5 transcripts are significantly enriched in the abdominal ganglia of adult mosquitoes relative to peripheral tissues and other regions of the CNS, with no sex-specific differences. Although low levels of GPA2 and GPB5 transcripts were detected in the thoracic ganglia and brain using RT-qPCR, fluorescence *in situ* hybridization (FISH) techniques used to localize GPA2 and GPB5 transcripts, along with immunohistochemical detection of GPB5, did not identify specific cells in these regions of the nervous system. Instead, GPA2 and GPB5 transcripts, as well as GPB5 immunoreactivity was identified in 2-3 laterally-localized bilateral pairs of neuroendocrine cells within the first five abdominal ganglia. These findings are consistent with previous findings in the fruit fly *Drosophila melanogaster*, where GPA2 and GPB5 subunit transcripts were localized to four bilateral pairs of neuroendocrine cells in the fused ventral nerve cord, that were distinct from cells expressing other neuropeptides including leucokinin, bursicon, crustacean cardioactive peptide or calcitonin-like diuretic hormone (16).

Both FISH and immunohistochemical techniques revealed GPA2 and GPB5 expression within two bilateral pairs of neuroendocrine cells, that were positioned slightly posterior to where the lateral nerves emanate. In some abdominal ganlia preparations, a third bilateral pair of cells immunoreactive for GPB5 protein was detected; however these additional cells were not detected using FISH, which suggests GPB5 transcript may be differentially regulated between

different bilateral pairs of cells. Given that the same number of cells were observed to express

GPA2 transcript, and these cells localized to similar positions as GPB5 expressing cells, GPA2

and GPB5 were believed to be co-expressed in the same cells. To verify cellular co-expression of

GPA2/GPB5, abdominal ganglia were simultaenously treated with both GPA2- and GPB5-

targeted anti-sense RNA probes. From this analysis, again only two bilateral pairs of cells were

detected and these were more intensely stained compared to preparations treated with either probe

alone, which confirms that GPA2 and GPB5 are indeed co-expressed within the same

neurosecretory cells of the first five abdominal ganglia in adult mosquitoes. The cellular co-

expression of GPA2 and GPB5 proteins implies that, upon a given stimulus, both subunits are

regulated in a similar manner and are likely simultaneously released following the appropriate

stimulus. Importantly, since co-expression and heterodimerization of the classic vertebrate

glycoprotein hormone subunits takes place within the same cells (21, 22), these findings indicate

that the mosquito GPA2/GPB5 subunits may be produced and released as heterodimers *in vivo*.

Heterodimerization and Homodimerization of GPA2/GPB5

To study the interactions of *A. aegypti* GPA2 and GPB5 subunits *in vitro*, hexa-histidine

tagged proteins secreted into the culture media of transfected HEK 293T cells were collected and

analysed under denaturing conditions after cross-linker treatments, which had been utilized

previously to show GPA2/GPB5 heterodimerization in other organisms (3, 6, 29). Cross-linked

protein samples were then deglycosylated to identify whether the removal of N-linked sugars

affected dimerization. Treatment of mosquito GPA2 and GPB5 subunits individually with cross-

linker resulted in the detection of bands with sizes corresponding to homodimers of GPA2 (~32

kDa) and GPB5 (48 kDa). GPA2 homodimer bands migrated lower to ~30 kDa following

deglycosylation treatment with PNGase. However, experiments performed with cross-linked

GPA2/GPB5 protein were not able to confirm heterodimeriation since the detected band sizes

could also reflect GPA2 and GPB5 homodimeric interactions. Previous cross-linking studies that

demonstrated GPA2/GPB5 heterodimerization in human (3, 6) and fruit fly (29) did not provide

evidence on the interactions of each subunit alone to determine if homodimerization is possible.

In these earlier studies, molecular weight band sizes that were identified as heterodimers could

have been the result of homodimeric interactions (3, 6, 29). As a result, the findings herein with

mosquito GPA2/GPB5 indicate additional experiments are required to confirm GPA2/GPB5

heterodimerization in these organisms.

To clarify whether *A. aegypti* (mosquito) GPA2/GPB5 heterodimerize, each subunit was

differentially tagged (GPA2-FLAG and GPB5-His), and immunoblots containing various

combinations of cross-linked subunits were probed with either anti-FLAG or anti-His antibody.

As a positive control, experiments were first performed using *H. sapiens* (human) GPA2/GPB5

subunit proteins. Similar to mosquito GPA2 and GPB5 subunits, results showed human GPA2

and GPB5 subunits are capable of homodimerization. To study heterodimeric interactions, GPA2

and GPB5 subunit proteins were expressed separately in HEK 293T cells. Upon combining and

treating protein samples with DSS, a molecular weight band size at 38 kDa, corresponding to the

molecular weight of GPA2/GPB5 heterodimers (human GPA2-FLAG/ GPB5-His), was detected

and migrated differently than bands corresponding to GPA2 (40 kDa) and GPB5 (36 kDa)

homodimers. Taken together, our results confirm human GPA2 and GPB5 is indeed capable of

heterodimerization *in vitro*. Further, an induction of cAMP was observed when both subunits

were present for TSHR functional activation, whereas treatments with individual subunits failed

to significantly increase cAMP-mediated luminescence. As a result, human GPA2/GPB5 is

capable of heterodimerization, and since a combination of both subunits were required to signal a

TSHR-mediated elevation in cAMP, these heterodimers are required to functionally activate its

cognate glycoprotein hormone receptor (TSHR).

The classic glycoprotein hormones, which include FSH, LH, TSH and CG, are formed by

a common alpha subunit non-covalently linked to a hormone-specific beta subunit. Each subunit

is co-expressed and assembled as a heterodimer in the endoplasmic reticulum within cells before

being secreted as heterodimer into circulation (21). Heterodimerization is required for secretion

and receptor binding activity of each hormone (1, 31, 32). Our results confirm human

GPA2/GPB5 heterodimerizes, as detected following chemical cross-linking treatments and, even

without cross-linking, both subunits are required to activate TSHR. In our experimental

approach, human GPA2 and GPB5 subunits were not co-expressed within the same cells using a

dual promoter expression construct. Rather, subunits were separately expressed by different

batches of cells and mixed post harvesting of conditioned media to demonstrate

heterodimerization and TSHR activation. Thus, unlike FSH, TSH, CG, and LH (1), our results

provide novel information that demonstrate co-expression is not necessarily required for

heterodimerization of human GPA2/GPB5 *in vitro*.

Analogous experiments using the same concentration of DSS cross-linker (data not

shown) as well as three-fold higher concentrations were performed with mosquito GPA2-FLAG

and GPB5-His subunit proteins. Irrespective of the DSS concentration used, no bands

corresponding to the expected molecular weights of GPA2/GPB5 heterodimers (37-40 kDa)

were observed, but rather, only GPA2 and GPB5 homodimers were detected (Fig. 4B). The

inability of mosquito GPA2 and GPB5 subunits to successfully heterodimerize could result from

improper protein folding of insect-derived secretory proteins in HEK 293T cells used for

heterologous expression. As a result, future experiments should test the heterologous expression

and heterodimerization of *A. aegypti* GPA2/GPB5 in insect cell lines that could provide a more

appropriate environment for tertiary and quaternary protein structure formation.

Mosquito GPA2/GPB5 (each subunit alone, mixed from different cell batches or co-

expressed in the same cells using a dual promoter vector) was incapable of stimulating a cAMP-

mediated luminescent response in HEK 293T cells expressing LGR1. Since we identified *A.*

aegypti LGR1 is predicted to couple a Gi/o signalling pathway (Table S3), we next examined if

mosquito GPA2/GPB5 could inhibit a forskolin-induced cAMP response. Sole treatments of

GPA2 and GPB5 subunit proteins inhibited a forskolin-induced rise in cAMP, however these

inhibitory actions were not owed to G protein signalling events related to *A. aegypti* LGR1, since

inhibition was observed in control cell lines in the absence of LGR1. These results suggest

mosquito GPA2 and GPB5 subunit proteins may non-specifically interact with other

endogenously expressed proteins, like the orphan glycoprotein hormone receptors LGR4 and

LGR5 that are highly expressed in HEK 293T cells (33).

***A. aegypti* GPA2/ GPB5 heterodimers activate LGR1 and initiate a switch from Gs to Gi**

coupling

Activation of the human thyrotropin receptor was only observed when both human GPA2

and GPB5 subunits were present which, unlike data obtained involving mosquito subunits,

demonstrated human GPA2/GPB5 subunit heterodimerization *in vitro* using the mammalian

heterologous system. Nonetheless, given the observed co-localization of the subunits within

bilateral pairs of cells in the first five abdominal ganglia in *A. aegypti*, which is comparable to

cellular colocalization shown earlier in *D. melanogaster* (16), we proposed that *A. aegypti*

GPA2/GPB5 heterodimers would be required to functionally activate LGR1 *in vitro*. To confirm

this possibility, a tethered construct was designed linking the C-terminus of GPB5 to the N-

terminus of GPA2 using a histidine tagged glycine/serine-rich linker sequence. Natural and

synthetic linkers function as spacers that connect multidomain proteins, and are commonly used

to study unstable or weak protein-protein interactions (34). The incorporation of a linker

sequence between glycoprotein hormone subunits has been performed previously, and does not

affect the assembly, secretion or bioactivity of human FSH (35), TSH (36) and CG (37). The

conversion of two independent glycoprotein hormone subunits into a single polypeptide chain

using a glycine-serine repeat linker sequence has also been performed recently with lamprey

GPA2/GPB5 (38), which was shown to induce a cAMP response. Interestingly, similar proteins

involving TSH alpha fused to TSH beta with carboxyl-terminal peptide (CTP) as a linker

promoted a three-fold higher induction of cAMP compared to wild-type TSH, likely because the

addition of a CTP linker increases protein stability and flexibility (39).

Thus, tethered *A. aegypti* GPA2/GPB5 was expressed in HEK 293T cells and cell lysates

and secreted protein fractions were collected for expression studies. Immunoblot studies revealed

the detection of a strong 32-40 kDa band in lanes loaded with cell lysate proteins, and two less

intense distinct bands at 40 kDa and 37 kDa in secreted protein fractions of GPA2/GPB5-

transfected cells. Molecular weight band sizes at 40 kDa and 37 kDa in secreted protein fractions

matched the expected band sizes of tethered GPA2/GPB5 heterodimers, corresponding to

nonglycosylated 13 kDa or glycosylated 16 kDa GPA2 plus 24 kDa GPB5. Moreover, treatments

with PNGase verified the tethered GPA2/GPB5 proteins are glycosylated, as observed for GPA2

expressed independantly in earlier experiments herein and in previous studies (14). Highly

intense bands in cell lysates indicate the retention of immature and mature GPA2/GPB5

heterodimers. For this reason, cell lysates and secreted protein fractions of tethered GPA2/GPB5-

expressing cells were individually tested for their ability to activate *A. aegypti* LGR1.

In humans, GPA2/GPB5-TSHR signalling stimulates adenylyl cyclase activity to increase

intracellular cAMP via interaction with a Gs protein (3, 6, 11), and these results were confirmed

in our studies. Comparatively, GPA2/GPB5 signalling was also shown to increase levels of

cAMP upon binding LGR1 in *D. melanogaster* (29). Interestingly, our experiments demonstrate

low level constitutive activity of adenylyl cyclase in LGR1 expressing cells since cAMP

luminescent response was moderately greater in LGR1-transfected cells compared to cells not

expressing LGR1. Constitutive activity of glycoprotein hormone receptors is well known and has

been demonstrated to be stronger for the thyrotropin receptor than for the LH/CG receptor (40).

Surprisingly, our experiments indicate that incubations of LGR1-expressing cells with

GPA2/GPB5 tethered heterodimers triggers a switch from low level constitutive Gs coupling to

Gi coupling for *A. aegypti* LGR1, given that GPA2/GPB5 heterodimers signficantly inhibited the

forskolin-induced increase in cAMP in LGR1-transfected cells but not in cells lacking LGR1

expression. This finding, while highly interesting, is not entirely unusual since promiscuous G

protein coupling has been reported for glycoprotein hormone receptors like the TSH receptor (Gs

and Gq) and LH/CG (Gi and Gs) (41–43).

Regulation by GPA2/GPB5 heterodimers

To help stabilize heterodimerization, the beta subunit sequences of the classic

glycoprotein hormones (FSH, LH, TSH and CG) contain two additional cysteine residues that

form an additional disulfide bridge which wraps around and "buckles" the alpha subunit (26).

Though heterodimerization can occur with mutated forms of this "seatbelt" structure, there is a

dramatic decrease in heterodimer stability (21, 22). GPB5 in vertebrates and invertebrates lack

the seatbelt structure required to stabilize heterodimerization (26). Thus, the hypothesis that

GPA2/ GPB5 functions as a heterodimer in a physiological situation (i.e. without chemical cross-

linking) is challenged. The dissociation constant (Kd) associated with heterodimerization of the

classic glycoproteins hormone subunits and GPA2/GPB5 is 10^{-7} M to 10^{-6} M, which indicates

heterodimeric interactions are favoured at these concentrations (44, 45). However, since the

classic beta subunits contain an additional disulfide bridge that strengthens its association with

the common alpha subunit, heterodimeric interactions are stabilized in circulation at

physiological concentrations as low as 10^{-11} M to 10^{-9} M (22). Without this seatbelt structure,

GPA2/GPB5 heterodimeric interactions are posssible only at micromolar concentrations, which

are not typically observed in circulation (22, 26). Together, the limited evidence so far using

heterolougous expression challenges the possibility of endocrine regulation by GPA2/GPB5

heterodimers. Nonetheless, it was argued in *D. melanogaster* that the large neurosecretory cells

co-expressing the glycoprotein hormone subunits, along with their corresponding axonal

projections that localized distinctly from organs that express the GPA2/GPB5 receptor (LGR1),

did support that this system is indeed endocrine in nature (16). Alternatively, the subunits could

function independently or regulate physiology as a heterodimer in a paracrine/ autocrine fashion.

In rats, GPA2/GPB5 is expressed in oocytes and may act as a paracrine regulator of TSHR-

expressed granulosa cells in the ovary to regulate reproductive processes (6). Another possibility

to consider is that additional endogenous co-factors may be involved, but remain unidentified,

which help to strengthen interaction between the GPA2 and GPB5 subunits, since the tethered

mosquito GPA2/GPB5 was indeed capable of activating LGR1 *in vitro* inducing a Gi signalling

cascade.

GPA2 and GPB5 homodimerization

Our results establish that human and mosquito GPA2 and GPB5 subunits can weakly and strongly, respectively, homodimerize. However, whether these homodimers, pertain to a physiological function *in vivo* is unknown. Treatments of either mosquito or human GPA2 and GPB5 subunits alone did not stimulate specific downstream signalling in LGR1 or TSHR-expressing cells, indicating that only GPA2/GPB5 heterodimers can activate their cognate glycoprotein hormone receptors. However, it is possible that GPA2 and GPB5 homodimers may target other unidentified receptors. In insects, the moulting hormone bursicon is a heterodimer of two subunits called burs and pburs. Burs/pburs heterodimers act via a glycoprotein hormone receptor (i.e. LGR2) to regulate processes such as tanning and sclerotization of the insect cuticle as well as wing inflation after adult emergence (46). Recently, it was demonstrated that bursicon subunits can homodimerize (i.e. burs/burs and pburs/pburs) and these homodimers mediate actions independently of LGR2 to regulate immune responses in *A. aegypti* and *D. melanogaster* (47, 48).

In addition to the human/ mosquito GPA2/GPB5 homodimers observed in our studies, human GPA2 was also shown to interact with the beta subunits of CG and FSH (3). Lastly, the expression patterns of GPA2 and GPB5 in a number of organisms do not always strictly co-localize, since GPA2 expression exhibits a much wider distribution and is expressed more abundantly than GPB5 in a number of vertebrate and invertebrate organisms (7, 12, 23, 24, 49, 50). Taken together, this raises the possibility that GPA2 and GPB5 subunits may interact with other unknown proteins that could activate different receptors or signalling pathways and elicit distinct functions.

Concluding remarks

Although much is known about the classic vertebrate glycoprotein hormones including

LH, FSH, TSH and CG along with their associated receptors, little progress has been made thus

far towards better understanding the function of GPA2/GPB5, signalling and subunit

interactions, particularly for the invertebrate organisms. To our knowledge, this is the first study

to demonstrate *A. aegypti* and *H. sapiens* GPA2 and GPB5 subunit homodimerization *in vitro*.

Our results also confirm that heterodimerization of *A. aegypti* and *H. sapiens* GPA2/GPB5 are

required for the activation of their cognate receptors LGR1 and TSHR, respectively. Unlike

previous reports showing GPA2/GPB5-induced LGR1 activation elevates intracellular cAMP by

coupling a Gs pathway, the current findings provide novel information supporting that *A. aegypti*

LGR1 couples to a Gi protein to inhibit cAMP levels following application of heterodimeric

GPA2/GPB5. Further, our results revealed that mosquito LGR1 is constitutively active when

overexpressed in the absence of its ligand, GPA2/GPB5, inducing a Gs signalling pathway that

raises levels of cAMP levels, which is consistent with previous observations with overexpression

of fruit fly LGR1 (29, 51) as well as mammals including dog and human TSH receptor (52, 53).

In the mosquito nervous system, GPA2 and GPB5 subunits are co-expressed within the

the same neurosecretory cells of the first five abdominal ganlgia where their coordinated release

and regulation are likely. As a result, whether GPA2/GPB5 are secreted as heterodimers, like the

classic glycoprotein hormones, and/or as homodimers remains to be determined *in vivo*;

however, these results confirm GPA2 and GPB5 homodimers do not activate LGR1 and TSHR.

Whether these homodimers are functional *in vivo* and what physiological role they play (if any)

is a research direction that should be addressed in future studies. All in all, this investigation has

provided novel information for an invertebrate GPA2/GPB5 and LGR1 signalling system and

contributes towards advancing our understanding and the functional elucidation of this ancient

glycoprotein hormone signalling system common to nearly all bilaterian organisms.

Materials and methods

Animals

Adult *Aedes aegypti* (Liverpool) were derived from an established laboratory-reared colony raised under conditions described previously (18).

GPA2/ GPB5 transcript analysis by RT-qPCR

Total RNA was isolated and purified from select *A. aegypti* tissues and organs, reverse transcribed into cDNA. GPA2 and GPB5 transcript abundance was quantified using a StepOnePlus^{TM} Real-Time PCR System (Applied Biosystems) as described previously (55) (see SI Materials and Methods for details). GPA2 and GPB5 transcript abundance was normalized to the expression of two reference genes following the ΔΔCt method as previously described (14). Experiments were repeated using a total of three technical replicates per sample and three biological replicates for each tissue/ organ.

Fluorescence in situ hybridization

Using gene-specific primers (Table S1), *A. aegypti* GPA2 and GPB5 sequences were amplified from previously prepared constructs (14) that contained the GPA2 and GPB5 complete open reading frames (ORFs). Sense and antisense probes were then generated following a similar protocol as recently reported (19). Briefly, cDNA fragments were ligated to pGEM T Easy vector (Promega, Madison, WI, USA) and used to transform NEB 5-α competent *Escherichia coli* cells (New England Biolabs, Whitby, ON, Canada). After screening plasmid constructs for directionality using T7 promoter oligonucleotide and gene-specific primers (Table S1), template

sense or anti-sense cDNA strands for GPA2 and GPB5 probe synthesis were created by PCR

amplification and verified by Sanger sequencing for base accuracy (The Centre for Applied

Genomics, Sick Kids Hospital, Toronto, ON, Canada). Digoxigenin (DIG)-labelled anti-sense

and sense RNA probes corresponding to GPA2 and GPB5 subunits were synthesized using the

HiScribe T7 High Yield RNA Synthesis kit (New England Biolabs, Whitby, ON, Canada).

Fluorescence *in situ* hybridization (FISH) was then used to detect GPA2 and/or GPB5 transcript

in the mosquito central nervous system using 4 ng μl^{-1} (GPB5) and/or 6 ng μl^{-1} (GPA2) RNA

sense/ antisense probes, following a previously established protocol (54). Preparations were

analyzed with a Lumen Dynamics X-CiteTM 120Q Nikon fluorescence microscope (Nikon,

Mississauga, ON, Canada), or a Yokogowa CSU-XI Zeiss Cell Observer Spinning Disk confocal

microscope, and images were processed using Zeiss Zen and ImageJ software. All microscope

settings were kept identical when acquiring images of control and experimental preparations.

Wholemount immunohistochemistry

GPB5 immunoreactivity in the abdominal ganglia of adult mosquitoes was examined in

newly emerged and four-day old *A. aegypti* that were lightly CO_2 anesthetized, and dissected in

PBS at RT. Tissues were fixed, permeabilized and incubated in a custom rabbit polyclonal GPB5

antibody (1 $\mu g\ ml^{-1}$) designed against an antigen sequence (CDSNEISDWRFP) positioned at

residues 69-80 (14) for 48 h at 4°C rocking. Control treatments involved pre-incubated *A.*

aegypti GPB5 primary antibody solution containing 100:1 peptide antigen:antibody (mol:mol).

After several washes, tissues were incubated overnight at 4°C in Alexa Fluor 488-conjugated

goat anti-rabbit Ab (1:200) secondary antibody (Life Technologies) in PBS containing 10%

normal sheep serum. The next day, samples were washed and then mounted onto coverslips

using mounting media containing Diamidino-2-phenylindole dihydrochloride (DAPI), and

analyzed using a Lumen Dynamics X-CiteTM 120Q Nikon fluorescence microscope (Nikon,

Mississauga, ON, Canada), or optically sectioned using a Yokogowa CSU-XI Zeiss Cell

Observer Spinning Disk confocal microscope. All images were processed using Zeiss Zen and

ImageJ software. Further details concerning the wholemount immunohistochemical protocol

were reported in an earlier study (18).

Plasmid expression constructs

Plasmid expression constructs were designed to study *A. aegypti* and *H. sapiens*

GPA2/GPB5 subunit dimerization patterns and receptor signalling. Using previously available

hexa-histidine-tagged *A. aegypti* GPA2-His (14), as well as FLAG (DYKDDDDK)-tagged (-

FLAG) *H. sapiens* GPB5-FLAG (Genscript, Clone OHu31847D) plasmid vectors as template,

the full ORF of each (*A. aegypti* and *H. sapiens*) GPA2 and GPB5 subunit coding sequence,

including a consensus Kozak translation initiation sequence, was amplified and a hexa-histidine

or FLAG tag sequence was incorporated on the carboxyl-terminus of subunits to produce the

following fusion proteins; *A. aegypti* GPA2-FLAG, *H. sapiens* GPB5-His (Table S2). A

pcDNA3.1$^+$ mammalian expression construct containing mCherry, which was a gift from Scott

Gradia (Addgene plasmid # 30125), was utilized to verify cell transfection efficiency.

Experiments also utilized previously prepared pcDNA3.1$^+$ constructs with *A. aegypti* GPB5-His

and *A. aegypti* LGR1 coding sequences and dual promoter vector pBudCE4.1 containing both *A.*

aegypti GPA2-His and GPB5-His (55). Additionally, pcDNA 3.1$^+$ mammalian expression vector

construct containing FLAG tagged *H. sapiens* thyrotropin receptor (TSHR-FLAG) (Genscript

USA Inc., Clone OHu18318D), *H. sapiens* GPA2-FLAG (Genscript USA Inc., Clone

OHu31847D), *H. sapiens* GPB5-FLAG (Genscript USA Inc., Clone OHu55827D) and

pGlosensor^{TM} -22F cyclic adenosine monophosphate (cAMP) biosensor plasmid (Promega

Corp., Madison, WI), which were used for receptor activation and intracellular signalling assays.

Generation of tethered *A. aegypti* GPA2/GPB5 construct

The ORFs of *A. aegypti* GPA2 and GPB5 sequences were tethered together in order to

promote heterodimer interactions for testing in receptor activity assays with mammalian cell

lines. A hexa-histidine tagged artificial linker sequence involving three glycine-serine repeats

was used to fuse the amino-terminus of *A. aegypti* GPA2 propeptide sequence to the carboxyl-

terminus of *A. aegypti* GPB5 prepropeptide sequence, using multiple PCR amplifications with

several primer sets (Table S2) as performed previously using lamprey GPA2 and GPB5

sequences (38) (see SI Materials and Methods for details).

Transient transfection of HEK 293T cells

Human embryonic kidney (HEK 293T) cells were grown in complete growth media

(Dulbecco's modified eagles medium: nutrient F12 (DMEM) media, 10% heat inactivated fetal

bovine serum (Wisent, St. Bruno, QC) and 1X antimycotic-antibiotic) and maintained in a water-

jacketed incubator at 37°C, 5% CO_2. When cells reached ~80-90% confluency, they were

transfected with plasmid expression constructs in 6-well tissue culture plates (Thermo Fisher

Scientific, Burlington, ON) using Lipofectamine 3000 transfection reagent (Life Technologies,

Carlsbad, CA) with 3:1 (µL:µg) transfection reagent to DNA ratio. Before transfection, culture

media was replaced with either serum-free medium (DMEM and 1X antimycotic-antibiotic) for

experiments that collected secreted proteins, or fresh complete growth medium for experiments

that dually-transfected cells with pGlosensor^TM -22F cAMP biosensor plasmid and either *H.*

sapiens TSHR, *A. aegypti* LGR1, or mCherry.

Preparation of protein samples

At 48 hours post-transfection, serum-free culture media containing secreted proteins were

collected and concentrated in 0.5 mL 3-kDa molecular weight cut-off centrifugal filters (VWR

North America). In some experiments, cells were dislodged with PBS containing 5 mM

ethylenediaminetetraacetic acid (EDTA; Life Technologies) (PBS-EDTA) pH 8.0, pelleted at

400x*g* for 5 min, resuspended in PBS and transferred to 1.5 mL centrifuge tubes for a subsequent

centrifugation. Cell lysates were then prepared by resuspending and sonicating cells for 5 s in

cell lysis buffer containing 37.5 mM Tris, pH = 7.5, 1.5 mM EDTA, pH 8.0, 3% sodium dodecyl

sulfate, 1.5% protease inhibitor cocktail (v/v), and 1.5 mM dithiothreitol (DTT). For receptor

activity assays, in order to prevent carry-over of lysis buffer, cell lysates were concentrated in 3-

kDa molecular weight cut-off centrifugal filters and re-constituted back to initial volumes with

serum-free media for a total of three repetitions. Proteins were then used for cross-linking

analysis, deglycosylation and immunoblotting or used as ligands for functional receptor

activation using the cAMP signalling biosensor assays.

Dissuccinimidyl suberate (DSS), a chemical cross-linker that is primarily reactive

towards amino groups providing stabilization of weak or transient protein intermolecular

interactions, was employed to study GPA2 and GPB5 protein-protein interactions. *A. aegypti* and

H. sapiens GPA2-FLAG and GPB5-His proteins were separately, or combined together (GPA2-

FLAG/GPB5-His) and then treated with DSS (Sigma Aldrich, Oakville, ON). In some

experiments, *A. aegypti* GPA2-His/GPB5-His were co-expressed using a dual promoter plasmid

and as such, media containing both His-tagged subunits were directly treated with DSS. To treat

secreted concentrates of culture media containing *H.* sapiens GPA2 and GPB5 proteins, 0.68 mM

DSS was used whereas both 0.68 mM (data not shown) and 2.04 mM DSS was used to test the

dimerization characteristics of *A. aegypti* GPA2-His/GPB5-FLAG proteins. Cross-linking was

performed for 30 min at RT and reactions were quenched with 50 mM Tris, pH 7.4 for 10 min

under constant mixing. To remove N-linked sugars, protein samples were treated with peptide-N-

Glycosidase F (PNGase) (New England Biolabs, Whitby, ON) following manufacturers

guidelines, with the only modification being that protein samples were not heated to 100°C

before enzymatic deglycosylation. Experiments aimed to determine the effects of protein

glycosylation on cross-linking ability first treated *A. aegypti* GPA2-His/ GPB5-His subunits with

PNGase and subsequently cross-linked samples with DSS after deglycosylation .

Western blot analysis

Samples were prepared in 2x Laemmli buffer (Sigma Aldrich, Oakville, ON) containing

4% SDS, 20% glycerol, 10% 2-mercaptoethanol, 0.004% bromophenol blue and 0.125 M Tris

HCl, pH ~6.8, and resolved on 10% or 15% SDS-polyacrylamide gels under reducing conditions

at 120 V for 90-110 min. Using a wet transfer system, proteins were then transferred to

polyvinylidene difluoride (PVDF) membranes at 100 V for 75 min. For GPA2-FLAG/GPB5-His

heterodimerization experiments, samples were run in duplicate on the same gel and following

transfer, membranes were cut in half for separate primary antibody incubations. Membrane blots

containing protein samples were blocked for 1 h in PBS containing 0.1% Tween-20 (Bioshop,

Burlington, ON, Canada) and 5% skim milk powder (PBSTB) rocking at RT. After blocking,

membranes were incubated overnight at 4°C on a rocking platform in PBSTB containing mouse

monoclonal anti-His (1:500 dilution) or mouse monoclonal anti-FLAG (1:500) primary antibody

solutions. The next day, membranes were washed three times with PBS containing 0.1% Tween-

20 (PBST) for 15 min each wash and then were incubated in PBSTB containing anti-mouse HRP

conjugated secondary antibody (1:2000 – 1:3000 dilution) for 1 h rocking at RT before washing

again with PBST (3 x 15 min washes). Finally, blots were incubated with Clarity Western ECL

substrate and images were developed using a ChemiDoc MP Imaging System (Bio-Rad

Laboratories, Mississauga, ON) and molecular weight measurements and analysis were

performed using Image Lab 5.0 software (Bio-Rad Laboratories, Mississauga, ON).

Receptor Functional Activation Bioluminescence Assays

HEK 293T cells were co-transfected to express (i) *H. sapiens* TSHR, *A. aegypti* LGR1, or

mCherry along with (ii) pGlosensor^{TM} -22F cAMP biosensor plasmid (Promega Corp., Madison,

WI), which encodes a modified form of firefly luciferase with a fused cAMP binding moiety

providing a biosensor for the direct detection of cAMP signalling in live cells. At 48 hours post

transfection, recombinant cells were dislodged with PBS-EDTA, pelleted at 400x*g* for 5 min,

resuspended in assay media (DMEM:F12 media with 10% fetal bovine serum (v/v)) containing

cAMP GloSensor reagent (2% v/v), and incubated for 3 h rocking at RT shielded from light.

White 96-well luminescence plates (Greiner Bio-One, Germany) were loaded under low light

with previously prepared secreted or cell lysate protein concentrates (described above), forskolin

or assay media alone, and incubated at 37°C for 30 min prior to performing receptor activity

assays. For stimulatory G-protein (Gs) pathway detection, recombinant cells expressing either *H.*

sapiens TSHR, *A. aegypti* LGR1, or mCherry along with the cAMP biosensor were pre-treated

with 0.25 mM 3-isobutyl-1-methylxanthine (IBMX) for 30 min at RT with rocking and shielded

from light. Using an automatic injector unit (BioTek Instruments Inc., Winooski VT),

recombinant cells were then loaded into wells containing various treatments, including 250 nM

forskolin as a positive control or concentrates of protein fractions from mCherry-transfected cells

as a negative control. For inhibitory G-protein (Gi/o) pathway testing, *A. aegypti* GPA2 and/or

GPB5 was tested for the ability to inhibit a forskolin-induced cAMP-mediated bioluminescent

response. Using an automatic injector unit (BioTek Instruments Inc., Winooski VT), cells

expressing LGR1 or mCherry (LGR1 activation and signalling negative control) were loaded

into wells that contained various ligand treatments. Subsequently, 250 nM or 1 µM forskolin was

added to wells immediately after the addition of recombinant cells, using a second automatic

injector unit. For bioluminescent assays with tethered GPA2/GPB5 proteins, after the 3 h

incubation, recombinant cells expressing or not expressing LGR1 were equally divided to test Gs

and Gi signalling pathways simultaneously using the same batch of cells per biological replicate.

In all assays, luminescence was measured every 2 min for 20 min at 37°C using a Synergy 2

Multimode Microplate Reader (BioTek, Winooski, VT) and averaged over 4-8 technical

replicates for each treatment. To calculate the relative luminescent response, data was

normalized to luminescent values recorded from negative control ligand treatments involving

protein secretions derived from mCherry-transfected cells. Assays were performed repeatedly

and involved 3-6 independent biological replicates.

Acknowledgements

Research in this study was supported by a Natural Sciences and Engineering Research Council of

Canada (NSERC) Discovery Grant and Ontario Ministry of Research & Innovation Early

Researcher Award to JPP.

References

1. Pierce JG, Parsons TF (1981) Glycoprotein hormones: structure and function. *Annu Rev Biochem* 50:465–495.
2. Stockell Hartree A, Renwick AGCC (1992) Molecular structures of glycoprotein hormones and functions of their carbohydrate components. *Biochem J* 287(3):665–679.
3. Nakabayashi K, et al. (2002) Thyrostimulin, a heterodimer of two new human glycoprotein hormone subunits, activates the thyroid-stimulating hormone receptor. *J Clin Invest* 109(11):1445–1452.
4. Hsu SY (2002) Evolution of Glycoprotein Hormone Subunit Genes in Bilateral Metazoa: Identification of Two Novel Human Glycoprotein Hormone Subunit Family Genes, GPA2 and GPB5. *Mol Endocrinol* 16(7):1538–1551.
5. Santos S, Mazan S, Venkatesh B, Cohen-Tannoudji J, Quérat B (2011) Emergence and evolution of the glycoprotein hormone and neurotrophin gene families in vertebrates. *BMC Evol Biol* 11(1):332.
6. Sun SC, et al. (2010) Thyrostimulin, but not thyroid-stimulating hormone (TSH), acts as a paracrine regulator to activate the TSH receptor in mammalian ovary. *J Biol Chem* 285(6):3758–3765.
7. Okada SL, et al. (2006) A glycoprotein hormone expressed in corticotrophs exhibits unique binding properties on thyroid-stimulating hormone receptor. *Mol Endocrinol* 20(2):414–425.
8. Duncan Bassett JH, et al. (2015) Thyrostimulin Regulates Osteoblastic Bone Formation During Early Skeletal Development. *Endocrinology* (August):en.2014-1943.
9. van Zeijl CJJ, et al. (2014) Thyrostimulin deficiency does not alter peripheral responses to acute inflammation-induced nonthyroidal illness. *Am J Physiol Endocrinol Metab* 307(6):E527-37.
10. Suzuki C, et al. (2009) Inflammatory cytokines regulate glycoprotein subunit beta5 of thyrostimulin through nuclear factor-kappaB. *Endocrinology* 150(5):2237–43.
11. Huang W, et al. (2016) Thyrostimulin-TSHR signaling promotes the proliferation of NIH□: OVCAR-3 ovarian cancer cells via trans-regulation of the EGFR pathway. *Nat Publ Gr* (June):1–13.
12. Heyland A, et al. (2012) Distinct expression patterns of glycoprotein hormone subunits in the lophotrochozoan Aplysia: Implications for the evolution of neuroendocrine systems in animals. *Endocrinology* 153(11):5440–5451.
13. Vandersmissen HP, Van Hiel MB, Van Loy T, Vleugels R, Vanden Broeck J (2014) Silencing D. melanogaster lgr1 impairs transition from larval to pupal stage. *Gen Comp Endocrinol* 209:135–147.
14. Paluzzi J-PJPJ-P, Vanderveken M, O'Donnell MJMJ (2014) The heterodimeric glycoprotein hormone, GPA2/GPB5, regulates ion transport across the hindgut of the adult mosquito, Aedes aegypti. *PLoS One* 9(1):e86386.
15. Rocco DA, Paluzzi J-PV (2016) Functional role of the heterodimeric glycoprotein hormone, GPA2/GPB5, and its receptor, LGR1: An invertebrate perspective. *Gen Comp Endocrinol* 234. doi:10.1016/j.ygcen.2015.12.011.
16. Sellami A, Agricola HJ, Veenstra J a. (2011) Neuroendocrine cells in Drosophila melanogaster producing GPA2/GPB5, a hormone with homology to LH, FSH and TSH. *Gen Comp Endocrinol* 170(3):582–588.

17.   Jourjine N, Mullaney BC, Mann K, Scott K (2016) Coupled Sensing of Hunger and Thirst
Signals Balances Sugar and Water Consumption. *Cell* 166(4):855–866.
18.   Rocco DA, Kim DH, Paluzzi J-P V. (2017) Immunohistochemical mapping and transcript
expression of the GPA2 / GPB5 receptor in tissues of the adult mosquito , Aedes aegypti.
Cell Tissue Res 369(2):313–330.
19.   Rocco DA, et al. (2019) Glycoprotein Hormone Receptor Knockdown Leads to Reduced
Reproductive Success in Male Aedes aegypti. *Front Physiol* 10(March):1–11.
20.   Combarnous Y (1992) Molecular basis of the specificity of binding of glycoprotein
hormones to their receptors. *Endocr Rev* 13(4):670–91.
21.   Xing Y, et al. (2004) Glycoprotein hormone assembly in the endoplasmic reticulum: II.
Multiple roles of a redox sensitive (beta)-subunit disulfide switch. *J Biol Chem*
279(34):35437–35448.
22.   Galet C, Lecompte F, Combarnous Y (2004) Association/dissociation of gonadotropin
subunits involves disulfide bridge disruption which is influenced by carbohydrate moiety.
Biochem Biophys Res Commun 324(2):868–873.
23.   Nagasaki H, et al. (2006) Differential expression of the thyrostimulin subunits,
glycoprotein α2 and β5 in the rat pituitary. *J Mol Endocrinol* 37(1):39–50.
24.   Dos Santos S, et al. (2009) Distinct expression patterns of glycoprotein hormone-alpha2
and -beta5 in a basal chordate suggest independent developmental functions.
Endocrinology 150(8):3815–3822.
25.   Tando Y, Kubokawa K (2009) A homolog of the vertebrate thyrostimulin glycoprotein
hormone alpha subunit (GPA2) is expressed in Amphioxus neurons. *Zoolog Sci*
26(6):409–414.
26.   Alvarez E, Cahoreau C, Combarnous Y (2009) Comparative structure analyses of cystine
knot-containing molecules with eight aminoacyl ring including glycoprotein hormones
(GPH) alpha and beta subunits and GPH-related A2 (GPA2) and B5 (GPB5) molecules.
Reprod Biol Endocrinol 7(1):90.
27.   Krause G, Kreuchwig A, Kleinau G (2012) Extended and Structurally Supported Insights
into Extracellular Hormone Binding, Signal Transduction and Organization of the
Thyrotropin Receptor. *PLoS One* 7(12). doi:10.1371/journal.pone.0052920.
28.   Hauser F, Nothacker HP, Grimmelikhuijzen CJ (1997) Molecular cloning, genomic
organization, and developmental regulation of a novel receptor from Drosophila
melanogaster structurally related to members of the thyroid-stimulating hormone, follicle-
stimulating hormone, luteinizing hormone/choriogonadotropin. *J Biol Chem* 272(2):1002–
10.
29.   Sudo S, Kuwabara Y, Park J Il, Sheau YH, Hsueh AJW (2005) Heterodimeric fly
glycoprotein hormone-α2 (GPA2) and glycoprotein hormone-β5 (GPB5) activate fly
leucine-rich repeat-containing G protein-coupled receptor-1 (DLGR1) and stimulation of
human thyrotropin receptors by chimeric fly GPA2 and human GPB5. *Endocrinology*
146(8):3596–3604.
30.   Sgourakis NG, Bagos PG, Papasaikas PK, Hamodrakas SJ (2005) A method for the
prediction of GPCRs coupling specificity to G-proteins using refined profile Hidden
Markov Models. 12:1–12.
31.   Brownstein M (1985) The biochemistry of the polypeptide hormones. *Cell* 48(2):147–183.
32.   Szkudlinski MW, Fremont V, Ronin C, Weintraub BD (2002) Thyroid-stimulating
hormone and thyroid-stimulating hormone receptor structure-function relationships.

Physiol Rev 82(2):473–502.

33. Atwood B, Lopez J, Wager-Miller J, Mackie K, Straiker A (2011) Expression of G
protein-coupled receptors and related proteins in HEK293, AtT20, BV2, and N18 cell
lines as revealed by microarray analysis. *BMC Genomics*. doi:10.1186/1471-2164-12-14.

34. Priyanka V, Chichili R, Kumar V, Sivaraman J (2013) Linkers in the structural biology of
protein – protein interactions. 22:153–167.

35. Fares F, et al. (1992) Design of a long-acting follitropin agonist by fusing the C-terminal
sequence of the chorionic gonadotropin β subunit to the follitropin β subunit. *Proc Natl*
Acad Sci USA 89:4304–4308.

36. Joshi L, et al. (1995) Recombinant thyrotropin containing a β-subunit chimera with the
human chorionic gonadotropin-β carboxy terminal is biologically active with a prolonged
plasma half-life: role of carbohydrate in bioactivity and metabolic clearance.
Endocrinology 136:3839–3948.

37. Furuhashi M, et al. (1995) Fusing the carboxy-terminal peptide of the chorionic
gonadotropin (CG) β-subunit to the common α-subunit: retention of O-linked
glycosylation and enhanced in vivo bioactivity of chimeric human CG. *Mol Endocrinol*
9:54–63.

38. Sower SA, et al. (2015) Emergence of an Ancestral Glycoprotein Hormone in the Pituitary
of the Sea Lamprey, a Basal Vertebrate. *Endocrinology* 156:3026–3037.

39. Azzam N, Bar-Shalom R, Fares F (2012) Conversion of TSH Heterodimer to a Single
Polypeptide Chain Increases Bioactivity and Longevity. 153(2):954–960.

40. Cetani F, Tonacchera M, Vassart G (1996) Differential effects of NaCl concentration on
the constitutive activity of the thyrotropin and the luteinizing hormone/chorionic
gonadotropin receptors. *FEBS Lett* 378:27–31.

41. Herrlich A, et al. (1996) Involvement of Gs and Gi Proteins in Dual Coupling of the
Luteinizing Hormone Receptor to Adenylyl Cyclase and Phospholipase C. *J Biol Chem*
271(28):16764–16772.

42. Grasberger H, Van Sande J, Hag-Dahood M, Tenenbaum-Rakover Y, S R (2007) A
familial thyrotropin (TSH) receptor mutation provides in vivo evidence that the inositol
phosphates/Ca2+ cascade mediates TSH action on thyroid hormone synthesis. *J Clin*
Endocrinol Metab 92:2816–2820.

43. Kleinau G, et al. (2010) Principles and Determinants of G-Protein Coupling by the
Rhodopsin-Like Thyrotropin Receptor. *PLoS One* 5(3):e9745.

44. Strickland TW, Puett D (1982) The kinetic and equilibrium parameters of subunit
association and gonadotropin dissociation. *J Biol Chem* 257(6):2954–2960.

45. Ingham K, Weintraub B, Edelhock H (1976) Kinetics of recombination of the subunits of
human chorionic gonadotropin. Effect of subunit concentration. *Biochemistry* 15(8):1720–
1726.

46. Luo C-W, et al. (2005) Bursicon, the insect cuticle-hardening hormone, is a heterodimeric
cystine knot protein that activates G protein-coupled receptor LGR2. *Proc Natl Acad Sci*
U S A 102(8):2820–2825.

47. An S, et al. (2012) Insect neuropeptide bursicon homodimers induce innate immune and
stress genes during molting by activating the NF-κB transcription factor relish. *PLoS One*
7(3):1–9.

48. Zhang H, Dong S, Xi C, Stanley D, Beerntsen B (2017) Relish2 mediates bursicon
homodimer-induced prophylactic immunity in the mosquito Aedes aegypti. (January):1–9.

49.   Tando Y, Kubokawa K (2009) Expression of the gene for ancestral glycoprotein hormone
846   β subunit in the nerve cord of amphioxus. *Gen Comp Endocrinol* 162(3):329–339.
847   50.   Trudeau VL (2009) Really old hormones up to new tricks: glycoprotein hormone subunits
848   may have roles in development. *Endocrinology* 150(8):3446–7.
849   51.   Gondalia K, Qudrat A, Bruno B, Fleites Medina J, Paluzzi JP V. (2016) Identification and
850   functional characterization of a pyrokinin neuropeptide receptor in the Lyme disease
851   vector, Ixodes scapularis. *Peptides* 86:42–54.
852
853

Figure Legends

Fig. 1. GPA2/GPB5 subunit transcript expression and localization in adult *A. aegypti*. (*A*, *B*) RT-

qPCR examining GPA2 (*A*) and GPB5 (*B*) transcript expression in the central nervous system of

adult mosquitoes, with significant enrichment in the abdominal ganglia (AG). Subunit transcript

abundance is shown relative to their expression in the thoracic ganglia (TG). Mean ± SEM of

three biological replicates. Columns denoted with different letters are significantly different from

one another. Multiple comparisons two-way ANOVA test with Tukey's multiple comparisons

(P<0.05) to determine sex- and tissue-specific differences. Fluorescence *in situ* hybridization

anti-sense (*C*, *D*, *G-J*) and sense (*E, F*) probes to determine GPA2 and GPB5 transcript

localization (GPA2 and/or GPB5 transcript, red; nuclei, blue). Unlike sense probe controls (*E*,

F), two bilateral pairs of cells (arrowheads) were detected with GPA2 (*C*, *G*) and GPB5 (*D*, *H*)

anti-sense probes in the first five abdominal ganglia. Co-localization of the GPA2 (*G*) and GPB5

(*H*) transcript was verified by treating abdominal ganglia dually with a combination of GPA2

and GPB5 anti-sense probe (*I, J*) that revealed two, intensely-stained bilateral pairs of cells. In *C-*

F and *I-J*, microscope settings were kept identical when acquiring images of control and

experimental ganglia. Scale bars are 50 µm in *C-F*, *I-J* and 40 µm in *G* and *H*.

Fig. 2. Immunolocalization of GPB5 subunit expression in the abdominal ganglia of adult *A.*

aegypti. Experimental treatments demonstrated GPB5 immunoreactivity (green) in two (*A*), and

in some mosquitoes, three (*B*) bilateral pairs of neuroendocrine cells (arrowheads) within the first

five abdominal ganglia of the ventral nerve cord (DAPI, blue). Optical sections of ganglia

revealed axonal projections emanate from GPB5-immunoreactive cells through the lateral nerves

(arrows). No cells were detected in the sixth, terminal ganglion (*C*) and in control treatments

(CON) where GPB5 antibody was preabsorbed with GPB5 synthetic antigen (*D*). In *A*, *D* or *B-C*,

microscope settings were kept identical when acquiring images of control and experimental

ganglia. Scale bars are 25 µm in *A-D* and 20 µm in *B-C*.

Fig. 3. Western blot analyses to determine the effects of glycosylation on homo- and heterodimer

formation on the glycoprotein hormone (GPA2/GPB5) subunits in the mosquito, *A. aegypti*. (*A*)

In untreated conditions, western blot analysis of GPA2 subunit alone reveals two bands at 16 and

13 kDa. Whereas following treatment with PNGase, the higher molecular weight band

disappears and the 13 kDa band is intensified. A thick, additional band at ~32 kDa appears when

GPA2 protein is cross-linked with DSS, and this band migrates slightly lower to ~30 kDa when

GPA2 protein is treated with both DSS and PNGase. (*B*) A 24 kDa band is observed in lanes

loaded with untreated GPB5 subunit alone. Upon PNGase treatment, the 24 kDa band is not

affected; however, upon treatment with DSS, a second faint band appears at 48 kDa that is not

affected by deglycosylation. (*C*) Western blot analyses of co-expressed GPA2 and GPB5

subunits shows three distinct band sizes at 24 kDa, 16 kDa and 13 kDa, corresponding to the

GPB5 subunit and two forms of GPA2 subunit protein. Similar to (*A*), after treatment with

PNGase, the higher molecular weight form of GPA2 is eliminated and the 13 kDa band

intensifies. When GPA2/GPB5 protein is cross-linked, two additional bands are detected at ~48

kDa and ~ 32 kDa; however, following cross-linking and PNGase treatment, the ~32 kDa band is

eliminated leaving only the 30 kDa band along with the unaffected ~48 kDa band.

Fig. 4. Elucidating heterodimerization of *H. sapiens* (human) (*A*) and *A. aegypti* (mosquito) (*B*)

GPA2 and GPB5 subunits. Single promoter expression constructs for human and mosquito

GPA2-FLAG and GPB5-His were used for transient expression in HEK 293T cells. Protein was

harvested and subsequently concentrated, treated with DSS cross-linker, and probed with an anti-

His or an anti-FLAG antibody after SDS-PAGE. (*A*, *B*) No bands were detected in lanes loaded

with cross-linked GPA2-FLAG protein (Lane 1) or cross-linked GPB5-His protein (Lane 6),

probed with an anti-His antibody or anti-FLAG antibody, respectively. (*A*) Bands corresponding

to the monomeric form (18 kDa) and homodimer (36 kDa) of cross-linked human GPB5-His

(Lane 3) and to the monomeric form (20 kDa) and homodimer (40 kDa) of cross-linked human

GPA2-FLAG protein (Lane 4). A combination of the subunits with subsequent cross-linking of

separately-produced human GPA2-FLAG and GPB5-His protein (Lane 2, 5) revealed a band size

correlating to the human GPA2/ GPB5 heterodimer (38 kDa), detected using an anti-His (Lane

2) or anti-FLAG (Lane 5) antibody. (*B*) Bands corresponding to the mosquito GPB5 monomer

(24 kDa) (Lane 3), mosquito GPA2 glycosylated monomer (16 kDa) and homodimer pairs (30

kDa and 32 kDa) (Lane 4). No detection of bands correlating to mosquito GPA2/GPB5

heterodimer (37-40 kDa) were observed, when probed with either anti-His (Lane 2) or anti-

FLAG (Lane 5) antibodies.

Fig. 5. cAMP-mediated bioluminescence assays to determine the effect of GPA2, GPB5 and

GPA2/ GPB5 on G-protein signalling of *H. sapiens* (human) TSHR (*A*), *A. aegypti* (mosquito)

LGR1 (*B*, *C*), or cells expressing a red fluorescent protein, mCherry (*D*). Secreted protein

fractions for each subunit were prepared separately from HEK 293T cells expressing human

GPA2 (hA2), human GPB5 (hB5), mosquito GPA2 (A2), mosquito GPB5 (B5), mCherry, or co-

expressing mosquito GPA2 and GPB5 in a dual promoter plasmid (A2/B5 coexp). Secreted

protein fractions were then tested separately or combined (A2 + B5) and then incubated with

cells co-expressing the cAMP biosensor along with either (*A*) human TSHR, (*B*, *C*) mosquito

LGR1 or (*D*) mCherry, the latter of which was used as a negative control in the functional assay

and also served to verify transfection efficiency of HEK 293T cells. Luminescent values were

recorded and normalized to incubations with protein secretions collected from the media of

mCherry-transfected cells. Forskolin (FSK, 250 nM) was used as a positive control for

stimulatory G-protein (Gs) pathway (*A*, *B*) and inhibitory G-protein (Gi/o) pathway testing (*C*,

D). (*A*) Unlike treatments with human GPA2 (hA2) or human GPB5 (hB5) applied singly, a

significant increase in cAMP-mediated luminescence was observed when TSHR-expressing cells

were incubated with culture media containing both human GPA2 and human GPB5 (hA2 +

hB5), relative to incubations with mCherry controls. (*B*) No differences in luminescence were

observed when LGR1-expressing cells were incubated with media containing mosquito

GPA2/GPB5 subunits, compared to mCherry controls. (*C*) The addition of GPA2 and GPB5 on

LGR1-expressing cells significantly inhibited FSK-induced luminescent response, compared to

treatments with mCherry controls; (*D*) however, this inhibition was also observed when GPA2

and GPB5 proteins were incubated with HEK 293T cells in the absence of LGR1. Mean ± SEM

of three (*A*, *B*, *D*) or six (*C*) biological replicates. Columns denoted with different letters are

significantly different from one another. Multiple comparisons one-way ANOVA test with

Tukey's multiple comparisons (P<0.05).

Fig. 6. Western blot analysis and verification of *A. aegypti* GPA2/GPB5 tethered protein

expressed in HEK 293T cells. (*A*) Western blot analysis of secreted or cell lysate protein

fractions of HEK 293T expressing tethered GPA2/GPB5 (tA2/B5) or red fluorescent protein

(mCherry) as a control. Tethered GPA2/GPB5 is represented as a strong 32-40 kDa band in cell

lysate fractions, and as two less intense 37 kDa and 40 kDa bands in secreted fractions; however,

no bands were detected in lanes loaded with proteins from mCherry transfected cells. (*B*) Upon

treatment of tethered GPA2/GPB5 secreted protein fractions with PNGase, the 40 kDa band is

eliminated and the 37 kDa band intensifies, indicating removal of N-linked oligosaccharides.

Fig. 7. cAMP-mediated bioluminescence assay determining the effects of tethered GPA2/GPB5

on receptor activation and G protein signalling of LGR1. Secreted protein fractions (*A*, *C*) and

cell lysates (*B*, *D*) derived from cells transiently expressing tethered GPA2/GPB5 (tA2/B5) or

red fluorescent protein (mCherry) were tested for their ability to stimulate (Gs signalling) (*A*, *B*)

or inhibit 1 µM forskolin-induced (Gi/o signalling) (*C*, *D*) cAMP-mediated luminescence. (*A-D*)

Luminescence response was recorded and normalized to treatments involving mCherry proteins

with LGR1-expressing HEK 293T cells. In all treatments, the relative luminescence response

was greater in LGR1-expressing cells (+ LGR1) compared to cells not expressing LGR1 (-

LGR1). (*A*, *B*) Applications of 1 µM forskolin (FSK) to recombinant cells expressing and not

expressing LGR1 significantly increased cAMP-mediated luminescence relative to treatments

with tA2/B5 or mCherry. However, incubation of LGR1-expressing cells with tA2/B5 secreted

(*A*) or cell lysate (*B*) proteins, failed to increase cAMP-mediated luminescence above

background levels from incubations with mCherry proteins. (*C*, *D*) 1 µM forskolin along with

either mCherry proteins, tA2/B5 proteins or assay media (Assay) was added to cells in the

presence or absence of LGR1 expression. The ability for each ligand treatment to reduce a

forskolin-induced increase in cAMP luminescence was examined. (*C*) The tA2/B5 secreted

protein samples incubated with LGR1-expressing cells did not significantly affect the forskolin-

induced cAMP luminescence, compared to control treatments with secreted proteins from

mCherry expressing cells. (*D*) Relative to incubations with mCherry cell lysate proteins, cell

lysates containing tA2/B5 protein significantly inhibited forskolin-induced elevations of cAMP-

mediated luminescence, and this inhibition was specific to LGR1-expressing cells. Mean ± SEM

of three biological replicates. Columns denoted with different letters are significantly different

from one another. Multiple comparisons two-way ANOVA test with Tukey's multiple

comparisons (P<0.05).

978   **Fig. 1**

980 **Fig. 2**

Fig. 3

986 **Fig. 4**

988   **Fig. 5**

991 **Fig. 6**

994   **Fig. 7**