STARRPeaker: Uniform processing and accurate identification of

STARR-seq active regions

Donghoon Lee[1,2], Manman Shi[3], Jennifer Moran[3], Martha Wall[3], Jing Zhang[1,2], Jason

Liu[2], Dominic Fitzgerald[3], Yasuhiro Kyono[3], Lijia Ma[3,4], Kevin P White[3,5]*, Mark

Gerstein[1,2,6,7]*

[1] Program in Computational Biology and Bioinformatics, Yale University, New Haven,

CT 06520, USA

[2] Department of Molecular Biophysics and Biochemistry, Yale University, New Haven,

CT 06520, USA

[3] Institute for Genomics and System Biology, University of Chicago, Chicago, IL 60637,

USA

[4] School of Life Sciences, Westlake University, Hangzhou, Zhejiang 310024, China

[5] Tempus Labs, Inc., Chicago, IL 60654, USA

[6] Department of Computer Science, Yale University, New Haven, CT 06520, USA

[7] Department of Statistics and Data Science, Yale University, New Haven, CT 06520,

USA

* Corresponding authors

E-mail: pi@gersteinlab.org

<u>**Abstract**</u>

Background: High-throughput reporter assays, such as self-transcribing active

regulatory region sequencing (STARR-seq), allow for unbiased and quantitative

assessment of enhancers at a genome-wide scale. Recent advances in STARR-seq

technology have employed progressively more complex genomic libraries and

increased sequencing depths, to assay larger sized regions, up to the entire human

genome. These advances necessitate a reliable processing pipeline and peak-calling

algorithm.

Results: Most STARR-seq studies have relied on chromatin immunoprecipitation

sequencing (ChIP-seq) processing pipelines. However, there are key differences in

STARR-seq versus ChIP-seq. First, STARR-seq uses transcribed RNA to measure the

activity of an enhancer, making an accurate determination of the basal transcription rate

important. Second, STARR-seq coverage is highly non-uniform, overdispersed, and

often confounded by sequencing biases, such as GC content and mappability. Lastly,

here, we observed a clear correlation between RNA thermodynamic stability and

STARR-seq readout, suggesting that STARR-seq may be sensitive to RNA secondary

structure and stability. Considering these findings, we developed a negative-binomial

regression framework for uniformly processing STARR-seq data, called STARRPeaker.

In support of this, we generated whole-genome STARR-seq data from the HepG2 and

K562 human cell lines and applied STARRPeaker to call enhancers.

Conclusions: We show STARRPeaker can unbiasedly detect active enhancers from

both captured and whole-genome STARR-seq data. Specifically, we report ~33,000 and

~20,000 candidate enhancers from HepG2 and K562, respectively. Moreover, we show

that STARRPeaker outperforms other peak callers in terms of identifying known

enhancers with fewer false positives. Overall, we demonstrate an optimized processing

framework for STARR-seq experiments can identify putative enhancers while

addressing potential confounders.

Keywords: STARR-seq, peak caller, enhancer, non-coding, regulatory element

52 **Background**

The transcription of eukaryotic genes is precisely coordinated by an interplay between

cis-regulatory elements. For example, enhancers and promoters serve as binding

platforms for transcription factors (TFs) and allow them to interact with each other via

three-dimensional looping of chromatin. Their interactions are often required to initiate

transcription [1,2]. Enhancers, which are often distant from the transcribed gene body

itself, play critical roles in the upregulation of gene transcription. Enhancers are cell-type

specific and can be epigenetically activated or silenced to modulate transcriptional

dynamics over the course of development. Enhancers can be found upstream or

downstream of genes, or even within introns [3–5]. They function independent of their

orientation, do not necessarily regulate the closest genes, and sometimes regulate

multiple genes at once [6,7]. In addition, several recent studies have demonstrated that

some promoters – termed E-promoters – may act as enhancers of distal genes [8,9].

Consensus sequences (or canonical sequences) have been identified at certain protein

binding sites, splice sites, and boundaries of protein-coding genes. However, there are

no known consensus sequences that characterize enhancer function, making it

challenging to identify enhancers based on sequence alone in an unbiased fashion. The

non-coding territory occupies over 98% of the genome landscape, making the search

space very broad. Moreover, the activity of enhancers depends on the physiological

condition and epigenetic landscape of the cellular environment, complicating a fair

assessment of enhancer function.

Previously, putative regulatory elements were computationally predicted, indirectly, by

profiling DNA accessibility (using DNase-seq, FAIRE-seq, or ATAC-seq) as well as

histone modifications (ChIP-seq) that are linked to regulatory functions [10–12]. More

recently, researchers have developed high-throughput episomal (exogenous) reporter

assays to directly measure enhancer activity across the whole genome, specifically

massively parallel reporter assays (MPRA) [13,14] and self-transcribing active

regulatory region sequencing (STARR-seq) [15,16]. These assays allow for quantitative

assessment of enhancer activity in a high-throughput fashion.

In STARR-seq, candidate DNA fragments are cloned downstream of a reporter gene

into the 3′ untranslated region (UTR). After transfecting the plasmid pool into host cells,

one can measure the regulatory potential by high-throughput sequencing of the 3′ UTR

of the expressed reporter gene mRNA. These exogenous reporters enable accurate

and unbiased assessment of enhancer activity at the whole-genome level, independent

of chromatin context. Unlike MPRA – which utilizes barcodes – STARR-seq produces

self-transcribed RNA fragments that can be directly mapped onto the genome (we call

this STARR-seq output hereafter). The activities of enhancers are measured by

comparing the amount of RNA produced from the relative amount of genomic DNA in

the STARR-seq library (we call this STARR-seq input hereafter). STARR-seq has

several technical advantages over MPRA. Library construction is relatively simple

because barcodes are not needed. In addition, candidate enhancers are cloned instead

of synthesized, allowing the assay to test extended sequence contexts (>500 bp) for

enhancer activity, which studies have shown to be critical for functional activity [17].

Importantly, STARR-seq can be scaled to the whole-genome level for unbiased

scanning of functional activities. However, scaling STARR-seq to the human genome is

still very challenging, primarily due to its massive size. A more complex genomic DNA

library, a higher sequencing depth, and increased transfection efficiency are required to

cover the whole human genome [16], which could ultimately introduce biases.

Furthermore, inserting a large fragment of DNA into the 3' UTR of the reporter gene

could inadvertently introduce regulatory sequences that might affect mRNA abundance

and stability, which could lead to both false positives and false negatives. MPRA is

more robust in this regard because the activity of each candidate enhancer is quantified

by multiple molecular barcodes associated with the fragment, making it less prone to

such artifacts than STARR-seq.

The processing of STARR-seq data is somewhat similar to that of ChIP-seq, where

protein-crosslinked DNA is immunoprecipitated and sequenced. A typical ChIP-seq

processing pipeline identifies genomic regions over-represented by sequencing tags in

a ChIP sample compared to a control sample. STARR-seq data is compatible with most

ChIP-seq peak callers. Hence, previous studies on STARR-seq have largely relied on

peak-calling software developed for ChIP-seq such as MACS2 [16,18,19]. However,

one must be cautious using ChIP-seq peak callers, at least without re-tuning the default

parameters optimized for processing TF ChIP-seq [20].

In this paper, we describe key differences in the processing of STARR-seq versus

ChIP-seq data. Due to increased complexity of the genomic screening library and

sequencing depth requirements, STARR-seq coverage is highly non-uniform. This leads

to a lower signal-to-noise ratio than a typical ChIP-seq experiment and makes

estimating the background model more challenging, which could ultimately lead to false-

positive peaks. In addition, STARR-seq measures more of a continuous activity, similar

to quantification in RNA-seq, than a discrete binding event. Therefore, STARR-seq

peaks should be further evaluated using a notion of activity score. These differences

necessitate a unique approach to processing STARR-seq data.

We propose an algorithm optimized for processing and identifying functionally active

enhancers from STARR-seq data, which we call STARRPeaker. This approach

statistically models the basal level of transcription, accounting for potential confounding

factors, and accurately identifies reproducible enhancers. We applied our method to two

whole human STARR-seq datasets and evaluated its performance against previous

methods. We also compared an R package, BasicSTARRseq, developed to process

peaks from the first STARR-seq data [15], which models enrichment of sequencing

reads using a binomial distribution. We benchmarked our peak calls against known

human enhancers. Thus, our findings support that STARRPeaker will be a useful tool

for uniformly processing STARR-seq data.

Results and Discussion

Precise measurement of STARR-seq coverage

We binned the genome using a sliding window of length, *l*, and step size, *s*. Based on

the average size of the STARR-seq library, we defined a 500 bp window length with a

100 bp step size to be the default parameter. Based on the generated genomic bins, we

calculated the coverage of both STARR-seq input and output mapped to each bin. For

calculating the sequence coverage, other peak callers and many visualization tools

commonly use the start position of the read [15,21,22]. However, given that the average

size of the fragments inserted into the STARR-seq libraries were approximately 500 bp,

we expected that the read coverage using the read start position may shift the estimate

of the summit of signal and dilute the enrichment. Some peak callers have used read

densities of forward and reverse strands separately to overcome this issue [23,24]. To

precisely measure the coverage of STARR-seq input and output, we first inferred the

size of the fragment insert from paired-end reads and used the center of the fragment

insert, instead of start position of the read, to calculate coverage. For inferring the size

of the fragment insert, we first strictly filtered out reads that were not properly paired and

chimeric. Chimeric alignments are reads that cannot be linearly aligned to a reference

genome, implying a potential discrepancy between the sequenced genome and the

reference genome and indicative of a structural variation or a PCR artifact [25]. We also

filtered out read pairs that had a fragment insert size greater than l_{max} and less than

l_{min}. By default, we filtered out fragment insert sizes less than 200 bp and greater than

1,000 bp. After filtering out spurious read-pairs, we estimated the center of the fragment

insert and counted the fragment depth for each genomic bin. To assess the benefit of

using fragment-based coverage, we compared the coverage calculated using the center

of fragment insert to an alternate model using the start position of the sequencing read.

We found that the position of the peaks shifted up approximately 200 bp when we used

the alternate model (Figure 1A, Supplementary Figure 1A). Such a shift caused by the

read-based coverage could lead to the omission of TF binding sites located at the

boundary. Moreover, we observed that the read-based coverage diluted the overall

STARR-seq signal; as a result, peaks calculated based on the alternate model had

lower fold enrichment and were less confident and broader in size (Figure 1B-D,

Supplementary Figure 1B-D). Overall, the fragment-based coverage offered more

concentrated and robust peak signal compared to the read-based coverage counting

scheme. The benefit of using the center of the fragment is highlighted in Figure 1E,

where we find more concise and precise peak with a higher fold enrichment using

fragment-based coverage.

Controlling for potential systemic bias in the STARR-seq assay

To unbiasedly test for the regulatory activity, a model needs to control for potential

systemic biases inherent to generating STARR-seq data. STARR-seq measures the

ratio of transcribed RNA to DNA for a given test region and determines whether the test

region can facilitate transcription at a higher rate than the basal level. This is based on

the assumption that (1) the basal transcriptional level stays relatively constant across

the genome and (2) the transcriptional rate is a reflection of the regulatory activity of the

DNA insert. However, these assumptions may not always be true, and one needs to

consider potential systemic biases that can interfere with the quantification of regulatory

activity when analyzing the data.

We next tested whether potential sequencing biases and other covariates confounded

STARR-seq readouts (Figure 2). We found that STARR-seq RNA coverage was

significantly correlated with GC content (PCC 0.61; P-val 1E-299) and mappability (PCC

0.45; P-val 2.9E-148). This could be attributed to intrinsic sequencing biases in library

preparation. A genome-wide reporter library is made from randomly sheared genomic

DNA, but DNA fragmentation is often non-random [26]. Studies also have suggested

that epigenetic mechanisms and CpG methylation may influence fragmentation [27].

Furthermore, the isolated polyadenylated RNAs are reverse transcribed and PCR

amplified before sequencing, and this process can further confound the sequenced

candidate fragments.

Notably, we found that STARR-seq coverage was also significantly confounded by RNA

thermodynamic stability (PCC -0.55; P-val 0). Unlike ChIP-seq, where both the

experiment and input controls derive from the same DNA origin, STARR-seq

experiments measure the regulatory potential from the abundance of transcribed RNA,

which adds a layer of complexity. For example, RNA structure and co-transcriptional

folding might potentially influence the readout of STARR-seq experiments [28]. Single-

stranded RNA starts to fold upon transcription and the resulting RNA structure might

influence the measurement of regulatory activity. Previously, researchers suggested a

potential linkage between RNA secondary structure and transcriptional regulation [29].

In addition, the resulting transcribed RNA undergoes a series of post-transcriptional

regulation, and RNA stability might play a critical role. Moreover, previous reports have

shown that the degradation rates – the main determinant of cellular RNA levels [30] –

vary significantly across the genome and that RNA stability correlates with functionality

[31,32].

Based on these findings, we built a regression-based model that accounts for various

confounding variables of test sequence fragments to unbiasedly identify potential

enhancer regions from STARR-seq data. Note that many of the covariates have

appreciable correlation with each other. However, we did find, using stepwise forward

selection, that each of them contributes substantially and independently to the model fit

as assessed by Akaike information criterion (AIC) and Bayesian information criterion

(BIC) (Supplementary Figure 2).

Accurate modelling of STARR-seq using negative binomial regression

To model the fragment coverage data from STARR-seq using discrete probability

distribution, we assumed that each genomic bin is independent and identically

distributed, as specified in the Bernoulli trials [33]. That is, each test fragment can only

map to a single fixed-length bin. Therefore, we only considered a non-overlapping

subset of bins for modeling and fitting the distribution. We also excluded bins not

covered by any genomic input or those in which the normalized input coverage was less

than a minimum quantile t_{min}, since these regions do not have sufficient power to detect

enrichment. We selected the bin size and the minimum coverage based on the

experimental design of STARR-seq. We simulated and fitted various discrete probability

distributions to STARR-seq output coverage. We observed that the STARR-seq output

coverage data was overdispersed and fit the best with a negative binomial distribution

(Figure 3A). Moreover, a Q-Q plot of simulated coverage further demonstrated that the

negative binomial model provides the best fit for the data (Figure 3B).

We observed a slight negative enrichment in the STARR-seq output coverage,

suggesting that some candidate fragments can repress the basal transcriptional activity.

However, these regions may contain sequences that can destabilize mRNAs. Therefore,

additional experiments are necessary to demonstrate that STARR-seq can reliably

detect silencers. In the meantime, we suggest opting for a system specifically designed

for identifying silencers for this task [34].

Peak-calling algorithm

To accurately model the ratio of STARR-seq output fragment coverage (RNA) to input

fragment coverage (DNA) while controlling for potential confounding factors, we applied

a negative binomial regression. The overview of our model is outlined in Figure 4. Our

model starts by fitting an analytical distribution to the observed fragment coverage

across fixed non-overlapping genomic bins. In doing so, we use covariates to model

expected counts in the form of multiple regression. Subsequently, once a model is fitted,

we evaluate the likelihood of obtaining the observed fragment counts and assign p-

values using the null negative binomial distribution. In this testing phase, we use flexible

genomic bins with a sliding window in order to find enrichment peaks at a higher

resolution. Genomic bins with significant enrichments are selected based on their

adjusted p-values using multiple testing correction. Finally, peak locations are fine-tuned

to the summit of the direct fragment coverage. Note that the adjusted p-value should be

regarded as the likelihood of a candidate region being an enhancer while the fold

enrichment can be directly interpreted as a quantitative measure of enhancer activity.

Let Y be a vector of STARR-seq output (RNA) coverage, then y_i for $1 \leq i \leq n$ denotes

the number of RNA fragments from a STARR-seq experiment mapped to the i-th bin

from the total of n genomic bins. Let t_i be the number of input library fragments (DNA)

mapped to the i-th bin. We define X to be the matrix of covariates, where $\vec{x_i}$ is the vector

of covariates corresponding to the i-th bin and x_{ij} is the j-th covariate for the i-th bin.

Negative binomial distribution

A negative binomial distribution, which arises from a Gamma-Poisson mixture, can be

parametrized as follows [35–37] (see Methods for derivation).

$$f_Y(y_i|\mu_i, \theta) = \frac{\Gamma(y_i + \theta)}{\Gamma(y_i + 1) \cdot \Gamma(\theta)} \cdot \left(\frac{\theta}{\theta + \mu_i}\right)^{\theta} \cdot \left(\frac{\mu_i}{\theta + \mu_i}\right)^{y_i} \tag{1}$$

A negative binomial is a generalization of a Poisson regression that allows the variance

to be different from the mean, shaped by the dispersion parameter θ. There are two

alternative forms of parametrization for a negative binomial – NB1 and NB2 – which

were first introduced by Cameron and Trivedi [36]. The difference between NB1 and

NB2 is in the conditional variance of y_i. Assuming y_i has mean λ_i, the general variance

function follows the form $\omega_i = \lambda_i + \alpha \lambda_i^{p}$, where α is a scalar parameter. NB1 uses $p = 1$,

whereas NB2 uses the quadratic form of variance with $p = 2$. We use the most common

implementation of the negative binomial, NB2, hereafter. The variance for the NB2

model is given as

$$\sigma^2 = \mu + \frac{\mu^2}{\theta}$$ (Error! Bookmark not defined.)

273   We assume that the majority of genomic bins will have a basal level of transcription, the

274   expected fragment counts at each i-th bin, $E(y_i)$, represents the mean incidence, μ_i,

275   and the count of RNA fragments Y follows the traditional negative binomial (NB2)

276   distribution.

$$E(y_i) = \mu_i$$
$$Y \sim NB(\mu, \theta)$$ (2)

279   *Negative binomial regression model*

280   The regression term for the expected RNA fragment count can be expressed in terms of

281   a linear combination of explanatory variables, a set of m covariates ($\vec{x}$). We use the

282   input library variable t_i as one covariate. For simplicity, we denote t_i as x_{0i} hereafter.

$$\ln \mu_i = \beta_0 x_{0i} + \beta_1 x_{1i} + \cdots + \beta_m x_{mi}$$
$$\mu_i = \exp(\beta_0 x_{0i} + \beta_1 x_{1i} + \cdots + \beta_m x_{mi})$$ (3)
$$\mu_i = \exp\left(\vec{x}_i^\top \beta\right)$$

Alternatively, instead of using the input library variable t_i as one covariate, we can

directly use it as an offset variable. Generally, a fractional observation cannot be

modeled using discrete probability. However, an offset variable in a generalized linear

model can be used to correct the response term to behave like a fraction. One

advantage of using the input variable as an "exposure" to the RNA output coverage is

that it allows us to directly model the basal transcription rate (the ratio of RNA to DNA)

as a rate response variable. More details on this alternative parametrization are

included in the Methods section. In our STARRPeaker model, we used four covariates;

fragment coverage of input genomic libraries, GC content, mappability, and the

thermodynamic stability of genomic libraries.

Maximum-likelihood estimation

We fit the model and estimate regression coefficients using the maximum likelihood

method, where log-likelihood function is shown as follows.

$$\mathcal{L}_{NB}(\mu|y,\theta) = \sum_{i=1}^{n} y_i \ln\left(\frac{\mu_i}{\theta + \mu_i}\right) + \theta \ln\left(\frac{\theta}{\theta + \mu_i}\right)$$
$$+ \ln\left(\frac{\Gamma(y_i + \theta)}{\Gamma(y_i + 1) \cdot \Gamma(\theta)}\right)$$

(Error! Bookmark not defined.)

301   Substituting μ_i with the regression term, the log-likelihood function can be parametrized

302   in terms of regression coefficients, β.

$$\mathcal{L}_{NB}(\beta|y,\theta) = \sum_{i=1}^{n} y_i \ln\left(\frac{e^{\overline{x_i}^\top \beta}}{\theta + e^{\overline{x_i}^\top \beta}}\right) + \theta \ln\left(\frac{\theta}{\theta + e^{\overline{x_i}^\top \beta}}\right)$$

$$+ \ln\left(\frac{\Gamma(y_i + \theta)}{\Gamma(y_i + 1) \cdot \Gamma(\theta)}\right)$$

(Error! Bookmark not defined.)

We can determine the maximum likelihood estimates of the model parameters by

setting the first derivative of the log-likelihood with respect to β, the gradient, to zero,

and there is no analytical solution for $\hat{\beta}$. Numerically, we iteratively solve for the

regression coefficients β and the dispersion parameter θ, alternatively, until both

parameters converge.

Estimation of P-value

The P-value is defined as the probability of observing equal or more extreme value than

the observed value at the i-th bin, y_i, under the null hypothesis.

$$P\text{--}value_i = \Pr(Y \geq y_i|H)$$

(4)

316 As defined earlier, we assume the random variable Y comes from a negative binomial

317 distribution with the fitted mean at the i-th bin, μ_i, as the expected value, and θ as the

318 dispersion parameter. Then, we can estimate the P-value from the cumulative

319 distribution function CDF, which is the sum of the probability mass function f_Y from 0 to

320 $y_i - 1$.

$$\Pr(Y \geq y_i | H) = 1 - CDF(y_i - 1) = 1 - \sum_{k=0}^{y_i - 1} f_Y(k | \mu_i, \theta) \tag{5}$$

323   Substituting (1) gives

$$P\text{-}value_i = 1 - \sum_{k=0}^{y_i - 1} \frac{\Gamma(k + \theta)}{\Gamma(k + 1) \cdot \Gamma(\theta)} \cdot \left(\frac{\theta}{\theta + \mu_i}\right)^\theta \cdot \left(\frac{\mu_i}{\theta + \mu_i}\right)^k \tag{6}$$

Finally, we calculate the false discovery rate using *Benjamini & Hochberg* method [38].

Application of STARRPeaker

We applied STARRPeaker to two whole human genome STARR-seq experiments,

K562 and HepG2, utilizing origin of replication (ORI)-based plasmids [39]. Based on

peaks identified from these datasets, we evaluated the quality and characteristics of the

identified enhancers as well as the performance of the peak caller by comparing to

external enhancer resources.

Initial evaluation of STARRPeaker enhancers

We processed two biological replicates from each cell type independently and assessed

the correlation between each pair. Overall, we observed high correlation between two

replicates (PCC=0.99 for both HepG2 and K562; see Supplementary Figure 3). By

intersecting peaks from two replicates, we identified 32,929 and 20,471 reproducible

candidate enhancers from HepG2 and K562, respectively (Supplementary Table S1).

Although the total number of peaks varied between HepG2 and K562, we observed a

comparable number of peaks within the accessible region of the genome. We found

12,019 (36.34%) and 11,420 (55.57%) candidate enhancers from HepG2 and K562,

respectively, within the open chromatin defined by ENCODE DNase-seq hotspots.

Consistent with previous findings [39], a substantial fraction of candidate enhancers was

epigenetically silenced at the chromatin level. However, as demonstrated previously

using a histone deacetylase inhibitor (HDAC) [16], these poised enhancers can become

functional under a more transcriptionally permissive environment. Therefore, episomal

reporter assays like STARR-seq have the unique advantage of detecting potential

enhancer activity independent from chromatin context. We would like to note that it is

important to identify poised enhancers located in heterochromatic regions of the

genome, which could become functional during developmental or pathological time

courses.

Assessment of robustness and reproducibility of the method

A reliable peak-calling method should be able to identify peaks from suboptimal

datasets. To evaluate the robustness of STARRPeaker, we used subsets of the whole-

genome STARR-seq library to call peaks and compared the results. We subsampled

randomly at various rates from 20 to 80% of the total dataset and compared the quality

of peaks. We found that STARRPeaker was able to reliably identify the peaks using

approximately 60% of the original sequencing library (Supplementary Figure 4).

However, the quality of the peak calls started to deteriorate when 40% or less were

used.

Evaluation of potential orientation bias in candidate enhancers

In general, enhancers are thought to function independent of orientation [40]. However,

the fragment counts in one orientation could be skewed over the other due to

orientation-specific activities, PCR, or sequencing artifacts. To test for potential

orientation-based biases, we ran a binomial test on the candidate enhancers we

identified. We observed a small fraction of candidate enhancers showing strand bias

[3.19% for HepG2 rep1 (n=1,605); 3.76% for HepG2 rep2 (n=1,991); 7.77% for K562

rep1 (n=2,347); 5.25% for K562 rep2 (n=2,195); FDR ≤ 0.01] (Supplementary Figure 5).

Less than one third of the enhancers (n=690) showed strand-specific activity in both

replicates. Thus, we conclude that there is insufficient evidence to show that orientation-

dependent biases are present in our STARR-seq data. Furthermore, this finding

provides further support that enhancers function independent of orientation.

Performance comparison to other peak-calling algorithms

We evaluated the performance of STARRPeaker by comparing it to previously used

methods, namely BasicSTARRseq and MACS2.

First, we qualitatively assessed the peak-calling algorithms using a simulated dataset

where the ground truth exists. We created a STARR-seq dataset that consists of four

spike-in controls (hybrid of DNA input library and RNA output library of known specific

location). All three methods successfully identified the four control peaks with high

confidence (Supplementary Figure 6). However, we noticed that BasicSTARRseq peaks

were fragmented due to its limitation of fixed peak size. Moreover, the peaks were

shifted toward the enrichment of sequencing reads. Furthermore, BasicSTARRseq

identified a false-positive peak, and as a result, identified a total of eight regions instead

of four.

Second, we quantitatively assessed the peak-calling algorithms using the whole human

genome STARR-seq dataset. After uniformly calling peaks from each method using the

recommended default settings, we evaluated the quality of the candidate enhancers

identified. We found that both BasicSTARRseq and MACS2 called significantly more

peaks (4 to 20-fold higher) than STARRPeaker (Supplementary Table S4). While it is

uncertain how many true enhancers were present in each sample, we had to ensure

that we made a fair comparison across different methods due to the tradeoff between

sensitivity and specificity. An increase in sensitivity is generally achieved at the expense

of a decrease in specificity, as described in receiver operating characteristic curves. In

our context, a method having higher specificity suffers from having less overlap with

open chromatin and previously identified enhancers from other assays. Therefore, we

used a uniform P-value threshold of 0.001 and subsampled the peaks before the

comparison. After uniformly processing the dataset using each method, we measured

the level of epigenetic profile enrichment around the peaks. We observed higher

enrichment of DNase-hypersensitive sites, as well as more distinct double-peak

patterns of H3K27ac and H3K4me1, using STARRPeaker compared to

BasicSTARRseq or MACS2 (Figure 5, Supplementary Figure 7). Furthermore,

STARRPeaker peaks had significantly higher enrichment of TF binding events (based

on the number of TF ChIP-seq binding sites) compared to the peaks identified using

other methods.

Comparison to previously characterized enhancers

First, we compared the peaks identified by STARRPeaker to previously characterized

enhancers from HepG2 or K562 cell lines by CAGE [41], MPRA [17,42], and STARR-

seq [19] (Figure 6, Supplementary Table S2). Overall, we observed a higher fraction of

STARRPeaker peaks overlapping with external datasets compare to other methods.

Moreover, we found higher overlaps when peaks from both replicates were merged, due

to fewer but more precise candidate enhancers from merging replicates. However, we

noticed reduced agreement across different types of enhancer assays. Low overlap

between assays may arise from different formats or layouts of reporter plasmids, such

as differing enhancer cloning sites or promoters, or differences in the complexity of the

screening library. Furthermore, CAGE is an entirely different assay from episomal

reporter assays like MPRA and STARR-seq, with enhancers defined based on

bidirectional transcripts originating from an eRNA.

Second, we examined the nine distal enhancers from the GATA1 and MYC loci

characterized in-depth by CRISPRi tiling screen (Supplementary Figure 8). We found

that STARRPeaker accurately called peaks for 6 of 9 enhancers from both replicates.

For the remaining three regions, we observed insufficient enrichment of STARR-seq

output and, therefore, we concluded that this is not a shortcoming of the peak caller.

Application to external STARR-seq datasets

To ensure that STARRPeaker can be generally applied to different variants of STARR-

seq assays, we tested STARRPeaker on previously published STARR-seq datasets.

First, we applied STARRPeaker to the whole-genome ORI-STARR-seq dataset on

HeLa-S3 [39] and assessed the quality of the peaks identified. Consistent with the

previous claim that IFN-I signaling may induce false-positive enhancers, we identified

more peaks in untreated HeLa-S3 samples (n=28,381) compared to inhibitor-treated

samples (n=16,150). Furthermore, peaks from untreated samples had lower enrichment

of chromatin accessibility (DNase-seq) than those from inhibitor-treated samples,

supporting that TBK1/IKK/PKR inhibition reduces false-positive enhancer signals related

to IFN-I signaling (Supplementary Figure 9A). Moreover, STARRPeaker covered 77.5%

(n=7,451) of published peaks, which were called using BasicSTARRseq and then

further shortlisted using a stringent threshold (P-value 1E-5 with corrected enrichment ≥

4). Furthermore, STARRPeaker found 6,540 additional peaks from a HeLa-S3 sample

that was highly enriched with chromatin accessibility signals (Supplementary Figure 9B).

Second, we tested if STARRPeaker can be reliably applied to captured STARR-seq

datasets (Cap-STARR-seq). We applied STARRPeaker to a previously characterized

GM12878 STARR-seq dataset based on an ATAC-seq-capture technique called HiDRA

[43] and compared its performance with published results. The HiDRA dataset was

reported to have ~65,000 regions with enhancer function. In the STARRPeaker run, we

identified only 20,852 regions with significant enhancer activities from the five replicates

they produced. Approximately 73.6% of peaks overlapped with the published results

(n=15,347). While it is debatable to claim that one method is superior to the other, this

result demonstrates that STARRPeaker can be reliably used against the Cap-STARR-

seq dataset.

Third, we further evaluated the performance of the peak-calling methods by applying

STARRPeaker and two other peak-calling methods to another published Cap-STARR-

seq dataset [19]. The dataset covers approximately 91% of the surrounding 3 Mb of the

MYC locus. Consistent with the earlier analysis, we observed that STARRPeaker is

highly specific and identifies fewer candidate enhancers (n=26) compared to the other

methods (BasicSTARRseq n=223; MACS2 n=136). Furthermore, a four-way

comparison (STARRPeaker, BasicSTARRseq, MACS2, and published peaks) showed

that all of the STARRPeaker peaks overlapped with peaks from other methods but not

the other way around (Supplementary Figure 10). These results indicate that

STARRPeaker is more robust and reliable at identifying reproducible candidate

enhancers from various STARR-seq datasets than previous methods.

Conclusions

In summary, we developed a reliable peak-calling analysis pipeline named

STARRPeaker that is optimized for large-scale STARR-seq experiments. To illustrate

the utility of our method, we applied it to two whole human genome STARR-seq

datasets from K562 and HepG2 cell lines, utilizing ORI-based plasmids.

STARRPeaker has several key improvements over previous approaches including (1)

precise and efficient calculation of fragment coverage; (2) accurate modeling of the

basal transcription rate using negative binomial regression; and (3) accounting for

potential confounding factors, such as GC content, mappability, and the thermodynamic

stability of genomic libraries. We demonstrate the superiority of our method over

previously used peak callers, supported by strong enrichment of epigenetic marks

relevant to enhancers and overlap with previously known enhancers.

To fully understand how noncoding regulatory elements can modulate transcriptional

programs in human, STARR-seq active regions must be further characterized and

validated within different cellular contexts. For example, recent applications of CRISPR-

dCas9 to genome editing have allowed researchers to epigenetically perturb and test

these elements in their native genomic context [44,45]. The next step for CRISPR-

based functional screens is to overcome the current limitation of small scale by

leveraging barcodes and single-cell sequencing technology [46]. In the meantime, we

envision that the STARRPeaker framework could be utilized to detect and quantify

enhancers at the whole-genome level, thereby aiding in prioritizing candidate regions in

an unbiased fashion to maximize functional characterization efforts.

Methods

Cell culture

We cultured K562 cells (ATCC) in IMDM (Gibco #12440) supplemented with 10% fetal

bovine serum (FBS) and 1% pen/strep and maintained in a humidified chamber at 37°C

with 5% CO_2. We cultured HepG2 cells (ATCC) in EMEM (ATCC #30-2003)

supplemented with 10% FBS and 1% pen/strep, maintained in a humidified chamber at

37°C with 5% CO_2.

Generating an ORI-STARR-seq input plasmid library

We sonicated human male genomic DNA (Promega #G1471) using a Covaris S220

sonicator (duty factor – 5%; cycle per burst – 200; 40 sec) and ran it on a 0.8% agarose

gel to size-select 500 bp fragments. After gel purification using a MinElute Gel

Extraction kit (Qiagen), we end-repaired, ligated custom adaptors, and PCR-amplified

DNA fragments using Q5 Hot Start High-Fidelity DNA polymerase (NEB) (98°C for 30

sec; 10 cycles of 98°C for 10 sec, 65°C for 30 sec, and 72°C for 30 sec; 72°C for 2 min)

to add homology arms for Gibson assembly cloning.

We used AgeI-HF (NEB) and SalI-HF (NEB) to linearize the hSTARR-seq_ORI plasmid

(gift from Alexander Stark; Addgene plasmid #99296) and cloned the PCR products into

the vector using Gibson Assembly Master Mix (NEB); we set up 60 replicate reactions

to maintain complexity. We purified the assembly reactions using SPRI beads

(Beckman Coulter), dialyzed them using Slide-A-Lyzer MINI dialysis devices

(ThermoScientific), and concentrated them using an Amicon Ultra-0.5 device (Amicon).

We transformed the reaction into MegaX DH10BTM T1 electrocompetent cells (Thermo

Fisher Scientific) (with 25 replicate transformations to maintain complexity) and let them

grow in 12.5L LB-Amp medium until they reached an optical density of ~1.0. We

extracted the plasmids using a Plasmid Gigaprep Kit (Qiagen) and dialyzed the plasmid

prep using Slide-A-Lyzer MINI dialysis devices before electroporation.

Electroporation-mediated transfection of ORI-STARR-seq input plasmid library

into K562 and HepG2 cell lines

We electroporated the ORI-STARR-seq library using an AgilePulse Max (Harvard

Apparatus) and generated two biological replicates for each cell line. For K562 cells, we

electroporated 5.6 mg of input plasmid library into 700 million cells per biological

replicate by delivering three 500 V pulses (1 ms duration with a 20 ms interval). For

HepG2 cells, we electroporated 8 mg of input plasmid library into one billion cells in one

replicate, and 5.6 mg into 700 million cells in another replicate by delivering three 300 V

pulses (5 ms duration with a 20 ms interval).

Generation of an Illumina sequencing library

Output RNA library: We harvested cells 24 hr after electroporation, and extracted total

RNA using an RNeasy Maxi kit (Qiagen). We further isolated polyA-plus mRNA using

Dynabeads® Oligo (dT) kit (ThermoFisher Scientific), treated it with TURBO DNase

(Invitrogen), and purified the reaction using an RNeasy MinElute Kit (Qiagen). We

synthesized cDNA using SuperScript III (ThermoFisher Scientific) with a custom primer

that specifically recognizes mRNAs that had been transcribed from the ORI-STARR-seq

library. After reverse transcription, we treated the reactions with a cocktail of RNase A

and RNase T1 (ThermoFisher Scientific). We split cDNA samples into 160 replicate

sub-reactions, and PCR-amplified each sub-reaction with a primer with a unique index

(helping to identify PCR duplicates) using Q5 Hot Start High-Fidelity DNA polymerase

(NEB) with the following program: 98°C for 30 s; cycles of 98°C for 10 s, 65°C for 30 s,

72°C for 30 s (until they reached mid-log amplification phase; we cycled 18 cycles for

K562 Rep.1; 16 cycles for K562 Rep. 2; 18 cycles for HepG2 Rep. 1; and 15 cycles for

HepG2 Rep2); 72°C for 2 min). After PCR, we re-combined all sub-reactions into one

and purified it with Agencourt Beads. We generated 100 bp paired-end reads for each

biological replicate on an Illumina Hiseq4000 at the University of Chicago Genome

Facility.

Input DNA library: We PCR-amplified a total of 200 ng of input plasmid library (in 16

replicate reactions) using Q5 Hot Start High-Fidelity DNA polymerase (NEB) with the

following program: 98°C for 30 s; 4 cycles of 98°C for 10 s, 65°C for 30 s, and 72°C for

20 s; 8 cycles of 98°C for 10 s and 72°C for 50 s; 72°C for 2 min). After PCR, we

combined all products into one and purified it with Agencourt Beads. We generated 100

bp paired-end reads on an Illumina Hiseq4000 at the University of Chicago Genome

Facility.

Sequencing and preprocessing

For each of 160 replicates, paired-end sequencing reads were aligned to the human

reference genome GRCh38 downloaded from the ENCODE portal (ENCSR425FOI)

using BWA-mem (v0.7.17). Alignments were filtered against unmapped, secondary

alignments, mapping quality score less than 30, and PCR duplicates using SAMtools

(v1.9) and Picard (v2.9.0). All of the replicates were pooled and sorted for downstream

analysis.

Negative binomial distribution

A negative binomial distribution, which arises from Gamma-Poisson mixture, can be

parametrized for y>=0 as follows.

$$Pr(Y = y_i | \mu_i, \theta) = f_Y(y_i; \mu_i, \theta) = \binom{y_i + \theta - 1}{y_i} \cdot \left(\frac{\theta}{\theta + \mu_i}\right)^{\theta} \cdot \left(\frac{\mu_i}{\theta + \mu_i}\right)^{y_i}$$

567   where

$$\binom{y_i + \theta - 1}{y_i} = \frac{\Gamma(y_i + \theta)}{y_i! \cdot \Gamma(\theta)} = \frac{\Gamma(y_i + \theta)}{\Gamma(y_i + 1) \cdot \Gamma(\theta)}$$

569   Substituting gives:

$$f_Y(y_i; \mu_i, \theta) = \frac{\Gamma(y_i + \theta)}{\Gamma(y_i + 1) \cdot \Gamma(\theta)} \cdot \left(\frac{\theta}{\theta + \mu_i}\right)^\theta \cdot \left(\frac{\mu_i}{\theta + \mu_i}\right)^{y_i}$$

571   Rearranging gives:

$$f_Y(y_i; \mu_i, \theta) = \frac{\Gamma(y_i + \theta)}{\Gamma(y_i + 1) \cdot \Gamma(\theta)} \cdot \left(\frac{1}{1 + \frac{\mu_i}{\theta}}\right)^\theta \cdot \left(\frac{\frac{\mu_i}{\theta}}{1 + \frac{\mu_i}{\theta}}\right)^{y_i}$$

$$f_Y(y_i; \theta, \mu_i) = \frac{\Gamma(y_i + \theta)}{\Gamma(y_i + 1) \cdot \Gamma(\theta)} \cdot \left(\frac{\mu_i}{\theta}\right)^{y_i} \left(\frac{1}{1 + \frac{\mu_i}{\theta}}\right)^{\theta + y_i}$$

$$f_Y(y_i; \theta, \mu_i) = \frac{\Gamma(y_i + \theta)}{\Gamma(y_i + 1) \cdot \Gamma(\theta)} \cdot \left(\frac{\mu_i}{\theta}\right)^{y_i} \left(\frac{\theta}{\theta + \mu_i}\right)^{\theta + y_i}$$

$$f_Y(y_i; \theta, \mu_i) = \frac{\Gamma(y_i + \theta)}{\Gamma(y_i + 1) \cdot \Gamma(\theta)} \cdot \frac{\mu_i^{y_i} \theta^\theta}{(\theta + \mu_i)^{\theta + y_i}}$$

Alternative parametrization of negative binomial regression using a rate model

Alternative parametrization allows STARR-seq data to be modelled as a rate model. In

contrast to using input coverage as one of the covariates, we can consider it as

"exposure" to output coverage. This "trick" allows us to directly model the basal

transcription rate (the ratio of RNA to DNA) as a rate response variable. We defined the

transcription rate (RNA to DNA ratio) as a new variable, π_i.

$$\frac{y_i}{t_i} = \pi_i$$

582   If we assume the majority of genomic bins will have the basal transcription rate, we can

583   model the transcription rate at each i-th bin following the traditional negative binomial

584   (NB2) distribution.

$$\pi_i \sim NB\left(\frac{\mu_i}{t_i}, \theta\right)$$

587   The expected basal transcription, $E(\pi_i)$, becomes the mean incidence rate of y_i per unit

588   of exposure, t_i.

$$E\left(\frac{y_i}{t_i}\right) = \frac{\mu_i}{t_i}$$

591   By normalizing μ_i by t_i, we are modeling a rate instead of a discrete count using the

592   negative binomial distribution. The regression term for the expected transcription rate

593   can be expressed in terms of a linear combination of explanatory variables, j covariates

594    $(\vec{x})$.

$$\ln\frac{\mu_i}{t_i} = \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_j x_{ij}$$

597 Rearranging in terms of the expected value of y, or μ, gives

$$\ln\mu_i - \ln t_i = \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_j x_{ij}$$

$$\ln\mu_i = \ln t_i + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_j x_{ij}$$

$$\mu_i = \exp(\ln t_i + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_j x_{ij})$$

The natural log of t_i on the RHS ensures μ_i is normalized in the model, acting as an

offset variable. In STARRPeaker software, we allow users to optionally choose this

alternative rate model (implemented as "mode 2") instead of the default covariate model

described in the main text. This alternate model is useful if constant basal transcription

is expected throughout the genome or if covariates are available for directly modelling

the basal transcription rate π.

BasicSTARRseq

We used BasicSTARRseq R package version 1.10.0 downloaded from Bioconductor

(https://bioconductor.org/packages/release/bioc/html/BasicSTARRseq.html). We used

default setting as described in the software manual, except for disabling deduplication

(minQuantile = 0.9, peakWidth = 500, maxPval = 0.001, deduplicate = FALSE, model =

1), to call peaks.

MACS2

We used MACS2 version 2.1.1 [23] at the recommended default setting, except for

allowing duplicates in read (--keep-dup all), since our STARR-seq dataset was

multiplexed. We called peaks with an FDR cutoff of 0.01, as recommended by the

author of the software.

Calculating folding free energy

We used the LinearFold [47] algorithm to estimate the folding energy of each genomic

bin iteratively across the whole genome. Specifically, we used the Vienna RNAfold

thermodynamic model [48] with parameters from Mathews et al. 2004 [49]. We

implemented a parallel processing scheme to leverage multicore processors to expedite

the calculation of folding free energy.

Declarations

Availability of data and source codes

We implemented the method described in this article as a Python software package

called STARRPeaker. The software package can be downloaded, installed, and readily

used to call peaks from any STARR-seq dataset. The STARRPeaker package, as well

as source code and documentation, is freely available at:

http://github.com/gersteinlab/starrpeaker. All raw data used in the analysis as well as

derived resources are available to download from the ENCODE portal

(https://www.encodeproject.org/) with accession code ENCSR135NXN for HepG2 and

ENCSR858MPS for K562. DNase-seq and ChIP-seq data used for the analysis is also

publicly available from the ENCODE portal. The specific accession codes used for the

analysis are listed in Supplementary Table S3. GC content was downloaded from the

UCSC Genome Browser (http://hgdownload.cse.ucsc.edu/gbdb/hg38/bbi/gc5BaseBw/),

and the mappability track was created using gem-library software [50] with a k-mer size

of 100 bp and the reference human genome build hg38.

Competing Interests

The authors declare that they have no competing interests

Funding

We acknowledge support from the NIH and from the AL Williams Professorship funds.

Author Contributions

D.L., M.S., K.W., and M.G. conceived the project. D.L. and M.G. drafted the manuscript.

D.L. developed the STARRPeaker software package. M.S., J.M., M.W., D.F., Y.K., and

L.M. performed experimental works. M.W. and Y.K. performed experimental validations.

D.L., J.Z., and J.L. performed the downstream analyses. M.G. and K.W. provided

funding and supervised the project.

Acknowledgements

We thank Jinrui Xu and Joel Rozowsky for thoughtful discussion about ChIP-seq

processing, Michael Rutenberg Schoenberg and Zhen Chen for thoughtful discussion

about RNA-folding biology, and all other members of the Gerstein and White

laboratories for advice and critical feedback on the manuscript.

References

1. Muerdter F, Boryń ŁM, Arnold CD. STARR-seq — Principles and applications.

Genomics. Academic Press; 2015;106:145–50.

2. Yáñez-Cuna JO, Kvon EZ, Stark A. Deciphering the transcriptional cis-regulatory

code. Trends Genet. 2013;29:11–22.

3. Lettice LA, Heaney SJH, Purdie LA, Li L, de Beer P, Oostra BA, et al. A long-range

Shh enhancer regulates expression in the developing limb and fin and is associated

with preaxial polydactyly. Hum Mol Genet. 2003;12:1725–35.

4. Banerji J, Rusconi S, Schaffner W. Expression of a beta-globin gene is enhanced by

remote SV40 DNA sequences. Cell. 1981;27:299–308.

5. Sagai T, Hosoya M, Mizushina Y, Tamura M, Shiroishi T. Elimination of a long-range

cis-regulatory module causes complete loss of limb-specific Shh expression and

truncation of the mouse limb. Development. 2005;132:797–803.

6. Melo CA, Drost J, Wijchers PJ, van de Werken H, de Wit E, Vrielink JAFO, et al.

eRNAs Are Required for p53-Dependent Enhancer Activity and Gene Transcription. Mol

Cell. 2013;49:524–35.

7. Sanyal A, Lajoie BR, Jain G, Dekker J. The long-range interaction landscape of gene

promoters. Nature. 2012;489:109–13.

8. Dao LTM, Galindo-Albarrán AO, Castro-Mondragon JA, Andrieu-Soler C, Medina-

Rivera A, Souaid C, et al. Genome-wide characterization of mammalian promoters with

distal enhancer functions. Nat Genet. Nature Publishing Group; 2017;49:1073–81.

9. Diao Y, Fang R, Li B, Meng Z, Yu J, Qiu Y, et al. A tiling-deletion-based genetic

screen for cis-regulatory element identification in mammalian cells. Nat Methods.

2017;14:629–35.

10. Ernst J, Kellis M. ChromHMM: Automating chromatin-state discovery and

characterization. Nat Methods. Nature Research; 2012;9:215–6.

11. Hoffman MM, Buske OJ, Wang J, Weng Z, Bilmes JA, Noble WS. Unsupervised

pattern discovery in human chromatin structure through genomic segmentation. Nat

Methods. Nature Publishing Group; 2012;9:473–6.

12. Sethi A, Gu M, Gumusgoz E, Chan L, Yan K-K, Rozowsky J, et al. A cross-organism

framework for supervised enhancer prediction with epigenetic pattern recognition and

targeted validation. bioRxiv. Cold Spring Harbor Laboratory; 2018;385237.

13. Patwardhan RP, Hiatt JB, Witten DM, Kim MJ, Smith RP, May D, et al. Massively

parallel functional dissection of mammalian enhancers in vivo. Nat Biotechnol. Nature

Publishing Group; 2012;30:265–70.

14. Melnikov A, Murugan A, Zhang X, Tesileanu T, Wang L, Rogov P, et al. Systematic

dissection and optimization of inducible enhancers in human cells using a massively

parallel reporter assay. Nat Biotechnol. Nature Publishing Group; 2012;30:271–7.

15. Arnold CD, Gerlach D, Stelzer C, Boryń ŁM, Rath M, Stark A. Genome-wide

quantitative enhancer activity maps identified by STARR-seq. Science (80-).

2013;339:1074–7.

16. Liu Y, Yu S, Dhiman VK, Brunetti T, Eckart H, White KP. Functional assessment of

human enhancer activities using whole-genome STARR-sequencing. Genome Biol.

2017;18:219.

17. Klein JC, Agarwal V, Inoue F, Keith A, Martin B, Kircher M, et al. A systematic

evaluation of the design, orientation, and sequence context dependencies of massively

parallel reporter assays. bioRxiv. Cold Spring Harbor Laboratory; 2019;576405.

18. Johnson GD, Barrera A, McDowell IC, D'Ippolito AM, Majoros WH, Vockley CM, et

al. Human genome-wide measurement of drug-responsive regulatory activity. Nat

Commun. 2018;9:5317.

19. Rathert P, Roth M, Neumann T, Muerdter F, Roe J-S, Muhar M, et al.

Transcriptional plasticity promotes primary and acquired resistance to BET inhibition.

Nature. 2015;525:543–7.

20. Koohy H, Down TA, Spivakov M, Hubbard T. A comparison of peak callers used for

DNase-Seq data. Helmer-Citterich M, editor. PLoS One. 2014;9:e96303.

21. Uren PJ, Bahrami-Samani E, Burns SC, Qiao M, Karginov F V, Hodges E, et al. Site

identification in high-throughput RNA-protein interaction data. Bioinformatics. Oxford

University Press; 2012;28:3013–20.

22. Strbenac D, Armstrong NJ, Yang JYH. Detection and classification of peaks in 5'

cap RNA sequencing data. BMC Genomics. BioMed Central; 2013;14 Suppl 5:S9.

23. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-

based Analysis of ChIP-Seq (MACS). Genome Biol. BioMed Central; 2008;9:R137.

24. Kharchenko P V, Tolstorukov MY, Park PJ. Design and analysis of ChIP-seq

experiments for DNA-binding proteins. Nat Biotechnol. Nature Publishing Group;

2008;26:1351–9.

25. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25:2078–9.

26. Poptsova MS, Il'icheva IA, Nechipurenko DY, Panchenko LA, Khodikov M V,

Oparina NY, et al. Non-random DNA fragmentation in next-generation sequencing. Sci

Rep. 2014;4:4532.

27. Lazarovici A, Zhou T, Shafer A, Dantas Machado AC, Riley TR, Sandstrom R, et al.

Probing DNA shape and methylation state on a genomic scale with DNase I. Proc Natl

Acad Sci U S A. 2013;110:6376–81.

28. Lai D, Proctor JR, Meyer IM. On the importance of cotranscriptional RNA structure

formation. RNA. 2013;19:1461–73.

29. Ringnér M, Krogh M. Folding free energies of 5'-UTRs impact post-transcriptional

regulation on a genomic scale in yeast. PLoS Comput Biol. 2005;1:e72.

30. Rabani M, Levin JZ, Fan L, Adiconis X, Raychowdhury R, Garber M, et al. Metabolic

labeling of RNA uncovers principles of RNA production and degradation dynamics in

mammalian cells. Nat Biotechnol. 2011;29:436–42.

31. Yang E, van Nimwegen E, Zavolan M, Rajewsky N, Schroeder M, Magnasco M, et

al. Decay rates of human mRNAs: Correlation with functional characteristics and

sequence attributes. Genome Res. 2003;13:1863–72.

32. Tani H, Mizutani R, Salam KA, Tano K, Ijiri K, Wakamatsu A, et al. Genome-wide

determination of RNA stability reveals hundreds of short-lived noncoding transcripts in

mammals. Genome Res. 2012;22:947–56.

33. Papoulis A, Athanasios. Probability, random variables and stochastic processes.

New York McGraw-Hill, 1984, 2nd ed. 1984;

34. Pang B, Snyder MP. Systematic identification of silencers in human cells. Nat Genet.

Nature Publishing Group; 2020;52:1–10.

35. Hilbe JM. Negative Binomial Regression [Internet]. Cambridge: Cambridge

University Press; 2011.

36. Cameron ACA, Trivedi PK. Regression Analysis of Count Data [Internet].

Cambridge: Cambridge University Press; 2013.

37. Hilbe JM. Modeling Count Data [Internet]. Cambridge: Cambridge University Press;

2014.

38. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. J R Stat Soc Ser B. John Wiley & Sons, Ltd

(10.1111); 1995;57:289–300.

39. Muerdter F, Boryń ŁM, Woodfin AR, Neumayr C, Rath M, Zabidi MA, et al.

Resolving systematic errors in widely used enhancer activity assays in human cells. Nat

Methods. Nature Publishing Group; 2018;15:141–9.

40. Pennacchio LA, Bickmore W, Dean A, Nobrega MA, Bejerano G. Enhancers: five

essential questions. Nat Rev Genet. NIH Public Access; 2013;14:288–95.

41. Kawaji H, Kasukawa T, Forrest A, Carninci P. The FANTOM 5 collection, a data

series underpinning mammalian transcriptome atlases in diverse cell types. Sci Data.

2017;2016–8.

42. Inoue F, Kircher M, Martin B, Cooper GM, Witten DM, McManus MT, et al. A

systematic comparison reveals substantial differences in chromosomal versus episomal

encoding of enhancer activity. Genome Res. 2017;27:38–52.

43. Wang X, He L, Goggin SM, Saadat A, Wang L, Sinnott-Armstrong N, et al. High-

resolution genome-wide functional dissection of transcriptional regulatory regions and

nucleotides in human. Nat Commun. Nature Publishing Group; 2018;9:1–15.

44. Dixit A, Parnas O, Li B, Chen J, Fulco CP, Jerby-Arnon L, et al. Perturb-Seq:

Dissecting Molecular Circuits with Scalable Single-Cell RNA Profiling of Pooled Genetic

Screens. Cell. Cell Press; 2016;167:1853-1866.e17.

45. Xie S, Duan J, Li B, Zhou P, Hon GC. Multiplexed Engineering and Analysis of

Combinatorial Enhancer Activity in Single Cells. Mol Cell. Cell Press; 2017;66:285-

299.e5.

46. Gasperini M, Hill AJ, McFaline-Figueroa JL, Martin B, Kim S, Zhang MD, et al. A

Genome-wide Framework for Mapping Gene Regulation via Cellular Genetic Screens.

Cell. Cell Press; 2019;176:377-390.e19.

47. Huang L, Zhang H, Deng D, Zhao K, Liu K, Hendrix DA, et al. LinearFold: linear-

time approximate RNA folding by 5'-to-3' dynamic programming and beam search.

Bioinformatics. Narnia; 2019;35:i295–304.

48. Lorenz R, Bernhart SH, Höner zu Siederdissen C, Tafer H, Flamm C, Stadler PF, et

al. ViennaRNA Package 2.0. Algorithms Mol Biol. BioMed Central; 2011;6:26.

49. Mathews DH, Disney MD, Childs JL, Schroeder SJ, Zuker M, Turner DH.

Incorporating chemical modification constraints into a dynamic programming algorithm

for prediction of RNA secondary structure. Proc Natl Acad Sci U S A. National Academy

of Sciences; 2004;101:7287–92.

50. Derrien T, Estellé J, Marco Sola S, Knowles DG, Raineri E, Guigó R, et al. Fast

Computation and Applications of Genome Mappability. Ouzounis CA, editor. PLoS One.

2012;7:e30377.

797 **Supplementary Tables**

Table S1 contains significant peaks called by STARRPeaker.

Table S2 contains overlap of various peak callers (STARRPeaker, BasicSTARRseq,

and MACS2) to published enhancers identified using other types of enhancer assays.

Table S3 contains a list of data sources and accession numbers used for the analysis.

Table S4 compares peaks identified by various peak callers (STARRPeaker,

BasicSTARRseq, and MACS2).

Supplementary Figures

Supplementary Figure 1 Comparison of STARR-seq output coverage calculated using

the center of the fragment to using the start position of the sequencing read. **(A)**

Distribution of shift in final peak locations resulting from using two alternative coverage

counting schemes in HepG2. Comparison of **(B)** overall fold enrichment level, **(C)** p-

value, and **(D)** size of resulting peaks.

Supplementary Figure 2 Contribution of covariates and model selection. **(A)** Q-Q plots

of various models with different sets of covariates showing the goodness of fit. **(B)** Both

AIC and BIC measure relative qualities of statistical models considering the trade-off

between the goodness of fit and the simplicity of the model. AIC: Akaike information

criterion; BIC: Bayesian information criterion.

Supplementary Figure 3 Correlation between replicates for **(A)** HepG2 or **(B)** K562 cell

lines.

Supplementary Figure 4 Comparison of peaks called from subsamples of the original

STARR-seq library, highlighting the robustness of STARRPeaker.

Supplementary Figure 5 Orientation biases analysis for **(A-B)** HepG2 or **(C-D)** K562

cell lines. The ratio between forward and reverse stranded fragments was tested for

statistical significance using a binomial test. Orange dots represent peaks with

significant strand bias (FDR q-value < 0.01).

Supplementary Figure 6 Comparison of peaks identified by various methods using a

simulated STARR-seq dataset containing four spike-in control regions.

Supplementary Figure 7 Enrichment of epigenetic signals around peaks in K562. All

peaks were centered at the summit, uniformly thresholded using P-value < 0.001, and

10,000 peaks were randomly selected. Aggregated read depth at 2,000 bp upstream

and downstream were plotted for **(A)** DNase I hypersensitive sites (DHS), **(B)** H3K27ac,

(C) H3K4me1, and **(D)** aggregated TF ChIP-seq profile. For DNase-seq, enrichment

indicates unique read depth. For histone ChIP-seq, enrichment indicates fold change

over control. For TF ChIP-seq aggregate, enrichment indicates the number of TFs

binding.

Supplementary Figure 8 (A-C) Genome browser session comparing STARRPeaker to

other peak-calling methods at validated enhancers from CRISPRi.

Supplementary Figure 9 Application of STARRPeaker on an external HeLa-S3 dataset.

(A) Comparison of chromatin accessibility (DNase-seq) for STARRPeaker peaks

between untreated and inhibitor-treated samples. **(B)** Comparison of STARRPeaker

peaks to published results. STARRPeaker found 6,540 additional peaks that are

enriched with chromatin accessibility signals from a HeLa-S3 sample.

Supplementary Figure 10 Venn diagram for four-way comparison of peaks identified

by various methods using a published dataset from Rathert et al. 2015.

Figure legends

Figure 1 Comparison of STARR-seq output coverage calculated using the center of the

fragment to using the start position of the sequencing read. **(A)** Distribution of the shift in

final peak locations resulting from using two alternative coverage counting schemes in

HepG2. Comparison of **(B)** overall fold enrichment level, **(C)** p-value, and **(D)** size of

resulting peaks. **(E)** Example highlighting the difference between fragment-based and

read-based coverage counting schemes and their resulting peak calls from HepG2

STARR-seq data. Asterisks represents statistical significance using the Mann-Whitney-

Wilcoxon test two-sided with Bonferroni correction; (*) P <= 0.05, (**) P <= 0.01, (***) P

<= 0.001, (****) P <= 0.0001.

Figure 2 Confounding factors in the STARR-seq assay. STARR-seq output and input

coverages are significantly correlated with **(A)** input coverage, **(B)** GC content, **(C)**

mappability, and **(D)** RNA structure folding. PCC: Pearson Correlation Coefficient. Plots

were from a sampling of 5,000 random genomic bins.

Figure 3 STARR-seq output coverage is fitted against simulated coverage using three

distribution models; negative binomial, binomial, and Poisson. **(A)** Density histogram of

simulated distribution against STARR-seq output coverage. **(B)** Q-Q plot of simulated

distribution against STARR-seq output coverage. The red solid line represents where

the observed count equals the expected count.

Figure 4 Overview of STARRPeaker peak-calling scheme. **(A)** In contrast to using read

depth (grey), fragment depth (red) offers more precise and sharper STARR-seq output

coverage. Fragment inserts are directly inferred from properly paired-reads. **(B)**

Workflow of STARRPeaker describing how coverage is calculated for each genomic bin

and modelled using a negative binomial regression model. The analysis pipeline can

largely be divided into four steps: (1) Binning the genome; (2) calculating coverage and

computing covariate matrix; (3) fitting the STARR-seq data to the NB regression model;

and (4) peak calling, multiple hypothesis testing correction, and adjustment of the center

of peaks.

Figure 5 Enrichment of epigenetic signals around peaks in HepG2. All peaks were

centered at the summit, uniformly thresholded using P-value < 0.001, and 10,000 peaks

were randomly selected. Aggregated read depth at 2,000 bp upstream and downstream

were plotted for **(A)** DNase I hypersensitive sites (DHS), **(B)** H3K27ac, **(C)** H3K4me1,

and **(D)** aggregated TF ChIP-seq profile. For DNase-seq, enrichment indicates unique

read depth. For histone ChIP-seq, enrichment indicates fold change over control. For

TF ChIP-seq aggregate, enrichment indicates the number of TFs binding.

Figure 6 Comparison of peaks using an external dataset for **(A)** HepG2 or **(B)** K562 cell

lines. Peaks identified from STARRPeaker as well as BasicSTARRseq and MACS2

were compared against a published dataset. For a fair comparison, all peaks were

centered at the summit, uniformly thresholded using P-value < 0.001, and 20,000 peaks

were randomly drawn from peaks identified by each peak caller using the recommended

settings. The fraction of overlap was computed for each replicate. We considered it an

overlap when at least 50% of peaks intersected each other.

Figure 1

A

B HepG2

**** ****

fold change — rep1, rep2

C HepG2

**** ****

-logPval — rep1, rep2

D HepG2

**** ****

peak size — rep1, rep2

E

chr17:79,025,000-79,028,500

Coverage

Peak **FC=7.366** Fragment Center

Coverage

Peak **FC=2.726** Read Start

Fragment Center
Read Start

Figure 2

Figure 3

Figure 4

A

Read Depth

Read Distribution

Genomic Bin

Fragment Depth

Peak

B

Create Genomic Bins → Calculate Input Library Coverage → Fit model Y = NB(μ,θ) → FDR adjusted P-value → Fine-tune Peak Center

Calculate STARR-seq Coverage

Create Covariate Matrix

Figure 5

Figure 6

Supplementary Figure 1

A

B

C

D

Fragment Center
Read Start

Supplementary Figure 2

A

B

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

A

B

C

D

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 8

A

GATA1 locus chrX:48774179-48806695

B

MYC locus chr8:127830000-128090000

C

MYC locus chr8:129550000-129720000

Supplementary Figure 9

A

B

Supplementary Figure 10