Exploring the phylogeny of rosids with a five-locus supermatrix from GenBank

Miao Sun[1, 2, 3*], Ryan A. Folk[1], Matthew A. Gitzendanner[2, 6], Stephen A. Smith[4], Charlotte

Germain-Aubrey[1], Robert P. Guralnick[1, 6], Pamela S. Soltis[1, 5, 6], Douglas E. Soltis[1, 2, 5, 6],

Zhiduan Chen[3*]

1. Florida Museum of Natural History, University of Florida, Gainesville, FL 32611, USA

2. Department of Biology, University of Florida, Gainesville, FL 32611, USA

3. State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, the

Chinese Academy of Sciences, Beijing 100093, China

4. Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI

48109, USA

5. Genetics Institute, University of Florida, Gainesville, FL 32608, USA

6. Biodiversity Institute, University of Florida, Gainesville, FL 32611, USA

*Corresponding authors:

Miao Sun: cactusresponsible@gmail.com

Zhiduan Chen: zhiduan@ibcas.ac.cn

Abstract: 217

Text: total words: 3553

Introduction: 720

Materials and Methods: 932

Results: 1247

Discussion: 724

Acknowledgements: 96

Author contributions: 62

Figures: 3

Colored: 3

Tables: 2

Supporting information:

Supplementary tables: 3

Supplementary file: 1

Running headline: Phylogeny and dating of rosids

Abstract

Current advances in sequencing technology have greatly increased the availability of

sequence data from public genetic databases. With data from GenBank, we assemble and

phylogenetically investigate a 19,740-taxon, five-locus supermatrix (i.e., *atpB*, *rbcL*, *matK*,

matR, and ITS) for rosids, a large clade containing over 90,000 species, or approximately a

quarter of all angiosperms (assuming an estimate of 400,000 angiosperm species). The

topology and divergence times of the five-locus tree generally agree with previous estimates

of rosid phylogeny, and we recover greater resolution and support in several areas along the

rosid backbone, but with a few significant differences (e.g., the placement of the COM clade,

as well as Myrtales, Vitales, and Zygophyllales). Our five-locus phylogeny is the most

comprehensive DNA data set yet compiled for the rosid clade. Yet, even with 19,740 species,

current sampling represents only 16-22% of all rosids, and we also find evidence of strong

phylogenetic bias in the accumulation of GenBank data, highlighting continued challenges

for species coverage. These limitations also exist in other major angiosperm clades (e.g.,

asterids, monocots) as well as other large, understudied branches of the Tree of Life,

highlighting the need for broader molecular sampling. Nevertheless, the phylogeny presented

here improves upon sampling by more than two-fold and will be an important resource for

macroevolutionary studies of this pivotal clade.

Keywords: Mega-phylogeny, rosids, dating, GenBank, sampling

I. Introduction

Given their size, rosids (*Rosidae*; Cantino et al., 2007; Wang et al., 2009; APG IV, 2016) have great potential for understanding the evolution and diversification of angiosperms. A clade of 90,000–120,000 species (estimated from the Open Tree of Life and Open Tree Taxonomy database (OTT); Hinchliff et al., 2015), the rosid clade represents more than a quarter of all angiosperms (based on an estimated 400,000 species of angiosperms; Govaerts, 2001). Rosids comprise two large subclades, fabids (i.e., eurosids I, *Fabidae*) and malvids (i.e., eurosids II, *Malvidae*) and are further divided into 17 orders and 135 families (APG IV, 2016). The clade originated in the Early to Late Cretaceous (115 to 93 Million years ago, Myr), followed by rapid diversification yielding the crown groups of fabids (112 to 91 Myr) and malvids (109 to 83 Myr; Wang et al., 2009; Bell et al., 2010; Magallón et al., 2015). The rosid clade diversified rapidly to form the two major lineages in perhaps as little as 4 to 5 million years (Wang et al., 2009; Bell et al., 2010).

Most rosid families have high species diversity in the tropics, but many extend from the tropics to subtropical and temperate areas (e.g., Wang et al., 2009; Soltis et al., 2010). Most ecologically dominant forest trees are found within the clade, as well as diverse aquatics, parasites, arctic, alpine, and desert lineages; the clade also exhibits tremendous diversity in chemistry, reproductive strategy, and life history (Magallón et al., 1999; Wang et al., 2009; Stevens, 2001 onwards). Unique ecological traits are prevalent in the rosids, including nodular association with nitrogen-fixing bacteria (Soltis et al., 1995; Li et al., 2015), chemical defense mechanisms including glucosinolate production in Brassicales (Rodman et al., 1998; Soltis et al., 2005; Edger et al., 2015), and independent origins of parasitism, sometimes associated with rampant horizontal gene transfer (e.g., *Rafflesia*; Davis & Wurdack, 2004; Xi et al., 2012). Many important crops are rosids, including cotton and cacao (Malvaceae), hops (Cannabaceae), legumes (Fabaceae), rubber (Euphorbiaceae), and numerous vegetable and fruit crops (Brassicaceae, Caricaceae, Cucurbitaceae, Rosaceae, Rutaceae, and Vitaceae). Some rosids have been selected as genetic models, including *Arabidopsis thaliana* (*Arabidopsis* Genome Initiative, 2000), *Brassica rapa* (Wang et al., 2011), and various legumes (Sato et al., 2008; Schmutz et al., 2010, 2014; Varshney et al., 2012, 2013; Young et al., 2011).

The "rise of the rosids" yielded most angiosperm-dominated forests present today. Many other lineages of life radiated in the shadow of these rosid-dominated forests (e.g., ants: Moreau et al., 2006; Moreau & Bell, 2013; beetles: Farrell, 1998; Wilf et al., 2000;

amphibians: Roelants et al., 2007; mammals: Bininda-Emonds et al., 2007; fungi: Hibbett &

Matheny, 2009; liverworts: Feldberg et al., 2014; ferns: Schneider et al., 2004; Watkins &

Cardelús, 2012). The initial rise of the rosids and subsequent repeated cycles of radiations

within the rosid clade (Soltis & Soltis, 2004; Soltis et al., 2004) have profoundly shaped

much of current terrestrial biodiversity (Wang et al., 2009; Boyce et al., 2010).

Although rosids have long been the focus of phylogenetic research (e.g., Wang et al.,

2009; Soltis et al., 2011; Zeng et al., 2017; and references therein), the enormous size of this

clade has thus far precluded achieving the sampling required for macroevolutionary

inferences (Ricklefs, 2007; Smith et al., 2011; Thomas et al., 2013; Folk et al., 2018). Hence,

a robust, time-calibrated phylogeny with large-scale species-level sampling is needed for

future diversity studies. Additionally, the rosid clade also provides an opportunity to evaluate

the implications of taxon and gene sampling. Using data from GenBank, we constructed a 5-

locus phylogenetic tree having more than twice the taxon sampling used in earlier studies

(e.g., the 4-locus study of Sun et al., 2016)—and compared this tree to all rosid names in the

OpenTree of Life (Hinchliff et al., 2015) to quantify bias in DNA sampling across the clade.

We hypothesized that (1) taxon sampling remains highly biased across the large rosid clade,

and that (2) the use of more genes and increased taxon sampling impacts phylogenetic

resolution and divergence time estimation. We tested these hypotheses via a series of

comparisons across two trees: a previously published 4-locus, 8,855-taxon supermatrix (Sun

et al, 2016) and the more densely sampled 5-locus, 19,740-taxon supermatrix generated here.

We then quantified patterns of taxon sampling bias, phylogenetic resolution, and time

calibration.

II.   Materials and Methods

Data mining, alignment, and phylogeny reconstruction

We mined GenBank (Release 214: June 15, 2016) for the chloroplast genes *atpB*,

matK, and *rbcL*, the mitochondrial gene *matR*, and the nuclear ribosomal ITS (including ITS-

1, 5.8S, and ITS-2 regions) using the PHyLogeny Assembly With Databases pipeline

(PHLAWD, version 3.4a, https://github.com/blackrim/phlawd; Smith et al., 2009). These

genes represent those most commonly used in phylogenetic studies of plants and therefore the

most numerous loci for plants deposited in GenBank; they also represent all three plant

genomes.

We employed PHLAWD data mining using three bait sequences of each target locus

that represent the phylogenetic diversity of the rosid clade (Hinchliff & Smith, 2014). The

quality of sequence data from the five sampled loci was investigated by calculating the best-

hit scores from BLAST and plotting the distribution of identity scores ([0, 1]) against

coverage scores ([0, 1]). Low-quality and outlier sequences were removed based on these

scores (see Results). For all resulting alignments, we (1) validated the species names

following The Plant List (TPL; http://www.theplantlist.org/), using the R package

Taxonstand v2.0 (Cayuela et al., 2012), and then (2) pruned all taxa with "subsp.", "var.",

"f.", "cf." and "aff." designations in taxon names. Names for orders and families follow APG

IV (2016), and those for major supra-ordinal clades follow Soltis et al. (2011) and Cantino et

al. (2007).

We curated the 5-locus data set iteratively by screening individual loci and

concatenated matrices (below) for rogue taxon behavior through manual inspection of initial

phylogenies for spurious taxon placement and through the RAxML dropsets algorithm

(Pattengale et al., 2010a; Sun et al., 2016; Smith & Brown, 2018). The final 5-locus data set

contained 19,740 ingroup species (135 families and 17 orders) and 20,294 species, including

outgroup taxa (i.e., 554 outgroup species in 17 families and three orders) from Saxifragales,

Proteales, and Trochodendrales.

We also compared results from our 5-locus (*atpB*, *rbcL*, *matK*, *matR*, and ITS) data

set with those from a previously published 4-locus rosid data set comprising chloroplast and

mitochondrial loci (Sun et al., 2016; *atpB*, *rbcL*, *matK*, and *matR*). The 4-locus data set

contained 8,855 ingroup taxa (9,300 taxa including outgroups; i.e., 445 outgroup species

from the same three orders; Sun et al., 2016). Hereafter, all tree sampling statistics will only

concern the ingroup.

Taxon sampling analyses

To evaluate sampling gaps among all taxonomically recorded rosid species and the

species included in the 5-locus data set, we mapped the 19,740 validated rosid names in our

phylogeny against all rosid species present in OTT v3.0

(https://devtree.opentreeoflife.org/about/taxonomy-version/ott3.0; Hinchliff et al., 2015).

Generic names of this complete list were also manually curated via an online tool, Index

Nominum Genericorum (Farr & Zijlstra, 1996 onwards), and then invalid, rejected (nom.

rej.), illegitimate, and synonymous generic names were all removed, as well as any taxon

names with "sp.", "subsp.", "var.", "x", "cf." and "aff.", or other non-species designations

(e.g., "spp.", "clone", "environmental sample", "group").

Additionally, to evaluate whether sampling in rosid DNA data from GenBank is

phylogenetically biased, we scored DNA data presence and absence by mapping our

phylogeny names to OpenTree (Smith & Brown, 2018) and then executing a λ test on this

"trait" under an equal rates model (R packge geiger V.2.0.6.2; Pennell et al., 2014).

Significance was assessed with a likelihood ratio test.

Phylogenetics

The edited and pruned alignments of each locus were concatenated into a single

supermatrix using FASconCAT v.1.0 (Kück & Meusemann, 2010). We ran maximum

likelihood (ML) analyses for each individual locus alignment and for the concatenated

matrices using RAxML v.8.2.10 (Stamatakis, 2014) with 100 bootstrap (BS) replicates for

topology examination, under an unpartitioned GTRCAT model. For the 5-locus concatenated

matrix, the best ML tree was constructed with RAxML using the extended Majority Rule

Criterion (autoMRE) as a bootstrap stopping rule (Pattengale et al., 2010b; reached at 352

replicates). We visually examined potential topological conflicts by concatenating different

data sets and evaluating strongly supported differences among trees at the family level

inferred from the combined supermatrix and of each individual data set (i.e., ITS, *matR*, and

the plastid genes; see Results and Discussion). Trees were manipulated for display using

Newick utilities (Junier & Zdobnov, 2010), Dendroscope 3 (Huson & Scornavacca, 2012),

MEGA (Tamura et al., 2013), and iTOL v3.0 (Letunic & Bork, 2016).

Divergence Time Estimation

Divergence time estimation was conducted using both the previously published 4-locus

(Sun et al., 2016) phylogeny and newly constructed 5-locus phylogeny. In total, 59

calibration points (covering 15/17 rosid orders) were used as time constraints, based on

validated fossils frequently used as calibration points in previous molecular dating studies

(Davis et al., 2005; Wang et al., 2009; Bell et al., 2010; Sauquet et al., 2012; Magallón et al.,

2015; Table S1). The root was constrained to a maximum age of 125 Myr following Wang et

al. (2009). For both phylogenies, we used the penalized likelihood program treePL v.1.0

(Smith & O'Meara, 2012) to generate a time-calibrated ultrametric tree. We initially

conducted random cross-validation procedures with three options ("randomcv", "thorough",

and "prime") to determine the best smoothing value and optimization parameters for both the

4-locus and 5-locus ML trees and then ran 200,000 annealing iterations (default = 5,000) for

divergence time estimation.

To estimate variation in the timing of the rosid divergence, we also employed PATHd8

v.1.0 (Britton et al., 2006) using the same 59 calibration points and root constraint as above.

Unlike treePL, PATHd8 is a faster heuristic method that sequentially takes averages over

path lengths from an internode to all its descending terminals, one pair of sister groups at a

time (Ericson et al., 2006; Anderson, 2007).

III.   Results

Limitations in Taxon and Locus Sampling

For the commonly sequenced locus ITS, PHLAWD initially recovered 42,890 rosid

sequences; after removing sequences with non-species designations (cf. Materials and

Methods), 39,735 sequences remained. Removal of sequences with low identity and coverage

scores (coverage score ≤ 0.1 and identity score ≤ 0.1 were considered low quality) and further

duplicate removal and non-ortholog cleaning, 15,100 sequences remained in the single locus

matrix; however, only 13,157 sequences were retained in the final combined supermatrix for

phylogeny reconstruction, due to two reasons: 1) some ITS sequences still exhibiting

characteristics of rogue taxa identified by initial RAxML analyses in primary 5-locus

supermatrix (Pattengale et al., 2010; Sun et al., 2016); and 2) in this combined matrix, some

species have only one fragment of either ITS1, 5.8S, or ITS2, and the other four genes are not

available; therefore, these short single ITS fragments were removed from the combined

matrix to avoid introducing large amounts of missing data. Updating the remaining loci

(*atpB*, *rbcL*, *matK*, and *matR*) with new GenBank data resulted in 1,257, 6,960, 8,489, and

721 sequences, respectively. The alignment lengths for *atpB*, *rbcL*, *matK*, *matR,* and ITS

were, respectively, 1,500, 1,401, 1,815, 2,349, and 835 bp, with a concatenated length of

7,900 bp, and 70.55% missing data.

To better understand sampling patterns in our data sets, we matched our recovered

phylogenetic tips (species) with those in the Open Tree of Life taxonomic database (OTT

v3.0; Hinchliff et al., 2015), which includes rosid clades—e.g., Rutaceae, Francoaceae, and

Kirkiaceae—yet to be integrated into the Open Tree topology itself. We sampled 135 families

(100% coverage of the rosid families recognized in APG IV, 2016), 3,070 genera (matching

66.34% of OTT), and 19,740 species (matching 16.25% of OTT). The unsampled genera and

species mainly reflect absence of DNA data (Fig. 1 and Tables 1, S2), but some are due to

taxonomic issues such as synonyms and invalid names. Among these mismatches are names

in our 5-gene tree unaccounted for in OTT, comprising 1,134 species (5.74%) and 72 genera

(2.35%). Total coverage and taxon representation compared with the rosids present in OTL

and OTT are summarized in Table 1, and Table S2, respectively (order and family

circumscription adjusted to comply with APG IV [2016]; marked with asterisks in Table S2).

Sampling coverage in our tree shows a strong phylogenetic bias (p-value for λ test, p

≈ 0; Fig. 1 and Tables 1, S2). Overall, larger orders (> 10,000 species, e.g., Rosales and

Myrtales) tend to be more poorly sampled with > 90% of the species unsampled (Table 1).

Several large families (> 1,000 species) also have poor coverage; Polygalaceae, Rosaceae,

Myrtaceae, Malvaceae, Rutaceae, and Phyllanthaceae have 3.14% to 15.27% sampling (Table

S2). Geraniales, Crossosomatales, Brassicales, Cucurbitales, and Huerteales have better

sampling, yet no order or family exceeds 50% coverage of known species richness.

Phylogenetic Analyses

The topology of our 5-locus rosid tree (Fig. S1) generally agrees with that of the 4-

locus tree inferred in a previous GenBank mining effort (Sun et al., 2016), but provides

greater resolution and support in several areas along the backbone (Fig. S1) as well as greatly

improved species-level sampling. The median BS value of the 5-locus rosid tree is lower than

that obtained in the 4-locus, 8,855-taxon rosid phylogeny (Fig. 2a), but within

methodological expectations (see Discussion).

The topologies inferred from single locus partitions and the concatenated data set are

generally consistent, with the exception of the following conflicting phylogenetic placements:

(1) For the COM clade (Celastrales-Oxalidales-Malpighiales), we observed the same

conflicting placements when trees from nuclear, plastid, and mitochondrial data are compared

as observed in Sun et al. (2015, 2016). Nuclear and mtDNA data favor a placement of the

COM clade with malvids, whereas plastid data indicate a placement with fabids. (2) The

placement of Myrtales and Geraniales was unstable across locus partitions. In the chloroplast

tree, Myrtales and Geraniales were sequential sisters to fabids with strong support, but in the

matR tree and total evidence tree, Myrtales and Geraniales were sisters to the rest of the

rosids (cf. Sun et al., 2015, 2016). The monophyly of Myrtales was not supported in the ITS

tree. (3) The placement of Zygophyllales varied among locus partitions. In the *matR* tree,

Zygophyllales were resolved as sister to malvids (cf. Sun et al., 2016; Zhao et al., 2016), a

different result from the chloroplast and total evidence trees, where the clade was placed

within fabids with low to moderate support. (4) The placement of Vitales with respect to the

rest of the rosids and Saxifragales was unstable across analyses. In the *matR* and total

evidence trees, the three groups were resolved as (rosids + Vitales) + Saxifragales (cf. Sun et

al., 2016), a topology that has also been recovered in most studies (Soltis et al., 2007, 2011;

Worberg et al., 2007; Zhu et al., 2007; APG, 2009, 2016; Wang et al., 2009; Smith et al.,

2010; Barniske et al., 2012; Ruhfel et al., 2014). Our chloroplast data set, by contrast,

recovers the relationship among these three clades as rosids + (Saxifragales + Vitales), a

result also seen previously (cf. Moore et al., 2010; Ruhfel et al., 2014; Zhang et al., 2012,

2016; Sun et al., 2016). (5) Non-monophyly of some families was occasionally seen within

single-gene trees. Two families (Cannabaceae and Euphorbiaceae) are resolved as non-

monophyletic in the combined plastid tree, while they are recovered as monophyletic in the

total evidence tree as expected; similarly, 12 families are non-monophyletic in the *matR* gene

tree and 18 in the ITS tree (see Table S3).

Dating Analyses

The crown age of rosids is estimated as 117.93 Myr by treePL (89.80 Myr by PATHd8)

using the 5-locus tree, and 122.62 Myr by treePL (104.20 Myr by PATHd8) for the 4-locus

tree. The ages of other major rosid clades are reported in Table 2. Comparisons of crown ages

obtained for all major rosid clades (17 orders and 135 families, *sensu* APG IV) are provided

in Figs. 2b and 3, covering both treePL and PATHd8 and both 4-locus and 5-locus trees, as

well as previous studies (Wikström et al., 2001; Wang et al., 2009; Bell et al., 2010; Zanne et

al., 2014; Magallón et al., 2015; Zeng et al., 2017). The ages of major rosid lineages

estimated using the two dating methods and the two trees largely overlap with uncertainty

intervals reported in previous studies (Fig 3). However, we did find that some clade ages

(e.g., Celastrales, Crossosomatales, and Picramniales; Fig. 3) estimated in our study are

younger and outside of the range of uncertainty reported in previous studies (Wikström et al.,

2001; Wang et al., 2009; Bell et al., 2010; Zanne et al., 2014; Magallón et al., 2015), even

though their placements agree with earlier studies (e.g., Wang et al., 2009; Magallón et al.,

2015; APG IV, 2016). This discrepancy is likely due to poor taxon sampling of smaller rosid

orders in previous studies; particularly, inclusion of a single species (e.g., Picramniales in the

dating analyses of Zanne et al. 2014; Magallón et al. 2015) only allows stem age estimation.

The greatly increased rosid sampling in the present study compared to earlier investigations

(e.g., Wang et al., 2009; Bell et al., 2010; Magallón et al., 2015) could impact age estimation.

Hence, we favor the results from treePL (both 4- and 5-locus; Table 2, Fig 2b). Although

differing in overall scaling, estimated divergence times for all nodes estimated from treePL

and PATHd8 were highly correlated ($R^2 = 0.762$).

IV. Discussion

Our 5-locus supermatrix represents the most comprehensive DNA data set yet compiled for the rosid clade. However, even with 19,740 ingroup species (out of 114,477 species estimated from OTT), our matrix is far from complete. Only 30,234 species of rosids (ca. 34% estimated from Hinchliff et al., 2015) have any type of DNA data in GenBank (see also Folk et al., 2018), and after a series of filtering steps, our topology represents only 16.25% of all rosid species recorded in OTT (Table 1). This relative sampling level (less than 20%) typifies most major clades of flowering plants (Eiserhardt et al., 2018; Folk et al., 2018).

Taxon sampling within rosids exhibits a strong phylogenetic bias (Fig. 1) in accumulation of molecular data ($p \approx 0$). Among large families (> 1,000 species), the five with the poorest sampling (Polygalaceae, Myrtaceae, Rosaceae, Phyllanthaceae, and Malvaceae) have only 3.14% to 13.97% of species with at least one of the five loci sampled here in GenBank after matrix assembly and cleaning (cf. Table S2). For the five best-sampled families (Euphorbiaceae, Fabaceae, Passifloraceae, Brassicaceae, Cucurbitaceae), only 21.95% to 40.56% of species have one of the five loci. Hence, no large (> 1,000 species) family of rosids exceeds 45% species coverage, and most are below 30% coverage (Table S2), consistent with molecular sampling patterns across the angiosperms (Eiserhardt et al. 2018, Folk et al., 2018).

Our 5-locus topology is generally in close agreement with previous work (Wang et al., 2009; Soltis et al., 2011; Ruhfel et al., 2014; Sun et al., 2016), but with better overall resolution and without any cases of non-monophyletic families in the total-evidence trees (Fig. S1). Although the overall support across our tree is lower than that obtained in a previously published GenBank mining effort (Sun et al., 2016; Fig 2a), this is not surprising because: (1) studies have shown that BS values tend globally to decrease as the number of taxa increases (e.g., Sanderson & Donoghue, 1996; Sanderson & Wojciechowski, 2000; Soltis & Soltis, 2003; here the 5-locus tree has ~ 2.2 fold the species sampling of the 4-locus tree); and (2) the standard phylogenetic bootstrap method (e.g., as implemented by RAxML) tends to yield particularly low support for deep branches with large sampling scales (Lemoine et al., 2018).

While our ITS tree yielded low overall resolution for relationships within the rosids, adding ITS data improved the phylogenetic resolution within 14 families compared to that obtained in the 4-locus tree in Sun et al. (2016; Table S3). Additionally, some nodes remain

unresolved across data partitions (e.g., placement of Zygophyllales, Myrtales, and Vitales; Table S3), which likely reflect the rapid radiation of the rosid clade (Zhang et al., 2012, 2016; Zhao et al., 2016; Zeng et al., 2017; Green Plant Consortium, submitted). The divergence times here are generally consistent within methods and broadly congruent with the previous literature (Fig. 3). Between the two estimation methods, divergence times from PATHd8 were younger than those from treePL (Figs. 2b and 3), but highly correlated.

Despite the importance of the rosids to terrestrial landscapes, our knowledge of this clade remains limited (Folk et al., 2018), with species sampling gaps and bias that have likewise persisted across flowering plants and in many other major clades of life (Smith & Brown, 2018). Transparently assessing and quantifying data gaps is crucial for studies using large biodiversity data sets (Folk et al., 2018). Our exploration of phylogenetic sampling here highlights the essential role of taxonomic resources like Open Tree for assessing sampling gaps. We anticipate that similar analyses of cladewise sampling and explicit tests of phylogenetic bias will become standard approaches in large-scale studies as these become more numerous, and as discussion continues over their construction and use (e.g., Rabosky, 2015; Beaulieu & O'Meara, 2018; Folk et al., 2018; Donoghue & Edwards, 2019).

While limitations in sampling continue to hinder our understanding of angiosperm evolution, growth of sequence databases continues to be rapid, and our updated supermatrix effort has increased rosid species coverage by more than two-fold while recovering a backbone that is largely robust. Given the importance of deeply sampled global phylogenies for comparative biology (see Folk et al., 2018; Beaulieu & O'Meara, 2018; Smith & Brown, 2018; Allen et al., 2019), the data set we have assembled here will be an important resource for macroevolutionary synthesis across a globally important clade.

Acknowledgements

This work was supported by the National Science Foundation (DEB-1208809 to D.E.S.), Dimensions of Biodiversity US-China (DEB-1442280 to P.S.S. and D.E.S.), ABI Innovation (DBI-1458640 to P.S.S. and D.E.S.), and National Natural Science Foundation of China (Grant no. 31590822 to Z.D.C.). We thank Dr. Greg Stull for valuable suggestions on the choice of fossil constraints, and Dr. Mark Miller and Dr. Pfeiffer Wayne from Cyber-Infrastructure for Phylogenetic Research (CIPRES) Science Gateway for their extended computation support. We thank the staff at the HiPerGator cluster at the University of Florida and CIPRES for providing us extensive computational resources.

Author contributions

The authors declare no conflict interests. D.E.S., P.S.S., Z.D.C., and M.S. designed the study; M.S., S.A.S., and M.A.G. conducted GenBank data mining; M.S. and C.G.-A. performed OpenTree mapping; M.S. did the phylogeny and dating analyses; M.S., C.G.-A., and D.E.S. drafted the manuscript; R.A.F., D.E.S., S.A.S., M.A.G., P.S.S., Z.D.C., and R.P.G. revised the manuscript. All authors contributed to and approved the final manuscript.

References

354 Allen J, Folk RA, Soltis PS, Soltis DE, Guralnick RP. 2019. Big data and biodiversity: Big
355 challenges and broad applications. *Nature Plants* 5: 11–13.

356
Anderson CL. 2007. *Dating divergence times in phylogenies*. Ph.D. Dissertation. Uppsala,
Sweden: Uppsala University.

APG III. 2009. An update of the Angiosperm Phylogeny Group classification for the orders
and families of flowering plants: APG III. *Botanical Journal of the Linnean Society*
161: 105–121.

APG IV. 2016. An update of the Angiosperm Phylogeny Group classification for the orders
and families of flowering plants: APG IV. *Botanical journal of the Linnean Society*
181: 1–20.

Arabidopsis Genome Initiative. 2000. Analysis of the genome sequence of the flowering
plant *Arabidopsis thaliana*. *Nature* 408: 796.

Barniske AM, Borsch T, Mueller K, Krug M, Worberg A, Neinhuis C, Quandt D. 2012.
Phylogenetics of early branching eudicots: comparing phylogenetic signal across
plastid introns, spacers, and genes. *Journal of Systematics and Evolution* 50: 85–108.

Beaulieu JM, O'Meara BC. 2018. Can we build it? Yes we can, but should we use it?
Assessing the quality and value of a very large phylogeny of campanulid
angiosperms. *American Journal of Botany* 105: 417–432.

Bell C, Soltis D, Soltis P. 2010. The age and diversification of the angiosperms re-revisited.
American Journal of Botany 97: 1296–1303.

Bininda-Emonds OR, Cardillo M, Jones KE, MacPhee RD, Beck RM, Grenyer R, Price SA,
Vos RA, Gittleman JL, Purvis A. 2007. The delayed rise of present-day mammals.
Nature 446: 507.

Boyce CK, Lee JE, Feild TS, Brodribb TJ, Zwieniecki MA. 2010. Angiosperms helped put
the rain in the rainforests: The impact of plant physiological evolution on tropical
biodiversity. *Annals of the Missouri Botanical Garden* 97: 527–540.

Britton T, Anderson CL, Jaquet D, Lundqvist S, Bremer K. 2006. PATHd8—a new method
for estimating divergence times in large phylogenetic trees without a molecular clock
[online]. Available from www.math.su.se/PATHd8 [accessed 1 May 2017].

Cantino PD, Doyle JA, Graham SW, Judd WS, Olmstead RG, Soltis DE, Soltis PS,
Donoghue MJ. 2007. Towards a phylogenetic nomenclature of *Tracheophyta*. *Taxon*
56: 822–846.

Cayuela L, Granzow-de la CÍ, Albuquerque FS, Golicher DJ. 2012. Taxonstand: An R

package for species names standardisation in vegetation databases. *Methods in*

Ecology and Evolution 3: 1078–1083.

Davis CC, Wurdack KJ. 2004. Host-to-parasite gene transfer in flowering plants:

phylogenetic evidence from Malpighiales. *Science* 305: 676–678.

Davis CC, Webb CO, Wurdack KJ, Jaramillo CA, Donoghue MJ. 2005. Explosive radiation

of Malpighiales supports a Mid-Cretaceous origin of modern tropical rain forests. *The*

American Naturalist 165: E36–E65.

Donoghue MJ, Edwards E J. 2019. Model clades are vital for comparative biology, and

ascertainment bias is not a problem in practice: a response to Beaulieu and O'Meara.

American Journal of Botany 106: 327–330.

Edger PP, Heidel-Fischer HM, Bekaert M, Rota J, Glöckner G, Platts AE, Heckel DG, Der

JP, Wafula EK, Tang M *et al.* 2015. The butterfly plant arms-race escalated by gene

and genome duplications. *Proceedings of the National Academy of Sciences, USA*

112: 8362–8366.

Edgar RC. 2004. MUSCLE: A multiple sequence alignment method with reduced time and

space complexity. *BMC Bioinformatics* 5: 113.

Eiserhardt WL, Antonelli A, Bennett DJ, Botigué LR, Burleigh JG, Dodsworth S, Enquist

BJ, et al. 2018. A roadmap for global synthesis of the plant tree of life. *American*

Journal of Botany 105: 614–622.

Ericson PG, Anderson CL, Britton T, Elzanowski A, Johansson US, Källersjö M, Ohlson JI,

Parsons TJ, Zuccon D, Mayr G. 2006. Diversification of Neoaves: integration of

molecular sequence data and fossils. *Biology Letters* 2: 543–547.

Farr ER, Zijlstra G. 1996. Index Nominum Genericorum (Plantarum) [online]. Available

from http: //botany.si.edu/ing/ [accessed 16 April 2017].

Farrell BD. 1998. "Inordinate fondness" explained: Why are there so many beetles? *Science*

281: 555.

Feldberg K, Schneider H, Stadler T, Schäfer-Verwimp A, Schmidt AR, Heinrichs J. 2014.

Epiphytic leafy liverworts diversified in angiosperm-dominated forests. *Scientific*

Reports 4: 5974.

Folk RA, Sun M, Soltis PS, Smith SA, Soltis DE, Guralnick RP. 2018. Challenges of

comprehensive taxon sampling in comparative biology: Wrestling with rosids.

American Journal of Botany 105:433–445. doi: 10.1002/ajb2.1059

Green Plant Consortium (One Thousand Plant Transcriptomes Initiative). 2018. A

phylogenomic view of evolutionary complexity in green plants. *Nature*, in review.

Govaert R. 2001. How many species of seed plants are there? *Taxon* 50: 1085-1090.

Hibbett DS, Matheny PB. 2009. The relative ages of ectomycorrhizal mushrooms and their

plant hosts estimated using Bayesian relaxed molecular clock analyses. *BMC Biology*

7: 1.

Hinchliff CE, Smith SA. 2014. Some limitations of public sequence data for phylogenetic

inference (in plants). *PLoS ONE* 9: e98986.

Hinchliff CE, Smith SA, Allman JF, Burleigh JG, Chaudhary R, Coghill LM, Crandall KA,

Deng J, Drew BT, Gazis R. 2015. Synthesis of phylogeny and taxonomy into a

comprehensive tree of life. *Proceedings National Academy of Sciences, USA* 112:

12764–12769.

Huson DH, Scornavacca C. 2012. Dendroscope 3: an interactive tool for rooted phylogenetic

trees and networks. *Systematic Biology* 61: 1061–1067.

Lemoine F, Domelevo Entfellner JB, Wilkinson E, Correia D, Dávila Felipe M, De Oliveira

T, Gascuel O. 2018. Renewing Felsenstein's phylogenetic bootstrap in the era of big

data. *Nature* 556: 452–456.

Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and

annotation of phylogenetic and other trees. *Nucleic Acids Research* 44: W242–W245.

Li HL, Wang W, Mortimer PE, Li RQ, Li DZ, Hyde KD, Xu JC, Soltis DE, Chen, Z. D.

2015. Large-scale phylogenetic analyses reveal multiple gains of actinorhizal

nitrogen-fixing symbioses in angiosperms associated with climate change. *Scientific*

Reports 5: 14023.

Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. 2012. The global diversity of birds

in space and time. *Nature* 491: 444–448.

Junier T, Zdobnov EM. 2010. The Newick utilities: high-throughput phylogenetic tree

processing in the UNIX shell. *Bioinformatics* 26: 1669–1670.

Katoh K, Kuma K, Toh H, Miyata T. 2005. MAFFT version 5: improvement in accuracy of

multiple sequence alignment. *Nucleic Acids Research* 33: 511–518.

Kück P, Meusemann K. 2010. FASconCAT: convenient handling of data matrices.

Molecular Phylogenetics and Evolution 56: 1115–1118.

Magallón S, Crane PR, Herendeen PS. 1999. Phylogenetic pattern, diversity, and

diversification of eudicots. *Annals of the Missouri Botanical Garden* 86: 297–372.

Magallón S, Gómez-Acevedo S, Sánchez-Reyes LL, Hernández-Hernández T. 2015. A

meta-calibrated timetree documents the early rise of flowering plant phylogenetic

diversity. *New Phytologist* 207: 437–453.

Moreau CS, Bell CD, Vila R, Archibald SB, Pierce NE. 2006. Phylogeny of the ants:

diversification in the age of angiosperms. *Science* 312: 101–104.

Moreau CS, Bell CD. 2013. Testing the museum versus cradle tropical biological diversity

hypothesis: phylogeny, diversification, and ancestral biogeographic range evolution of

the ants. *Evolution* 67: 2240–2257.

Moore MJ, Soltis PS, Bell CD, Burleigh JG, Soltis DE. 2010. Phylogenetic analysis of 83

plastid genes further resolved the early diversification of eudicots. *Proceedings of the*

National Academy of Sciences, USA 107: 4623–4628.

Pattengale ND, Alipour M, Bininda-Emonds OR, Moret BM, Stamatakis A. 2010a. How

many bootstrap replicates are necessary? *Journal of Computational Biology* 17: 337–

354.

Pattengale ND, Swenson KM, Moret BME. 2010b. Uncovering hidden phylogenetic

consensus. In: Borodovsky M, Gogarten JPeter, Przytycka TM, Rajasekaran S eds.

Bioinformatics research and applications. Berlin, Germany: Springer-Verlag. 128–

139.

Pennell MW, Eastman JM, Slater GJ, Brown JW, Uyeda JC, FitzJohn RG, Alfaro ME,

Harmon LJ. 2014. geiger v2.0: an expanded suite of methods for fitting

macroevolutionary models to phylogenetic trees. *Bioinformatics* 15: 2216–2218.

Rabosky DL. 2015. No substitute for real data: A cautionary note on the use of phylogenies

from birth–death polytomy resolvers for downstream comparative analyses. *Evolution*

69: 3207–3216.

Ricklefs RE. 2007. Estimating diversification rates from phylogenetic information. *Trends*

in Ecology & Evolution 22: 601–610.

Rodman JE, Soltis PS, Soltis DE, Sytsma K, Karol K. 1998. Parallel evolution of

glucosinolate biosynthesis inferred from congruent nuclear and plastid gene

phylogenies. *American Journal of Botany* 85: 997–997.

Roelants K, Gower DJ, Wilkinson M, Loader SP, Biju SD,Guillaume K, Moriau L, Bossuyt

F. 2007. Global patterns of diversification in the history of modern amphibians.

Proceedings of the National Academy of Sciences, USA 104: 887–892.

Ruhfel BR, Gitzendanner MA, Soltis DE, Soltis PS, Burleigh JG. 2014. From algae to

angiosperms – inferring the phylogeny of green plants (*Viridiplantae*) from 360

plastid genomes. *BMC Evolutionary Biology* 14: 23.

Sanderson MJ, Donoghue MJ. 1996. The relationship between homoplasy and confidence in

a phylogenetic tree. In: Sanderson MJ, Hufford L, eds. *Homoplasy: The recurrence of*
similarity in evolution. San Diego, USA: Academic Press. 67–89.

Sanderson MJ, Wojciechowski MF. 2000. Improved bootstrap confidence limits in large-
scale phylogenies with an example from Neo-Astragalus (Leguminosae). *Systematic*
Biology 49: 671–685.

Sato S, Nakamura Y, Kaneko T, Asamizu E, Kato T, Nakao M, Sasamoto S, Watanabe A,
Ono A, Kawashima K et al. 2008. Genome structure of the legume, *Lotus japonicus*.
DNA research 15: 227–239.

Sauquet H, Ho S, Gandolfo M, Jordan G, Wilf P, Cantrill D, Bayly M, Bromham L, Brown
G, Carpenter R. 2012. Testing the impact of calibration on molecular divergence
times using a fossil-rich group: the case of *Nothofagus* (Fagales). *Systematic Biology*
61: 298–313.

Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, Hyten DL, Song Q, Thelen
JJ, Cheng J *et al*. 2010. Genome sequence of the palaeopolyploid soybean. *Nature*
463: 178–183.

Schmutz J, McClean PE, Mamidi S, Wu GA, Cannon SB, Grimwood J, Jenkins J, Shu S,
Song Q, Chavarro C *et al*. 2014. A reference genome for common bean and genome-
wide analysis of dual domestications. *Nature Genetics* 46: 707–713.

Schneider H, Schuettpelz E, Pryer KM, Cranfill R. 2004. Ferns diversified in the shadow of
angiosperms. *Nature* 428: 553.

Smith SA, Beaulieu JM, Donoghue MJ 2009. Mega-phylogeny approach for comparative
biology: An alternative to supertree and supermatrix approaches. *BMC Evolution*
Biology 9: 37.

Smith SA, Beaulieu JM, Donoghue M. 2010. An uncorrelated relaxed-clock analysis
suggests an earlier origin for flowering plants. *Proceedings of the National Academy*
of Sciences, USA 107: 5897–5902.

Smith SA, Beaulieu JM, Stamatakis A, Donoghue MJ. 2011. Understanding Angiosperm
diversification using small and large phylogenetic trees. *American Journal of Botany*
98: 404–414.

Smith SA, O'Meara BC. 2012. treePL: divergence time estimation using penalized
likelihood for large phylogenies. *Bioinformatics* 28: 689–2690.

Smith SA, Brown JW. 2018. Constructing a broadly inclusive seed plant phylogeny.
American Journal of Botany 105: 1–13. doi: 10.1002/ajb2.1019

Soltis DE, Soltis PS, Morgan DR, Swensen SM, Mullin BC, Dowd JM, Martin P G. 1995.

Chloroplast gene sequence data suggest a single origin of the predisposition for
symbiotic nitrogen fixation in angiosperms. *Proceedings of the National Academy of*
Sciences, USA 92: 2647–2651.

Soltis DE, Soltis PS, Endress PK, Chase MW. 2005. *Phylogeny and evolution of*
angiosperms. Sunderland, MA, USA: Sinauer Associates.

Soltis DE, Gitzendanner MA, Soltis PS. 2007. A 567-taxon data set for angiosperms: The
challenges posed by Bayesian analyses of large data sets. *International Journal of*
Plant Sciences 168: 137–157.

Soltis DE, Moore MJ, Burleigh JG, Bell CD, Soltis PS. 2010. Assembling the angiosperm
tree of life: progress and future prospects. *Annals of the Missouri Botanical Garden*
97: 514–526.

Soltis DE, Smith SA, Cellinese N, Wurdack KJ, Tank DC, Brockington SF, Refulio-
Rodriguez NF, Walker JB, Moore MJ, Carlsward BS *et al*. 2011. Angiosperm
phylogeny: 17 genes, 640 taxa. *American Journal of Botany* 98: 704–730.

Soltis DE, Soltis PS. 2016. Mobilizing and integrating big data in studies of spatial and
phylogenetic patterns of biodiversity. *Plant Diversity* 38: 264–270.

Soltis PS, Soltis DE. 2003. Applying the bootstrap in phylogeny reconstruction. *Statistical*
Science 8: 256–267.

Soltis PS, Soltis DE. 2004. The origin and diversification of angiosperms. *American Journal*
of Botany 91: 1614–1626.

Soltis PS, Soltis DE, Chase MW, Endress PK, Crane PR. 2004. The diversification of
flowering plants. In: Cracraft J, Donoghue M, eds. *The tree of life*. Oxford, UK &
New York, USA: Oxford University Press. 154–167.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* 30: 1312–1313.

Stevens PF. 2001 onwards. Angiosperm phylogeny website, Version 14, July 2017 [online].
Available from http://www.mobot.org/MOBOT/research/APweb/ [accessed 1 January
2018].

Sun M, Soltis DE, Soltis PS, Zhu X, Burleigh JG, Chen ZD. 2015. Deep phylogenetic
incongruence in the angiosperm clade Rosidae. *Molecular Phylogenetics and*
Evolution 83: 156–166.

Sun M, Naeem R, Su JX, Cao ZY, J. Burleigh G, Soltis PS, Soltis DE, Chen ZD. 2016.
Phylogeny of the *Rosidae*: A dense taxon sampling analysis. *Journal of Systematic*
and Evolution 54: 363–391.

Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular

558   Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution* 30:

559   2725–2729.

560   Testo W, Sundue M. 2016. A 4000-species dataset provides new insight into the evolution

561   of ferns. *Molecular Phylogenetics and Evolution* 105: 200–211.

562   Thomas GH, Hartmann K, Jetz W, Joy JB, Mimoto A, Mooers AO. 2013. PASTIS: an R

563   package to facilitate phylogenetic assembly with soft taxonomic inferences. *Methods*

564   *in Ecology and Evolution* 4: 1011–1017. doi:10.1111/2041-210X.12117

565   Varshney RK, Chen WB, Li YP, Bharti AK, Saxena RK, Schlueter JA, Donoghue MTA,

566   Azam S, Fan GY, Whaley AM *et al*. 2012. Draft genome sequence of pigeonpea

567   (*Cajanus cajan*), an orphan legume crop of resource-poor farmers. *Nature*

568   *Biotechnology* 30: 83–89.

569   Varshney RK, Song C, Saxena RK, Azam S, Yu S, Sharpe AG, Cannon S, Baek J, Rosen

570   BD, Tar'an B *et al*. 2013. Draft genome sequence of chickpea (*Cicer arietinum*)

571   provides a resource for trait improvement. *Nature biotechnology* 31: 240–246.

572   Wang H, Moore MJ, Soltis PS, Bell CD, Brockington SF, Alexandre R, Davis CC, Latvis

573   M, Manchester SR, Soltis DE. 2009. Rosid radiation and the rapid rise of angiosperm-

574   dominated forests. *Proceedings of the National Academy of Sciences, USA* 106:

575   3853–3858.

576   Wang X, Wang H, Wang J, Sun R, Wu J, Liu S, Bai Y, Mun J, Bancroft I, Cheng F *et al*.

577   2011. The genome of the mesopolyploid crop species *Brassica rapa*. *Nature Genetics*

578   43: 1035–1039.

579   Watkins JJE, Cardelús CL. 2012. Ferns in an angiosperm world: Cretaceous radiation into

580   the epiphytic niche and diversification on the forest floor. *International Journal of*

581   *Plant Sciences* 173: 695–710.

582   Wikström N, Savolainen V, Chase MW. 2001. Evolution of the angiosperms: Calibrating the

583   family tree. *Proceedings of the Royal Society of London, B, Biological Sciences* 268:

584   2211–2220.

585   Wilf P, Labandeira CC, Kress WJ, Staines CL, Windsor DM, Allen AL, Johnson KR. 2000.

586   Timing the radiations of leaf beetles: Hispines on gingers from latest Cretaceous to

587   recent. *Science* 289: 291–294.

588   Worberg A, Quandt D, Barnkske AM, Lohne C, Hilu KW, Borsch T. 2007. Phylogeny of

589   basal eudicots: Insights from non-coding and rapidly evolving DNA. *Organisms*

590   *Diversity and Evolution* 7: 55–77.

Xi Z, Ruhfel BR, Schaefer H, Amorim AM, Sugumaran M, Wurdack KJ, Endress PK,

Matthews M, Stevens PF, Mathews S *et al*. 2012. Phylogenomics and a posteriori data

partitioning resolve the Cretaceous angiosperm radiation in Malpighiales.

Proceedings National Academy of Sciences, USA 109: 17519–1752.

Young ND, Debellé F, Oldroyd GE, Geurts R, Cannon SB, Udvardi MK, Benedito VA,

Mayer KFX, Gouzy J, Schoof H *et al*. 2011. The *Medicago* genome provides insight

into the evolution of rhizobial symbioses. *Nature* 480: 520–524.

Zanne AE, Tank DC, Cornwell WK, Eastman JM, Smith SA, FitzJohn RG, McGlinn DJ,

O'Meara BC, Moles AT, Reich PB *et al*. 2014. Three keys to the radiation of

angiosperms into freezing environments. *Nature* 506: 89–92.

Zeng L, Zhang N, Zhang Q, Endress PK, Huang J, Ma H. 2017. Resolution of deep eudicot

phylogeny and their temporal diversification using nuclear genes from transcriptomic

and genomic datasets. *New Phytologist* 214: 1338–1354.

Zhang N, Zeng LP, Shan HY, Ma H. 2012. Highly conserved low-copy nuclear genes as

effective markers for phylogenetic analyses in angiosperms. *New Phytologists* 195:

923–937.

Zhang N, Wen J, Zimmer EA. 2016. Another look at the phylogenetic position of the grape

order Vitales: Chloroplast phylogenomics with an expanded sampling of key lineages,

Molecular Phylogenetics and Evolution 101: 216–223.

Zhao L, Li X, Zhang N, Zhang SD, Yi TS, Ma H, Guo ZH, Li DZ. 2016. Phylogenomic

analyses of large-scale nuclear genes provide new insights into the evolutionary

relationships within the rosids. *Molecular Phylogenetics and Evolution* 105: 166–176.

Zhu XY, Chase MW, Qiu YL, Kong HZ, Dilcher DL, Li JH, Chen ZD. 2007. *MatR*

sequences help to resolve deep phylogenetic relationships in rosids. *BMC*

Evolutionary Biology 7: 217.

Table 1. Ordinal-level summary sampling table for the 5-locus rosid supermatrix ("Matrix") compared to the rosid clade of the Open Tree Taxonomy ("OTT") database v. 3.0 (https://devtree.opentreeoflife.org/about/taxonomy-version/ott3.0; Hinchliff et al., 2015) and matching taxon names between these data sets. Orders follow APG IV (2016). A summary table at the family level is available in Table S2.

Order	Match (Matrix genera)/OTT genera	Matched genus %	Match (Matrix species)/OTT species	Matched species %
Brassicales	357(369)/502	71.12%	1693(1843)/5940	28.50%
Celastrales	68(69)/111	61.26%	272(281)/1499	18.15%
Crossosomatales	13(13)/14	92.86%	24(24)/82	29.27%
Cucurbitales	117(125)/133	87.97%	823(863)/3094	26.60%
Fabales	657(668)/864	76.04%	5311(5678)/24191	21.95%
Fagales	36(38)/74	48.65%	496(540)/2263	21.92%
Geraniales	15(18)/20	75.00%	295(305)/962	30.67%
Huerteales	6(6)/6	100.00%	7(7)/30	23.33%
Malpighiales	586(596)/891	65.77%	3703(3868)/21316	17.37%
Malvales	255(257)/405	62.96%	1285(1349)/7771	16.54%
Myrtales	257(263)/475	54.11%	1286(1373)/15529	8.28%
Oxalidales	43(45)/69	62.32%	182(193)/2207	8.25%
Picramniales	2(2)/3	66.67%	5(5)/57	8.77%
Rosales	217(227)/359	60.45%	1694(1787)/20620	8.22%
Sapindales	342(347)/551	62.07%	1361(1434)/7421	18.34%
Vitales	9(9)/15	60.00%	110(119)/1155	9.52%
Zygophyllales	18(18)/27	66.67%	60(71)/340	17.65%
Total	2998(3070)/4519	66.34%	18607(19740)/114477	16.25%

Table 2. Summary table of ages estimated for rosid major clades by treePL and PATHd8 based on the tree inferred here ("5-locus") and that inferred in Sun et al. 2016 ("4-locus"). The age unit is million years ago (Myr).

Clade	5-locus		4-locus	
	Age treePL	Age PATHd8	Age treePL	Age PATHd8
Fabids	117.03	89.80	121.11	94.95
Fagales	95.60	83.50	100.83	83.50
Betulaceae	60.64	59.80	59.80	59.80
Casuarinaceae	27.57	46.63	27.06	41.62
Fagaceae	66.27	61.12	67.04	69.46
Juglandaceae	72.55	69.39	69.66	66.60
Myricaceae	30.08	27.63	26.18	27.97
Nothofagaceae	18.53	11.42	18.26	11.30
Ticodendraceae	66.67	59.80	64.52	63.23
Cucurbitales	111.95	65.06	113.39	53.48
Anisophylleaceae	48.13	35.30	49.44	39.52
Apodanthaceae	70.73	65.06	74.69	48.60
Begoniaceae	52.11	35.40	23.74	20.39
Cucurbitaceae	34.91	28.17	42.07	29.20
Tetramelaceae	19.21	9.01	22.57	12.15
Corynocarpaceae	6.95	3.62	6.96	2.86
Coriariaceae	14.04	7.82	22.97	10.15
Datiscaceae	8.47	4.54	9.54	5.24
Rosales	107.52	89.80	111.03	89.80
Cannabaceae	67.73	65.50	67.26	65.50
Elaeagnaceae	27.41	20.06	28.70	27.79
Moraceae	46.03	28.78	40.49	16.31

Rhamnaceae	74.55	48.60	80.11	61.70
Rosaceae	89.80	89.80	89.80	89.80
Ulmaceae	59.41	32.53	88.93	21.49
Urticaceae	60.28	51.05	68.70	33.25
Dirachmaceae	88.51	70.60	96.98	70.60
Barbeyaceae	84.47	53.33	90.03	70.60
Fabales	96.58	59.90	105.41	75.15
Fabaceae	83.90	55.80	93.77	65.37
Polygalaceae	60.89	55.80	72.51	13.46
Surianaceae	47.39	15.50	53.52	21.96
Quillajaceae	90.75	26.59	98.52	36.49
Malpighiales	114.67	89.30	114.47	89.30
Achariaceae	54.72	14.37	64.21	25.88
Bonnetiaceae	70.08	58.52	71.27	58.67
Calophyllaceae	55.73	21.85	47.53	17.35
Caryocaraceae	20.12	5.20	22.18	8.73
Centroplacaceae	74.12	15.13	77.14	23.34
Chrysobalanaceae	22.13	10.69	27.61	16.17
Clusiaceae	52.63	27.13	52.05	22.91
Dichapetalaceae	17.65	5.96	22.97	10.87
Elatinaceae	50.37	17.24	51.62	21.99
Erythroxylaceae	46.31	17.55	55.76	26.26
Euphorbiaceae	63.30	38.14	99.38	41.95
Humiriaceae	33.90	33.90	33.90	33.90
Hypericaceae	81.66	63.84	81.46	63.49
Irvingiaceae	5.19	1.57	6.10	2.61
Ixonanthaceae	23.76	8.15	31.28	13.56
Lacistemataceae	7.82	2.19	9.54	3.73

Linaceae	58.84	25.10	60.48	37.36
Malpighiaceae	53.17	33.00	47.75	33.00
Ochnaceae	51.42	15.36	54.39	25.49
Pandaceae	38.67	8.97	42.54	15.19
Passifloraceae	63.15	33.74	83.25	3.24
Peraceae	53.84	13.68	68.14	22.94
Phyllanthaceae	68.83	28.13	101.43	50.67
Picrodendraceae	35.15	12.86	44.13	21.99
Podostemaceae	80.18	89.30	82.99	89.30
Putranjivaceae	18.11	6.86	16.72	6.28
Rafflesiaceae	81.48	41.45	70.24	41.13
Rhizophoraceae	59.88	33.90	59.74	33.90
Salicaceae	64.37	48.00	71.33	48.00
Trigoniaceae	38.07	15.32	50.40	26.28
Violaceae	57.17	27.34	68.43	47.89
Balanopaceae	17.87	3.61	24.51	6.52
Ctenolophonaceae	80.30	34.51	109.65	52.48
Euphroniaceae	46.86	23.35	64.26	37.35
Goupiaceae	75.00	32.02	87.00	54.34
Lophopyxidaceae	60.81	19.39	77.63	31.51
Celastrales	89.44	45.93	94.99	54.88
Lepidobotryaceae	12.11	3.25	12.84	4.99
Celastraceae	67.89	37.82	76.39	38.90
Oxalidales	112.85	79.20	114.28	79.20
Connaraceae	35.38	6.59	24.76	10.10
Cunoniaceae	43.50	79.20	32.67	65.56
Elaeocarpaceae	61.70	61.70	61.70	61.70
Huaceae	14.23	4.30	16.56	6.62

Oxalidaceae	52.72	21.00	46.59	30.05
Cephalotaceae	77.57	53.13	70.61	79.20
Brunelliaceae	2.65	3.41	3.71	3.27
Zygophyllales	96.67	42.02	96.95	74.54
Zygophyllaceae	91.38	41.07	90.94	69.52
Krameriaceae	31.36	8.25	18.16	7.87
Malvids	116.49	89.30	120.33	91.47
Brassicales	86.82	86.66	95.92	89.30
Akaniaceae	4.10	1.86	3.07	1.19
Brassicaceae	36.20	30.70	45.27	32.55
Capparaceae	38.94	17.68	43.31	22.21
Caricaceae	26.58	13.58	28.74	16.33
Cleomaceae	38.58	24.01	47.85	29.53
Gyrostemonaceae	6.80	3.34	9.87	5.39
Limnanthaceae	13.78	8.26	11.54	7.52
Resedaceae	53.78	44.91	58.24	43.56
Salvadoraceae	26.97	14.59	40.37	24.73
Tropaeolaceae	40.20	20.73	13.06	7.64
Bataceae	32.77	20.86	36.62	24.73
Emblingiaceae	52.88	24.59	61.28	25.27
Koeberliniaceae	67.42	50.48	77.00	55.39
Moringaceae	12.17	5.91	14.61	6.99
Pentadiplandraceae	63.01	62.42	70.52	70.43
Setchellanthaceae	78.46	79.01	87.32	89.30
Tovariaceae	52.88	24.59	61.28	25.27
Malvales	100.59	69.12	101.97	68.91
Bixaceae	50.29	24.36	54.37	26.98
Cistaceae	45.68	34.47	50.40	45.35

Cytinaceae	73.43	67.35	73.58	55.17
Dipterocarpaceae	35.80	17.07	36.05	19.22
Malvaceae	50.85	36.94	44.04	25.61
Muntingiaceae	42.72	21.12	49.82	25.44
Neuradaceae	10.59	5.59	11.18	5.71
Sarcolaenaceae	12.94	6.04	13.37	6.31
Sphaerosepalaceae	11.85	4.89	13.20	5.20
Thymelaeaceae	48.55	40.20	47.79	12.59
Picramniales/Picramniaceae	36.78	9.65	36.91	14.80
Huerteales	61.57	37.20	67.93	37.20
Dipentodontaceae	38.06	10.16	37.79	15.23
Tapisciaceae	37.20	37.20	37.20	37.20
Petenaeaceae	56.88	14.16	61.53	22.34
Gerrardinaceae	56.88	14.16	61.53	22.34
Sapindales	88.96	65.50	82.21	65.50
Anacardiaceae	45.59	32.35	50.69	47.00
Burseraceae	43.87	31.43	31.32	32.09
Meliaceae	48.60	48.60	49.04	48.60
Nitrariaceae	62.91	17.63	32.49	16.91
Rutaceae	65.50	65.50	65.50	65.50
Sapindaceae	64.76	55.80	60.52	55.80
Simaroubaceae	46.19	12.04	47.98	21.83
Kirkiaceae	3.46	0.92	3.97	1.49
Biebersteiniaceae	19.11	5.12	17.58	7.59
Crossosomatales	88.66	28.40	90.75	34.68
Crossosomataceae	17.62	4.88	16.78	8.65
Staphyleaceae	28.40	28.40	28.40	28.40
Strasburgeriaceae	20.74	4.68	23.87	8.25

Aphloiaceae	75.92	15.79	78.94	27.26
Geissolomataceae	46.62	9.63	56.37	17.05
Guamatelaceae	40.90	6.30	31.80	10.86
Stachyuraceae	2.84	0.63	3.67	1.01
Geraniales + Myrtales	116.49	88.20	121.39	96.30
Geraniales	107.69	47.69	117.77	81.03
Geraniaceae	74.66	32.08	97.14	77.94
Francoaceae	97.34	29.59	104.62	36.60
Myrtales	95.00	88.20	95.00	88.20
Combretaceae	51.58	16.24	42.78	12.45
Crypteroniaceae	18.04	9.91	17.77	9.68
Lythraceae	72.66	70.60	71.98	70.60
Melastomataceae	49.25	51.74	47.35	60.23
Myrtaceae	64.10	55.80	64.27	55.80
Onagraceae	49.56	40.09	60.65	53.07
Penaeaceae	34.38	17.22	26.29	21.19
Vochysiaceae	39.18	26.95	36.65	27.24
Alzateaceae	52.22	26.13	32.38	26.29
Vitales/Vitaceae	75.77	66.00	78.27	66.37
Rosids	117.93	89.80	122.62	104.20

Figure Legends:

Fig. 1. The 5-locus rosid phylogeny showing sampling coverage of sequence data for the 5 loci. This comparison shows substantial phylogenetic bias in each of the 5 loci sampled in the rosid matrix based on presence/absence heatmap layers. The five layers are labeled as *matR*, *atpB*, *rbcL*, *matK*, and ITS from inside toward the outer edge. Species with black tips at each layer mean sequence data are available for a specific locus. Each order is labeled and colored, so phylogenetic bias of DNA data can be viewed by the rough distribution of the black and gray tips within and/or among each order (cf. Tables 1, S2 for further percentage details).

Fig. 2. Comparison of phylogenetic resolution (a) and divergence time estimation (b) between the 4-locus, 8,855-taxon rosid phylogeny and the 5-locus, 19,740-taxon rosid phylogeny. BS stands for bootstrap. Orange denotes the treePL method, and blue denotes the PATHd8 method in panel (b).

Fig. 3. Comparison of crown ages for major rosid clades reported in this study, Wikström et al. (2001), Wang et al. (2009), Bell et al. (2010), Zanne et al. (2014), and Magallón et al. (2015). Error bars represent age ranges reported for the given node. Red hollow circles and diamonds stand for ages estimated from the 4-locus tree; solid symbols are estimated from the 5-locus tree; the color bars at the bottom of the plot correspond to the orders in Fig. 1. In several previous studies, only a single species was sampled for small clades such as Picramniales, preventing the estimation of crown ages; in these cases only the stem age is given here.

Fig. 1.

Fig. 2.

Fig. 3.

Supporting Information

Fig. S1 The best ML tree obtained from the RAxML analysis.

Table S1. Information on the 59 rosid calibration constraints used in this study.

Table S2. Family-level summary sampling table for the 5-locus supermatrix ("Matrix") compared to the rosid clade of the Open Tree Taxonomy ("OTT") database v.3.0 (https://devtree.opentreeoflife.org/about/taxonomy-version/ott3.0; Hinchliff et al., 2015) and matched taxon names between these data sets.

Table S3. List of non-monophyletic families in the *matR* and ITS locus trees.