Novel insights into the taxonomic diversity and molecular mechanisms of bacterial Mn(III)

reduction

Nadia Szeinbaum[1,2,6], Brook L. Nunn[3], Amanda R. Cavazos[2], Sean A. Crowe[2,4], Frank J.

Stewart[1,5], Thomas J. DiChristina[1], Christopher T. Reinhard[2,6], and Jennifer B. Glass[2,6]*

[1]School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA, USA

[2]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

[3]Department of Genome Sciences, University of Washington, Seattle, WA, USA

[4]Department of Microbiology & Immunology and Department of Earth, Ocean, & Atmospheric

Sciences, University of British Columbia, Vancouver, Canada

[5]Department of Microbiology and Immunology, Montana State University, Bozeman, Montana

[6]NASA Astrobiology Institute, Alternative Earths Team, Mountain View, CA

*Corresponding author: jennifer.glass@eas.gatech.edu

Running title: Novel undecaheme in Betaproteobacteria

Originality-Significance Statement: Recent observations suggest that Mn(III)-ligand

complexes are geochemically important in diverse aquatic environments. Thus far, microbially-

driven Mn(III) reduction has only been associated with *Gammaproteobacteria* encoding three-

component outer-membrane porin-cytochrome c conduits. Here, we demonstrate that

Betaproteobacteria dominate in abundance and with respect to protein expression during

biologically-mediated Mn(III) reduction in an enrichment culture from an anoxic lacustrine

system. Using metaproteomics, we detect for the first time that *Betaproteobacteria* express a

two-component porin-cytochrome c conduit, and an uncharacterized extracellular undecaheme

(11-heme) c-type cytochrome. Although this is the first definitive report of an undecaheme

within the *Betaproteobacteria,* we find evidence that they are widespread in uncultivated strains.

These results widen the phylogenetic diversity of Mn(III)-reducing bacteria, and provide new

insights into potential molecular mechanisms for soluble Mn(III) reduction.

Summary: Soluble ligand-bound Mn(III) can support anaerobic microbial respiration in diverse

aquatic environments. Thus far, Mn(III) reduction has only been associated with certain

Gammaproteobacteria. Here, we characterized microbial communities enriched from Mn-replete

sediments of Lake Matano, Indonesia. Our results provide the first evidence for biological

reduction of soluble Mn(III) outside the *Gammaproteobacteria*. Metagenome assembly and

binning revealed a novel betaproteobacterium, which we designate "*Candidatus* Dechloromonas

occultata." This organism dominated the enrichment and expressed a porin-cytochrome c

complex typically associated with iron-oxidizing *Betaproteobacteria* and a novel cytochrome c-

rich protein cluster (Occ), including an undecaheme putatively involved in extracellular electron

transfer. This *occ* gene cluster was also detected in diverse aquatic bacteria, including

uncultivated *Betaproteobacteria* from the deep subsurface. These observations provide new

insight into the taxonomic and functional diversity of microbially-driven Mn(III) reduction in

natural environments.

Introduction. Manganese(III) is a strong oxidant with a reduction potential close to molecular oxygen (Kostka et al., 1995). Mn(III) is short-lived and unstable, but its stability is greatly increased when bound to ligands (Luther III et al., 2015). Ligand-bound Mn(III) is often the most abundant dissolved Mn species in sediment porewaters (Madison et al., 2013; Oldham et al., 2019) and soils (Heintze and Mann, 1947), with the potential to facilitate one-electron redox reactions in a variety of biogeochemical cycles (Luther III et al., 2015). Microbes accelerate the oxidation and reduction of Mn by orders of magnitude compared to abiotic mechanisms (Hem, 1963; Diem and Stumm, 1984; Morgan, 2005; Tebo et al., 2005; Learman et al., 2011; Luther et al., 2018; Jung et al., 2020). Yet, despite clear evidence for the environmental importance of Mn(III), knowledge about microbial Mn(III) cycling pathways remains fragmented.

To date, only *Shewanella* spp. (*Gammaproteobacteria*) have been confirmed to respire soluble Mn(III) (Kostka et al., 1995; Szeinbaum et al., 2014). *Shewanella* respire Mn(III) using the Mtr pathway (Szeinbaum et al., 2017), a porin-cytochrome (PCC) conduit that transports electrons across the periplasm for extracellular respiration of Mn(III/IV), Fe(III), and other metals (Richardson et al., 2012; Shi et al., 2016). Many Fe(II)-oxidizing *Betaproteobacteria* also contain PCCs (MtoAB, generally lacking the C subunit), which are proposed to oxidize Fe(II) to Fe(III) by running the PCC in reverse (Emerson et al., 2013; Kato et al., 2015; He et al., 2017). In some metal-reducing *Gammaproteobacteria* and *Deltaproteobacteria*, extracellular undecaheme (11-heme) UndA is thought to play a key functional role in soluble Fe(III) reduction (Fredrickson et al., 2008; Shi et al., 2011; Smith et al., 2013; Yang et al., 2013). UndA's crystal structure shows a surface-exposed heme surrounded by positive charges, which may bind negatively-charged soluble iron chelates (Edwards et al., 2012). Environmental omics suggest that metal reduction by *Betaproteobacteria* may be widespread in the deep subsurface

(Anantharaman et al., 2016; Hernsdorf et al., 2017). However, only a few Fe(III)-reducing

Betaproteobacteria isolates have been characterized to date (Cummings et al., 1999; Finneran et

al., 2003), and little is known about metal reduction pathways in *Betaproteobacteria*.

Manganese reduction coupled to methane (CH_4) oxidation is a novel metabolism only

recently discovered in cultures enriched in Archaea (Ettwig et al., 2016; Leu et al., 2020).

Biological and geochemical evidence suggest that this metabolism may be found in a variety of

environments (Beal et al., 2009; Crowe et al., 2011; Riedinger et al., 2014), including Fe-rich

Lake Matano, Indonesia. In an attempt to explore whether CH_4 can fuel microbial Mn(III)

reduction in enrichments inoculated with sediments from Lake Matano, Indonesia, which has

active and pronounced microbial Mn and CH_4 cycles (Jones et al., 2011), we uncovered a novel

betaproteobacterium as the most dominant and active member of our Mn(III)-reducing

enrichment culture. Our results provide the first evidence for biological reduction of soluble

Mn(III) outside *Gammaproteobacteria* and provide evidence for a new biochemical pathway

involved in extracellular electron transfer.

Results and discussion

Enrichment of Mn(III)-reducing populations. Lake Matano, Indonesia, is a permanently

stratified ultraoligotrophic lake (Crowe et al., 2008). Below its oxic surface waters, Lake

Matano's permanently anoxic and stratified waters are highly enriched in iron and manganese,

and support the activity of Mn cycling organisms with organic carbon and CH_4 as potential

sources of electrons (Crowe et al., 2011; Jones et al., 2011; Kuntz et al., 2015; Sturm et al.,

2019). We designed an enrichment strategy to select for microbes capable of anaerobic CH_4

oxidation coupled to soluble Mn(III) reduction by incubating anoxic Lake Matano sediment

communities with soluble Mn(III)-pyrophosphate as the electron acceptor (with 2% O_2 in a

subset of bottles), and CH_4 as the sole electron donor and carbon source after pre-incubation to

deplete endogenous organic carbon (see **Supporting Information** for enrichment details).

Enrichment cultures were transferred into fresh media after Mn(III) was completely reduced to

Mn(II), for a total of five transfers over 395 days. By the fourth transfer, cultures with CH_4

headspace (with or without 2% O_2) reduced ~80% of soluble Mn(III) compared to ~30% with N_2

headspace (**Fig. 1**). 16S rRNA gene sequences were dominated by *Betaproteobacteria*

(*Rhodocyclales*; 8-35%) and *Deltaproteobacteria* (*Desulfuromonadales*; 13-26%; **Fig. S1**).

$^{13}CH_4$ oxidation to $^{13}CO_2$ was undetectable (**Fig. S2**).

Figure 1. Consumption of Mn(III) in Lake Matano enrichments in the presence and absence of methane.
Sediment-free cultures (transfer 4) from 335 days after the initial enrichment were incubated for 45 days with 1 mM
Mn(III) pyrophosphate as the sole electron acceptor. One set was incubated with Mn^{3+} and 2% O_2. Initial bottle
headspace contained 50% CH_4 + 50% N_2 (black circles), 50% CH_4+48% N_2+2% O_2 (gray circles), 100% N_2 (white
circles), and 50% CH_4+50% N_2 heat killed controls (black triangles). Error bars are standard deviations from
duplicate experiments. Color change from red to clear indicates Mn(III) reduction.
103
Samples for metagenomic and metaproteomic analysis were harvested from the fifth

transfer (**Fig. 1; Fig. S1**). Out of 2,952 proteins identified in the proteome, 90% were assigned to

Betaproteobacteria; of those, 72% mapped to a 99.5% complete metagenome-assembled genome

(MAG; *Rhodocyclales* bacterium GT-UBC; NCBI accession QXPY01000000) with 81-82%

average nucleotide identity (ANI) and phylogenetic affiliation to *Dechloromonas* spp. (**Table**

S1; Fig. S3). This MAG is named here "*Candidatus* Dechloromonas occultata" sp. nov.;

etymology: occultata; (L. fem. adj. 'hidden'). The remaining 10% of proteins mapped to

Deltaproteobacteria; of those, 70% mapped to a nearly complete MAG (*Desulfuromonadales*

bacterium GT-UBC; NCBI accession RHLS01000000) with 80% ANI to *Geobacter*

sulfurreducens. This MAG is named here "*Candidatus* Geobacter occultata".

Cytochrome expression during Mn(III) reduction. Cytochromes containing multiple *c*-

type hemes are key for electron transport during microbial metal transformations, and therefore

also expected to play a role in Mn(III) reduction. Numerous mono-, di-, and multi (>3)-heme

cytochromes (MHCs) were expressed by "*Ca.* D. occultata" in Mn(III)-reducing cultures. Nine

out of 15 MHCs encoded by the "*Ca.* D. occultata" MAG were expressed, including two

decahemes similar to MtoA in Fe(II)-oxidizing *Betaproteobacteria* (**Tables 1, S2, S3; Figs. 2A,**

S4). Several highly expressed MHCs were encoded on a previously unreported 19-gene cluster

with 10 cytochrome-*c* proteins, hereafter *occA-S* (**Table 1; Figs. 2B, S5, S6**). OccP was

predicted to be an extracellular undecaheme protein of ~100 kDa (922 amino acids). "*Ca.*

Dechloromonas occultata" may reduce Mn(III) using the novel extracellular undecaheme OccP

as the terminal Mn(III) reductase. Experimental verification of the function of the putative Occ

complex is currently limited by the scarcity of genetically tractable *Betaproteobacteria.*

Proteins with 40-60% identity to the expressed "*Ca.* D. occultata" OccP protein were

widely distributed in *Betaproteobacteria* from diverse freshwaters and deep subsurface

groundwaters*,* as well as in several *Gammaproteobacteria* and one alphaproteobacterium (**Fig.**

2D; Table S3). Most *occP*-containing bacteria also possessed *mtoA* and denitrification genes

(**Fig. 2D**; **Figs. S7**, **S8**). These results widen the phylogenetic diversity of candidate extracellular

MHCs that may be involved in microbial Mn(III) reduction.

132
Figure 2. Gene arrangement, predicted protein location, and taxonomic distribution of major expressed
respiratory complexes in "*Ca*. D. occultata". **A:** MtoDAB(Y)X porin-cytochrome c electron conduit; **B:** OccA-S;
C: denitrification complexes (Nap, Nir, Nor and cNos); **D:** Occurrence of key marker genes in *Betaproteobacteria*
and *Gammaproteobacteria* with >95% complete genomes that encode OccP. Protein sequences from "*Ca*. D.
occultata" were used as query against a genome database and searched using PSI BLAST. Matches with identities
>40%, query coverage >80% and E values <10[-5] were considered positive. Red fill around genes and proteins
indicate cytochrome-*c* proteins. Black outlines around blue circles in D indicate type I nitrous oxide reductase to
distinguish from blue dots (type II/cytochrome-nitrous oxide reductase). Gray-shaded genes on the *occ* gene cluster
indicate 6-NHL repeat proteins. Protein locations shown are based on P-sort predictions. Numbers above genes
indicate number of CxxCH motifs predicted to bind cytochrome *c*. IM: inner membrane; OM: outer membrane. For
more details, see **Table 1** and **Table S3**.

Heme-copper oxidases in "Ca. D. occultata". "*Ca.* D. occultata" expressed high-affinity

cbb$_3$-type cytochrome c oxidase (CcoNOQP) associated with microaerobic respiration (**Table**

S4). Features of the "*Ca*. D. occultata" *occS* gene product, including conserved histidine residues

(H-94, H-411, and H-413) that bind hemes a and a$_3$, as well as the H-276 residue that binds Cu$_B$

(**Fig. S6**), suggest that OccS may function similarly to CcoN, the terminal heme-copper oxidase

proton pump in aerobic respiration. All identified OccS amino acid sequences lack Cu$_B$ ligands

Y-280 and H-403, and most lack Cu$_B$ ligands H-325 and H-326. OccS sequences also lack polar

and ionizable amino acids that comprise the well-studied D and K channels involved in proton

translocation in characterized cytochrome c oxidases (Blomberg and Siegbahn, 2014), but

contain conserved H, C, E, D, and Y residues that may serve in alternate proton translocation

pathways, similar to those recently discovered in qNOR (Gonska et al., 2018). OccS homologs

were also found in *Azoarcus* spp. and deep subsurface *Betaproteobacteria* (**Fig. S6**).

Expression of denitrification proteins and possible sources of oxidized nitrogen species.

Periplasmic nitrate reductase (NapA), cytochrome nitrite reductase (NirS), and type II atypical

nitrous oxide reductase (cNosZ; **Fig. S7**) were highly expressed by "*Ca.* D. occultata" (**Table 1**).

Expression of the denitrification pathway was not expected because oxidized nitrogen species

were not added to the medium, to which the only nitrogen supplied was 0.2 mM NH$_4$Cl (along

with headspace N$_2$). The close redox potential of Mn^{3+}-pyrophosphate (~0.8 V; Yamaguchi and

Sawyer, 1985)) to oxidized nitrogen species (0.35-0.75 V) at circumneutral pH and the lack of

oxygen in the media could have induced the expression of denitrification genes simultaneously

with Mn(III)-reduction genes. *Gammaproteobacteria*, for example, reduce Mn(III) even in the

presence of nitrate (Kostka et al., 1995), and there is precedent for microbial use of multiple

electron acceptors, e.g. "co-respiration" of oxygen and nitrate during aerobic denitrification

(Chen and Strous, 2013; Ji et al., 2015).

Because solid-phase Mn(III) is known to chemically oxidize NH_4^+ (Aigle et al., 2017;

Boumaiza et al., 2018), we tested for abiotic NH_4^+ oxidation by soluble Mn(III) (1 mM).

Ammonium concentrations remained unchanged, and no N_2O or NO_x^- production was observed

(**Fig. S8**), likely because our experiments lacked solid surfaces to mediate electron transfer.

These findings are consistent with lack of detectable ammonium oxidation by Mn(III)

pyrophosphate in estuarine sediments (Crowe et al., 2012).

Electron donors. Methane was the only electron donor added intentionally to the

enrichment cultures. Because we did not detect $^{13}CO_2$ after addition of $^{13}CH_4$ (**Fig. S2**), we

considered alternative electron donors that might have been trace contaminants, but sufficiently

abundant to drive the observed ~300-600 μM Mn(III) reduction (**Fig. 1**). The ethanol catabolism

pathway (PQQ-dependent methanol/ethanol dehydrogenase (RIX45050), quinoprotein alcohol

dehydrogenase (RIX45053), and an NAD^+-dependent aldehyde dehydrogenase-II (RIX45061))

were all highly expressed in "Ca. *D. occultata*" (**Table 1**). Ethanol could have been introduced to

the bottles during culture preparation during sterilization of bottle stoppers. Based on the

stoichiometry of ethanol oxidation coupled to Mn(III) reduction:

$\qquad 2CH_3CH_2OH + 10H_2O + 5Mn(III)_2(P_2O_7)_3 \rightarrow 4HCO_3^- + 10Mn^{2+} + 24H^+ + 15(P_2O_7)^{2-}$

150 μM ethanol would be required to reduce 600 μM of Mn(III), which equates to ~1 μL of 70%

ethanol (12 M) into 100 mL culture medium. We conclude that trace contamination of ethanol

was likely the major electron donor to our cultures.

It is also possible that other substrates, such as H_2 from fermentation by other microbes in

the enrichment or from impurities in the headspace gas, could have supplied another source of

electrons. Indeed, an NAD-reducing hydrogenase (RIX44099-100) was expressed by "*Ca*. D.

occultata" (**Table 1**). Based on the stoichiometry of H_2 oxidation coupled to Mn(III) reduction:

$$H_2 + Mn(III)_2(P_2O_7)_3 \rightarrow 2H^+ + 2Mn^{2+} + 3(P_2O_7)^{2-}$$

600 μM H_2 would be required to reduce 600 μM of Mn(III). Thus, H_2 may have contributed

electrons to Mn(III) reduction, but is not likely sole electron donor. There is precedent for other

metal-reducers simultaneously using H_2 and an organic electron donor (Brown et al., 2005).

Another trace source of organics to our cultures could have been leaching from the rubber

stoppers, which were black bromobutyl and pre-boiled in 0.1 N NaOH. A previous study

reported that organics leaked an array of n-alkanes (C_{16}–C_{34}) and unidentified organic

contaminants in black bromobutyl stoppers (Niemann et al., 2015).

Finally, we considered the possibility that 0.2 mM NH_4^+, added to the cultures as a

nitrogen source, could have provided the electron donor, via an unknown pathway. Based on the

stoichiometry of NH_4^+ oxidation coupled to Mn(III) reduction:

$$NH_4^+ + 3H_2O + 4Mn(III)_2(P_2O_7)_3 \rightarrow NO_3^- + 8H^+ + 8Mn^{2+} + 4(P_2O_7)^{2-}$$

0.2 mM NH_4^+ would supply 1.6 mM electron equivalents, which is more than enough to account

for the observed reduction of 600 μM of Mn(III). This process could operate cryptically if the

oxidized products were reduced to N_2 via denitrification enzymes, such as nitrous oxide

reductase (cNosZ), which was one of the most abundant proteins expressed in Mn(III)-reducing

cultures **(Fig. 2c, Table 1).**

Carbon metabolism. "*Ca*. D. occultata" appeared to be growing mixotrophically. "Ca. *D.*

occultata" encoded several central metabolic pathways, including a complete TCA cycle with a

glyoxylate bypass, an incomplete (acetate-dependent) 3-hydroxypropionate bicycle (3-HP), a

modified Calvin-Benson-Bassham (CBB) pathway, and a pathway for synthesis of

polyhydroxybutyrate **(Fig. S11)**. In addition, "Ca. *D. occultata*" encoded genes for organic

carbon transport, and lactate, acetate, and propionate utilization **(Fig. S11)**. Like *D. agitata* and

D. denitrificans, the CBB pathway of "*Ca.* D. occultata" did not encode RuBisCO and

sedoheptulose-1,7-bisphosphatase (SHbisPase; **Fig. S10**); SHbisPase may be replaced by 6-

phosphofructokinase and an energy-generating pyrophosphatase (RIX41248; Kleiner et al.,

2012; Zorz et al., 2018). The presence of incomplete carbon fixation pathways and for organic

carbon utilization pathways suggests that "Ca. *D. occultata*" relies on organic carbon to fix

inorganic carbon mixotrophically. The source of this organic carbon could have been ethanol,

which is converted to acetate via the pathway discussed in the previous section.

Effect of methane. Although we did not measure appreciable $^{13}CH_4$ oxidation to $^{13}CO_2$,

CH_4 stimulated Mn(III) reduction and cytochrome expression in "*Ca*. D. occultata" enrichment

cultures. While the specific role of CH_4 in Mn(III) reduction remains unknown, the addition of

CH_4 appeared to significantly stimulate expression of many cytochrome *c* proteins, including

OccABGJK, MtoD-2, and cytochrome-*c*4 and -*c*5 proteins associated with anaerobic respiration

($p < 0.05$; **Table 1; Fig. 2C**). Expression of several "*Ca.* D. occultata" proteins involved in outer

membrane structure and composition - including an extracellular DUF4214 protein located next

to an S-layer protein similar to those involved in manganese binding and deposition (Wang et al.,

2009), a serine protease possibly involved in Fe(III) particle attachment (Burns et al., 2009), an

extracellular PEP-CTERM sorting protein for protein export (Haft et al., 2006), and a Tol-Pal

system for outer membrane integrity - was higher in the presence of CH_4 (**Table 1**).

Transporters and sensors. Numerous transporters were present in the "*Ca*. D. occultata"

genome, including 26 TonB-dependent siderophore transporters, 13 TRAP transporters for

dicarboxylate transport, as well as ABC transporters for branched-chained amino acids and

dipeptides and polypeptides (**Table S4**). "*Ca*. D. occultata" also contained a large number of

environmental sensing genes: 52 bacterial hemoglobins with PAS-PAC sensors, 8 TonB-

dependent receptors, and 8 NO responsive regulators (Dnr: Crp/fr family; **Table S4**). Uniquely

in "*Ca*. D. occultata", PAC-PAS sensors flanked accessory genes *nosFLY* on the *c-nosZ* operon

(**Fig. S7**). Comparison of these flanking PAC-PAS sensors in "*Ca*. D. occultata" with O_2-binding

sensors revealed that an arginine ~20 aa upstream from the conserved histidine as the distal

pocket ligand for O_2-binding is not present in either sensor (**Fig. S11**), suggesting that the sensor

may bind a different ligand, possibly NO, consistent with the placement of these genes next to

cNosZ (Shimizu et al., 2015).

244   ***Nutrient storage.*** Active synthesis of storage polymers suggested that "*Ca*. D. occultata"

was experiencing electron acceptor starvation at the time of harvesting, consistent with Mn(III)

depletion in the bottles (Liu et al., 2015; Guanghuan et al., 2018). Polyphosphate-related

proteins, including phosphate transporters, polyphosphate kinase, polyphosphatase, and poly-3-

hydroxybutyrate synthesis machinery were detected in the proteome (**Table S4**). Polyphosphate-

accumulating organisms store polyphosphates with energy generated from organic carbon

oxidation during aerobic respiration or denitrification. These stored compounds are later

hydrolyzed when respiratory electron acceptors for ATP production are limiting. Cyanophycin

was actively synthesized for nitrogen storage.

253   ***Geobacter.*** "*Ca.* Geobacter occultata" expressed proteins in the TCA cycle at moderate

abundance. "*Ca.* G. occultata" contained 17 multiheme c-type cytochromes, none of which were

detected in the proteome. The lack of expression of electron transport and metal-reducing

pathways makes it unlikely that "*Ca.* G. occultata" was solely responsible for Mn(III) reduction

observed in the incubations. A periplasmic group I Ni-Fe hydrogenase (RNC64340; 91% identity

to a protein (RLB64899) from *Geobacter* MAG from terrestrial hot spring sediment) and a type

IV pilin (RNC67631; 10% aromatics, 87% identity to *Geobacter pickeringii* (Holmes et al.,

2016)) were significantly more expressed in the presence of CH_4 than N_2 in the "*Ca.* G.

occultata" proteome (p < 0.05; **Table 1**). It is possible that "*Ca.* G. occultata" transferred

electrons to "*Ca*. D. occultata" via e-pilins (e.g. direct interspecies electron transfer),

contributing to the higher rates of Mn(III) reduction in the presence of CH_4 vs. N_2. The possible

involvement of *Geobacter* e-pilins in Mn(III) reduction remains an open question, due to the lack

of studies examining the possibility of Mn(III) reduction in *Deltaproteobacteria*.

Conclusions. To our knowledge, this study provides the first evidence for biological

reduction of soluble Mn(III) by a bacterium outside of the *Gammaproteobacteria* class. The

dominant bacterium in Mn(III)-reducing enrichment cultures was "*Ca*. D. occultata", a member

of the *Rhodocyclales* order of *Betaproteobacteria*. "*Ca*. D. occultata" expressed decahemes

similar to the Mto pathway, and *occ* genes, including a novel extracellular undecaheme (OccP),

which are predicted to encode a new respiratory electron transport pathway. The novel *occ*

operon was found to be widespread in *Betaproteobacteria* from the deep subsurface, where metal

cycling can fuel microbial metabolism. We also found highly expressed peptides from various

central metabolic cycles and organic substrate utilization pathways, suggesting that "Ca. *D.*

occultata" may have been using multiple pathways simultaneously for energy generation and

carbon assimilation during Mn(III) reduction.

Puzzles remain about whether "*Ca*. D. occultata" can transform two potent greenhouse

gases: methane and nitrous oxide. Although "*Ca*. D. occultata" was enriched with CH_4 as the

sole electron donor and cultures reduced Mn(III) more rapidly in the presence of CH_4, no CH_4

oxidation activity was measured in Mn(III)-reducing cultures, and proteomic data suggested that

"*Ca*. D. occultata" was growing mixotrophically rather than assimilating CH_4. Further, although

we did not add oxidized nitrogen compounds to our media, and Mn(III) did not chemically

oxidize NH_4^+ under our culture conditions, type II nitrous oxide reductase (cNosZ) was one of

the most abundant proteins expressed in Mn(III)-reducing cultures. The role of cNosZ and other

denitrification enzymes in "*Ca*. D. occultata" metabolism, and their possible connection to

Mn(III) reduction, remain to be investigated.

Acknowledgements. This research was funded by NASA Exobiology grant NNX14AJ87G.

Support was also provided by a Center for Dark Energy Biosphere Investigations (NSF-CDEBI

OCE-0939564) small research grant and supported by the NASA Astrobiology Institute

(NNA15BB03A) and a NASA Astrobiology Postdoctoral Fellowship to NS. SAC was supported

through NSERC CRC, CFI, and Discovery grants. We thank Ellery Ingall for dissolved organic

carbon analyses. We thank Marcus Bray, Andrew Burns, Caleb Easterly, Pratik Jagtap, Cory

Padilla, Angela Peña, Johnny Striepen, and Rowan Wolschleger for technical assistance. We

thank Karen Lloyd, Nagissa Mahmoudi, and Emily Weinert for helpful discussions.

Competing Interests: The authors declare no competing interests.

Table 1. Expression levels for select "*Ca*. Dechloromonas occultata" and "*Ca*. Geobacter occultata" proteins
in the presence of CH_4 and N_2. Gray boxes indicate membrane proteins[1]. Bold proteins indicate proteins that were
significantly more expressed with CH_4 than N_2 ($CH_4/N_2>1$; $p<0.05$). P values indicate significance of abundance
difference between CH_4 and N_2 treatments.

---

[1]SP: signal peptide (Y:present/N:absent); TMH: numbers of transmembrane helices; CxxCH: number of heme-binding motifs; P-sort: predicted cellular location based on Psortb v.3.0. MCP: methyl-accepting chemotaxis protein; PPIase: Peptidyl-proline isomerase; P: periplasm, C: cytoplasm; OM: outer membrane; IM: inner membrane, E: extracellular; U: unknown. MtoX and MtoY were predicted to be an inner membrane cytochrome-b protein and a methyl-accepting chemotaxis protein, respectively. Membrane proteins may be underrepresented by mass spectrometry-based metaproteomic analyses, which inherently favor soluble over insoluble membrane-bound or hydrophobic proteins.

Protein Sequence Predictions				Motifs				Normalized Peptide Abundance				Differential Peptide		
Enzyme Complex/ Category	Function	Proteins	NCBI ID	SP	TMH	CxxCH	P-sort	CH₄	SD	N₂	SD	Avg	SD	P value

Enzyme Complex	Function	Proteins	NCBI ID	SP	TMH	CxxCH	P-sort	CH4	SD	N2	SD	Avg	SD	P value
		---------Ca. Dechloromonas occultata---------												
Mto-1	Outer membrane porin-cytochrome c electron conduit	MtoX-1 (cyt-b)	RIX49676	N	5	0	IM							
		MtoY-1 (MCP)	RIX49677	N	2	1	IM	2.7	0.5	3.6	0.2	0.8	0.2	0.2
		MtoB-1 (porin)	RIX49678	Y	0	0	OM	10	2	15	2	0.6	0.1	0.004
		MtoA-1	RIX49874	Y	1	10	P	5	1	2.5	0.1	1.9	0.4	0.1
		MtoD-1	RIX49875	N	0	1	P							
Mto-2	Outer membrane porin-cytochrome c electron conduit	MtoX-2 (cyt-b)	RIX48942	N	4	0	IM							
		MtoB-2 (porin)	RIX48943	Y	0	0	OM	8	1	16	0.2	0.5	0.1	0.04
		MtoA-2	RIX48944	Y	1	10	P	7.3	0.8	4	2	2.1	1.3	0.2
		MtoD-2	RIX48945	Y	1	1	U	2.6	0.3	0.7	0.3	4.0	1.4	0.003
Occ	Membrane-spanning electron transport cytochromes	OccA	RIX49688	Y	1	3	P	4	0.5	0.7	0.6	7.8	5.7	0.01
		OccB	RIX49689	Y	0	3	U	41	4	19	2	2.2	0.0	0.03
		OccC	RIX49877	N	0	1	U							
		OccD	RIX49878	N	0	3	U							
		OccE (6-NHL)	RIX49690	N	1	0	U	22	2.1	20.5	0.2	1.1	0.1	0.2
		OccF	RIX49691	Y	2	4	E	13	0.7	10.1	0.1	1.3	0.1	0.06
		OccG (PPIase)	RIX49692	N	0	0	U	14	1	3.3	0.5	4.2	0.3	0.01
		OccH	RIX49693	N	0	0	OM/E	6.0	0.2	7.7	0.6	0.8	0.1	0.10
		OccI	RIX49694	N	1	3	U	7	2.5	2.3	0.0	2.9	1.1	0.1
		OccJ	RIX49879	Y	0	4	U	44	0.2	19	3	2.4	0.4	0.03
		OccK	RIX49880	N	0	0	C	39	6	13	1	3.0	0.2	0.04
		OccL	RIX49695	N	1	3	U							
		OccM	RIX49881	N	0	3	U							
		OccN (6 NHL)	RIX49696	N	2	0	U	5.7	0.3	6	1	0.9	0.1	0.2
		OccO (6 NHL)	RIX49882	N	0	0	U	1.2	0.8	4.2	0.4	0.3	0.2	0.03
		OccP	RIX49697	N	0	11	E	14	2	12	3	1.2	0.5	0.4
		OccQ	RIX49698	Y	4	0	IM							
		OccR	RIX49883	N	8	0	IM							
		OccS	RIX49699	N	12	0	IM							
Cyt c	Mono- and diheme c-type cytochromes involved in electron transfer	Cyt c5	RIX47670	N	1	1	U	27	2	9	3	3.2	0.8	0.01
		Cyt c5	RIX40984	Y	1	2	P	19	2	6	1	3.3	1.0	0.06
		Cyt c'/C_2	RIX44710	Y	1	1	P	5	3.6	0.8	4.8	2.3	0.09
		Cyt c'/C_2	RIX49630	Y	1	1	P	7	1	1.2	0.9	8.2	6.6	0.07
		Cyt c551/c552	RIX49087	Y	0	1	P	13	3	2.8	0.0	4.8	1.1	0.06
		Cyt c4	RIX48804	Y	0	2	P	16	0.8	9.8	0.8	1.6	0.2	0.06
		Cyt c4	RIX44782	Y	0	2	P	4	2	1.7	0.7	2.6	0.1	0.08
		Cyt c4	RIX45018	Y	0	2	P	7	0.6	2.2	0.2	3.0	0.0	0.02
Nap	Periplasmic nitrate reductase	NapA	RIX41011	Y	0	0	P	76	2	67	3	1.1	0.1	0.1
		NapB	RIX41010	Y	1	2	P	15	1	5	2	3.2	0.9	0.02
		NapC	RIX41009	N	1	4	IM	12	3	13	1	1.0	0.2	0.1
Nir	Nitrite reductase	NirS	RIX44719	Y	0	1	P	58	2	44	4	1.3	0.2	0.1
		NirB	RIX44720	Y	1	2	P	14	3	10	2	1.5	0.6	0.2
		NirC	RIX44788	N	0	1	P							
		NirF	RIX44721	Y	1	0	P or C	2	1	7	1	0.3	0.1	0.02
Nor	Nitric oxide reductase	NorC	RIX45182	N	1	1	IM	3.5	0.7	3.2	0.7	1.1	0.0	0.1
		NorB	RIX45183	N	12	1	IM							
cNos	Type II nitrous oxide reductase	cNosZ	RIX42539	Y	0	0	P	77	17	66	8	1.2	0.3	0.1
		cNosC1	RIX42538	Y	1	1	P	16	2	4	2	4.9	3.3	0.08
		cNosC2	RIX42537	Y	1	2	P	10	0.1	3.9	0.3	2.6	0.1	0.02
		cNosB	RIX42536	N	6	0	IM							
		cNosD	RIX42535	N	0	0	P							
		cNosG	RIX42534	N	1	0	C							
		cNosH	RIX42533	N	4	0	IM							
Qcr	Menaquinol-cytochrome c reductase complex	QcrA	RIX41976	N	9	0	CM							
		QcrB	RIX41977	N	9	0	CM							
		QcrC	RIX41978	N	1	0	CM							
Proteases	Serine protease		RIX49468	N	0	0	P	27	2	1.0	0.3	29.0	9.9	0.02
	Carboxyl-terminal protease (S41)		RIX48818	N	1	0	CM	18.5	0.8	8.0	0.9	2.3	0.1	0.0002
Membrane/Extracellular	DUF4214 protein		RIX44180	N	0	0	OM/E	146	25	43	0.6	3.4	0.5	0.05
	S-layer protein		RIX44181	N	0	0	U	8	0.5	10	0.6	0.8	0.1	0.14
	PEP-CTERM sorting protein		RIX45463	Y	1	0	E	68	6	33	10	2.1	0.5	0.03
	Tol-Pal system protein	TolB	RIX44015	Y	0	0	P	20	2	12	1	1.7	0.0	0.03
	Peptidoglycan-associated lipoprotein	Pal	RIX44016	N	0	0	OM	27.3	0.2	10	3	2.7	0.7	0.04
	Tol-Pal system protein	YbgF	RIX44017	N	0	0	U	10.8	0.4	4	2	3.7	2.2	0.06
	Pilus assembly protein		RIX46961	N	0	0	U	54	5	30	5	1.8	0.1	0.001
Other	Ethanol/methanol dehydrogenase		RIX45050	Y	0	0	P	37	4	17	1	2.2	0.1	0.03
	Alcohol dehydrogenase		RIX45053	Y	0	0	P	12.4	1.4	14.2	1.7	0.9	0.0	0.04
	Aldehyde dehydrogenase		RIX45061	Y	0	0	P	125	31	221	75	0.6	0.1	0.10
	Phasin family granule-associated protein		RIX40682	N	0	0	U	49	2	22	1	2.2	0.2	0.03
	Phasin family granule-associated protein		RIX40683	Y	0	0	U	34	4	16	1	2.1	0.0	0.03
	High potential iron-sulfur protein		RIX49681	Y	0	0	U	10.79	0.01	6.5	0.4	1.7	0.1	0.02
	Electron transfer flavoprotein	FixA	RIX43544	N	0	0	C	16	3	10	2	1.7	0.0	0.04
	NAD-reducing hydrogenase	HoxH	RIX46736	N	0	0	C	22.9	0.7	37	3	0.6	0.1	0.07
		---------Ca. Geobacter occultata---------												
Hydrogenase	[Ni/Fe] hydrogenase, group 1, small subunit	HyaA	RNC64339	Y	0	0	P	11.1	0.4	3	1	5	2	0.06
	[Ni/Fe] hydrogenase, group 1, large subunit	HyaB	RNC64340	N	0	0	P	32	0	11	5	3	1	0.01
E-pilus	Type IV pilin	PilA	RNC67631	N	1	0	E	93	3	18	3	5.6	0.6	0.02

References

Aigle, A., Bonin, P., Iobbi-Nivol, C., Mejean, V., and Michotey, V. (2017) Physiological and transcriptional approaches reveal connection between nitrogen and manganese cycles in *Shewanella algae* C6G3. *Sci Rep* **7**: 44725.

Anantharaman, K., Brown, C.T., Hug, L.A., Sharon, I., Castelle, C.J., Probst, A.J., Thomas, B.C., Singh, A., Wilkins, M.J., Karaoz, U., Brodie, E.L., Williams, K.H., Hubbard, S.S., and Banfield, J.F. (2016) Thousands of microbial genomes shed light on interconnected biogeochemical processes in an aquifer system. *Nat Commun* **7**: 13219.

312 Beal, E.J., House, C.H., and Orphan, V.J. (2009) Manganese- and iron-dependent marine methane
313 oxidation. *Science* **325**: 184-187.

314 Blomberg, M.R., and Siegbahn, P.E. (2014) Proton pumping in cytochrome c oxidase: Energetic
315 requirements and the role of two proton channels. *Biochim Biophys Acta* **1837**: 1165-1177.

316 Boumaiza, H., Coustel, R., Despas, C., Ruby, C., and Bergaoui, L. (2018) Interaction of ammonium with
317 birnessite: Evidence of a chemical and structural transformation in alkaline aqueous medium. *J Solid*
318 *State Chem* **258**: 543-550.

319 Brown, D.G., Komlos, J., and Jaffé, P.R. (2005) Simultaneous utilization of acetate and hydrogen by
320 Geobacter sulfurreducens and implications for use of hydrogen as an indicator of redox conditions.
321 *Environmental science & technology* **39**: 3069-3076.

322 Burns, J.L., Ginn, B.R., Bates, D.J., Dublin, S.N., Taylor, J.V., Apkarian, R.P., Amaro-Garcia, S., Neal,
323 A.L., and DiChristina, T.J. (2009) Outer membrane-associated serine protease involved in adhesion of
324 *Shewanella oneidensis* to Fe(III) oxides. *Environ Sci Technol* **44**: 68-73.

325 Chen, J., and Strous, M. (2013) Denitrification and aerobic respiration, hybrid electron transport chains
326 and co-evolution. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **1827**: 136-144.

327 Crowe, S.A., Canfield, D.E., Mucci, A., Sundby, B., and Maranger, R. (2012) Anammox, denitrification
328 and fixed-nitrogen removal in sediments from the Lower St. Lawrence Estuary. *Biogeosciences* **9**: 4309-
329 4321.

330 Crowe, S.A., O'Neill, A.H., Katsev, S., Hehanussa, P., Haffner, G.D., Sundby, B., Mucci, A., and Fowle,
331 D.A. (2008) The biogeochemistry of tropical lakes: A case study from Lake Matano, Indonesia.
332 *Limnology and Oceanography* **53**: 319-331.

333 Crowe, S.A., Katsev, S., Leslie, K., Sturm, A., Magen, C., Nomosatryo, S., Pack, M.A., Kessler, J.D.,
334 Reeburgh, W.S., Roberts, J.A., Gonzalez, L., Douglas Haffner, G., Mucci, A., Sundby, B., and Fowle,
335 D.A. (2011) The methane cycle in ferruginous Lake Matano. *Geobiology* **9**: 61-78.

336 Cummings, D.E., Caccavo, F., Spring, S., and Rosenzweig, R.F. (1999) Ferribacterium limneticum, gen.
337 nov., sp. nov., an Fe(III)-reducing microorganism isolated from mining-impacted freshwater lake
338 sediments. *Arch Microbiol* **171**: 183-188.

339 Diem, D., and Stumm, W. (1984) Is dissolved Mn2+ being oxidized by O2 in absence of Mn-bacteria or
340 surface catalysts? *Geochimica et Cosmochimica Acta* **48**: 1571-1573.

341 Edwards, M.J., Hall, A., Shi, L., Fredrickson, J.K., Zachara, J.M., Butt, J.N., Richardson, D.J., and
342 Clarke, T.A. (2012) The crystal structure of the extracellular 11-heme cytochrome UndA reveals a
343 conserved 10-heme motif and defined binding site for soluble iron chelates. *Structure* **20**: 1275-1284.

344 Emerson, D., Field, E.K., Chertkov, O., Davenport, K.W., Goodwin, L., Munk, C., Nolan, M., and
345 Woyke, T. (2013) Comparative genomics of freshwater Fe-oxidizing bacteria: implications for
346 physiology, ecology, and systematics. *Front Microbiol* **4**: 254.

347 Ettwig, K.F., Zhu, B., Speth, D., Keltjens, J.T., Jetten, M.S.M., and Kartal, B. (2016) Archaea catalyze
348 iron-dependent anaerobic oxidation of methane. *Proceedings of the National Academy of Sciences* **113**:
349 12792-12796.

Finneran, K.T., Johnsen, C.V., and Lovley, D.R. (2003) *Rhodoferax ferrireducens* sp. nov., a
psychrotolerant, facultatively anaerobic bacterium that oxidizes acetate with the reduction of Fe(III). *Int J*
Syst Evol Microbiol **53**: 669-673.

Fredrickson, J.K., Romine, M.F., Beliaev, A.S., Auchtung, J.M., Driscoll, M.E., Gardner, T.S., Nealson,
K.H., Osterman, A.L., Pinchuk, G., and Reed, J.L. (2008) Towards environmental systems biology of
Shewanella. *Nat Rev Microbiol* **6**: 592.

Gonska, N., Young, D., Yuki, R., Okamoto, T., Hisano, T., Antonyuk, S., Hasnain, S.S., Muramoto, K.,
Shiro, Y., and Tosha, T. (2018) Characterization of the quinol-dependent nitric oxide reductase from the
pathogen *Neisseria meningitidis*, an electrogenic enzyme. *Sci Rep* **8**: 3637.

Guanghuan, G., Jianqiang, Z., Aixia, C., Bo, H., Ying, C., Kun, G., Xiaoling, L., and Xiaoqian, D. (2018)
Nitrogen removal and nitrous oxide emission in an anaerobic/oxic/anoxic sequencing biofilm Batch
reactor. *Environ Eng Sci* **35**: 19-26.

Haft, D.H., Paulsen, I.T., Ward, N., and Selengut, J.D. (2006) Exopolysaccharide-associated protein
sorting in environmental organisms: the PEP-CTERM/EpsH system. Application of a novel phylogenetic
profiling heuristic. *BMC Biol* **4**: 29.

He, S., Barco, R.A., Emerson, D., and Roden, E.E. (2017) Comparative genomic analysis of neutrophilic
iron(II) oxidizer genomes for candidate genes in extracellular electron transfer. *Front Microbiol* **8**: 1584.

Heintze, S., and Mann, P. (1947) Soluble complexes of manganic manganese. *J Agric Sci* **37**: 23-26.

Hem, J.D. (1963) *Chemical equilibria and rates of manganese oxidation*: US Government Printing
Office.

Hernsdorf, A.W., Amano, Y., Miyakawa, K., Ise, K., Suzuki, Y., Anantharaman, K., Probst, A., Burstein,
D., Thomas, B.C., and Banfield, J.F. (2017) Potential for microbial H2 and metal transformations
associated with novel bacteria and archaea in deep terrestrial subsurface sediments. *ISME J* **11**: 1915-
1929.

Holmes, D.E., Dang, Y., Walker, D.J., and Lovley, D.R. (2016) The electrically conductive pili of
Geobacter species are a recently evolved feature for extracellular electron transfer. *Microb Genom* **2**.

Ji, B., Yang, K., Zhu, L., Jiang, Y., Wang, H., Zhou, J., and Zhang, H. (2015) Aerobic denitrification: a
review of important advances of the last 30 years. *Biotechnology and bioprocess engineering* **20**: 643-
651.

Jones, C., Crowe, S.A., Sturm, A., Leslie, K.L., MacLean, L.C.W., Katsev, S., Henny, C., Fowle, D.A.,
and Canfield, D.E. (2011) Biogeochemistry of manganese in ferruginous Lake Matano, Indonesia.
Biogeosciences **8**: 2977-2991.

Jung, H., Taillefert, M., Sun, J., Wang, Q., Borkiewicz, O.J., Liu, P., Yang, L., Chen, S., Chen, H., and
Tang, Y. (2020) Redox Cycling Driven Transformation of Layered Manganese Oxides to Tunnel
Structures. *Journal of the American Chemical Society*.

Kato, S., Ohkuma, M., Powell, D.H., Krepski, S.T., Oshima, K., Hattori, M., Shapiro, N., Woyke, T., and
Chan, C.S. (2015) Comparative genomic insights into ecophysiology of neutrophilic, microaerophilic iron
oxidizing bacteria. *Front Microbiol* **6**: 1265.

Kleiner, M., Wentrup, C., Lott, C., Teeling, H., Wetzel, S., Young, J., Chang, Y.-J., Shah, M.,
VerBerkmoes, N.C., Zarzycki, J., Fuchs, G., Markert, S., Hempel, K., Voigt, B., Becher, D., Liebeke, M.,
Lalk, M., Albrecht, D., Hecker, M., Schweder, T., and Dubilier, N. (2012) Metaproteomics of a gutless
marine worm and its symbiotic microbial community reveal unusual pathways for carbon and energy use.
Proc Natl Acad Sci **109**: 1173-1182.

Kostka, J.E., Luther, G.W., and Nealson, K.H. (1995) Chemical and biological reduction of Mn(III)-
pyrophosphate complexes - potential importance of dissolved Mn(III) as an environmental oxidant.
Geochim Cosmochim Acta **59**: 885-894.

Kuntz, L.B., Laakso, T.A., Schrag, D.P., and Crowe, S.A. (2015) Modeling the carbon cycle in Lake
Matano. *Geobiology* **13**: 454-461.

Learman, D., Wankel, S., Webb, S., Martinez, N., Madden, A., and Hansel, C. (2011) Coupled biotic–
abiotic Mn (II) oxidation pathway mediates the formation and structural evolution of biogenic Mn oxides.
Geochem Cosmochim Acta **75**: 6048-6063.

Leu, A.O., Cai, C., McIlroy, S.J., Southam, G., Orphan, V.J., Yuan, Z., Hu, S., and Tyson, G.W. (2020)
Anaerobic methane oxidation coupled to manganese reduction by members of the *Methanoperedenaceae*.
The ISME Journal **14**: 1030-1041.

Liu, Y., Peng, L., Guo, J., Chen, X., Yuan, Z., and Ni, B.-J. (2015) Evaluating the role of microbial
internal storage turnover on nitrous oxide accumulation during denitrification. *Sci Rep* **5**: 15138.

Luther, G.W., Thibault de Chanvalon, A., Oldham, V.E., Estes, E.R., Tebo, B.M., and Madison, A.S.
(2018) Reduction of Manganese Oxides: Thermodynamic, Kinetic and Mechanistic Considerations for
One- Versus Two-Electron Transfer Steps. *Aquatic Geochemistry* **24**: 257-277.

Luther III, G.W., Madison, A.S., Mucci, A., Sundby, B., and Oldham, V.E. (2015) A kinetic approach to
assess the strengths of ligands bound to soluble Mn (III). *Marine Chemistry* **173**: 93-99.

Madison, A.S., Tebo, B.M., Mucci, A., Sundby, B., and Luther, G.W., 3rd (2013) Abundant porewater
Mn(III) is a major component of the sedimentary redox system. *Science* **341**: 875-878.

Morgan, J.J. (2005) Kinetics of reaction between O2 and Mn (II) species in aqueous solutions.
Geochimica et Cosmochimica Acta **69**: 35-48.

Niemann, H., Steinle, L., Blees, J., Bussmann, I., Treude, T., Krause, S., Elvert, M., and Lehmann, M.F.
(2015) Toxic effects of lab□grade butyl rubber stoppers on aerobic methane oxidation. *Limnol Oceangr:*
Methods **13**: 40-52.

Oldham, V.E., Siebecker, M.G., Jones, M.R., Mucci, A., Tebo, B.M., and Luther, G.W. (2019) The
speciation and mobility of Mn and Fe in estuarine sediments. *Aquat Geochem* **25**: 3-26.

Richardson, D.J., Butt, J.N., Fredrickson, J.K., Zachara, J.M., Shi, L., Edwards, M.J., White, G., Baiden,
N., Gates, A.J., and Marritt, S.J. (2012) The 'porin–cytochrome' model for microbe□to□mineral electron
transfer. *Mol Microbiol* **85**: 201-212.

Riedinger, N., Formolo, M.J., Lyons, T.W., Henkel, S., Beck, A., and Kasten, S. (2014) An inorganic
geochemical argument for coupled anaerobic oxidation of methane and iron reduction in marine
sediments. *Geobiology* **12**: 172-181.

Shi, L., Belchik, S.M., Wang, Z., Kennedy, D.W., Dohnalkova, A.C., Marshall, M.J., Zachara, J.M., and
Fredrickson, J.K. (2011) Identification and characterization of UndAHRCR-6, an outer membrane
endecaheme c-type cytochrome of *Shewanella* sp. strain HRCR-6. *Appl Environ Microbiol* **77**: 5521-
5523.

Shi, L., Dong, H., Reguera, G., Beyenal, H., Lu, A., Liu, J., Yu, H.-Q., and Fredrickson, J.K. (2016)
Extracellular electron transfer mechanisms between microorganisms and minerals. *Nat Rev Microbiol* **14**:
651.

Shimizu, T., Huang, D., Yan, F., Stranava, M., Bartosova, M., Fojtíková, V., and Martínková, M.t. (2015)
Gaseous O2, NO, and CO in signal transduction: structure and function relationships of heme-based gas
sensors and heme-redox sensors. *Chem Rev* **115**: 6491-6533.

Smith, J.A., Lovley, D.R., and Tremblay, P.L. (2013) Outer cell surface components essential for Fe(III)
oxide reduction by Geobacter metallireducens. *Appl Environ Microbiol* **79**: 901-907.

Sturm, A., Fowle, D.A., Jones, C., Leslie, K., Nomosatryo, S., Henny, C., Canfield, D.E., and Crowe,
S.A. (2019) Rates and pathways of CH4 oxidation in ferruginous Lake Matano, Indonesia. *Geobiology*
17: 294-307.

Szeinbaum, N., Burns, J.L., and DiChristina, T.J. (2014) Electron transport and protein secretion
pathways involved in Mn(III) reduction by Shewanella oneidensis. *Environ Microbiol Rep* **6**: 490-500.

Szeinbaum, N., Lin, H., Brandes, J.A., Taillefert, M., Glass, J.B., and DiChristina, T.J. (2017) Microbial
manganese(III) reduction fuelled by anaerobic acetate oxidation. *Environ Microbiol* **19**: 3475-3486.

Tebo, B.M., Johnson, H.A., McCarthy, J.K., and Templeton, A.S. (2005) Geomicrobiology of
manganese(II) oxidation. *Trends Microbiol* **13**: 421-428.

Wang, X., Schröder, H.C., Schloßmacher, U., and Müller, W.E. (2009) Organized bacterial assemblies in
manganese nodules: evidence for a role of S-layers in metal deposition. *Geo-Marine Letters* **29**: 85-91.

Yamaguchi, K.S., and Sawyer, D.T. (1985) The redox chemistry of manganese(III) and manganese(IV)
complexes. *Isr J Chem* **25**: 164-176.

Yang, Y., Chen, J., Qiu, D., and Zhou, J. (2013) Roles of UndA and MtrC of *Shewanella putrefaciens*
W3-18-1 in iron reduction. *BMC Microbiol* **13**: 267.

Zorz, J.K., Kozlowski, J.A., Stein, L.Y., Strous, M., and Kleiner, M. (2018) Comparative proteomics of
three species of ammonia-oxidizing bacteria. *Front Microbiol* **9**: 938.

455
456